Short Title: Exploring functional diversity of tobacco aquaporins.

Author for contact: Annamaria De Rosa (annamaria.derosa@anu.edu.au)

Exploring functional diversity of *Nicotiana tabacum* Aquaporins

Annamaria De Rosa^{1*}, John R Evans¹ and Michael Groszmann¹

¹ARC Centre of Excellence for Translational Photosynthesis, Research School of Biology, Australian National University, Canberra, ACT 2601, Australia

* For correspondence (email annamaria.derosa@anu.edu.au)

Sentence Summary:

Diverse tobacco PIP, TIP and NIP aquaporin isoforms were functionally characterised using high-throughput yeast-based assays, assessing their transport capabilities for key plant solutes: water, H₂O₂, boric acid and urea.

Author Contributions:

MG and JRE conceived original research plans; ADR performed yeast screening experiments and subcellular localisation analyses; ADR and MG performed 3D protein homology modeling analyses; ADR, JRE and MG analyzed the data and wrote the manuscript. ADR agrees to serve as the author responsible for contact and ensures communication.

Funding Information:

This work was funded by the Australian Government through the Australian Research Council Centre of Excellence for Translational Photosynthesis (CE140100015).

Abstract

Aquaporins (AQPs) are involved in a variety of vital plant physiological processes, including water relations, development, stress responses and photosynthesis. *Nicotiana tabacum* (tobacco) has 84 AQP genes due to its recent hybridisation and allotetraploid genome. We functionally characterised a diverse subset of tobacco AQPs spanning the 3 largest AQP subfamilies, selecting nine isoforms from the PIPs, TIPs, and NIPs. Using high-throughput yeast-based functional assays, we determined AQP permeability to water, hydrogen peroxide (H₂O₂), boric acid (BA) and urea. For each AQP, subcellular localisations *in planta* were established using GFP translational fusions. From 3D protein homology modelling, we found that the monomeric pore shape/size, selectivity filter region and NPA motifs is insufficient to comprehensively predict their transport capabilities. PIPs had the narrowest pore diameter and were permeable to water, H₂O₂ and BA. The pore in TIPs was wider and more cylindrical in shape than for the PIPs. TIP1;1s was permeable to all four substrates tested, and is highly expressed in leaves and flowers, suggesting it functions in multiple roles. By contrast, NIP5;1t, with a larger pore size than the NtTIPs, is only expressed in young flowers and enhanced permeability only to BA. Its homolog in Arabidopsis (*AtNIP5;1*) has the same substrate specificity and functions as a boron channel.

Introduction

Aquaporins (AQPs) constitute a major family of integral membrane channel proteins found across all kingdoms of life (Abascal et al. 2014), becoming most diversified in number and subfamilies in plants (Groszmann et al. 2017; Abascal et al. 2014). Plant AQPs play a vital role in diverse physiological processes, including water relations, growth and development, stress responses, and photosynthesis (Hachez et al. 2006; Groszmann et al. 2017; Chaumont and Tyerman 2017). This range in cellular functions reflects a capability for transporting a wide variety of substrates including water, nitrogen compounds (e.g. ammonia, urea and nitrate), gases (e.g. carbon dioxide, oxygen), hydrogen peroxide, metalloids (e.g. boron, silicon), and ions (Gomes et al. 2009; Pommerrenig et al. 2015; Hove and Bhavé 2011; Choi and Roberts 2007; Zwiazek et al. 2017; Byrt et al. 2017; Bienert et al. 2013; Liu et al. 2020). However, functional characterisation of AQPs is quite limited. Elucidating substrate transport profiles of AQPs is a key component for understanding their potential roles in plants and for enabling their use in crop improvement.

Aquaporins assemble as tetrameric complexes, with each monomer forming a functional pore created by six membrane spanning helices, five connecting loops and two shorter helices. At the centre of the AQP tetramer there is a fifth central and functioning pore (Pommerrenig et al. 2015; Kirscht et al. 2016; Törnroth-Horsefield et al. 2006). Although the gross tertiary structure of AQP is highly conserved across organisms, slight deviations in structural and functional characteristics between isoforms contribute to differences in their transport selectivity. Such characteristics include pore dimensions (pore diameter and overall morphology), chemical properties and flexibilities of pore-lining residues, and specific configurations of residues at key constriction points (Luang and Hrmova 2017). Higher plant AQPs divide into five phylogenetically distinct sub-families, namely the Plasma membrane Intrinsic Proteins (PIPs), Tonoplast Intrinsic Proteins (TIPs), Small basic Intrinsic Proteins (SIPs), Nodulin 26-like Intrinsic Proteins (NIPs), and X Intrinsic Proteins (XIPs) (Danielson and Johanson 2008; Johanson and Gustavsson 2002; Kaldenhoff and Fischer 2006). Within each of these sub-families, there can be diversity in permeating substrate selectivity and specific organelle membrane integration (Maurel et al. 2008). Subfamily-specific substrate specificities in plant AQPs have been attributed to diversity in the aromatic arginine (ar/R) selectivity filter (SF) which forms the first constriction site towards the extracellular side of the pore (Hove and Bhavé 2011). Variation in this site largely determines the substrates able to permeate across the membrane through the AQP (Hove and Bhavé 2011; Sui et al. 2001). Dual Asn-Pro-Ala (NPA) motifs located at the centre of the pore act as a second constriction, with variation in residue composition contributing to selectivity for substrates such as ammonia, boric acid (BA) and urea (Wu and Beitz 2007; Hove and Bhavé 2011).

Nicotiana tabacum (tobacco) arose from a recent hybridization and its allotetraploid genome contains 84 AQP genes (De Rosa et al. 2020; Ahmed et al. 2020; Groszmann et al. 2021). The multiplicity of AQP gene isoforms and relatedness to horticulturally important crops such as tomato, potato, eggplant and capsicum, makes tobacco a favourable study species for translating key findings into crops (De Rosa et al. 2020).

Teasing apart the complexities of aquaporin biology *in planta* can be difficult. Plants have many AQP isoforms, with some having redundancy of function under certain environmental conditions (Abascal et al. 2014; Fox et al. 2017). As such, modification of aquaporin expression within the plant, via over expression or down regulation of a specific AQP, may also affect the expression of closely related isoforms (Bi et al. 2015; Kaldenhoff et al. 2007). Functional characterisation of plant AQPs is often accomplished through heterologous expression systems, such as yeast (Kaldenhoff et al. 2007), allowing for assignment of substrate specificity to an AQP homo-tetramer in isolation from other isoforms and other confounding external regulatory mechanisms. We developed a high-throughput micro-cultivation-based yeast assay to study AQP function and used it to develop substrate profiles

for the entire Arabidopsis PIP family (Groszmann et al. in preparation). Here we extend the application of this platform to screen a diverse subset of tobacco AQPs spanning the 3 largest AQP subfamilies (PIPs, TIPs and NIPs). These AQPs isoforms were chosen based on homology to already characterised AQPs in other species or on gene expression characteristics within the plant which might implicate diverse functional roles. Three PIP1 genes were chosen: *NtPIP1;5s* (NtAQP1), *NtPIP1;1t* and *NtPIP1;3t*, each sharing more than 90% homology in gene sequence. *NtPIP1;5s* (NtAQP1) is an established CO₂-permeable AQP isoform, that enhanced CO₂ diffusion and photosynthetic efficiency *in planta* (Uehlein et al. 2003; Flexas et al. 2006). Here we assessed permeability to water, H₂O₂, BA and urea, but not CO₂. Two PIP2 genes, *NtPIP2;4s* and *NtPIP2;5t*, were chosen as representative isoforms from distinct phylogenetic sub-clades within the PIP2 phylogeny. From the TIPs, *NtTIP1;1s* was chosen as a gene highly expressed throughout the plant (De Rosa et al. 2020) and potentially permeable to range of solutes, while *NtTIP2;5t*, a TIP predominantly expressed in roots (De Rosa et al. 2020), had high homology to *AtTIP2;1*, an established transporter of nitrogen compounds (urea and ammonium) (Loqué et al. 2005; Liu et al. 2003). Representative isoforms from two distinct subclasses within the NIP subfamily were selected. *NtNIP5;1s* belongs to the NIP II class and is homologous to the boron-permeable *AtNIP5;1* (Takano et al. 2006; Pommerrenig et al. 2015). *NtNIP2;1s* belongs to a distinct class of NIPs (NIP III), which transports a diverse range of metalloid compounds including the important micronutrients boron and silicon.

We used our high-throughput yeast-based functional system to test for water, hydrogen peroxide (H₂O₂), BA and urea permeability. Subcellular localisation of each AQP was visualised *in planta* with an Arabidopsis expression system. Substrate permeability and sub-cellular localisation data were collated with gene expression data and 3D homology modelling to characterise these tobacco AQPs.

Results

Tobacco Aquaporin localisation in Yeast

Nine diverse NtAQP isoforms, from the PIP, NIP and TIP sub-families (Figure 1) were functionally characterised. N-terminal GFP-AQP fusions were used to establish whether NtAQP proteins integrated into the plasma membrane (PM) of yeast (Figure 2A-I), a necessary condition for attributing AQP-facilitated diffusion into the cell. Optically thin focal plane images of each yeast construct show the localisation of the GFP signal. These images were further processed using surface profiling and line scans of signal intensity to better assess the distribution of GFP-AQP within the yeast cells. Yeast expressing GFP alone (Figure 2J), showed a uniform signal throughout the cell with the exception of the vacuole. The surface and line signal scans show a relatively equal distribution of intensity, consistent with cytosolic localisation. The fusion of NtAQPs to GFP resulted in the redistribution of GFP fluorescence to different yeast sub-cellular compartments including; PM, endoplasmic reticulum (ER), and/or the tonoplast (vacuolar membrane). The GFP-NtPIP1 fusions localised to the periphery of the cell and the ER. Although signal intensity in the periphery varied, likely due to co-localisation in peripheral ER, the signal remained continuous around the cell, consistent with PM integration (Figure 2 A-C). In addition to localisation in the yeast PM, images for *NtPIP1;5s* frequently contained bright spots in the periphery of the cell which are characteristic of ER localisation (Figure 2C). The *NtPIP2* proteins integrated into the PM, with clearly defined peaks present in the line scans, but GFP signal also localised to the ER and faintly inside the vacuole (Figure 2E-D). *NtNIP2;1s* localised to the PM and ER, similar to that observed for the *NtPIP1s* (Figure 2F). *NtNIP5;1* weakly localised to the PM and ER, with signal predominantly associated with integration into the tonoplast (Figure 2G). The NtTIPs had strong signals distributed between the ER, tonoplast and notably the PM (Figure 2H-I). We confirmed PM integration for all NtAQP constructs tested.

Water permeability “Freeze-thaw” assay

Growth curves of *aqy1 aqy2* yeast expressing the Empty vector did not survive exposure to two freeze-thaw cycles, failing to grow after treatments (Figure 3A). By contrast, the growth *aqy1 aqy2* yeast expressing *NtPIP2;4s* subjected to two freeze-thaw cycles was only slightly delayed compared to untreated yeast (Figure 3B). Substantial differences in growth were observed between treated and untreated yeast expressing either a particular NtAQP or empty vector (Figure 3C). Following exposure to two freeze-thaw cycles, the *aqy1 aqy2* yeast expressing the empty vector had only 2% growth relative to the untreated empty vector culture (Figure 3C). *NtPIP2;4s* and *NtPIP2;5t* expression resulted in the greatest yeast growth following the freeze-thaw treatments, achieving 70% of the untreated growth (Figure 3C). *NtTIP1;1s* and *NtTIP2;5t* grew 62% and 30%, respectively, relative to untreated controls. Thus, four out of the nine NtAQPs tested were able to increase the permeability of the PM to water sufficiently to allow the yeast to survive two freeze-thaw treatments.

H₂O₂ toxicity assay

Dose-dependent differences upon exposure to increasing H₂O₂ treatments were observed in growth curves of yeast expressing the empty vector (Figure 4A) or a “H₂O₂-permeable” AQP (Figure 4B). Yeast expressing the empty vector and exposed to 0.25mM or 0.5mM H₂O₂ treatments, had no significant reduction on growth relative to untreated yeast, while 1mM H₂O₂ treatment caused a 37% decrease in growth relative to the untreated control (Figure 4C). By contrast, growth was dramatically reduced in the presence of 0.25mM H₂O₂ for yeast expressing *NtPIP2;4s* (growth reductions were 66%, 86% and 80% at 0.25mM, 0.5mM and 1mM H₂O₂, respectively, Figure 4C). Five NtAQPs increased yeast sensitivity to H₂O₂ exposure compared to empty vector, consistent with these AQPs being classified as H₂O₂ permeable: *NtPIP2;4s*, *NtPIP2;5t*, *NtPIP1;1t*, *NtTIP1;1s* and *NtNIP2;1s*. The lowest concentration of H₂O₂ that resulted in a significant reduction in growth compared to untreated was 0.25mM for *NtPIP2;4s* and *NtPIP2;5t* and 0.5mM for *NtPIP1;1t*, *NtNIP2;1s* and *NtTIP1;1s*.

Boric acid toxicity assay

Yeast expressing the empty vector (Figure 5A) or a “BA-permeable” AQP (Figure 5B) showed reduced growth in the presence of BA. Exposure of yeast expressing the empty vector control to 10mM BA did not impair growth (Figure 5A and 5C). However, greater BA concentrations progressively reduced growth (by 33% and 64% at 20 and 30mM BA, respectively). Three phenotypes were observed across NtAQP-expressing yeast exposed to 20mM and 30mM BA concentrations. The first phenotype displayed a 20-30% reduction in growth relative to the empty vector yeast (*NtTIP1;1s* and *NtTIP2;5t*, Figure 5C), consistent with these AQPs being classified as BA-permeable, enhancing a toxicity response. The second phenotype had a BA toxicity response that was within 10-20% of the empty vector (*NtNIP2;1s*, *NtNIP5;1t* and *NtPIP1;5s*, Figure 5C). Yeast expressing *NtNIP2;1s* had a significant reduction in growth at 10mM or 20mM BA (p<0.05), suggesting moderate sensitivity to BA exposure. The third phenotype, observed in 4 of the 5 PIP-expressing yeasts (*NtPIP1;1t*, *NtPIP1;3t*, *NtPIP2;4s*, *NtPIP2;5t*), resulted in a greater tolerance to BA exposure (average growth 20% greater than empty vector at 20mM and 30mM BA, Figure 5C). The reduced toxicity response associated with expressing any of these 4 NtPIPs could result from increased protein abundance in the PM reducing space available for free membrane diffusion of BA across the PM, thereby decreasing cell permeability to BA. Therefore, although *NtPIP1;5s* did not have a drastic decline in growth compared to the Empty vector control, growth was significantly reduced (~25% at 20mM BA) compared to yeast expressing the other PIPs, likely due to boron permeability.

Urea growth-based assay

Permeability of AQPs to urea was assessed by enhanced growth phenotypes. For *ynvwl* yeast, 12mM urea provided sufficient nitrogen for yeast cultures to reach a growth curve plateau within a ~50 hour incubation (Figure 6A-B, black lines). Growth curves of yeast expressing the empty vector (Figure 6A) or a “urea-permeable” AQP (Figure 6B) show that the expression of the latter enhanced yeast growth at low urea concentrations (2mM and 4mM urea). Yeast expressing the empty vector exhibited a linear growth response to increasing urea concentrations (Figure 6C). Expression of *NtTIP1;1s*, *NtNIP2;1s* or *NtTIP2;5t* resulted in a 50% growth enhancement at both 2mM and 4mM urea compared to yeast expressing an empty vector (Figure 6C). Growth responses for the other 6 NtAQPs (*NtPIP1;1t*, *NtPIP1;5s*, *NtPIP1;3t*, *NtPIP2;4s*, *NtPIP2;5t*, *NtNIP5;1t*), were similar to the yeast expressing the empty vector and these AQPs are presumed not permeable to urea.

In planta sub-cellular localisation of tobacco AQPs

Confocal images of root cortical cells from Arabidopsis plants expressing GFP-NtAQP constructs were obtained (Figure 7). To enhance interpretation, surface plots of a region of GFP intensity near the cell wall are shown at greater magnification (indicated by white dashed box). Organelle-specific marker lines for the plasma membrane, ER and tonoplast are shown in magenta (Figure 7A-C). We observed diversity in AQP membrane integration across the PIP, TIP and NIP subfamilies. The PIPs localised to the PM, with the PIP1s (PIP1;1t, PIP1;3t and PIP1;5s) appearing to have a weaker and more diffuse PM integration when compared to the PIP2s (PIP2;4s and PIP2;5t) which had a sharp and defined GFP signal around the cell’s periphery (Figure 7D-H). The NtNIPs (NIP2;1s and NIP5;1t) also localised to the cell’s periphery. However, their GFP signal was speckled in appearance with distinct localised spots of brighter fluorescence which are characteristic of ER localisation (indicated by white arrow on NIP2;1s surface plot profile of GFP intensity, Figure 7I-J). There was also a wider spread in GFP signal, arising from adjacent PM and ER. The localisation of NtTIPs (TIP1;1s and TIP2;5t) was consistent with integration in the tonoplast, showing a uniform yet diffuse localisation with a wavy topology; also denoted by the presence of internal membranes resembling transvacuolar strands (V, Figure 7K-L).

Protein modelling of aquaporin pores

Tertiary structure homology modelling was used to compare pore diameter and the physico-chemical properties between the 9 NtAQPs functionally characterised in this study. We made pair-wise comparisons between homology models based on the SoPIP2;1 (open; PDB: 2b5f.1.A) and AtTIP2;1 (PDB:5i32.1.A) crystal structure templates to ascertain the most appropriate (see Supplemental Methods S1).

Pore diameter profiles and physico-chemical characteristics of the NtAQPs across the different subfamilies were compared using the highest-confidence 3D homology models (Figure 8-9). All 5 pore diameter profiles for the NtPIPs closely overlapped (blue line, Figure 8A), with the SF region being the narrowest point along the pore (diameter of 2.4Å). The SF residue composition of the NtPIPs was conserved, having Phe-His-Thr-Arg composition in Helix 2 (H2), Helix 5 (H5), Loop E position 1 (LE1) and Loop E position 2 (LE2), respectively. The TIPs also have a conserved pore diameter profile, with the SF region being the narrowest point (dark and light purple lines, Figure 8A). However, the pore shape of the NtTIPs was less undulating in comparison to the PIPs, with the SF region being wider (2.75 Å diameter) and the NPA region being slightly narrower (4.26 Å diameter in TIPs vs. the 4.48 Å of the PIPs; Figure S1 A, B). The SF residue composition differed between the 2 NtTIPs; with NtTIP1;1s having His- Ile- **Ala- Val** vs. NtTIP2;5t having His-Ile-**Gly-Arg** at H2-H5-LE1-LE2 positions (Figure 8C). The NtNIPs that we characterised showed some variation in pore diameter profiles; with NtNIP2;1s having a SF diameter ranging from **3.5-4 Å**, vs. NtNIP5;1t’s SF diameter of **2.6-3.5 Å** (Figure 8A). The remainder of the pore toward the cytosolic side was similar in profile between these two genes. The differences in

SF pore diameter between the NtNIPs were matched with variation in residue composition in this region, with NtNIP2;1s having SF residues: **Gly-Ser-Gly-Arg** and NtNIP5;1t: **Ala-Ile-Ala-Arg** at H2-H5-LE1-LE2 positions respectively. The small size of the residues at the H2 and H5 position (Ser and Gly, respectively) in NtNIP2;1s, contribute to the larger pore diameter at this constriction site (Figure 8Civ). The composition of the NPA motifs (NPA1 and NPA2) is conserved across all of the NtPIPs, NtTIPs and NtNIP2;1s, with the exception of NtNIP5;1t which has NPS at NPA1 and NPV at NPA2 (Figure 8D).

We also characterised physico-chemical properties of the NtAQP pores (Figure 9). The NtPIPs have high homology in residue hydrophobicity and flexibility (Figure 9D). For both PIP1s and PIP2s, conserved regions around the cytosolic pore mouth and the narrowest part of the pore (the SF region) are lined with hydrophilic residues, whereas the remainder of the pore is lined with hydrophobic residues. The flexibility of the PIPs' pore lining residues was also conserved with a band of low-flexibility residues in their SF constriction region while the remainder of the pore having medium-high flexibility residues (Figure 9D). The pore profiles for NtTIPs were nearly cylindrical without the hourglass shape at the selectivity filter, being lined with mostly hydrophobic and less flexible residues (Figure 9B). Variation in flexibility was observed between the 2 NtNIPs, with pore lining residues in NtNIP5;1t being more flexible at the SF than in NtNIP2;1 (Figure 9C). The apoplastic pore entrance of NtNIP2;1 was very flexible and more hydrophilic than NtNIP5;1t (Figure 9C). All the NIPs and TIPs had low flexibility at the NPA central bottleneck region.

Discussion

NtAQP integration into the plasma membrane of yeast cells

A fundamental assumption in permeability assays is that the expressed AQP is present in the plasma membrane. We established this by using GFP translational fusions to visualise NtAQP subcellular localisation for each construct in yeast. Having confirmed that all AQP isoforms integrated into the PM in yeast (Figure 2), we could then infer that alterations to the growth of yeast upon exposure to specific substrates were due to AQP-related changes in PM permeability (Bienert and Chaumont 2014). Although we detected fluorescence consistent with incorporation of GFP tagged NtPIP1s in the PM of yeast (Fig. 2), there are several cases where PIP1 permeability to water has only been observed when PIP1 was co-expressed with a PIP2 (Xenopus oocytes, (Fetter et al. 2004), Yeast, (Groszmann et al. in preparation), reviewed in (Groszmann et al. 2017)). In future work, it would be useful to assess whether co-expression of PIP1s with PIP2s maximises detection of permeability to water or other substrates. Observing both GFP fluorescence consistent with localisation in the PM together with a positive increase in apparent permeability for one substrate enabled us to assign a 'non-permeable' result for other substrates for a given AQP. We therefore propose that confirmation of PM integration should be a necessary checkpoint when functionally testing AQPs in heterologous expression systems.

Relating functional characterisation, sub-cellular localisation, and gene expression of NtAQPs

Having established that the expressed NtAQP were incorporated into the yeast PM, we present the combined permeability results, *in planta* sub-cellular localisation in Arabidopsis root cortical cells, gene expression localisation and protein modelling results in Table 1. The NtAQP data are placed in context with multiple permeabilities published for other AQPs. *In planta* subcellular localisation of the NtAQPs was consistent with their respective AQP subfamilies.

NtPIPs

The PIP subfamily in plants generally has the largest number of isoforms (Anderberg et al. 2012). They are involved in plant water homeostasis through highly selective water transport activity, as well as facilitating diffusion of other small molecules such as glycerol, urea, BA, arsenous acid, H₂O₂, gases and ions (Bienert et al. 2018; Liu et al. 2020). Certain isoforms enhance permeability to multiple molecules e.g. AtPIP2;1 permeable to water, H₂O₂, CO₂ and cations, with functionality dictated by its location within the plant, interacting proteins and associated post-translational modifications (Tyerman et al. 2021).

Both NtPIP2s tested were permeable to water and H₂O₂, with NtPIP2;4s seemingly more efficient at transporting H₂O₂ than PIP2;5t. H₂O₂ permeability was predicted for these two PIP2s by Ahmed et al. (2020) based on substrate specific signature sequences (Azad et al. 2016) (see (Groszmann et al. 2021) to convert between equivalent NtAQP gene names). None of the three NtPIP1s examined here enhanced permeability to water, but were permeable to H₂O₂ (NtPIP1;1t) or BA (NtPIP1;5s) (Table 1). As permeability to H₂O₂ and BA was not predicted for these PIP1s by Ahmed et al. (2020), subtle sequence differences or pore dynamics of these isoforms clearly can modify an AQP's preferential transport to a given substrate (Qiu et al. 2020). Various PIP1 and 2 isoforms from other species are permeable to BA (e.g. ZmPIP1;1 (Dordas et al. 2000), VvPIP1;4 and VvPIP2;3 (Sabir et al. 2014), HvPIP1;3 and 1;4 (Fitzpatrick and Reid 2009), OsPIP2;4 and 2;7 (Kumar et al. 2014)). NtPIP1;5s has been reported to have a low permeability to H₂O₂ using a fluorescence dye-based assay (Navarro-Ródenas et al. 2015), but as we could not detect permeability to H₂O₂, this needs to be confirmed using co-expression of both PIP1 and PIP2.

Combining substrate permeabilities with *in planta* subcellular localisation and tissue specific expression analysis helps elucidate possible physiological roles for these NtAQPs. The results implicate NtPIP2;4s and NtPIP2;5t having roles in regulating water transport across cell membranes in roots and leaves, respectively. Three of the NtPIPs tested here were permeable to H₂O₂, so are likely to be involved with ROS signalling in response to stress (Hachez et al. 2006) by facilitating H₂O₂ diffusion between cells. NtPIP1;5s (NtAQP1) was the first plant AQP shown to permeate CO₂ (Uehlein et al. 2003), facilitating diffusion of CO₂ into the chloroplast during photosynthesis (Flexas et al. 2006). We suggest NtPIP1;5s might have an additional role in boron uptake and distribution throughout the plant (expressed in roots, stems, leaves and flowers), similar to the functional roles reported for other boron-permeable PIPs, HvPIP1;3 and HvPIP1;4 (Fitzpatrick and Reid 2009) and OsPIP2;4 and OsPIP2;7 (Kumar et al. 2014). The boron-transport capability of the PIPs, and their narrow inflexible pore constriction site (2.4 Å SF size vs. 5.14 Å size of BA) suggests that permeability of this substrate could potentially occur through the tetrameric central pore.

NtTIPs

The TIP subfamily regulates the diffusion of water, ammonia, urea and metalloids across the tonoplast (Maurel et al. 1993; Loqué et al. 2005; Liu et al. 2003). Five specialised TIP subgroups have evolved in higher plants (TIP1-TIP5), differing in ar/R filter (SF) composition and substrate specificities (Anderberg et al. 2012; Kirscht et al. 2016). Exemplars from both TIP1 (NtTIP1;1s) and TIP2 (NtTIP2;5t) subgroups localised to the tonoplast *in planta*, where they could increase permeability to water, BA and urea. NtTIP1;1s was also moderately permeable to H₂O₂ but NtTIP2;5t was not. The substrate permeabilities observed here for NtTIP1;1s exactly match those for ZmTIP1;1 (Fox et al. 2017).

The SF region of the NtTIPs was found to be wider than that of the NtPIPs. An extended selectivity filter has been characterised for the TIP subfamily, containing an additional contact residue in Loop C of the AQP monomer (Kirscht et al. 2016), with NtTIP1;1s and NtTIP2;5t having a Phe and His at this position, respectively (De Rosa et al. 2020). The NtTIP1;1 SF composition of a Phe in Loop C and a Val in Loop E2 creates a more hydrophobic environment in the SF compared to that of NtTIP2;5s which

has His and Arg in the same two positions respectively. This results in TIP1;1s being more hydrophobic than TIP2;5t in its constriction region (Figure 9B). While Ahmed *et al.* (2020) correctly predicted TIP1;1s should be permeable to H₂O₂ and urea, our results did not match their prediction for TIP2;5t for H₂O₂ and ammonia. Instead we found TIP2;5t was permeable to urea, but not H₂O₂. The differences in SF composition and hydrophobicity between NtTIP1;1s and NtTIP2;5t might explain their divergences in permeability to H₂O₂ (Table 1) and potentially also ammonia (Kirscht *et al.* 2016). *NtTIP1;1s* is expressed in leaves and flowers whereas *NtTIP2;5t* is predominantly expressed in the roots (having low expression in leaves and flowers). Root-specific gene expression of *TIP2* isoforms was also observed in closely related gene ortholog, tomato *TIP2;5* (De Rosa *et al.* 2020) and more distantly related maize *TIP2s* (Fox *et al.* 2017), indicating potential conservation of function of TIP2s across closely related and diverse species. The proposed functional roles for NtTIP1;1s and NtTIP2;5t include the loading and unloading of urea from vacuolar storage, the storage and translocation of boron, and equilibration of water in tissues where they are expressed (Maurel *et al.* 2015). Furthermore, NtTIP1;1s could also be involved in ROS signalling in response to stress, as it has been suggested that TIPs are involved in cellular detoxification of H₂O₂ (Bienert and Chaumont 2014).

NtNIPs

NIP aquaporins are known to facilitate the transport of small uncharged solutes, such as glycerol, urea and metalloids (Wallace *et al.* 2006). NIPs have a more hydrophobic ar/R selectivity filter, which reduces water permeability in favour of other substrates such as ammonia, urea and metalloids (Wu and Beitz 2007; Hove and Bhavé 2011). There are three sub-classes (NIP I-III), based on ar/R selectivity filter and NPA motif composition (Mitani *et al.* 2008). A representative from each of NIP II (*NtNIP5;1t*) and NIP III (*NtNIP2;1s*) sub-classes were characterised here. GFP tagging demonstrated that both NIP2;1s and NIP5;1s were incorporated into the PM as well as accumulating in the ER *in planta*. The integration in the plant cell PM (versus tonoplast localisation of the TIPs) implies transport of solutes in and out of cells, rather than simply storage/translocation from the vacuole.

NIP II aquaporins tend to have a larger pore diameter than those found in the NIP I sub class, having a substitution of the highly conserved and bulky Trp at the ar/R H2 position for a smaller Ala (Wallace and Roberts 2004). *NtNIP5;1t* had a wider pore diameter compared to that of the tobacco PIP and TIP isoforms (Figure 8). NIP II aquaporins have been shown to permeate BA, glycerol and urea, together with reduced water permeability compared to the NIP Is (Wallace *et al.* 2006; Takano *et al.* 2006; Hanaoka *et al.* 2014; Tanaka *et al.* 2008). By contrast, while *NtNIP5;1t* enhanced permeability to BA, it did not enhance permeability to urea despite their similar size (Table 1) *NtNIP5;1t* expression is highly targeted to young flowers and could be involved in boron redistribution during flower development, similar to the orthologous gene in Arabidopsis (*AtNIP5;1*), which has an established role in boron transport and flower development (Takano *et al.* 2006). Notably, expression of NIP II isoforms, *AtNIP5;1* (in flowers) and *AtNIP6;1* (in basal shoot), is induced in boron-limiting conditions (Takano *et al.* 2006; Tanaka *et al.* 2008). The variation in expression patterns (localisation and stress-responsiveness) reported for PIPs vs. NIP IIs, allude to differences in physiological relevance of their boron transport *in planta*. PIPs could mediate a broad boron uptake and distribution (Fitzpatrick and Reid 2009), however their co-function as water channels subjects them to tight regulation (Chaumont and Tyerman 2014). Boron-permeable PIPs would also provide tolerance to boron toxicity by enabling efflux of excess boron from roots and shoots (Kumar *et al.* 2014; Kumar *et al.* 2018). Unlike the PIPs, water-tight NIP IIs enable highly targeted boron transport in boron-limiting conditions, and are down-regulated in boron sufficient concentrations (Takano *et al.* 2006; Chaumont and Tyerman 2017).

NIP III aquaporins (such as *NtNIP2;1s*) are characterised by an ar/R filter composed of smaller residues (Gly-Ser-Gly-Arg), resulting in an even wider, flexible and more hydrophilic SF (Bansal and Sankaramakrishnan 2007; Mitani-Ueno *et al.* 2011). Our 3D homology modelling showed that *NtNIP2;1s* indeed has a wider pore than all the other NtAQP isoforms characterised (Figure 8),

consistent with this sub-class being permeable to larger substrates, such as silicic acid (4.38 Å diameter) and lactic acid (Mitani-Ueno et al. 2011). Unlike NIP5;1t, NIP2;1s was permeable to multiple substrates, urea, BA and H₂O₂ (low), implicating this NIP isoform in multiple functional roles.

NIP III isoforms occur widely among Graminae, but are not found in all dicots (e.g. absent in Arabidopsis), with evidence suggesting their principal role as facilitators of silicon uptake in plants (Chaumont and Tyerman 2017). In addition to silicon, *ZmNIP2;1* (NIP III) is also permeable to water, urea, BA and H₂O₂ (low) (Fox et al. 2017), suggesting some functional homology in the maize and tobacco isoforms. Expression of *NtNIP2;1* is restricted to young flowers where it is likely to be involved in strategic translocation of small molecules in this target tissue.

Conclusions

This study characterised a diverse set of isoforms in the tobacco AQP family assessing permeability to key solutes for plant growth: water, H₂O₂, urea and boric acid. The functional diversity observed between the NtAQP isoforms highlights complexity in assigning *in planta* function to specific isoforms, with monomeric pore shape/size, SF and NPA motifs alone insufficient to comprehensively predict their transport capabilities.

We observed permeability to each of the substrates tested across three largest AQP subfamilies, with substrate selectivity ranging from none to all substrates tested, indicative of specific or broad functional roles. By using isoforms with sequence homology to well characterised AQPs from other species (*NtTIP2;5t* to *AtTIP2;1* and *NtNIP5;1t* to *AtNIP5;1*), we could correlate sequence similarity to functional homology, as well as identify novel permeating substrates to extend their transport profiles.

We propose that testing transport capability to several substrates could be used to assign *in planta* roles to multifunctional aquaporins.

Methods

Generation of NtAQP phylogeny

MUSCLE-aligned nucleotide sequences identified in De Rosa et al. 2020 and Ahmed et al. 2020, (consensus NtAQP family in (Groszmann et al. 2021)), were used to construct a phylogenetic tree using neighbour-joining method (pair-wise deletion; bootstrap=1000) in MEGA7 software (Kumar et al. 2016).

Generation of NtAQP expression constructs and transformation into yeast

Native sequences of NtAQPs sequences, *NtPIP1;1t* (BK011393), *NtPIP1;3t* (BK011396), *NtPIP1;5s* (BK011398), *NtPIP2;4s* (BK011406), *NtPIP2;5t* (BK011409), *NtTIP1;1s* (BK011426), *NtTIP2;5t* (BK011440), *NtNIP2;1s* (BK011379), *NtNIP5;1t* (BK011387) were commercially synthesised in Gateway-enabled destination vectors. Entry vectors were cloned into three destination vectors from (Alberti et al. 2007): pRS423-GPD with a Histidine3 (HIS3) marker gene for yeast selection, and pRS423-GPD-ccdB-ECFP and pRS426-GPD-eGFP-ccdB both containing Uracil3 (URA3) yeast selection gene. Yeast expression vectors were transformed in respective yeast strains required for functional assays (described below), using the “Frozen-EZ yeast Transformation Kit II” (Zymo Research, Los Angeles, USA). Transformed colonies were grown in Yeast Nitrogen Base, YNB, media (Standard drop out, DO, -URA or -HIS) and spotted on agar YNB (DO -URA or -HIS) selection plates for incubation at 30°C for 2 days, then stored at 4°C. Spotted plates were used for the starting cultures of functional assays.

Confirming plasma membrane integration of NtAQPs in yeast cells

We assessed NtAQP subcellular localisations in yeast cells with AQP-GFP translational fusions to confirm incorporation of the expressed AQP in the yeast plasma membrane (PM). Tobacco AQP:GFP translational fusions were generated via gateway cloning of pUC57 entry vectors with NtAQP coding sequences into pRS426-GPD-EGFP-ccdB yeast expression vector (Alberti et al. 2007); producing N-terminal GFP::NtAQP fusion proteins driven by the constitutive GPD promoter. The GFP-only yeast expression was obtained using the empty vector (no GOI fusion), with eGFP alone constitutively expressed via the GPD promoter. Yeast was grown overnight in 2mL YNB (-URA) (OD_{600} 1-1.5). 1mL aliquots of overnight cultures were sub-cultured and grown 3-4 hours in 2mL of fresh YNB (-URA) media, to ensure imaging of newly formed cells. Yeast (10 μ L) was mounted on a polysine slide with a coverslip sealed with nail polish. Yeast cells were visualised with a Zeiss LSM 780 Confocal microscope using a 40x oil immersion objective (1.2 NA). Light micrographs of yeast cells were acquired using Differential Interference Contrast (DIC), with GFP fluorescence captured using excitation at 488 nm and emission detection across the 490-526 nm range. Images were processed using Fiji (ImageJ) software (Schindelin et al. 2012).

Assessing water, H₂O₂, boric acid and urea permeability using high-throughput yeast-based assays

Yeast-based microculture assays (Groszmann et al. in preparation), were used to test membrane permeability to specific substrates associated with the expression of foreign AQPs which resulted in enhanced or impaired growth. H₂O₂, BA and Urea treatments were previously optimised in (Groszmann et al. in preparation). Yeast growth was monitored using a SPECTROStar nano absorbance microplate reader (BMG Labtech, Germany) at 10-20 minute intervals over 42 to 60 hours. Data collection and processing was consistent between each growth or toxicity-based assay.

Water permeability. We used the *aqy1 aqy2* yeast strain (null *aqy1 aqy2*; background Σ 1278b; genotype: Mat α ; *leu2::hisG*; *trp1::hisG*, *his3::hisG*; *ura352 aqy1D::KanMX aqy2D::KanMX*, provided by Peter Dahl of the S. Hohmann lab) (Tanghe et al. 2002) exploiting the property of yeast cells that show increased freezing tolerance when they express functional water AQPs (Deshmukh et al. 2016). Yeast expressing NtAQPs and the empty vector were grown for 24-28 hours (OD_{650} of 0.5-1) in 1.25mL YNB(-HIS), at 30°C, shaking at 250rpm. Cultures were diluted to 0.6x10⁷ cells/mL in YPD medium and incubated at 30°C for 60 mins. 250 μ L of each culture was aliquoted to 2 Eppendorf tubes; one tube was placed on ice (untreated control) the other was used for two 'Freeze-thaw' cycles. Each 'Freeze thaw' cycle consisted of yeast culture aliquots being frozen in liquid nitrogen for 30 seconds, and thawed in a water bath at 30°C for 20 mins. For each construct, 'Untreated' and 'Treated' yeast were transferred into a 96 well plate (200 μ L aliquots) for growth monitoring.

H₂O₂ permeability was assessed using a reactive oxygen species (ROS) hypersensitive yeast strain, Δ *skn7* (null *skn7*; background BY4741 genotype: Mat α ; *his3 Δ 1 leu2 Δ 0 met15 Δ 0 ura3 Δ 0 Δ SKN7). Obtained from ATCC) (Bienert et al. 2007; Halliwell and Gutteridge 2015; Lee et al. 1999). Yeast's survival was further compromised if AQPs facilitated the diffusion (and accumulation) of H₂O₂ into the cell, which enhanced the toxicity response. Δ *skn7* yeast expressing NtAQPs and Empty vector were grown and diluted to 0.6x10⁷ cells/mL as per the Freeze-thaw assay (above). 200 μ L microcultures of each NtAQP/Empty vector were distributed in 96-well plates with 190 μ L of yeast and 10 μ L H₂O₂ treatments: 0mM/water, 0.25mM, 0.5mM and 1mM H₂O₂.*

Boric acid (BA) toxicity assays were used to assess AQP permeability to BA. Enhanced BA diffusion into yeast expressing AQP lead to greater sensitivity to BA treatments. *aqy1 aqy2* yeast expressing NtAQPs or Empty vector were grown and diluted to 0.6x10⁷ cells/mL. 200 μ L microcultures each NtAQP/Empty vector were distributed in 96-well plates with 180 μ L of yeast and 20 μ L BA treatments: 0mM/water, 10 mM, 20 mM and 30 mM BA.

Urea transport capability. *ynvwI* yeast (null *dur3*; background $\Sigma 23346c$; genotype: Mat α , $\Delta ura3$, $\Delta dur3$, provided by Patrick Bienert of the Nicolaus von Wirén lab) is limited in growth due to a deletion of the DUR3 urea transporter (Liu et al. 2003). Expression of urea-permeable AQPs in *ynvwI* yeast provided a growth advantage when exposed to media containing urea as the sole nitrogen source. *ynvwI* yeast spots were resuspended in 1.25mL of Yeast Basic media (YB, culture medium without nitrogen source) with 2% Glucose. Yeast cultures were diluted to 1.2×10^7 cells/mL. 200 μ L microcultures for each NtAQP/Empty vector construct were distributed in 96-well plates with 190 μ L of yeast and 10 μ L urea treatments: 0mM/water, 1 mM, 4 mM and 12 mM urea.

The yeast microculture OD₆₅₀ readings for the water, H₂O₂, BA and urea assay were processed using a method developed by (Groszmann et al. in preparation). Growth curves were integrated using the natural log of OD₆₅₀/initial OD₆₅₀ ($\ln(OD_t/OD_i)$ vs time) up to a time when the growth rate of the Untreated culture had declined to 5% of its maximum. Area Under the Curve (AUC), was calculated as a proxy that captured the potential growth characteristics affected, regardless of the treatment, in a single parameter.

Characterising *in planta* subcellular localisation of NtAQPs

Tobacco AQP-GFP constructs were generated via Gateway cloning of NtAQP coding sequences from pZeo entry vectors into the pMDC43 destination vector (Curtis and Grossniklaus 2003); N-terminal GFP-NtAQP fusion proteins were driven by the constitutive 2x35S CaMV promoter. Arabidopsis transgenic lines were generated via agrobacterium (GV3101) floral dipping transformation method (Clough and Bent 1998). Seeds were liquid sterilised, washed and sown on Gamorg's B5 medium (0.8% Agar, hygromycin). 8-day-old Arabidopsis seedlings were gently removed from the agar, mounted in phosphate Buffer (100mM NaPO₄ buffer, pH 7.2) on a standard slide, covered with coverslip, and visualised with a Zeiss LSM 780 Confocal microscope using a 40x water immersion objective (1.2 NA). Light micrographs of cortical cells in the root elongation zone were visualised using Differential Interference Contrast (DIC), GFP fluorescence was captured using excitation at 488 nm and emission detection across 490-526 nm. Autofluorescence was detected across 570-674 nm and excluded from GFP detection channel. Images were processed using Fiji (ImageJ) software (Schindelin et al. 2012). Organelle-specific marker lines, established in Nelson et al. (2007) and previously published in De Rosa et al. (2020), were used to guide our interpretation of AQP subcellular localisations.

3D protein homology modelling and characterisation of NtAQP pores

We generated 3D models of the 9 NtAQPs functionally characterised in this study. The Spinach PIP2;1 resolved crystal structure (Törnroth-Horsefield et al. 2006) and Arabidopsis TIP2;1, PDB:5i32.1.A (Kirscht et al. 2016), were used as templates for our modelling analyses. See Supplemental Methods S1 for more background detail.

Acknowledgements

We thank Rosemary White from CSIRO for providing seeds of the Tonoplast:GFP and ER:GFP marker lines (Nelson et al. 2007). The authors acknowledge the facilities and the scientific and technical assistance (Darryl Webb) of Microscopy Australia at the Advanced Imaging Precinct at the Australian National University; a facility funded by the ANU, and State and Federal Governments of Australia. We acknowledge that National Collaborative Research Infrastructure Strategy (NCRIS) of the Australian Government, providing The Australian National University with the growth facilities utilised as part of the Australian Plant Phenomics Facility.

Table 1. Summary of NtAQP properties and other AQP isoforms characterised in other studies.

Permeability to water, H₂O₂, boric acid and urea, molecular diameter (Å) shown in parenthesis; in planta sub-cellular localisation in root cortical cells, Selectivity filter (SF), pore diameter (Å) and gene expression localisations (reported in De Rosa et al. 2020). Red tick (✓) denotes a positive permeability for a specific substrate with the number of ticks indicating the magnitude of phenotypic responses. One, two and three ticks represent a small, medium and large effect, respectively. C denotes water permeable when PIP2 co-expressed with PIP1 and permeability for other substrates when co-expressed. Cross (✖) denotes a negative permeability. Blanks indicate no data is available. Three sub-cellular localisations for the NtAQPs were tested: plasma membrane (PM), endoplasmic reticulum (ER) and tonoplast.

	Water (2.75 Å)	H ₂ O ₂ (2.8 Å)	Boric Acid (5.14 Å)	Urea (5.22 Å)	In planta subcellular localisation	SF pore diameter (Å)	Expression localisation	Ref.
ZmPIP1	C	×	×	×				(Fox et al. 2017)
AtPIP1;1	C	✓	✓✓	×				(Groszmann et al. in preparation)
NtPIP1;1t	×	✓	×	×	PM	2.40	Roots	
NtPIP1;3t	×	×	×	×	PM	2.40	Whole plant	
AtPIP1;4	C	✓✓	×	×				(Groszmann et al. in preparation)
NtPIP1;5s	×	×	✓	×	PM	2.40	Leaves, stem, roots, flowers	
AtPIP2;4	✓	✓	×	×				(Groszmann et al. in preparation)
NtPIP2;4s	✓✓	✓✓✓	×	×	PM	2.40	Roots, flowers (low)	
NtPIP2;5t	✓✓	✓✓	×	×	PM	2.40	Leaves	
ZmPIP2;5	✓✓	✓✓	-	-			Roots, stem, leaves,	(Fox et al. 2017)
AtPIP2;7	✓✓	✓✓	✓	×				(Groszmann et al. in preparation)
ZmNIP1;1	✓✓	✓	✓	×				(Fox et al. 2017)
AtNIP1;2		✓						(Dynowski et al. 2008)
ZmNIP2;1	✓✓	✓	✓✓	✓✓				(Fox et al. 2017)
NtNIP2;1s	×	✓	✓	✓✓	ER + PM	3.50	Young flowers	
AtNIP5;1	✓		✓✓					(Takano et al. 2006; Wang et al. 2017)
NtNIP5;1t	×	×	✓	×	ER + PM	3.26	Young flowers	
AtNIP6;1	×			✓✓				(Wallace and Roberts 2005)
AtTIP1;1	✓✓	✓✓						(Bienert et al. 2007)
NtTIP1;1s	✓✓	✓	✓✓	✓	Tonoplast	2.75	Leaves, flowers	
ZmTIP1;1	✓✓	✓✓	✓✓	✓✓				(Fox et al. 2017)
AtTIP1;3	✓✓		×	✓✓				(Soto et al. 2008)
AtTIP2;1	✓			✓				(Liu et al. 2003)
NtTIP2;5t	✓	×	✓✓	✓	Tonoplast	2.75	Roots, leaves (low), flowers (low)	

542

543

544

545

Figure 1. Phylogeny of the NtAQP family, highlighting genes selected for functional characterisation in this study. The phylogenetic tree was generated using the neighbour-joining method from MUSCLE aligned protein sequences. Confidence levels (%) of branch point generated through bootstrapping analysis (n=1000). Red arrows point to PIP (PIP1;1t, PIP1;3t, PIP1;5s, PIP2;4s, PIP2;5t), TIP (TIP1;1s, TIP2;5t) and NIP (NIP2;1s and NIP5;1t) isoforms functionally characterised in this study.

Figure 2. Subcellular localisation of GFP tagged aquaporins expressed in yeast. Confocal microscopy images of yeast expressing GFP::NtAQP fusions of **A.** NtPIP1;1t, **B.** NtPIP1;3t, **C.** NtPIP1;5s, **D.** NtPIP2;4s, **E.** NtPIP2;5t, **F.** NtNIP2;1s, **G.** NtNIP5;1t, **H.** NtTIP1;1s, **I.** NtTIP2;5t and **J.** Free GFP localisation. For each construct we report a Brightfield + GFP overlay image of a yeast cell; a GFP only image; a surface plot profile of GFP signal intensity at the imaged focal plane; and a line scan of signal intensity traversing the cell (indicated by white arrow in GFP only image). Grey shading in GFP signal line scan corresponds to regions which align with the plasma membrane (PM). PM, endoplasmic reticulum (ER) and vacuole (V) are labelled. Scale bar 2µm.

Figure 3. NtAQP water permeability assessed with the ‘Freeze-thaw’ assay. Yeast growth curves, Ln(OD/OD_i) vs. time, of aqy1 aqy2 yeast expressing **A.** Empty vector control or **B.** a freeze-thaw tolerant AQP (NtPIP2;4s), exposed to freeze-thaw treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** Yeast culture growth following the freeze-thaw treatment (AUC relative to untreated yeast control) for aqy1 aqy2 yeast expressing an Empty vector or one of the 9 NtAQPs. Asterisks denote significantly greater growth following the freeze – thaw treatment compared against Empty vector from an ANOVA with Fishers LSD test: “*” p<0.01 and “****” p<0.001, N=6, Error bars = SE.

Figure 4. NtAQP H₂O₂ permeability assay. Yeast growth curves, Ln(OD/OD_i) vs. time, of snk7 yeast expressing **A.** Empty vector control or **B.** an H₂O₂-sensitive AQP (NtPIP2;4s), exposed to 0.25mM, 0.5mM and 1mM H₂O₂ treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** Yeast culture growth relative to ‘Untreated’ control (AUC relative to untreated), for snk7 yeast expressing an Empty vector or one the 9 NtAQPs. Asterisks denote One-way ANOVA with Fishers LSD test results comparing H₂O₂-treated yeast growth against Empty vector; “*” p<0.05, “***” p<0.01 and “****” p<0.001. N=6, Error bars=SE.

Figure 5. Boric acid permeability assay for yeast expressing NtAQPs. Yeast growth curves, Ln(OD/OD_i) vs. time, of aqy1 aqy2 yeast expressing **A.** Empty vector control or **B.** a boric acid-sensitive AQP (NtTIP1;1s), exposed to 10mM, 20mM and 30mM boric acid treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** Yeast culture growth relative to untreated control (AUC) of aqy1 aqy2 yeast expressing either an Empty vector or one of the 9 NtAQPs exposed to boric acid. Asterisks denote growth that was significantly different to Empty Vector using a One-Way ANOVA with Fishers LSD test; “*” p<0.05, “***” p<0.01 and “****” p<0.001, N=6, Error bars=SE.

Figure 6. Urea permeability assays for yeast expressing NtAQPs. Yeast growth curves, Ln(OD/OD_i) vs. time, of ynvwl yeast expressing **A.** Empty vector control or **B.** a urea-permeable AQP (NtNIP2;1s), exposed to 0mM, 2mM, 4mM or 12mM urea treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** AUC relative to 0mM Urea treatment of ynvwl yeast expressing either an Empty vector or one of the 9 screened NtAQPs. Asterisks denote growth that was significantly greater than Empty vector using a One-way ANOVA with Fishers LSD test, “*” p<0.05, “***” p<0.01 and “****” p<0.001. N=6, Error bars=SE.

Figure 7. In planta sub-cellular localisation of NtAQPs. Confocal images of root cortical cells of transgenic 8-day old Arabidopsis seedlings. GFP marker lines; false coloured purple: **A.** plasma membrane, GFP:PM, **B.** endoplasmic reticulum, GFP:ER, **C.** tonoplast, GFP:tono. **GFP::NtAQP lines: D.** NtPIP1;1t, **E.** NtPIP1;3t, **F.** NtPIP1;5s, **G.** NtPIP2;4s, **H.** NtPIP2;5t, **I.** NtNIP2;1s, **J.** NtNIP5;1t, **K.** NtTIP1;1s and **L.** NtTIP2;5t. A region of the membrane (indicated by white dashed boxes, 5µm x 20µm

dimension) is magnified in the panel below each confocal image to show surface profiles. Transvacuolar strands are denoted by V. White arrows highlight peak intensity discrepancies present in the NIPs assigned to AQP integration into the ER and PM. Scale bar 5µm.

Figure 8. Modelled NtAQP pore features. **A.** Pore profiles of PIPs (PIP1;1t, PIP1;3t, PIP1;5s, PIP2;4s, PIP2;5t; blue), TIP1;1s (light purple), TIP2;5t (dark purple), NIP2;1s (dark green) and NIP5;1t (light green). **B.** A 3D protein model highlighting the Selectivity Filter region (SF, orange residue in 3D Protein model) and NPA region (dark red residues in 3D protein model). **C.** Amino acid residues forming the selectivity filter and the diameter at its narrowest point, viewed perpendicular to the membrane plane from the extracellular side. **D.** NPA motifs: NPA 1 and NPA 2 composition of PIPs, TIP1;1s, TIP2;5t, NIP2;1s and NIP5;1t.

Figure 9. Hydrophobicity and flexibility profiles of NtAQP pores. **A.** 3D Pore model illustrates orientation of the pore profile with the apoplastic entrance (red ball, top), cytosolic entrance (green ball, bottom), residues contacting the pore (yellow balls) and the area inside the AQP pore (blue). Pore profile diameters from ChexVis software (note maximum diameter and pore length scales for each AQP), superimposed with Hydrophobicity (Left hand, Blue to Red indicating low to high hydrophobicity, respectively) and Flexibility (Right hand, Blue to Red indicating low to high flexibility, respectively): **B.** NtTIPs (NtTIP1;1s and NtTIP25t.) **C.** NtNIPs (NtNIP2;1s, NtNIP5;1t) and **D.** NtPIPs (NtPIP1;1t, NtPIP1;3t, NtPIP1;5s, NtPIP2;4s, NtPIP2;5t). Black dots to the left of each pore profile correspond to contact residue interactions for which hydrophobicity and flexibility outputs were generated. Red brackets indicate Selectivity Filter (SF) region and grey brackets indicate NPA region.

References

- Abascal F, Irisarri I, Zardoya R (2014) Diversity and evolution of membrane intrinsic proteins. *Biochimica et Biophysica Acta (BBA)-General Subjects* 1840 (5):1468-1481
- Ahmed J, Mercx S, Boutry M, Chaumont F (2020) Evolutionary and predictive functional insights into the aquaporin gene family in the allotetraploid plant *Nicotiana tabacum*. *International journal of molecular sciences* 21 (13):4743
- Alberti S, Gitler AD, Lindquist S (2007) A suite of Gateway® cloning vectors for high-throughput genetic analysis in *Saccharomyces cerevisiae*. *Yeast* 24 (10):913-919
- Anderberg HI, Kjellbom P, Johanson U (2012) Annotation of *Selaginella moellendorffii* major intrinsic proteins and the evolution of the protein family in terrestrial plants. *Frontiers in plant science* 3:33
- Azad AK, Ahmed J, Alum MA, Hasan MM, Ishikawa T, Sawa Y, Katsuhara M (2016) Genome-wide characterization of major intrinsic proteins in four grass plants and their non-aqua transport selectivity profiles with comparative perspective. *PLoS One* 11 (6):e0157735
- Bansal A, Sankararamakrishnan R (2007) Homology modeling of major intrinsic proteins in rice, maize and *Arabidopsis*: comparative analysis of transmembrane helix association and aromatic/arginine selectivity filters. *BMC Structural Biology* 7 (1):27
- Bi Z, Merl-Pham J, Uehlein N, Zimmer I, Mühlhans S, Aichler M, Walch AK, Kaldenhoff R, Palme K, Schnitzler J-P (2015) RNAi-mediated downregulation of poplar plasma membrane intrinsic proteins (PIPs) changes plasma membrane proteome composition and affects leaf physiology. *Journal of proteomics* 128:321-332
- Bienert GP, Chaumont F (2014) Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochimica et Biophysica Acta (BBA)-General Subjects* 1840 (5):1596-1604
- Bienert GP, Desguin B, Chaumont F, Hols P (2013) Channel-mediated lactic acid transport: a novel function for aquaglyceroporins in bacteria. *Biochemical Journal* 454 (3):559-570
- Bienert GP, Møller AL, Kristiansen KA, Schulz A, Møller IM, Schjoerring JK, Jahn TP (2007) Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *Journal of Biological Chemistry* 282 (2):1183-1192
- Bienert MD, Diehn TA, Richet N, Chaumont F, Bienert GP (2018) Heterotetramerization of plant PIP1 and PIP2 aquaporins is an evolutionary ancient feature to guide PIP1 plasma membrane localization and function. *Frontiers in plant science* 9:382
- Byrt CS, Zhao M, Kourghi M, Bose J, Henderson SW, Qiu J, Gilliam M, Schultz C, Schwarz M, Ramesh SA (2017) Non-selective cation channel activity of aquaporin AtPIP2; 1 regulated by Ca²⁺ and pH. *Plant, cell & environment* 40 (6):802-815
- Chaumont F, Tyerman S (2017) *Plant Aquaporins*. Springer,
- Chaumont F, Tyerman SD (2014) Aquaporins: Highly regulated channels controlling plant water relations. *Plant Physiology* 164 (4):1600-1618. doi:10.1104/pp.113.233791
- Choi W-G, Roberts DM (2007) *Arabidopsis* NIP2; 1, a major intrinsic protein transporter of lactic acid induced by anoxic stress. *Journal of Biological Chemistry* 282 (33):24209-24218
- Curtis MD, Grossniklaus U (2003) A gateway cloning vector set for high-throughput functional analysis of genes in planta. *Plant physiology* 133 (2):462-469
- Danielson JA, Johanson U (2008) Unexpected complexity of the aquaporin gene family in the moss *Physcomitrella patens*. *BMC Plant Biology* 8 (1):45
- De Rosa A, Watson-Lazowski A, Evans JR, Groszmann M (2020) Genome-wide identification and characterisation of Aquaporins in *Nicotiana tabacum* and their relationships with other Solanaceae species. *BMC Plant Biology* 20 (1):1-29
- Deshmukh RK, Sonah H, Bélanger RR (2016) Plant Aquaporins: genome-wide identification, transcriptomics, proteomics, and advanced analytical tools. *Frontiers in plant science* 7:1896

Dordas C, Chrispeels MJ, Brown PH (2000) Permeability and channel-mediated transport of boric acid across membrane vesicles isolated from squash roots. *Plant physiology* 124 (3):1349-1362

Dynowski M, Schaaf G, Loque D, Moran O, Ludewig U (2008) Plant plasma membrane water channels conduct the signalling molecule H₂O₂. *Biochemical Journal* 414 (1):53-61

Fetter K, Van Wilder V, Moshelion M, Chaumont F (2004) Interactions between plasma membrane aquaporins modulate their water channel activity. *The Plant Cell* 16 (1):215-228

Fitzpatrick KL, Reid RJ (2009) The involvement of aquaglyceroporins in transport of boron in barley roots. *Plant, cell & environment* 32 (10):1357-1365

Flexas J, Ribas-Carbó M, Hanson DT, Bota J, Otto B, Cifre J, McDowell N, Medrano H, Kaldenhoff R (2006) Tobacco aquaporin NtAQP1 is involved in mesophyll conductance to CO₂ in vivo. *The Plant Journal* 48 (3):427-439

Fox AR, Maistriaux LC, Chaumont F (2017) Toward understanding of the high number of plant aquaporin isoforms and multiple regulation mechanisms. *Plant Science* 264:179-187

Gomes D, Agasse A, Thiébaud P, Delrot S, Gerós H, Chaumont F (2009) Aquaporins are multifunctional water and solute transporters highly divergent in living organisms. *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1788 (6):1213-1228

Groszmann M, De Rosa A, Ahmed J, Chaumont F, Evans JR (2021) A consensus on the Aquaporin Gene Family in the Allotetraploid Plant, *Nicotiana tabacum*. *Plant Direct* doi:10.1002/PLD3.321

Groszmann M, De Rosa A, Chen W, Qiu J, Byrt CS, Evans JR (in preparation) Characterising the permeability of all 13 Arabidopsis PIP genes using a high throughput yeast approach. *Plant Physiology*

Groszmann M, Osborn HL, Evans JR (2017) Carbon dioxide and water transport through plant aquaporins. *Plant, Cell & Environment* 40 (6):938-961

Hachez C, Zelazny E, Chaumont F (2006) Modulating the expression of aquaporin genes in planta: a key to understand their physiological functions? *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1758 (8):1142-1156

Halliwell B, Gutteridge JM (2015) Free radicals in biology and medicine. Oxford University Press, USA,

Hanaoka H, Uruguchi S, Takano J, Tanaka M, Fujiwara T (2014) OsNIP3; 1, a rice boric acid channel, regulates boron distribution and is essential for growth under boron-deficient conditions. *The Plant Journal* 78 (5):890-902

Hove RM, Bhavé M (2011) Plant aquaporins with non-aqua functions: deciphering the signature sequences. *Plant Molecular Biology* 75 (4):413-430. doi:10.1007/s11103-011-9737-5

Johanson U, Gustavsson S (2002) A new subfamily of major intrinsic proteins in plants. *Molecular Biology and Evolution* 19 (4):456-461

Kaldenhoff R, Bertl A, Otto B, Moshelion M, Uehlein N (2007) Characterization of plant aquaporins. In: *Methods in enzymology*, vol 428. Elsevier, pp 505-531

Kaldenhoff R, Fischer M (2006) Aquaporins in plants. *Acta Physiologica* 187 (1-2):169-176

Kirscht A, Kaptan SS, Bienert GP, Chaumont F, Nissen P, de Groot BL, Kjellbom P, Gourdon P, Johanson U (2016) Crystal structure of an ammonia-permeable aquaporin. *PLoS biology* 14 (3):e1002411

Kumar K, Mosa KA, Chhikara S, Musante C, White JC, Dhankher OP (2014) Two rice plasma membrane intrinsic proteins, OsPIP2; 4 and OsPIP2; 7, are involved in transport and providing tolerance to boron toxicity. *Planta* 239 (1):187-198

Kumar K, Mosa KA, Meselhy AG, Dhankher OP (2018) Molecular insights into the plasma membrane intrinsic proteins roles for abiotic stress and metalloids tolerance and transport in plants. *Indian Journal of Plant Physiology* 23 (4):721-730

Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular biology and evolution* 33 (7):1870-1874

- Lee J, Godon C, Lagniel G, Spector D, Garin J, Labarre J, Toledano MB (1999) Yap1 and Skn7 control two specialized oxidative stress response regulons in yeast. *Journal of Biological Chemistry* 274 (23):16040-16046
- Liu L-H, Ludewig U, Gassert B, Frommer WB, von Wirén N (2003) Urea transport by nitrogen-regulated tonoplast intrinsic proteins in Arabidopsis. *Plant physiology* 133 (3):1220-1228
- Liu S, Fukumoto T, Gena P, Feng P, Sun Q, Li Q, Matsumoto T, Kaneko T, Zhang H, Zhang Y (2020) Ectopic expression of a rice plasma membrane intrinsic protein (OsPIP1; 3) promotes plant growth and water uptake. *The Plant Journal* 102 (4):779-796
- Loqué D, Ludewig U, Yuan L, von Wirén N (2005) Tonoplast intrinsic proteins AtTIP2; 1 and AtTIP2; 3 facilitate NH₃ transport into the vacuole. *Plant physiology* 137 (2):671-680
- Luang S, Hrmova M (2017) Structural basis of the permeation function of plant aquaporins. In: *Plant Aquaporins*. Springer, pp 1-28
- Maurel C, Boursiac Y, Luu D-T, Santoni V, Shahzad Z, Verdoucq L (2015) Aquaporins in plants. *Physiological reviews* 95 (4):1321-1358
- Maurel C, Reizer J, Schroeder JI, Chrispeels MJ (1993) The vacuolar membrane protein gamma-TIP creates water specific channels in *Xenopus* oocytes. *The EMBO Journal* 12 (6):2241-2247
- Maurel C, Verdoucq L, Luu D-T, Santoni V (2008) Plant aquaporins: membrane channels with multiple integrated functions. *Annu Rev Plant Biol* 59:595-624
- Mitani-Ueno N, Yamaji N, Zhao F-J, Ma JF (2011) The aromatic/arginine selectivity filter of NIP aquaporins plays a critical role in substrate selectivity for silicon, boron, and arsenic. *Journal of experimental botany* 62 (12):4391-4398
- Mitani N, Yamaji N, Ma JF (2008) Characterization of substrate specificity of a rice silicon transporter, Lsi1. *Pflügers Archiv-European Journal of Physiology* 456 (4):679-686
- Navarro-Ródenas A, Xu H, Kempainen M, Pardo AG, Zwiazek JJ (2015) Laccaria bicolor aquaporin LbAQP1 is required for H₂O net development in trembling aspen (*Populus tremuloides*). *Plant, cell & environment* 38 (11):2475-2486
- Nelson BK, Cai X, Nebenführ A (2007) A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. *The Plant Journal* 51 (6):1126-1136
- Pommerrenig B, Diehn TA, Bienert GP (2015) Metalloids-porins: Essentiality of Nodulin 26-like intrinsic proteins in metalloid transport. *Plant Science* 238:212-227
- Qiu J, McGaughey SA, Groszmann M, Tyerman SD, Byrt CS (2020) Phosphorylation influences water and ion channel function of AtPIP2; 1. *Plant, Cell & Environment* 43 (10):2428-2442
- Sabir F, Leandro MJ, Martins AP, Loureiro-Dias MC, Moura TF, Soveral G, Prista C (2014) Exploring three PIPs and three TIPs of grapevine for transport of water and atypical substrates through heterologous expression in aqy-null yeast. *PloS one* 9 (8):e102087
- Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B (2012) Fiji: an open-source platform for biological-image analysis. *Nature methods* 9 (7):676
- Soto G, Alleva K, Mazzella MA, Amodeo G, Muschietti JP (2008) AtTIP1; 3 and AtTIP5; 1, the only highly expressed Arabidopsis pollen-specific aquaporins, transport water and urea. *FEBS letters* 582 (29):4077-4082
- Sui H, Han B-G, Lee JK, Walian P, Jap BK (2001) Structural basis of water-specific transport through the AQP1 water channel. *Nature* 414 (6866):872
- Takano J, Wada M, Ludewig U, Schaaf G, Von Wirén N, Fujiwara T (2006) The Arabidopsis major intrinsic protein NIP5; 1 is essential for efficient boron uptake and plant development under boron limitation. *The Plant Cell* 18 (6):1498-1509
- Tanaka M, Wallace IS, Takano J, Roberts DM, Fujiwara T (2008) NIP6; 1 is a boric acid channel for preferential transport of boron to growing shoot tissues in Arabidopsis. *The Plant Cell* 20 (10):2860-2875

Tanghe A, Van Dijck P, Dumortier F, Teunissen A, Hohmann S, Thevelein JM (2002) Aquaporin expression correlates with freeze tolerance in baker's yeast, and overexpression improves freeze tolerance in industrial strains. *Appl Environ Microbiol* 68 (12):5981-5989

Törnroth-Horsefield S, Wang Y, Hedfalk K, Johanson U, Karlsson M, Tajkhorshid E, Neutze R, Kjellbom P (2006) Structural mechanism of plant aquaporin gating. *Nature* 439 (7077):688

Tyerman SD, McGaughey SA, Qiu J, Yool AJ, Byrt CS (2021) Adaptable and multifunctional ion-conducting aquaporins. *Annual Review of Plant Biology* 72

Uehlein N, Lovisolo C, Siefritz F, Kaldenhoff R (2003) The tobacco aquaporin NtAQP1 is a membrane CO₂ pore with physiological functions. *Nature* 425 (6959):734-737

Wallace IS, Choi W-G, Roberts DM (2006) The structure, function and regulation of the nodulin 26-like intrinsic protein family of plant aquaglyceroporins. *Biochimica et Biophysica Acta (BBA)-Biomembranes* 1758 (8):1165-1175

Wallace IS, Roberts DM (2004) Homology modeling of representative subfamilies of Arabidopsis major intrinsic proteins. Classification based on the aromatic/arginine selectivity filter. *Plant Physiology* 135 (2):1059-1068

Wallace IS, Roberts DM (2005) Distinct transport selectivity of two structural subclasses of the nodulin-like intrinsic protein family of plant aquaglyceroporin channels. *Biochemistry* 44 (51):16826-16834

Wang S, Yoshinari A, Shimada T, Hara-Nishimura I, Mitani-Ueno N, Ma JF, Naito S, Takano J (2017) Polar localization of the NIP5; 1 boric acid channel is maintained by endocytosis and facilitates boron transport in Arabidopsis roots. *The Plant Cell* 29 (4):824-842

Wu B, Beitz E (2007) Aquaporins with selectivity for unconventional permeants. *Cellular and molecular life sciences* 64 (18):2413-2421

Zwiazek JJ, Xu H, Tan X, Navarro-Ródenas A, Morte A (2017) Significance of oxygen transport through aquaporins. *Scientific reports* 7:40411

Figure 1. Phylogeny of the NtAQP family, highlighting genes selected for functional characterisation in this study. The phylogenetic tree was generated using the neighbour-joining method from MUSCLE aligned protein sequences. Confidence levels (%) of branch point generated through bootstrapping analysis (n=1000). Red arrows point to PIP (PIP1;1t, PIP1;3t, PIP1;5s, PIP2;4s, PIP2;5t), TIP (TIP1;1s, TIP2;5t) and NIP (NIP2;1s and NIP5;1t) isoforms functionally characterised in this study.

Figure 2. Subcellular localisation of GFP tagged aquaporins expressed in yeast. Confocal microscopy images of yeast expressing GFP::NtAQP fusions of **A.** NtPIP1;1t, **B.** NtPIP1;3t, **C.** NtPIP1;5s, **D.** NtPIP2;4s, **E.** NtPIP2;5t, **F.** NtNIP2;1s, **G.** NtNIP5;1t, **H.** NtTIP1;1s, **I.** NtTIP2;5t and **J.** Free GFP localisation. For each construct we report a Brightfield + GFP overlay image of a yeast cell; a GFP only image; a surface plot profile of GFP signal intensity at the imaged focal plane; and a line scan of signal intensity traversing the cell (indicated by white arrow in GFP only image). Grey shading in GFP signal line scan corresponds to regions which align with the plasma membrane (PM). PM, endoplasmic reticulum (ER) and vacuole (V) are labelled. Scale bar 2 μ m.

Figure 3. NtAQP water permeability assessed with the 'Freeze-thaw' assay. Yeast growth curves, Ln(OD/OD_i) vs. time, of aqy1 aqy2 yeast expressing **A.** Empty vector control or **B.** a freeze-thaw tolerant AQP (NtPIP2;4s), exposed to freeze-thaw treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** Yeast culture growth following the freeze-thaw treatment (AUC relative to untreated yeast control) for aqy1 aqy2 yeast expressing an Empty vector or one of the 9 NtAQPs. Asterisks denote significantly greater growth following the freeze – thaw treatment compared against Empty vector from an ANOVA with Fishers LSD test: “***” p<0.01 and “****” p<0.001, N=6, Error bars = SE.

Figure 4. NtAQP H₂O₂ permeability assay. Yeast growth curves, Ln(OD/OD_i) vs. time, of *snk7* yeast expressing **A.** Empty vector control or **B.** an H₂O₂-sensitive AQP (NtPIP2;4s), exposed to 0.25mM, 0.5mM and 1mM H₂O₂ treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** Yeast culture growth relative to 'Untreated' control (AUC relative to untreated), for *snk7* yeast expressing an Empty vector or one the 9 NtAQPs. Asterisks denote One-way ANOVA with Fishers LSD test results comparing H₂O₂-treated yeast growth against Empty vector; "*" p<0.05, "***" p<0.01 and "****" p<0.001. N=6, Error bars=SE.

Figure 5. Boric acid permeability assay for yeast expressing NtAQPs. Yeast growth curves, $\text{Ln}(\text{OD}/\text{OD}_i)$ vs. time, of aqy1 aqy2 yeast expressing **A.** Empty vector control or **B.** a boric acid-sensitive AQP (NtTIP1;1s), exposed to 10mM, 20mM and 30mM boric acid treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** Yeast culture growth relative to untreated control (AUC) of aqy1 aqy2 yeast expressing either an Empty vector or one of the 9 NtAQPs exposed to boric acid. Asterisks denote growth that was significantly different to Empty Vector using a One-Way ANOVA with Fishers LSD test; “*” $p < 0.05$, “***” $p < 0.01$ and “****” $p < 0.001$, $N=6$, Error bars=SE.

Figure 6. Urea permeability assays for yeast expressing NtAQPs. Yeast growth curves, Ln(OD/OD_i) vs. time, of ynvwl yeast expressing **A.** Empty vector control or **B.** a urea-permeable AQP (NtNIP2;1s), exposed to 0mM, 2mM, 4mM or 12mM urea treatments. Growth was assessed from the area under the curves (AUC) until the vertical dashed lines. **C.** AUC relative to 0mM Urea treatment of ynvwl yeast expressing either an Empty vector or one of the 9 screened NtAQPs. Asterisks denote growth that was significantly greater than Empty vector using a One-way ANOVA with Fishers LSD test, “*” p<0.05, “**” p<0.01 and “***” p<0.001. N=6, Error bars=SE.

Figure 7. In planta sub-cellular localisation of NtAQPs. Confocal images of root cortical cells of transgenic 8-day old Arabidopsis seedlings. GFP marker lines; false coloured purple: **A.** plasma membrane, GFP:PM, **B.** endoplasmic reticulum, GFP:ER, **C.** tonoplast, GFP:tono. **GFP:NtAQP** lines: **D.** NtPIP1;1t, **E.** NtPIP1;3t, **F.** NtPIP1;5s, **G.** NtPIP2;4s, **H.** NtPIP2;5t, **I.** NtNIP2;1s, **J.** NtNIP5;1t, **K.** NtTIP1;1s and **L.** NtTIP2;5t. A region of the membrane (indicated by white dashed boxes, 5µm x 20µm dimension) is magnified in the panel below each confocal image to show surface profiles. Transvacuolar strands are denoted by V. White arrows highlight peak intensity discrepancies present in the NIPs assigned to AQP integration into the ER and PM. Scale bar 5µm.

Figure 8. Modelled NtAQP pore features. **A.** Pore profiles of PIPs (PIP1;1 t, PIP1;3t, PIP1;5s, PIP2;4s, PIP2;5t; blue), TIP1;1s (light purple), TIP2;5t (dark purple), NIP2;1s (dark green) and NIP5;1t (light green). **B.** A 3D protein model highlighting the Selectivity Filter region (SF, orange residue in 3D Protein model) and NPA region (dark red residues in 3D protein model). **C.** Amino acid residues forming the selectivity filter and the diameter at its narrowest point, viewed perpendicular to the membrane plane from the extracellular side. **D.** NPA motifs: NPA 1 and NPA 2 composition of PIPs, TIP1;1s, TIP2;5t, NIP2;1s and NIP5;1t.

Figure 9. Hydrophobicity and flexibility profiles of NtAQP pores. **A.** 3D Pore model illustrates orientation of the pore profile with the apoplastic entrance (red ball, top), cytosolic entrance (green ball, bottom), residues contacting the pore (yellow balls) and the area inside the AQP pore (blue). Pore profile diameters from ChexVis software (note maximum diameter and pore length scales for each AQP), superimposed with Hydrophobicity (Left hand, Blue to Red indicating low to high hydrophobicity, respectively) and Flexibility (Right hand, Blue to Red indicating low to high flexibility, respectively): **B.** NtTIPs (NtTIP1;1s and NtTIP25t.) **C.** NtNIPs (NtNIP2;1s, NtNIP5;1t) and **D.** NtPIPs (NtPIP1;1t, NtPIP1;3t, NtPIP1;5s, NtPIP2;4s, NtPIP2;5t). Black dots to the left of each pore profile correspond to contact residue interactions for which hydrophobicity and flexibility outputs were generated. Red brackets indicate Selectivity Filter (SF) region and grey brackets indicate NPA region.