Generation and first characterization of TRDC-Knockout pigs lacking $\gamma\delta$ T cells

Bjoern Petersen^{1*}, Robert Kammerer^{2*}, Antje Frenzel¹, Petra Hassel¹, Tung Huy Dau², Roswitha Becker¹, Angele Breithaupt³, Reiner Georg Ulrich⁴, Andrea Lucas-Hahn¹ and Gregor Meyers²

¹Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut, Mariensee, Neustadt, Germany

²Institute of Immunology, Friedrich-Loeffler-Institut, Greifswald, Insel Riems, Germany

³Department of Experimental Animal Facilities and Biorisk Management, Friedrich-Loeffler-Institut, Greifswald, Insel Riems, Germany

⁴Institute for Veterinary-Pathology, Leipzig University, Leipzig, Germany

* Authors contributed equally

Correspondence to:

Bjoern Petersen
Institute of Farm Animal Genetics
Friedrich-Loeffler-Institut
Hoeltystasse 10
31535 Neustadt am Rbge./Mariensee,
Germany
bjoern.petersen@fli.de

or Robert Kammerer
Institute of Immunology
Friedrich-Loeffler-Institut
Suedufer 10
17493 Greifswald-Insel Riems,
Germany
robert.kammerer@fli.de

AF: antje.frenzel@fli.de
PH: petra.hassel@fli.de
THD: HuyTung.Dau@fli.de
RB: roswitha.becker@fli.de
AB: angele.breithaupt@fli.de
RU: reiner.ulrich@vetmed.uni-leipzig.de
ALH: andrea.lucas-hahn@fli.de
GM: gregor.meyers@fli.de

Running title: $\gamma\delta$ T cell knockout pigs

Keywords: TRDC-knockout, $\gamma\delta$ T cell, TCR, Immunity, SCNT, CRISPR/Cas

1 **Abstract**

2 The TRDC-Locus encodes the T cell receptor delta constant region, one component of the
3 $\gamma\delta$ T cell receptor which is essential for development of $\gamma\delta$ T cells. In contrast to peptide
4 recognition by $\alpha\beta$ T cells, antigens activating $\gamma\delta$ T cells are mostly MHC independent and not
5 well characterized. Therefore, the function of $\gamma\delta$ T cells and their contribution to protection
6 against infections is still unclear. Higher numbers of circulating $\gamma\delta$ T cells compared to mice,
7 render the pig a suitable animal model to study $\gamma\delta$ T cells. Knocking-out the porcine TRDC-
8 locus by intracytoplasmic microinjection and somatic cell nuclear transfer resulted in healthy
9 living $\gamma\delta$ T cell deficient offspring. Flow cytometric analysis revealed that TRDC-KO pigs lack
10 $\gamma\delta$ T cells in peripheral blood mononuclear cells (PBMC) and spleen cells. The composition
11 of the remaining leucocyte subpopulations was not affected by the depletion of $\gamma\delta$ T cells.
12 Genome-wide transcriptome analyses in PBMC revealed a pattern of changes reflecting the
13 impairment of known or expected $\gamma\delta$ T cell dependent pathways. Histopathology did not
14 reveal developmental abnormalities of secondary lymphoid tissues. However, in a
15 vaccination experiment the KO pigs stayed healthy but had a significantly lower neutralizing
16 antibody titer as the syngenic controls.

17

18

19 Introduction

20 The adaptive immune system is composed of three lymphocyte subsets, B cells, T cells
 21 expressing the $\alpha\beta$ T cell receptor (TCR) and T cells expressing $\gamma\delta$ TCRs. The tripartite
 22 organization of the lymphocytic immune system seems to be fundamental since it evolved
 23 independently in jawed and jawless vertebrates¹. These cells generate their antigen specific
 24 receptors using somatic V(D)J recombination, which enables them to recognize a vast
 25 spectrum of different antigens. While the nature of antigen recognition by B cells and $\alpha\beta$ T
 26 cells is well established, a unifying concept for $\gamma\delta$ T cell antigen recognition and function is
 27 still pending. The majority of $\gamma\delta$ T cells are activated in an MHC-independent manner, which
 28 is in striking contrast to MHC-restricted $\alpha\beta$ T cells. They can attack target cells directly
 29 through their cytotoxic activity or indirectly through activation of other immune cells². Until
 30 recently, $\gamma\delta$ T cells were thought to be simply innate immune cells with limited or redundant
 31 functions³. The current view is that these cells complement many different players of the
 32 immune system⁴, and it is becoming obvious that they represent a heterogenous population
 33 of cells with important unique features in many infections, autoimmune diseases, allergies
 34 and in immunoregulation. A wide range of $\gamma\delta$ T cell functions have been described in humans
 35 and mice, including skin and mucosal epithelial wound repair, induction of tolerance,
 36 cytotoxicity and the production of various cytokines that regulate immune responses⁵⁻⁸. The
 37 increase in $\gamma\delta$ T cells following vaccinia virus infection, high vaccinia virus replication in $\gamma\delta$ T
 38 cell knockout mice, and the selective lysis of $\gamma\delta$ T cells against vaccinia virus infected target
 39 cells indicate some role for $\gamma\delta$ T cells during virus infections⁹⁻¹¹. To understand what they do
 40 and what role they play in different challenging situations of the immune system, a knockout
 41 animal model would be indispensable. A $\gamma\delta$ T cell deficient mouse model was successfully
 42 established¹², however $\gamma\delta$ T cells show a remarkable degree of diversification in function,
 43 anatomical localization and TCR usage that also differs largely between species¹³.
 44 Therefore, care has to be taken when extrapolating results from one species to the other. In
 45 humans and mice $\gamma\delta$ T cells constitute only 0.5–10% of T cells in peripheral blood but are
 46 substantially enriched in epithelial tissues (e.g. in skin, lungs, intestine). A prominent

population of $\gamma\delta$ T cells in human blood, V γ 9V δ 2 T cells which recognize phosphoantigens (PAgs) do not exist in mice but were recently described in the new world camelid *Vicugna pacos* (alpaca)¹⁴. Furthermore, while in human and mice a limited number of $\gamma\delta$ VDJ cassettes exist, species like cattle, pigs and chicken have larger numbers of $\gamma\delta$ VDJ cassettes and a high percentage of circulating $\gamma\delta$ T cells (20 - 50 %), thus called $\gamma\delta$ T cell-high species in contrast to $\gamma\delta$ T cell-low species (humans and mice)¹⁵. In addition, in some species the co-receptor WC1 family, which belong to the group B scavenger receptor cysteine-rich molecules, is expanded¹⁶. WC1 receptors are exclusively expressed by $\gamma\delta$ T cells and function as hybrid pattern recognition receptors and $\gamma\delta$ TCR co-receptors in cattle¹⁷. Bovine WC1 receptors can directly bind to pathogenic bacteria and transduce activation signals to $\gamma\delta$ T cells, thus, $\gamma\delta$ T cells may be crucial for the control of certain systemic infections¹⁷. Based on the remarkable lack of conservation between $\gamma\delta$ T cells of different species, additional animal models are needed to understand the function of $\gamma\delta$ T cells and to establish a unifying concept for their role during immune responses. In this report we describe the generation of TRDC-KO pigs and their genetically identical wild type controls. Since the number and quality of inbred pig strains are limited, the identity at the MHC locus of the KO and control animals is of utmost importance for the comparison of the immune response to infections. We did not observe any abnormalities in health and behavior of the KO pigs under standard housing conditions. In addition, the immune system in KO pigs developed normally, with the exception of the absence of $\gamma\delta$ T cells. The availability of a $\gamma\delta$ T cells deficient animal model from a $\gamma\delta$ T cell-high species may add important new insights in the role of this mysterious part of the adaptive immune system.

69

70 **Results**

71 **Intracytoplasmic microinjection of porcine *in vitro* derived zygotes with TRDC-** 72 **CRISPR/Cas and transfer to recipients**

73 After transfer of 35 microinjected embryos, both recipients (#695, #696) were determined
74 pregnant on day 25 after surgical embryo transfer. Sow 695 delivered 5 liveborn piglets on
75 day 112 of gestation and sow 696 gave birth to 9 liveborn piglets on day 114 of gestation
76 (Table 1). All piglets were healthy and showed no aberrations regarding birth weight
77 compared to age-matched counterparts or wild type littermates.

78 **Genotyping of TRDC-KO piglets**

79 Genomic DNA from tail tips was used for PCR-based detection of genetic modifications at
80 the TRDC locus in the piglets. The genomic DNA of the piglets originating from
81 intracytoplasmic microinjection of the pX330-TRDCEx4 #1 and #2 plasmids was employed
82 for PCR (Figure 1) and subsequent sequencing (Figure 2) to detect genetic modifications at
83 the TRDC gene. In total, two out of five piglets (40%) from sow 695 showed a biallelic 40 bp
84 deletion within exon 4 of the TRDC gene, while four out of nine piglets (44.4%) from sow 696
85 carried a biallelic 40 bp deletion. The remaining piglets were detected to be wild type. The
86 overall efficiency to generate a 40 bp biallelic deletion in the exon 4 of the TRDC gene was
87 42.9% (6/14).

88 **Transfection of CRISPR/Cas9 plasmids**

89 As an alternative to the intracytoplasmic microinjection approach, we employed somatic cell
90 nuclear transfer to produce genetically identical TRDC knockout pigs and syngeneic wild
91 type control pigs. Therefore, 3×10^6 cells in total were co-transfected with the plasmids
92 pX330-TRDCEx4#1 and #2 and subsequently seeded on three T25 flasks (25 cm²). After
93 reaching 70-80 % confluency, cells were detached by EDTA/Trypsin treatment and seeded
94 at a concentration of 7-10 cells per well on a 96-well plate. Individual cell clones were
95 analyzed by PCR. Cells with a biallelic deletion of the TRDC gene (- 40 bp) showed only the
96 lower band, while cell cultures with a mixed population of wild type, monoallelic and biallelic
97 genetically modified cells showed an upper wild type band at 499 bp and a lower band at 459
98 bp (Figure 3). Cell clones D12 (mixture) and H2 (almost pure lower band) were chosen as

99 donor cells for SCNT. D12 was chosen to produce both TRDC knockout and syngeneic wild
100 type control pigs.

101 **Somatic cell nuclear transfer of TRDC-KO embryos and syngeneic controls**

102 H2-derived cloned embryos were surgically transferred to two hormonally synchronized
103 recipients. Each of the two recipients (#8106 and #738) received 90 cloned embryos and
104 were detected to be pregnant on day 25 after embryo transfer. D12-derived cloned embryos
105 were also transferred to two recipients. One recipient (#8115) received 97 embryos, while the
106 second one received 96 embryos (#737). Again, both recipients were determined to be
107 pregnant by ultrasound scanning on day 25 after embryo transfer (Table 2). All recipients
108 were allowed to go to term and delivered in total 9 healthy liveborn (9/17:52.9%) and 8
109 stillborn piglets (8/17:47.1%). The stillborn piglets did not show any abnormalities but were
110 not checked for deletions in the exon 4 of the TRDC gene.

111 PCR analysis of the cloned piglets revealed that 4 out of the 7 piglets (57.1%) originating
112 from the mixed cell clone D12 (737-1-7) carried a biallelic 40 bp deletion of the TRDC gene,
113 while the remaining 3 piglets remained wild type (42.9%). The two piglets (738-1, -2)
114 originating from cell clone H2 carried a biallelic 40 bp deletion of the TRDC gene, as
115 expected (Figure 4).

116 **Fitness and viability of TRDC-knockout pigs**

117 The TRDC knockout pigs did not differ to syngenic wild type control animals and age-
118 matched wild type pigs in our facility in regard of their health status and growing performance
119 (Figure 5). TRDC-KO pigs are kept for 2 years in our facility under standard conditions and
120 never showed any health impairment.

121 **FACS analysis of TRDC-knockout blood and spleen cells**

122 To determine the effect of the 40 bp deletion in the TRDC gene on lymphocyte composition
123 in TRDC-KO pigs we performed flow cytometry analysis of PBMC and spleen cells from 8 wt
124 pigs and 5 TRDC-KO pigs. We used two mAb that specifically detect independent antigens
125 of porcine $\gamma\delta$ T cells. One binds to the δ -chain of the T cell receptor and the other to the CD3

antigen of $\gamma\delta$ T cells. In average 20 % (ranging from 8 % – 47 %) of peripheral blood T cells were double-stained by these mAbs in wild type pigs while no double-stained cells were detected in TRDC-KO pigs, indicating that no $\gamma\delta$ T cells were present in peripheral blood of TRDC-KO pigs (Figure 6A). In addition, we isolated spleen cells from wild type and TRDC-KO pigs and analyzed the content of $\gamma\delta$ T cell populations. Typically, in wild type pigs the majority (80 %) of $\gamma\delta$ T cells in the blood belong to the CD2 negative phenotype, while in the spleen the majority (66 %) of $\gamma\delta$ T cells are CD2 positive. No $\gamma\delta$ CD3 positive cells of either type were detected in the spleen of TRDC-KO pigs (Figure 6B). These data indicate that no $\gamma\delta$ T cells developed in TRCD-KO pigs. The majority of $\gamma\delta$ T cells in pigs are CD4 and CD8 negative, reflected by minimal numbers of CD4/CD8 double negative T cells in TRCD-KO pigs (Figure 6C). Comparing the numbers of various lymphocyte subpopulations between TRDC-KO pigs and wild type pigs we did not find significant differences (Figure 6D). These results indicated that the absence of $\gamma\delta$ T cells had no major impact on the development of other lymphocyte populations.

Comparison of gene expression profiles of TRDC-KO versus control animals via microarray analysis

Since we did not detect a major influence on the composition of lymphocyte subpopulations in the peripheral blood by the complete loss of $\gamma\delta$ T cells, we asked whether the absence of $\gamma\delta$ T cells has an impact on the transcriptional profile of the remaining lymphocyte population. Thus, we performed microarray analysis of PBMC isolated from wild type pigs and TRDC-KO pigs. Only 0.7 % of genes were regulated more than 2-fold (0.4 % upregulated and 0.3 % downregulated) in PBMC of TRDC-KO pigs (Figure 7A). Hierarchical clustering identified gene clusters with a significant expression level in either the wild type or the TRDC-KO pigs, only these genes were further analyzed (Figure 7B). A high portion of down-regulated genes are known to be preferentially expressed by $\gamma\delta$ T cells, such as δ T cell receptor genes, GATA3, WC1.1, SOX13, IL-1R and IL-18R. Genes which were upregulated in TRDC-KO pigs encompass genes which are preferentially expressed by granulocytes and monocytes in human PBMC like Zyxin, CXCL8 and CXCL2. Thus, the

transcriptional profile of PBMC in TRDC-KO pigs reflect the loss of $\gamma\delta$ T cells and as a result the proportional increase of non-lymphoid cells.

No differences between wild type and TRDC-KO pigs were detected by gross pathological, histological and immunohistological examination of lymphoid tissues

We wondered if the development of lymphoid tissues was influenced by the absence of $\gamma\delta$ T cells. At autopsy the size and morphology of all lymphoid tissues did not differ between wild-type or knockout pigs. Further, neither the detailed histopathologic evaluation on H&E stained slides nor the application of T-cell (CD3) and B-cell (CD20) markers revealed differences in the overall architecture, distribution and quantity of the immune cells labelled. Representative slides of the thymus, ileal Peyer's patches and tonsil are shown in Figure 8 (A-R). Both mAbs directed against the specific $\gamma\delta$ T cell-antigens (PGBL22A, PPT16) cross react on tissue slides with cytoplasmic structures of an unknown cell type, resulting in unspecific reactions. Thus both antibodies we had at hand, were unfortunately not suited for the use in immunohistology on cryosections.

Possible role of $\gamma\delta$ T cells in vaccine-mediated protection

As a first approach towards testing the immunological competence of the TRDC-KO pigs we decided to vaccinate the animals with a commercial vaccine against the pestivirus classical swine fever virus (CSFV). The vaccine contains a highly attenuated live virus of the so-called C-strain type. In normal pigs, the vaccine is known to be completely apathogenic but very effective with regard to induction of a protective immunity against CSFV field virus infection¹⁸⁻²¹. Three knockout pigs [group 1, numbers 1314, 1318, 1319] and two syngenic wild type controls [group 2, numbers 1315, 1320] were vaccinated at day 0 with the commercial live virus vaccine via the intramuscular route as recommended by the supplier. Rectal temperature of the animals was recorded daily starting 10 days before vaccination. The animals were monitored daily for general health status. Neither in group 1 nor 2 any signs of disease were detected during the observation period. Body temperatures of all animals remained in the normal range. Blood for serum production was taken on day 0 and on day 21

p.i. The sera were tested for the presence of neutralizing antibodies against the homologous CSFV C-strain. As expected, the results were negative for all animals at day 0, proving that they had not had any contact to CSFV antigens before. In contrast, neutralization titers were recorded on day 21 for all animals (Figure 9). The values were significantly higher for the wild type pigs compared to TRDC-KO (P value = 0.0428).

Discussion

Mice deficient for $\gamma\delta$ T cells have been established almost 30 years ago²². In the meantime, they have proven to be extraordinary helpful to elucidate functions of $\gamma\delta$ T cells. In particular, the role and contribution of certain murine $\gamma\delta$ T cell compartments during infections with viral, bacterial and parasitic pathogens could be defined using these mice²³⁻²⁵. Taken into account that $\gamma\delta$ T cell populations can differ largely between species it is of considerable importance to have $\gamma\delta$ T deficient animals from additional species. One major variation of $\gamma\delta$ T cell compartments between species is the amount of circulating $\gamma\delta$ T cells. The proportion of circulating $\gamma\delta$ T cells is surprisingly high in artiodactyls (e.g. cattle, sheep and pigs) and chicken¹⁵. Thus, the animal model described in the present report is the first of a $\gamma\delta$ T cell-high species. Pigs with a biallelic knockout of the TRDC gene were efficiently produced either by intracytoplasmic microinjection or somatic cell nuclear transfer using modified cells as donor cells. While healthy offspring could be obtained after embryo transfer of microinjected zygotes, somatic cell nuclear transfer (SCNT) resulted in 47.1 % stillborn piglets. As healthy TRDC knockout piglets were obtained by both techniques, we conclude that the relatively high number of stillborn piglets was associated with SCNT and false or incomplete reprogramming of the donor cell nucleus²⁶⁻²⁸. TRDC knockout pigs developed normally and reached sexual maturity. After breeding TRDC deficient pigs, all offspring carried a monoallelic knockout of the TRDC gene as expected. Though mosaicism cannot be excluded in the pigs originating from intracytoplasmic microinjection²⁹, we found the 40 bp deletion in all tissues analyzed (skin, heart, liver, kidney, muscle), rendering mosaicism unlikely. Besides, no off-target events could be detected in the 10 most possible genomic sites in microinjected and cloned offspring (data not shown). As previously observed in

Tcrd-/- mice no $\gamma\delta$ T cells remain in the TRDC-KO pigs while the studies suggest the presence of a normal $\alpha\beta$ T cell repertoire in the periphery of the $\gamma\delta$ T cell deficient pigs²². Furthermore, microarray analysis did not indicate that there is a major impact on the transcriptome of PBMC by the loss of $\gamma\delta$ T cells. These observations may explain why the immune phenotype of TRDC-KO pigs is surprisingly mild. The vaccination experiment described above was conducted with a commercial vaccine containing a live virus that is known to be completely apathogenic in wild type pigs¹⁸⁻²¹. We have chosen this approach to check whether the knockout pigs are hampered in a way that they cannot control a virus infection at all, even when the infecting virus is highly attenuated. In fact, we saw no signs of disease in the vaccinated pigs which could be explained in different ways, namely that the so far unknown attenuation principle of CSFV C-strain is sufficient even in immunologically compromised animals, that the loss of $\gamma\delta$ T cells has only mild effects on virus control by the immune system or a mixture of both reasons. The results of the antibody neutralization test showed that the knockout pigs were able to mount a measurable antibody response to an infecting virus, but the titers were lower than in the wild type animals. Even though the statistical significance of this result is only low due to the small number of animals tested, it represents a first indication for a reduced immunological competence of the knockout animals. This may indicate that different outcomes of infections are due to the missing contribution of $\gamma\delta$ T cells during the immune response not to other deficiencies of the immune system. This point will have to be followed up using different infection and challenge systems. There are various reports pointing to a protective role of $\gamma\delta$ T cells against important pig pathogens such as African swine fever virus^{30,31}, Foot and mouth disease virus^{32,33}, Mycobacterium bovis³⁴, Porcine respiratory and reproductive syndrome virus^{35,36}, and Influenza A virus³⁷. A specific role of $\gamma\delta$ T cells has not been definitively proven so far, but with the TRDC-KO pigs at hand, the large variety of swine pathogens displaying different ways of pathogenesis and interplay with the hosts immune system can be tested to finally elucidate the role of $\gamma\delta$ T cells in immunological defense against infectious agents. The fact

that some of these pathogens are zoonotic and responsible for severe human diseases
render this approach even more attractive.

Material and Methods

Animals

German Landrace pigs served as recipient animals for genetically modified embryos
derived by somatic cell nuclear transfer (SCNT).

CRISPR/Cas vector and single-guide RNA

The CRISPR/Cas9 system was used to induce defined deletions within the exon 4 of the
porcine TRDC gene (Ensembl transcript: ENSSSCT00000026772). Guide RNAs (gRNAs)
were designed using the web-based design tool *CRISPOR* (<http://crispor.tefor.net/>) (Fig. X).
Target sequences were analyzed via BLAST to reduce the probability for off-target events.
Two oligo duplexes including the target sequence (TRDCEX4#1: 5' -CAC CGT GAT GTC
TGT CAC AGT GCT T-3' and TRDCEX4#2: 5'-CAC CGG CAG TCA AGA GAA AAT TGA-
3') and BbsI overhangs were designed, and each gRNA was cloned into a linearized
CRISPR/Cas9 vector pX330-U6-Chimeric_BB-CBh-hSpCas9 (Addgene plasmid # 42230)³⁸.
The final plasmids pX330-TRDCEX4#1 and pX330-TRDCEX4#2 were then used for
intracytoplasmic microinjection into IVF zygotes and for transfecting porcine fetal fibroblasts.

Culture and lysis of primary cell cultures from wild type fetuses

Porcine fibroblasts were isolated from ear tissue of wild type piglets and cultured in
Dulbecco's modified Eagle's medium (DMEM) with 2 % penicillin/streptomycin, 1 % non-
essential amino acids and sodium pyruvate and 30 % fetal calf serum (FCS) (Gibco, 10270-
106). After the first passage the antibiotic concentration was reduced to 1%.

Transfection of CRISPR/Cas9 plasmids

In total, 3×10^6 cells were transfected when they reached 70-80% confluency. The two
CRISPR/Cas9 plasmids (pX330-TRDCEX4#1+2) were co-transfected (at a final
concentration of 5 µg/µl) into porcine fibroblasts by electroporation (Neon™ Transfection

System, ThermoFisher Scientific) to test the efficacy of the plasmids to induce a 40bp deletion at the targeted locus (Figure X). Electroporation conditions were as follows: 1350 V, 20 mm, and two pulses. After lysis of transfected cells in cell lysis buffer followed by ethanol extraction, the DNA was analyzed using TRDC specific primer (TRDCEx4-2F: 5'-CTGGGTGTAAGTAGCAGCCT-3' and TRDCEx4-2R: 5'-ACACGAGTTTTGAGTCTGGC-3'). The purified 499bp PCR product (10 ng/μl) (Invisorb® Fragment CleanUp – Startec) was Sanger sequenced to detect mutations at the target site. To produce TRDC-KO cell clones for serving as donor cells in SCNT, cells were trypsinized and subsequently diluted to a final concentration of 7-10 cells per well.

In-Vitro-Fertilization and In-Vitro-Maturation.

In-vitro-maturation of porcine oocytes was performed as previously described³⁹. Briefly, porcine oocytes were collected from ovaries derived from slaughterhouse and matured for 40 hours in FLI medium. For in vitro fertilization, frozen boar semen from a fertile landrace boar was thawed for 30 seconds in a water bath (37 °C). The sperm motility was microscopically checked (Olympus, BH-2). After washing with Androhep® Plus (Minitube) and centrifugation for 6 minutes at 600 g, approx. 75 to 100 sperm per oocyte (depending on semen capacity) were used for fertilization (no sexed sperm were utilized for fertilization). After four hours of co-incubation, the fertilized oocytes were cultured in porcine-zygote-medium (PZM-3 medium).

Intracytoplasmic Microinjection

Plasmids were prepared in 10 mM Tris-HCl pH 7.6 and 0.25 mM EDTA pH 8.0, and backfilled in glass injection capillaries and diluted to a final concentration of 2.5 ng/μl. Individual zygotes were fixed by suction to a holding pipette. The plasmids pX330-TRDCEX4#1 and pX330-TRDCEX4#2 were intracytoplasmically co-injected into IVF-produced zygotes derived from oocytes collected from slaughterhouse ovaries 20 hours after fertilization. To this end, approx. 10 pl plasmid solution was injected with a pressure of 600 hPa into IVF-produced zygotes (FemtoJet, Eppendorf). The injected zygotes were cultured in

PZM-3 medium at 39 °C, 5 % CO₂ and 5 % O₂. At day 5, when embryos had reached the morula/blastocyst stage, 35 embryos were surgically transferred into each of the two recipients.

Somatic cell nuclear transfer

SCNT was performed as previously described⁴⁰. Fetal fibroblasts transfected with px330-TRDCEx4#1 and px330-TRDC-Ex4#2 targeting the exon 4 of the porcine TRDC gene were used as donor cells. In order to produce syngeneic wild type control pigs simultaneously, we chose cell clone D12 that gave two bands with a 50:50 ration on the PCR gel (Fig. 3) and cell clone H2 that gave an almost clean -40bp PCR band. In total, 90 H2-derived one- to two-cell embryos were surgically transferred into each of two hormonally synchronized German Landrace gilts (7 to 9-months old). For D12-derived embryos, 97 and 96 one- to two-cell embryos were surgically transferred into two hormonally synchronized German Landrace gilts mg/day/gilt Altrenogest (Regumate® 4mg/ml, MSD Germany) for 12 days, followed by an injection of 1,000 IU PMSG (pregnant mare serum gonadotropin, Pregmagon®, IDT Biologika) on day 13 and induction of ovulation by intramuscular injection of 500 IU hCG (human chorion gonadotropin, Ovogest®300, MSD Germany) 72 h after PMSG administration. The surgical embryo transfer was performed the day after the hCG administration.

Genotyping of transfected cells and TRDC-KO pigs

Genomic DNA of the pigs was extracted from tail tips. Transfected cells and tail tips were lysed in cell lysis buffer (10%SDS, Proteinase K (20mg/ml), 10xPCR Buffer, aqua dest.) and purified by ethanol extraction. The DNA concentration was determined using the NanoDrop™ (Kikser-Biotech) system. For genotyping the pigs, polymerase chain reaction (PCR) was employed using specific primers (TRDCEx4-2F: 5'-CTGGGTGTAAGTAGCAGCCT-3' and TRDCEx4-2R: 5'-ACACGAGTTTTGAGTCTGGC-3') flanking a 499 bp segment of the TRDC gene (Fig. X). PCR amplification was performed in a total volume of 50 µl: 20 ng DNA, 0.6 µM reverse and forward primer, 1.5 mM MgCl₂, 0.2 mM dNTPs and 1.25 U *Taq* Polymerase.

Cycling conditions were as follows: 32 cycles with denaturation at 94°C for 30 sec, annealing at 60 °C for 45 sec, extension at 72°C for 30 sec and a final extension at 72°C for 5 minutes. The standard conditions for gel electrophoresis were set up to 80 V, 400 mA and 60 min using a 1 % agarose gel. The PCR-product was purified (Invisorb®Fragment CleanUp-Kit, Startec) and Sanger sequenced.

Allele-specific sequencing of the TRDC gene of cloned piglets

The PCR product of the targeted region within exon 4 of the TRDC gene was subcloned into the pGEM-T Easy Vector system (Promega, Germany) in accordance with the manufacturer's protocol and then transformed into XL10 bacteria. Bacteria were plated on Ampicillin containing Agar dishes (100 µg/ml) and allowed to grow for 16 hours. For each piglet, six colonies were picked and sequenced by using the T7prom primer.

FACS analysis of TRDC-knockout blood and spleen cells

PBMC were isolated by density gradient centrifugation using Lympholyte®-Mammal Cell Separation Media (Cedarlane, Canada). Spleen cells were isolated by generation of a single cell suspension and erythrocyte lysis with ammonium chloride. For flow cytometry 1×10^5 cells were incubated with primary antibodies followed by an incubation with secondary antibodies. The following primary antibodies were used anti-TCRγδ (PGBL22A, IgG1; PPT16, IgG2b), anti-CD2 (MSA4, IgG2a), anti-CD3 (PPT3, IgG1), anti-CD4 (74-12-4, IgG2b) and anti-CD8 (11/295/33, IgG2a). Detection of binding primary antibodies was performed using anti-mouse IgG1-PE, anti-mouse IgG2a-FITC, anti-mouse IgG2b-APC. All incubations were done in acid buffer at 4°C and dead cells were excluded by adding propidium iodide (PI) (dilution 1:1000 in acid buffer) to the cells before measurement. Flow cytometry was performed with the MACSQuant Analyser and the "MACS Quantify" software.

Transcriptome analysis

RNA from isolated PBMC was extracted using TRIzol® Reagent (Thermo Fisher Scientific). RNA from 4 TRDC-knockout pigs and 2 wild type pig were pooled, respectively. Pools were

used to assess the quality of the RNA using an agilent bioanalyser. Labeled fragmented single-stranded cDNAs (ss-cDNA) were synthesized by using purified total RNA (100–500 ng) as template following Affymetrix WT PLUS Labeling Assay protocols. Porcine Gene 1.1 ST Arrays (Affymetrix, Santa Clara, CA, USA) were hybridized to the biotinylated ss-cDNA targets. After 20 h of hybridization at 48 °C, arrays were washed by a fluidics station and then scanned by an imaging station in a GeneAtlas System (Affymetrix, Santa Clara, CA, USA). After scanning, the intensity data (CEL files) of Porcine Gene 1.1 ST arrays (Affymetrix) were extracted from the image data (DAT files) by the Affymetrix Command Console Software Version 1.4, and then normalized and analyzed by the Affymetrix Transcriptome Analysis Console (TAC) Software 4.0 for gene expression profiles and DEGs. The DEGs were selected by a cutoff of fold change >2.

Autopsy, Histology and immune fluorescence

Full autopsy was performed on all pigs. Samples of the thymus, tonsil, spleen, femoral bone marrow, ileal Peyer's patches, and lymph nodes (mandibular, tracheobronchial, cecal, popliteal) were immersion-fixed in neutral buffered, 10% formalin for at least 72 hours, and additional samples were snap frozen in liquid nitrogen. Formalin-fixed tissues were trimmed, embedded in paraffin, cut in 2 µm sections and stained with hematoxylin and eosin (H&E) according to standardized procedures. For immunohistochemistry (IHC), consecutive slides were mounted on adhesive glass slides, dewaxed in xylene, followed by rehydration in descending graded alcohols. Endogenous peroxidase was quenched with 3% H₂O₂ in distilled water for 10 minutes at room temperature (RT). Heat induced antigen retrieval was performed in a decloaking chamber for 10 minutes at 110°C in 10mM citrate buffer (pH 6). Nonspecific antibody binding was blocked with 1:2 diluted goat normal serum in Tris-buffered saline (TBS) for 30 minutes at RT. A polyclonal rabbit anti-CD3 (#A0452, diluted 1:100 in TBS, Dako Agilent, Santa Clara, CA, USA) or rabbit anti-CD20 (#RB-9013-P1, diluted 1:200 in TBS, Thermo Fisher Scientific, Waltham, MA, USA) was applied over night at 4°C. A secondary biotinylated goat-anti rabbit antibody was used (#BA-1000, diluted 1:200 in TBS, Vector Laboratories, Burlingame, CA,

USA) for 30 minutes at RT. The red-brown antigen labelling was developed by application of avidin-biotin-peroxidase complex (ABC) solution (Vectastain ABC Kit, #PK 6100, Vector Laboratories), followed by exposure to 3-amino-9-ethylcarbazole substrate (AEC, Dako, Agilent, Santa Clara, CA, USA). Sections were counter-stained with Mayer's hematoxylin, dehydrated in ascending graded alcohols, cleared in xylene, and coverslipped. All slides were digitized using the Aperio CS2 slide scanner (Leica Biosystems Imaging Inc., CA, USA) and image files were generated using the NDP.view2 Software (Hamamatsu Photonics, Hamamatsu City, Japan). The establishment of immunofluorescence double-labelling using a polyclonal rabbit anti-CD3 and a monoclonal mouse anti-TCR $\gamma\delta$ /PGBL22A antibody on snap frozen lymphoid tissues failed (not details shown). All examinations were performed by a board-certified pathologist (DipIECVP).

Vaccination experiment

Five pigs at an age of 3 months were included in a vaccination study. Three of these animals had a $\gamma\delta$ knockout phenotype (numbers 1314, 1318, 1319 (738-1, 737-4, 737-5) = group 1] and two represented syngenic wild type controls [group 2, numbers 1315, 1320 (737-1, 737-6)]. The two groups were kept in separate rooms. Vaccination was done after 14 days of acclimatization with a commercial live CSFV vaccine (Pestiffa CL, Boehringer Ingelheim Vetmedica, Ingelheim, Germany) according to the providers' recommendation via the intramuscular route into the muscle brachiocephalicus. After vaccination, the animals were monitored daily. Rectal temperatures were recorded daily from -10 dpv to 24 dpv. Blood samples for serum production were taken on days 0 dpv and 21 dpv. Determination of neutralizing antibody levels was done as essentially as described before^{41,42}

Declarations

Ethics approval and consent to participate

Animal experiments were approved by the supervisory authorities (LAVES, AZ 33.19-42502-04-17/2532 and LALLF, 7221.3-2-042/17) and conducted in compliance with the German animal welfare law, the German guidelines for animal welfare and the EU Directive

2010/63/EU. All experiments were performed in accordance with relevant guidelines and regulations.

Consent for publication

Not applicable.

Availability of data and materials

Not applicable

Competing interests

The authors declare that they have no financial and non-financial competing interests.

Funding

This study was supported by DFG (HE 6249/4-1) to R.K. This funding source had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Authors' Contributions

B.P., R.K. and G.M. conceived and designed the study and carried out data analysis as well as writing and revising the manuscript. B.P., A.F., P.H., R.B. and A.L.-H. established the TRDC-KO pigs, T.H.D., A.B. and R.G.U. performed experiments and data analysis.

Acknowledgements

We thank Julia Sehl for the support with the autopsy and Gaby Stooß, Franziska Grenkowitz, Freja Pfirske, Gabriele Czerwinski and Maren Ziegler for excellent technical assistance.

Figures

Figure 1: PCR-Analysis of the porcine TRDC locus in samples from pigs generated by intracytoplasmic microinjection. Pigs 695-2, -3 and 696-1, -5, -6, -7 display a lower band of about 460 bp in contrast to wild type pigs (500 bp) indicating a 40 bp deletion caused application of the CRISPR/Cas9 construct.

Figure 2: Sequencing data of pigs generated by intracytoplasmic microinjection of the CRISPR/Cas9 construct targeting the porcine TRDC-locus. Sequencing results confirmed the PCR results (Fig.1) that pigs 695-2, -3 and pigs 696-1, -5, -6, -7 carry a homozygous knockout of the TRDC locus consisting of a clear 40 bp deletion. For each piglet, six samples were sequenced. None of the piglets were monoallelic knockouts. (The binding sequences of the guide RNAs are indicated by green arrows).

Figure 3: PCR analysis of the TRDC locus of CRISPR/Cas9 transfected cells. The lower band of 460bp indicates a 40bp deletion caused by the CRISPR/Cas9 construct. Cell clone D12 displayed two bands of 500 and 460bp of size which indicates a mixed population of cells consisting of TRDC-knockout cells, wild type cells and/or cells with a monoallelic knockout of the TRDC locus. In contrast, cell clone H2 shows a single band of 460 bp indicating an almost pure population of TRDC knockout cells.

Figure 4: Sequencing results of the cloned liveborn TRDC knockout pigs. Pig 737-4 differed regarding the genetic modification compared to the other TRDC knockout pigs with a larger deletion encompassing 45bp, indicating its origin from a separate cell subclone. The binding sequences of the guide RNAs are indicated by red arrows.

Figure 5: Three TRDC-knockout pigs from intracytoplasmic microinjection.

Figure 6: No $\gamma\delta$ T cells are present in blood and spleen of TRDC-Knockout pigs. (A)

PBMC of TRDC-Knockout and wild type pigs were stained with anti-CD3, anti- δ chain and anti- $\gamma\delta$ T cell-specific CD3 mABs. CD3 positive cells were gated and δ chain vs. $\gamma\delta$ T cell-specific CD3 dotblots are displayed. About 20% of peripheral blood T cells were $\gamma\delta$ T cells in wild type pigs while no $\gamma\delta$ T cells were detected in TRDC-Knockout pigs. **(B)** PBMC and spleen cells were isolated and stained for CD2 and $\gamma\delta$ T cell-specific CD3. Typically in wild type pigs the majority (80%) of $\gamma\delta$ T cells in the blood belong to the CD2 negative phenotype (left dot blot), while in the spleen the majority (66%) of $\gamma\delta$ T cells are CD2 positive (middle dot blot). No $\gamma\delta$ T cells of either type were detected in the spleen of TRDC-Knockout pigs. **(C)** Most $\gamma\delta$ T cells are CD4 and CD8 negative. Accordingly TRDC-Knockout pigs have minimal numbers of CD4/CD8 double negative T cells. **(D)** The percentage of lymphocyte subpopulations of all lymphocytes without $\gamma\delta$ T cells were compared of wild type and TRDC-Knockout pigs. No significant difference of any lymphocyte subpopulation was detected.

Figure 7: Transcriptional changes in PBMC of TRDC-Knockout pig reflect the loss of $\gamma\delta$ T cells. (A) Only 0.7 % of genes were regulated more than 2 fold in PBMC of TRDC-Knockout pigs. (B) Using hierarchical clustering up-regulated and down-regulated genes were sorted according to the expression levels. (C) down-regulated (1,2,3) and upregulated (4) genes from the hierarchical clustering (B) were characterized in more detail. (1) all annotated genes of this cluster were expressed preferentially by T cells. GATA3 is highly expressed by $\gamma\delta$ T cells. (2) Cluster 2 contains genes like WC1.1 and SOX13 which are preferentially expressed by $\gamma\delta$ T cells to a various degree. (3) T cell receptor genes were found in cluster 3. In addition, IL-1R (3) and IL-18R (2) were downregulated in PBMC of TRDC-Knockout pigs. (4) Genes which were upregulated contain genes which were preferentially expressed by granulocytes and monocytes in human PBMC like Zyxin, CXCL8 and CXCL2.

Figure 8: No differences between wildtype and TRDC-Knockout pigs were detected by histological examination of lymphoid tissues. Lymphoid tissues from wild type (WT: A, C, D, G, I, J, M, O, P) and knockout (KO: B, E, F, H, K, L, N, Q, R) domestic pigs. Thymus (A, B) with cortex (black arrow) and medulla (green arrow), Peyer's Patches (G, H) with mucosa associated lymphoid follicle (black arrow) and absorptive epithelium (green arrow) containing intraepithelial, and transmigrating lymphocytes, Tonsil (M, N) with mucosa associated lymphoid follicle (black arrow) and mucosal epithelium (green arrow) containing intraepithelial and transmigrating lymphocytes. H&E stain (A, B, G, H, M, N), immunohistochemistry (C-F, I-L, O-R), using anti-CD3 or anti-CD20 antibody as indicated, ABC method, AEC chromogen

Figure 9: Diagram showing the titers of virus-neutralizing serum antibodies of 2 syngenic wt (wt) and 3 TRDC-Knockout (ko) pigs at day 21 post vaccination, given as the maximal reciprocal dilution of the sera that is able to neutralize 50% of 100 TCID₅₀ infective doses of CSFV C-strain calculated according to Spaerman-Kaerber. The mean values for both groups are shown as horizontal lines. The P value determined by statistical analysis is 0.0428.

References

- Hirano, M. *et al.* Evolutionary implications of a third lymphocyte lineage in lampreys. *Nature* **501**, 435-438, doi:10.1038/nature12467 (2013).

- 525 2 Lawand, M., Dechanet-Merville, J. & Dieu-Nosjean, M. C. Key Features of Gamma-Delta T-Cell
526 Subsets in Human Diseases and Their Immunotherapeutic Implications. *Frontiers in*
527 *immunology* **8**, 761, doi:10.3389/fimmu.2017.00761 (2017).
- 528 3 Deroost, K. & Langhorne, J. Gamma/Delta T Cells and Their Role in Protection Against
529 Malaria. *Frontiers in immunology* **9**, 2973, doi:10.3389/fimmu.2018.02973 (2018).
- 530 4 Vantourout, P. & Hayday, A. Six-of-the-best: unique contributions of gammadelta T cells to
531 immunology. *Nature reviews. Immunology* **13**, 88-100, doi:10.1038/nri3384 (2013).
- 532 5 Hanninen, A. & Harrison, L. C. Mucosal tolerance to prevent type 1 diabetes: can the
533 outcome be improved in humans? *Rev Diabet Stud* **1**, 113-121, doi:10.1900/RDS.2004.1.113
534 (2004).
- 535 6 Jameson, J. *et al.* A role for skin gammadelta T cells in wound repair. *Science* **296**, 747-749,
536 doi:10.1126/science.1069639 (2002).
- 537 7 Born, W. *et al.* Immunoregulatory functions of gamma delta T cells. *Adv Immunol* **71**, 77-144
538 (1999).
- 539 8 Hayday, A. C. [gamma][delta] cells: a right time and a right place for a conserved third way of
540 protection. *Annual review of immunology* **18**, 975-1026,
541 doi:10.1146/annurev.immunol.18.1.975 (2000).
- 542 9 Bukowski, J. F., Morita, C. T. & Brenner, M. B. Recognition and destruction of virus-infected
543 cells by human gamma delta CTL. *J Immunol* **153**, 5133-5140 (1994).
- 544 10 Johnson, R. M. *et al.* A murine CD4-, CD8- T cell receptor-gamma delta T lymphocyte clone
545 specific for herpes simplex virus glycoprotein I. *J Immunol* **148**, 983-988 (1992).
- 546 11 Welsh, R. M. *et al.* Alpha beta and gamma delta T-cell networks and their roles in natural
547 resistance to viral infections. *Immunological reviews* **159**, 79-93, doi:10.1111/j.1600-
548 065x.1997.tb01008.x (1997).
- 549 12 Han, H. J. *et al.* Murine gammadelta T Cells Render B Cells Refractory to Commitment of IgA
550 Isotype Switching. *Immune Netw* **18**, e25, doi:10.4110/in.2018.18.e25 (2018).
- 551 13 Castro, C. D., Boughter, C. T., Broughton, A. E., Ramesh, A. & Adams, E. J. Diversity in
552 recognition and function of human gammadelta T cells. *Immunological reviews* **298**, 134-152,
553 doi:10.1111/imr.12930 (2020).
- 554 14 Fichtner, A. S. *et al.* Alpaca (*Vicugna pacos*), the first nonprimate species with a
555 phosphoantigen-reactive Vgamma9Vdelta2 T cell subset. *Proceedings of the National*
556 *Academy of Sciences of the United States of America* **117**, 6697-6707,
557 doi:10.1073/pnas.1909474117 (2020).
- 558 15 Holderness, J., Hedges, J. F., Ramstead, A. & Jutila, M. A. Comparative biology of gammadelta
559 T cell function in humans, mice, and domestic animals. *Annual review of animal biosciences*
560 **1**, 99-124, doi:10.1146/annurev-animal-031412-103639 (2013).
- 561 16 Herzig, C. T., Waters, R. W., Baldwin, C. L. & Telfer, J. C. Evolution of the CD163 family and its
562 relationship to the bovine gamma delta T cell co-receptor WC1. *BMC evolutionary biology* **10**,
563 181, doi:10.1186/1471-2148-10-181 (2010).
- 564 17 Hsu, H. *et al.* WC1 is a hybrid gammadelta TCR coreceptor and pattern recognition receptor
565 for pathogenic bacteria. *J Immunol* **194**, 2280-2288, doi:10.4049/jimmunol.1402021 (2015).
- 566 18 Dong, X. N. & Chen, Y. H. Marker vaccine strategies and candidate CSFV marker vaccines.
567 *Vaccine* **25**, 205-230, doi:10.1016/j.vaccine.2006.07.033 (2007).
- 568 19 Leifer, I. *et al.* Characterization of C-strain "Riems" TAV-epitope escape variants obtained
569 through selective antibody pressure in cell culture. *Veterinary research* **43**, 33,
570 doi:10.1186/1297-9716-43-33 (2012).
- 571 20 Lin, T. T. C. L., R.C.T. *An overall report on the development of a highly safe and potent*
572 *lapinized hog cholera virus strain for hog cholera control in Taiwan* 40-42 (National Science
573 Council, 1981).
- 574 21 van Oirschot, J. T. Vaccinology of classical swine fever: from lab to field. *Veterinary*
575 *microbiology* **96**, 367-384, doi:10.1016/j.vetmic.2003.09.008 (2003).

576 22 Itohara, S. *et al.* T cell receptor delta gene mutant mice: independent generation of alpha
577 beta T cells and programmed rearrangements of gamma delta TCR genes. *Cell* **72**, 337-348
578 (1993).

579 23 Sell, S. *et al.* Control of murine cytomegalovirus infection by gammadelta T cells. *PLoS*
580 *pathogens* **11**, e1004481, doi:10.1371/journal.ppat.1004481 (2015).

581 24 McKenna, K. C. *et al.* gammadelta T cells are a component of early immunity against
582 preerythrocytic malaria parasites. *Infection and immunity* **68**, 2224-2230,
583 doi:10.1128/iai.68.4.2224-2230.2000 (2000).

584 25 Omar, T., Ziltener, P., Chamberlain, E., Cheng, Z. & Johnston, B. Mice Lacking gammadelta T
585 Cells Exhibit Impaired Clearance of *Pseudomonas aeruginosa* Lung Infection and Excessive
586 Production of Inflammatory Cytokines. *Infection and immunity* **88**, doi:10.1128/IAI.00171-20
587 (2020).

588 26 Kurome, M. *et al.* Factors influencing the efficiency of generating genetically engineered pigs
589 by nuclear transfer: multi-factorial analysis of a large data set. *BMC Biotechnol* **13**, 43,
590 doi:10.1186/1472-6750-13-43 (2013).

591 27 Liu, Y. *et al.* Developmental Competence and Epigenetic Profile of Porcine Embryos Produced
592 by Two Different Cloning Methods. *Cell Reprogram* **19**, 171-179, doi:10.1089/cell.2016.0055
593 (2017).

594 28 Schmidt, M., Winther, K. D., Secher, J. O. & Callesen, H. Postmortem findings in cloned and
595 transgenic piglets dead before weaning. *Theriogenology* **84**, 1014-1023,
596 doi:10.1016/j.theriogenology.2015.05.037 (2015).

597 29 Tanihara, F. *et al.* Effects of concentration of CRISPR/Cas9 components on genetic mosaicism
598 in cytoplasmic microinjected porcine embryos. *J Reprod Dev* **65**, 209-214,
599 doi:10.1262/jrd.2018-116 (2019).

600 30 Huhr, J. *et al.* Impaired T-cell responses in domestic pigs and wild boar upon infection with a
601 highly virulent African swine fever virus strain. *Transboundary and emerging diseases* **67**,
602 3016-3032, doi:10.1111/tbed.13678 (2020).

603 31 Post, J., Weesendorp, E., Montoya, M. & Loeffen, W. L. Influence of Age and Dose of African
604 Swine Fever Virus Infections on Clinical Outcome and Blood Parameters in Pigs. *Viral*
605 *Immunol* **30**, 58-69, doi:10.1089/vim.2016.0121 (2017).

606 32 Takamatsu, H. H. *et al.* Porcine gammadelta T cells: possible roles on the innate and adaptive
607 immune responses following virus infection. *Veterinary immunology and immunopathology*
608 **112**, 49-61, doi:10.1016/j.vetimm.2006.03.011 (2006).

609 33 Toka, F. N. & Golde, W. T. Cell mediated innate responses of cattle and swine are diverse
610 during foot-and-mouth disease virus (FMDV) infection: a unique landscape of innate
611 immunity. *Immunology letters* **152**, 135-143, doi:10.1016/j.imlet.2013.05.007 (2013).

612 34 Lee, J., Choi, K., Olin, M. R., Cho, S. N. & Molitor, T. W. Gammadelta T cells in immunity
613 induced by *Mycobacterium bovis* bacillus Calmette-Guerin vaccination. *Infection and*
614 *immunity* **72**, 1504-1511, doi:10.1128/iai.72.3.1504-1511.2004 (2004).

615 35 Olin, M. R. *et al.* Gammadelta lymphocyte response to porcine reproductive and respiratory
616 syndrome virus. *Viral Immunol* **18**, 490-499, doi:10.1089/vim.2005.18.490 (2005).

617 36 Kick, A. R. *et al.* The T-Cell Response to Type 2 Porcine Reproductive and Respiratory
618 Syndrome Virus (PRRSV). *Viruses* **11**, doi:10.3390/v11090796 (2019).

619 37 Schwaiger, T. *et al.* Experimental H1N1pdm09 infection in pigs mimics human seasonal
620 influenza infections. *PloS one* **14**, e0222943, doi:10.1371/journal.pone.0222943 (2019).

621 38 Cong, L. *et al.* Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**, 819-
622 823, doi:10.1126/science.1231143 (2013).

623 39 Lucas-Hahn, A. *et al.* 122 A New Maturation Medium Improves Porcine Embryo Production *In*
624 *Vitro* %J Reproduction, Fertility and Development. **30**, 200-201,
625 doi:https://doi.org/10.1071/RDv30n1Ab122 (2018).

626 40 Holker, M. *et al.* Duration of in vitro maturation of recipient oocytes affects blastocyst
627 development of cloned porcine embryos. *Cloning Stem Cells* **7**, 35-44,
628 doi:10.1089/clo.2005.7.35 (2005).

- 629 41 Meyers, G., Saalmuller, A. & Buttner, M. Mutations abrogating the RNase activity in
630 glycoprotein E(rns) of the pestivirus classical swine fever virus lead to virus attenuation.
631 *Journal of virology* **73**, 10224-10235, doi:10.1128/JVI.73.12.10224-10235.1999 (1999).
632 42 von Freyburg, M., Ege, A., Saalmuller, A. & Meyers, G. Comparison of the effects of RNase-
633 negative and wild-type classical swine fever virus on peripheral blood cells of infected pigs. *J*
634 *Gen Virol* **85**, 1899-1908, doi:10.1099/vir.0.79988-0 (2004).

635

636

637

638 **Table 1: Results of the intracytoplasmic microinjection of CRISPR/Cas9 targeting exon**
639 **4 of the porcine TRDC locus in porcine zygotes.**

Recipient	Microinjected plasmids	Number of transferred embryos	Pregnant*	Born piglets	No. of TRDC-KO
#695	pX330-TRDCEX4#1, pX330TRDCEX4#2	35	+	5	2
#696	pX330-TRDCEX4#1, pX330TRDCEX4#2	35	+	9	4
Total: 2		70	2/2 (100%)	14 (eff. 20%)	6/14 (42.9%)

640 *Pregnancy was detected on day 25

641 **Table 2: Results from somatic cell nuclear transfer employing genetically modified**
642 **cells as donor cells.**

Recipient	Cell clone*	Number of transferred embryos	Pregnant	Born piglets (stillborn)	No. of TRDC-KO of liveborn
#8106	H2	90	+	2 (2)	n.d.
#738	H2	90	+	3 (1)	2/2
#8115	D12	97	+	3 (3)	n.d.
#737	D12	96	+	9 (2)	4/7
Total: 4		373 (avg. 93.3)	4/4 (100%)	17 (cloning efficiency 4.6%)	6/9 (66.7%)

643 * cell clone D12: mixed population of knockout and wild type cells, cell clone H2: pure
644 knockout cells.

645