Age-dependent appearance of SARS-CoV-2 entry cells in mouse chemosensory systems reflects COVID-19 anosmia and ageusia symptoms.

Julien Brechbühl¹, Dean Wood¹, Sofiane Bouteiller¹, Ana Catarina Lopes¹, Chantal Verdumo¹, Marie-Christine Broillet^{1*}.

¹Department of Biomedical Sciences, Faculty of Biology and Medicine, University of Lausanne, Bugnon 27, CH-1011 Lausanne, Switzerland.

*Correspondence to: Marie-Christine Broillet, Department of Biomedical Sciences, Faculty of Biology and Medicine, University of Lausanne, Bugnon 27, CH-1011 Lausanne, Switzerland.

Email: mbroille@unil.ch

ORCID to Marie-Christine Broillet: 0000-0002-0487-6638.

ORCID to Julien Brechbühl: 0000-0002-2335-3058.

Abstract

COVID-19 pandemic has given rise to a collective scientific effort to study its viral causing agent SARS-CoV-2. Research is focusing in particular on its infection mechanisms and on the associated-disease symptoms. Interestingly, this environmental pathogen directly affects the human chemosensory systems leading to anosmia and ageusia. Evidence for the presence of the cellular entry sites of the virus, the ACE2/TMPRSS2 proteins, has been reported in non-chemosensory cells in the rodent's nose and mouth, missing a direct correlation between the symptoms reported in patients and the observed direct viral infection in human sensory cells. Here, mapping the gene and protein expression of ACE2/TMPRSS2 in the mouse olfactory and gustatory cells, we precisely identified the virus target cells to be of basal and sensory origin and we revealed their age-dependent appearance. Our results not only clarify human viral-induced sensory symptoms but also propose new investigative perspectives based on ACE2-humanized mouse models.

Introduction

The Corona Virus Disease 2019 (COVID-19) has federated worldwide scientific efforts for understanding the viral epidemiological mechanisms of the coronavirus 2 (SARS-CoV-2) that causes this severe acute respiratory syndrome. In humans, the viral syndrome is characterized by an increased mortality rate in aged and/or comorbidity patients associated with the upper respiratory infection symptoms, such as severe respiratory distress¹⁻³. In addition to its major impact, COVID-19 is associated by its direct alteration of human olfaction and gustation, in absence of substantial nasal inflammation or coryzal signs, resulting to anosmia and ageusia in up to 77% of the patients⁴⁻⁷. While these sensory symptoms are well established and intensely affect everyday behaviors^{8,9}, the precise related mechanisms remain elusive¹⁰.

The target cells of the virus share a molecular signature: the concomitant cellular expression of the angiotensin-converting enzyme 2 (ACE2) and of its facilitating transmembrane serine protease 2 (TMPRSS2), which plays a crucial role for the interaction of viral spike proteins with the host cell¹¹⁻¹³. Paradoxically, these entry sites seem to be lacking in sensory cells¹⁴⁻¹⁸, while a direct SARS-CoV-2 contamination has been observed both in humans and rodents^{19,20}, requesting further investigations to explain the sensory-associated symptoms²¹⁻²⁴. Therefore, the characterization of the animal model is necessary prior to its use to understand the causality underling the viral-induced sensory symptoms.

The use of mice is indeed limited for epidemiological studies due to their absence of hands, which, with aerosols, are the foremost passages of inter-individual viral transmission²⁵, as well as their published lack of SARS-CoV-2 ACE2-spike protein affinity^{26,27}. Nevertheless, the ease of production of genetically-modified mice and their scientific availability, as well as their well-studied and specialized chemosensory

systems²⁸⁻³⁰, make them a valuable ally for the development of potential prophylactic and protective treatments related to these sensory symptoms.

Thus, we aimed here at characterizing the potential viral entry sites across mouse sensory systems. We found SARS-CoV-2 entry cells to be of different origins depending on the sensory systems. In summary, the virus could target cells involved in tissue regulation such as the supporting cells of the olfactory receptor neurons and the regenerative basal cells but also, specifically, the chemosensory cells of both gustatory and olfactory systems. We finally revealed that the emergence of viral entry sites in sensory and basal cells only occurs with age, which could explain both, the observed COVID-19 long-lasting effects and the age-dependent sensory-symptomatology in human.

Results

SARS-CoV-2 entry genes are differentially expressed in the mouse sensory systems. Focusing on the mouse, where chemosensing takes place in different sensory systems (Fig. 1a), we took advantage of our previous studies³¹⁻³⁵ to first assess their overall expression of the major SARS-CoV-2 entry sites. Interestingly, we found a differential expression of the *Ace2* and *Tmprss2* transcripts (Fig. 1b). *Ace2* is strongly expressed in a specific area of the main olfactory epithelium (MOE), the dorsal part (MOE_D) which is directly exposed to the environment and specialized in sensing volatile chemical cues^{14,15}. Moreover, we observed a previously unreported expression of *Ace2*, in the most rostral sensory subsystem, the Grueneberg ganglion (GG), mostly implicated in volatile danger cues detection³⁴⁻³⁶ and in the circumvallate taste papilla (CV) involved in water-soluble tastant perception²⁹. Interestingly, this specific pattern of *Ace2* expression seems to be correlated with the mode of viral dissemination (volatile suspensions of viral droplets²⁵), as only a limited expression of *Ace2* is found in the vomeronasal organ (VNO) and in the septal organ of Masera (SO), considered to be implicated in pheromonal and retronasal communications via indirect and only limited access to the environment³⁰. Concerning the associated facilitating cleavage protease, we found that the *Tmprss2* transcript was expressed in all olfactory subsystems and to a limited extent at the CV level (Fig. 1b). Thus, this apparent disparity of expression requires further investigations to identify the precise cells expressing these entry sites in the different mouse subsystems.

SARS-CoV-2 entry cells in the MOE_D are of non-neuronal and multipotent types.

Taking advantage of a genetically modified mouse model in which, the olfactory marker protein³⁷ (OMP) drives the expression of the green fluorescent protein (GFP) in all mature olfactory neurons^{38,39}, we first examined the MOE_D sensory epithelium (Fig. 2) which is continuously exposed to inhaled air (Fig. 2a). Remarkably, we found by immunohistochemical stainings that the ACE2 protein was not only expressed at the apical surface and in the Bowman's glands^{14-16,21} but also in the basal layer (Fig. 2b). Moreover, we found that, in addition to being irregularly localized in different regions of the neuroepithelium¹⁵ (Supplementary Fig. 1), TMPRSS2 was mostly co-expressed with ACE2 in these OMP-negative basal cells (Fig. 2c, d). Focusing on the identification of these ACE2-expressing cells (Fig. 3), we first confirmed that the cytokeratin 18 (CK18)-positive sustentacular cells^{16,21}, operating as supporting cells for the olfactory neurons, were indeed harboring, at their luminal surface, the observed ACE2 protein (Fig. 3a). We next established the multipotency characteristics of the ACE2-expressing basal cells, as they also expressed two proteins specific to their identity of stem cells, the perinuclear cytokeratin 5 (CK5; Fig. 3b) and the nuclear transcription factor sex determining region Y-box 2 (SOX2; Fig. 3c), markers of horizontal basal cells (HBCs) or of the pear-shaped globose basal cells (GBCs) respectively. These two populations of basal cells are both involved in the regeneration of the neuroepithelium by acting as short- and long-lasting reservoir cells^{14,40,41}. Taken together, we confirmed our initial RT-PCR results (Fig. 1b) by precisely profiling the ACE2 and TMPRSS2 expression in the MOE_D neuroepithelium. We found that the cell candidates for viral entry are ACE2-expressing cells of non-neuronal and multipotent origin.

SARS-CoV-2 entry cells in the CV are mature sensory cells. Wondering about the apparent limited level of *Tmprss2* transcript in the CV (Fig. 1b), we next decided to exploit our histological methodology (Figs. 2, 3) to precisely localize the potential SARS-CoV-2 entry sites in sensory cells of the CV taste buds (Fig. 4). For that, we first used G_α gustacin (GUST) as a marker of sensory cells in the tongue³², to identify taste buds in close contact with the oral cavity (Fig. 4a). Surprisingly, using a double immunostaining approach, we distinctly observed ACE2 expression, not only in apical keratinocytes^{17,42}, but also in sensory cells (Fig. 4b). Moreover, ACE2 is predominantly expressed in the microvilli of sensory cells, gathered in the so-called taste pore, which are directly exposed to substances entering the oral cavity²⁹, such as viral droplets (Fig. 4b). Furthermore, we observed a restricted localization of TMPRSS2 in the taste pore region, strikingly co-expressed with ACE2 (Fig. 4c, d) which might also explain the observed low-level signal for the *Tmprss2* transcript (Fig. 1b). We next confirmed the mature status of these ACE2-expressing sensory cells as they preferentially expressed the CK18 marker⁴³ (Fig. 5a). Moreover, and contrary to the MOE_D, we found only sporadic ACE2 expression in basal CK5-positive cells (Fig. 5b) and SOX2-positive cells (Fig. 5c, d). In summary, we identified here the co-expression of the main SARS-CoV-2 entry sites in mature taste sensory cells.

SARS-CoV-2 entry cells in the GG are sensory neurons. In the mouse, there is an ancestral sensory subsystem, the GG (Fig. 6), which has the particularity of displaying both olfactory and gustatory traits^{31,32,44}. Morphologically, GG neurons are covered with a keratinized epithelium permeable to volatile water-soluble molecules coming from the nasal cavity³⁵ (Fig. 6a). Molecularly, it expresses both main sensory markers, OMP and GUST³² and uses common sensory signals such as olfactory and taste

receptors³² to respond to both danger-associated odorants and tastants^{32,35,36}. Performing histological approaches, we first noticed that the apical keratinocyte cell layer was, as in the CV, positive for ACE2 expression (Fig. 6b). But more surprising, we next found ACE2 expression in the GG cells (Fig. 6b). Interestingly and contrary to the MOE_D, this expression was distinctly restricted to the sensory GG neurons and not found in the supporting S100 calcium-binding protein β (S100B)-expressing glial cells³⁵ (Supplementary Fig. 2a). Remarkably, we furthermore localized TMPRSS2 both in apical keratinocytes and in GG neurons indicating that the two main SARS-CoV-2 entry sites are indeed found in the GG sensory subsystem (Fig. 6c, d). Looking for a precise characterization of ACE2-expressing cells in the GG, we next discovered that OMP-positive GG neurons were also CK18-positive (Fig. 7a) giving the GG a double cellular affiliation (neuronal and fibroblastic) that also highlights the discovery of CK18 as a reliable marker of ACE2 expressing cells, the putative target cells of viral infection⁴⁵. We next observed that in the apical keratinocyte layer of the GG, ACE2-expressing cells are mostly mature as they rarely co-expressed CK5 and SOX2 (Supplementary Fig. 2b, c), a characteristic shared with taste tissue (Fig. 5). Interestingly, and as previously shown, no regeneration happened throughout the lifespan in the GG subsystem^{31,46} as basal cells are absent which was further confirmed here by a lack of SOX2 staining (Supplementary Fig. 2c). However, we observed in GG neurons, a punctiform-like staining for ACE2 and CK5 (Supplementary Fig. 2b) indicating a co-expression in a precise cytoskeleton structure, as for the taste buds (Fig. 4b). We found that these cytoskeleton regions were associated to GG primary cilia (Fig. 7b, c), an organelle implicated in chemosensing^{33,36}. Thanks to the particulate guanylyl cyclase G (pGCG; Fig. 7b), a marker of the axonemes of the GG-cilia³³, we observed that ACE2 was located in the so-called basal body structures (Fig. 7c), where it was also

co-expressed with the gamma-tubulin marker³⁵ (γ -TUB; Fig. 7c). In summary, we found that the GG sensory subsystem possesses both the viral target cells and the specific protein profiles of expression displayed by both the olfactory and taste systems. Moreover, we bring here evidences of the presence of viral entry sites in chemosensory neurons.

SARS-CoV-2 entry site expression in mouse sensory systems is age-dependent.

During the course of our experiments, we noticed interindividual variations of ACE2 expression not only in its intensity but also in its general arrangement (e.g., apical *versus* basal ACE2 expression in the MOE_D; (Fig. 2b, c). Moreover, this observation was not only done for the MOE_D (Figs. 2, 3), but also for the CV (Figs. 4, 5) as well as for the GG (Figs. 6, 7). We considered a potential age-dependent correlation⁴⁷⁻⁵⁰ and thus we challenged the expression of ACE2 through the different sensory subsystems using mice of chosen ages. (Fig. 8). We saw first an intense variation of ACE2 expression in the MOE_D (Fig. 8a). Indeed, in young mice, this expression was restricted to the apical region, and then gradually intensifies with age (Fig. 8a). At the basal level, it appears to be absent first and only shows up in adult mice (Fig. 8a). In the sensory cells of the CV, ACE2 appears to be absent in young mice and only emerges with age (Fig. 8b). Interestingly, ACE2 expression initiates in the microvilli, at around 5 months of age, and then gradually extents throughout the cell body with aging (Fig. 8b). Concerning the GG, we first noticed that its apical keratinocyte cell layer constitutively expressed ACE2, while its expression was distinctly age-dependent in the sensory neurons (Fig. 8c). Indeed, and as for the sensory cells of the CV (Fig. 8b), a subcellular expression was first observed which spread into the soma with time (Fig. 8c). Overall,

we demonstrated here a striking age-dependent effect on ACE2 expression across mouse sensory systems, correlating with an increase of viral entry sites.

Discussion

Our ability to perceive and interact with our environment is directly linked to our senses. From an evolutionary point of view, our sensory performance is often associated with the way we communicate with our surroundings. In humans, although olfaction and taste have lost importance in comparison with mice, they remain essential not only to our well-being but also to protect us from dangers such as intoxication¹⁰. Their disorders are correlated with our general and mental health and can be early indicators of central nervous system impairments^{51,52}. Our sensory systems have developed a variety of protective mechanisms to guard us against environmental attacks such as the presence of protective keratinous epithelia, the combined action of mucus and cilia of sustentacular cells, ensuring the drainage of environmental microorganisms via mucociliary fluid clearance⁵³ or even the presence of regenerative basal cells^{40,41}. SARS-CoV-2 seems to have found a mechanism to thwart our sensory defenses. Indeed, we have shown here that, in mice, these very protective cells express viral entry sites. Although the direct action by the virus on these cells remains to be demonstrated, the underlying inflammatory^{22,24,54} or cytopathic destruction mechanisms²³ would strongly impact the senses of smell and taste. It therefore seems obvious that their targeted impairment can directly contribute to long-term anosmia and ageusia^{7,55}. Moreover, we have demonstrated here that the sensory cells themselves also express the SARS-CoV-2 entry sites which could contribute to both the sensory symptoms observed on the short term^{14,15,19,21} and the ability of coronaviruses to directly infect human sensory cells²⁰.

From a clinical point of view, anosmia and ageusia have a low prevalence in infected children and increase with the age of the COVID-19 patient^{50,56}, which seems to be consistent with the protein expression of ACE2 that we have observed in the mice olfactory system. Indeed, after we first confirmed the age-dependent trend made of ACE2 expression in sustentacular cells⁴⁷, we focused on the dorsal part of the MOE and found that the basal cells of the neuronal epithelium also expressed ACE2 as a function of age. Interestingly, TMPRSS2 is also expressed in these SARS-CoV-2 target cells. However, further studies, using viral inoculation for example, are still necessary to link this specific protein expression with viral sensitivity. Moreover, we also observed this age-effect in the taste system. As an absence of expression of the viral entry sites is found in the sensory cells of young mice¹⁷, but an increased expression of ACE2 and TMPRSS2 appears in sensory cells when the mouse advances in age. Thus, SARS-CoV-2 could therefore also directly target these sensory cells in aged mice and, possibly, in aged humans⁴², altering them and consequently disabling the taste sensory ability leading to ageusia symptoms. Thus, the use of mice sensitized to the virus^{45,57-60} appears to be a promising strategy but the age of the mice has to be carefully considered and older mice should preferentially be used (from 9 months of age). Interestingly, based on our results, additional studies performed on the lower respiratory airways could show whether a similar increased ACE2 expression as a function of age occurs, not only in respiratory ciliated cells^{49,61} but also in basal cells which could therefore contribute to the differences in pulmonary infectiousness observed between young and older people⁶².

Mice possess multiple sensory systems, separated into specialized subsystems³⁰, that we have now precisely characterized for their expression of the viral entry proteins. Interestingly, our study identified the GG to express the SARS-CoV-2

entry sites on its sensory neurons. We also found that GG sensory neurons expressed CK18, confirming that the cellular expression of this protein is an excellent marker for the localization of putative viral entry cells⁴⁵. Moreover, this observation is remarkable and gives to the GG a double cellular affiliation, namely neuronal and fibroblastic by the expression of both OMP and CK18. In this period of race for collective immunity⁹, we desperately need animal models to study this disease. Our results not only confirm the unique primary origin of the GG⁴⁴ but, also make it useful as a model of sensory systems to study, in *ex-vivo* and *in-vivo* preparations³⁵, the mechanism of viral entry into sensory cells as well as the testing of potential protective treatments against viral infection, such as intranasal drugs and / or COVID-19 vaccine candidate delivery^{13,63,64}. Reinforcing this idea, it should be noted that another SARS-CoV-2 portal of entry, neuropilin-1^{65,66} is also expressed in the GG⁶⁷, thus increasing its infectious susceptibility.

In summary, we have precisely characterized the SARS-CoV-2 entry cells in the multiple sensory systems of the mouse. These cells are of different origins. They are either involved in tissue function and regeneration but also, specifically, in chemosensing. We have revealed that the expression of viral entry sites in sensory cells increases with age, in this experimental model, which might be an important factor for viral infectivity studies. Thus, we have suggested a direct correlation between human sensory-symptomatology and mice SARS-CoV-2-expressing entry cells providing a putative explanation for the observed anosmia and ageusia in COVID-19 patients.

Methods

Animals. C57BL/6 mice (*Mus musculus*; Janvier Labs) and heterozygote OMP-GFP mice³⁹, of both sexes, were used at the indicated ages. In this gene-targeted mouse strain, mature olfactory sensory neurons³⁷ expressed the histological reporter green fluorescent protein (GFP) under the control of the olfactory marker protein (OMP) promoter^{38,39}. Mice were grouped-housed between 21 to 23 °C under a 12 hrs light/dark cycle with ad libitum access to food and water. Mice were euthanized by CO₂ inhalation and the experimental procedures were in accordance with Swiss legislation and approved by the EXPANIM committee of the Lemanique Animal Facility Network and the veterinary authority of Canton de Vaud (SCAV).

Chemosensory epithelia isolation. Prior to tissue isolation processes, phosphate buffered saline (PBS; 138 mM NaCl, 2.7 mM KCl, 1.76 mM KH₂PO₄, and 10 mM Na₂HPO₄, pH 7.4), dissection tools and equipment were sterilized and RNase removing agent (RNaseZAP™; Sigma) treatments were applied. The chemosensory epithelia were localized under the fluorescent stereomicroscope (M165 FC; Leica) thanks to the GFP expression for the olfactory subsystems^{32,33}. The dorsal GFP-positive prolongation of the main olfactory epithelium (MOE_D), corresponding to the ethmoid turbinate 1E adherent to the nasal bone, was separated from the rest of the MOE for the analysis. The other olfactory subsystems, the vomeronasal organ (VNO), the septal organ of Masera (SO) and the Grueneberg ganglion (GG) were delicately separated from the neighboring GFP negative tissues. The circumvallate papilla (CV) was extracted from the surface of the tongue.

Gene expression profile by RT-PCR. RNA purification was completed according to the instructions of the manufactured kit (RNeasy® Plus Mini kit; Qiagen). Briefly, the chemosensory epithelia obtained from 5-10 adult mice (4-11 months with a comparable sex ratio) were isolated and pooled in buffer RLT Plus supplemented with β -mercaptoethanol. Homogenization was performed through high-speed tissue disruption (TissueLyser II; Qiagen) and genomic DNA was removed from lysate by using the gDNA Eliminator spin column. Centrifugation and loaded processes were performed onto RNeasy spin columns to finally elute the total RNA in 30 μ l of RNase-free water. Reverse transcription (RT) was then initiated with the cDNA synthesis kit (PrimeScript™ 1st strand cDNA Synthesis Kit; Takara) using the random hexamers option. Polymerase chain reaction (PCR) experiments were subsequently performed using 140 ng of cDNA and 800 nM of each specific primers for *Ace2* (forward 5'-CTACAGGCCCTTCAGCAAAG-3'; reverse 5'-TGCCCAGAGCCTAGAGTTGT-3'; product size of 204 bp), *Tmprss2* (forward 5'-ACTGACCTCCTCATGCTGCT-3'; reverse 5'-TGACAGATGTTGAGGCTTGC-3'; product size of 225 bp) and *Gapdh* (forward 5'-AACTTTGGCATTGTGGAAGG-3'; reverse 5'-ACACATTGGGGGTAGGAACA-3'; product size of 223 bp). Primers were designed using the Primer3plus resource³². Amplification was done with 0.5 units of DNA Polymerase (GoTaq® DNA Polymerase; Promega) using a thermocycler (Veriti™; Applied Biosystems) programmed at 95°C 30 sec, 55°C 30 sec and 68°C 45 sec for 30 cycles. Visualization of the amplification products was done with ethidium bromide on a 3% electrophoresis gel and sizes were assessed (Bench Top 100bp DNA Ladder; Promega). Negative controls were performed with H₂O or by omitting the reverse transcription phase.

Protein expression profile by immunohistochemistry. Floating immunohistochemistry procedures were performed to precisely localize protein expression in the different chemosensory epithelia^{31,32}. Accordingly, and prior to tissue isolation, mouse heads were chemically fixated in 4% paraformaldehyde (PAF 4%, pH 7.4). After 24 h of fixation, chemosensory epithelia were rinsed in PBS and included in low melting 4% agar. Serial coronal slices of 80-120 μ m were generated with a vibroslicer (VT1200S; Leica) and collected in ice-cold PSB. Slices were then selected under a stereomicroscope (M165 FC; Leica) based on their general morphology and / or GFP expression. Selected slices were then blocked for 3 h at room temperature in a PBS solution containing 5% normal donkey serum (NDS; Jackson ImmunoResearch) and 1% of non-ionic detergent (Triton[®] X-100; Fluka). A double and indirect immunostaining approach using specific primary antibodies was then used to simultaneously localize ACE2 expression with other marker proteins^{14,15}. The primary antibodies used were directed against ACE2 (Goat anti-ACE2; PA5-47488; Invitrogen; 1:40), TMPRSS2 (Rabbit anti-TMPRSS2; ab109131; Abcam; 1:200), CK18 (Rabbit anti-CK18; PA5-14263; Invitrogen; 1:50), CK5 (Rabbit anti-CK5; ab52635; Abcam; 1:160), SOX2 (Rabbit anti-SOX2; PA1-094; Invitrogen; 1:200), GUST (Rabbit anti-G_α gust; sc-395; Santa Cruz Biotechnology; 1:250), S100B (Rabbit anti-S100 β ; ab41548; Abcam; 1:500), pGCG (Rabbit anti-PGCG; PGCG-701AP; FabGennix; 1:250) and γ TUB (Rabbit anti-gamma Tubulin; ab179503; Abcam; 1:250). Detection of primary antibodies was performed using fluorochrome-conjugated secondary antibodies against goat (Alexa Fluor Plus 647-conjugated, donkey anti-Goat; A32849; Invitrogen; 1:200) and rabbit (Cy3-conjugated, donkey anti-Rabbit; 711-165-152; Jackson ImmunoResearch; 1:200). Slices were then rinsed in 1% NDS solution; nuclei were counterstained with DAPI thanks to the antifade mounting medium (Vectashield[®]; H-

1200; Vector Labs). Control experiments were performed by omitting primary antibodies and ran in parallel on C57BL/6 mice. Observations were made by LED-fluorescence microscopy (EVOS M5000; Invitrogen) and acquisitions performed with confocal microscopy (SP5; Leica). Maximum projections, reconstructions and colocalization analysis were made with computer assistance (IMARIS 6.3; Bitplane).

References

- 1 Guan, W. J. *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* **382**, 1708-1720, doi:10.1056/NEJMoa2002032 (2020).
- 2 Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).
- 3 Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506, doi:10.1016/S0140-6736(20)30183-5 (2020).
- 4 Wolfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**, 465-469, doi:10.1038/s41586-020-2196-x (2020).
- 5 Menni, C. *et al.* Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nat Med* **26**, 1037-1040, doi:10.1038/s41591-020-0916-2 (2020).
- 6 Spinato, G. *et al.* Alterations in Smell or Taste in Mildly Symptomatic Outpatients With SARS-CoV-2 Infection. *JAMA* **323**, 2089-2090, doi:10.1001/jama.2020.6771 (2020).
- 7 Hannum, M. E. *et al.* Objective Sensory Testing Methods Reveal a Higher Prevalence of Olfactory Loss in COVID-19-Positive Patients Compared to Subjective Methods: A Systematic Review and Meta-Analysis. *Chemical senses* **45**, 865-874, doi:10.1093/chemse/bjaa064 (2020).
- 8 Pierron, D. *et al.* Smell and taste changes are early indicators of the COVID-19 pandemic and political decision effectiveness. *Nat Commun* **11**, 5152, doi:10.1038/s41467-020-18963-y (2020).
- 9 Aschwanden, C. Five reasons why COVID herd immunity is probably impossible. *Nature* **591**, 520-522, doi:10.1038/d41586-021-00728-2 (2021).
- 10 Marshall, M. COVID's toll on smell and taste: what scientists do and don't know. *Nature* **589**, 342-343, doi:10.1038/d41586-021-00055-6 (2021).
- 11 Bertram, S. *et al.* Cleavage and activation of the severe acute respiratory syndrome coronavirus spike protein by human airway trypsin-like protease. *J Virol* **85**, 13363-13372, doi:10.1128/JVI.05300-11 (2011).
- 12 Bertram, S. *et al.* Influenza and SARS-coronavirus activating proteases TMPRSS2 and HAT are expressed at multiple sites in human respiratory and gastrointestinal tracts. *PloS one* **7**, e35876, doi:10.1371/journal.pone.0035876 (2012).
- 13 Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278, doi:10.1016/j.cell.2020.02.052 (2020).
- 14 Brann, D. H. *et al.* Non-neuronal expression of SARS-CoV-2 entry genes in the olfactory system suggests mechanisms underlying COVID-19-associated anosmia. *Science advances* **6**, doi:10.1126/sciadv.abc5801 (2020).
- 15 Fodoulou, L. *et al.* SARS-CoV-2 Receptors and Entry Genes Are Expressed in the Human Olfactory Neuroepithelium and Brain. *iScience* **23**, 101839, doi:10.1016/j.isci.2020.101839 (2020).
- 16 Chen, M. *et al.* Elevated ACE-2 expression in the olfactory neuroepithelium: implications for anosmia and upper respiratory SARS-CoV-2 entry and replication. *Eur Respir J* **56**, doi:10.1183/13993003.01948-2020 (2020).
- 17 Wang, Z. *et al.* SARS-CoV-2 Receptor ACE2 Is Enriched in a Subpopulation of Mouse Tongue Epithelial Cells in Nongustatory Papillae but Not in Taste Buds or Embryonic Oral Epithelium. *ACS Pharmacol Transl Sci* **3**, 749-758, doi:10.1021/acsptsci.0c00062 (2020).
- 18 Klingenstein, M. *et al.* Evidence of SARS-CoV2 Entry Protein ACE2 in the Human Nose and Olfactory Bulb. *Cells, tissues, organs*, 1-10, doi:10.1159/000513040 (2021).
- 19 Sia, S. F. *et al.* Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature* **583**, 834-838, doi:10.1038/s41586-020-2342-5 (2020).
- 20 Meinhardt, J. *et al.* Olfactory transmucosal SARS-CoV-2 invasion as a port of central nervous system entry in individuals with COVID-19. *Nature neuroscience* **24**, 168-175, doi:10.1038/s41593-020-00758-5 (2021).
- 21 Bryche, B. *et al.* Massive transient damage of the olfactory epithelium associated with infection of sustentacular cells by SARS-CoV-2 in golden Syrian hamsters. *Brain Behav Immun* **89**, 579-586, doi:10.1016/j.bbi.2020.06.032 (2020).
- 22 Cazzolla, A. P. *et al.* Taste and Smell Disorders in COVID-19 Patients: Role of Interleukin-6. *ACS Chem Neurosci* **11**, 2774-2781, doi:10.1021/acschemneuro.0c00447 (2020).
- 23 Glezer, I., Bruni-Cardoso, A., Schechtman, D. & Malnic, B. Viral infection and smell loss: The case of COVID-19. *Journal of neurochemistry*, doi:10.1111/jnc.15197 (2020).

- 24 Merad, M. & Martin, J. C. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat Rev Immunol* **20**, 355-362, doi:10.1038/s41577-020-0331-4 (2020).
- 25 Lewis, D. COVID-19 rarely spreads through surfaces. So why are we still deep cleaning? *Nature* **590**, 26-28, doi:10.1038/d41586-021-00251-4 (2021).
- 26 Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* **581**, 215-220, doi:10.1038/s41586-020-2180-5 (2020).
- 27 Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature* **581**, 221-224, doi:10.1038/s41586-020-2179-y (2020).
- 28 Firestein, S. How the olfactory system makes sense of scents. *Nature* **413**, 211-218, doi:10.1038/35093026 (2001).
- 29 Mombaerts, P. Genes and ligands for odorant, vomeronasal and taste receptors. *Nature reviews. Neuroscience* **5**, 263-278, doi:10.1038/nrn1365 (2004).
- 30 Munger, S. D., Leinders-Zufall, T. & Zufall, F. Subsystem organization of the mammalian sense of smell. *Annual review of physiology* **71**, 115-140, doi:10.1146/annurev.physiol.70.113006.100608 (2009).
- 31 Brechbühl, J., de Valliere, A., Wood, D., Nenniger Tosato, M. & Broillet, M. C. The Grueneberg ganglion controls odor-driven food choices in mice under threat. *Commun Biol* **3**, 533, doi:10.1038/s42003-020-01257-w (2020).
- 32 Moine, F., Brechbühl, J., Nenniger Tosato, M., Beaumann, M. & Broillet, M. C. Alarm pheromone and kairomone detection via bitter taste receptors in the mouse Grueneberg ganglion. *BMC biology* **16**, 12, doi:10.1186/s12915-017-0479-y (2018).
- 33 Brechbühl, J. *et al.* Morphological and physiological species-dependent characteristics of the rodent Grueneberg ganglion. *Frontiers in neuroanatomy* **8**, doi:10.3389/fnana.2014.00087 (2014).
- 34 Brechbühl, J. *et al.* Mouse alarm pheromone shares structural similarity with predator scents. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 4762-4767, doi:10.1073/pnas.1214249110 (2013).
- 35 Brechbühl, J., Klaey, M. & Broillet, M. C. Grueneberg ganglion cells mediate alarm pheromone detection in mice. *Science* **321**, 1092-1095, doi:10.1126/science.1160770 (2008).
- 36 Chao, Y. C., Fleischer, J. & Yang, R. B. Guanylyl cyclase-G is an alarm pheromone receptor in mice. *The EMBO journal* **37**, 39-49, doi:10.15252/embj.201797155 (2018).
- 37 Margolis, F. L. A brain protein unique to the olfactory bulb. *Proceedings of the National Academy of Sciences of the United States of America* **69**, 1221-1224, doi:10.1073/pnas.69.5.1221 (1972).
- 38 Mombaerts, P. *et al.* Visualizing an olfactory sensory map. *Cell* **87**, 675-686, doi:10.1016/S0092-8674(00)81387-2 (1996).
- 39 Potter, S. M. *et al.* Structure and emergence of specific olfactory glomeruli in the mouse. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **21**, 9713-9723 (2001).
- 40 Fletcher, R. B. *et al.* Deconstructing Olfactory Stem Cell Trajectories at Single-Cell Resolution. *Cell Stem Cell* **20**, 817-830 e818, doi:10.1016/j.stem.2017.04.003 (2017).
- 41 Lin, B. *et al.* Injury Induces Endogenous Reprogramming and Dedifferentiation of Neuronal Progenitors to Multipotency. *Cell Stem Cell* **21**, 761-774 e765, doi:10.1016/j.stem.2017.09.008 (2017).
- 42 Sakaguchi, W. *et al.* Existence of SARS-CoV-2 Entry Molecules in the Oral Cavity. *Int J Mol Sci* **21**, doi:10.3390/ijms21176000 (2020).
- 43 Zhang, C. *et al.* Keratin 18 is associated with a subset of older taste cells in the rat. *Differentiation* **59**, 155-162, doi:10.1046/j.1432-0436.1995.5930155.x (1995).
- 44 Brechbühl, J., Moine, F. & Broillet, M. C. Mouse Grueneberg ganglion neurons share molecular and functional features with *C. elegans* amphid neurons. *Front Behav Neurosci* **7**, 193, doi:10.3389/fnbeh.2013.00193 (2013).
- 45 Winkler, E. S. *et al.* SARS-CoV-2 infection of human ACE2-transgenic mice causes severe lung inflammation and impaired function. *Nat Immunol* **21**, 1327-1335, doi:10.1038/s41590-020-0778-2 (2020).
- 46 Roppolo, D., Ribaud, V., Jungo, V. P., Luscher, C. & Rodriguez, I. Projection of the Grueneberg ganglion to the mouse olfactory bulb. *The European journal of neuroscience* **23**, 2887-2894, doi:10.1111/j.1460-9568.2006.04818.x (2006).
- 47 Bilinska, K., Jakubowska, P., Von Bartheld, C. S. & Butowt, R. Expression of the SARS-CoV-2 Entry Proteins, ACE2 and TMPRSS2, in Cells of the Olfactory Epithelium: Identification of Cell

- Types and Trends with Age. *ACS Chem Neurosci* **11**, 1555-1562, doi:10.1021/acscchemneuro.0c00210 (2020).
- 48 Lingappan, K., Karmouty-Quintana, H., Davies, J., Akkanti, B. & Harting, M. T. Understanding the age divide in COVID-19: why are children overwhelmingly spared? *Am J Physiol Lung Cell Mol Physiol* **319**, L39-L44, doi:10.1152/ajplung.00183.2020 (2020).
- 49 Yee, M., David Cohen, E., Haak, J., Dylag, A. M. & O'Reilly, M. A. Neonatal hyperoxia enhances age-dependent expression of SARS-CoV-2 receptors in mice. *Sci Rep* **10**, 22401, doi:10.1038/s41598-020-79595-2 (2020).
- 50 Steinman, J. B., Lum, F. M., Ho, P. P., Kaminski, N. & Steinman, L. Reduced development of COVID-19 in children reveals molecular checkpoints gating pathogenesis illuminating potential therapeutics. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 24620-24626, doi:10.1073/pnas.2012358117 (2020).
- 51 Riera, C. E. *et al.* The Sense of Smell Impacts Metabolic Health and Obesity. *Cell Metab* **26**, 198-211 e195, doi:10.1016/j.cmet.2017.06.015 (2017).
- 52 Rochet, M., El-Hage, W., Richa, S., Kazour, F. & Atanasova, B. Depression, Olfaction, and Quality of Life: A Mutual Relationship. *Brain Sci* **8**, doi:10.3390/brainsci8050080 (2018).
- 53 Gentzsch, M. & Rossier, B. C. A Pathophysiological Model for COVID-19: Critical Importance of Transepithelial Sodium Transport upon Airway Infection. *Function (Oxf)* **1**, zqaa024, doi:10.1093/function/zqaa024 (2020).
- 54 Muus, C. *et al.* Single-cell meta-analysis of SARS-CoV-2 entry genes across tissues and demographics. *Nat Med*, doi:10.1038/s41591-020-01227-z (2021).
- 55 Nguyen, N. N., Hoang, V. T., Lagier, J. C., Raoult, D. & Gautret, P. Long-term persistence of olfactory and gustatory disorders in COVID-19 patients. *Clin Microbiol Infect*, doi:10.1016/j.cmi.2020.12.021 (2021).
- 56 Somekh, I., Yakub Hanna, H., Heller, E., Bibi, H. & Somekh, E. Age-Dependent Sensory Impairment in COVID-19 Infection and its Correlation with ACE2 Expression. *Pediatr Infect Dis J* **39**, e270-e272, doi:10.1097/INF.0000000000002817 (2020).
- 57 Bao, L. *et al.* The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature* **583**, 830-833, doi:10.1038/s41586-020-2312-y (2020).
- 58 Dinno, K. H., 3rd *et al.* A mouse-adapted model of SARS-CoV-2 to test COVID-19 countermeasures. *Nature* **586**, 560-566, doi:10.1038/s41586-020-2708-8 (2020).
- 59 Munoz-Fontela, C. *et al.* Animal models for COVID-19. *Nature* **586**, 509-515, doi:10.1038/s41586-020-2787-6 (2020).
- 60 Sun, S. H. *et al.* A Mouse Model of SARS-CoV-2 Infection and Pathogenesis. *Cell Host Microbe* **28**, 124-133 e124, doi:10.1016/j.chom.2020.05.020 (2020).
- 61 Lee, I. T. *et al.* ACE2 localizes to the respiratory cilia and is not increased by ACE inhibitors or ARBs. *Nat Commun* **11**, 5453, doi:10.1038/s41467-020-19145-6 (2020).
- 62 Zimmermann, P. & Curtis, N. Why is COVID-19 less severe in children? A review of the proposed mechanisms underlying the age-related difference in severity of SARS-CoV-2 infections. *Arch Dis Child*, doi:10.1136/archdischild-2020-320338 (2020).
- 63 de Vries, R. D. *et al.* Intranasal fusion inhibitory lipopeptide prevents direct-contact SARS-CoV-2 transmission in ferrets. *Science*, doi:10.1126/science.abf4896 (2021).
- 64 Hassan, A. O. *et al.* A Single-Dose Intranasal ChAd Vaccine Protects Upper and Lower Respiratory Tracts against SARS-CoV-2. *Cell* **183**, 169-184 e113, doi:10.1016/j.cell.2020.08.026 (2020).
- 65 Daly, J. L. *et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection. *Science* **370**, 861-865, doi:10.1126/science.abd3072 (2020).
- 66 Cantuti-Castelvetri, L. *et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity. *Science* **370**, 856-860, doi:10.1126/science.abd2985 (2020).
- 67 Matsuo, T., Rossier, D. A., Kan, C. & Rodriguez, I. The wiring of Grueneberg ganglion axons is dependent on neuropilin 1. *Development* **139**, 2783-2791, doi:10.1242/dev.077008 (2012).

Acknowledgments

We thank A. de Vallière and E. Corset for technical support; B. Rossier for his thoughtful discussions on the manuscript. **Funding:** This work was supported by the Department of Biomedical Sciences, University of Lausanne, and by the Swiss National Science Foundation Grant 310030_185161 (to M.-C.B.).

Author contributions

J.B. and M.C.B designed the project. J.B., D.W., S.B., A.C.L. and C.V. carried out experimental procedures. All authors discussed the results and analyzed data. J.B. and M.C.B. wrote the manuscript.

Figures

Figure 1. Differential expression of the SARS-CoV-2 entry site transcripts in the mouse chemosensory systems. **a** Schematic representation of a mouse head with its chemosensory systems. Olfactory neurons are distributed in different olfactory subsystems: the main olfactory epithelium (MOE) and its dorsal region (MOE_D), the vomeronasal organ (VNO) and the septal organ of Masera (SO). Chemosensory cells for gustation are found in different tongue regions, grouped in taste buds which are presented by taste papilla such as the circumvallate papilla (CV). The Grueneberg ganglion (GG), the most rostral olfactory subsystems, displays both olfactory and gustatory properties. **b** Gene expression profiles of the SARS-CoV-2 entry sites: *Ace2* and *Tmprss2* in the different mouse chemosensory systems performed by RT-PCR. *Gapdh* is used as a reporter gene and H₂O as a negative control of transcript expression. Gene expression profiles obtained from 5-10 heterozygote OMP-GFP mice of 4 to 11 months old. Ladder of 100 base pairs (bp).

Figure 2. Expression profiles of the ACE2 and TMPRSS2 proteins in the MOE_D. Immunohistochemical investigations on the MOE_D for SARS-CoV-2 entry sites (ACE2, in pink; TMPRSS2, in red). Here, the olfactory marker protein (OMP, in green) allows the precise localization of the mature olfactory neurons. **a** Large view of OMP+ neurons of the MOE_D. Chemosensory cells are in direct contact with the nasal cavity. **b** Enlarged view of (**a**, dashed rectangle), showing expression of ACE2 in the apical surface, Bowman's gland and basal layer. A zoom in view of the apical surface is shown. **c** Co-expression profiles of TMPRSS2 and ACE2 in OMP- basal cells (white arrowhead, highlighted with zoom in view). **d** Control experiment (Ctrl neg) illustrating the absence of fluorescent background expression. (**a-d**) Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profiles obtained from heterozygote OMP-GFP mice of 20 (**a** and **b**) and 6 months old (**c** and **d**). Scale bars are 50 μm (**a**), 15 μm (**b**) and 10 μm (**c** and **d**).

Figure 3. Non-neuronal and immature olfactory cell localization of ACE2 in MOE_D. Cellular profile of ACE2 expression in the MOE_D by immunohistochemistry. **a** Co-expression of ACE2 (in pink) with apical CK18+ sustentacular cells (CK18, in red). **b, c** Basal co-expression of ACE2 with CK5+ HBCs (CK5, in red; **b**) and SOX2+ GBCs (SOX2, in red; **c**). Nuclei of sustentacular cells also expressed SOX2 (**c**). (**a-c**) Nuclei are counterstained with Dapi (DAPI, in blue). (**b** and **c**) Horizontal basal cells (HBCs), globose basal cells (GBCs). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 9 (**a** and **b**) and 7 months old (**c**). Scale bars are 5 μm (**a** and **b**) and 10 μm (**c**).

Figure 4. Expression profile of the ACE2 and TMPRSS2 proteins in the taste circumvallate papilla. Immunohistochemical investigations on the taste buds for SARS-CoV-2 entry sites. **a, b** Here, the gustducin marker protein (GUST; in red) allows the localization of the gustatory sensory cells. **(a)** Large view of Gust⁺ sensory cells of the circumvallate papilla (CV). **(b)** Enlarged view of **(a)**, dashed rectangle), showing expression of ACE2 (in pink) in the apical keratinocytes and in the microvilli of the chemosensory cells forming the taste pores, which are in direct contact with the oral cavity. A zoom in view of the microvilli of a taste pore is shown. **c** Co-expression profile of TMPRSS2 (in red) and ACE2 in the taste pores (highlighted with a zoom in view). **d** Control experiment (Ctrl neg) illustrating the absence of fluorescent background expression. **(a-d)** Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profiles were obtained from heterozygote OMP-GFP mice of 6 **(a and b)**, 5 **(c)** and 7 months old **(d)**. Scale bars are 50 μ m **(a)**, 10 μ m **(b-d)**.

Figure 5. Precise protein expression profile of ACE2 in CV taste buds reveals its presence in mature sensory cells. Cellular profile of ACE2 expression in the CV taste buds by immunohistochemistry. **a** Co-expression of ACE2 (in pink) with CK18+ sensory cells (CK18, in red). Enlarged view (dashed white circle) shows episodic expression of ACE2 in sensory cell soma. **b** Limited basal co-expression of ACE2 in immature CK5+ cells (CK5, in red). **c, d** Somatic expression of ACE2 is associated with nuclear expression of SOX2+ (SOX2, in red). **(d)** Enlarged view of a nuclear expression of SOX2 (indicated by white asterisks) in a ACE2+ sensory cell (white arrowhead in **(c)**). **(a-d)** Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 11 (**a** and **b**) and 13 months old (**c-d**). Scale bars are 10 μ m (**a-c**) and 5 μ m (**d**).

Figure 6. Expression profile of ACE2 and TMPRSS2 proteins in the GG. Immunohistochemical investigations on the GG for SARS-CoV-2 entry sites (ACE2, in pink; TMPRSS2, in red). Here, the olfactory marker protein (OMP, in green) allows the precise localization of the mature olfactory neurons. **a** General view of OMP+ neurons of the GG. Chemosensory cells are separated from the nasal cavity by a keratinocyte layer. **b** Enlarged view of (**a**, white dashed rectangle), showing expression of ACE2 in the apical keratinocytes and in GG neurons. A zoom in view of GG neurons is shown (white dashed circle). **c** Co-expression profile of TMPRSS2 and ACE2 in apical keratinocytes and OMP+ cells (zoom in view, white dashed circle). **d** Control experiment (Ctrl neg) illustrating the absence of fluorescent background expression. (**a-d**) Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 9 (**a**, **b** and **d**) and 11 months old (**c**). Scale bars are 50 μ m (**a**), 15 μ m (**b-d**).

Figure 7. Precise protein expression profile of ACE2 in the GG reveals its neuronal type. OMP+ cellular profile of ACE2 expression in the GG by immunohistochemistry. **a** Co-expression of ACE2 (in pink) with CK18+ GG neurons (CK18, in red). **b** ACE2 is expressed together with pGCG+ GG primary cilia (pGCG, in red). **c** Co-expression of ACE2 with γ TUB+ punctiform staining in GG neurons (γ TUB, in red) indicating a localization in the basal body of GG primary cilia. (**a-c**) Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 11 (**a**) and 5 months old (**b** and **c**). Scale bars are 10 μ m (**a-c**).

Figure 8. Age-dependent ACE2 expression in the MOE_D, CV and GG. Assessment of ACE2 expression (in pink) in chemosensory systems by immunohistochemistry at the indicated age in months (M). **a** A gradual increase in ACE2 expression is observed in the apical and basal cell layer of the MOE_D. **b** In taste buds of the CV, ACE2 is sequentially expressed in the taste pore microvilli and in the sensory somas. **c** In the GG region, apical keratinocytes constantly express ACE2 while this expression is progressively observed in GG neurons (punctiform to somatic staining). Colocalization (Coloc, in yellow) in sensory cells of ACE2 with OMP (in **a** and **c**) or GUST (in **b**) markers are highlighted. (**a-c**) Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 0.5, 5 and 20 months old are indicated. Scale bars are 10 μ m (**a** and **b**) and 20 μ m (**c**).

Supplementary Figures

Supplementary Figure 1. Irregular expression of TMPRSS2 in the MOE_D. TMPRSS2 (in red) is irregularly localized in different regions of the neuroepithelium. In addition to OMP- basal cell expression, sporadic cells express the TMPRSS2 protein. Here, TMPRSS2+ sustentacular cells co-stained with ACE2 (in pink) are shown. Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 7 months old. Scale bars are 10 μ m.

Supplementary Figure 2. ACE2 is expressed in GG neurons and not in supporting glial cells. a ACE2 (in pink) is expressed in OMP+ GG neurons (in green) and absent in their surrounding S100B+ glial cells (in red). White arrow head highlights a precise wrapping glial cell where a clear absence of ACE2 staining is observed. **b** Co-expression of ACE2 with CK5+ punctiform staining in GG neurons (CK5, in red). Absence of co-expression in the apical keratinocyte cell layer (zoom in view, white dashed circle). **c** Co-expression of ACE2 and SOX2 (SOX2, in red) in apical keratinocyte cells (white arrowhead). Absence of SOX2 expression in OMP+ GG neurons (zoom in view, white dashed circle). Nuclei are counterstained with Dapi (DAPI, in blue). Representative protein expression profile obtained from heterozygote OMP-GFP mice of 13 (a), 6 (b) and 7 months old (c). Scale bars are 10 μ m (a) and 15 μ m (b and c).