Calcium-independent astrocytic lipid release modulates neuronal excitability

Fushun Wang^{1,2}, Heather B. Bradshaw³, Salvador Pena⁴, Beata Jablonska⁵, Julia Xavier⁶, Sheng Gong⁷, Baoman Li¹, Devin Chandler-Militello¹, Lane K. Bekar⁸, and Nathan A. Smith^{5,9,10*}

¹Departments of Neurosurgery and Neurology, University of Rochester School of Medicine and Dentistry, Rochester, NY 14642, USA.

²Institute of Brain and Psychological Sciences, Sichuan Normal University, Chengdu, Sichuan 610060, China.

³Department of Psychological and Brain Sciences, Indiana University, Bloomington, IN 47405, USA

⁴Department of Pathology, University of Rochester School of Medicine and Dentistry, Rochester, NY 14642, USA

⁵Center for Neuroscience Research, Children's National Research Institute, Children's National Hospital, Washington DC 20010, USA.

⁶George Washington University, Washington DC 20052, USA.

⁷Department of Physiology and Biophysics, Case Western Reserve University School of Medicine, Cleveland, OH 44106, USA

⁸Department of Anatomy, Physiology, and Pharmacology, University of Saskatchewan, Saskatoon, SK S7N 5E5.

⁹George Washington University School of Medicine and Health Sciences, Washington DC 20052, USA.

¹⁰Lead Contact

*Correspondence should be addressed to Dr. Nathan Anthony Smith,
Email: nasmith2@childrensnational.org, Twitter: @SmithLab2018

ABSTRACT

Accumulating data point to a key role of Ca^{2+} -dependent gliotransmitter release as a modulator of neuronal networks. Here, we tested the hypothesis that astrocytes in response to agonist exposure also release lipid modulators through activation of Ca^{2+} -independent phospholipase A_2 (iPLA $_2$) activity. We found that cultured rat astrocytes treated with selective ATP and glutamatergic agonists released arachidonic acid (AA) and/or its derivatives, including the endogenous cannabinoid 2-arachidonoyl-sn-glycerol (2AG) and prostaglandin E_2 (PGE $_2$). Surprisingly, buffering of cytosolic Ca^{2+} resulted in a sharp increase in agonist-induced astrocytic lipid release. In addition, astrocytic release of PGE $_2$ enhanced miniature excitatory post-synaptic potentials (mEPSPs) by inhibiting the opening of neuronal Kv channels in brain slices. This study provides the first evidence for the existence of a Ca^{2+} -independent pathway regulating the release of PGE $_2$ from astrocytes, and furthermore demonstrates a functional role for astrocytic lipid release in the modulation of synaptic activity.

Significance Statement

Until now, the majority of studies implicating astrocytes in modulating synaptic activity have focused on Ca^{2+} -dependent release of traditional gliotransmitters such as D-serine, ATP, and glutamate. Mobilization of intracellular stores of Ca^{2+} occurs within a matter of seconds, but this novel Ca^{2+} -independent lipid pathway in astrocytes could potentially occur on a still faster time scale and thus participate in the rapid signaling processes involved in synaptic potentiation, attention, and neurovascular coupling.

INTRODUCTION

Comprising the major part of the dry weight of adult brain, lipids are an essential component of the phospholipid bilayer, mainly occurring as long-chain polyunsaturated fatty acids (PUFA) such as arachidonic acid (AA) and docosahexanoic acid (DHA) (Sinclair, 1975). The liver is the major site of AA synthesis, but the brain can produce AA and DHA *in situ* from their precursor fatty acids, linoleic and linolenic acids (Dhopeshwarkar and Subramanian, 1976). Astrocytes play a central role in the synthesis of AA and DHA in brain. Through their vascular endfeet, astrocytes have prime access to fatty acid precursors arriving across the blood-brain barrier (BBB), and serve as the major site for processing the essential fatty acids in the central nervous system (CNS) (Moore, 1993). Astrocytes also play a key role in macroscopic distribution of lipids in the brain parenchyma via perivascular glymphatic flux (Rangroo Thrane et al., 2013a; Plog and Nedergaard, 2018).

Lipids have gained much attention for their role as bioactive mediators in the CNS (Carta et al., 2014; Ledo et al., 2019). Numerous studies have focused on lipids in relation to functional hyperemia and synaptic activity. For instance, PGE₂ is a potent vasodilator and vasoconstrictor regulating CNS blood flow (Zonta et al., 2003; Takano et al., 2006; Gordon et al., 2007; Dabertrand et al., 2013; MacVicar and Newman, 2015; Czigler et al., 2019), and modulates the membrane excitability of CA1 pyramidal neurons during synaptic activity (Chen and Bazan, 2005). Furthermore, AA and its derivatives are important intracellular second messengers that can modulate the activities of various ion channels (Piomelli, 1993; Meves, 1994; Horimoto et al., 1997; Boland and Drzewiecki, 2008; Cordero-Morales and Vasquez, 2018). In addition, PGE₂ can suppress the outwardly rectifying Kv current in sensory neurons (Nicol et al., 1997; Evans et al., 1999), whereas AA suppresses Kv channels in the soma or dendrites of pyramidal neurons, and consequently broadens their presynaptic action potentials (Carta et al., 2014) and enhances EPSPs (Ramakers and Storm, 2002). However, those studies focused on the effects of neuronal lipid release effects on synaptic activity and paid scant attention to receptor

mediated pathways by which astrocytic lipids might influence synaptic activity. Given that the majority of AA, DHA, and other lipids present in the extracellular fluid is produced by astrocytes (Moore et al., 1991), receptor-mediated release could be a significant factor in the modulation of synaptic activity.

In culture, astrocytes can release AA in a cPLA₂ Ca²⁺-dependent pathway upon activation of metabotropic glutamate (mGluR) and P2Y purine (ATP) receptors (Bruner and Murphy, 1990; Stella et al., 1994; Stella et al., 1997; Chen and Chen, 1998). In addition, astrocytes also express the Ca²⁺-independent PLA₂ (iPLA₂) enzyme (Sun et al., 2005). Both of these isoforms are activated by the G-protein $\beta\gamma$ subunit (Jelsema and Axelrod, 1987; Murayama et al., 1990; van Tol-Steys et al., 1999), but iPLA₂ does not require Ca²⁺ or PKC phosphorylation for its activation (Winstead et al., 2000). Receptor-stimulated iPLA₂ activation can release AA and DHA in numerous cell types (Gross et al., 1993; Portilla et al., 1994; Akiba et al., 1998; Seegers et al., 2002; Tay and Melendez, 2004), but has not been fully explored in the case of astrocytes.

Previous work in our laboratory and elsewhere has demonstrated that astrocytes are capable of releasing gliotransmitters upon stimulation with mGluR or ATP receptor agonists, but the impact of this signaling on synaptic activity remains controversial. Recent studies with genetically encoded calcium indicators enabled us to identify localized Ca²⁺ signals within astrocytic fine processes and have confirmed Ca²⁺-dependent astrocytic effects on synaptic activity (Yu et al., 2018). However, the existence of Ca²⁺-independent lipid signaling in astrocytes is still not clearly established. Therefore, in this study we tested the hypothesis that astrocytes could support Ca²⁺-independent lipid signaling to modulate synaptic transmission. We used Ca²⁺ chelation to show that ATP and mGluR agonism can release lipids through iPLA₂ activation, resulting in the potentiation of synaptic activity. Because we currently lack the tools to assess Ca²⁺-independent signaling in astrocyte fine processes adjacent to synapses, we use Ca²⁺ inhibition on a grand scale to unmask this phenomenon (see discussion).

MATERIALS and METHODS

Culture

Cultured neocortical astrocytes were prepared from postnatal day 1 or 2 Wistar rat pups (Taconic Farms, Inc.) of either sex, as previously described (Lin et al., 1998). In brief, cerebral cortices were dissected on ice, and the meninges removed. The tissue was washed three times in Ca²⁺-free Hanks' balanced salt solution, and then triturated, filtered through a 70 µm nylon mesh, and centrifuged. The pellet was resuspended in 10 % fetal bovine serum in Dulbecco's modified Eagle's medium (DMEM)/F12 containing penicillin (100 IU ml⁻¹) and streptomycin (100 µg ml⁻¹), and transferred to culture flasks. Cells were maintained at 37 °C in an incubator containing humidified air and 5% CO₂. The medium was changed after 24 hrs and twice a week thereafter. More than 95% of the cells immunostained positive for GFAP. When the cells became confluent, they were rinsed two times in Ca²⁺-free Hanks' balanced salt solution, suspended in 0.05 % trypsin-containing PBS for 1 min, resuspended in DMEM/F12, centrifuged to remove the trypsin-containing supernatant, and then plated in 24-well plates. Experiments were performed when the cells were 95% confluent.

Viral Vectors and Viral Transductions

Viral vectors driving GFAP cyto-GCaMP6f (Baljit S. Khakh) were obtained from the University of Pennsylvania Vector Core (AAV 2/5 serotype). Secondary rat astrocytic cultures were transduced with AAV GFAP cyto-GCaMP6f. After transduction, the cultures were incubated at 37 °C for 5 days prior to Ca²⁺ imaging experiments.

Ca²⁺ imaging

Cultured cells in 24-well plates were transduced with AAV GFAP cyto-GCaMP6f and incubated with various pharmacological agents for 30 minutes at 37 °C. Using confocal microscopy (Olympus FV500), calcium wave activity was evoked by adding an equal volume of medium containing 100 µM of ATP to each well. Relative increases in fluorescence signal evoked by P2Y

receptor agonist exposure over baseline fluoresce ($\Delta F/F$) were calculated as previously described (Nedergaard, 1994; Smith et al., 2018).

Radiolabeling and Assessment of AA Release

Confluent rat astrocytic cultures were incubated with 100 nCi [5,6,8,9,11,12,14,15- ^3H]-AA (PerkinElmer) overnight before the experiments. The cells were washed three times with serum-free medium, and then allowed to recover for 20 minutes. Before stimulation of P2Y receptors with 100 μM ATP, the cells were incubated with appropriate inhibitors for 10 to 12 minutes. Aliquots of medium were taken 15 minutes after agonist stimulation, and ^3H -AA and/or its metabolites were measured by liquid scintillation counting.

HPLC/MS/MS analysis of lipids

Confluent rat astrocytes were washed with serum-free media and then allowed to recover for 1 hr in serum-free media. At this time, cells were incubated with vehicle or 20 μM cyclopiazonic acid (CPA) for 15 minutes and then challenged with 100 μM of ATP or vehicle for 15 minutes, as in the above [^3H]-AA experiments. Media was then removed and put aside, and 2 ml of HPLC-grade methanol was added to the flask for 5 minutes, and then removed and out aside. An additional 2 ml of HPLC-grade methanol was added, cells were scraped from the sides of the flask, and the contents added to the previous two retained fractions. Deuterium standards were added to a final concentration of 200 pM, and the samples were centrifuged at 19,000 x g and 24 °C. Lipids in the eluent were partially purified on 500 mg C18 Bond Elut solid phase extraction columns (Varian), concentrated into fractions of 40, 60, 85, and 100% methanol, then analyzed using HPLC/MS/MS (Shimadzu SCL10Avp-Wilmington, DE, USA; API 3000 Applied Biosystems/MDS SCIEX, Foster City, CA, USA) as previously described (Leishman et al., 2018). Over 50 lipids were targeted in these analyses, including 40 lipoamines, 3 acylglycerols, 2 free fatty acids, and 2 prostaglandins (namely X and Y). It was determined in pilot studies that

the analyte concentrations were too low for reliable detection in the medium alone, but the extraction of medium and cells together gave sufficient lipid concentrations for reliable assay.

PGE₂ Release Assessment via PGE₂ Immunoassay

Confluent rat astrocytic cultures were washed 3 times with serum-free medium, and then allowed to recover for 20 minutes. Before stimulation with 100 μ M ATP, the cells were incubated with appropriate P2Y inhibitors for 10 to 12 minutes. Aliquots of the medium were collected 15 minutes after stimulation and analyzed for PGE₂ content using an immunoassay kit (Cayman Chemicals) following the manufacturer's instructions.

Western Blot

Protein in samples harvested from the 24-well plates were separated by SDS-PAGE and transferred to a nitrocellulose membrane, which was then blocked with Tris-buffered saline containing 0.05% (wt/vol) Tween 20 and 5% nonfat dry milk. The primary antibodies were anti-iPLA2 (Sigma, St. Louis, MO), anti- β -actin (Cell Signaling, Danvers, MA), at 1:1000 to 1:2000 dilutions in blocking buffer. Detection of chemiluminescence from horseradish peroxidase-linked secondary antibodies was performed using the ChemiDoc™ XRS+ System and running Image Lab™ software.

Isolation of human fetal astrocytes

Human fetal forebrain tissues were obtained from second-trimester aborted fetuses of 20 weeks gestational age. Tissues were obtained from aborted fetuses, with informed consent and tissue donation approval from the mothers, under protocols approved by the Research Subjects Review Board of the University of Rochester Medical Center. No patient identifiers were made available to or known by the investigators; no samples with known karyotypic abnormalities were included. The forebrain tissue samples were collected and washed 2-3 times with sterile Hank's balanced salt solution containing $\text{Ca}^{2+}/\text{Mg}^{2+}$ (HBSS^{+/+}). The cortical plate region (CTX) of the fetal forebrain was dissected and separated from the ventricular zone/subventricular zone (VZ/SVZ) portion. The CTX was then dissociated with papain as previously described (Keyoung

et al., 2001). The cells were resuspended at a density of $2-4 \times 10^6$ cells/ml in DMEM/F12 supplemented with N2, 0.5 % FBS, and 10 ng/ml bFGF and plated in suspension culture dishes. The day after dissociation, cortical cells were recovered and subjected to magnetic activated cell sorting (MACS) to purify the astrocyte progenitor population. The recovered cells were briefly incubated with CD44 microbeads as per the manufacturer's recommendations (Miltenyi Biotech). The cells were then washed, resuspended in Miltenyi Washing buffer, and bound to a magnetic column (Miltenyi Biotech). The bound CD44+ astrocyte progenitor cells were eluted, collected, and then washed with DMEM/F12. The purified human fetal astrocyte progenitors were cultured in DMEM/F12 supplemented with N2, and 5% FBS to differentiate them further. To prepare culture dishes for PGE2 immunoassays or immunocytochemistry, the fetal cortical astrocytes were dissociated with TrypLE (Invitrogen) into single cells and then plated onto poly-L-ornithine/laminin-coated 24-well plates (50,000 cells per well).

shRNA Lentiviral knockdown of iPLA₂ in astrocytes

Rat astrocytes cultures were plated in a 24-well plate and grown to approximately 50% confluence. Following the manufacturer's instructions with minor modifications, the cultures were transduced overnight by adding either group VI iPLA₂ shRNA (r) lentiviral particles (sc-270117-V) or control shRNA lentiviral particles-A (sc-108080) directly to the culture medium containing polybrene (sc-134220) (all from Santa Cruz Biotechnology, Santa Cruz). At 24 hours after transfection, the culture medium was removed, and fresh culture medium without polybrene was added. Experiments and Western blot analysis were performed 7 days after transduction.

Acute Hippocampal Slice Preparation and Electrophysiology

Unless otherwise noted, 15-21 day old C57BL/6 (Charles River, Wilmington, MA), MrgA1^{+/-} transgenic, and littermate control MrgA1^{-/-} pups (courtesy of Dr. Ken McCarthy) (Fiacco et al., 2007) of either sex were used for the preparation of hippocampal slices as previously described (Wang et al., 2012). The pups were anesthetized in a closed chamber with isoflurane (1.5%)

and decapitated. The brains were rapidly removed and immersed in an ice-cold cutting solution that contained (in mM): 230 sucrose, 2.5 KCl, 0.5 CaCl₂, 10 MgCl₂, 26 NaHCO₃, 1.25 NaH₂PO₄, and 10 glucose, pH=7.2-7.4. Coronal slices (400 μm) were cut using a vibratome (Vibratome Company, St. Louis) and transferred to oxygenated artificial cerebrospinal fluid (aCSF) that contained (in mM): 126 NaCl, 4 KCl, 2 CaCl₂, 1 MgCl₂, 26 NaHCO₃, 1.25 NaH₂PO₄, and 10 glucose, pH = 7.2-7.4 (osmolarity = 310 mOsm). Slices were incubated in this aCSF for 1-5 hours at room temperature before electrophysiological recording. Experiments were performed at room temperature (21-23 °C). During the recordings, the slices were placed in a perfusion chamber and superfused with aCSF gassed with 5% CO₂ and 95% O₂ at room temperature. Cells were visualized with a 40X water-immersion objective and differential interference contrast (DIC) optics (BX51 upright microscope, Olympus Optical, New York, NY). Patch electrodes were fabricated from filament thin-wall glass (World Precision Instruments) on a vertical puller; the pipette's resistance was around 6-9 MΩ with addition of intracellular pipette solution. The pipette solution contained (in mM) 140 K-gluconate, 5 Na-phosphocreatine, 2 MgCl₂, 10 HEPES, 4 Mg-ATP, 0.3 Na-GTP (pH adjusted to 7.2 with KOH). The current-voltage relationship (I-V curve) of voltage-gated potassium currents was recorded under voltage-clamp using an AxoPatch MultiClamp 700B amplifier (Axon Instruments, Foster City, CA). When measuring outward currents, QX314 (0.5 mM) was added to the pipette solution to block Na⁺ currents. For recordings of miniature excitatory post-synaptic potentials (mEPSPs), 0.5 μM TTX was added to the aCSF. The junction potential between the patch pipette and bath solution was zeroed before forming a giga-seal. Patches with seal resistances less than 1 GΩ were rejected. Data were low-pass filtered at 2 kHz and digitized at 10 kHz with a Digidata 1440 interface controlled by pClamp Software (Molecular Devices, Union City, CA).

Pharmacological agents used in cultures and slice experiments

Adenosine 5 -triphosphate (ATP, 100 μM); Cyclopiazonic acid (CPA, 20 μM); *trans*- (1S, 3R)-1-Amino-1, 3-cyclopentanedicarboxylic acid (*t*-ACPD, 100 μM); (±)-α-Amino-3-hydroxy-5-

methylisoxazole-4-propionic acid hydrobromide ((±)-AMPA, 100 µM); Prostaglandin E₂ (PGE₂, 50 µM, Tocris); Phe-Met-Arg-Phe amide (FMRF, 15 µM); Thr-Phe-Leu-Leu-Arg-NH₂ (TFLLR-NH₂, 30 µM, Tocris); N-Acetyl-Asp-Glu (NAAG, 100 µM); (1*R*, 4*R*, 5*S*, 6*R*)-4-Amino-2-oxabicyclo [3.1.0] hexane-4,6-dicarboxylic acid disodium salt (LY379268, 100 µM, Tocris); calmidazolium chloride (CMZ, 2 µM, Tocris); methyl arachidonyl fluorophosphonate (MAFP, 10 µM); bromoenol lactone (Bel, 10 µM); AA (50 µM); *N*-(2,6-dimethylphenylcarbamoylmethyl) triethylammonium chloride (QX314, 1 mM, Tocris); 4-(4-,9-diethoxy-1,3-dihydro-1-oxo-2*H*-benz[*f*]isoindol-2-yl)-*N*-(phenylsulfonyl) benzeneacetamide (GW627368X 3 µM, Tocris); 6-Isopropoxy-9-xanthone-2-carboxylic acid (AH6809, 10 µM, Tocris); *N*-(piperidin-1-yl)-5-(4-iodophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide (AM251, 5 µM, Tocris); Tetrodotoxin (TTX, 0.5 µM, Tocris); 1,2-bis(2-aminophenoxy)ethane-*N,N,N,N*-tetraacetic acid (BAPTA, 50 µM, Tocris); and 1,2-Bis(2-aminophenoxy)ethane-*N,N,N,N*-tetra-acetic acid tetrakis (acetoxymethylester) (BAPTA-AM, 20 µM). All chemicals were from Sigma unless otherwise noted.

Statistical analysis of data

Statistical significance was evaluated by one-way ANOVA and *post hoc* Tests (Tukey and Dunn) using Prism software and deemed significant when $P < 0.05$ for the [³H]-AA and PGE₂ assay experiments. Normality of the data was evaluated by the Shapiro-Wilk test with $\alpha = 0.05$. For electrophysiology experiments, significance was determined by paired or unpaired t-tests or Tukey-Kramer *post hoc* multiple comparison tests. HPLC/MS/MS lipidomics data were analyzed with ANOVA and Fishers LSD *post-hoc* test using SPSS when $P < 0.05$ or $P < 0.10$. All results are reported as mean $\pm$ s.e.m.

RESULTS

Ca²⁺-independent release of [³H]-AA and its metabolites from cultured astrocytes

We first assessed the efficiency by which preloading with an inhibitor of the endoplasmic reticulum (ER) Ca²⁺ pump, cyclopiazonic acid (CPA) (20 µM), or the cytosolic Ca²⁺ chelator 1,2-

bis(o-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid (acetoxymethyl ester) (BAPTA-AM) (20 μ M) was able to block ATP (100 μ M) induced cytosolic Ca^{2+} increases in cultured rat astrocytes. Imaging cytosolic Ca^{2+} (AAV GFAP cyto-GCaMP6f) showed that the purine agonist ATP (100 μ M) induced a prompt increase in Ca^{2+} that was completely blocked in CPA and BAPTA loaded cultures, whereas 10 μ M methylarachidonyl fluorophosphate (MAFP), a nonspecific inhibitor of both cPLA₂ and iPLA₂, or 10 μ M of bromoenol lactone (Bel) (Cornell-Bell et al.), a specific inhibitor of iPLA₂, did not affect ATP induced Ca^{2+} rises in astrocytic culture (**Figure 1A-C**).

As a broad-based approach to AA-specific lipidomics, Ca^{2+} -independent release of AA and/or its metabolites was performed using a [³H]-AA assay (**Figure 2A**). Cultured rat astrocytes were pre-incubated overnight with [³H]-AA, providing sufficient time for its incorporation into multiple biosynthetic and metabolic pathways. This would include incorporation into membrane phospholipids, which are precursors for endocannabinoids, and related lipids that are themselves precursors for AA release (Chen and Chen, 1998; Strokin et al., 2003). Therefore, this assay serves to determine if any AA precursors or metabolites derived from the [³H]-AA incorporated into the cell are being released; hereafter we refer to the composite of [³H]-labeled AA metabolites as [³H]-AA. ATP alone failed to induce a detectable release of [³H]-AA (**Figure 2B**); however, in cultures pretreated for 10 to 12 minutes with CPA or BAPTA-AM, ATP evoked a robust increase in the release of [³H]-AA, whereas CPA alone had no such effect (**Figure 2B**). Similarly, we found that upon stimulation with a combination of the non-selective mGluR agonist tACPD (100 μ M) and the ionotropic glutamate receptor agonist AMPA (100 μ M), we observed a significant release of [³H]-AA when cytosolic Ca^{2+} was blocked with CPA, but not in cultures without CPA pretreatment (**Figure 2C**). These observations confirm that rat astrocytes can release AA derivatives, key precursors of bioactive eicosanoids (Strokin et al., 2003; Rosenegger et al., 2015), but that the AA release is, surprisingly, inhibited by increases in cytosolic Ca^{2+} .

To explore the mechanism of Ca^{2+} -independent release of $[^3\text{H}]\text{-AA}$, we next evaluated whether inhibition of the Ca^{2+} -sensitive cPLA₂ or the Ca^{2+} -insensitive iPLA₂ enzymes would reduce $[^3\text{H}]\text{-AA}$ release. Pretreatment of astrocytes with 10 μM MAFP significantly decreased the ATP-evoked release of $[^3\text{H}]\text{-AA}$ in cultures exposed to CPA (**Figure 2D**). Similarly, 10 μM BEL treatment suppressed the release of $[^3\text{H}]\text{-AA}$, thus confirming a role for iPLA₂ in Ca^{2+} -independent lipid release (**Figure 2D**). The $[^3\text{H}]\text{-AA}$ release was only observed when agonist-induced increases in cytosolic Ca^{2+} were blocked (**Figures 2B-C**), suggesting that intracellular Ca^{2+} inhibits the Ca^{2+} -independent iPLA₂.

Calmodulin is a potent Ca^{2+} -dependent inhibitor of iPLA₂ (Wolf and Gross, 1996). To assess the interaction between calmodulin and iPLA₂, the cells were treated with calmidazolium (CMZ), an inhibitor of Ca^{2+} /calmodulin interactions that has been shown to remove the calmodulin block of iPLA₂ (Wolf and Gross, 1996). In the presence of CMZ (2 μM), ATP treatment led to a significant release of $[^3\text{H}]\text{-AA}$, which was comparable to the release upon blocking increases in cytosolic Ca^{2+} by preloading astrocytes with either CPA or BAPTA (**Figure 2D**). This observation suggests that Ca^{2+} acts primarily as a brake through calmodulin, which effectively inhibits iPLA₂ activity (Wolf and Gross, 1996; Wolf et al., 1997). These findings are consistent with previous studies showing that iPLA₂ is involved in receptor-mediated AA release from pancreatic islet cells (Gross et al., 1993), smooth muscle cells (Lehman et al., 1993), and endothelial cells (Seegers et al., 2002). Taken together, these data provide evidence of a new signaling mechanism in astrocytes through Ca^{2+} -independent iPLA₂, which is the major PLA₂ isoform in the brain, accounting for 70% of total PLA₂ activity (Yang et al., 1999).

Targeted lipidomics reveals Ca^{2+} -independent lipid production in cultured astrocytes

Using lipid extraction and partial purification methods coupled to HPLC/MS/MS, we performed targeted lipidomics screening on cultured astrocytes that were preincubated with CPA and then challenged with ATP, versus vehicle control astrocytes. Of the 50 lipids screened, 30 were present in each of the samples and could thus be used for comparative analyses.

Figure 3A lower panel lists all of the concentrations as mean $\pm$ SEM (**Figure 3A**). **Figure 3A** summarizes those lipids showing significant concentration differences, as well as the magnitudes of the differences. Of the 30 lipids detected, 16 (including AA and PGE₂) increased upon ATP challenge in the presence of CPA. **Figures 3B-C** are representative chromatograms from the HPLC/MS/MS methods used to detect PGE₂. Notably, 5 of the 8 AA-derivatives in the set were significantly increased by ATP challenge with CPA preincubation, including docosahexaenoyl ethanolamine, although the most dramatic increases were of the prostaglandin, PGE₂. **Figure 3D** shows representative concentration differences in PGE₂, DEA, and AA.

Ca²⁺-independent release of PGE₂ from cultured astrocytes

Given that Ca²⁺-dependent astrocyte lipid release has been previously implicated in vasoregulation (Zonta et al., 2003; Takano et al., 2006; Gordon et al., 2008), it was surprising that PGE₂ was released via a Ca²⁺-independent mechanism. Our HPCL/MS/MS method required large quantities of cells to detect PGE₂, whereas the PGE₂ ELISA assay accurately detects PGE₂ release in 24-well cultures. Therefore, we used ELISA to explore and define the Ca²⁺ dependence of astrocytic PGE₂ release. Because iPLA₂ is essential for Ca²⁺-independent liberation of AA-derived lipids, we first assessed whether knockdown of iPLA₂ via viral transduction would inhibit astrocytic release of PGE₂. In the presence of CPA (20 μ M), ATP (100 μ M) failed to induce PGE₂ release in shRNA-transduced astrocytic cultures, whereas there was a significant increase in the release of PGE₂ by ATP in the presence of CPA in control shRNA cultures (**Figure 4A**). Knockdown of iPLA₂ via shRNA viral transduction was confirmed with western blot analysis (**Figure 4B**).

Since cultured astrocytes express mGluR5 (Balazs et al., 1997; Silva et al., 1999; Gebremedhin et al., 2003), we next tested whether the agonists tACPD (100 μ M) and AMPA (100 μ M) in the presence/absence of CPA evoked PGE₂ release. In the absence of CPA, these agents evoked little to no release of PGE₂ (**Figure 4C**). However, when CPA was used to block

the release of Ca^{2+} from internal stores, tACPD and AMPA significantly increased PGE_2 release (**Figure 4C**).

The most abundant mGluR receptors expressed by astrocytes are mGluR5 and mGluR3 (Petreria et al., 1996; Aronica et al., 2000; Tamaru et al., 2001). However, mGluR5 is developmentally regulated and is not expressed by astrocytes in the adult brain, whereas mGluR3 is persistently expressed at high levels throughout adulthood (Sun et al., 2013). It is important to establish whether activation of mGluR3 can induce Ca^{2+} -independent PGE_2 release, since activation of this receptor has recently been shown to induce Ca^{2+} transients in adult hippocampal astrocytes (Haustein et al., 2014; Tang et al., 2015). In the presence of CPA, the mGluR3 agonists NAAG and LY379268 (Wroblewska et al., 1997; Bond et al., 1999) evoked a significant increase in the release of PGE_2 , whereas the same agonists failed to release PGE_2 in the absence of CPA (**Figure 4D**). Notably, this result shows that activation of an astrocytic Gi-coupled receptor can release gliotransmitters by a Ca^{2+} -independent mechanism.

To extend the observation to human astrocytes, we performed the same experiments in primary cultured astrocytes harvested from human embryonic tissue (Windrem et al., 2004; Windrem et al., 2008; Han et al., 2013). After pharmacologically assessing iPLA₂ activity in rat astrocytes (**Figure 2D**), we evaluated iPLA₂ expression in all the culture models. Western blot analysis showed that iPLA₂ expression was not limited to rat astrocytic cultures, but was also expressed in human and mouse astrocytes (**Figure 4E**). In the presence of CPA, ATP evoked a significant PGE_2 release from cultured human astrocytes, whereas little to no release was observed with ATP alone (**Figure 4F**). Likewise, co-application of tACPD and AMPA showed the same effect (**Figure 4F**). Taken together, these findings show that ATP or mGluR3 activation leads to Ca^{2+} -independent PGE_2 release and that iPLA₂ is expressed in mouse, rat, and human astrocyte cultures.

Ca^{2+} -independent astrocytic lipid release enhances mEPSPs via Kv channel blockade

Thus far, experiments have demonstrated that cultured astrocytes in response to agonist exposure can indeed release lipids in a Ca^{2+} -independent pathway. In fact, the *in vitro* analysis showed that agonist induced Ca^{2+} increases served to impede PGE_2 release, whereas CPA, BAPTA, and CMZ pretreatment potentiated PGE_2 release.

Earlier studies have shown that AA and its metabolite PGE_2 inhibit neuronal Kv currents (Horimoto et al., 1997; Nicol et al., 1997; Evans et al., 1999) and thereby enhance excitability (Sekiyama et al., 1995; Chen and Bazan, 2005; Sang et al., 2005). We therefore evaluated whether astrocytic lipid release also inhibits neuronal Kv current and enhances neuronal excitability in acute hippocampal slices. We performed dual patch-clamp recordings of pairs of CA1 pyramidal neurons and astrocytes in acute hippocampal brain slices prepared from 12–18-day old mice (**Figure 5A**). To isolate transiently active potassium currents, we used 100 ms voltage ramps, which is sufficiently fast to capture transient A-type currents (Phillips et al., 2018). It is worth noting that the I-V ramp enables one to discern what voltage deflection in outward current a drug might affect (Jackson and Bean, 2007). Kv currents in CA1 neurons are an assay for astrocytic lipid release since these currents are sensitive to AA and/or its metabolites (Villarroel and Schwarz, 1996; Carta et al., 2014). We isolated the Kv currents by adding 1 mM QX314 in the patch pipette to block sodium channels (Talbot and Sayer, 1996; Kim et al., 2010) and by imposing a voltage ramp (from -100 mV to 50 mV) every 5 seconds to continuously monitor changes in Kv channel current (Ji et al., 2000; Rangroo Thrane et al., 2013b; Carta et al., 2014) as an assay for agonist-induced astrocytic lipid release (Figure 4A). As expected, direct puffing of PGE_2 (50 μM) or AA (50 μM) from a micropipette significantly reduced the Kv current, in keeping with previous studies (Evans et al., 1999; Carta et al., 2014) (**Figure 5B-C**).

To assess the effects of astrocytic Ca^{2+} signaling on neighboring neuronal Kv channels, we first stimulated astrocytes with ATP or TFLLR- NH_2 , an agonist of protease-activated receptor-1 (PAR1) that is primarily expressed by astrocytes (Shigetomi et al., 2008). ATP (100 μM) induced

a comparable change in neuronal Kv when BAPTA (50 μ M) was present in the astrocyte patch pipette, but not in its absence (**Figure 5B-C**). A similar transient decrease in Kv current was observed when astrocytes were activated by TFLLR-NH₂ (30 μ M). Again, the reduction in neuronal Kv currents was only detected when TFLLR was applied, and when BAPTA was added to the astrocytic pipette solution (**Fig. 5B-C**). To selectively activate astrocytic Ca²⁺ signaling, we next employed acute hippocampal slices from MrgA1 transgenic animals (MrgA1^{+/+}), which express the exogenous Gq-coupled MRG receptor (MrgA1) under control of the GFAP promoter. The MrgA1 agonist Phe-Met-Arg-Phe amide (FMRF) mobilizes intracellular astrocytic Ca²⁺ stores, enabling assessment of the effects of astrocytic Ca²⁺ signaling on neighboring neuronal Kv currents. Although FMRF (15 μ M) induces potent and selective increases in astrocytic cytosolic Ca²⁺ in hippocampal slices (Fiacco et al., 2007; Agulhon et al., 2010; Wang et al., 2012), we saw no detectable changes in neuronal Kv current when the pipette used to patch the astrocyte did not contain BAPTA (**Figure 5B-C**). However, when BAPTA was present in the astrocyte patch pipette, there was a marked decrease in neuronal Kv current evoked by FMRF exposure (**Figure 5B-C**). Thus, astrocytes can modulate neuronal Kv current via a previously undocumented Ca²⁺-independent lipid release.

To assess whether a decrease in neuronal Kv current is a consequence of Ca²⁺-independent astrocytic lipid release, we next employed specific lipid receptor antagonists for PGE₂ and endocannabinoids. Since PGE₂ receptors 1, 2, 3, and 4 are expressed on hippocampal pyramidal neurons (Andreasson, 2010; Maingret et al., 2017), we employed AH6809, a PGE₂ EP1, 2, and 3 antagonist (Abramovitz et al., 2000; Ganesh, 2014), and GW627368, a PGE₂ EP4 antagonist (Jones and Chan, 2005; Wilson et al., 2006). With AH6809 (10 μ M) and GW627368X (3 μ M) in the perfusion solution, the TFLLR, FMRF, and ATP induced Ca²⁺-independent PGE₂ decrease in neuronal Kv current was abolished (**Figure 5C**); In contrast, the CB1 antagonist AM251 (5 μ M) failed to abolish the decrease in Kv current (**Figure 5C**). Taken

together, these data suggest that the observed decrease in neuronal Kv current is a result of astrocytic Ca^{2+} -independent release of PGE_2 .

Interestingly, the presence or absence of BAPTA in the astrocytic pipette solution also affected agonist-induced changes in neuronal membrane potential (**Figure 6A**). Without BAPTA in astrocyte pipettes, TFLLR induced a hyperpolarization (-2.4 ± 0.26 mV) (**Figure 6A**), which has been linked previously to a decrease in extracellular potassium (Wang et al., 2012). However, with BAPTA present in astrocyte pipettes, TFLLR induced depolarization (2.1 ± 0.25 mV) (**Figure 6A**), an effect possibly attributable to the blockage of potassium current.

PGE_2 has previously been shown to enhance neuronal mEPSPs (Sekiya et al., 1995) (Sang et al., 2005). We found that TFLLR-induced activation of astrocytes triggered a decrease in mEPSP amplitude and frequency in normal mouse acute hippocampal slices (**Figure 6B**). The opposite effect, an increase in the amplitude and frequency of mEPSPs, was observable only when BAPTA was added to the astrocytic pipette solution (**Figure 6C**). This observation supports the notion that astrocytic Ca^{2+} -independent lipid release may function as a signaling mechanism capable of modulating proximal synaptic activity by blocking Kv channels, which in turn may increase the frequency and amplitude of mEPSPs. Interestingly, TFLLR's opposing effects on mEPSP amplitude and frequency were dependent upon astrocytic Ca^{2+} levels (**Figure 6D-E**), suggesting that Ca^{2+} depletion provides a pathway favoring the enhancement by astrocytic PGE_2 of neuronal excitability.

DISCUSSION

Lipidomics has the potential to open exciting new avenues within the field of gliotransmission. In the present study, we utilized a series of lipidomics methodologies to show that Ca^{2+} -chelation followed by metabotropic glutamate or purinergic receptor stimulation provokes the formation of a variety of lipids (**Figures 2, 3**) in astrocytes and drives the release of PGE_2 in rat

and human embryonic astrocyte cultures (**Figure 4**). In addition, we demonstrated that receptor-mediated Ca^{2+} -independent PGE_2 release serves to modulate neuronal Kv channels, resulting in enhanced synaptic activity in slices detected as increased mEPSP amplitude and frequency (**Figures 5, 6**). Multiple lines of evidence presented herein also support our hypothesis that this agonist-induced and Ca^{2+} -independent lipid release from astrocytes depends on activation of iPLA_2 . These observations represent, to our knowledge, the first demonstration of Ca^{2+} -independent release of lipid gliotransmitters, and thus add a novel dimension to our understanding of glial-neuronal communication.

Importance of iPLA_2 pathway in Ca^{2+} -independent astrocytic lipid production

Although astrocytes are known to constitutively release AA and other fatty acids (Moore, 2001; Bouyakdan et al., 2015), the pathways involved in constitutive AA release are currently not known, but inverse agonists of purinergic receptors (Ding et al., 2006) and mGluR (Carroll et al., 2001) might serve as useful pharmacological tools for investigating the constitutive PLA_2 activity in relation to gliotransmission. To date, many studies have shown the multifaceted functions of PLA_2 in the CNS, but iPLA_2 accounts for 70% of the PLA_2 activity in the rat brain (Yang et al., 1999). Although the effects of cPLA_2 activation are best documented (Malaplate-Armand et al., 2006; Schaeffer and Gattaz, 2007; Kim et al., 2008), iPLA_2 has also been shown to participate in phospholipid remodeling (Sun et al., 2004), and to regulate hippocampal AMPA receptors involved in learning and memory (Menard et al., 2005). Furthermore, iPLA_2 regulates store-operated Ca^{2+} entry in cerebellar astrocytes (Singaravelu et al., 2006), and provides neuroprotection in an oxygen-glucose deprivation model (Strokin et al., 2006). Astrocytes express both isoforms of PLA_2 (Sun et al., 2005), and numerous studies document the release of AA and its metabolites under regulation of Ca^{2+} -dependent PLA_2 (cPLA_2) signaling in astrocytes (Bruner and Murphy, 1990; Stella et al., 1994; Stella et al., 1997; Chen and Chen, 1998). Previous studies have demonstrated astrocytic release of DHA via iPLA_2 (Strokin et al., 2003, 2007). Here, we focused on the largely unexplored iPLA_2 lipid pathway mediating PGE_2

release from astrocytes. Given the ubiquitous expression of iPLA₂ in astrocytes throughout the brain, we contend that various receptor-activated pathways converge on iPLA₂ for the regulation of astrocyte lipid release (**Figure 7**).

Ca²⁺-independent astrocytic PGE₂ release for rapid modulation of neuronal Kv channels

An important aspect of astrocytic gliotransmitter release is its timing. Agonist-induced astrocytic Ca²⁺ increases occur on a slow time scale of seconds (Cornell-Bell et al., 1990; Wang et al., 2006; Srinivasan et al., 2015), which allows integration of many factors on that time scale. Increased intracellular Ca²⁺ is a key step in the release of gliotransmitters (Parpura et al., 1994; Bezzi et al., 1998; Kang et al., 1998), such as ATP (Coco et al., 2003; Parpura and Zorec, 2010; Illes et al., 2019) and D-Serine (Mothet et al., 2000; Yang et al., 2003; Li et al., 2018; Neame et al., 2019). We have recently shown that agonist-induced astrocytic Ca²⁺ signaling can modulate synaptic activity by promoting K⁺ uptake, resulting in a transient lowering of extracellular K⁺ and depression of synaptic activity (Wang et al., 2012). More recently, we demonstrated that astrocytes can modulate a rapid form of synaptic activity (≥ 500 ms) via 2AG release upon agonism of mGluR3 receptors (Smith et al., 2019). Because receptor-mediated astrocytic Ca²⁺ drives K⁺ uptake and release of gliotransmitters, these processes must, therefore, occur over a relatively prolonged time course (>500 ms). However, agonist-induced Ca²⁺-independent iPLA₂ lipid release does not require the mobilization of intracellular Ca²⁺ stores, thus imparting the potential to signal on a much faster time scale, possibly within 10s of milliseconds. We speculate that iPLA₂-mediated lipid release may act as a feedback system that enhances fast synaptic transmission in the short-term or under conditions of minimal neuronal activity, whereas subsequent activity-mediated increases in Ca²⁺ serve as a brake in the form of the calmodulin-dependent inhibition of iPLA₂. The slower Ca²⁺-dependent release of gliotransmitters and stimulation of K⁺ uptake seems more suited toward the slow and widespread modulation of

brain activity (typically inhibition) that occurs in the setting of, for example, tonic activation of the locus coeruleus and consequent norepinephrine release (Bekar et al., 2008; Ding et al., 2013).

Voltage-gated potassium channels are located on the dendrites of hippocampal pyramidal neurons (Johnston et al., 2000), where they play a major role in controlling dendritic excitability by modulating the amplitude of EPSPs. A morphological study found that the density of Kv channels in the dendrites of pyramidal neurons increased 5-fold proceeding from the soma to the most distal point measured in the apical dendrites (Hoffman et al., 1997). Inhibition of voltage-gated K⁺ currents will consequently enhance EPSPs, possibly explaining why PGE₂ enhances synaptic transmission and LTP in the hippocampus (Sang et al., 2005).

Lipidomics models as a guide for future gliotransmitter discoveries

Lipidomics is one of the fastest-growing branches of the metabolomics field, and offers the possibility of describing the enormous diversity of lipid species throughout the body, and especially in the brain, which is largely composed of lipids (Sinclair, 1975). The complex and fine structure of astrocytic processes gives astrocytes a larger surface area to volume ratio than most other cell types, which makes them uniquely responsive to changes in the extracellular milieu. HPLC/MS/MS lipidomics techniques have provided a means to quantify specific lipid species with an accuracy and sensitivity not possible with more traditional methods employing radio-labeled fatty acids and ELISA. Here, we have combined these classical techniques with lipidomics to test the hypothesis that lipids are produced and released from astrocytes in a Ca²⁺-independent manner. Although our present focus was on the release of PGE₂, we found upregulated release of 15 additional lipids. Notably, among the 30 lipids measured in the present targeted HPLC/MS/MS assay, 14 did not change upon activation of glial receptors, supporting the specificity of GPCR stimulated, Ca²⁺-independent lipid release. Interestingly, two of the lipids released in response to agonist exposure, namely *N*-arachidonyl taurine and *N*-palmitoyl tyrosine, were recently shown to activate TRPV4, a cation channel involved in osmotic

sensitivity and mechanosensitivity (Raboune et al., 2014). Emerging evidence suggests that a TRPV4/AQP4 complex regulates the response to hypo-osmotic stress in astrocytes (Benfenati et al., 2011). The present lipidomics data thus casts light on unpredicted signaling mechanisms that involve astrocytic-derived lipid modulators (**Figure 3**).

Potential physiological relevance of calcium-independent lipid release

Astrocytes are now known to respond to single experimental stimulation events with small calcium increases in their distal fine processes (Panatier et al., 2011). This results from synchronous firing of as many as 50-1000 synapses in the astrocytes microdomain, as may occur when a large stimulation is applied as far as 500 μm away. However, the spontaneous activity of asynchronous synaptic events need not elicit astrocyte calcium events. Metabotropic GluR- and AMPA-mediated activation of astrocytes may elicit iPLA₂ activation and the concerted liberation of AA metabolites and PGE₂ to affect pre- and post-synaptic potassium channels on nearby neurons. When stimulation frequency and/or amplitude increase, astrocytic calcium rises would then serve to suppress over-activation of synapses. In this way, astrocytes may help maintain the strength of relatively quiescent synapses in a calcium-independent fashion, while employing multiple calcium-dependent mechanisms to suppress over-activation. Substantiation of this model requires rigorous testing of the spatial and temporal dynamics of astrocytic and individual dendritic activation, but are limited by current methodological capabilities that do not allow electrical recording of individual dendritic spines.

In conclusion, Ca²⁺-independent astrocytic lipid release constitutes a largely unexplored factor in the regulation of complex neuro-glial signaling interactions. Our present analysis adds a new dimension to the understanding of agonist-induced Ca²⁺ signaling by demonstrating that agonism of several Gq and Gi-linked astrocytic receptors can promote the release of lipid modulators, and that increases in cytosolic Ca²⁺ act as a brake to prevent PGE₂ release.

Author contributions: F.W., H.B.B., and N.A.S. performed the experiments. F.W., H.B.B., J.X., and N.A.S. analyzed data. S.P. and B.J. performed Western Blots. S.G. and B.L. made cultures. L.B. and N.A.S. planned the experiments. D.C.M. provided Human Astrocytes. F.W., H.B.B., and N.A.S. wrote the paper.

Acknowledgments: We thank Dr. Ken McCarthy for generously sharing transgenic mice. This work was supported by the National Institutes of Health Grant K01NS110981, NSFNCs-FR 1926781, and Department of Defense Army Research Office Award W91NF2020189 to N.A.S. We thank Vittorio Gallo, Baljit Khakh, Bartosz Kula, Stefano Vicini, Alexander S. Thrane, Vinita Rangroo Thrane, Takahiro Takano, Paul Cumming, and Fernando R. Fernandez for comments and critical discussion for this manuscript. The authors declare no competing financial interest.

Figure Legends

Figure 1: GPCR-evoked Ca^{2+} transients in astrocytic cultures.

A.) Cultured astrocytes were transduced with AAV GFAP cyto-GCaMP6f and the fluorescent changes associated with intracellular Ca^{2+} increases upon ATP stimulation were measured using confocal microscopy. B.) Representative individual traces of GCaMP6f fluorescence changes ($\Delta F/F_0$) in response to ATP stimulation (upper panel). C.) BAPTA-AM (20 μM , n=5 wells) and CPA (20 μM , n=5 wells) eliminated the ATP (100 μM , n=5 wells) evoked Ca^{2+} increases in fluorescent. MAFP (10 μM , n=5 wells), Bel (10 μM , n=5 wells) did not inhibit ATP (100 μM) induced Ca^{2+} rises in Astrocytes. *** $P \leq 0.0001$, Tukey's *post-hoc* test. All bar graphs show means $\pm$ s.e.m.

Figure 2: GPCR-mediated Ca^{2+} independent release of ^3H -AA and/ or its metabolites from astrocytic cultures.

A.) Schematic of the ^3H -AA Radioactive Assay. B.) The effects of ATP (100 μM , n=45 wells), CPA (20 μM , n=9 wells), ATP/CPA (n=44 wells), and ATP/BAPTA-AM (n=24 wells) on Ca^{2+} independent release of [^3H]-AA and its derivatives in astrocytic cultures compared to control and each other. *** $P \leq 0.0001$; ** $P \leq 0.001$; * $P \leq 0.05$, Tukey's *post-hoc* test. C.) The effects of mGluR agonists tACPD/AMPA (100 μM , n=7), tACPD/AMPA/CPA (n=7 wells), and ATP/CPA (n=7 wells) on Ca^{2+} -independent release of [^3H]-AA and its derivatives in astrocytic cultures compared to control and individual effects. *** $P \leq 0.0001$; ** $P \leq 0.001$; * $P \leq 0.05$, Tukey's *post-hoc* test. D.) Effects of the iPLA₂ inhibitor, BEL (10 μM , n=12 wells), cPLA₂ inhibitor, MAFP (10 μM , n=12 wells), or Calmodulin/ Ca^{2+} complex inhibitor, CMZ (2 μM , n=20 wells) co-applied with ATP on Ca^{2+} independent release of [^3H]-AA and its derivatives in astrocytic cultures compared to ATP alone. *** $P \leq 0.0001$; ** $P \leq 0.001$, Tukey's *post-hoc* test. All bar graphs show means $\pm$ s.e.m.

Figure 3. Targeted lipidomics of rat astrocytic cultures using HPLC/MS/MS.

A.) Summary of the significant differences and the magnitude of change between the vehicle controls and the CPA/ATP stimulated cells for the 30 lipids in the screen that were detected across samples. Dark green denotes significant differences ($p \leq 0.05$); light green denotes trending increases ($0.10 \geq p \geq 0.05$). No significant difference is signified by “*n.d.*”. Arrows indicate the magnitude of effects, as indicated in the legend. *Raw data and statistical analyses for all lipids detected are shown in figure table.* B.) Example chromatograph of a PGE₂ standard (1 pmol) as analyzed by the HPLC/MS/MS method optimized for this standard in negative ion mode. C.) Example of an analysis of a partially purified astrocytic extract using the HPLC/MS/MS method that is optimized for PGE₂. D.) Differences in astrocytic production of AA and two of its derivatives, PGE₂ and docosahexaenoyl ethanolamine (DEA), expressed as the average moles/sample; vehicle n=3 flask, CPA/ATP n=4 flask.

Figure 4: GPCR-mediated Ca²⁺ independent release of PGE₂ from astrocytic cultures.

A.) The effects of ATP (100 μ M, n=4 wells) and ATP/CPA (n=4 wells) on Ca²⁺ independent PGE₂ release in rat astrocytic cultures virally transduced with shRNA to knockdown iPLA₂ compared to control shRNA cultures. * $P \leq 0.05$, Dunn post-test. B.) Western blot analysis of iPLA₂ knockdown via shRNA. β -actin specific antibody was used to standardize for equal protein loading. C.) The effects of GPCR agonists ATP (100 μ M, n=12 wells), tACPD/AMPA (100 μ M, n=7 wells), tACPD/AMPA/CPA (n=7), and ATP/CPA (n=12 wells) on Ca²⁺-independent PGE₂ release in rat astrocytic cultures compared to control. **** $P \leq 0.0001$; *** $P \leq 0.001$, Tukey’s post-test. D.) The effects of ATP/CPA (100 μ M, n=4) and mGluR3 agonist NAAG (100 μ M, n=4 wells), LY379268 (100 μ M, n=4 wells), NAAG/CPA (n=4), and LY379268/CPA (n=4 wells) on Ca²⁺-independent PGE₂ release in rat astrocytic cultures compared to control and each other. *** $P \leq 0.001$, Tukey’s post-test. E.) Western blot analysis of iPLA₂ protein for human, rat, and mouse cultures. β -actin specific antibody was used to standardize for equal protein loading. F.) The effects of GPCR agonists ATP (100 μ M, n=8 wells), ATP/CPA (n=8 wells), tACPD/AMPA (100 μ M, n=8 wells), and tACPD/AMPA/CPA (n=8 wells) on Ca²⁺-independent PGE₂ release in

human astrocytic cultures compared to control and each other. *** $P \leq 0.0001$; ** $P \leq 0.001$; * $P \leq 0.05$, Tukey's *post-hoc* test. All bar graphs show means $\pm$ s.e.m.

Figure 5: Ca^{2+} -independent astrocytic lipid release inhibits Kv channel blockade.

A.) Image of a patched astrocyte and a patched neuron, filled with fluorescent dyes. B.) Direct application of AA (50 μM) and PGE2 (50 μM) decreased Kv current. ATP (100 μM) and TFLLR (30 μM) together with BAPTA also decreased Kv current, as can be seen in the peak currents. FMRF (15 μM) in the MrgA1^{+/-} mice applied in the presence of BAPTA decreased Kv current, but not in the absence of BAPTA. Traces to the left are representative recordings of changes to membrane voltages in the presence of drugs described above. C.) Quantitation of effects of different drugs on the holding voltage ramp induced voltage gated K^+ currents (*, $p < 0.05$, **, $p < 0.01$, $n = 5-7$ mice). All bar graphs show mean $\pm$ s.e.m.

Figure 6: Ca^{2+} -independent astrocytic lipid release increase mEPSP amplitude and frequency.

A.) Representative traces showing changes to membrane potentials VM upon TFLLR application in the presence or absence of BAPTA, and quantitations are shown in inset (**, $p < 0.01$, t-test, $n = 5-6$ mice). B.) Representative trace showing hyperpolarization and reduction in mEPSP amplitude and frequency in excitatory hippocampal neurons upon TFLLR (30 μM) application without BAPTA being present in neighboring astrocytes. C.) Representative trace showing depolarization and an increase in mEPSP amplitude and frequency in excitatory hippocampal neurons upon TFLLR (30 μM) application with BAPTA present in neighboring astrocytes. Lower traces are extensions of parts of recordings in upper trace. D.) Comparison of the cumulative mEPSP amplitude and cumulative mEPSP frequency distribution recorded in B). E.) Statistical analysis of the amplitude and inter-event intervals before and after TFLLR application, with and without BAPTA present (**, $p < 0.01$, paired t-test, $N = 5-7$ mice). All bar graphs show mean $\pm$ s.e.m.

Figure 7: Models of GPCR-mediated Ca^{2+} -independent and Ca^{2+} -dependent lipid release in astrocytes.

A.) Schematic of receptor mediated Ca^{2+} -Independent lipid release across the astrocytic membrane. B.) Schematic of receptor mediated Ca^{2+} -Dependent lipid release across the astrocytic membrane.

References

- Abramovitz M, Adam M, Boie Y, Carriere M, Denis D, Godbout C, Lamontagne S, Rochette C, Sawyer N, Tremblay NM, Belley M, Gallant M, Dufresne C, Gareau Y, Ruel R, Juteau H, Labelle M, Ouimet N, Metters KM (2000) The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs. *Biochim Biophys Acta* 1483:285-293.
- Agulhon C, Fiacco TA, McCarthy KD (2010) Hippocampal short- and long-term plasticity are not modulated by astrocyte Ca²⁺ signaling. *Science* 327:1250-1254.
- Akiba S, Hayama M, Sato T (1998) Inhibition of Ca²⁺-independent phospholipase A2 by bromoenol lactone attenuates prostaglandin generation induced by interleukin-1 beta and dibutyryl cAMP in rat mesangial cells. *FEBS Lett* 437:225-228.
- Andreasson K (2010) Emerging roles of PGE2 receptors in models of neurological disease. *Prostaglandins Other Lipid Mediat* 91:104-112.
- Aronica E, van Vliet EA, Mayboroda OA, Troost D, da Silva FH, Gorter JA (2000) Upregulation of metabotropic glutamate receptor subtype mGluR3 and mGluR5 in reactive astrocytes in a rat model of mesial temporal lobe epilepsy. *The European journal of neuroscience* 12:2333-2344.
- Balazs R, Miller S, Romano C, de Vries A, Chun Y, Cotman CW (1997) Metabotropic glutamate receptor mGluR5 in astrocytes: pharmacological properties and agonist regulation. *J Neurochem* 69:151-163.
- Bekar LK, He W, Nedergaard M (2008) Locus coeruleus alpha-adrenergic-mediated activation of cortical astrocytes in vivo. *Cerebral cortex* 18:2789-2795.
- Benfenati V, Caprini M, Dovizio M, Mylonakou MN, Ferroni S, Ottersen OP, Amiry-Moghaddam M (2011) An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4) complex is essential for cell-volume control in astrocytes. *Proc Natl Acad Sci U S A* 108:2563-2568.
- Bezzi P, Carmignoto G, Pasti L, Vesce S, Rossi D, Rizzini BL, Pozzan T, Volterra A (1998) Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. *Nature* 391:281-285.
- Boland LM, Drzewiecki MM (2008) Polyunsaturated fatty acid modulation of voltage-gated ion channels. *Cell biochemistry and biophysics* 52:59-84.
- Bond A, Ragumoorthy N, Monn JA, Hicks CA, Ward MA, Lodge D, O'Neill MJ (1999) LY379268, a potent and selective Group II metabotropic glutamate receptor agonist, is neuroprotective in gerbil global, but not focal, cerebral ischaemia. *Neurosci Lett* 273:191-194.
- Bouyakdan K, Taib B, Budry L, Zhao S, Rodaros D, Neess D, Mandrup S, Faergeman NJ, Alquier T (2015) A novel role for central ACBP/DBI as a regulator of long-chain fatty acid metabolism in astrocytes. *J Neurochem* 133:253-265.
- Bruner G, Murphy S (1990) ATP-evoked arachidonic acid mobilization in astrocytes is via a P2Y-purinergic receptor. *J Neurochem* 55:1569-1575.
- Carroll FY, Stolle A, Beart PM, Voerste A, Brabet I, Mauler F, Joly C, Antonicek H, Bockaert J, Muller T, Pin JP, Prezeau L (2001) BAY36-7620: a potent non-competitive mGlu1 receptor antagonist with inverse agonist activity. *Mol Pharmacol* 59:965-973.
- Carta M, Lanore F, Rebola N, Szabo Z, Da Silva SV, Lourenco J, Verraes A, Nadler A, Schultz C, Blanchet C, Mulle C (2014) Membrane Lipids Tune Synaptic Transmission by Direct Modulation of Presynaptic Potassium Channels. *Neuron*.

- Chen C, Bazan NG (2005) Endogenous PGE2 regulates membrane excitability and synaptic transmission in hippocampal CA1 pyramidal neurons. *J Neurophysiol* 93:929-941.
- Chen WC, Chen CC (1998) ATP-induced arachidonic acid release in cultured astrocytes is mediated by Gi protein coupled P2Y1 and P2Y2 receptors. *Glia* 22:360-370.
- Coco S, Calegari F, Pravettoni E, Pozzi D, Taverna E, Rosa P, Matteoli M, Verderio C (2003) Storage and release of ATP from astrocytes in culture. *J Biol Chem* 278:1354-1362.
- Cordero-Morales JF, Vasquez V (2018) How lipids contribute to ion channel function, a fat perspective on direct and indirect interactions. *Curr Opin Struct Biol* 51:92-98.
- Cornell-Bell AH, Finkbeiner SM, Cooper MS, Smith SJ (1990) Glutamate induces calcium waves in cultured astrocytes: long-range glial signaling. *Science* 247:470-473.
- Czigler A, Toth L, Szarka N, Szilagyi K, Kellermayer Z, Harci A, Vecsernyes M, Ungvari Z, Szolics A, Koller A, Buki A, Toth P (2019) Prostaglandin E2, a postulated mediator of neurovascular coupling, at low concentrations dilates whereas at higher concentrations constricts human cerebral parenchymal arterioles. *Prostaglandins Other Lipid Mediat* 146:106389.
- Dabertrand F, Hannah RM, Pearson JM, Hill-Eubanks DC, Brayden JE, Nelson MT (2013) Prostaglandin E2, a postulated astrocyte-derived neurovascular coupling agent, constricts rather than dilates parenchymal arterioles. *J Cereb Blood Flow Metab* 33:479-482.
- Dhopeshwarkar GA, Subramanian C (1976) Biosynthesis of polyunsaturated fatty acids in the developing brain: I. Metabolic transformations of intracranially administered 1-14C linolenic acid. *Lipids* 11:67-71.
- Ding F, O'Donnell J, Thrane AS, Zeppenfeld D, Kang H, Xie L, Wang F, Nedergaard M (2013) α 1-Adrenergic receptors mediate coordinated Ca^{2+} signaling of cortical astrocytes in awake, behaving mice. *Cell calcium* 54:387-394.
- Ding Z, Kim S, Kunapuli SP (2006) Identification of a potent inverse agonist at a constitutively active mutant of human P2Y12 receptor. *Mol Pharmacol* 69:338-345.
- Evans AR, Vasko MR, Nicol GD (1999) The cAMP transduction cascade mediates the PGE2-induced inhibition of potassium currents in rat sensory neurones. *J Physiol* 516 (Pt 1):163-178.
- Fiacco TA, Agulhon C, Taves SR, Petravic J, Casper KB, Dong X, Chen J, McCarthy KD (2007) Selective stimulation of astrocyte calcium in situ does not affect neuronal excitatory synaptic activity. *Neuron* 54:611-626.
- Ganesh T (2014) Prostanoid receptor EP2 as a therapeutic target. *J Med Chem* 57:4454-4465.
- Gebremedhin D, Yamaura K, Zhang C, Bylund J, Koehler RC, Harder DR (2003) Metabotropic glutamate receptor activation enhances the activities of two types of Ca^{2+} -activated K^{+} channels in rat hippocampal astrocytes. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 23:1678-1687.
- Gordon GR, Mulligan SJ, MacVicar BA (2007) Astrocyte control of the cerebrovasculature. *Glia* 55:1214-1221.
- Gordon GR, Choi HB, Rungta RL, Ellis-Davies GC, MacVicar BA (2008) Brain metabolism dictates the polarity of astrocyte control over arterioles. *Nature* 456:745-749.
- Gross RW, Ramanadham S, Kruszka KK, Han X, Turk J (1993) Rat and human pancreatic islet cells contain a calcium ion independent phospholipase A2 activity selective for hydrolysis of arachidonate which is stimulated by adenosine triphosphate and is specifically localized to islet beta-cells. *Biochemistry* 32:327-336.

- Han X, Chen M, Wang F, Windrem M, Wang S, Shanz S, Xu Q, Oberheim NA, Bekar L, Betstadt S, Silva AJ, Takano T, Goldman SA, Nedergaard M (2013) Forebrain engraftment by human glial progenitor cells enhances synaptic plasticity and learning in adult mice. *Cell Stem Cell* 12:342-353.
- Haustein MD, Kracun S, Lu XH, Shih T, Jackson-Weaver O, Tong X, Xu J, Yang XW, O'Dell TJ, Marvin JS, Ellisman MH, Bushong EA, Looger LL, Khakh BS (2014) Conditions and constraints for astrocyte calcium signaling in the hippocampal mossy fiber pathway. *Neuron* 82:413-429.
- Hoffman DA, Magee JC, Colbert CM, Johnston D (1997) K⁺ channel regulation of signal propagation in dendrites of hippocampal pyramidal neurons. *Nature* 387:869-875.
- Horimoto N, Nabekura J, Ogawa T (1997) Arachidonic acid activation of potassium channels in rat visual cortex neurons. *Neuroscience* 77:661-671.
- Illes P, Burnstock G, Tang Y (2019) Astroglia-Derived ATP Modulates CNS Neuronal Circuits. *Trends in neurosciences* 42:885-898.
- Jackson AC, Bean BP (2007) State-dependent enhancement of subthreshold A-type potassium current by 4-aminopyridine in tuberomammillary nucleus neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 27:10785-10796.
- Jelsema CL, Axelrod J (1987) Stimulation of phospholipase A2 activity in bovine rod outer segments by the beta gamma subunits of transducin and its inhibition by the alpha subunit. *Proc Natl Acad Sci U S A* 84:3623-3627.
- Ji J, Salapatek AM, Diamant NE (2000) Inwardly rectifying K(+) channels in esophageal smooth muscle. *Am J Physiol Gastrointest Liver Physiol* 279:G951-960.
- Johnston D, Hoffman DA, Magee JC, Poolos NP, Watanabe S, Colbert CM, Migliore M (2000) Dendritic potassium channels in hippocampal pyramidal neurons. *J Physiol* 525 Pt 1:75-81.
- Jones RL, Chan KM (2005) Investigation of the agonist activity of prostacyclin analogues on prostanoid EP4 receptors using GW 627368 and taprostene: evidence for species differences. *Prostaglandins Leukot Essent Fatty Acids* 72:289-299.
- Kang J, Jiang L, Goldman SA, Nedergaard M (1998) Astrocyte-mediated potentiation of inhibitory synaptic transmission. *Nature neuroscience* 1:683-692.
- Kim DH, Fitzsimmons B, Hefferan MP, Svensson CI, Wancewicz E, Monia BP, Hung G, Butler M, Marsala M, Hua XY, Yaksh TL (2008) Inhibition of spinal cytosolic phospholipase A(2) expression by an antisense oligonucleotide attenuates tissue injury-induced hyperalgesia. *Neuroscience* 154:1077-1087.
- Kim JH, Kushmerick C, von Gersdorff H (2010) Presynaptic resurgent Na⁺ currents sculpt the action potential waveform and increase firing reliability at a CNS nerve terminal. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 30:15479-15490.
- Ledo A, Rocha BS, Laranjinha J (2019) Bioactive Lipids and the Gut-Brain Axis: Diet as a Modulator of Bioactivity and Diversity of Lipids in the Brain. *Adv Exp Med Biol* 1127:147-168.
- Lehman JJ, Brown KA, Ramanadham S, Turk J, Gross RW (1993) Arachidonic acid release from aortic smooth muscle cells induced by [Arg8]vasopressin is largely mediated by calcium-independent phospholipase A2. *J Biol Chem* 268:20713-20716.
- Leishman E, Manchanda M, Thelen R, Miller S, Mackie K, Bradshaw HB (2018) Cannabidiol's Upregulation of N-acyl Ethanolamines in the Central Nervous System Requires N-acyl

- Phosphatidyl Ethanolamine-Specific Phospholipase D. *Cannabis Cannabinoid Res* 3:228-241.
- Li S, Uno Y, Rudolph U, Cobb J, Liu J, Anderson T, Levy D, Balu DT, Coyle JT (2018) Astrocytes in primary cultures express serine racemase, synthesize d-serine and acquire A1 reactive astrocyte features. *Biochem Pharmacol* 151:245-251.
- Lin JH, Weigel H, Cotrina ML, Liu S, Bueno E, Hansen AJ, Hansen TW, Goldman S, Nedergaard M (1998) Gap-junction-mediated propagation and amplification of cell injury. *Nat Neurosci* 1:494-500.
- MacVicar BA, Newman EA (2015) Astrocyte regulation of blood flow in the brain. *Cold Spring Harb Perspect Biol* 7.
- Maingret V, Barthet G, Deforges S, Jiang N, Mulle C, Amedee T (2017) PGE2-EP3 signaling pathway impairs hippocampal presynaptic long-term plasticity in a mouse model of Alzheimer's disease. *Neurobiol Aging* 50:13-24.
- Malaplate-Armand C, Florent-Bechard S, Youssef I, Koziel V, Sponne I, Kriem B, Leininger-Muller B, Olivier JL, Oster T, Pillot T (2006) Soluble oligomers of amyloid-beta peptide induce neuronal apoptosis by activating a cPLA2-dependent sphingomyelinase-ceramide pathway. *Neurobiol Dis* 23:178-189.
- Menard C, Valastro B, Martel MA, Chartier E, Marineau A, Baudry M, Massicotte G (2005) AMPA receptor phosphorylation is selectively regulated by constitutive phospholipase A(2) and 5-lipoxygenase activities. *Hippocampus* 15:370-380.
- Meves H (1994) Modulation of ion channels by arachidonic acid. *Prog Neurobiol* 43:175-186.
- Moore SA (1993) Cerebral endothelium and astrocytes cooperate in supplying docosahexaenoic acid to neurons. *Adv Exp Med Biol* 331:229-233.
- Moore SA (2001) Polyunsaturated fatty acid synthesis and release by brain-derived cells in vitro. *J Mol Neurosci* 16:195-200; discussion 215-121.
- Moore SA, Yoder E, Murphy S, Dutton GR, Spector AA (1991) Astrocytes, not neurons, produce docosahexaenoic acid (22:6 omega-3) and arachidonic acid (20:4 omega-6). *J Neurochem* 56:518-524.
- Mothet J, Parent A, Wolosker H, Jr RB, Linden D, Ferris C, Rogawski M, Synder S (2000) D-serine is an endogenous ligand for the site of N-methyl D-aspartate receptor. *Proc Natl Acad Sci U S A* 97:4926-4931.
- Murayama T, Kajiyama Y, Nomura Y (1990) Histamine-stimulated and GTP-binding proteins-mediated phospholipase A2 activation in rabbit platelets. *J Biol Chem* 265:4290-4295.
- Neame S, Safory H, Radziszewsky I, Touitou A, Marchesani F, Marchetti M, Kellner S, Berlin S, Foltyn VN, Engelender S, Billard JM, Wolosker H (2019) The NMDA receptor activation by d-serine and glycine is controlled by an astrocytic Phgdh-dependent serine shuttle. *Proceedings of the National Academy of Sciences of the United States of America* 116:20736-20742.
- Nedergaard M (1994) Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science* 263:1768-1771.
- Nicol GD, Vasko MR, Evans AR (1997) Prostaglandins suppress an outward potassium current in embryonic rat sensory neurons. *J Neurophysiol* 77:167-176.
- Panatier A, Vallee J, Haber M, Murai KK, Lacaille JC, Robitaille R (2011) Astrocytes are endogenous regulators of basal transmission at central synapses. *Cell* 146:785-798.
- Parpura V, Zorec R (2010) Gliotransmission: Exocytotic release from astrocytes. *Brain Res Rev* 63:83-92.

- Parpura V, Basarsky TA, Liu F, Jęftinija K, Jęftinija S, Haydon PG (1994) Glutamate-mediated astrocyte-neuron signalling. *Nature* 369:744-747.
- Petralia RS, Wang YX, Niedzielski AS, Wenthold RJ (1996) The metabotropic glutamate receptors, mGluR2 and mGluR3, show unique postsynaptic, presynaptic and glial localizations. *Neuroscience* 71:949-976.
- Phillips WS, Del Negro CA, Rekling JC (2018) Dendritic A-Current in Rhythmically Active PreBotzinger Complex Neurons in Organotypic Cultures from Newborn Mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 38:3039-3049.
- Piomelli D (1993) Arachidonic acid in cell signaling. *Current opinion in cell biology* 5:274-280.
- Plog BA, Nedergaard M (2018) The Glymphatic System in Central Nervous System Health and Disease: Past, Present, and Future. *Annu Rev Pathol* 13:379-394.
- Portilla D, Shah SV, Lehman PA, Creer MH (1994) Role of cytosolic calcium-independent plasmalogen-selective phospholipase A2 in hypoxic injury to rabbit proximal tubules. *The Journal of clinical investigation* 93:1609-1615.
- Ramakers GM, Storm JF (2002) A postsynaptic transient K(+) current modulated by arachidonic acid regulates synaptic integration and threshold for LTP induction in hippocampal pyramidal cells. *Proc Natl Acad Sci U S A* 99:10144-10149.
- Rangroo Thrane V, Thrane AS, Plog BA, Thiyagarajan M, Iliff JJ, Deane R, Nagelhus EA, Nedergaard M (2013a) Paravascular microcirculation facilitates rapid lipid transport and astrocyte signaling in the brain. *Sci Rep* 3:2582.
- Rangroo Thrane V, Thrane AS, Wang F, Cotrina ML, Smith NA, Chen M, Xu Q, Kang N, Fujita T, Nagelhus EA, Nedergaard M (2013b) Ammonia triggers neuronal disinhibition and seizures by impairing astrocyte potassium buffering. *Nature medicine* 19:1643-1648.
- Rosenegger DG, Tran CH, Wamsteeker Cusulin JJ, Gordon GR (2015) Tonic Local Brain Blood Flow Control by Astrocytes Independent of Phasic Neurovascular Coupling. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 35:13463-13474.
- Sang N, Zhang J, Marcheselli V, Bazan NG, Chen C (2005) Postsynaptically synthesized prostaglandin E2 (PGE2) modulates hippocampal synaptic transmission via a presynaptic PGE2 EP2 receptor. *J Neurosci* 25:9858-9870.
- Schaeffer EL, Gattaz WF (2007) Requirement of hippocampal phospholipase A2 activity for long-term memory retrieval in rats. *J Neural Transm* 114:379-385.
- Seegers HC, Gross RW, Boyle WA (2002) Calcium-independent phospholipase A(2)-derived arachidonic acid is essential for endothelium-dependent relaxation by acetylcholine. *J Pharmacol Exp Ther* 302:918-923.
- Sekiyama N, Mizuta S, Hori A, Kobayashi S (1995) Prostaglandin E2 facilitates excitatory synaptic transmission in the nucleus tractus solitarius of rats. *Neurosci Lett* 188:101-104.
- Shigetomi E, Bowser DN, Sofroniew MV, Khakh BS (2008) Two forms of astrocyte calcium excitability have distinct effects on NMDA receptor-mediated slow inward currents in pyramidal neurons. *J Neurosci* 28:6659-6663.
- Silva GA, Theriault E, Mills LR, Pennefather PS, Feeney CJ (1999) Group I and II metabotropic glutamate receptor expression in cultured rat spinal cord astrocytes. *Neurosci Lett* 263:117-120.
- Sinclair AJ (1975) Long-chain polyunsaturated fatty acids in the mammalian brain. *The Proceedings of the Nutrition Society* 34:287-291.

- Singaravelu K, Lohr C, Deitmer JW (2006) Regulation of store-operated calcium entry by calcium-independent phospholipase A2 in rat cerebellar astrocytes. *J Neurosci* 26:9579-9592.
- Smith NA, Bekar LK, Nedergaard M (2019) Astrocytic Endocannabinoids Mediate Hippocampal Transient Heterosynaptic Depression. *Neurochem Res*.
- Smith NA, Kress BT, Lu Y, Chandler-Militello D, Benraiss A, Nedergaard M (2018) Fluorescent Ca(2+) indicators directly inhibit the Na,K-ATPase and disrupt cellular functions. *Sci Signal* 11.
- Srinivasan R, Huang BS, Venugopal S, Johnston AD, Chai H, Zeng H, Golshani P, Khakh BS (2015) Ca(2+) signaling in astrocytes from *Ip3r2(-/-)* mice in brain slices and during startle responses in vivo. *Nat Neurosci* 18:708-717.
- Stella N, Tence M, Glowinski J, Premont J (1994) Glutamate-evoked release of arachidonic acid from mouse brain astrocytes. *J Neurosci* 14:568-575.
- Stella N, Estelles A, Siciliano J, Tence M, Desagher S, Piomelli D, Glowinski J, Premont J (1997) Interleukin-1 enhances the ATP-evoked release of arachidonic acid from mouse astrocytes. *J Neurosci* 17:2939-2946.
- Strokin M, Sergeeva M, Reiser G (2003) Docosahexaenoic acid and arachidonic acid release in rat brain astrocytes is mediated by two separate isoforms of phospholipase A2 and is differently regulated by cyclic AMP and Ca2+. *Br J Pharmacol* 139:1014-1022.
- Strokin M, Sergeeva M, Reiser G (2007) Prostaglandin synthesis in rat brain astrocytes is under the control of the n-3 docosahexaenoic acid, released by group VIB calcium-independent phospholipase A2. *J Neurochem* 102:1771-1782.
- Strokin M, Chechneva O, Reymann KG, Reiser G (2006) Neuroprotection of rat hippocampal slices exposed to oxygen-glucose deprivation by enrichment with docosahexaenoic acid and by inhibition of hydrolysis of docosahexaenoic acid-containing phospholipids by calcium independent phospholipase A2. *Neuroscience* 140:547-553.
- Sun GY, Xu J, Jensen MD, Simonyi A (2004) Phospholipase A2 in the central nervous system: implications for neurodegenerative diseases. *Journal of lipid research* 45:205-213.
- Sun GY, Xu J, Jensen MD, Yu S, Wood WG, Gonzalez FA, Simonyi A, Sun AY, Weisman GA (2005) Phospholipase A2 in astrocytes: responses to oxidative stress, inflammation, and G protein-coupled receptor agonists. *Mol Neurobiol* 31:27-41.
- Sun W, McConnell E, Pare JF, Xu Q, Chen M, Peng W, Lovatt D, Han X, Smith Y, Nedergaard M (2013) Glutamate-dependent neuroglial calcium signaling differs between young and adult brain. *Science* 339:197-200.
- Takano T, Tian GF, Peng W, Lou N, Libionka W, Han X, Nedergaard M (2006) Astrocyte-mediated control of cerebral blood flow. *Nat Neurosci* 9:260-267.
- Talbot MJ, Sayer RJ (1996) Intracellular QX-314 inhibits calcium currents in hippocampal CA1 pyramidal neurons. *Journal of neurophysiology* 76:2120-2124.
- Tamaru Y, Nomura S, Mizuno N, Shigemoto R (2001) Distribution of metabotropic glutamate receptor mGluR3 in the mouse CNS: differential location relative to pre- and postsynaptic sites. *Neuroscience* 106:481-503.
- Tang W, Szokol K, Jensen V, Enger R, Trivedi CA, Hvalby O, Helm PJ, Looger LL, Sprengel R, Nagelhus EA (2015) Stimulation-evoked Ca2+ signals in astrocytic processes at hippocampal CA3-CA1 synapses of adult mice are modulated by glutamate and ATP. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 35:3016-3021.

- Tay HK, Melendez AJ (2004) FcγRI-triggered generation of arachidonic acid and eicosanoids requires iPLA2 but not cPLA2 in human monocytic cells. *J Biol Chem* 279:22505-22513.
- van Tol-Steye H, Lodder JC, Mansvelder HD, Planta RJ, van Heerikhuizen H, Kits KS (1999) Roles of G-protein βγ, arachidonic acid, and phosphorylation in convergent activation of an S-like potassium conductance by dopamine, Ala-Pro-Gly-Trp-NH₂, and Phe-Met-Arg-Phe-NH₂. *J Neurosci* 19:3739-3751.
- Villarroel A, Schwarz TL (1996) Inhibition of the Kv4 (Shal) family of transient K⁺ currents by arachidonic acid. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 16:2522-2532.
- Wang F, Smith NA, Xu Q, Fujita T, Baba A, Matsuda T, Takano T, Bekar L, Nedergaard M (2012) Astrocytes modulate neural network activity by Ca²⁺-dependent uptake of extracellular K⁺. *Sci Signal* 5:ra26.
- Wang X, Lou N, Xu Q, Tian GF, Peng WG, Han X, Kang J, Takano T, Nedergaard M (2006) Astrocytic Ca²⁺ signaling evoked by sensory stimulation in vivo. *Nature neuroscience* 9:816-823.
- Wilson RJ, Giblin GM, Roomans S, Rhodes SA, Cartwright KA, Shield VJ, Brown J, Wise A, Chowdhury J, Pritchard S, Coote J, Noel LS, Kenakin T, Burns-Kurtis CL, Morrison V, Gray DW, Giles H (2006) GW627368X ((N-{2-[4-(4,9-diethoxy-1-oxo-1,3-dihydro-2H-benzo[f]isoindol-2-yl)phenyl]acetyl} benzene sulphonamide): a novel, potent and selective prostanoid EP4 receptor antagonist. *Br J Pharmacol* 148:326-339.
- Windrem MS, Nunes MC, Rashbaum WK, Schwartz TH, Goodman RA, McKhann G, 2nd, Roy NS, Goldman SA (2004) Fetal and adult human oligodendrocyte progenitor cell isolates myelinate the congenitally dysmyelinated brain. *Nat Med* 10:93-97.
- Windrem MS, Schanz SJ, Guo M, Tian GF, Washco V, Stanwood N, Rasband M, Roy NS, Nedergaard M, Havton LA, Wang S, Goldman SA (2008) Neonatal chimerization with human glial progenitor cells can both remyelinate and rescue the otherwise lethally hypomyelinated shiverer mouse. *Cell stem cell* 2:553-565.
- Winstead MV, Balsinde J, Dennis EA (2000) Calcium-independent phospholipase A(2): structure and function. *Biochimica et biophysica acta* 1488:28-39.
- Wolf MJ, Gross RW (1996) The calcium-dependent association and functional coupling of calmodulin with myocardial phospholipase A2. Implications for cardiac cycle-dependent alterations in phospholipolysis. *J Biol Chem* 271:20989-20992.
- Wolf MJ, Wang J, Turk J, Gross RW (1997) Depletion of intracellular calcium stores activates smooth muscle cell calcium-independent phospholipase A2. A novel mechanism underlying arachidonic acid mobilization. *J Biol Chem* 272:1522-1526.
- Wroblewska B, Wroblewski JT, Pshenichkin S, Surin A, Sullivan SE, Neale JH (1997) N-acetylaspartylglutamate selectively activates mGluR3 receptors in transfected cells. *J Neurochem* 69:174-181.
- Yang HC, Mosior M, Ni B, Dennis EA (1999) Regional distribution, ontogeny, purification, and characterization of the Ca²⁺-independent phospholipase A2 from rat brain. *J Neurochem* 73:1278-1287.
- Yang Y, Ge W, Chen Y, Zhang Z, Shen W, Wu C, Poo M, Duan S (2003) Contribution of astrocytes to hippocampal long-term potentiation through release of D-serine. *Proc Natl Acad Sci U S A* 100:15194-15199.

- Yu X, Taylor AMW, Nagai J, Golshani P, Evans CJ, Coppola G, Khakh BS (2018) Reducing Astrocyte Calcium Signaling In Vivo Alters Striatal Microcircuits and Causes Repetitive Behavior. *Neuron* 99:1170-1187 e1179.
- Zonta M, Angulo MC, Gobbo S, Rosengarten B, Hossmann KA, Pozzan T, Carmignoto G (2003) Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. *Nat Neurosci* 6:43-50.

A **B** **C**

Figure 1

A **B** **C** **D** **Figure 2**

A

Relative differences of lipid levels in astrocytes treated with CPA and ATP compared to vehicle

Prostaglandins	
PGE ₂	↑↑↑
Free fatty acids	
Arachidonic acid	↑
Linoleic acid	n.d.
2-arachidonoyl-sn-glycerol	
2-arachidonoyl-sn-glycerol	↑
2-linoleoyl-sn-glycerol	n.d.
2-oleoyl-sn-glycerol	n.d.
N-acyl ethanolamine	
N-palmitoyl ethanolamine	↑↑
N-stearoyl ethanolamine	↑↑
N-oleoyl ethanolamine	↑↑
N-linoleoyl ethanolamine	n.d.
N-arachidonoyl ethanolamine	↑
N-docosahexaenoyl ethanolamine	↑↑
N-acyl glycine	
N-palmitoyl glycine	n.d.
N-stearoyl glycine	↑
N-oleoyl glycine	↑
N-linoleoyl glycine	n.d.
N-arachidonoyl glycine	n.d.
N-acyl serine	
N-palmitoyl serine	n.d.
N-stearoyl serine	n.d.
N-oleoyl serine	n.d.
N-arachidonoyl serine	n.d.
N-acyl taurine	
N-arachidonoyl taurine	↑
N-acyl tyrosine	
N-palmitoyl tyrosine	↑
N-stearoyl tyrosine	↑
N-oleoyl tyrosine	↑
N-arachidonoyl tyrosine	n.d.
N-docosahexaenoyl tyrosine	↑
N-acyl tryptophan	
N-palmitoyl tryptophan	n.d.
N-oleoyl tryptophan	n.d.
N-docosahexaenoyl tryptophan	n.d.

Magnitude Differences	
2-2.99 times higher	↑↑↑
1.50-1.99 times higher	↑↑
1-1.49 times higher	↑
Significant Differences	
Significant increase p<0.05	
Trending increase p=0.05-0.10	
no difference between groups	n.d.

B

C

D

Figure 3

Figure 4

A

B

C

Figure 5

Figure 6