ADNKA overcomes SARS-CoV2-mediated NK cell inhibition through non-spike antibodies

Fielding CA¹, Sabberwal P¹, Williamson JC², Greenwood EJD², Crozier TWM², Zelek W¹, Seow J³, Graham C³, Huettnner I³, Edgeworth JD^{3,4}, Morgan BP¹, Ladell K¹, Eberl M¹, Humphreys IR¹, Merrick B^{3,4}, Doores K³, Wilson SJ⁵, Lehner PJ², Wang ECY¹, Stanton RJ^{1*}

1. Division of Infection and Immunity, School of Medicine, Cardiff University, CF14 4XN

2. Cambridge Institute for Therapeutic Immunology & Infectious Disease, Jeffrey Cheah Biomedical Centre, Cambridge Biomedical Campus, University of Cambridge, Cambridge CB2 0AW, UK

3. Department of Infectious Diseases, School of Immunology & Microbial Sciences, King's College London, London, UK

4. Department of Infectious Diseases, Guy's and St Thomas' NHS Foundation Trust, London

5. MRC - University of Glasgow Centre for Virus Research, 464 Bearsden Road, Glasgow, G61 1QH

*Corresponding author, StantonRJ@cardiff.ac.uk

Abstract

SARS-CoV-2 antagonises the cellular interferon response, but whether the virus manipulates cellular immunity is unclear. An unbiased proteomic approach to determine how cell surface protein expression is altered on SARS-CoV-2-infected lung epithelial cells showed downregulation of activating NK cell ligands: B7-H6, MICA, ULBP2, and Nectin1, but no effect on surface MHC-I expression. NK ligand downregulation correlated with a reduction in NK cell activation by infected cells, and was overcome by antibody-dependent NK cell activation (ADNKA). Depletion of spike-specific antibodies confirmed their dominant role in virus neutralisation, but these antibodies played only a minor role in ADNKA compared to antibodies to other viral proteins, including ORF3a, Membrane, and Nucleocapsid. In contrast, ADNKA induced following vaccination was focussed solely on spike, was weaker than ADNKA following natural infection, and was not boosted by the second dose. These insights have important implications for understanding disease progression, vaccine efficacy, and vaccine design.

Introduction

Both innate and adaptive immunity are critical for control of viral infection. As a result, the capacity for many pathogens to cause disease is dependent on their ability to antagonise immune pathways, with the most successful pathogens carrying a vast array of genes capable of inhibiting all arms of the immune system^{1,2}. In the case of SARS-CoV2, the virus antagonises interferon induction via the activity of at least 10 genes³⁻⁶. However, whether SARS-CoV2, or indeed any coronavirus, is capable of counteracting other arms of immunity is currently unknown.

Many successful viruses interfere with cellular immunity; for example, downregulation of MHC-I inhibits CD8+ T-cell mediated cytotoxicity². The cellular immune system counters this through Natural Killer (NK) cells, which display a number of dominant inhibitory receptors to MHC-I, and are therefore more likely to activate in its absence. However, NK cell activation also requires activation signals from a wide range of activating receptors⁷. The ligands for these can be induced on the surface of target cells in response to stress, infection, or transformation. For example, the stress ligands MICA, MICB, and ULBP1-6, all bind to the ubiquitously expressed activating receptor NKG2D. To limit NK cell activation, some viruses have therefore evolved to also manipulate levels of these NK activating ligands, reducing NK-cell mediated control¹. Whether similar manipulations underlie the ability of SARS-CoV2 to establish infection and cause disease in humans has not been investigated to date.

NK cells intrinsically function early in infection, limiting pathogen spread prior to the induction of adaptive immunity, but can also operate within the adaptive phase via antibody dependent NK activation (ADNKA) which leads to antibody dependent cellular cytotoxicity (ADCC). Here, antibodies bind to viral antigens expressed on the infected cell surface, and activate NK cells through binding of Fc to the activating receptor CD16 on the NK cell. This leads to release of cytotoxic granules, and the production of TNF α and IFN γ , contributing significantly to a pro-inflammatory environment. Multiple lines of evidence indicate that ADCC is an important function in COVID-19; ADCC is strongly induced following infection with SARS-CoV2⁸⁻¹¹, and delayed induction of ADCC correlates with severe disease¹². Furthermore, Fc mediated functions correlate with protection in Rhesus models of vaccination¹³, and are induced by vaccination in humans^{14,15}. Many neutralising antibodies targeting the SARS-COV2 spike

protein also demonstrate an ability to activate ADCC¹⁶⁻¹⁸, and the ability of these antibodies to control SARS-CoV2 infection in animal models is enhanced by Fc-dependent mechanisms of action¹⁹⁻²². Finally, ‘adaptive’ NK cells, marked by expression of NKG2C, CD57, and LIR-1²³, are exceptionally potent ADCC-mediators, and expansion of these cells has been observed in severe COVID-19²⁴, while genetic loss of NKG2C is a risk factor for severe disease²⁵.

Taken together, ADCC is likely to play an important role in COVID-19. However significant limitations remain in our understanding of this process. Most importantly, studies to date have focussed on spike protein as the dominant ADCC target, using either plate-bound protein or transfected cells^{8-12,14-18}. While viral entry glycoproteins are often found on the infected cell surface and can mediate ADCC when expressed alone, other viral proteins may be the principle mediators of ADCC during infection. Indeed, we have previously shown, in the context of other viruses, that non-structural proteins mediate more potent ADCC responses than entry glycoproteins²⁶. It is therefore important to evaluate responses in the context of productive viral infection.

To address these critical gaps in our understanding, we used a proteomic approach to quantitate the changes in cell surface protein expression of a lung epithelial cell line, that is permissive to efficient SARS-CoV2 infection²⁷. We found that SARS-CoV2-infected cells show downregulation of multiple activating NK ligands, resulting in a reduced ability of NK cells to respond to infected cells, with no effect on cell surface MHC-I. Following natural infection, NK inhibition was overcome through ADNKA. However this was not primarily driven by spike antibody, but dominated by antibody to other viral proteins such as nucleocapsid, membrane and ORF3a. Critically, spike was not only a poor ADNKA target following natural infection, but was also a poor activator of ADNKA following vaccination. Thus, the breadth of ADNKA responses is limited in current vaccination strategies. The addition of antigens such as nucleocapsid would recruit additional effector mechanisms, and may reduce reduction in vaccine efficacy following mutation of spike in naturally circulating virus variants.

Results

SARS-CoV-2 remodels the Plasma Membrane Proteome

Immune cells recognise and interact with infected cells through cell surface ligands. We therefore wanted to gain a comprehensive and unbiased overview of how SARS-CoV-2 infection changes the plasma membrane protein landscape. We performed plasma membrane enrichment through selective aminooxy-biotinylation (Plasma Membrane Profiling; PMP), a methodology that we previously developed²⁸ and have used to interrogate infection by HIV²⁹, HCMV^{26,30-33}, and HSV³⁴.

High infection rates are critical to minimise confounding effects from bystander cells. We therefore applied PMP to SARS-CoV-2 infected A549-ACE2-TMPRSS2 (AAT) cells, which are highly permissive to SARS-CoV-2²⁷ and infection at a high multiplicity of infection (MOI) results in over 90% infection by 24h (Fig. 1A). We utilised TMT based quantification to compare PM protein abundance in uninfected cells, and cells infected for 12, 24, and 48 h (Fig 1B). In total, 4953 proteins were quantitated, including 2227 proteins with gene ontology annotation associated with the cell surface and plasma membrane (Fig S1A). These proteins made up nearly 80% of the total protein abundance, indicating a high degree of enrichment for plasma membrane proteins, comparable with our previous PMP datasets^{28,29}.

Five SARS-CoV-2 viral proteins were detected, including spike, ORF3A, and membrane, which have previously been annotated as having a cell surface localisation (Fig. 1C). Consistent with prior reports, the viral receptor ACE2 was strongly downregulated after infection³⁵, while TMPRSS2 expression was unaffected (Fig. 1C). Overall, of the 2227 PM proteins quantified with more than one peptide, 914 showed changes which were both significant and with a greater than 1.5-fold over the course of the experiment. Clustering of proteins by their temporal pattern of change has previously shown utility in defining classes of regulated proteins and predicting underlying mechanisms^{29,36,37}. Here, we used K-means clustering to define 5 groups of significantly altered proteins by their temporal profile.

To assess the innate immune response triggered by SARS-CoV-2 infection, the clusters were interrogated for the presence of a predefined list of interferon alpha-inducible genes. Thirteen of the 23 detected PM genes associated with the interferon response fell into cluster

5 (Fig S1B), which describes proteins with an increase in expression at 48 h, examples shown in Fig S1C. As the SARS-CoV-2 viral replication cycle is completed in less than 24 h⁶, SARS-CoV-2 is able to evade innate immune recognition and/or suppress IFN signalling until long after virus egress has been achieved.

To identify potential innate and adaptive immune evasion strategies employed by SARS-CoV-2, we focused on cluster 1, which describes proteins downregulated from early in the time course. This cluster was subject to gene ontology and pathway analysis to define proteins related by functional and structural similarities (Fig 2). Several classifications enriched in this cluster are particularly relevant, including the downregulation of several cytokine receptors, including IFNAR1. Much of the SARS-CoV-2 genome is dedicated to antagonising a type I interferon response⁶ and the plasma membrane downregulation of IFNAR1 presents a previously undescribed strategy to prevent exogenous signalling.

Also of note is the downregulation of heparan sulfate proteoglycans, the syndecan (SDC), glypican (GPC) and neurophilin (NRP) proteins, which together encompass the families of heparan sulfate modified proteins expressed on epithelial cells³⁸. NRP1 is a cell entry factor for SARS-CoV-2^{39,40}, and SARS-CoV-2 spike is also reported to bind heparan sulphated proteins⁴¹. The decrease in cell surface expression of heparan sulfated proteins may therefore enable the egress of viral particles and prevent super-infection, and is reminiscent of the capacity of other viruses to downregulate both primary viral receptors and co-receptors⁴².

SARS-CoV2 modulates multiple NK cell ligands, and inhibits NK cell activation

Our experience with PMP has emphasised the modulation of NK ligands as an important immune evasion mechanism^{30,37}. As these ligands are poorly annotated in public databases, we investigated the PMP dataset for previously described ligands⁷, of which 18 were detected (Fig S2). MHC-I proteins HLA-B to HLA-E were upregulated in cluster 5, likely reflecting a late IFN-mediated upregulation, while HLA-A was unchanged. By contrast, the activating NK-cell ligands MICA, ULBP2, Nectin1, and B7-H6 (NCR3LG1) were significantly downregulated in cluster 1, and was confirmed by flow cytometry (Fig 3A). To determine how these changes affected NK cell activation, we incubated SARS-CoV2 infected cells with PBMC, and measured (i) degranulation of CD3⁺CD56⁺ NK cells (Fig 3B), (ii) production of IFN γ (Fig 3C) and (iii) TNF α (Fig 3D). Consistent with data using other viruses²⁶, CD107a was activated on a higher

proportion of cells than IFN γ and TNF α . Nevertheless, all three markers of NK activation were significantly reduced in SARS-CoV2 infected versus mock infected cells, suggesting that the virus has evolved to inhibit NK cell responses. Interestingly, all 4 ligands have orthologues in bats (the likely reservoir species for most coronaviruses), with one ligand (Nectin1) showing extremely high identity (>96%) to the human gene (Table 1). Thus, control of NK activation may display similarities between host species, with some NK-evasins functioning across species barriers. This is consistent with reports of SARS-CoV2 being a ‘generalist’ virus that did not need to undergo extensive evolution to be capable of human-to-human transmission⁴³.

Gene	Homo sapiens ID	Rhinolophus ferrumequinum ID	% Identity
B7-H6	XP_011518377	XP_032975102	52.19
Nectin 1	NP_002846	XP_032976402	96.12
MICA	NP_000238	XP_032955521	43.41
ULBP2	NP_079493	XP_032959100 (ULBP1)	42.26
		XP_032958641 (ULBP3)	44.86

Table 1. Homology of NK ligands from humans with those of the horseshoe bat

NK-cell evasion is overcome by antibody-dependent activation

In addition to their role in innate immunity through interactions with activating and inhibitory NK receptors, NK cells play additional roles in adaptive immunity via ADNKA/ADCC, in which CD16 interacts with the Fc portion of antibodies bound to targets on the surface of infected cells. To analyse this aspect of NK function, PBMC were incubated with infected cells in the presence of sera from individuals that were seronegative or seropositive for SARS-CoV2. Degranulation (Fig 4A), as well as production of IFN γ (Fig 4B) and TNF α (Fig 4C), were all significantly increased in the presence of serum containing anti-SARS-CoV2 antibodies, across a range of different NK donors (Fig 4D-F). Viral immune evasion can therefore be overcome

through antibody dependent mechanisms, suggesting that in addition to neutralising virus, SARS-CoV2 antibodies may aid virus clearance through ADCC.

The magnitude of the neutralising antibody response to SARS-CoV2 correlates with severity of disease; individuals with more severe disease generate higher levels of neutralising antibody⁴⁴⁻⁴⁶. Although memory B-cell responses persist, levels of circulating antibody can decrease over time following resolution of disease, such that in some mildly infected people they become undetectable after a few months⁴⁴⁻⁴⁶. To determine whether the same was true of ADCC responses, we assessed ADNKA in a cohort of donors with known disease status, for whom longitudinal sera were available (Fig 4G). ADNKA responses were generally of greater magnitude in individuals with more severe disease, with responses remaining detectable during the 150-200 day follow-up. Amongst those with milder disease, strong initial responses were maintained over time, while weaker responses decreased, becoming virtually undetectable by around 200 days. There was a significant correlation between the ability of antibodies to neutralise SARS-CoV2, and to activate ADNKA in response to SARS-CoV2 (Fig 4H), with a R^2 value of 0.7. Thus, although the antibodies mediating these activities may be induced in a similar manner, the antibodies mediating the two responses may not be identical.

Multiple SARS-CoV2 ORFs mediate ADCC

There has been a major focus on spike as an activator of ADCC^{8-12,14-18}. It is certainly a potential target during infection as both our PMP and flow cytometric analysis detected substantial levels of spike protein on the infected cell surface (Fig 5A). However, our studies with other viruses have demonstrated that although viral proteins involved in virion binding and entry are found on the cell surface, and can be bound by antibody, they are not necessarily the strongest mediators of ADCC activity²⁶. PMP also detected Membrane (M), Nucleocapsid (N) and ORF3a on the cell surface, while bioinformatic analysis of the entire viral genome revealed a further five viral proteins that contained potential transmembrane domains or signal peptides. We therefore screened cells transfected with each of these genes for their ability to activate NK cells in an antibody dependent manner. We used sera from patients with severe covid19 disease, that had previously generated the strongest ADNKA signals against infected cells (Fig 5B). As expected, spike primed a substantial ADNKA response. However, in

support of our PMP data, expression of ORF3a, Membrane, and Nucleocapsid, were also capable of priming ADNKA in the presence of serum containing SARS-CoV2 antibodies.

Interestingly, nucleocapsid readily activated ADNKA despite the extremely low levels of this protein on the surface of transfected cells. We have previously observed the exquisite sensitivity of ADNKA, which can be activated in response to levels of virus protein that are difficult to detect by flow cytometry²⁶. We have also shown that antigen levels can be very different in the context of virus infection, compared with transfection²⁶. In support of this, nucleocapsid levels were much higher on the surface of infected cells (Fig 5C). Given how well even very low levels of nucleocapsid activated ADNKA, this suggested that multiple proteins other than spike could be major contributors to ADCC during SARS-CoV2 infection.

Monoclonal anti-spike antibodies only weakly mediate ADNKA despite binding strongly to infected cells

To determine the relative contribution of spike, and of different antigenic sites on Spike, to ADNKA during SARS-CoV2 infection, we tested a panel of 26 human anti-spike monoclonal antibodies, which had been isolated from naturally infected donors, and cloned as IgG1 constructs⁴⁷. The panel included antibodies targeting the RBD, S1, NTD, and S2 domains, neutralising and non-neutralising, and those targeting multiple distinct epitopes (Table S3).

All 26 antibodies bound efficiently to infected cells, at levels comparable to polyclonal serum (Fig S3). However, only five of these antibodies were capable of triggering degranulation (Fig 6A), of which two triggered TNF α (Fig 6B), and one IFN γ (Fig 6C). Importantly, these NK responses were at significantly lower levels than seen with polyclonal serum from a naturally infected donor. This suggested that, despite high levels of antibody binding, and a substantial ability to activate ADCC when expressed in isolation, spike may be a relatively poor ADCC target in the context of natural infection.

Following natural infection, ADNKA responses are dominated by non-spike antibodies

Although our data using monoclonal antibodies indicated that spike was a comparatively weak activator of ADNKA, the antibodies that mediate ADNKA may represent a minor proportion of the overall anti-spike response; we simply could not have captured those antibodies in our panel. To directly address this concern, we specifically depleted anti-spike

antibodies from the serum of naturally infected individuals, and assessed changes in activity. ELISA analysis confirmed the effective depletion of spike-specific antibodies (Fig 7A), which was accompanied by a significant loss of neutralisation activity (Fig 7B), across multiple donors (Fig 7C). However, there was no loss of ADNKA activity from any of these samples following spike-antibody depletion (Fig 7D, E), consistent with the results of our monoclonal antibody data. We conclude that spike-specific antibodies dominate the neutralising antibody response, but play a minor roles in the ADCC response following natural infection.

Spike-specific antibodies following vaccination are weak mediators of ADNKA

Since non-spike antigens must prime ADCC following natural infection, we investigated individuals who have been vaccinated against spike, but not exposed to the virus - since they will only have spike-specific antibodies. As with serum from naturally infected individuals, ELISA demonstrated effective depletion of spike antibodies (Fig 7F), which was accompanied by a marked loss of virus neutralisation activity (Fig 7G, H). In functional NK assays, these sera were capable of mediating relatively weak ADNKA (Fig 7I, J) and, unlike sera from naturally infected individuals, ADNKA activity was abrogated following depletion of spike antibodies. When we compared responses following natural infection or vaccination, vaccine mediated ADNKA was significantly weaker than responses seen following natural infection, and was not boosted following the second dose of the vaccine (Fig 7K). This was in stark contrast with the neutralisation activity of those same serum samples, which were dramatically increased following the second dose, with activity comparable to patients with severe covid-19 disease (Fig 7L).

Thus, anti-spike antibodies can mediate ADNKA, but this activity appears weaker than responses seen during natural infection, which targets a broader repertoire of viral antigens.

Discussion

The host innate immune system plays a critical role in both early SARS-CoV2 infection, and later COVID-19 disease. This is emphasised by the identification of multiple interferon antagonists encoded by SARS-CoV2, and the increased disease severity in patients with genetic and acquired interferon deficiencies³⁻⁶. To determine whether the virus evades other components of the host immune system we took a proteomic approach to systematically analyse how SARS-CoV2 infection manipulates the cell surface during the time course of SARS-CoV2 infection. We find that SARS-CoV-2 remodelling of the plasma membrane leads to the downregulation of multiple cell surface immune ligands. This likely enables SARS-CoV2 to evade different effector arms of the host immune system. Downregulation of NK activating ligands inhibits NK cell activation early in infection, but this effect is countered later with the production of antibody and subsequent ADCC. Somewhat surprisingly, we find that spike antibody is not responsible for the robust ADNKA which develops after natural infection, and only weak ADNKA is generated following spike vaccination.

Genetic variation in both viruses and their hosts affects the ability of viruses to modulate NK responses, and influence pathogenesis^{1,48}. SARS-CoV2 appears to have evolved to antagonise NK activation by targeting the expression levels of multiple activating NK ligands. All four of the NK ligands targeted by the virus are also found in bat species, thought to be the reservoir for most coronaviruses, implying an element of cross-species functionality to viral immune-evasins, which may be important for zoonotic transmission. Indeed, the 96% sequence identity between the nectin-1/CD111 bat and human genes is striking. Nectin-1 is a ligand for CD96/TACTILE⁴⁹, which activates human NK cells⁵⁰. SARS-CoV-2 also targets at least 3 additional NK activating ligands (B7-H6, ULBP2 and MICA). Of these, MICA is highly polymorphic (100+ alleles), with A549 cells expressing MICA*001/004⁵¹, while the truncated MICA*008 is the most common allele worldwide. Further work will be needed to determine whether MICA genetic diversity affects the extremely variable outcome of SARS-CoV-2 infection.

In contrast to the effect on NK ligands, our data imply that the virus cannot actively antagonise adaptive cellular immunity. Our evidence for this includes the lack of significant MHC-I cell surface downregulation, the emergence of ADNKA with humoral immunity, and the inability

of infected cells to bind human IgG from seronegative donors (i.e. the virus does not encode Fc receptors as ADNKA decoys). This is consistent with SARS-CoV2 acting as an acute 'hit and run' virus, which has replicated and transmitted before host adaptive immunity develops. It will be interesting to determine whether coronaviruses evade innate and adaptive immunity in their original bat host, and whether this contributes to their persistence in that species.

Animal models show that in both vaccination¹³, and monoclonal antibody administration¹⁹⁻²², Fc-mediated activation of cellular immunity plays an important protective role. However, inflammation dominates severe COVID-19 disease^{52,53}. The high levels of TNF α and IFN γ observed following ADNKA are likely to exacerbate disease under these circumstances. In support of this, TNF α and IFN γ correlate with severe disease⁵⁴, and critically ill patients have a higher proportion of afucosylated antibodies that promote a stronger ADCC response. Given that ADCC can potentially contribute to both protection and immunopathology, it is important to understand what determines the magnitude and quality of the ADCC response. Viral entry glycoproteins are a common focus as mediators of ADCC. E.g. ADCC mediated by Env antibodies have been a major focus in HIV⁵⁵, and gB-specific antibodies have been studied in HCMV⁵⁶. Although antibodies targeting these antigens can clearly bind infected cells, using similar approaches to the present study we recently showed that non-structural accessory proteins were more potent ADCC targets during HCMV infection²⁶. This conclusion is reinforced here, where the neutralising antibody response is dominated by spike, but spike antibodies are poor activators of ADNKA, with other viral antigens driving efficient ADNKA following natural infection. Nucleocapsid, ORF3a, and membrane are all present on the infected cell surface, and can promote ADNKA if appropriate antibodies are present. However, antibodies targeting ORF3a and membrane are only detected in a small subset of patients, while almost everyone infected with SARS-CoV2 develops antibodies targeting⁵⁷. Nucleocapsid is therefore likely to be the major target for ADCC during natural SARS-CoV2 infection in most people. Influenza nucleoprotein is also detected at the plasma membrane⁵⁸⁻⁶⁰, and antibodies targeting nucleoprotein can mediate ADCC^{61,62}. Structural proteins involved in genome packaging may therefore represent a common target for ADCC across multiple virus families.

The identification of the antigens that drive ADCC is important to our understanding of the mechanisms that underlie pathogenesis and disease. A skewing of the spike:nucleocapsid antibody ratio is associated with severe disease⁶³⁻⁶⁵, where ADCC-mediating nucleocapsid antibodies might be promoting inflammation. Both the breadth of the nucleocapsid antibody response, and the particular epitopes targeted, have also been correlated with disease severity^{57,66}. These correlations now need to be followed up functionally, to determine how the epitopes bound by anti-nucleocapsid antibodies affect ADCC. It will also be important to determine the contribution of these ADCC-capable antibodies to inflammation and myeloid cell activation⁶⁷⁻⁶⁹, via antibody-dependent cellular phagocytosis (ADCP) and complement dependent cytotoxicity (CDC).

Although Fc-dependent mechanisms of cellular immune activation could contribute to immunopathology via multiple mechanisms in severe disease, all patients with mild/asymptomatic disease also developed strong and persistent ADNKA responses. Furthermore, lineage defining mutations in the viral variants of concern B.1.1.7, B.1.351, and P.1, all include non-synonymous mutations in nucleocapsid, which is one of the most variable genes amongst circulating strains⁷⁰, and these nucleocapsid mutations can result in reduced antibody binding⁷¹. Virus variants may therefore evolve to evade ADCC responses as well as neutralising responses, supporting an important role for ADCC in protection. It is therefore significant that spike-vaccine generated antibody responses were poor ADNKA inducers. The addition of other viral proteins such as nucleocapsid to vaccines would engage a wider range of immune effector pathways. This approach might improve efficacy against both viral transmission and disease, resulting in vaccines that are more resistant to viral variants containing mutations that diminish antibody neutralisation⁷².

Acknowledgements

This work was supported in part by the UK Coronavirus Immunology Consortium (CiC). RS was supported by the MRC (MR/S00971X/1), and Wellcome Trust (204870/Z/16/Z), PJL was funded by the Wellcome Trust through a Principal Research Fellowship [210688/Z/18/Z], the MRC (MR/V011561/1), the Addenbrooke's Charitable Trust and the NIHR Cambridge Biomedical Research Centre. KLD was supported by King's Together Rapid COVID-19 Call, Huo Family Foundation Award, and a Wellcome Trust Multi-User Equipment Grant 208354/Z/17/Z. CG was supported by the MRC-KCL Doctoral Training Partnership in Biomedical Sciences (MR/N013700/1). BM was supported by an NIHR Academic Clinical Fellowship in Combined Infection Training

Materials and methods

Cells and viruses

A549 were transduced with lentiviruses expressing human ACE2, and TMPRSS2 (AAT cells), as previously described²⁷. The England2 strain of SARS-CoV2 was obtained from Public Health England (PHE), grown on VeroE6 cells, and titrated by plaque assay on both VeroE6 and AAT, as previously described²⁷. All cell lines were grown in DMEM containing 10% FCS (Gibco), at 37°C and in 5% CO₂. For assays, cells were plated out the day before, then infected at a multiplicity of infection (MOI) of 5, for 1h on a rocking platform. The inoculum was removed, and fresh DMEM containing 2% FCS was added, then cells were incubated until the assay.

Plasma membrane profiling

Cell surface proteins were labelled essentially as described⁷³. Briefly, cells were incubated in a solution containing sodium periodate, aniline and aminoxy biotin to label predominantly sialic acid containing glycans at the cell surface. Cell lysates were then enriched for labelled proteins using streptavidin-agarose beads. After extensive washing trypsin was added to liberate peptides of the enriched proteins. The resulting peptide pools were dried prior to labelling with TMT reagents.

TMT Labelling and clean-up

Samples were resuspended in 21µL 100mM TEAB pH 8.5. After allowing to come to room temperature, 0.2mg TMT reagents (Thermo Fisher) were resuspended in 9µL anhydrous ACN which was added to the respective samples and incubated at room temperature for 1h. A 3µL aliquot of each sample was taken and pooled to check TMT labelling efficiency and equality of loading by LC-MS. After checking each sample was at least 98% TMT labelled total reporter ion intensities were used to normalise the pooling of the remaining samples such that the final pool was as close to a 1:1 ratio of total peptide content between samples as possible. This final pool was then dried in a vacuum centrifuge to evaporate the majority of ACN from labelling. The sample was acidified to a final 0.1% Trifluoroacetic Acid (TFA) (~200µL volume) and FA was added until the SDC visibly precipitated. 4 volumes of ethyl acetate were then added and the sample vortexed vigorously for 10s. Sample was then centrifuged at 15'000g for 5mins at RT to effect phase separation. A gel loading pipette tip was used to withdraw the lower (aqueous) phase to a fresh low adhesion microfuge tube. The sample was then partially

dried in a vacuum centrifuge and brought up to a final volume of 1mL with 0.1% TFA. FA was added until the pH was <2, confirmed by spotting onto pH paper. The sample was then cleaned up by SPE using a 50mg tC18 SepPak cartridge (Waters). The cartridge was wetted with 1mL 100% Methanol followed by 1mL ACN, equilibrated with 1mL 0.1% TFA and the sample loaded slowly. The sample was passed twice over the cartridge. The cartridge was washed 3x with 1mL 0.1% TFA before eluting sequentially with 250µL 40% ACN, 70% ACN and 80% ACN and dried in a vacuum centrifuge.

Basic pH reversed phase fractionation

TMT labelled samples were resuspended in 40µL 200mM Ammonium formate pH10 and transferred to a glass HPLC vial. BpH-RP fractionation was conducted on an Ultimate 3000 UHPLC system (Thermo Scientific) equipped with a 2.1 mm × 15 cm, 1.7µ Kinetex EVO column (Phenomenex). Solvent A was 3% ACN, Solvent B was 100% ACN, solvent C was 200 mM ammonium formate (pH 10). Throughout the analysis solvent C was kept at a constant 10%. The flow rate was 500 µL/min and UV was monitored at 280 nm. Samples were loaded in 90% A for 10 min before a gradient elution of 0–10% B over 10 min (curve 3), 10–34% B over 21 min (curve 5), 34–50% B over 5 mins (curve 5) followed by a 10 min wash with 90% B. 15s (100µL) fractions were collected throughout the run. Fractions containing peptide (as determined by A280) were recombined across the gradient to preserve orthogonality with on-line low pH RP separation. For example, fractions 1, 25, 49, 73, 97 are combined and dried in a vacuum centrifuge and stored at -20°C until LC-MS analysis. 24 Fractions were generated in this manner.

Mass Spectrometry

Samples were analysed on an Orbitrap Fusion instrument on-line with an Ultimate 3000 RSLC nano UHPLC system (Thermo Fisher). Samples were resuspended in 10µL 5% DMSO/1% TFA and all sample was injected. Trapping solvent was 0.1% TFA, analytical solvent A was 0.1% FA, solvent B was ACN with 0.1% FA. Samples were loaded onto a trapping column (300µm x 5mm PepMap cartridge trap (Thermo Fisher)) at 10µL/min for 5 minutes at 60 degrees. Samples were then separated on a 75cm x 75µm i.d. 2µm particle size PepMap C18 column (Thermo Fisher) at 55 degrees. The gradient was 3–10% B over 10mins, 10–35% B over 155 minutes, 35–45% B over 9 minutes followed by a wash at 95% B for 5minutes and equilibration at 3% B.

Eluted peptides were introduced by electrospray to the MS by applying 2.1kV to a stainless-steel emitter (5cm x 30µm (PepSep)). During the gradient elution, mass spectra were acquired with the parameters detailed in Fig S4 using Tune v3.3 and Xcalibur v4.3 (Thermo Fisher).

Data Processing

Data were processed with PeaksX+, v10.5 (Bioinformatic Solutions). Processing parameters are shown in detail in Supplementary Fig S5. Briefly, .raw files were searched iteratively in three rounds, with unmatched *de novo* spectra (at 1% PSM FDR) from the previous search used as the input for the next. The three iterations were as follows 1) Swissprot Human (27/03/2020) + common contaminants 2) The same databases as search 1 but permitting semi-specific cleavage 3) trEMBL Human (27/03/2020), with specific cleavage rules. Proteins were then quantified using the parameters outlined in Fig S5. Identified proteins and their abundances were output to .csv format and further subjected to statistical analysis. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository⁷⁴ with the dataset identifier PXD025000 and 10.6019/PXD025000. Table S1 contains the analysed data.

Statistical analysis and K-means clustering.

Prior to statistical analysis, proteins were filtered for those quantified across all TMT reporter channels and with more than 1 unique. Only proteins with a plasma membrane related GO:CC annotation as previously described²⁸ were then carried forward for statistical analysis. All SARS-CoV-2 proteins quantified with more than one unique peptide were also included. Statistical tests were performed with the aov and p.adjust functions in the R base stats package (version 4.0.3) to calculate Benjamini-Hochberg corrected p-values for changes in protein abundance across the measured time-points.

Proteins with a p-value < 0.05 and a maximum fold-change across the time-course of > 1.5-fold were selected for k-means clustering. Mean intensities of each time-point were scaled using the scale function in base R and utilised to cluster proteins into 5 groups using the kmeans function in R base stats package.

Functional Annotation Clustering

Assessment of enriched gene annotation terms in temporal cluster one was carried out using the Functional Annotation Clustering tool at DAVID (david.ncifcrf.gov) v6.8⁷⁵, using the default clustering settings for medium stringency and the following libraries: Uniprot UP_Keyword, GOTERM:MF_ALL, BIOCARTA,, KEGG_PATHWAY and REACTOME_PATHWAY. Output clusters were curated with representative names based on the enriched terms within. The background for enrichment was a list of proteins detected in the PMP dataset with more than one peptides and a plasma membrane associated GO:CC term. Functional annotation clusters with an enrichment score of >1.5 are shown, and any clusters in which no individual annotation term had a Benjamini-Hochberg correct p-value of <0.05 were excluded.

Flow cytometry

Cells were dissociated using TrypLE, then stained with primary antibody for 30 min at 4°C. Following washing, they were incubated with secondary antibody, again for 30 min at 4°C. Cells were washed, fixed in 4% paraformaldehyde, and data collected on an Attune cytometer (Thermofisher). Antibodies used were against HLA-ABC (W632; AbD Serotec), NCR3LG1/B7-H6 (MAB7144, Biotechne R&D Systems), Nectin 1 (R1.302; Biolegend), MICA (AMO1-100; BAMOMAB), MICB (BMO2-100; BAMOMAB), ULBP2 (BUMO1; BAMOMAB), Spike (1A9; Insight), Nucleocapsid (1C7; Stratech), anti-mouse IgG AF647 (Thermofisher).

Plasmids and transfections

Lentivirus plasmids encoding each SARS-CoV2 ORF individually, with a C-terminal twin-strep tag, were obtained from Addgene⁷⁶. Plasmids were midiprep (Nucleobond Xtra Midi; Machery-Nagel), and transfected into 293T cells using GeneJuice (Merck) according to manufacturers' instructions.

NK activation assays

PBMCs from healthy donors were thawed from liquid N2 storage, rested overnight in RPMI supplemented with 10% FCS, and L-glutamine (2 mM) in the absence or presence of IFN- α (1,000 U/ml). Target cells were harvested using TrypLE Express (Gibco), preincubated for 30 min with the relevant antibody or serum preparations, then mixed with PBMCs at an effector:target (E:T) ratio of 10:1 in the presence of GolgiStop (0.7 μ l/ml, BD Biosciences), Brefeldin-A (1:1000, Biolegend) and anti-CD107a-FITC (clone H4A3, BioLegend). Cells were

incubated for 5 h, washed in cold PBS, and stained with live/dead Fixable Aqua (Thermo Fisher Scientific), anti-CD3–PECy7 (clone UCHT1, BioLegend) or anti-CD3-BV711 (Clone SK7, BioLegend), anti-CD56–BV605 (clone 5.1H11, BioLegend), anti-CD57–APC (clone HNK-1, BioLegend) or anti-CD57-PECy7 (clone HNK-1, biolegend), and anti-NKG2C–PE (clone 134591, R&D Systems). In some experiments, cells were also fixed/permeabilized using Cytofix/Cytoperm (BD Biosciences) and stained with anti-TNF α –BV421 (clone MAb11, BioLegend) and anti-IFN γ –APC (clone 4S.B3, BioLegend). Data were acquired using an AttuneNxT (Thermo Fisher) and analyzed with Attune NxT software or FlowJo software version 10 (Tree Star). To ensure inter-assay variation did not affect results, a donor serum demonstrating moderate ADCC activity against SARS-CoV2 was included as a positive standard in every assay, while a serum collected before 2020 was included as a negative control in every assay. Where sera were tested at a range of dilutions, the area under the curve (AUC) was calculated using Graphpad Prism 9. This value was then normalised to the AUC for the standard serum in that particular assay.

Serums

Serums were collected from healthy donors vaccinated with the BNT162b2 vaccine, a minimum of 3 weeks after the first dose, or 1 week after the second. A pre-vaccine sample was taken from every donor, and an ELISA for SARS-CoV2 RBD performed as previously described⁷⁷, to determine pre-exposure to live SARS-CoV2. Longitudinal serums from naturally infected individuals have been described previously, clinical characteristics are in Table S2⁴⁴. Those labelled ‘mild’ had severity scores of 0, 1, 2 or 3, and ‘severe’ was 4 or 5.

Virus neutralisation assays

600PFU of SARS-CoV2 was incubated with appropriate dilutions of serum, for 1h, at 37°C. The mixes were then added to pre-plated VeroE6 cells for 48h. After this time, monolayers were fixed with 4% PFA, permeabilised for 15 min with 0.5% NP-40, then blocked for 1h in PBS containing 0.1% Tween (PBST) and 3% non-fat milk. Primary antibody (anti-nucleocapsid 1C7, Stratech, 1:500 dilution) was added in PBST containing 1% non-fast milk and incubated for 1h at room temperature. After washing in PBST, secondary antibody (anti-mouse IgG-HRP, Pierce, 1:3,000 dilution) was added in PBST containing 1% non-fat milk and incubated for 1h. Monolayers were washed again, developed using Sigmafast OPD according to manufacturers’

instructions, and read on a Clariostar Omega plate reader. Wells containing no virus, virus but no antibody, and a standardised serum displaying moderate activity were included as controls in every experiment. NT50 were calculated in Graphpad Prism 9.

Antibody depletions

Anti-spike antibody was depleted from sera using magnetic bead conjugated spike trimer protein (Acrobiosystems). Beads were resuspended in PBS+0.05% BSA, then 50µl serum was mixed with 150µl beads. Mixtures were incubated on a rotating mixer at 4°C overnight. Serum diluted in buffer alone was used as a control. Magnetic beads were then removed using a 3D printed magnetic stand. All values given in assays are corrected for this initial 4-fold dilution. Levels of anti-spike antibody were measured using a spike trimer ELISA (Acrobiosystems).

Ethics

PBMC were extracted from apheresis cones obtained from the Welsh Blood Service (WBS) via an ad-hoc agreement or from blood samples from healthy volunteers and stored in liquid N₂ until use. Use of healthy volunteer PBMC for this project, including those from WBS, was ethically approved by the Cardiff University School of Medicine Research Ethics Committee (SMREC) nos. 20/55 and 20/101. Recruitment of healthy volunteers after vaccination was covered by the Cardiff University School of Medicine Research Ethics Committee under reference no. 18/04.

References

- 1 Patel, M. *et al.* HCMV-Encoded NK Modulators: Lessons From in vitro and in vivo Genetic Variation. *Frontiers in immunology* **9**, 2214, (2018).
- 2 Berry, R., Watson, G. M., Jonjic, S., Degli-Esposti, M. A. & Rossjohn, J. Modulation of innate and adaptive immunity by cytomegaloviruses. *Nat Rev Immunol* **20**, 113-127, (2020).
- 3 Sa Ribero, M., Jouvenet, N., Dreux, M. & Nisole, S. Interplay between SARS-CoV-2 and the type I interferon response. *PLoS Pathog* **16**, e1008737, (2020).
- 4 Miorin, L. *et al.* SARS-CoV-2 Orf6 hijacks Nup98 to block STAT nuclear import and antagonize interferon signaling. *Proc Natl Acad Sci U S A* **117**, 28344-28354, (2020).
- 5 Schroeder, S. *et al.* Interferon antagonism by SARS-CoV-2: a functional study using reverse genetics. *The Lancet Microbe*, (2021).
- 6 Lei, X. *et al.* Activation and evasion of type I interferon responses by SARS-CoV-2. *Nature communications* **11**, 3810, (2020).
- 7 Kumar, S. Natural killer cell cytotoxicity and its regulation by inhibitory receptors. *Immunology* **154**, 383-393, (2018).
- 8 Dufloo, J. *et al.* Asymptomatic and symptomatic SARS-CoV-2 infections elicit polyfunctional antibodies. *medRxiv*, 2020.2011.2012.20230508, (2020).
- 9 Herman, J. D. *et al.* Functional Antibodies in COVID-19 Convalescent Plasma. *medRxiv*, 2021.2003.2008.21253157, (2021).
- 10 Lee, W. S. *et al.* Decay of Fc-dependent antibody functions after mild to moderate COVID-19. *medRxiv*, 2020.2012.2013.20248143, (2020).
- 11 Anand, S. P. *et al.* Longitudinal analysis of humoral immunity against SARS-CoV-2 Spike in convalescent individuals up to 8 months post-symptom onset. *bioRxiv*, 2021.2001.2025.428097, (2021).
- 12 Zohar, T. *et al.* Compromised Humoral Functional Evolution Tracks with SARS-CoV-2 Mortality. *Cell* **183**, 1508-1519 e1512, (2020).
- 13 Gorman, M. J. *et al.* Collaboration between the Fab and Fc contribute to maximal protection against SARS-CoV-2 in nonhuman primates following NVX-CoV2373 subunit vaccine with Matrix-M vaccination. *bioRxiv*, 2021.2002.2005.429759, (2021).
- 14 Barrett, J. R. *et al.* Phase 1/2 trial of SARS-CoV-2 vaccine ChAdOx1 nCoV-19 with a booster dose induces multifunctional antibody responses. *Nat Med* **27**, 279-288, (2021).
- 15 Tauzin, A. *et al.* A single BNT162b2 mRNA dose elicits antibodies with Fc-mediated effector functions and boost pre-existing humoral and T cell responses. *bioRxiv*, 2021.2003.2018.435972, (2021).
- 16 Tortorici, M. A. *et al.* Ultrapotent human antibodies protect against SARS-CoV-2 challenge via multiple mechanisms. *Science* **370**, 950-957, (2020).
- 17 McCallum, M. *et al.* N-terminal domain antigenic mapping reveals a site of vulnerability for SARS-CoV-2. *bioRxiv*, 2021.2001.2014.426475, (2021).
- 18 Cathcart, A. L. *et al.* The dual function monoclonal antibodies VIR-7831 and VIR-7832 demonstrate potent in vitro and in vivo activity against SARS-CoV-2. *bioRxiv*, 2021.2003.2009.434607, (2021).
- 19 Chan, C. E. Z. *et al.* The Fc-mediated effector functions of a potent SARS-CoV-2 neutralizing antibody, SC31, isolated from an early convalescent COVID-19 patient, are essential for the optimal therapeutic efficacy of the antibody. *bioRxiv*, 2020.2010.2026.355107, (2020).
- 20 Schafer, A. *et al.* Antibody potency, effector function, and combinations in protection and therapy for SARS-CoV-2 infection in vivo. *J Exp Med* **218**, (2021).
- 21 Winkler, E. S. *et al.* Human neutralizing antibodies against SARS-CoV-2 require intact Fc effector functions and monocytes for optimal therapeutic protection. *bioRxiv*, 2020.2012.2028.424554, (2020).

- 22 Suryadevara, N. *et al.* Neutralizing and protective human monoclonal antibodies recognizing the N-terminal domain of the SARS-CoV-2 spike protein. *bioRxiv*, 2021.2001.2019.427324, (2021).
- 23 Hammer, Q. & Romagnani, C. OMIP-039: Detection and analysis of human adaptive NKG2C(+) natural killer cells. *Cytometry A* **91**, 997-1000, (2017).
- 24 Maucourant, C. *et al.* Natural killer cell immunotypes related to COVID-19 disease severity. *Sci Immunol* **5**, eabd6832, (2020).
- 25 Vietzen, H. *et al.* Deletion of the NKG2C receptor encoding KLRC2 gene and HLA-E variants are risk factors for severe COVID-19. *Genet Med*, (2021).
- 26 Vlahava, V. M. *et al.* Monoclonal antibodies targeting nonstructural viral antigens can activate ADCC against human cytomegalovirus. *J Clin Invest* **131**, (2021).
- 27 Rihn, S. J. *et al.* A plasmid DNA-launched SARS-CoV-2 reverse genetics system and coronavirus toolkit for COVID-19 research. *PLoS Biol* **19**, e3001091, (2021).
- 28 Weekes, M. P. *et al.* Comparative analysis of techniques to purify plasma membrane proteins. *J Biomol Tech* **21**, 108-115, (2010).
- 29 Matheson, N. J. *et al.* Cell Surface Proteomic Map of HIV Infection Reveals Antagonism of Amino Acid Metabolism by Vpu and Nef. *Cell host & microbe* **18**, 409-423, (2015).
- 30 Fielding, C. A. *et al.* Control of immune ligands by members of a cytomegalovirus gene expansion suppresses natural killer cell activation. *Elife* **6**, (2017).
- 31 Hsu, J. L. *et al.* Plasma membrane profiling defines an expanded class of cell surface proteins selectively targeted for degradation by HCMV US2 in cooperation with UL141. *PLoS Pathog* **11**, e1004811, (2015).
- 32 Weekes, M. P. *et al.* Quantitative temporal viromics: an approach to investigate host-pathogen interaction. *Cell* **157**, 1460-1472, (2014).
- 33 Weekes, M. P. *et al.* Latency-associated degradation of the MRP1 drug transporter during latent human cytomegalovirus infection. *Science* **340**, 199-202, (2013).
- 34 Soh, T. K. *et al.* Temporal Proteomic Analysis of Herpes Simplex Virus 1 Infection Reveals Cell-Surface Remodeling via pUL56-Mediated GOPC Degradation. *Cell reports* **33**, 108235, (2020).
- 35 Patra, T. *et al.* SARS-CoV-2 spike protein promotes IL-6 trans-signaling by activation of angiotensin II receptor signaling in epithelial cells. *PLoS Pathog* **16**, e1009128, (2020).
- 36 Greenwood, E. J. *et al.* Temporal proteomic analysis of HIV infection reveals remodelling of the host phosphoproteome by lentiviral Vif variants. *eLife* **5**, (2016).
- 37 Weekes, M. P. *et al.* Quantitative temporal viromics: an approach to investigate host-pathogen interaction. *Cell* **157**, 1460-1472, (2014).
- 38 Sarrazin, S., Lamanna, W. C. & Esko, J. D. Heparan sulfate proteoglycans. *Cold Spring Harb Perspect Biol* **3**, (2011).
- 39 Cantuti-Castelvetri, L. *et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity. *Science* **370**, 856-860, (2020).
- 40 Daly, J. L. *et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection. *Science* **370**, 861-865, (2020).
- 41 Clausen, T. M. *et al.* SARS-CoV-2 Infection Depends on Cellular Heparan Sulfate and ACE2. *Cell* **183**, 1043-1057 e1015, (2020).
- 42 Landi, A., Iannucci, V., Nuffel, A. V., Meuwissen, P. & Verhasselt, B. One protein to rule them all: modulation of cell surface receptors and molecules by HIV Nef. *Curr HIV Res* **9**, 496-504, (2011).
- 43 MacLean, O. A. *et al.* Natural selection in the evolution of SARS-CoV-2 in bats created a generalist virus and highly capable human pathogen. *PLoS Biol* **19**, e3001115, (2021).
- 44 Seow, J. *et al.* Longitudinal observation and decline of neutralizing antibody responses in the three months following SARS-CoV-2 infection in humans. *Nat Microbiol* **5**, 1598-1607, (2020).
- 45 Dan, J. M. *et al.* Immunological memory to SARS-CoV-2 assessed for up to 8 months after infection. *Science* **371**, eabf4063, (2021).

- 46 Wajnberg, A. *et al.* SARS-CoV-2 infection induces robust, neutralizing antibody responses that are stable for at least three months. *medRxiv*, 2020.2007.2014.20151126, (2020).
- 47 Graham, C. *et al.* Impact of the B.1.1.7 variant on neutralizing monoclonal antibodies recognizing diverse epitopes on SARS-CoV-2 Spike. *bioRxiv*, 2021.2002.2003.429355, (2021).
- 48 Vietzen, H. *et al.* Extent of Cytomegalovirus Replication in the Human Host Depends on Variations of the HLA-E/UL40 Axis. *MBio* **12**, (2021).
- 49 Seth, S. *et al.* The murine pan T cell marker CD96 is an adhesion receptor for CD155 and nectin-1. *Biochemical and Biophysical Research Communications* **364**, 959-965, (2007).
- 50 Holmes, V. M. *et al.* Interaction between nectin-1 and the human natural killer cell receptor CD96. *PLoS One* **14**, e0212443, (2019).
- 51 Seidel, E. *et al.* Dynamic Co-evolution of Host and Pathogen: HCMV Downregulates the Prevalent Allele MICA*008 to Escape Elimination by NK Cells. *Cell reports* **10**, 968-982, (2015).
- 52 Horby, P. W. *et al.* Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): preliminary results of a randomised, controlled, open-label, platform trial. *medRxiv*, 2021.2002.2011.21249258, (2021).
- 53 Group, R. C. *et al.* Dexamethasone in Hospitalized Patients with Covid-19. *N Engl J Med* **384**, 693-704, (2021).
- 54 Karki, R. *et al.* Synergism of TNF-alpha and IFN-gamma Triggers Inflammatory Cell Death, Tissue Damage, and Mortality in SARS-CoV-2 Infection and Cytokine Shock Syndromes. *Cell* **184**, 149-168 e117, (2021).
- 55 Forthal, D. N. & Finzi, A. Antibody-dependent cellular cytotoxicity in HIV infection. *AIDS* **32**, 2439-2451, (2018).
- 56 Nelson, C. S. *et al.* HCMV glycoprotein B subunit vaccine efficacy mediated by nonneutralizing antibody effector functions. *Proc Natl Acad Sci U S A* **115**, 6267-6272, (2018).
- 57 Shrock, E. *et al.* Viral epitope profiling of COVID-19 patients reveals cross-reactivity and correlates of severity. *Science* **370**, (2020).
- 58 Yewdell, J. W., Frank, E. & Gerhard, W. Expression of influenza A virus internal antigens on the surface of infected P815 cells. *J Immunol* **126**, 1814-1819, (1981).
- 59 Fujimoto, Y. *et al.* Cross-protective potential of anti-nucleoprotein human monoclonal antibodies against lethal influenza A virus infection. *J Gen Virol* **97**, 2104-2116, (2016).
- 60 Virelizier, J. L., Allison, A. C., Oxford, J. S. & Schild, G. C. Early presence of ribonucleoprotein antigen on surface of influenza virus-infected cells. *Nature* **266**, 52-54, (1977).
- 61 Vandervén, H. A. *et al.* What Lies Beneath: Antibody Dependent Natural Killer Cell Activation by Antibodies to Internal Influenza Virus Proteins. *EBioMedicine* **8**, 277-290, (2016).
- 62 Jegaskanda, S. *et al.* Induction of H7N9-Cross-Reactive Antibody-Dependent Cellular Cytotoxicity Antibodies by Human Seasonal Influenza A Viruses that are Directed Toward the Nucleoprotein. *The Journal of infectious diseases* **215**, 818-823, (2017).
- 63 Roltgen, K. *et al.* Defining the features and duration of antibody responses to SARS-CoV-2 infection associated with disease severity and outcome. *Sci Immunol* **5**, eabe0240, (2020).
- 64 de Campos Mata, L. *et al.* SARS-CoV-2-Specific Antibody Profiles Distinguish Patients with Moderate from Severe COVID-19. *medRxiv*, 2020.2012.2018.20248461, (2020).
- 65 Atyeo, C. *et al.* Distinct Early Serological Signatures Track with SARS-CoV-2 Survival. *Immunity* **53**, 524-532 e524, (2020).
- 66 Voss, C. *et al.* Epitope-resolved serology test differentiates the clinical outcome of COVID-19 and identifies defects in antibody response in SARS-CoV-2 variants. *medRxiv*, 2021.2003.2016.21253716, (2021).
- 67 Schulte-Schrepping, J. *et al.* Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell Compartment. *Cell* **182**, 1419-1440 e1423, (2020).
- 68 Szabo, P. A. *et al.* Longitudinal profiling of respiratory and systemic immune responses reveals myeloid cell-driven lung inflammation in severe COVID-19. *Immunity*, (2021).

- 69 Merad, M. & Martin, J. C. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat Rev Immunol* **20**, 355-362, (2020).
- 70 Moore, C. *et al.* Localised community circulation of SARS-CoV-2 viruses with an increased accumulation of single nucleotide polymorphisms that adversely affect the sensitivity of real-time reverse transcription assays targeting Nucleocapsid protein. *medRxiv*, 2021.2003.2022.21254006, (2021).
- 71 Del Vecchio, C. *et al.* Emergence of N antigen SARS-CoV-2 genetic variants escaping detection of antigenic tests. *medRxiv*, 2021.2003.2025.21253802, (2021).
- 72 Madhi, S. A. *et al.* Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. *N Engl J Med*, (2021).
- 73 Weekes, M. P. *et al.* Proteomic plasma membrane profiling reveals an essential role for gp96 in the cell surface expression of LDLR family members, including the LDL receptor and LRP6. *Journal of proteome research* **11**, 1475-1484, (2012).
- 74 Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* **47**, D442-D450, (2019).
- 75 Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57, (2009).
- 76 Gordon, D. E. *et al.* A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature* **583**, 459-468, (2020).
- 77 Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nature Medicine* **26**, 1033-+, (2020).

Fig 1. SARS-CoV-2 remodels the Plasma Membrane Proteome. (A) AAT cells were infected with SARS-CoV2 in biological triplicate. 24h later they were detached with trypsin, fixed and permeabilised, stained for Spike protein, and analysed by flow cytometry. (B) Schematic of plasma membrane profiling and analysis pathway. (C) Examples of temporal profiles of viral and cellular genes, fold change is compared to 0 h timepoint. Data points show mean $\pm$ SD. (D) Left – heat map of the 914 significantly changing proteins clustered by k-mean, colour indicates log₂ fold change compared to 0 h, right, Z-score normalised temporal profiles of proteins within each cluster.

Figure 2. SARS-CoV2 manipulates multiple classes of plasma membrane proteins. (A) Enriched gene ontology and pathway annotations of proteins downregulated in temporal cluster 1 were condensed to groups of related terms, bar chart shows the enrichment score of these groups. Examples of annotation terms falling within each group are shown underlined with Bejamani-Hochberg adjusted P-values in parentheses, alongside four examples of proteins for each annotation.

Figure 3. SARS-CoV2 infection leads to downregulation of multiple NK ligands, inhibiting NK activation. (A) AAT cells were either mock infected, or infected with SARS-CoV2 for 24h, detached using TrypLE, then stained for the indicated NK cell receptors before being analysed by flow cytometry (bottom). Plots for the same proteins from PMP are included for reference (top) (B) AAT cells were either mock infected, or infected with SARS-CoV2 for 24h, detached with TrypLE and mixed with PBMC for 5h in the presence of golgistop, golgiplug, and CD107a antibody, before staining for CD3/CD56/CD57 and Live/Dead Aqua. Cells were then fixed, permeabilised, and stained for TNF α and IFN γ . Cells were gated on live CD57+ NK cells, and the percentage of cells positive for CD107a, TNF α and IFN γ calculated. Individual assays were run in triplicate, with data combined from multiple donors as indicated. Kruskal–Wallis, * $p < 0.05$, ** $p < 0.01$.

Figure 4. SARS-CoV2 inhibition of NK activation can be overcome via ADNKA. AAT cells were either mock infected, or infected with SARS-CoV2 for 24h, detached using TrypLE, then mixed with PBMC in the presence of golgistop, CD107a antibody, and serum from donors who were seronegative or seropositive for SARS-CoV2. After 5h, cells were stained for CD3, CD56, CD57, and live/dead aqua, then analysed by flow cytometry. (A-F) assays were performed using 1% serum, and the percentage of CD57⁺ NK cells positive for CD107a (A, D), TNF α (B, E), and IFN γ (C, F) were calculated. Individual assays were performed in triplicate (A-C), 2-way ANOVA **p < 0.01, ****p < 0.0001. Assays from multiple donors were also compared (D-E), Kruskal–Wallis **p < 0.01, ***p < 0.001. (G-H) assays were performed using a serial 3-fold dilution of serum, starting from 3.3%, then the area under the curve (AUC) calculated. (G) longitudinal serums from donors who experienced mild (black) or severe (grey) were used. (H) In addition to AUC for ADCC activity, serums were tested for their ability to neutralise SARS-COV2 infection of VeroE6 cells, and the NT50 calculated, then compared to the AUC for ADCC; Spearman rank correlation analysis is shown.

A

B

C

Figure 5. Multiple SARS-CoV2 proteins mediate ADCC. (A) Plots of viral proteins detected in the PMP analysis. (B) 293T cells were transfected with plasmids expressing the indicated SARS-CoV2 proteins for 48h, then mixed with PBMC in the presence of golgistop, CD107a antibody, and 1% serum from donors who were seronegative or seropositive for SARS-CoV2. After 5h, cells were stained for CD3, CD56, CD57, and live/dead aqua, then analysed by flow cytometry. Assays were performed in triplicate. Data is shown from two donor serums. ** $p < 0.01$, **** $p < 0.0001$, 2-way ANOVA. (C) 293T cells were transfected with plasmid expression SARS-CoV2 nucleocapsid for 48h, or AAT cells were infected with SARS-CoV2 for 24h, then cells were detached with TrypLE, stained for nucleocapsid, and analysed by flow cytometry.

Figure 6. Monoclonal anti-spike antibodies bind infected cells strongly, but only weakly activate ADNKA. AAT cells were either mock infected, or infected with SARS-CoV2 for 24h, detached with TrypLE and mixed with PBMC for 5h in the presence of golgistop, golgiplug, and CD107a antibody, before staining for CD3, CD56, CD57 and Live/Dead Aqua. Cells were then fixed, permeabilised, and stained for TNF α and IFN γ . Cells were gated on live CD57+ NK cells, and the percentage of cells positive for CD107a (A), TNF α (B) and IFN γ (C) calculated. Assays were run in triplicate. 2-way ANOVA ***p<0.001, ****p<0.0001.

Figure 7. Antibodies targeting spike are weak activators of ADNKA, and the ADNKA response is dominated by non-spike antibodies following natural infection. Serums from individuals naturally infected with SARS-CoV2 (A-E), or donors that had been vaccinated against SARS-CoV2 but were seronegative prior to vaccination (F-J) were depleted of anti-spike antibodies using spike Trimer protein conjugated to magnetic beads. (A, F) ELISA for spike trimer was

used to measure levels of antibodies before and after depletion. (B, C, G, H) the ability of the original, or anti-spike depleted, serums to neutralise the ability of SARS-CoV2 to infect VeroE6 cells was determined across a range of concentrations, then NT50 values calculated. Example plots (B, G) and NT50 values for multiple donors (C, H) are shown. (D, E, I, J) AAT cells were either mock infected, or infected with SARS-CoV2 for 24h, detached using TrypLE, then mixed with PBMC in the presence of golgistop, CD107a antibody, and serial dilutions of serum. After 5h, cells were stained for CD3, CD56, CD57, and live/dead aqua, then analysed by flow cytometry for the percentage of CD107a positive CD57+ NK cells. AUC values were then calculated. Example plots (D, I) and AUC values for multiple donors (E, J) are shown. NT50 (K) and AUC (L) values are also given for samples categorised according to disease or vaccination status. Kruskal–Wallis * $p < 0.05$, ** $p < 0.001$.

Figure S1. Plasma Membrane Proteins are enriched in the PMP dataset. (A) Pie charts showing the proportion of proteins falling into previously defined annotation categories for plasma membrane proteins by left, protein abundance, or right, protein identifiers. (B) Pie chart showing the distribution of proteins defined by the Molecular Signatures Database (MSigDB) gene set “Hallmark Interferon Alpha Response” into the defined temporal clusters. C. Example temporal profiles of type I interferon inducible genes upregulated in cluster 5, y-axis shows fold change compared to 0 h timepoint.

Figure S2. NK ligands detected in the PMP Dataset. Temporal profiles of NK-cell ligands detected in the PMP dataset, y-axis shows fold change compared to 0 h timepoint.

Figure S3. Anti-spike antibodies bind to SARS-CoV2 infected cells. AAT cells were mock infected (light grey), or infected with SARS-CoV2 (black) for 24h, then detached with TrypLE,

stained with the indicated human anti-spike antibodies, then analysed by flow cytometry.

<p>Start Time (min): 0 End Time (min): 190 Cycle Time (sec): 3</p> <p>Master Scan:</p> <p>MS OT</p> <p>Detector Type: Orbitrap Orbitrap Resolution: 120000 Mass Range: Normal Use Quadrupole Isolation: True Scan Range (m/z): 400-1500 RF Lens (%): 60 AGC Target: Custom Normalized AGC Target (%): 125 Maximum Injection Time Mode: Custom Maximum Injection Time (ms): 50 Microscans: 1 Data Type: Profile Polarity: Positive Source Fragmentation: Disabled Scan Description:</p> <p>Filters:</p> <p>MIPS</p> <p>Monoisotopic Peak Determination: Peptide</p> <p>Charge State</p> <p>Include charge state(s): 2-7 Include undetermined charge states: False</p> <p>Dynamic Exclusion</p> <p>Use Common Settings: False Exclude after n times: 1 Exclusion duration (s): 90 Mass Tolerance: ppm Low: 10 High: 10 Exclude Isotopes: True Perform dependent scan on single charge state per precursor only: True</p> <p>Intensity</p> <p>Filter Type: Intensity Threshold Intensity Threshold: 5.0e3</p> <p>Data Dependent</p> <p>Data Dependent Mode: Cycle Time Time between Master Scans (sec): 3</p> <p>Scan Event Type 1:</p> <p>Scan:</p> <p>ddMS² IT CID</p> <p>Isolation Mode: Quadrupole Isolation Window (m/z): 0.7 Isolation Offset: Off</p>	<p>Activation Type: CID Collision Energy Mode: Fixed CID Collision Energy (%): 35 CID Activation Time (ms): 10 Activation Q: 0.25 Multistage Activation: False Detector Type: Ion Trap Ion Trap Scan Rate: Rapid Mass Range: Normal Scan Range Mode: Auto AGC Target: Custom Normalized AGC Target (%): 80 Maximum Injection Time Mode: Auto Microscans: 1 Data Type: Centroid Scan Description:</p> <p>Filters:</p> <p>Precursor Selection Range</p> <p>Selection Range Mode: Mass Range Mass Range (m/z): 400-2000</p> <p>Precursor Ion Exclusion</p> <p>Exclusion mass width: m/z Low: 19 High: 7</p> <p>Isobaric Tag Loss Exclusion</p> <p>Reagent: TMT</p> <p>Data Dependent</p> <p>Data Dependent Mode: Scans Per Outcome</p> <p>Scan Event Type 1:</p> <p>Scan:</p> <p>ddMS² OT HCD</p> <p>MSⁿ Level: 3 Synchronous Precursor Selection: True Number of SPS Precursors: 10 MS Isolation Window (m/z): 2 MS2 Isolation Window (m/z): 2 Isolation Offset: Off Activation Type: HCD HCD Collision Energy (%): 65 Detector Type: Orbitrap Orbitrap Resolution: 50000 Mass Range: Normal Scan Range Mode: Define m/z range Scan Range (m/z): 100-1000 AGC Target: Custom Normalized AGC Target (%): 40 Maximum Injection Time Mode: Custom Maximum Injection Time (ms): 120 Microscans: 1 Data Type: Profile Use EASY-IC™: False Scan Description: Number of Dependent Scans: 3</p>
---	---

Figure S4. MS Method parameters

The figure displays four overlapping software windows used for MS data processing, showing various search parameters.

Data Refinement window:

- Predefined parameters: TheOne
- ☒ Merge Scans [DDA]
- ☒ Correct Precursor [DDA]
 - ☒ Mass only
 - ☐ Mass and Charge states
 - Min charge: 1 Max charge: 3
- ☒ Associate feature with chimera scan [DDA]
- ☒ Filter Features
 - Only keep features satisfying:
 - ☐ m/z between [] and []
 - ☐ Retention time between [] and [] min
 - ☒ Charge between 2 and 8
 - ☐ Abundance $\geq$ []
- Data refinement is performed on each fraction separately
- Buttons: OK, Cancel, Help

De Novo window:

- Predefined parameters: []
- Error Tolerance**
 - Precursor mass: 15.0 ppm
 - Fragment ion: 0.5 Da
- Enzyme**
 - Specified by each sample
 - Buttons: New
- PTM**
 - Buttons: Set PTM, Remove, Switch type
 - PTM list: Beta-methylthiolation, TMT16plex, Acetylation (Protein N-term), Deamidation (NQ), Oxidation (M)
- Maximum allowed variable PTM per peptide: 3
- General Options**
 - Report up to 5 candidates per spectrum
- Buttons: OK, Cancel, Help

PEAKS Search window:

- Predefined parameters: []
- Error Tolerance**
 - Precursor mass: 10.0 ppm using monoisotopic mass
 - Fragment ion: 0.6 Da
- Enzyme**
 - Trypsin/LysC
 - Buttons: View
- Digest mode: Specific
- Maximum missed cleavages per peptide: 3
- PTM**
 - Buttons: Set PTM, Remove, Switch type
 - PTM list: Beta-methylthiolation, TMT11plex, Deamidation (NQ), Acetylation (Protein N-term), Oxidation (M)
- Maximum allowed variable PTM per peptide: 3
- Database**
 - ☒ Select database Database: SP_Human+CoV2 Buttons: View
 - ☐ Paste sequence Taxa: all species Set/View taxa...
 - ☒ Contaminant database Contaminants Buttons: View
- De Novo Tag Options**
 - Available de novo tags: de novo with current parameter
- General Options**
 - ☒ Estimate FDR with decoy-fusion.
 - ☐ Find unspecified PTMs with PEAKS PTM Advanced Settings
 - ☐ Find more mutations with GPM

Quantification window:

- Predefined parameters: []
- Reporter Ion Quantification**
 - Select Methods: TMT-16plex (CID/HCD) Buttons: View
- Basic Options**
 - Mass Error Tolerance: 0.02 Da
 - Reporter Ion Type: ☐ MS2 ☒ MS3
 - ☐ -10logP Threshold 21.6
 - ☒ FDR Threshold(%) 1.0
- Purity Correction**
 - ☒ Perform Purity Correction Buttons: Edit Factors ...

Figure S5. Search Parameters for MS data processing

Table S2. Clinical characteristics of patients giving longitudinal serum samples

Study ID	Severity	Age	Gender	Past medical history
1	Mild	82	M	dementia, ESRF, HTN, IHD, PVD
2	Mild	77	F	ESRF, HTN, T2DM
3	Mild	64	F	AF, CVA, dementia, ESRF, HTN, T2DM
4	Mild	27	F	ESRF, obesity
5	Mild	48	F	ESRF, HTN, obesity
6	Severe	52	F	Asthma, obesity
7	Severe	62	M	colon ca, epilepsy, HTN, obesity, T2DM
8	Severe	65	F	HTN, hypothyroidism, obesity
9	Severe	61	M	ESRF, obesity, VTE
10	Severe	57	M	
11	Mild	23	F	CKD, HTN, T1DM
12	Mild	62	M	ESRF, HTN, renal transplant
13	Mild	25	F	AML
14	Mild	52	M	CVA, prostate ca, T2DM
15	Mild	88	F	AF, CCF, CKD, COPD, IHD, RA, T2DM
16	Mild	64	M	
17	Mild	69	M	AF, HTN, T2DM, VTE
18	Mild	22	M	NHL
19	Severe	49	M	
20	Severe	67	F	AF, HTN, obesity, T2DM

AF=atrial fibrillation, AML=acute myeloid leukaemia, ca=cancer, CCF=congestive cardiac failure, CKD=chronic kidney disease, COPD=chronic obstructive pulmonary disease, CVA cerebrovascular accident, ESRF=end-stage renal failure, HTN=hypertension, IHD=ischemic heart disease, NHL=non-Hodgkin lymphoma, RA=rheumatoid arthritis, T1DM/T2DM=type 1/2 diabetes mellitus, VTE=venous thromboembolism.

mAb name	Specificity	Competition group	S EC50	RBD EC50	NTD EC50	S2 ELISA	SARS-CoV-2 WT FL IC50	SARS-CoV-2 WT PV IC50
P008_060	NON-S1	7	0.031	>10	>10	n.d.	32.90	0.095
P008_004	NON-S1	n.d.	0.045	>10	>10	+	>100	>100
P008_005	NON-S1	n.d.	0.20	>10	>10	+	>100	>100
P008_006	NON-S1	n.d.	0.10	>10	>10	+	>100	>100
P008_016	NON-S1	n.d.	0.11	>10	>10	+	>100	>100
P008_023	NON-S1	n.d.	0.40	>10	>10	+	>100	>100
P008_032	NON-S1	n.d.	0.060	>10	>10	+	>100	>100
P008_051	NTD	5	0.091	>10	0.056	n.d.	48.82	0.30
P008_052	NTD	5	0.085	>10	0.063	n.d.	48.65	0.13
P003_027	NTD	6	0.0039	>10	0.0065	n.d.	1.64	12.92
P008_056	NTD	6	0.060	>10	0.049	n.d.	0.014	>100
P008_001	NTD	n.d.	0.82	>20	1.20	n.d.	>100	>100
P008_002	NTD	n.d.	0.0084	>20	0.015	n.d.	>100	>100
P008_014	NTD	n.d.	0.066	>20	0.021	n.d.	1.81	>100
P008_099	NTD	n.d.	0.022	>20	0.15	n.d.	>100	>100
P054_044	NTD	n.d.	0.031	>20	4.00	n.d.	0.94	0.74
P003_016	NTD	n.d.	0.13	>20	2.00	n.d.	0.15	0.45
P003_014	RBD	1	0.0076	0.0086	n.d.	n.d.	>100	>100
P054_027	RBD	1	0.0077	0.013	n.d.	n.d.	>100	0.7
P008_018	RBD	3	0.071	0.082	n.d.	n.d.	0.055	0.00081
P008_090	RBD	3	0.018	0.026	n.d.	n.d.	0.071	0.0042
P008_108	RBD	3	0.0092	0.014	n.d.	n.d.	0.0023	0.0031
P008_015	RBD	4	0.034	0.036	n.d.	n.d.	0.055	0.00012
P008_096	RBD	4	0.030	0.055	n.d.	n.d.	9.09	0.41

Table S3. Monoclonal Ant-spike Antibodies used, data taken from⁴⁴