Title: The volatile cedrene from plant beneficial *Trichoderma guizhouense* modulates *Arabidopsis* root development through auxin transport and signaling

Running title: Fungal cedrene reprograms root architecture

One-sentence Summary: Cedrene, a high- abundance sesquiterpenes produced by plant beneficial *Trichoderma guizhouense* NJAU 4742, stimulates *Arabidopsis* lateral root formation and primary root elongation by relying on auxin signaling pathway and auxin transporter PIN2 and AUX1.

Authors: Yucong Li^{1¶}, Jiahui Shao^{1¶}, Yansong Fu¹, Yu Chen¹, Hongzhe Wang², Zhihui Xu¹, Haichao Feng¹, Weibing Xun¹, Yunpeng Liu³, Nan Zhang¹, Qirong Shen¹, Wei Xuan^{2,†}, Ruifu Zhang^{1,3,†}

[¶]These authors contributed equally to this work.

Addresses: ¹Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, National Engineering Research Center for Organic-based Fertilizers, Jiangsu Collaborative Innovation Center for Solid Organic Waste Resource Utilization, Nanjing Agricultural University, Nanjing 210095, China

²State Key Laboratory of Crop Genetics and Germplasm Enhancement and MOA Key Laboratory of Plant Nutrition and Fertilization in Lower-Middle Reaches of the Yangtze River, Nanjing Agricultural University, Nanjing 210095, China

³Key Laboratory of Microbial Resources Collection and Preservation, Ministry of Agriculture, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing 100081, China

†Corresponding authors:

Ruifu Zhang, rfzhang@njau.edu.cn. Tel:86-25-84396477, Fax:86-25-84396260;

Wei Xuan, wexua@njau.edu.cn.

Footnotes:

This work was funded by the National Natural Science Foundation of China (31972512 and 32072665), the Agricultural Science and Technology Innovation Program of CAAS (No. CAAS-ZDRW202009) and Key R&D Program of Shandong Province (2019JZZY020614).

Author contributions:

31 R.Z. planned and designed the research. Y.C.L. performed the experiments and wrote
32 the manuscript. R.Z. and W.X. polished the manuscript. J.H.S. and Y.S.F. and helped
33 with root phenotype analysis and imaging. Y.C. and H.Z.W. analyzed the data. All
34 authors discussed the results. R.Z. and W.X. agree to serve as the authors responsible
35 for contact and ensure communication.
36

ABSTRACT

Rhizosphere microorganisms interact with plant roots by producing chemical signals to regulate root development. However, the involved distinct bioactive compounds and the signal transduction pathways are remaining to be identified. Here, we show that sesquiterpenes (SQTs) are the main volatile compounds produced by plant beneficial *Trichoderma guizhouense* NJAU 4742, inhibition of SQTs synthesis in this strain indicated their involvement in plant-fungus cross-kingdom signaling. SQTs component analysis further identified the cedrene, a high abundant SQT in strain NJAU 4742, could stimulate plant growth and root development. Genetic analysis and auxin transport inhibition showed that auxin receptor TIR1, AFB2, auxin-responsive protein IAA14, and transcription factor ARF7, ARF19 affect the response of lateral roots to cedrene. Moreover, auxin influx carrier AUX1, efflux carrier PIN2 were also indispensable for cedrene-induced lateral root formation. Confocal imaging showed that cedrene affected the expression of *pPIN2:PIN2:GFP* and *pPIN3:PIN3:GFP*, which may be related to the effect of cedrene on root morphology. These results suggest that a novel SQT molecule from plant beneficial *T. guizhouense* can regulate plant root development through auxin transport and signaling.

KEY WORDS

sesquiterpenes (SQTs), cedrene, volatile compounds, auxin, lateral root, *Trichoderma*

INTRODUCTIONS

The periodic formation of lateral roots (LR) is a post-embryonic process that is regulated by both endogenous and environmental cues (Xie et al., 2019; Li et al., 2021). Auxin plays the central role in all stages of LR formation, which consists of LR priming, LR initiation and patterning and LR emergence (Lavenus et al., 2013; Santos Teixeira and Ten Tusscher, 2019). Following the act of lateral root cap (LRC) derived auxin on the oscillation zone (OZ), an oscillatory signal is generated to control xylem pole pericycle cells (XPP) ‘priming’ and pre-branch sites forming (Xuan et al., 2020). Auxin regulates LR initiation by controlling LR founder cell divisions and LR founder cell polarity and/or identity acquisition (Dubrovsky et al., 2008). In the LR emergence stage, auxin induces the expression of cell wall-remodeling enzymes to promote cell separation, changes the cell turgor pressure in the outer tissue layers and in the lateral root primordia (LRP), these progresses will help LRPs to go through three overlaying cell layers (endodermis, cortex, and epidermis) to emerge as the LR (Neuteboom et al., 1999; Laskowski et al., 2006; Swarup et al., 2008; Peret et al., 2012; Lee et al., 2013).

Environmental factors, which include water, salt, drought, light, nitrate and phosphate, affect LR formation by interfering with auxin synthesis, conjugation, and degradation, as well as auxin transport or response (Santos Teixeira and Ten Tusscher, 2019). The rhizosphere, defined as the narrow zone influenced by plant roots and characterized by their intense association with microbial activity (Mendes et al., 2013; van Dam and Bouwmeester, 2016), is relatively rich in nutrients, because about 20-40% of the photosynthetic products can be lost from the root in the form of root exudates, including ions, oxygen, water, enzymes, mucus and primary and secondary metabolites (Bais et al., 2006; Philippot et al., 2013; Venturi and Keel, 2016). Consequently, a large number and variety of microorganisms, including bacteria and fungi, were inhabited in the rhizosphere. The rhizosphere microorganisms affect root system architecture (RSA) by the production of phytohormones and secondary metabolites, such as auxin, cytokinin and 2,4-diacetylphloroglucinol (DAPG), which interfere with auxin-dependent signaling pathways in plants (Vacheron et al., 2013).

89 Microbes also produce many volatile compounds (VCs) with the ability to reprogram
 90 RSA (Zhang et al., 2007; Kanchiswamy et al., 2015; Werner et al., 2016; Tyc et al.,
 91 2017; Fincheira and Quiroz, 2018), which can evaporate and diffuse easily far from
 92 their original point and migrate in soil and aerial environments for their low molecular
 93 masses and low polarity, and as well as a high vapor pressure (Schulz and Dickschat,
 94 2007; Schmidt et al., 2015). VC component of indole emitted by soil-borne bacteria
 95 affected LR development in *Arabidopsis* through auxin signaling, depend on polar
 96 auxin transport system (Bailly et al., 2014). (–)-Thujopsene, a kind of sesquiterpenes
 97 (SQTs), produced by *Laccaria bicolor*, stimulated LR formation in *Arabidopsis* by
 98 inducing superoxide anion radicals in roots, independent on auxin signaling pathways
 99 (Ditengou et al., 2015). 6-pentyl-2H-pyran-2-one (6-PP) detected in *Trichoderma*
 100 *atroviride* regulated LR development through auxin signaling and transport in the root
 101 system (Garnica-Vergara et al., 2016). The VCs produced by *T. viride* with the main
 102 ingredients of isobutyl alcohol, isopentyl alcohol, and 3-methylbutanal, increased
 103 biomass in the shoot and the root system (Hung et al., 2013). So far, considerable
 104 progress has been made in elucidating the mode of action of VCs; however, it is still
 105 poorly understood, especially in the identification of distinct bioactive compounds.

106 *Trichoderma* species are able to colonize the root surface to promote the root
 107 development and plant growth, which also represent excellent biocontrol agents in
 108 agriculture because of their strong ability to fight with plant pathogens (Druzhinina et
 109 al., 2011). *Trichoderma* species have broad VC profiles (Hung et al., 2013; Muller et
 110 al., 2013), the produced SQTs were supposed to be good candidates for underground
 111 microbe-plant signaling as SQTs were the representative diffusing compounds in
 112 complex environment (Hiltpold and Turlings, 2008). In this study, we identified the
 113 cedrene from *Trichoderma guizhouense* NJAU 4742, the plant beneficial fungus with
 114 efficient promotion for plant growth and root development, as well as the soil-borne
 115 pathogen suppression (Zhang et al., 2016; Meng et al., 2019; Zhang et al., 2019). The
 116 cedrene can enhance shoot and root biomass and stimulate root development in a
 117 dose-dependent manner. Cedrene induced auxin response in primary root tips and
 118 early stages LRP (I-III), but not the late stages LRP (IV-VII). Furthermore, cedrene

119 differentially modulated expression of auxin transporters PIN1, PIN2, PIN3 and PIN7.
 120 A genetic screen for cedrene resistance established that this compound required auxin
 121 receptors TIR1 and AFB2, and downstream auxin-responsive protein IAA14, and
 122 transcription factors ARF7 and ARF19 to stimulate lateral root development.
 123 Moreover, auxin influx carrier AUX1 and efflux carrier PIN2 were both indispensable
 124 for promoting lateral root formation.

125

126

RESULTS

VCs released by *T. guizhouense* NJAU 4742 promote *Arabidopsis* growth and root development

Arabidopsis plants were grown in the presence of fungal VCs in bi-compartmented Petri dishes (Fig. 1a). After 8 d of co-cultivation, despite the absence of direct contact with *T. guizhouense* NJAU 4742, strong stimulation of growth and root development was observed in *Arabidopsis*, which should have been caused by VCs of NJAU 4742. *T. guizhouense* VCs slightly promoted primary root elongation, but significantly increased the lateral root number and lateral root density by 37% and 39%, respectively (Fig.1a-c). In addition, *T. guizhouense* VCs can also increase the root and shoot biomass by 101% and 56% (Fig.1e, f), respectively, and increase the total biomass production of plants by 71% (Fig.1g).

Cedrene mimics fungal VCs effects in promoting LR formation

To identify the VCs component responsible for LR induction, VCs of *T. guizhouense* NJAU 4742 in the headspace was measured by SPME-GC-MS. Table 1 shows that sesquiterpenes (SQTs) are the major compounds within the VC profile (75.93%) from *T. guizhouense* NJAU 4742. Previous studies have shown that SQTs from ectomycorrhizal fungi *Laccaria bicolor* can reprogramme root architecture (Ditengou et al., 2015). To test whether SQTs in *T. guizhouense* VCs are related to the induction of lateral root formation, we inhibited SQT biosynthesis using inhibitor lovastatin (Rodriguez-Concepcion, 2006; Ditengou et al., 2015) in *T. guizhouense* NJAU 4742 and investigated the effect on LR stimulation in *Arabidopsis*. The result showed that the effect of *T. guizhouense* VCs on promoting root development was abolished when fungal SQTs were suppressed by lovastatin (Fig. 2a-c). To identify distinct bioactive compounds, we tested the effect of a pure product (cedrene) that can be obtained and contained in a large amount in *T. guizhouense* VCs profile on the plant growth-promoting activity. The *Arabidopsis* seedlings were treated with DMSO (as control) or with 10-2000 μ M cedrene dissolved in DMSO (Fig. 3a). After 8 d of growth in medium supplied with 20-100 μ M cedrene, a significant increase in shoot, root and total plant biomass was observed (Fig. 3b-d). By contrast, the higher

Figure 1. Effect of *Trichoderma guizhouense* NJAU 4742 volatile compounds on *Arabidopsis* biomass production and root development. (a) Three-day-old *Arabidopsis thaliana* (Col-0) seedlings were germinated and grown for 8 d in the presence of *T. guizhouense* NJAU 4742 in a bi-compartmented Petri dish avoiding direct contact and solute exchange between the plant and the fungus. Scale bar, 2 cm. (b and c) Quantification of primary root elongation (b) and number of emerged lateral roots (c). $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). (d) Lateral root density (number of emerged lateral roots cm^{-1}). $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). (e-g) Quantification of root biomass (e), shoot biomass (f) and total biomass (g) production per *Arabidopsis* seedling. $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Error bars indicate $\pm$ SD of the mean.

Figure 2. Sesquiterpenes in *Trichoderma guizhouense* NJAU 4742 volatile compounds are the main bioactive compounds for inducing *Arabidopsis* root development. (a) Three-day-old *Arabidopsis* seedlings were grown for 7 d in bi-compartmented Petri dishes (10cm × 10cm), which was inoculated with *T. guizhouense* (Tg) in the presence or absence of Lov (10 μM) in the adjacent compartment. Scale bar, 2 cm. (b and c) Quantification of primary root elongation (b) and emerged lateral roots number of plants shown in (a). Different letters indicate significant differences of the mean values at $P < 0.05$ (One-way ANOVA, $n = 12$). Error bars indicate $\pm$ SD of the mean.

Figure 3. Effect of cedrene on *Arabidopsis* biomass production and root development.

(a) Representative photographs of *Arabidopsis* seedlings grown for 8 d in 1/2 MS salts agar medium supplied with the solvent (DMSO) or increasing concentrations of cedrene. Scale bar, 2 cm. (b-d) Quantification of shoot biomass (b), root biomass (c), and total biomass (d) per seedling of plants shown in (a). Different letters indicate significant differences of the mean values at $P < 0.05$ (One-way ANOVA, $n = 12$). (e-g) Quantification of primary root elongation (e), emerged lateral roots number (f), and lateral root density (g, number of emerged lateral roots cm^{-1}) of plants shown in (a). Different letters indicate significant differences of the mean values at $P < 0.05$ (One-way ANOVA, $n = 12$). Error bars indicate $\pm$ SD of the mean.

(20-500 μ M) and lower concentration (20-100 μ M) of cedrene could induce the increase of lateral root number and primary root elongation, respectively (Fig. 3e, f), but higher concentration cedrene would have no or even inhibitory effect on that (Fig. 3e, f). Moreover, the cedrene treatments increased lateral root density in a dose-dependent manner (Fig. 3g).

Cedrene affects early LRP stages to induce lateral root formation

To further investigate the stage of LRP that is impacted by cedrene treatment, the developmental stage of LRP in the newly formed PR of control and cedrene-treated plants were quantified. Cedrene induced more LRP formation at very early stages (Stages I) compared with control treatment (Fig. 4c). However, we did not observe changes on later stages LRP (from Stage II to Stage VII) in cedrene-treated seedling roots (Fig. 4c). It is noteworthy that cedrene treatment did not affect the total LRP number (Fig. 4c). Next, we analyzed auxin distribution using the *pDR5:GUS* reporter and an enhanced DR5-directed GUS activity was detected in early stages (Stages I to III) LRP and meristematic protoxylem pole (Fig. 4a, b). Moreover, a strong and diffuse increase of *GUS* activity in primary root of auxin-inducible LRP specific marker line *pGATA23:nls-GUS* (De Rybel et al., 2010) was observed when treated with cedrene (Fig. 4d). Together, these results demonstrated that cedrene might stimulate early lateral roots initiation and following LRP development by affecting auxin distribution and its downstream signaling.

Cedrene does not alter DR5 oscillation in roots

In order to investigate whether cedrene is involved in the regulation of ‘lateral root clock’, which generates an oscillatory signal that is translated into a developmental cue to specify a set of founder cells for LR formation (Xuan *et al.*, 2020), we used *pDR5:LUC* (Fig. 5a), which fused *DR5* promoter to the *luciferase* (LUC) gene to mark early LR founder cell positions and allow visualization of its behavior *in vivo* (Moreno-Risueno et al., 2010; Van Norman et al., 2013; Laskowski and Ten Tusscher, 2017; Xuan et al., 2020). Expression of *pDR5:LUC* in the root tip showed oscillatory activity, and a static point of expression, which is the future site of LRP and LR and therefore is called as prebranch sites (Moreno-Risueno *et al.*, 2010;

Figure 4. The effects of cedrene on lateral root primordium development in *Arabidopsis*. (a) Different stages of lateral root primordia expressing *pDR5:GUS* under solvent (DMSO, Ctrl) or 100 μ M cedrene-treated conditions. Scale bar, 100 μ m. (b) Expression pattern of *DR5:GUS* in the root tips of 3-day-old seedling treated with or without cedrene for six more days. Percentages indicate the proportion of seedlings showing the identical *DR5* expression pattern within a population ($n = 12$). PX, protoxylem pole. Scale bar, 100 μ m. (c) Distribution of lateral root primordia (LRP) in seven developmental classes as defined by the *pCYCB1;1:GUS* activity after treatment with 100 μ M cedrene for 7 more days. $n = 8$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Total LRP, total number of lateral root primordia including all seven developmental stages. ELR, emerged lateral roots. (d) *pGATA23:nls-GUS* expression in primary root under control and cedrene-treated (100 μ M) conditions. Scale bar, 200 μ m.

Figure 5. The effects of cedrene on prebranch site formation and *DR5* oscillation. (a and b) Luciferase imaging and quantification of prebranch site number of 3-day-old *pDR5:Luciferase* seedlings grown for 6d on medium supplied with or without 100 μ M cedrene. The prebranch site in the newly formed primary root after transfer was measured. The red triangle indicates *pDR5:Luciferase* signal in the OZ, and white arrow indicates prebranch site revealed by persistent *pDR5:Luciferase* signal. Yellow dotted line indicates the position of root tip after 4 d co-cultivation. Scale bar, 1 cm. (c) Quantification of prebranch site number of *pDR5:Luciferase* seedlings within 48 h under control and cedrene-treated (100 μ M) conditions. $n \geq 10$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Error bars indicate $\pm$ SD of the mean.

192 treatment comparing with control treatment (Fig. 5b), which is in accordance with the
 193 observation of total LRP number under cedrene treatment (Fig. 4c). Since the
 194 periodicity and signal intensity of *DR5* oscillation are two important factors required
 195 for prebranch site formation, we further examined the effects of cedrene on those. We
 196 measured the periodic production of the prebranch sites in 48 h and found that
 197 cedrene treatment did not affect the formation of prebranch sites (Fig. 5c), indicating
 198 the periodicity of *DR5* oscillation remain unchanged between control and cedrene
 199 treatments. Also, we did not observed a change in the expression intensity of the *DR5*
 200 in OZ for plants grown in the presence of cedrene (Fig. 5a). Therefore, cedrene
 201 induces LR formation is not through affecting *DR5* oscillation.

202 **Effect of cedrene on root development of auxin-related *Arabidopsis* mutants**

203 Since cedrene can enhance the auxin response of early LRP, implying that
 204 auxin plays a role in cedrene-induced lateral roots formation, we further analyzed the
 205 response of wild-type (Col-0) and *Arabidopsis* mutants deficiented in genes related to
 206 auxin transport or response (*tir1afb2*, *slr-1/iaa14*, *arf7arf19*, *aux1-7*, *pin2*, *pin3*) to
 207 cedrene treatments (Fig. 6a). Col-0 and mutant lines were grown in medium
 208 supplemented with the solvent only or with 100 μ M cedrene, and LR formation were
 209 analyzed after 8 d treatment. The results showed that in all these auxin signaling
 210 mutants, including *tir1afb2*, in which oscillation amplitude and prebranch site are
 211 drastically compromised (Xuan et al., 2015), *arf7arf19* and *slr-1/iaa14*, which
 212 completely abolished LR formation (Fukaki et al., 2002; Okushima et al., 2007),
 213 cedrene treatment did not influence LR formation and primary root elongation as
 214 compared with the control treatment (Fig. 6b, c). Moreover, the LR formation and
 215 primary root elongation promoting effect of cedrene on the auxin influx mutant
 216 *aux1-7* and auxin efflux mutant *pin2* were also disappeared (Fig. 6b, c). By contrast,
 217 the promoting effects of cedrene on LR formation and primary root elongation were
 218 not affected in the auxin efflux mutant *pin3*. These results suggested that
 219 cedrene-triggered LR formation and primary root elongation operates via a canonical
 220 auxin-response pathway and auxin transport system is necessary for that.

221 To further address the role of polar auxin transport in cedrene-mediated LR

Figure 6. Influence of auxin signaling and transport on root system architecture modified by cedrene. (a) Representative photographs of wild-type (Col-0) and mutant seedlings of auxin signaling (*tir1afb2*, *slr-1/iaa14* and *arf7arf19*) and transport (*aux1-7*, *pin2* and *pin3*) under control and cedrene (100 μM) treatments for 8 d. Scale bar, 2 cm. (b and c) Quantification of emerged lateral root number (b) and primary root elongation (c) of plants shown in (a). n = 12. **, P < 0.01; *, P < 0.05 (Student's t-test). Error bars indicate ± SD of the mean.

Figure 7. Effects of the polar auxin transport inhibitor NPA on cedrene-induced primary root elongation and LR formation. (a) Representative photographs of Arabidopsis seedlings grown under solvent (Ctrl) and cedrene (100 μ M) treatments in presence of varying concentrations of NPA. Scale bar, 2 cm. (b and c) Quantification of emerged lateral root number (c) and primary root elongation (b) of plants shown in (a). n = 12. **, P < 0.01; *, P < 0.05 (Two-way ANOVA). Error bars indicate $\pm$ SD of the mean.

conditions (Fig. 7a). At 1 μ M NPA, the ability of cedrene to stimulate LR formation was basically unaffected, whereas in the presence of 3 and 5 μ M NPA, LR formation in response to cedrene was completely abolished (Fig. 7c). Moreover, the stimulation effect of cedrene on primary root elongation was completely abolished in the presence of NPA (Fig. 7b). Thus, it can be concluded that a functional auxin efflux machinery is required for cedrene-induced LR formation.

Since the polar auxin transport is mediated by polarly localized PIN proteins (van Berkel et al., 2013), we analyzed the expression pattern of PIN1, PIN2, PIN3 and PIN7 in primary root tips of seedlings expressing *pPIN1:PIN1:GFP*, *pPIN2:PIN2:GFP*, *pPIN3:PIN3:GFP* and *pPIN7:PIN7:GFP* (Fig. 8a-d), to test whether cedrene could regulate LR formation and/or primary root elongation through differential expression of the PINs family of auxin transporters. The results showed that the expression of *pPIN1:PIN1:GFP*, which was only detected in the stele of primary roots under control condition, and *pPIN7:PIN7:GFP*, which was detected in the stele and columella of primary roots under control condition, were not affected by cedrene treatment (Fig. 8a, e, d, h). By contrast, the expression of *pPIN2:PIN2:GFP*, which was detected in the cortex and epidermal cells under control condition, was significantly increased when treated with cedrene (Fig. 8b, f). Furthermore, the expression of *pPIN3:PIN3:GFP*, which was detected in the stele and columella of primary roots under control condition, displayed a weak expression in the stele when treated with cedrene (Fig. 8c, g). These findings suggest that cedrene affects the expression and distribution of the PIN proteins in primary roots and the root response to cedrene does not occur in all tissues, but shows a clear preference for specific tissues and transport components.

DISCUSSION

Previous studies on rhizosphere microbe's beneficial function usually emphasized the first step of rhizospheric colonization, resulting in a direct contact with the root (Bais et al., 2004; Beauregard et al., 2013). However, volatile compounds (VCs) produced by microbes can spread to distant places through the air, porous soil, and liquid environment, so it should be an ideal signal substance to

Figure 8. Expression of auxin efflux transporters in response to cedrene in primary roots. (a-d) Confocal images of *pPIN1:PIN1:GFP* (a), *pPIN2:PIN2:GFP* (b), *pPIN3:PIN3:GFP* (c) and *pPIN7:PIN7:GFP* (d) signal in the primary root tips of 3-day-old seedlings grown with or without cedrene for 6 d. The seedlings were stained with propidium iodide and analyzed by confocal microscopy. Scale bar, 100 μ m. (e-h) Quantification of *pPIN1:PIN1:GFP* (e), *pPIN2:PIN2:GFP* (f), *pPIN3:PIN3:GFP* (g) and *pPIN7:PIN7:GFP* (g) signal intensity at primary root tip shown in (a-d). a.u., arbitrary units. $n \geq 8$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Error bars indicate $\pm$ SD of the mean.

255 a comparative analysis of experimental data has shown that volatile metabolites made
256 an enormous contribution to the microbial interactions than non-volatile ones
257 (Tirranen and Gitelson, 2006). Our results clearly demonstrated that the exchange of
258 molecules between *T. guizhouense* and *Arabidopsis* via intimate contact is not
259 necessary (Fig.1a), and *T. guizhouense* VCs can act as signals to modulate plant
260 growth and development.

261 Actually, it is well recognized that plant-associated microorganisms produce a
262 large number of VCs, which have the potential to play an important role in mediating
263 the interaction between plants and microbes (Kanchiswamy et al., 2015; Piechulla et
264 al., 2017; Fincheira and Quiroz, 2018). So far, most of the effects mediated by
265 microbial VCs have been obtained through co-culture experiments, that is, plants are
266 exposed to complex mixtures of inorganic and organic volatiles released by
267 microorganisms. This makes it difficult to disentangle the functions of individual
268 compounds. On the other hand, the substrate availability and metabolic activities of
269 the microorganisms also will affect the composition of microbial blends to a large
270 extent (Fiddaman and Rossall, 1994). Regarding plants, the observation results
271 obtained by different co-culture experiments vary greatly from strong growth
272 inhibition to significant growth promotion (Splivallo et al., 2007; Li et al., 2021). In
273 order to confirm and provide functional evidence for the microbial VCs action
274 potential, it is necessary to identify distinct biologically active compounds and test
275 them separately or in prescribed mixtures (Piechulla et al., 2017). Here, we discovered
276 cedrene, a high-abundant SQT in the emission profile of *T. guizhouense* NJAU 4742,
277 was sufficient to stimulate LR formation. Cedrene was previously identified in
278 ascomycete *Muscodor albus*, and has the potential to kill a broad range of plant- and
279 human-pathogenic fungi and bacteria synergistically with other sesquiterpenes
280 volatiles (Strobel et al., 2001). Its function as a signaling compound in root branching
281 as demonstrated in this study, has not been reported before. This evidence is closely
282 related to the results previously reported by Ditengou et al. (2015) that (–)-thujopsene,
283 as a low-abundant SQT, can also stimulate lateral root formation. Nevertheless, not all
284 SQTs can induce lateral roots (Ditengou et al., 2015), indicating that the plant

response to these chemicals is specific. Furthermore, along with other environmental factors, such as water, temperature and light, various distinct bioactive microbial volatiles that can affect root system architecture have been identified in bacteria and fungi (Bailly et al., 2014; Ditengou et al., 2015; Garnica-Vergara et al., 2016; Perez-Flores et al., 2017), emphasizing the high plasticity of plant roots in response to heterogeneous macro- and micro-conditions.

The phytohormone auxin fulfils multiple roles throughout LR development (Du and Scheres, 2018). Tryptophan treatment initiated additional LRPs were mostly originated from the auxin-producing pericycle sectors, suggesting that a local auxin input is able to specify lateral root founder cells (LRFCs) (Dubrovsky et al., 2008). In this study, an enhanced expression of *DR5*, which is an established marker for auxin response and indirectly for auxin accumulation (Ulmasov et al., 1997; Sabatini et al., 1999), was observed at early stages (I-III) LRPs and in the columella and meristematic protoxylem pole of root tip under cedrene treatment (Fig. 4a and b). These results suggest that cedrene promotes the formation of early stage LRP by regulating auxin homeostasis in LRPs. Before LRP initiation, the cell files of the xylem pole pericycle (XPP) at one side of the root will be specified and activated to form LRFCs, so that to regulate the spatial distribution of LRPs (Van Norman et al., 2013; Du and Scheres, 2018). The auxin-regulatory GATA23 transcription factor is considered as the first molecular marker for specification of LRFCs (De Rybel et al., 2010). Using a GATA23 RNAi line, in which the expression of GATA23 was reduced to about 30% of normal levels, a strong reduction in the number of stage I-II primordia and an overall decrease in the number of emerged primordia was observed. In contrast, overexpression of GATA23 resulted in a strong increase of stage I and II primordia (De Rybel et al., 2010). This result is basically consistent with our data, cedrene treatment enhanced the expression of GATA23 in root and increased the number of very early stage LRPs (stage I) and ELR, although the total number of LRP was not affected by cedrene treatment (Fig. 4b-d). These results imply that cedrene may stimulate the initiation of LRP by controlling founder cell identity of XPP cells via enhancing GATA23 expression. SOLITARY-ROOT (SLR)/IAA14-ARF7/ARF19

module is an important auxin signal component for LR initiation, in which auxin-induced degradation of unstable SLR protein that de-repressed ARF7 and ARF19 transcription factors, thus activating downstream gene expression (Fukaki et al., 2002; Fukaki et al., 2005; Okushima et al., 2005). Our results showed cedrene treatment could not rescue the lack of LR in *slr-1/iaa14* and *arf7arf19* mutants and failed to induce LR formation in the auxin perception double mutant *tir1afb2* (Fig. 6a, b). These results suggest that canonical auxin-response pathway is indispensable for cedrene induced lateral root formation.

In plants, auxin is generally transported by two types of carriers (i.e. influx carriers and efflux carriers). So far, some transmembrane proteins, such as AUX1/LIKE AUX1 (AUX1/LAX) family, with specific auxin influx functions have been described in *Arabidopsis* (Bennett et al., 1996; Marchant et al., 2002; Swarup et al., 2008). For auxin efflux, PIN protein family and ATP-binding cassette subfamily B (ABCB)-type transporters of the multidrug resistance/phosphoglycoprotein (ABCB/MDR/PGP) protein family have been identified to play the roles (Galweiler et al., 1998; Noh et al., 2001). Both auxin influx and efflux carriers can affect the formation of lateral roots (Marchant et al., 2002; Swarup et al., 2008; Peret et al., 2013). The LR reprogram ability of VCs produced by microbes does not only influence auxin signaling but also its transport through the root system. In *T. atroviride*, 6-PP enhanced the expression of the auxin transporters PIN1, 2 and 3 in the primary root of *Arabidopsis* (Garnica-Vergara et al., 2016). In *F. oxysporum*, the enhanced LR formation by VCs is abolished in auxin transport mutants (Bitas et al., 2015). Moreover, stimulation of LR in *Arabidopsis* by the ectomycorrhizal fungus *L. bicolor* requires PIN2-mediated auxin transport (Felten et al., 2009). Consistent with these observations, treatment with the polar auxin transport inhibitor NPA compromised the ability of cedrene to promote LR formation, suggesting that cedrene is subjected to polar transport. Furthermore, stimulation of LR formation by cedrene was completely abolished in mutant *aux1-7*, suggesting that cedrene-triggered LR formation also requires a complete auxin influx system.

The cedrene-induced LR formation and primary root elongation shown here

opens new avenues for biotechnological application of microbial VCs, and demonstrates clearly that the operation of distinct bioactive microbial VCs is a practical tool to decode the underlying cellular and molecular reactions and mechanism occurring in microbial VC-mediated interactions. Understanding those mechanisms will be a prerequisite for the development of strategies for applying microbial VCs in plant growth in the future.

MATERIALS AND METHODS

Plant materials and growth conditions

Arabidopsis thaliana accessions Col-0 is the wild-type genotype. The marker lines *pDR5:GUS* (Ulmasov et al., 1997), *pDR5:Luciferase* (Moreno-Risueno et al., 2010), *pCYCB1;1:GUS* (Himanen et al., 2002), *pGATA23:nls-GUS* (De Rybel et al., 2010), *pPIN1:PIN1:GFP* (Benkova et al., 2003), *pPIN2:PIN2:GFP* (Blilou et al., 2005), *pPIN3:PIN3:GFP* (Zadnikova et al., 2010), *pPIN7:PIN7:GFP* (Blilou et al., 2005) and the mutant lines *tirlafb2* (Dharmasiri et al., 2005), *slr-1/iaa14* (Fukaki et al., 2002), *arf7arf19* (Okushima et al., 2007), *aux1-7* (Pickett et al., 1990), *pin2* (Roman et al., 1995), *pin3* (*salk_005544*) were used in this study. After 2-3 d of stratification at 4°C in the dark, *Arabidopsis* seeds were surface-sterilized with 30% (v/v) NaClO solution for 10 min. The seeds were germinated and grown on agar plates containing Murashige and Skoog Basal Salts Mixture (MS salts, PhytoTech LABS) in square Petri plates (10 × 10 cm). Standard growth medium consisted of 0.5 × MS salts (2.15 g l⁻¹), 0.1 g l⁻¹ Myo-inositol, 0.5 g l⁻¹ 2-(N-morpholino) ethanesulfonic acid (MES), 1% sucrose (pH 5.7), and 1% Agar (Solarbio). Plants were vertically placed at an angle of 65° in a plant growth chamber, under a long-day photoperiod (16 h: 8 h, light: dark), with a light intensity of 100 μmol m⁻² s⁻¹, at 22 °C. After 3 d of growth, the seedlings were applied for further experiments.

Growth of *T. guizhouense* NJAU 4742 and co-cultivation with plants

For experiments involving fungal VCs, bi-compartmented Petri dishes (9 cm diameter) were used. One of the compartments was filled with MS salts agar medium, and the other one was placed with a patch of MS salts agar medium (1.5 cm × 1.5 cm)

incubated with 3 ul *T. guizhouense* NJAU 4742 (maintained in the Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, China) spores solution. After 3-4 d of fungal growth at 28°C, 3-day-old seedlings (4 plants) were transferred to the compartment filled with MS salts agar. The plates were sealed with breathable tape (MBT, a pressure-sensitive adhesive type usually used in medicine) and incubated for 8 d in a growth chamber at 22°C. At the end of this period, primary root length, lateral root number, and biomass production were recorded.

Analysis of *T. guizhouense* NJAU 4742 VCs

T. guizhouense NJAU 4742 was grown in a head-space bottle containing MS salts agar medium for 6 d at 28 °C. At a constant temperature of 40°C, the head-space bottle was shaken for 60 min at a shaking speed of 450 rpm (5s on and 2s off). Then solid-phase microextraction (SPME) fiber (50/30 µm DVB/CAR on PDMS) was inserted into the headspace of the sample, and the sample was extracted in the headspace for 60 min. The SPME fiber was removed and desorbed at 250°C for 5min, and then was separated and identified by gas chromatography-mass spectrometry (GC-MS). GC-MS analysis was carried out by Agilent 7890B-5977B (GC-MS)_PAL RSI 120, equipped with a chromatographic column (Agilent DB-wax, 30m×0.25mm×0.25µm). Helium (> 99.999%) was used as the carrier gas (1.0 ml/min), and the injection temperature was 260 °C. The column was held at 40 °C for 3 min and then was programmed to increase by 4 °C per min to a final temperature of 220 °C, which was maintained for 10 min. The interface temperature, ion source temperature and quadrupole temperature was 280 °C, 230 °C and 150 °C respectively; the ionization mode was EI⁺, 70ev; the detector voltage was 901V; the scanning mode was full-scan, and the mass range was 20-650 (m/z). These compounds were identified by comparison with mass spectra from a library of NIST (<https://webbook.nist.gov/chemistry/>). The identification of cedrene was performed by comparing retention time (Rt) and the mass spectra from an authentic standard ((-)-α-cedrene, Sigma-Aldrich) with those obtained in the sample.

Chemical preparation and treatments

1-N-naphthylphthalamic acid (NPA) and (-)-α-cedrene (≥ 95%, Sigma-Aldrich)

were dissolved in DMSO to make a 50 mM and 200 mM stock solution, respectively. For treatment, the required amount of the stock solutions was added into MS salts agar and mixed in uniform before being poured into Petri dishes. For lovastatin treatment, lovastatin (PHR1285, Supelco), was dissolved in ethanol to make a 100 mM stock solution, sterile-filtered and added to the MS salts media at the indicated concentration. The plants and *T. guizhouense* were cultivated in a square bi-compartment Petri dish (10 cm × 10 cm), in which a small Petri dish (3cm diameter) was placed to inoculate *T. guizhouense*. *T. guizhouense*, not *Arabidopsis*, grown with lovastatin. LR formation and primary root length was measured after treatment for 7 days.

Phenotypic data analysis

After co-cultivation with fungus or with cedrene for an indicated time period, the length of the newly elongated PR during the treatment was quantified and the emerged LRs in the whole PR were counted. The plates with seedlings were scanned using EPSON XL11000 for the measurement of PR elongation with Fiji software (<http://fiji.sc/>) and the emerged LRs were recorded under a microscope. The LR density was determined by dividing the emerged LR number by the new formed PR length for each analyzed seedling. The biomass production of each seedling was measured on an analytical balance. LRPs were quantified 6 d after co-cultivation. The *pCYCB1;1:GUS* seedlings were stained and cleared to visualize the LRPs at early stages of development and each LRP developmental stage was classified according to Malamy and Benfey (1997) as follows. Stage I, LRP initiation, in the longitudinal plane, approximately 8 to 10 “short” pericycle cells are formed. Stage II, the formed LRP is divided into two layers by a periclinal division. Stage III, the outer layer of the primordium is divided periclinally, generating a three-layer primordium. Stage IV, LRP with four cell layers. Stage V-VII, the midway between the LRP departing from the parent cortex to the LRP appears to be just about to emerge from the parent root.

GUS histochemical staining

For histochemical analysis of GUS activity, the roots of 3-day-old *pDR5:GUS*, *pCYCB1;1:GUS* and *pGATA23:nls-GUS* marker lines were incubated overnight at 37 °C

in a GUS reaction buffer after 6 d of cedrene treatment. The stained roots were cleared using the method of Malamy and Benfey (1997). For each marker line and each treatment, at least 8 transgenic plants were analyzed. A representative sample was chosen and photographed using Leica DM2500 microscope.

Luciferase assay

pDR5:Luciferase expression along the primary root was analyzed by using Lumazon (Xuan et al., 2018). After 6 d of treatment with cedrene, *pDR5:Luciferase* plants were sprayed with a 1mM potassium luciferin (Gold Biotechnology) and reacted for 15 min in darkness, then were imaged immediately with a 15 min exposure time. For investigating the effects of cedrene on the periodicity of *DR5* oscillation, the position of root tip was marked after 4 d of co-cultivation and the prebranch sites formed in the following 2 d was counted. The picture series were saved as TIFF format by IVScopeEQ software for further analysis in Fiji (<http://fiji.sc/>). The Fiji lookup tables "Fire" was used to convert black and white images into color scales based on pixel intensity.

Fluorescence microscopy

For confocal microscopy, control or cedrene-treated transgenic *Arabidopsis* seedlings (*pPIN1:PIN1:GFP*, *pPIN2:PIN2:GFP*, *pPIN3:PIN3:GFP* and *pPIN7:PIN7:GFP*) were mounted in 10 mg ml⁻¹ propidium iodide solution on microscope slides. A Leica SP8 laser-scanning microscope was used for fluorescence imaging of the *Arabidopsis* roots. Each sample was analyzed separately for propidium iodide (with a 568-nm wavelength argon laser for excitation, and an emission window of 585–610 nm) and GFP fluorescence (488 nm excitation/505–550 nm emission). More than eight independent seedlings were analyzed per line, and treatment representative images were selected for figure construction.

Figure legends

Figure 1. Effect of *Trichoderma guizhouense* NJAU 4742 volatile compounds on *Arabidopsis* biomass production and root development. (a) Three-day-old *Arabidopsis thaliana* (Col-0) seedlings were germinated and grown for 8 d in the presence of *T.*

465 *guizhouense* NJAU 4742 (Tg) in a bi-compartmented Petri dish avoiding direct
466 contact and solute exchange between the plant and the fungus. Scale bar, 2 cm. (b and
467 c) Quantification of primary root elongation (b) and number of emerged lateral roots
468 (c). $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). (d) Lateral root density
469 (number of emerged lateral roots cm^{-1}). $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's
470 *t*-test). (e-g) Quantification of root biomass (e), shoot biomass (f) and total biomass (g)
471 production per *Arabidopsis* seedling. $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's
472 *t*-test). Error bars indicate $\pm$ SD of the mean.

473 **Figure 2.** Sesquiterpenes in *Trichoderma guizhouense* NJAU 4742 volatile
474 compounds are the main bioactive compounds for inducing *Arabidopsis* root
475 development. (a) Three-day-old *Arabidopsis* seedlings were grown for 7 d in
476 bi-compartmented Petri dishes (10cm $\times$ 10cm), which was inoculated with *T.*
477 *guizhouense* in the presence (Tg+Lov) or absence (Tg) of 10 μM lovastatin (Lov) in
478 the adjacent compartment. Scale bar, 2 cm. (b and c) Quantification of primary root
479 elongation (b) and emerged lateral roots number of plants shown in (a). Different
480 letters indicate significant differences of the mean values at $P < 0.05$ (One-way
481 ANOVA, $n = 12$). Error bars indicate $\pm$ SD of the mean.

482 **Figure 3.** Effect of cedrene on *Arabidopsis* biomass production and root
483 development. (a) Representative photographs of *Arabidopsis* seedlings grown for 8 d
484 in 1/2 MS salts agar medium supplied with the solvent (DMSO) or increasing
485 concentrations of cedrene. Scale bar, 2 cm. (b-d) Quantification of shoot biomass (b),
486 root biomass (c), and total biomass (d) per seedling of plants shown in (a). Different
487 letters indicate significant differences of the mean values at $P < 0.05$ (One-way
488 ANOVA, $n = 12$). (e-g) Quantification of primary root elongation (e), emerged lateral
489 roots number (f), and lateral root density (g, number of emerged lateral roots cm^{-1}) of
490 plants shown in (a). Different letters indicate significant differences of the mean
491 values at $P < 0.05$ (One-way ANOVA, $n = 12$). Error bars indicate $\pm$ SD of the mean.

492 **Figure 4.** The effects of cedrene on lateral root primordium development in
493 *Arabidopsis*. (a) Different stages of lateral root primordia expressing *pDR5:GUS*
494 under solvent (DMSO, Ctrl) or 100 μM cedrene-treated conditions. Scale bar, 100 μm .

(b) Expression pattern of *DR5::GUS* in the root tips of 3-day-old seedling treated with or without cedrene for six more days. Percentages indicate the proportion of seedlings showing the identical *DR5* expression pattern within a population ($n = 12$). PX, protoxylem pole. Scale bar, 100 μm . (c) Distribution of lateral root primordia (LRP) in seven developmental classes as defined by the *pCYCB1;1::GUS* activity after treatment with 100 μM cedrene for 7 more days. $n = 8$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Total LRP, total number of lateral root primordia including all seven developmental stages. ELR, emerged lateral roots. (d) *pGATA23::nls-GUS* expression in primary root under control and cedrene-treated (100 μM) conditions. Scale bar, 200 μm .

Figure 5. The effects of cedrene on prebranch site formation and *DR5* oscillation. (a and b) Luciferase imaging and quantification of prebranch site number of 3-day-old *pDR5::Luciferase* seedlings grown for 6d on medium supplied with or without 100 μM cedrene. The prebranch site in the newly formed primary root after transfer was measured. The red triangle indicates *pDR5::Luciferase* signal in the OZ, and white arrow indicates prebranch site revealed by persistent *pDR5::Luciferase* signal. Yellow dotted line indicates the position of root tip after 4 d co-cultivation. Scale bar, 1 cm. (c) Quantification of prebranch site number of *pDR5::Luciferase* seedlings within 48 h under control and cedrene-treated (100 μM) conditions. $n \geq 10$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Error bars indicate $\pm$ SD of the mean.

Figure 6. Influence of auxin signaling and transport on root system architecture modified by cedrene. (a) Representative photographs of wild-type (Col-0) and mutant seedlings of auxin signaling (*tir1afb2*, *slr-1/iaa14* and *arf7arf19*) and transport (*aux1-7*, *pin2* and *pin3*) under control and cedrene (100 μM) treatments for 8 d. Scale bar, 2 cm. (b and c) Quantification of emerged lateral root number (b) and primary root elongation (c) of plants shown in (a). $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Error bars indicate $\pm$ SD of the mean.

Figure 7. Effects of the polar auxin transport inhibitor NPA on cedrene-induced primary root elongation and LR formation. (a) Representative photographs of *Arabidopsis* seedlings grown under solvent (Ctrl) and cedrene (100 μM) treatments in

525 presence of varying concentrations of NPA. Scale bar, 2 cm. (b and c) Quantification
526 of emerged lateral root number (c) and primary root elongation (b) of plants shown in
527 (a). $n = 12$. **, $P < 0.01$; *, $P < 0.05$ (Two-way ANOVA). Error bars indicate $\pm$ SD of
528 the mean.

529 **Figure 8.** Expression of auxin efflux transporters in response to cedrene in
530 primary roots. (a-d) Confocal images of *pPIN1:PIN1:GFP* (a), *pPIN2:PIN2:GFP* (b),
531 *pPIN3:PIN3:GFP* (c) and *pPIN7:PIN7:GFP* (d) signal in the primary root tips of 3-
532 day-old seedlings grown with or without cedrene for 6 d. The seedlings were stained
533 with propidium iodide and analyzed by confocal microscopy. Scale bar, 100 μ m. (e-h)
534 Quantification of *pPIN1:PIN1:GFP* (e), *pPIN2:PIN2:GFP* (f), *pPIN3:PIN3:GFP* (g)
535 and *pPIN7:PIN7:GFP* (h) signal intensity at primary root tip shown in (a-d). $n \geq 8$. **,
536 $P < 0.01$; *, $P < 0.05$ (Student's *t*-test). Error bars indicate $\pm$ SD of the mean.

537

Table 1 Volatile compounds produced by *Trichoderma guizhouense* NJAU 4742 after 6 d of growth in 1/2 MS salts agar medium, analyzed by solid-phase microextraction (SPME)- GC-MS

	Metabolite name	Normalized amount of volatile compound (%)
Sesquiterpene s	1,4-Cadinadiene	25.50 ± 4.08
	Cedrene	17.11 ± 4.29
	Dauca-4(11),8-diene	8.66 ± 2.69
	Cis-calamenene	9.31 ± 4.54
	(+)-Cuparene	4.28 ± 1.76
	Copaene	3.57 ± 1.54
	(+)-Acoradiene	2.39 ± 0.57
	g-Muurolene	1.89 ± 0.35
	4,9-Cadinadiene	0.93 ± 0.11
	β-Cubebene	0.74 ± 0.08
	Cycloisolongifolene	0.70 ± 0.34
	Aromandendrene	0.21 ± 0.09
	β-Curcumene	0.19 ± 0.09
other VCs	Benzene, 1,4-diethyl-	17.24 ± 4.29
	3,3-Dimethylphthalide	1.59 ± 0.86
	D-alanine, n-(4-butylbenzoyl)-, isohexyl ester	1.05 ± 0.16
	4-hepten-3-one, 5-ethyl-4-methyl-	1.31 ± 0.84
	6,6-dimethyl-2-vinylidenebicyclo[3.1.1]heptane	0.94 ± 0.71
	Spiro[4.5]dec-6-en-8-one, 1,7-dimethyl-4-(1-methylethyl)-	0.52 ± 0.26
	Benzene, 1-(1,5-dimethylhexyl)-4-methyl-	0.46 ± 0.14
	1-bromo-3,7-dimethyl-2,6-octadiene	0.32 ± 0.13
	Spiro[4.5]dec-8-en-7-ol, 4,8-dimethyl-1-(1-methylethyl)-	0.26 ± 0.11

Normalized amount of volatile compound = (peak area of volatile compound) / (total peak area of all volatile compounds). Compounds were qualitative analyzed on the basis of NIST library (<https://webbook.nist.gov/chemistry/>). Values represent means of three independent determinations.

Parsed Citations

- Bailly A, Groenhagen U, Schulz S, Geisler M, Eberl L, Weisskopf L (2014)** The inter-kingdom volatile signal indole promotes root development by interfering with auxin signalling. *Plant J* 80: 758-771
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bais HP, Fall R, Vivanco JM (2004)** Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. *Plant Physiology* 134: 307-319
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM (2006)** The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology* 57: 233-266
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Beauregard PB, Chai Y, Vamakis H, Losick R, Kolter R (2013)** *Bacillus subtilis* biofilm induction by plant polysaccharides. *Proceedings of the National Academy of Sciences of the United States of America* 110: E1621-E1630
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Benkova E, Michniewicz M, Sauer M, Teichmann T, Seifertova D, Jurgens G, Friml J (2003)** Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* 115: 591-602
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bennett MJ, Marchant A, Green HG, May ST, Ward SP, Millner PA, Walker AR, Schulz B, Feldmann KA (1996)** *Arabidopsis* AUX1 gene: A permease-like regulator of root gravitropism. *Science* 273: 948-950
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Bitas V, McCartney N, Li NX, Demers J, Kim JE, Kim HS, Brown KM, Kang S (2015)** *Fusarium oxysporum* Volatiles Enhance Plant Growth Via Affecting Auxin Transport and Signaling. *Frontiers in Microbiology* 6
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Blilou I, Xu J, Wildwater M, Willemsen V, Paponov I, Friml J, Heidstra R, Aida M, Palme K, Scheres B (2005)** The PIN auxin efflux facilitator network controls growth and patterning in *Arabidopsis* roots. *Nature* 433: 39-44
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- De Rybel B, Vassileva V, Parizot B, Demeulenaere M, Grunewald W, Audenaert D, Van Campenhout J, Overvoorde P, Jansen L, Vanneste S, Moller B, Wilson M, Holman T, Van Isterdael G, Brunoud G, Vuylsteke M, Vernoux T, De Veylder L, Inze D, Weijers D, Bennett MJ, Beeckman T (2010)** A novel aux/IAA28 signaling cascade activates GATA23-dependent specification of lateral root founder cell identity. *Current Biology* 20: 1697-1706
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dharmasiri N, Dharmasiri S, Estelle M (2005)** The F-box protein TIR1 is an auxin receptor. *Nature* 435: 441-445
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Ditengou FA, Muller A, Rosenkranz M, Felten J, Lasok H, van Doorn MM, Legue V, Palme K, Schnitzler JP, Polle A (2015)** Volatile signalling by sesquiterpenes from ectomycorrhizal fungi reprogrammes root architecture. *Nature Communications* 6: 6279
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger S, Grigoriev IV, Kubicek CP (2011)** *Trichoderma*: the genomics of opportunistic success. *Nature Reviews Microbiology* 9: 749-759
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Du Y, Scheres B (2018)** Lateral root formation and the multiple roles of auxin. *Journal of Experimental Botany* 69: 155-167
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Dubrovsky JG, Sauer M, Napsucialy-Mendivil S, Ivanchenko MG, Friml J, Shishkova S, Celenza J, Benkova E (2008)** Auxin acts as a local morphogenetic trigger to specify lateral root founder cells. *Proceedings of the National Academy of Sciences of the United States of America* 105: 8790-8794
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Felten J, Kohler A, Morin E, Bhalerao RP, Palme K, Martin F, Ditengou FA, Legue V (2009)** The Ectomycorrhizal Fungus *Laccaria bicolor* Stimulates Lateral Root Formation in Poplar and *Arabidopsis* through Auxin Transport and Signaling. *Plant Physiology* 151: 1991-2005
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fiddaman PJ, Rossall S (1994)** Effect of substrate on the production of antifungal volatiles from *Bacillus subtilis*. *Journal of Applied Bacteriology* 76: 395-405
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fincheira P, Quiroz A (2018)** Microbial volatiles as plant growth inducers. *Microbiological Research* 208: 63-75
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)
- Fincheira P, Quiroz A (2018)** Microbial volatiles as plant growth inducers. *Microbiol Res* 208: 63-75
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fukaki H, Nakao Y, Okushima Y, Theologis A, Tasaka M (2005) Tissue-specific expression of stabilized SOLITARY-ROOT/IAA14 alters lateral root development in Arabidopsis. Plant Journal 44: 382-395

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fukaki H, Tameda S, Masuda H, Tasaka M (2002) Lateral root formation is blocked by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of Arabidopsis. Plant Journal 29: 153-168

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Galweiler L, Guan CH, Muller A, Wisman E, Mendgen K, Yephremov A, Palme K (1998) Regulation of polar auxin transport by AtPIN1 in Arabidopsis vascular tissue. Science 282: 2226-2230

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Garnica-Vergara A, Barrera-Ortiz S, Munoz-Parra E, Raya-Gonzalez J, Mendez-Bravo A, Macias-Rodriguez L, Francisco Ruiz-Herrera L, Lopez-Bucio J (2016) The volatile 6-pentyl-2H-pyran-2-one from Trichoderma atroviride regulates Arabidopsis thaliana root morphogenesis via auxin signaling and ETHYLENE INSENSITIVE 2 functioning. New Phytologist 209: 1496-1512

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Garnica-Vergara A, Barrera-Ortiz S, Munoz-Parra E, Raya-Gonzalez J, Mendez-Bravo A, Macias-Rodriguez L, Ruiz-Herrera LF, Lopez-Bucio J (2016) The volatile 6-pentyl-2H-pyran-2-one from Trichoderma atroviride regulates Arabidopsis thaliana root morphogenesis via auxin signaling and ETHYLENE INSENSITIVE 2 functioning. New Phytol 209: 1496-1512

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hiltpold I, Turlings TCJ (2008) Belowground chemical signaling in maize: When simplicity rhymes with efficiency. Journal of Chemical Ecology 34: 628-635

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Himanen K, Boucheron E, Vanneste S, Engler JD, Inze D, Beeckman T (2002) Auxin-mediated cell cycle activation during early lateral root initiation. Plant Cell 14: 2339-2351

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hung R, Lee S, Bennett JW (2013) Arabidopsis thaliana as a model system for testing the effect of Trichoderma volatile organic compounds. Fungal Ecology 6: 19-26

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kanchiswamy CN, Mainoy M, Maffei ME (2015) Chemical diversity of microbial volatiles and their potential for plant growth and productivity. Frontiers in Plant Science 6

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kanchiswamy CN, Malnoy M, Maffei ME (2015) Chemical diversity of microbial volatiles and their potential for plant growth and productivity. Front Plant Sci 6: 151

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Laskowski M, Biller S, Stanley K, Kajstura T, Prusty R (2006) Expression profiling of auxin-treated Arabidopsis roots: Toward a molecular analysis of lateral root emergence. Plant and Cell Physiology 47: 788-792

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Laskowski M, Ten Tusscher KH (2017) Periodic lateral root priming: What makes it tick? Plant Cell 29: 432-444

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lavenus J, Goh T, Roberts I, Guyomarc'h S, Lucas M, De Smet I, Fukaki H, Beeckman T, Bennett M, Laplace L (2013) Lateral root development in Arabidopsis: fifty shades of auxin. Trends Plant Sci 18: 450-458

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee HW, Kim MJ, Kim NY, Lee SH, Kim J (2013) LBD18 acts as a transcriptional activator that directly binds to the EXPANSIN14 promoter in promoting lateral root emergence of Arabidopsis. Plant Journal 73: 212-224

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li Y, Shao J, Xie Y, Jia L, Fu Y, Xu Z, Zhang N, Feng H, Xun W, Liu Y, Shen Q, Xuan W, Zhang R (2021) Volatile compounds from beneficial rhizobacteria Bacillus spp. promote periodic lateral root development in Arabidopsis. Plant, cell & environment

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Maffei ME, Gertsch J, Appendino G (2011) Plant volatiles: Production, function and pharmacology. Natural Product Reports 28: 1359-1380

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Malamy JE, Benfey PN (1997) Organization and cell differentiation in lateral roots of Arabidopsis thaliana. Development 124: 33-44

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Marchant A, Bhalarao R, Casimiro I, Eklof J, Casero PJ, Bennett M, Sandberg G (2002) AUX1 promotes lateral root formation by facilitating indole-3-acetic acid distribution between sink and source tissues in the Arabidopsis seedling. Plant Cell 14: 589-597

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mendes R, Garbeva P, Raaijmakers JM (2013) The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. Fems Microbiology Reviews 37: 634-663

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Meng X, Miao Y, Liu Q, Ma L, Guo K, Liu D, Ran W, Shen Q (2019) TgSWO from *Trichoderma guizhouense* NJAU4742 promotes growth in cucumber plants by modifying the root morphology and the cell wall architecture. *Microbial Cell Factories* 18

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Moreno-Risueno MA, Van Norman JM, Moreno A, Zhang J, Ahnert SE, Benfey PN (2010) Oscillating gene expression determines competence for periodic *Arabidopsis* root branching. *Science* 329: 1306-1311

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Muller A, Faubert P, Hagen M, Castell WZ, Polle A, Schnitzler JP, Rosenkranz M (2013) Volatile profiles of fungi - Chemotyping of species and ecological functions. *Fungal Genetics and Biology* 54: 25-33

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Neuteboom LW, Ng JMY, Kuyper M, Clijdesdale OR, Hooykaas PJJ, van der Zaal BJ (1999) Isolation and characterization of cDNA clones corresponding with mRNAs that accumulate during auxin-induced lateral root formation. *Plant Molecular Biology* 39: 273-287

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Noh B, Murphy AS, Spalding EP (2001) Multidrug resistance-like genes of *Arabidopsis* required for auxin transport and auxin-mediated development. *Plant Cell* 13: 2441-2454

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Okushima Y, Fukaki H, Onoda M, Theologis A, Tasaka M (2007) ARF7 and ARF19 regulate lateral root formation via direct activation of LBD/ASL genes in *Arabidopsis*. *Plant Cell* 19: 118-130

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Okushima Y, Overvoorde PJ, Arima K, Alonso JM, Chan A, Chang C, Ecker JR, Hughes B, Lui A, Nguyen D, Onodera C, Quach H, Smith A, Yu GX, Theologis A (2005) Functional genomic analysis of the AUXIN RESPONSE FACTOR gene family members in *Arabidopsis thaliana*: Unique and overlapping functions of ARF7 and ARF19. *Plant Cell* 17: 444-463

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Peret B, Li GW, Zhao J, Band LR, Voss U, Postaire O, Luu DT, Da Ines O, Casimiro I, Lucas M, Wells DM, Lazzerini L, Nacry P, King JR, Jensen OE, Schaffner AR, Maurel C, Bennett MJ (2012) Auxin regulates aquaporin function to facilitate lateral root emergence. *Nature Cell Biology* 14: 991-+

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Perez-Flores P, Valencia-Cantero E, Altamirano-Hernandez J, Pelagio-Flores R, Lopez-Bucio J, Garcia-Juarez P, Macias-Rodriguez L (2017) *Bacillus methylotrophicus* M4-96 isolated from maize (*Zea mays*) rhizosphere increases growth and auxin content in *Arabidopsis thaliana* via emission of volatiles. *Protoplasma* 254: 2201-2213

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH (2013) Going back to the roots: the microbial ecology of the rhizosphere. *Nature Reviews Microbiology* 11: 789-799

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pickett FB, Wilson AK, Estelle M (1990) The aux1 mutation of *Arabidopsis* confers both auxin and ethylene resistance. *Plant Physiology* 94: 1462-1466

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Piechulla B, Lemfack MC, Kai M (2017) Effects of discrete bioactive microbial volatiles on plants and fungi. *Plant Cell and Environment* 40: 2042-2067

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rodriguez-Concepcion M (2006) Early steps in isoprenoid biosynthesis: Multilevel regulation of the supply of common precursors in plant cells. *Phytochemistry Reviews* 5: 1-15

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene signal transduction in *Arabidopsis thaliana*: five novel mutant loci integrated into a stress response pathway. *Genetics* 139: 1393-1409

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sabatini S, Beis D, Wolkenfelt H, Murfett J, Guilfoyle T, Malamy J, Benfey P, Leyser O, Bechtold N, Weisbeek P, Scheres B (1999) An auxin-dependent distal organizer of pattern and polarity in the *Arabidopsis* root. *Cell* 99: 463-472

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Santos Teixeira JA, Ten Tusscher KH (2019) The Systems Biology of Lateral Root Formation: Connecting the Dots. *Mol Plant* 12: 784-803

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schmidt R, Cordovez V, de Boer W, Raaijmakers J, Garbeva P (2015) Volatile affairs in microbial interactions. *Isme Journal* 9: 2329-2335

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schulz S, Dickschat JS (2007) Bacterial volatiles: the smell of small organisms. *Natural Product Reports* 24: 814-842

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Splivallo R, Novero M, Berteaux CM, Bossi S, Bonfante P (2007) Truffle volatiles inhibit growth and induce an oxidative burst in *Arabidopsis thaliana*. New Phytologist 175: 417-424

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Strobel GA, Dirkse E, Sears J, Markworth C (2001) Volatile antimicrobials from *Muscodor albus*, a novel endophytic fungus. Microbiology-Sgm 147: 2943-2950

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Swarup K, Benkova E, Swarup R, Casimiro I, Peret B, Yang Y, Parry G, Nielsen E, De Smet I, Vanneste S, Levesque MP, Carrier D, James N, Calvo V, Ljung K, Kramer E, Roberts R, Graham N, Marillonnet S, Patel K, Jones JDG, Taylor CG, Schachtman DP, May S, Sandberg G, Benfey P, Friml J, Kerr I, Beeckman T, Laplace L, Bennett MJ (2008) The auxin influx carrier LAX3 promotes lateral root emergence. Nature Cell Biology 10: 946-954

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tirranen LS, Gitelson II (2006) The role of volatile metabolites in microbial communities of the LSS higher plant link. Advances in Space Research 38: 1227-1232

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tyc O, Song C, Dickschat JS, Vos M, Garbeva P (2017) The Ecological Role of Volatile and Soluble Secondary Metabolites Produced by Soil Bacteria. Trends Microbiol 25: 280-292

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ulmasov T, Murfett J, Hagen G, Guilfoyle TJ (1997) Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. Plant Cell 9: 1963-1971

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vacheron J, Desbrosses G, Bouffaud ML, Touraine B, Moenne-Loccoz Y, Muller D, Legendre L, Wisniewski-Dye F, Prigent-Combaret C (2013) Plant growth-promoting rhizobacteria and root system functioning. Front Plant Sci 4: 356

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

van Berkel K, de Boer RJ, Scheres B, ten Tusscher K (2013) Polar auxin transport: models and mechanisms. Development 140: 2253-2268

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

van Dam NM, Bouwmeester HJ (2016) Metabolomics in the Rhizosphere: Tapping into Belowground Chemical Communication. Trends in Plant Science 21: 256-265

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Van Norman JM, Xuan W, Beeckman T, Benfey PN (2013) To branch or not to branch: the role of pre-patterning in lateral root formation. Development 140: 4301-4310

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Venturi V, Keel C (2016) Signaling in the Rhizosphere. Trends in Plant Science 21: 187-198

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Werner S, Polle A, Brinkmann N (2016) Belowground communication: impacts of volatile organic compounds (VOCs) from soil fungi on other soil-inhabiting organisms. Appl Microbiol Biotechnol 100: 8651-8665

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xie Y, Wang J, Zheng L, Wang Y, Luo L, Ma M, Zhang C, Han Y, Beeckman T, Xu G, Cai Q, Xuan W (2019) Cadmium stress suppresses lateral root formation by interfering with the root clock. Plant, Cell & Environment 42: 3182-3196

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xuan W, Audenaert D, Parizot B, Moller BK, Njo MF, De Rybel B, De Rop G, Van Isterdael G, Mahonen AP, Vanneste S, Beeckman T (2015) Root cap-derived auxin pre-patterns the longitudinal axis of the *Arabidopsis* root. Current Biology 25: 1381-1388

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xuan W, De Gernier H, Beeckman T (2020) The dynamic nature and regulation of the root clock. Development 147: dev181446

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xuan W, Opdenacker D, Vanneste S, Beeckman T (2018) Long-term in vivo imaging of luciferase-based reporter gene expression in *Arabidopsis* roots. Methods in molecular biology (Clifton, N.J.) 1761: 177-190

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zadnikova P, Petrasek J, Marhavy P, Raz V, Vandenbussche F, Ding Z, Schwarzerova K, Morita MT, Tasaka M, Hejatko J, Van Der Straeten D, Friml J, Benkova E (2010) Role of PIN-mediated auxin efflux in apical hook development of *Arabidopsis thaliana*. Development 137: 607-617

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang H, Kim MS, Krishnamachari V, Payton P, Sun Y, Grimson M, Farag MA, Ryu CM, Allen R, Melo IS, Pare PW (2007) Rhizobacterial volatile emissions regulate auxin homeostasis and cell expansion in *Arabidopsis*. Planta 226: 839-851

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang J, Akcapinar GB, Atanasova L, Rahimi MJ, Przylucka A, Yang D, Kubicek CP, Zhang R, Shen Q, Druzhinina IS (2016) The neutral metallopeptidase NMP1 of *Trichoderma guizhouense* is required for mycotrophy and self-defence. *Environmental Microbiology* 18: 580-597

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang J, Miao Y, Rahimi MJ, Zhu H, Steindorff A, Schiessler S, Cai F, Pang G, Chenthamara K, Xu Y, Kubicek CP, Shen Q, Druzhinina IS (2019) Guttation capsules containing hydrogen peroxide: an evolutionarily conserved NADPH oxidase gains a role in wars between related fungi. *Environmental Microbiology* 21: 2644-2658

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)