1 **Functional genome and microbiome in blood of goats affected by the** 2 **gastrointestinal pathogen *Haemonchus contortus***

3
4 Yonathan Tilahun

5
6 *E-mail addresses:* yonathan.tilahun@langston.edu

7
8 ^a School of Agriculture and Applied Sciences, Langston University, Langston, Oklahoma
9 73050, USA

10 11 **Abstract**

12 The Alpine goat *Capra aegagrus hircus* is parasitized by the barber pole worm
13 (*Haemonchus contortus*). This relationship results in changes that affect the gene
14 expression of the host, the pest, and the microbiome of both. Hematological parameters
15 indicating genes that are expressed and/or the % Composition of abundant and diverse
16 microbial flora are reflective of infestation. We explored the similarity/dissimilarity
17 between and among blood samples of non-infected, infected, infected zoledronic acid-
18 treated, and infected antibody (anti- $\gamma\sigma$ T cells) treated wethers under controlled conditions.
19 We identified responses to barber pole worms using blood-based analysis of transcripts
20 and the microbiome. Seven (7) days post-inoculation (dpi) we identified 7,627 genes
21 associated with different treatment types. Across all treatments we identified fewer raw
22 read counts and a reduced diversity in microbial flora on 7 dpi than in 21 dpi wethers. We
23 also identified that there were differences in % Composition of microbial flora known to
24 be associated with inflammation. This study identifies treatment specific genes, and an
25 increase in microflora abundance and diversity as wethers age post infestation. Further,
26 *Firmicutes/Bacteroidetes* (F/B) ratio reflect metabolic health, based on depression or
27 elevation above thresholds defined by the baseline of non-infected hosts depending on the
28 type of intrusion exhibited by the pest.

29 **Keywords:** Small Ruminants, Gene Expression, Strongylida, Nematode, Parasite
30 Diagnostics

Introduction

Capra aegagrus hircus (the Alpine goat) is parasitized by many strongylid nematodes. Among these the barber pole worm (*Haemonchus contortus*) is particularly important [1]. They are the principal parasites of goats, causing global losses to agriculture estimated at over \$100 billion each year, increasing every year since 1987 [2]. This nematode is a blood-sucking parasite and can remove < 30 μ L of blood/day from the host, inducing anemia and often death.

Haemonchus contortus can prove fatal if untreated in cattle, sheep, and goats [3]. Adult worms are inhibited by goats until the environment becomes favorable. Such circumstances increase susceptibility in kids up to eight weeks after parturition [4]. Even moderate infections can reduce milk production and lead to stunted growth in kids. Other effects include anemia, low packed cell volume (PCV), diarrhea, dehydration, peripheral, and internal fluid accumulation, lower reproductive performance, higher mortality and more frequent illness [5]. Worm egg counts are the primary method to diagnose worm infections before localized production losses [6]. Significant blood loss leads to visual signs of infestation that may be confused with or due to a combined effect from other types of parasites and diseases.

H. contortus produces excretory and secretory products that depress the host immune response [7,8,9]. The host counters through immune responses by its genome and microbiome in a back-and-forth arms race [9]. The specific method that is used to avoid host surveillance is not fully understood, but one theory suggests helminth inflammation inhibition is completed by modulating butyrate biosynthesis [10,8].

The microbiome is composed of bacteria, archeal, viral, and fungal microbial taxa. These may be commensal, mutualistic, or pathogenic, serving roles ranging from beneficial to inconsequential or detrimental [11]. We described our performance and analysis to identify differences in microbiome composition displaying significant differences in abundance between uninfected control wethers, infected only wethers with *H. contortus*, infected

wethers treated with zoledronic acid (ZA), and infected wethers treated with anti- $\gamma\sigma$ T cell antibodies (AB). These different states or treatments will estimate “metabolic health” in small ruminant goat blood microbiomes (GBMs) following infection by the barber pole nematode *H. contortus*. A large array of products exuded from intestinal helminths modulate microbial community growth and metabolism [11]. Also, *H. contortus* compete with naturally occurring flora of the host for energy-rich nutrients or essential minerals [12]. Infection by *H. contortus* is known to impact intestinal physiology by increasing fluid secretion that alters the habitat of healthy bacterial communities [11]. We examine if this impacts the GBM.

Helminth infestations are known to quickly change the metabolic activity of the abomasum in hosts versus unafflicted small ruminants [7,13,8]. An array of structures, cells, and secretions respond to infestation [14-16]. Natural microbiotas provide resources for innate immunity [16]. This typically happens via altered microbiota diversity richness when compared to un-infested animals [10].

The genes expressed by an organism’s genome are major players in how and when the host responds to a parasite. Additionally, the microbiome modulates responses via internal and external environmental cues [17]. Infestation of a host by a parasite interact with surrounding environmental factors to form an intricate web of stimuli and responses by both entities. The microbiome influences many aspects of these diverse ecosystems [18]. A tri-directional interaction is predicted [19], whereby the host depends on its own genome and its vast microbiome content to defend itself against external stressors including parasite intrusions.

Although blood is assumed to be sterile, devoid of other types of cells [20,21], microorganisms often occur in blood without inducing disease [20,21,22]. Here, we describe the blood expressed transcriptomes, the abundance and diversity of resident microorganisms, their phylogenetic affiliations, and their relevance towards a metabolically healthy GBM of both uninfected and infected small ruminant *Capra hircus* wethers with *H. contortus*.

We hypothesize that the microflora composition responds to parasite infestation; therefore, the immune response is expressed through genes exhibited by the host genome and its microbiome. We hypothesized that the host genome and the microbiome respond to invasion by parasites by depressing the *Firmicutes/Bacteroidetes* ratio below threshold. We predicted that host genomes would express more immunological genes and that microfloral composition and populations would change over time.

Objectives

The objectives of the research described here are to identify if blood may be used as a parasite diagnostic tool by: 1) analyzing the different responses to different treatments (uninfected wethers, infected wethers with *H. contortus*, infected wethers treated with zoledronic acid (ZA), and infected wethers treated with anti- $\gamma\sigma$ T cell antibodies (AB)) as determined by transcriptomic and metagenomic analyses; 2) identifying genes expressed from these treatments after seven (7) days post inoculation (dpi); 3) comparing microbial flora abundance and diversity between seven 7 dpi and 21 dpi using operational taxonomic units (OTUs); 4) and by determining if there are significant differences between *Firmicutes/Bacteroidetes* (F/B) ratios as an indication of infection in order to determine differences of “metabolic health” in wethers.

Materials and Methods

Ethical statement

The treatment of animals in our research abided by the guidelines of the Langston University Institutional Animal Care and Use Committee (LUACUC) Approval # 2018-14.

Animals and treatments

Forty (40) Alpine wethers (114.2±0.92 d of age; 19.4±0.33 kg BW at the start of experiment) being raised in indoor pens at the Langston University farm were used. The animals were checked for fecal egg counts (FECs) and confirmed that they were nematode-parasite free. The animals were allocated randomly to four (4) groups of 10 animals each, and two (2) or three (3) animals from each group were assigned to one of the four (4) pens. All animals were allowed to acclimatize to pens and feeders for daily supplies of 200 g concentrate pellet per animal composed of 500 g of ground grass (50%) and alfalfa (50%) hays. The treatment groups were as:

Table 1. Experimental Set-up. Treatment Group (1-4) for infection L3 *H. contortus* (+) or non-infection *H. contortus* (-), with (+) or without (-) treatment type (Zoledronic acid injection or $\gamma\delta$ T depletion).

Group	L3 <i>H. contortus</i> infection	Zoledronic acid	$\gamma\delta$ T depletion
1	-	-	-
2	+	-	-
3	+	+	-
4	+	-	+

On the first (1) day prior to the L3 infection, the anti- $\gamma\delta$ T cells antibody were administered intravenously. ZA were administered intravenously 7 days prior to and 0, 7, and 14 days after the L3 infection. At the beginning of the experiment all kids except Group 1 were given 10,000 *H. contortus* infective larvae (L3; hatched and isolated from feces being collected from LU goats) by gavage. Five animals from each group were euthanized on seven (7) days post inoculation (dpi) for sampling and the other five (5) animals were euthanized 21 dpi.

Blood sample collection and processing

Blood samples were collected from five goats in four treatment groups (No infection, Infection, Infection ZA inject, and Infection AB inject), following 7 dpi. Blood samples

included red blood cells, white blood cells (total and differential), hemoglobin, platelets, and plasma protein, from the jugular vein in EDTA tubes. Quality assurance/quality control (QA/QC) parameters resulted in blood samples from 19/20 cDNA libraries that were used from samples collected 7 dpi. The cDNA libraries were sequenced on an illumina RNA-Seq Next-generation sequencing (NGS) instrument and filtered and normalized using Partek® Flow® software suites.

Also, methods of identifying naturally occurring microbial flora in nontreated and treated wethers were identified. Analytics for QA/QC for high-throughput barcoded illumina MiSeq NGS sequencing of 16S rRNA, resulted in 18/20 samples that were obtained for 7 dpi and 20/20 samples for 21 dpi.

Total RNA Purification

Total RNA was collected from blood samples using a modified TRIzol reagent procedure. The blood samples were lysed by using ice chilled TRIzol Reagent. The broken cells were homogenized by pipetting up and down many times. After transferring the homogenized and broken cells into Eppendorf, Chloroform was added to the lysed cells. It gave three layers. The upper aqueous layer contained extracted RNA, an interphase contained DNA, and proteins were dissolved in the bottom red organic layer. The pH was kept at around 4 for RNA purification, which held RNA in the aqueous phase preferentially. After centrifugation, the upper aqueous layer was pipetted out carefully and isopropanol was added to precipitate the RNA. Again, the RNA was precipitated by centrifugation to get RNA pellets which were washed with 70% ethanol (made with DEPC treated water), air-dried and suspended in DEPC treated (RNase free) water. Extracted RNA was quantified to calculate yield by a NanoDrop™ 2000 spectrophotometer.

A. Lysate Preparation from Blood

The cDNA libraries were constructed by initial first strand synthesis using the Protocol for Non-directional RNA-seq Workflows and NEBNext® Ultra™ II RNA First Strand Synthesis Module (E7771), according to a modified manufacturer's protocol as follows:

Input Amount Requirement

A 100 ng total RNA was quantified after the purification. The protocol was optimized for approximately 200 nt RNA inserts. The protocol was optimized using Universal Human Reference Total RNA.

RNA Fragmentation and Priming

The fragmentation and priming reactions were assembled on ice in a nuclease-free tube by adding the Fragmentation and Priming Mix for a total volume of 10 µL. They were mixed thoroughly by pipetting. The samples were placed in a thermocycler and incubated at 94 °C. The tube was immediately transferred to ice and First Strand cDNA Synthesis was begun immediately.

First Strand cDNA Synthesis Reaction

The first strand synthesis reaction was assembled on ice by adding components to the fragmented and primed RNA for a total volume of 20 µL. The reaction was mixed thoroughly by pipetting. The sample was incubated in a preheated thermocycler with the heated lid set at $\geq 80^{\circ}\text{C}$ as follows: Step 1: 10 minutes at 25°C ; Step 2: 15 minutes at 42°C ; Step 3: 15 minutes at 70°C ; and Step 4: Hold at 4°C . We then proceeded directly to Swift Biosciences™ ACCEL-NGS® 1S PLUS DNA LIBRARY KIT: Single, Dual Combinatorial and Unique Dual Indexing and prepared the DNA Libraries as follows:

Denaturation

216

217 Due to the short incubation time of the denaturation step, all of the reagents of the Adaptase
218 Reaction Mix were pre-assembled and placed on ice. The thermocycler was pre-heated to
219 95° C. The fragmented DNA sample was transferred to a 0.2 mL PCR tube and the volume
220 of the sample adjusted to a final volume of 15 µL using Low EDTA TE, if it was necessary.
221 The samples were placed in the thermocycler, programmed at 95 ° C for 2 minutes with lid
222 heating ON. Upon completion, the tube(s) were placed on ice immediately for 2 minutes.
223 We then proceeded directly to the Adaptase step to preserve the maximum amount of
224 ssDNA substrate.

225

226 **Adaptase**

227

228 The Adaptase Thermocycler Program was loaded on the thermocycler and paused at the
229 first step to pre-heat to 37 ° C until all samples were loaded. Twenty-five (25) µL of the
230 pre-assembled Adaptase Reaction Mix was added to each PCR tube containing a 15 µL
231 DNA sample and mixed by pipetting or gentle vortexing until homogeneous, after which
232 they were spun down. The samples in the thermocycler were run at the following
233 parameters with the lid heating ON. The thermocycler Program followed: 37° C, 15
234 minutes; 95° C, 2 minutes; and a 4° C hold.

235

236 **Extension**

237

238 The Extension Thermocycler Program was loaded on the thermocycler and paused at the
239 first step to pre-heat to 98° C until all samples were loaded. Forty-seven (47) µL of the
240 Adaptase Reaction was added, using reagents in the order listed in the manufacturers
241 protocol. The sample was mixed by pipetting or gentle vortexing until homogenous and
242 spun down. The samples were placed in the thermocycler and the following program was
243 run, with lid heating ON. The thermocycler Program followed: 98° C, 30 seconds; 63° C,
244 15 seconds; 68° C, 5 minutes; and a 4° C hold. Each sample was transferred to a 1.5 mL

tube and clean up the Extension Reaction using beads and freshly prepared 80% ethanol was completed.

Ligation

Twenty (20) μ L of the pre-mixed Ligation Reaction Mix was placed in a new PCR tube containing 20 μ L of the Post-Extension eluate. Samples were mixed by pipetting or gently vortexing until homogenous and spun down. The samples were placed in the thermocycler programmed at 25 ° C for 15 minutes with lid heating ON, followed by a 4° C hold. Each sample was transferred to a 1.5 mL tube and clean up the Ligation Reaction using beads and freshly prepared 80% ethanol was completed.

Indexing PCR

Five (5) μ L of the appropriate indexed adapter primer(s) were added directly to each sample. Twenty-five (25) μ L of the already pre-mixed Indexing PCR Reaction Mix were added to each PCR tube containing 25 μ L of sample, using reagents in the order listed in the manufacturers protocol. Samples were mixed by pipetting or gently vortexing until homogenous and spun down. The samples were placed in the thermocycler and the following program run with the proper recommended PCR cycles, with lid heating ON. The thermocycler Program followed: 98° C, 30 seconds; PCR Cycles: 98° C, 10 seconds; 60° C, 30 seconds; 68° C, 60 seconds; 4° C hold. Each sample was transferred to a 1.5 mL tube and clean-up of the Indexing PCR Reaction using beads and freshly prepared 80% ethanol was completed.

Size Selection/Clean-up

The following protocol was used for each clean-up step, substituting the correct Sample Volume, Bead Volume, and Elution Volume based on the table provided for each section.

The magnetic beads were at room temperature and vortexed the beads to homogenize the suspension before use. Each Sample Volume was transferred to a 1.5 mL tube. The specified Bead Volume was added to each sample, mixed by vortex, and quickly spun on a tabletop microcentrifuge. The samples were incubated for 5 minutes at room temperature (off the magnet) and placed on a magnet rack until the solution cleared and a pellet formed (~2 minutes). The supernatant was removed and discarded without disturbing the pellet (less than 5 µL may have been left behind). A freshly prepared 80% ethanol solution (500 µL) was added to the samples while still on the magnetic rack. Using care not to disturb the pellet, the samples were incubated for 30 seconds, and then the ethanol solution was removed. This step was repeated once more for a second wash with the 80% ethanol solution. The samples were spun in a tabletop microcentrifuge and placed back on the magnetic rack. Residual ethanol solution was removed from the bottom of the tube. The specified Elution Volume of Low TE buffer was added and the pellet re-suspended. Samples were mixed by pipetting up and down until homogenous. Incubation of the samples was completed at room temperature for 2 minutes off the magnet, then placed on the magnet. The entire eluate was transferred to a new 0.2 ml PCR tube. The eluate, without containing the magnetic beads (indicated by brown coloration in the eluate), was ensured to be pure by pipetting the samples into a new tube, placing on a magnet, and transferring the eluate again.

Microbiome DNA Isolation

Microbiome DNA was collected from blood samples using a modified Microbiome DNA Isolation Kit from NORGEN BIOTEK CORP. (Thorold, ON, Canada). The procedures are as follows:

A. For Samples Collected using Norgen's Preservation Devices

Transfer of 0.5 mL of whole blood sample to a 2 mL DNAase-free microcentrifuge tube were completed. Lysis Buffer was added and the tube vortexed. One hundred (100) µL of Lysis Additive was added to the mixture and vortexed briefly. The mixture was incubated

at 65° C for 5 minutes. The tubes were centrifuged for 2 minutes at 20,000 x g (~14,000 RPM). The supernatant was transferred to a DNAase-free microcentrifuge tube. One hundred (100) µL of Binding Buffer was added and mixed before incubation on ice. Centrifugation was completed for 2 minutes at 20,000 x g (~ 14,000 RPM). A pipette was used to transfer up to 700 µL of supernatant into a 2mL DNAase-free microcentrifuge tube. An equal volume of 70% ethanol was added to the lysate collected above and vortexed.

B. Binding to Column

A spin column with one of the provided collection tubes was assembled. Seven hundred (700) µL of the clarified lysate with ethanol was added onto the column and centrifuged for 1 minute at 10,000 x g (~ 10,000 RPM). The flowthrough was discarded, and the spin column reassembled with the collection tube. The step with the remaining volume of lysate mixture was repeated.

C. Column Wash

Five hundred (500) µL of Binding Buffer was added to the column and centrifuged for 1 minute at 10,000 x g (~ 10,000 RPM). The flowthrough was discarded, and the spin column reassembled with its collection tube. Five hundred (500) µL of Wash Solution was applied to the column and centrifuged for 1 minute at 10,000 x g (~ 10,000 RPM). The flowthrough was discarded and the spin column reassembled with its collection tube. Repeat the previous two steps. The column was centrifuged for 2 minutes at 20,000 x g (~ 14,000 RPM) in order to thoroughly dry the resin. The collection tube was discarded.

D. DNA Elution

The column was placed into a fresh 1.7 mL Elution tube provided with the kit. Fifty (50) µL of Elution Buffer was added to the column. Centrifugation was completed for 1 minute at 200 x g (~ 2,000 RPM), followed by a 1-minute spin at 20,000 xg (~14,000 RPM). An additional elution was performed using 50 µL of the Elution Buffer.

E. Storage of DNA

The purified genomic DNA was stored at -80° C.

Sequencing

Quality Assurance/Quality Control of bar-coded sequence prepped samples of cDNA were completed by The Genomics Core Facility at Oklahoma State University (Stillwater, OK). The Genomics Core Facility at Oklahoma State University (Stillwater, OK) completed the cDNA library sequencing with an illumina RNASeq NGS instrument.

Quality Assurance/Quality Control of bar-coded sequence prepped samples were completed and sequenced for 16S rRNA metagenomics of blood samples by Swift Biosciences™ (Ann Arbor, MI USA) using an illumina MiSeq NGS instrument.

Computational Analysis

Several methods were utilized to conduct bioinformatic analysis of the obtained sequence data. For gene expression analyses, we used the Partek® Flow® software suites pipelines that include, but are not limited to the STAR algorithm, Normalization, and the gene set differential analysis method (GSA).

Preliminary analyses included Qiime2 analysis for the metagenomic or microbiome analysis. We present here, the results obtained utilizing the Kraken pipeline through the Partek® Flow® software suites, a start-to-finish software analysis solution for next generation sequencing data applications. Inflammation comparisons were based on the ratio of *Firmicutes/Bacteroidetes* (*F/B* Ratios) that are evident as well as the reduction of microbial diversity.

Results

We analyzed gene expression in samples from 19/20 wethers (7 dpi). The microbiome portion resulted in analysis of 38/40 wethers separated into 7 dpi (18/20 samples) and 21 dpi (20/20 samples). Initially, ten (10) individuals were not infected with *H. contortus*, 10 received infection with *H. contortus* only, 10 were infected with *H. contortus* and received ZA injections, and 10 were infected with *H. contortus* and received AB injection. As mentioned in this study blood samples were collected 7 dpi, where 19/20 cDNA libraries passed QA/QC analytics for cDNA gene expression analysis. QA/QC analytics of high-throughput barcoded illumina MiSeq NGS sequencing for 16S rRNA metagenomics, resulted in the 18/20 samples being obtained for 7 dpi and the 20/20 samples for 21 dpi. The sequences used for the purposes of this study, involved the inclusion of gene expression and metagenomic analyses. The initial design for the study is shown for 20 individuals in Table 2 (Supplementary Documents).

Haemonchus contortus infection affected the metabolic system of wethers. A 95% confidence interval ($* = p < 0.05$) indicated likely significant changes in the expression of at least 184 genes (affecting treatment comparisons of samples that were infected with zoledronic acid injection versus infection with antibody injection Fig. 7) when using a numeric triad of p-values for no infection versus infection only, no infection versus infection zoledronic acid injection, and no infection versus infection with antibody injection. The hierarchical clustering/heat map (Figs. 1) generated after selection of three of the most highly significant specific differentially analyzed genes, depicts colored tiles showing differences in genomic features in the integration site data sets from the blood samples of each subject ($n = 19$). They indicate the intensity and direction of any significant departures from the distributions of random controls varying to a degree depending on the type of treatment and the subject [19].

Fig.1. Hierarchical clustering/heat map. Three selected genes (identified at the bottom of the image) following differential analysis using GSA counts of 19/20 samples (identified on the right side of the image belonging to the described treatment group identified on the left side of the image) indicate the most likely significant genes ($p < 0.05$) affecting samples when comparing No Infection samples to those that were only infected with the *H. contortus* pathogen, injected with zoledronic acid (ZA) or injected with antibodies (AB) and then infected with the *H. contortus* pathogen on 7 dpi.

The following Volcano plots (Figs. 2-7) show genes that were identified as being downregulated ($* = < -2$), having no fold change (NC: $* = > -2$, $* = < 2$), and/or being upregulated ($* = > 2$). The Volcano plots illustrates the related distribution of genes out of 7627 expressed, depending on subject and treatment type. The distribution of genes in this case were based on comparisons using a numeric triad of p-values for no infection versus infection only, no infection versus infection zoledronic acid (ZA) injection, and no

infection versus infection with antibody (AB) injection. Significance was based on a 95% confidence interval ($* = p < 0.05$) where fold changes of downregulated, NC, and upregulated gene distributions of treatment groups were assessed. In Fig. 2 a comparison of no infection versus infection only subjects, out of 7627 genes that were expressed, 523 genes were significant in expression during downregulation, NC, and upregulation. For the comparison of no infection vs infection ZA (Fig. 3), 290 genes that were significant in expression during downregulation, NC, and upregulation. In Fig. 4 289 genes were significant in expression during downregulation, NC, and upregulation when comparing no infection vs infection AB. In Fig. 5 examines the comparison of infection only vs infection ZA, resulting in the identification of 338 genes that were expressed with significance in expression during downregulation, NC, and upregulation. The Volcano plot for Fig. 6 indicates 275 genes that were expressed in fold changes for downregulated, NC, and upregulated genes that were significant.

Fig. 2. No Infection vs Infection only. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing No Infection samples to infection only samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 3. No Infection vs Infection ZA inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing No Infection samples to Infection ZA inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 4. No Infection vs Infection AB inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing No Infection samples to infection AB inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 5. Infection only vs Infection ZA inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing infection only samples to Infection ZA inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig.6. Infection only vs Infection AB inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing infection only samples to infection AB inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

483

484

485 **Fig. 7. Infection ZA inject vs Infection AB inject.** A Volcano plot for 19/20 blood
486 samples indicating likely significant expression of genes ($* = p < 0.05$) based on
487 treatment type of *Capra hircus* following STAR alignment and GSA differential analysis
488 for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change
489 indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing Infection ZA inject samples to infection AB
490 inject samples when expressed against a Numeric Triad of P-values for No Infection vs
491 Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs
492 infection AB inject (blue).
493
494

The average number of raw reads for metagenomic analyses of blood samples using Kraken is illustrated in Figs. 8 and 9. The distribution of raw read counts indicate the apparent difference in number between 7 dpi and 21 dpi subjects. There were 43.61 raw reads for metagenomic analysis of blood collected on 7 dpi as compared to 2,638.80 raw reads for metagenomic analysis of blood samples at 21 dpi.

Fig. 8. Distribution of raw read counts for all samples at 7 dpi. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Fig. 9. Distribution of raw read counts for all samples at 21 dpi. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

After QA/QC, we collected Alpha diversity reports for 18 individuals for 7 dpi using Shannon and Simpson distribution indices (Fig. 10) [12]. The Alpha diversity reports for the Shannon and Simpson distribution indices (Table 3; Supplementary Documents) were

evaluated. Our results indicate that *H. contortus* has a significant effect on species-level microbial diversity for blood that is infected and injected with ZA using both indices [P(T<=t) first-tail 0.012 and P(T<=t) second-tail 0.023] (Table 4).

Fig. 10. Shannon and Simpson indices of 7dpi. Alpha diversity reports depicted by Shannon and Simpson indices distributions for 18 samples of 7dpi (loss of sample numbers 17 and 50 due to poor quality) [23-25] where single samples and the variation of microbes in them are identified. The first number before each sample ID identifying the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Table 4. Significant values of 7 dpi Shannon and Simpson indices. Statistical comparison of a) Shannon and b) Simpson distribution indices for Alpha diversity reports of 7 dpi richness and diversity of microbial flora in host *Capra hircus* wethers.

a) Shannon index

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection ZA inject	n1 = 1.06	5	0.562	0.012	0.023
	n2 = 2.47	5	0.718		

b) Simpson index

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
	n1 = 0.389	5	0.0685		

No Infection vs Infection ZA inject	n2 = 0.782	5	0.0160	0.012	0.023
--	------------	---	--------	--------------	--------------

We also collected Alpha diversity reports for 20 individuals for 21 dpi (Table 4; Supplementary Documents; Fig. 11).

Fig. 11. Shannon and Simpson indices of 21dpi. Alpha diversity reports depicted by Shannon and Simpson distribution indices for 20 samples of 21dpi. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Species richness and diversity for non-infected controls, infected wethers, infected ZA injected wethers, and infected AB injected wethers indicated significant differences between 7dpi and 21 dpi microbial composition. This contrasts with 21 dpi where there are no significant differences when comparisons are made between non-infected and infected wethers (ZA/AB injected or not). This supports literature where the total bacterial load increases over time after infection with *H. contortus* [8].

Infection likely has a broad range of quantitative biological effects on each host. Our investigation of the composition of microbial flora shows a more similar pattern to other studies that support the notion as to why a host is better suited to withstand intrusion by external threats. The richer and more diverse the microbiota, the better the host may combat external pathogens [12]. The Alpha diversity report, depicting Shannon and Simpson distribution indices, identifies that treatments for 21 dpi (Table 5; Supplementary Documents), when compared with each other do not show significance. The non-infected wethers appear to have developed a pronounced, non-infected version of an abundant and diverse microbial flora. Thus, when infected 21 dpi wethers are statistically compared to non-infected 21 dpi controls for abundance and diversity using Alpha diversity reports with Shannon and Simpson distribution indices, there are no significant differences.

However, when comparing Shannon and Simpson Alpha indices between 7 dpi and 21 dpi (Table 6) there are significant differences. The Shannon indices for 7 dpi vs 21 dpi t-Test: Two-Sample Assuming Unequal Variances results showed that there are significant differences between the different treatment groups. There are differences between “non-infected” wethers based on age alone. Table 5 shows that there are significant differences between comparisons between “non-infected” wethers for 7 dpi and those wethers that were “non-infected” for 21dpi, only “infected” for 21 dpi, “infected” with ZA injection for 21 dpi, and “infected” with an AB injection for 21 dpi. Therefore, it is evident that the richness and diversity for microbial flora in blood changes over time, regardless of the treatment. Further evidence supports this when examining wethers subjected to “infection” only for 7 dpi as compared to those with “non-infected” 21 dpi following inoculation, “infection” with ZA injection for 21dpi, and “infection” with an AB injection for 21 dpi. This is further support that despite the condition, as the age of the wethers increases (dpi), there are significant changes in the richness and diversity of microbial flora in the blood.

Table 6. Statistically different comparison results of 7 dpi versus 21 dpi. Statistically different comparison results of 7 dpi versus 21 dpi a) Shannon and b) Simpson distribution indices for Alpha diversity reports of abundance and diversity for microbial flora in host *Capra hircus* wethers.

586

587 a) Shannon index 7 dpi vs 21 dpi t-Test: Two-Sample Assuming Unequal Variances

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) one-tail	P(T<=t) two-tail
No Infection 7 dpi vs No Infection 21 dpi	n1 = 1.05 n2 = 2.91	5 5	0.562 2.56	0.029	0.057
No Infection 7 dpi vs Infection only 21 dpi	n1 = 1.05 n2 = 3.37	5 5	0.562 2.83	0.015	0.031
No Infection 7 dpi vs Infection ZA inject 21 dpi	n1 = 1.05 n2 = 3.65	5 5	0.562 3.86	0.020	0.040
No Infection 7 dpi vs Infection AB inject 21 dpi	n1 = 1.05 n2 = 3.67	5 5	0.562 1.12	0.001	0.003
Infection only 7 dpi vs No Infection 21 dpi	n1 = 0.707 n2 = 2.91	5 5	0.714 2.56	0.017	0.035
Infection only 7 dpi vs Infection only 21 dpi	n1 = 0.707 n2 = 3.37	5 5	0.714 2.83	0.010	0.020
Infection only 7 dpi vs Infection ZA inject 21 dpi	n1 = 0.707 n2 = 3.65	5 5	0.714 3.86	0.014	0.028
	n1 = 0.707	5	0.714	0.001	0.001

Infection only 7dpi vs Infection AB inject 21 dpi	n2 = 3.67	5	1.12		
Infection ZA inject 7dpi vs Infection AB inject 21 dpi	n1 = 2.47	5	0.718		
	n2 = 3.67	5	1.12	0.041	0.082
Infection AB inject 7dpi vs Infection AB inject 21 dpi	n1 = 1.77	3	1.60		
	n2 = 3.67	5	1.12	0.047	0.094

588

Simpson index 7dpi vs 21 dpi t-Test: Two-Sample Assuming Unequal Variances

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) one-tail	P(T<=t) two-tail
No Infection 7 dpi vs Infection AB inject 21 dpi	n1 = 0.390	5	0.0685		
	n2 = 0.845	5	0.0172	0.007	0.013
Infection only 7 dpi vs No Infection 21 dpi	n1 = 0.196	5	0.0522		
	n2 = 0.699	5	0.132	0.017	0.034
Infection only 7 dpi vs Infection only 21 dpi	n1 = 0.196	5	0.0522		
	n2 = 0.758	5	0.130	0.011	0.022
Infection only 7 dpi vs Infection ZA inject 21 dpi	n1 = 0.196	5	0.0522		
	n2 = 0.762	5	0.161	0.017	0.034

Infection only 7dpi vs Infection AB inject 21 dpi	n1 = 0.196	5	0.0522	0.001	0.002
---	------------	---	--------	--------------	--------------

Gene amplicons for 16S rRNA Relative Abundance profiles are illustrated for 18 individuals on 7dpi (Fig. 12). The blood samples are dominated by 31 most abundant operational taxonomic units (OTUs).

Fig. 12. Relative Abundance of OTUs on 7 dpi for 18 samples. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

We identify that Fig. 12, illustrates the Relative Abundance of OTU profiles on 7 dpi that follow the approximate total “Mean” profile percentage of the most abundant phylum being *Proteobacteria* (~84.16%) (Table 7).

Table 7. OTU percentage composition. OTU percentage composition of the most prevalent Phylum after 7 days post inoculation (*Proteobacteria*).

Seven (7) days post inoculation		
Sample Number	<i>Proteobacteria</i> (%) Composition	Treatment
1-8_E1_S123	76.0	No Infection
1-28_F1_S205	94.6	No Infection

1-44_E2_S131	54.2	No Infection
1-52_E3_S139	94.3	No Infection
1-54_F2_S213	61.0	No Infection
Mean No Infection	76.0	
2-2_E4_S147	73.7	Infection only
2-13_F3_S221	94.4	Infection only
2-20_E5_S155	97.2	Infection only
2-51_F5_S237	67.8	Infection only
2-58_F4_S229	98.3	Infection only
Mean Infection only	86.3	
3-6_F6_S246	90.2	Infection ZA inject
3-27_F8_S262	95.4	Infection ZA inject
3-31_F7_S254	45.5	Infection ZA inject
3-38_F9_S270	95.2	Infection ZA inject
3-57_E6_S164	93.5	Infection ZA inject
Mean ZA inject	84.0	
4-25_E8_S180	95.8	Infection AB inject
4-30_E7_S172	94.2	Infection AB inject
4-33_E9_S188	93.5	Infection AB inject
Mean AB inject	94.5	

Table 8. Statistical comparison of *Proteobacteria* of 7 dpi. Statistical comparison of *Proteobacteria* (% Composition) after seven (7) dpi in host *Capra hircus* wethers.

Proteobacteria (% Composition) Seven Days Post Inoculation

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection AB inject	n1 = 76.026	5	346.0	0.046	0.091
	n2 = 94.51	3	1.374

Fig. 13. Relative Abundance of OTUs on 21 dpi for 20 samples. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject). Gene amplicons for 16S rRNA Relative Abundance of OTU profiles are illustrated for 20 individuals on 21 dpi (Fig. 13). The blood samples are dominated by 36 most abundant OTUs.

The 21 dpi blood samples accounted for a much more quantified and rich pool of OTUs compared to that of 7 dpi blood samples.

Table 9. OTU percentage composition of 21 dpi. OTU percentage composition of the most prevalent Phylums after 21 days post inoculation (*Proteobacteria*, *Bacteroidetes*, *Actinobacteria*, and *Firmicutes*).

21 days post inoculation					Treatment
Sample Number	<i>Proteobacteria</i>	<i>Bacteroidetes</i>	<i>Actinobacteria</i>	<i>Firmicutes</i>	
	(%Composition)	(%Composition)	(%Composition)	(%Composition)	
1-14_H9_S143	97.6	0.360	0.460	0.350	No Infection
1-32_H10_S151	92.5	0.500	0.800	0.290	No Infection
1-34_H3_S380	36.5	17.4	20.5	8.93	No Infection
1-45_H8_S134	45.4	15.6	18.1	6.86	No Infection
1-62_H7_S126	65.1	9.72	9.46	5.13	No Infection

Mean No Infection	67.4	8.71	9.86	4.31	
2-12_H1_S363	33.1	15.6	21.4	13.7	Infection only
2-24_H6_S381	32.6	15.9	20.4	14.3	Infection only
2-36_G8_S344	32.4	16.0	20.7	14.0	Infection only
2-42_H5_S373	92.0	2.00	2.61	1.15	Infection only
2-66_G6_S328	60.9	11.6	10.8	6.16	Infection only
Mean Infection only	50.2	12.2	15.2	9.87	
3-5_G10_S360	34.6	15.0	20.0	13.1	Infection ZA inject
3-29_G5_S319	34.1	15.7	20.6	12.9	Infection ZA inject
3-43_H4_S364	94.6	0.410	1.040	0.410	Infection ZA inject
3-46_H2_S371	29.8	17.2	21.7	13.6	Infection ZA inject
3-53_G4_S311	34.8	15.3	18.8	15.3	Infection ZA inject
Mean ZA inject	45.6	12.7	16.4	11.1	
4-10_G3_S303	91.5	0.380	0.860	0.390	Infection AB inject
4-19_G9_S352	40.5	14.5	19.2	10.9	Infection AB inject
4-23_G2_S295	30.9	16.5	21.1	13.9	Infection AB inject
4-39_G7_S336	22.9	16.5	20.7	23.6	Infection AB inject
4-49_G1_S287	55.7	10.1	15.6	7.24	Infection AB inject
Mean AB inject	48.3	11.6	15.5	11.2	

628

629 There is clear evidence that greater amounts of difference are present when prolonged

630 exposure to *H. contortus* infection are examined (7 dpi No Infection versus 21 dpi of all

631 other treatments). *Proteobacteria* (Table 10: No Infection 7 dpi vs Infection only 21 dpi;

No Infection 7 dpi vs Infection ZA inject 21 dpi; and No Infection 7 dpi vs Infection AB inject 21 dpi) shows the most significant values for relative abundance of OTUs.

Table 10. Statistical comparison of *Proteobacteria* between 7 dpi and 21 dpi. Statistical comparison of *Proteobacteria* (% Composition) after seven (7) dpi versus 21 dpi of host *Capra hircus* wethers.

Proteobacteria (% Composition) Seven dpi versus 21 dpi

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection 7 dpi vs Infection only 21 dpi	n1 = 76.0	5	346.0	0.018	0.036
	n2 = 50.2	5	696.0		
No Infection 7 dpi vs Infection ZA inject 21 dpi	n1 = 76.0	5	346.0	0.020	0.040
	n2 = 45.6	5	756.0		
No Infection 7 dpi vs Infection AB inject 21 dpi	n1 = 76.0	5	346.0	0.009	0.019
	n2 = 48.3	5	732.0		

When a comparison of % Composition were made among 21 dpi treatments, there were P(T<=t) first-tail values that were significant/near significant for *Firmicutes* (Table 14). This implies agreement with the occurrence of inflammation [17] during parasite intrusion.

Table 11. Statistical comparison of *Proteobacteria* of 21 dpi. Statistical comparison of *Proteobacteria* (% Composition) after 21 dpi of host *Capra hircus* wethers.

Proteobacteria (% Composition) 21 dpi versus 21 dpi

Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection only	n1 = 67.4	5	747.0	0.170	0.340
	n2 = 50.2	5	696.0		
	n1 = 67.4	5	747.0	0.122	0.244

No Infection vs Infection ZA inject	n2 = 45.6	5	756.0		
No Infection vs Infection AB inject	n1 = 67.4	5	747.0	0.150	0.299
	n2 = 48.3	5	733.0		

Table 12. Statistical comparison of *Bacteroidetes* of 21 dpi. Statistical comparison of *Bacteroidetes* (% Composition) after 21 dpi of host *Capra hircus* wethers

<i>Bacteroidetes</i> (% Composition) 21 dpi versus 21 dpi					
Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection only	n1 = 8.71	5	65.3	0.232	0.464
	n2 = 12.2	5	35.9		
No Infection vs Infection ZA inject	n1 = 8.71	5	65.3	0.213	0.430
	n2 = 12.7	5	48.0		
No Infection vs Infection AB inject	n1 = 8.71	5	65.3	0.265	0.530
	n2 = 11.6	5	37.0		

Table 13. Statistical comparison of *Actinobacteria* of 21 dpi. Statistical comparison of *Actinobacteria* (% Composition) after 21 dpi of host *Capra hircus* wethers.

<i>Actinobacteria</i> (% Composition) 21 dpi versus 21 dpi					
Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection only	n1 = 9.86	5	87.9	0.184	0.369
	n2 = 15.2	5	68.5		
No Infection vs Infection ZA inject	n1 = 9.86	5	87.9	0.142	0.283
	n2 = 16.4	5	75.2		
No Infection vs Infection AB inject	n1 = 9.86	5	87.9	0.174	0.349
	n2 = 15.5	5	71.7		

Table 14. Statistical comparison of *Firmicutes* of 21 dpi. Statistical comparison of *Firmicutes* (% Composition) after 21 dpi of host *Capra hircus* wethers.

<i>Firmicutes</i> (% Composition) 21 dpi versus 21 dpi					
Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection only	n1 = 4.31	5	15.1	0.0618	0.1237
	n2 = 9.87	5	35.4		
No Infection vs Infection ZA inject	n1 = 4.31	5	15.1	0.0368	0.0736
	n2 = 11.1	5	36.3		
No Infection vs Infection AB inject	n1 = 4.31	5	15.1	0.076	0.153
	n2 = 11.2	5	73.2		

The “Mean” profile percentages of the most abundant phyla shown in Table 9 for “No Infection” were *Proteobacteria* (~67.4%), followed by *Actinobacteria* (~9.86%), *Bacteroidetes* (~8.71%), and *Firmicutes* (~4.31%) of all OTUs. The “Mean” profile percentages of the most abundant phyla shown in Table 9 for “Infection only” were *Proteobacteria* (~50.2%), followed by *Actinobacteria* (~15.2%), *Bacteroidetes* (~12.2%), and *Firmicutes* (~9.87%) of all OTUs. The “Mean” profile percentages of the most abundant phyla shown in Table 9 for infection ZA inject were *Proteobacteria* (~45.6%), followed by *Actinobacteria* (~16.4%), *Bacteroidetes* (~12.7%), and *Firmicutes* (~11.1%) of all OTUs. The “Mean” profile percentages of the most abundant phyla shown in Table 9 for Infection AB inject were *Proteobacteria* (~48.3%), followed by *Actinobacteria* (~15.5%), *Bacteroidetes* (~11.6%), and *Firmicutes* (~11.2%) of all OTUs.

Table 15 shows the most statistically significant treatment values for *Firmicutes* % Composition/*Bacteroidetes* % Composition (*F/B*) ratios. When comparing No Infection versus the other treatments, statistical comparisons revealed that two treatments indicated significant differences (* = $p < 0.05$): No Infection 21 dpi vs Infection ZA inject 21 dpi and No Infection 21 dpi vs Infection AB inject 21 dpi for the P(T<=t) first-tail.

Table 15. Statistical comparison of *F/B* ratios. Statistical comparison of *F/B* ratios % Composition after 21 dpi in host *Capra hircus* wethers.

<i>F/B</i> ratio 21 dpi versus 21 dpi					
Sample Comparisons	Mean	Sample Size (n)	Sample Variance (s)	P(T<=t) first-tail	P(T<=t) second-tail
No Infection vs Infection ZA inject	n1 = 0.606	5	0.044	0.015	0.030
	n2 = 0.898	5	0.010		
No Infection vs Infection AB inject	n1 = 0.606	5	0.044	0.035	0.070
	n2 = 0.948	5	0.085		

The non-infected controls, after 21 dpi, revealed both less *Bacteroidetes* % Composition and *Firmicutes* % Composition than did all the other treatments. There is a 0.494 ratio for the “Mean” *Bacteroidetes* % Composition to *Firmicutes* % Composition, when non-infected controls are examined. The “Infection only” treatments had a 0.809 ratio for the “Mean” *Bacteroidetes* % Composition to *Firmicutes* % Composition. The treated ZA inject had a ratio of 0.874 and the treated AB inject had a ratio of 0.966 “Mean” *Bacteroidetes* % Composition to *Firmicutes* % Composition. This supports that there are measurable outcomes that can differentiate metabolic healthy wethers from parasitically compromised wethers based on changes in *Bacteroidetes* % Composition to *Firmicutes* % Composition [17].

Discussion

To characterize the nematode-infected host-transcriptome, host-microbiome, we examined the distribution of 19-7 dpi transcriptome sequencing (RNA-Seq; n = 40) samples and

microbial flora from 18- seven (7) dpi and 20- 21 dpi samples of four treatment types of 16S rRNA sequencing (MiSeq, n= 80). We focused on whole-blood or buffy coat derived extractions of mbDNA [26,27,28] to discriminate among non-infected, metabolically healthy controls and infected types. We performed microbial assignments using Kraken [29] and transcriptome analysis using STAR [30], completing differential analysis using GSA [31]. Our data identified 7627 genes that were expressed and shows differences in association across treatment types on 7 dpi and 21 dpi in specific microbiota profiles.

The evidence determined in this study shows that blood-based microbial DNA (mbDNA) can be used to discriminate between a non-infected, metabolically healthy or a *Haemonchus contortus*-attacked *Capra hircus wethers* [32]. The evidence also shows that the microbiome changes in abundance and diversity over age (dpi), even within the interval of a few days [33,34,35,36,37], further supporting that microbial flora abundance and diversity is associated with metabolic health or the indication of non-infection [44].

Factors, including antibiotics, can change the composition of microbiota [38,39,40,41] often destroying the composition of beneficial microbes along with pathological ones. Thus, causing dysbiosis or the development of unwanted microbes [41,42,43, 44]. Thus, the residual infection condition is still evident despite antibiotic or other type of treatment, indicated by inflammatory results or an increase in ratio of inflammation causing microbial flora. We identified evidence that the composition of inflammatory microbiota increases with zoledronic acid or anti- $\gamma\delta$ T cells treatment. With a greater increase being exhibited by anti- $\gamma\delta$ T cells treatment.

Unlike the results described by a previous study where it was surmised that infection by *H. contortus* did not affect caprine microbial diversity [8], we identified that in goat blood samples there were likely significant differences based especially on treatment types. There is also likely significance in evidence that microbial abundance and diversity differences are dependent on age (days post inoculation).

Our results imply that there are differences in gene expression as subjects become older or are exposed to the environment longer. Attributes of the different treatment types show that genes are expressed in response to the *H. contortus* presence. All pointing towards the implication of *H. contortus* to effectively change and disrupt the internal habitat of the host.

The host health may be determined based on *Firmicutes/Bacteroidetes* (*F/B*) ratios. The *F/B* Ratio is estimated by utilizing the lowest and highest values of the reference range (uninfected host) for individual organisms [44]. A high *F/B* ratio may be related to increased caloric extraction from food, fat deposition and lipogenesis, impaired insulin sensitivity, and increased inflammation. Low *Firmicutes/Bacteroidetes* (*F/B*) ratio, is an indicator of dysbiosis indicated by a decreased diversity of the microbiome compared to healthy cohorts. *F/B* is considered as “low” when the value falls below a threshold [44]. Therefore, “metabolic health” is determined by the actions taken by the parasite. If the parasite yields an *F/B* ratio that is depressed compared to the reference range of harmful intrusion, it may be postulated that the host does not have adequate means of defending itself, thus the parasite load is depleting host resources. Alternatively, if the parasite yields an *F/B* ratio that is elevated compared to the reference range of harmful intrusion, it indicates that the host is being hyperactive or inflamed as a response to the parasite burden.

Conclusions

The metabolic systems affected warrant further investigation to identify specific pathways where significant changes have resulted based on being exposed to infection. Furthermore, the development of computational algorithms for correlation of microbial abundance and diversity are warranted. The authors conjecture that blood samples are shown here to be a possible means to indicate *H. contortus* infection based on detection of microbial flora abundance and diversity as well as in gene expression profiles. Correlations can be drawn on statistical levels of microbial flora for this specific type of inflammation. The specificity in the range of microbial flora, dependent on the age of the *wethers*, can indicate the occurrence of depleting resources or inflammation due to *H. contortus* infection. In other words, this implies *H. contortus* does effectively change and disrupt the internal habitat

health of the host and the effects are measurable. The development of a standard laboratory diagnostic procedure using blood microbiota to detect gastrointestinal infection with *H. contortus* is the ideal course of action.

Availability of data and materials

The metagenomic data have been deposited with links to BioProject accession number PRJNA612987 in the NCBI BioProject database (<https://www.ncbi.nlm.nih.gov/bioproject/>).

The gene expression data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus [45] and are accessible through GEO Series accession number GSE169607 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE169607>).

Acknowledgements

The author is thankful to coinvestigators Zaisen Wang, Qunhui Yang, and Jessica Quijada-Pinango and the Director of the American Institute for Goat research (AIGR: School of Agriculture and Applied Sciences, Langston University, Langston, Oklahoma, USA), Mrs. Kelly Kyle and Mrs. Xiaowen Wang (Partek Incorporated). The Relationship Between the Microbiome and Internal Parasitism in Goats is supported by the Evans Allen Program Grant at the Agricultural Research and Extension Center at Langston University.

Literature Cited

- [1] Laing R, T. Kikuchi, A. Martinelli, I. J. Tsai, R. N. Beech, E. Redman, N. Holroyd, D. J. Bartley, H. Beasley, C. Britton, D. Curran, E. Devaney, A. Gilabert, M. Hunt, F. Jackson, S. L. Johnston, I. Kryukov, K. Li, A. A. Morrison, A. J. Reid, N. Sargison, G. I. Saunders, J. D. Wasmuth, A. Wolstenholme, M. Berriman, J. S. Gilleard, and J. A. Cotton. 2013. The genome and transcriptome of *Haemonchus contortus*, a key model parasite for drug and vaccine discovery. *Genome Biology*, 14:R88. <http://genomebiology.com/2013/14/8/R88>.
- [2] Sasser, J.N. and Freckman, D.W. 1987. A World Perspective on Nematology: The Role of the Society. In: Veech, J.A. and Dickson, D.W., Eds., *Vistas on Nematology*, Hyattsville, Maryland, 7-20.
- [3] Mamun, Md. A. A, M. Sandeman, P. Rayment, P. Brook-Carter, E. Scholes, N. Kasinadhuni, D. Piedrafita, and A. R. Greenhill. 2020. Variation in gut bacterial composition is associated with *Haemonchus contortus* parasite infection of sheep. *Animal Microbiome*, 2:3. <https://doi.org/10.1186/s42523-020-0021-3>
- [4] Mekonnen, S. M. 2007. Helminth Parasites of Sheep and Goats in Eastern Ethiopia: Epidemiology and Anthelmintic Resistance and its Management. Doctoral thesis Swedish University of Agricultural Sciences.
- [5] Leite-Browning, M. L. 2006. *Haemonchus contortus* (Barber Pole Worm) Infestation in Goats. www.aces.edu/urban.

- [6] Barber's pole worm, wormboss. www.wormboss.com/au/Sheep-goats/worms/roundworm-barbers-pole-worm.php. 2020, accessed Oct. 22, 2020.
- [7] Weinstock, J. V. and D. E. Elliott. 2009. Helminths and the IBD hygiene hypothesis. *Inflammatory Bowel Diseases*, 15:128-133.
- [8] Li, R. W., W. Li, J. Sun, P. Yu, R. L. Baldwin, and J. F. Urban. 2016. The effect of helminth infection on the microbial composition and structure of the caprine abomasal microbiome. *Scientific Reports*, 6:20606. 10.1038/srep20606
- [9] Gadahi, J. A., B. Yongqian, M. Ehsan, Z. Zhang, S. Wang, R. F. Yan, L. X. Xu, and Li, X. R. 2016. *Haemonchus contortus* excretory and secretory proteins (HcESPs) suppress functions of goat PBMCs in vitro. *Oncotarget*, 7(24):35670–35679. DOI:10.18632/oncotarget.9589
- [10] Li, C. J. and R. W. Li. 2008. Butyrate induced cell cycle arrest in bovine cells through targeting gene expression relevant to DNA replication apparatus. *Gene Relulation System Biology*, 2:113-23.
- [11] Zaiss, M. M., and Harris, N. L. 2016. Interactions between the intestinal microbiome and helminth parasites. *Parasite immunology*, 38(1):5–11. <https://doi.org/10.1111/pim.12274>
- [12] A. K. Pathak. 2017. Nutritional Bases to Control Gastrointestinal Parasites of Livestock. *Dairy and Veterinary Science Journal*, 4(2): 555632. doi: 10.19080/JDVS.2017.04.555632
- [13] Li, R. W., S. Wu, W. Li, Y. Huang, and L. C. Gasbarre. 2011. Metagenome plasticity of the bovine abomasal microbiota in immune animals in response to *Ostertagia ostertagi* infection. *PLoS One*, 6, e24417.

- [14] Morales-de la Nueza, C. Rodriguez, M. Santana, L. E. Hernandez-Castellano, T. Nino, R.J. Reyes-Cacon, I. Moreno-Indies, N. Castro, and J. Capote. 2012. Morphological measurements and indexes as a tool to support molecular genetic studies: an example in Canary Islands, *Journal of Applied Animal Research*, 40:3, 2015-221, DOI: 10.1080/09712119.2012.658062
- [15] Powell, N. and T.T. MacDonald. 2017. Recent advances in gut immunology. *Parasite Immunology*, 39:e12430. DOI: 10.1111/pim.12430
- [16] Ayaz, M. M., M. M. Nazir, N. Samad, M. Zubair, M. Hanif, M. Aziz, A. S. Sheikh, A. Akbar, A. Waheed, A. Zaman and N. Mahmood. 2018. Parasitism in Goats: Husbandry Management, Range Management, Gut Immunity and Therapeutics, Goat Science, Sándor Kukovics, *IntechOpen*, DOI: 10.5772/intechopen.74203
- [17] Belkaid, Y., and T. W. Hand. 2014. Role of the microbiota in immunity and inflammation. *Cell*, 157(1):121–141. <https://doi.org/10.1016/j.cell.2014.03.011>
- [18] Wagg, C., Schlaeppli, K., Banerjee, S. et al. 2019. Fungal-bacterial diversity and microbiome complexity predict ecosystem functioning. *Nature Communications*, 10:4841. <https://doi.org/10.1038/s41467-019-12798-y>
- [19] Zheng, D., Liwinski, T. and E. Elinav. 2020. Interaction between microbiota and immunity in health and disease. *Cell Research*, 30:492–506. <https://doi.org/10.1038/s41422-020-0332-7>
- [20] Castillo, D.J., R.F. Rifkin, D.A. Cowan, and M. Potgieter. 2019. The Healthy Human Blood Microbiome: Fact or Fiction? *frontiers in Cellular and Infection Microbiology*, 9(148):1-12. doi:10.3389/fcimb.2019.00148

- [21] Scarsella, e., M. Sandri, S.D. Monego, D. Licastro, and B.Stefanon. 2020. Blood Microbiome: A New Marker of Gut Microbial Population in Dogs? *Veterinary Sciences*, 7(198):1-13. doi:10.3390/vetsci7040198
- [22] Qiu, J., H. Zhou, Y. Jing, and C. Dong. 2019. Association between blood microbiome and type 2 diabetes mellitus: A nested case-control study. *Journal of Clinical Laboratory Analysis*, 33:e22842. <https://doi.org/10.1002/jcla.22842>
- [23] Johnson, K. V. -A. and P. W. J. Burnet. 2016. Microbiome: Should we diversify from diversity? *Gut Microbes*, 7(6):455-458. doi: 10.1080/19490976.2016.1241933
- [24] DeJong, TM. 1975. A comparison of three diversity indices based on their components of richness and evenness. *Oikos*, 26:222-7. <http://dx.doi.org/10.2307/3543712>
- [25] Li, H., T Li, D. E. Beasley, P. Hedenec, Z. Xiao, S. Zhang, J. Li, Q. Lin, and X. Li. 2016. Diet Diversity Is Associated with Beta but not Alpha Diversity of Pika Gut Microbiota. *Frontiers in Microbiology*, 7:1169. doi. 10.3389/fmicb.2016.01169
- [26] Bettgowda, C. et al. 2014. Detection of circulating tumor DNA in early- and late-stage human malignancies. *Scientific Translations in Medicine*, 6:224ra24.
- [27] Clark, T. A. et al. 2018. Analytical Validation of a Hybrid Capture-Based Next Generation Sequencing Clinical Assay for Genomic Profiling of Cell-Free Circulating Tumor DNA. *Journal of Molecular Diagnostics*, 20:686-702.
- [28] Poore, G. D., E. Kopylova, Q. Zhu, C. Carpenter, S. Fraraccio, S. Wandro, T. Kosciolk, S. Janssen, J. Metcalf, S. J. Song, J. Kanbar, S. Miller-Montgomery, R. Heaton, R. McKay, S.P. Patel, A.D. Swafford, & R. Knight. 2020. Microbiome analyses of blood and tissues suggest cancer diagnostic approach. *Nature*, 579(7800), 567–574. <https://doi.org/10.1038/s41586-020-2095-1>

[29] Wood, D.E., J. Lu, and B. Langmead. 2019. Improved metagenomic analysis with Kraken 2. *Genome Biology*, 20:257. <https://doi.org/10.1186/s13059-019-1891-0>

[30] Dobin, A. and T.R. Gingeras. 2015. Mapping RNA-seq Reads with STAR. *Current protocols in bioinformatics*, 51:11.14.1–11.14.19. <https://doi.org/10.1002/0471250953.bi1114s51>

[31] Mathur, R., D. Rotroff, J. Ma, et al. 2018. Gene set analysis methods: a systematic comparison. *BioData Mining*, 11: 8. <https://doi.org/10.1186/s13040-018-0166-8>

[32] Gregory, D. P., E. Kopylova, Q. Zhu, C. Carpenter, S. Fraraccio, S. Wandro, T. Kosciolk, S. Janssen, J. Melcalf, S. J. Song, J. Kanbar, S. Miller-Montgomery, R. Heaton, r. McKay, S. P. Patel, A. d. Swafford, and R. Knight. 2020. Microbiome analysis of blood and tissues suggest cancer diagnostic approach. *Nature*, 579(7800):567-574. doi:10.1038/s41586-020-2095-1

[33] McDonald D, E. Hyde, J.W. Debelius, J.T. Morton, A. Gonzalez, G. Ackermann, A.A. Aksenov, B. Behsaz, C. Brennan, Y. Chen, L. DeRight Goldasich, P.C. Dorrestein, R.R. Dunn, A.K. Fahimipour, J. Gaffney, J.A. Gilbert, G. Gogul, J.L. Green, P. Hugenholtz, G. Humphrey, C. Huttenhower, M.A. Jackson, S. Janssen, D.V. Jeste, L. Jiang, S.T. Kelley, D. Knights, T. Kosciolk, J. Ladau, J. Leach, C. Marotz, D. Meleshko, A.V. Melnik, J.L. Metcalf, H. Mohimani, E. Montassier, J. Navas-Molina, T.T. Nguyen, S. Peddada, P. Pevzner, K.S. Pollard, G. Rahnavard, A. Robbins-Pianka, N. Sangwan, J. Shorenstein, L. Smarr, S.J. Song, T. Spector, A.D. Swafford, V.G. Thackray, L.R. Thompson, A. Tripathi, Y. Vázquez-Baeza, A. Vrbanc, P. Wischmeyer, E. Wolfe, Q. Zhu, The American Gut Consortium, and R. Knight. 2018. American Gut: an open platform for citizen science microbiome research. *mSystems* 3:e00031-18. doi:10.1128/mSystems.00031-18.

- [34] G. Falony, M. Joossens, S. Vieira-Silva, J. Wang, Y. Darzi, K. Faust, A. Kurilshikov, M.J. Bonder, M. Valles-Colomer, D. Vandeputte, R.Y. Tito, S. Chaffron, L. Rymenans, C. Verspecht, L. De Sutter, G. Lima-Mendez, K. D'hoë, K. Jonckheere, D. Homola, R. Garcia, E.F. Tigchelaar, L. Eeckhaut, J. Fu, L. Henckaerts, A. Zhernakova, C. Wijmenga, and J. Raes. 2016. Population-level analysis of gut microbiome variation. *Science* 352:560–564. doi:10.1126/science.aad3503.
- [35] A. Zhernakova, A. Kurilshikov, M.J. Bonder, E.F. Tigchelaar, M. Schirmer, T. Vatanen, Z. Mujagic, A.V. Vila, G. Falony, S. Vieira-Silva, J. Wang, F. Imhann, E. Brandsma, S.A. Jankipersadsing, M. Joossens, M.C. Cenit, P. Deelen, M.A. Swertz, LifeLines cohort study, R.K. Weersma, E.K.M. Feskens, M.G. Netea, D. Gevers, D. Jonkers, L. Franke, Y.S. Aulchenko, C. Huttenhower, J. Raes, M.H. Hofker, R.J. Xavier, C. Wijmenga, and J. Fu. 2016. Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity. *Science* 352:565–569. doi:10.1126/science.aad3369.
- [36] Rinninella E, Raoul P, Cintoni M, et al. 2019. What is the Healthy Gut Microbiota Composition? A Changing Ecosystem across Age, Environment, Diet, and Diseases. *Microorganism*, 2019;7(1):14. doi:10.3390/microorganisms7010014
- [37] Huang, S., N. Haiminen, A.P. Carrieri, R. Hu, L. Jiang, L. Parida, B. Russell, C. Allaband, A. Zarrinpar, Y. Vázquez-Baeza, P. Belda-Ferre, H. Zhou, H.C. Kim, A.D. Swafford, R. Knight, and Z.Z. Xu. 2020. Human Skin, Oral, and Gut Microbiomes Predict Chronological Age. *mSystems*, 5(1), e00630-19. <https://doi.org/10.1128/mSystems.00630-19>
- [38] Goodrich, J.K., J.L. Waters, A.C. Poole, J.L. Sutter, O. Koren, R. Blekhman, M. Beaumont, W. Van Treuren, R. knight, J. T. Bell, T.D. Spector, A.G. Clark, and R.E. Ley. 2014. Human genetics shape the gut microbiome. *Cell*, 2014, 159(4):789-799.

- [39] Ley, R.E., F. Bäckhed, P. Turnbaugh, C.A. Lozupone, R.D. Knight, and J.I. Gordon. 2005. Obesity alters gut microbial ecology. *Proceedings of the National Academy of Science*, 102(31):11070-11075.
- [40] Turnbaugh, P.J., M. Hanady, T. Yatsunencko, B.L. Cantarel, A. Duncan, R. E. Ley, M.L. Sogin, W.J. Jones, B.A. Roe, J.P. Affourtity, M. Egholm, B. Henrissat, A.C. Heath, R. Knight, and J.I. Gordon. 2009. A core gut microbiome in obese and lean twins. *Nature*, 457(7228):480-484.
- [41] Hasan, N. and H. Yang. 2019. Factors affecting the composition of the gut microbiota, and its modulation. *PeerJ*, 7:e7502. doi:10.7717/peerj.7502
- [42] Klingensmith, N.J. and C.m. Coopersmith. 2016. The Gut as the Motor of Multiple Organ Dysfunction in Crytical Illness. *Critical Care Clinics*, 32(2):203-212
- [43] Gee, X., C. Ding, W. Zhao, L. Xu, H. Tian, J. Gong, M. Zhu, J. Li, and N. li. 2017. Antibiotics-induced depletion of mice microbiota induces change in host secretonin biosynthesis and intestinal motility. *Journal of Translational Medicine*, 15(1):13.
- [44] Rowin, J., Y. Xia, B. Jung, and J. Sun. 2017. Gut inflammation and dysbiosis in human motor neuron disease. *Physiological reports*, 5(18), e13443. <https://doi.org/10.14814/phy2.13443>
- [45] Edgar, Ron et al. 2002. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic acids research*, 30(1): 207-10. doi:10.1093/nar/30.1.207

Supplementary Documents

1003

1004 **Table 2. Detailed Experimental Set-up.** The Group number, Treatment type, identifying
1005 Tag, days post inoculation (dpi), Age in days, and Body Weight (BW) in kg.

Group	Treatment	Tag/Sample ID	dpi	Age, days	BW, kg
1	No Infection	8_E1_S123	7	118	21.05
1	No Infection	28_F1_S205	7	116	17.9
1	No Infection	44_E2_S131	7	115	18.95
1	No Infection	52_E3_S139	7	113	17.75
1	No Infection	54_F2_S213	7	114	22.8
2	Infection only	13_F3_S221	7	118	22.9
2	Infection only	2_E4_S147	7	120	19.05
2	Infection only	20_E5_S155	7	117	16.8
2	Infection only	58_F4_S229	7	100	17.75
2	Infection only	51_F5_S237	7	113	16.65
3	Infection ZA inject	6_F6_S246	7	119	17.65
3	Infection ZA inject	31_F7_S254	7	116	17.7
3	Infection ZA inject	27_F8_S262	7	116	20.9
3	Infection ZA inject	38_F9_S270	7	115	20.35
3	Infection ZA inject	57_E6_S164	7	96	19.7
4	Infection AB inject	17_F10_S18	7	117	16.8
4	Infection AB inject	25_E8_S180	7	116	19.95
4	Infection AB inject	33_E9_S188	7	116	19.8
4	Infection AB inject	30_E7_S172	7	116	19.7
4	Infection AB inject	50_E10_S9	7	114	21.65

1006

1007 **Table 3. Alpha diversity report for 7 dpi Shannon and Simpson index.**

1008

Sample name	Shannon index	Simpson index	dpi	treatment
1-8_E1_S123	1.05	0.360	7	No Infection
1-28_F1_S205	0.190	0.0747	7	No Infection
1-44_E2_S131	2.05	0.772	7	No Infection
1-52_E3_S139	0.492	0.254	7	No Infection

1-54_F2_S213	1.49	0.487	7	No Infection
Mean No Infection	1.06	0.3894		
2-13_F3_S221	0.1607	0.061	7	Infection only
2-2_E4_S147	1.40	0.401	7	Infection only
2-20_E5_S155	0.0866	0.0198	7	Infection only
2-58_F4_S229	0.0535	0.0133	7	Infection only
2-51_F5_S237	1.828	0.486	7	Infection only
Mean Infection only	0.707	0.196		
3-6_F6_S246	1.58	0.591	7	Infection ZA inject
3-31_F7_S254	1.61	0.800	7	Infection ZA inject
3-27_F8_S262	2.86	0.823	7	Infection ZA inject
3-38_F9_S270	3.52	0.940	7	Infection ZA inject
3-57_E6_S164	2.77	0.757	7	Infection ZA inject
Mean Infection ZA inject	2.47	0.782		
4-25_E8_S180	0.317	0.099	7	Infection AB inject
4-33_E9_S188	2.64	0.733	7	Infection AB inject
4-30_E7_S172	2.35	0.659	7	Infection AB inject
Mean Infection AB inject	1.770	0.497		

Table 5. Alpha diversity report for 21 dpi Shannon and Simpson index.

Sample name	Shannon index	Simpson index	dpi	treatment
1-14_H9_S143	0.240	0.0643	21	No Infection
1-32_H10_S151	2.64	0.729	21	No Infection
1-45_H8_S134	3.83	0.901	21	No Infection
1-34_H3_S380	4.20	0.911	21	No Infection
1-62_H7_S126	3.63	0.892	21	No Infection
Mean No Infection	2.91	0.699		
2-12_H1_S363	3.97	0.917	21	Infection only
2-24_H6_S381	4.21	0.925	21	Infection only
2-36_G8_S344	4.71	0.949	21	Infection only

2-42_H5_S373	0.461	0.113	21	Infection only
2-66_G6_S328	3.52	0.886	21	Infection only
Mean Infection only	3.37	0.758		
3-5_G10_S360	4.68	0.952	21	Infection ZA inject
3-29_G5_S319	4.72	0.950	21	Infection ZA inject
3-43_H4_S364	0.145	0.0438	21	Infection ZA inject
3-53_G4_S311	4.38	0.929	21	Infection ZA inject
3-46_H2_S371	4.31	0.934	21	Infection ZA inject
Mean Infection ZA inject	3.65	0.762		
4-10_G3_S303	2.76	0.789	21	infection AB inject
4-19_G9_S352	4.66	0.950	21	infection AB inject
4-23_G2_S295	4.83	0.958	21	infection AB inject
4-39_G7_S336	3.62	0.886	21	infection AB inject
4-49_G1_S287	2.51	0.644	21	infection AB inject
Mean Infection AB inject	3.68	0.845		

1013

1014

Fig.1. Hierarchical clustering/heat map. Three selected genes (identified at the bottom of the image) following differential analysis using GSA counts of 19/20 samples (identified on the right side of the image) belonging to the described treatment group identified on the left side of the image) indicate the most likely significant genes ($p < 0.05$) affecting samples when comparing No Infection samples to those that were only infected with the *H. contortus* pathogen, injected with zoledronic acid (ZA) or injected with antibodies (AB) and then infected with the *H. contortus* pathogen on 7 dpi.

Fig. 2. No Infection vs Infection only. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing No Infection samples to infection only samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 4. No Infection vs Infection AB inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing No Infection samples to infection AB inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 5. Infection only vs Infection ZA inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes ($* = p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated ($* = < -2$), no change (NC; $* = > -2$, $* = < 2$), and upregulated ($* = > 2$) gene distribution when comparing infection only samples to Infection ZA inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig.6. Infection only vs Infection AB inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes (* = $p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated (* = < -2), no change (NC; * = > -2 , * = < 2), and upregulated (* = > 2) gene distribution when comparing infection only samples to infection AB inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 7. Infection ZA inject vs Infection AB inject. A Volcano plot for 19/20 blood samples indicating likely significant expression of genes (* = $p < 0.05$) based on treatment type of *Capra hircus* following STAR alignment and GSA differential analysis for transcript sequences of 7627 identified genes expressed on 7 dpi. The fold change indicates downregulated (* = < -2), no change (NC; * = > -2 , * = < 2), and upregulated (* = > 2) gene distribution when comparing Infection ZA inject samples to infection AB inject samples when expressed against a Numeric Triad of P-values for No Infection vs Infection only (green), No Infection vs Infection ZA inject (red) and No Infection vs infection AB inject (blue).

Fig. 8. Distribution of raw read counts for all samples at 7 dpi. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Fig. 9. Distribution of raw read counts for all samples at 21 dpi. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Fig. 10. Shannon and Simpson indices of 7dpi. Alpha diversity reports depicted by Shannon and Simpson indices distributions for 18 samples of 7dpi (loss of sample numbers 17 and 50 due to poor quality) [23-25] where single samples and the variation of microbes in them are identified. The first number before each sample ID identifying the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Fig. 11. Shannon and Simpson indices of 21dpi. Alpha diversity reports depicted by Shannon and Simpson distribution indices for 20 samples of 21dpi. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Fig. 12. Relative Abundance of OTUs on 7 dpi for 18 samples. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).

Fig. 13. Relative Abundance of OTUs on 21 dpi for 20 samples. The first number before each sample ID identifies the treatment type (1 = No infection, 2 = Infection only, 3 = Infection ZA inject, and 4 = Infection AB inject).