Choppakatla et al., April 2021-preprint copy- BioRxiv

Linker histone H1.8 inhibits chromatin-binding of condensins and DNA topoisomerase II to tune chromosome length and individualization

Pavan Choppakatla¹, Bastiaan Dekker², Erin E. Cutts³, Alessandro Vannini^{3,4}, Job Dekker^{2,5} and Hironori Funabiki^{1,*}

¹Laboratory of Chromosome and Cell Biology, The Rockefeller University, New York, NY 10065, USA.

²Program in Systems Biology, Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, Worcester, MA 01605, USA.

³Division of Structural Biology, The Institute of Cancer Research, London SW7 3RP, UK.

⁴Fondazione Human Technopole, Structural Biology Research Centre, 20157 Milan, Italy

⁵Howard Hughes Medical Institute, Chevy Chase, MD 20815, USA.

*Correspondence: funabih@rockefeller.edu

Summary

DNA loop extrusion by condensins and decatenation by DNA topoisomerase II (topo II) are thought to drive mitotic chromosome compaction and individualization. Here, we reveal that the linker histone H1.8 antagonizes condensins and topo II to shape mitotic chromosome organization. *In vitro* chromatin reconstitution experiments demonstrate that H1.8 inhibits binding of condensins and topo II to nucleosome arrays. Accordingly, H1.8 depletion in *Xenopus* egg extracts increased condensins and topo II levels on mitotic chromatin. Chromosome morphology and Hi-C analyses suggest that H1.8 depletion makes chromosomes thinner and longer through shortening the average loop size and reducing the DNA amount in each layer of mitotic loops. Furthermore, excess loading of condensins and topo II to chromosomes by H1.8 depletion causes hyper-chromosome individualization and dispersion. We propose that condensins and topo II are essential for chromosome individualization, but their functions are tuned by the linker histone to keep chromosomes together until anaphase.

Keywords

Chromosome compaction, chromatin, mitosis, linker histone, nucleosome, condensin, DNA topoisomerase, Hi-C, *Xenopus*

Introduction

Genomic DNA in eukaryotes is compacted by orders of magnitude over its linear length. The extent and mode of the packaging change between interphase and mitosis to support the cell cycle dependent functions of the DNA. While loosely packed DNA allows efficient decoding of genetic

information in interphase, mitotic compaction of DNA enables efficient distribution of genetic information to daughter cells. In addition, chromosome individualization during mitosis ensures that all duplicated chromosomes are independently moved by microtubules and are equally distributed to daughter cells. Despite these common functional requirements of mitotic

chromosomes and evolutionary conservation of major known regulators of mitotic chromosome structures, the size and shape of chromosomes vary among species and developmental stages. For example, rod-like individual mitotic chromosomes can be readily visualized in fission yeast *Schizosaccharomyces pombe*, but not in budding yeast *Saccharomyces cerevisiae* (Guacci et al., 1994; Umesono et al., 1983). During early embryogenesis in *Xenopus* and *C. elegans*, mitotic chromosome lengths become shorter (Ladouceur et al., 2015; Micheli et al., 1993). The mechanistic basis of mitotic chromosome shape regulation remains largely speculative (Heald and Gibeaux, 2018).

Classical experiments on mitotic chromosomes indicated that DNA is organized into loops around a central protein scaffold (Earnshaw and Laemmli, 1983; Paulson and Laemmli, 1977). Two major chromosome-associated ATPases play pivotal roles in mitotic chromatid formation: DNA topoisomerase II (topo II) and the structural maintenance of chromosomes (SMC) family complex, condensin (Kinoshita and Hirano, 2017; Kschonsak and Haering, 2015). Data obtained with chromosome conformation capture assays are consistent with the model that mitotic chromosomes are arranged in a series of loops organized by condensins (Elbatsh et al., 2019; Gibcus et al., 2018; Naumova et al., 2013). Vertebrate cells express two forms of the condensin complex, condensin I and condensin II (Hirano et al., 1997; Ono et al., 2003). Condensin I is loaded onto chromatin exclusively during mitosis, whereas condensin II retains access to chromosomes throughout the cell cycle (Hirota et al., 2004; Ono et al., 2004; Walther et al., 2018). Condensin II plays a role in maintaining chromosome territories in *Drosophila* nuclei (Bauer et al., 2012; Rosin et al., 2018) and drives sister chromatid decatenation by topo II (Nagasaka et al., 2016). It has been proposed that condensin II acts first in prophase to anchor large outer DNA loops, which are further branched into shorter inner DNA loops by condensin I (Gibcus et al., 2018). This proposal is consistent with their localization determined by super-resolution microscopy (Walther et al., 2018). In chicken DT40 cells, condensin II drives the helical positioning of loops around a centrally located axis thus controls the organization of long distance interactions (6- 20 Mb), whereas condensin I appears to control shorter distance interactions (Gibcus et al., 2018). This organization of the condensin I and II loops is also consistent with their roles in maintaining lateral and axial

compaction respectively (Bakhrebah et al., 2015; Green et al., 2012; Samejima et al., 2012). In *Xenopus* egg extracts, in the presence of wildtype condensin I levels, condensin II depletion does not appear to change mitotic chromosome length, suggesting a reduced role for condensin II on these chromosomes (Shintomi and Hirano, 2011).

The prevailing model suggests that mitotic chromatin loops are formed by the dynamic loop extrusion activity of condensins (Alipour and Marko, 2012; Nasmyth, 2001; Riggs, 1990), although the molecular details of the process remain unclear (Banigan and Mirny, 2020; Cutts and Vannini, 2020; Datta et al., 2020). Single molecule experiments using purified recombinant yeast and human condensin complexes demonstrated ATP-dependent motor activity and loop extrusion by yeast and human condensins (Ganji et al., 2018; Kong et al., 2020; Terakawa et al., 2017). Condensin dependent loop extrusion in a more physiological *Xenopus* extract system has also been shown (Golfier et al., 2020). *In silico* experiments further suggest that a minimal combination of loop extruders (like condensin) and strand passage activity (such as topo II) can generate well resolved rod-like sister chromatids from entangled, interphase like DNA fibers (Goloborodko et al., 2016a).

These demonstrations of loop extrusion by purified condensins and cohesin were all largely performed on naked DNA substrates. However, loops that comprise metazoan mitotic chromosomes contain 100-200 kb of DNA (Banigan and Mirny, 2020), which represents hundreds of nucleosomes. Since condensins prefer to bind nucleosome-free DNA (Kong et al., 2020; Shintomi et al., 2017; Toselli-Mollereau et al., 2016; Zierhut et al., 2014), it remains unknown if loop extrusion can proceed on chromatin. Human condensin complexes are capable of loop extrusion through sparsely arranged nucleosomes *in vitro* (Kong et al., 2020). However, mitotic chromatin adopts a more compact fiber structure (Grigoryev et al., 2016; Ou et al., 2017). Furthermore, large protein complexes such as RNA polymerases are able to limit loop extrusion by SMC protein complexes, such as bacterial condensins and eukaryotic cohesins (Brandão et al., 2019; Hsieh et al., 2020; Krietenstein et al., 2020). Nucleosomes also restrict the diffusion of cohesin *in vitro* (Stigler et al., 2016). Therefore, the effect of higher order chromatin fiber structure and other mitotic chromatin proteins on processive loop extrusion by condensin remains unknown.

Interphase nuclei are segmented to chromosome territories, each of which contain highly entangled sister chromatids after replication (Cremer and Cremer, 2010; Farcas et al., 2011; Sundin and Varshavsky, 1981). Replicated pairs of chromatids are linked by cohesin during interphase, and cohesin removal during prophase (except at centromeres and other limited protected loci) promotes resolution of sister chromatids, together with actions of condensin II and topo II (Gandhi et al., 2006; Nagasaka et al., 2016). Different chromosomes are also largely unentangled in interphase HeLa cells (Goundaroulis et al., 2020; Tavares-Cadete et al., 2020), and Ki-67 localization on chromosome peripheries may act as a steric and electrostatic barrier to prevent interchromosomal entanglement during mitosis (Cuylen et al., 2016). However, some interchromosomal linkages were observed in metaphase (Marko, 2008; Potapova et al., 2019). Although it has been shown that active transcription at rDNA results in mitotic interchromosomal links (Potapova et al., 2019), it remains unclear if mitotic chromosome compaction is coupled to resolution of these interchromosomal linkages.

One of the most abundant chromatin proteins beside core histones is the linker histone, which binds to the dyad of the nucleosome and tethers the two linker DNAs emanating from the nucleosome (Arimura et al., 2020; Bednar et al., 2017; Zhou et al., 2015, 2021). In reconstitution experiments, linker histones cluster oligo- nucleosomes (Li et al., 2016; White et al., 2016), and promote liquid-liquid phase separation (Gibson et al., 2019; Shakya et al., 2020). *In vivo*, linker histones are also enriched in highly compact chromatin (Izzo et al., 2013; Parseghian et al., 2001; Th'ng et al., 2005). While core histones are evolutionarily highly conserved in eukaryotes, linker histones are much more diversified (Izzo et al., 2008). In the human and mouse genome, 11 H1 paralogs are found, among which some combination of six variants (H1.0-H1.5) are widely expressed in somatic cells (Hergeth and Schneider, 2015). In vertebrate oocytes and early embryos, H1.8 (also known as H1OO, H1foo, H1M and B4) is the major linker histone variant (Dworkin-Rastl et al., 1994; Wühr et al., 2014). Immuno-depletion of H1.8 from *Xenopus* egg extracts made mitotic chromosomes thinner and longer, causing defective chromosome segregation in anaphase (Maresca et al., 2005). However, the mechanism by which the linker histone affects large scale chromosome length changes remains unknown.

Here we demonstrate that the linker histone H1.8 suppresses enrichment of condensins and topo

II on mitotic chromosomes. In a reconstitution system with purified components, H1.8 inhibits binding of topo II and condensins to nucleosome arrays. Through a combination of chromosome morphological analysis and Hi-C, we show that H1.8 reduces chromosome length by limiting condensin I loading on chromosomes, while H1.8 limits chromosome individualization by antagonizing both condensins and topo II. This study establishes a mechanism by which the linker histone tunes the compaction and topology of mitotic chromosomes.

Results

Linker histone H1.8 limits enrichment of condensins and topo II on mitotic chromatin

Depletion of linker histone H1.8 in *Xenopus* egg extracts makes chromosomes thinner and elongated (see **Figure 3C**, Maresca et al., 2005). We asked if this phenotype may reflect the potential role of H1.8 in regulating condensins and TOP2A (the dominant topo II isoform in *Xenopus* egg extracts, Wühr et al., 2014), which are essential for mitotic chromosome compaction in *Xenopus* egg extracts (Adachi et al., 1991; Cuvier and Hirano, 2003; Hirano and Mitchison, 1994). *In silico* simulation analysis suggests that increasing the number of loop extruders (such as condensin I) on DNA, beyond a minimum threshold, makes chromosomes longer and thinner (Goloborodko et al., 2016a). Reducing condensin I levels on chromatin also made chromosomes shorter experimentally (Elbatsh et al., 2019; Fitz-James et al., 2020; Shintomi and Hirano, 2011). Although it has been reported that H1.8 depletion does not affect chromosomal enrichment of major chromatin proteins (Maresca et al., 2005), we therefore attempted to quantify chromatin-bound levels of condensins and TOP2A.

To investigate whether H1.8 regulates chromatin levels of condensins and topo II, we prepared mitotic chromosomes in *Xenopus laevis* egg extracts depleted of H1.8. Demembranated *Xenopus laevis* sperm nuclei were added to either mock (Δ IgG) or H1.8 depleted (Δ H1) extracts from eggs arrested at meiotic metaphase II by cyostatic factor (CSF extracts) (**Figure 1A, 1B**).

Figure 1. Linker histone H1.8 suppresses enrichment of condensins and TOP2A on mitotic chromatin

A) Experimental scheme to generate replicated chromosomes in *Xenopus* egg extracts. **B)** Western blots of total extracts showing depletion of H1.8 from *Xenopus* egg extracts and rescue with recombinant H1.8 (rH1.8). **C)** Representative images of DNA (Hoechst 33342), CAP-G (condensin I) and TOP2A immunofluorescence on chromosomes in metaphase extracts treated with nocodazole in the indicated conditions. Chromosomes in each nucleus remain clustered in the presence of nocodazole. Bar, 10 μ m. **D)** Quantification of CAP-G (condensin I) and TOP2A immunofluorescence signals normalized to the DNA (Hoechst) signal for the indicated conditions. Each grey or magenta dot represents the average signal intensity of a single chromosome cluster (from one nucleus). Each black dot represents the median signal intensity from a single experiment. Bars represent mean and range of the medians of two independent experiments. **E)** Western blots of mitotic chromatin purified from mock (Δ IgG) and H1.8-depleted (Δ H1) extracts (top) and quantification of band intensities normalized to H3 and H2B (below). Mean and SEM from 3 experiments. **F)** Representative images of CAP-G2 (condensin II) immunofluorescence on chromosomes in metaphase extracts with nocodazole in the indicated conditions. Bar, 10 μ m. **G)** Quantification of the CAP-G2 (condensin II) normalized to the DNA (Hoechst) signal for the indicated conditions. Each grey or purple dot represents the average signal intensity of a single chromosome cluster (from one nucleus). Each black dot represents the median signal intensity from a single experiment. Bars represent mean and range of the median of two independent experiments. **H)** Western blots of mitotic chromatin purified from mock (Δ IgG) and H1.8-depleted (Δ H1) extracts. The statistical significance of the changes in each immunofluorescence experiment was assessed by a Mann Whitney *U*-test and the *p*-values

shown in D, E and G are calculated by an unpaired Student's *t*-test of the aggregate medians (immunofluorescence) or blot intensities of each single experiment. The number of nuclei imaged in D and G in each condition for each experiment are indicated in the figure.

Figure S1- H1.8 depletion does not lead to global accumulation of DNA-binding proteins

Table S1- High abundance proteins with the highest increases in Δ H1 depleted chromosomes

Table S2- High abundance proteins with the highest decreases in Δ H1 depleted chromosomes

Calcium was added to cycle the extract into interphase and induce functional nuclear formation, in which chromosomes were replicated. The corresponding depleted CSF extract was then added to generate metaphase chromosomes (Shamu and Murray, 1992). To eliminate the microtubule-dependent change in chromosome morphology, which may affect quantitative analyses of chromatin proteins, spindle assembly was inhibited using nocodazole. Chromosomes were fixed and the levels of TOP2A and condensin I subunit CAP-G were measured by immunofluorescence (**Figure 1C**). Depletion of H1.8 increased the levels of both CAP-G and TOP2A on mitotic chromosomes, while adding back recombinant H1.8 rescued the phenotype (**Figure 1D**). Identical results were obtained when immunofluorescence signal normalization was done by the minor groove DNA-binding dye Hoechst 33342, or by fluorescent dUTP that was incorporated during replication (**Figure S1A-C**). The dUTP quantitation also confirmed that H1.8 depletion did not affect DNA replication (**Figure S1B**), as reported previously (Dasso et al., 1994). The apparent increased signal intensities of CAP-G and TOP2A on chromatin in Δ H1 extracts were not the general consequence of elongated chromosome morphology, as H1.8 depletion did not affect a panel of other chromatin proteins, regardless of their binding preference to nucleosomes or nucleosome-free DNA (Zierhut et al., 2014) (**Figure S1D**). Enhanced chromosome binding of condensin I and TOP2A in Δ H1.8 extracts was biochemically confirmed by quantifying their levels on purified metaphase chromosomes by western blotting and by mass spectrometry (**Figure 1E, S1E, Table S1, S2**). Mass spectrometry data also confirmed that H1.8 depletion did not result in a global enrichment of all chromatin bound proteins. Condensin II levels on chromosomes also showed similar increases in Δ H1 extracts by immunofluorescence and western blots on purified chromosomes, although peptides of condensin II specific subunits were not detected by mass spectrometry (**Figure 1F-H, S1E**).

Linker histone H1.8 reduces binding of condensins and TOP2A to nucleosome arrays

Since linker histone depletion does not change the nucleosome spacing in *Xenopus* egg extracts (Ohsumi, Katagiri, & Kishimoto, 1993), we hypothesized that H1.8 depletion results in an increase in linker DNA that becomes accessible to condensins and TOP2A. To test this possibility, we reconstituted nucleosome arrays with purified histones, and asked if H1.8 interferes with binding of recombinant human condensins and *X. laevis* TOP2A (Kong et al., 2020; Ryu et al., 2010) (**Figure 2A**). The nucleosome array was composed of 19 tandem repeats of 147 bp Widom 601 nucleosome positioning sequence and 53 bp linker DNA (Lowary and Widom, 1998), where full occupancy of the array by nucleosome core particle (NCP) and H1.8 to each repeat unit was confirmed by native polyacrylamide gel electrophoresis (PAGE) (**Figure S2A**). As previously shown with human condensin I purified from HeLa cells (Kimura et al., 2001), our recombinant human condensin I, but not the ATP-binding defective Q-loop mutant (Hassler et al., 2019; Hopfner et al., 2000; Kong et al., 2020; Löwe et al., 2001), was able to rescue mitotic chromosome morphology defects caused by condensin I depletion in *Xenopus* extracts (**Figure S2B**), demonstrating that the recombinant human condensin I can functionally replace *Xenopus* condensin I. The recombinant *Xenopus laevis* TOP2A used was also able to rescue chromatid formation in Δ TOP2A extracts and was able to perform ATP dependent kinetoplast decatenation *in vitro* (**Figure S2C, D**).

Condensin I binds weakly to mononucleosomes with linker DNA (Kong et al., 2020). We observed similar weak binding of condensin I to nucleosome arrays and this weak binding was stimulated by ATP (**Figure S3A**). Since this enhancement was not seen when magnesium concentration was higher than the ATP concentration (**Figure S3B**), we hypothesized that ATP might have enhanced condensin binding to the nucleosome arrays by chelating magnesium, which is known to induce chromatin compaction

Figure 2. Linker histone inhibits binding of condensins and TOP2A to nucleosome arrays

A) Experimental scheme for testing the effect of recombinant H1.8 (rH1.8) on binding of purified condensins and TOP2A to arrays of nucleosomes assembled on the Widom 601 nucleosome positioning sequence. **B**) Coomassie staining of SDS-PAGE gels, showing input (top) and nucleosome-array bound fraction (middle) of condensin I, rH1.8 and core histones. The right most lanes represent the streptavidin beads only negative control. Buffer contains 2.5 mM MgCl₂, 5 mM ATP and indicated concentrations of NaCl. The band intensities of condensin I subunits were normalized to the histone bands and the binding at 50 mM NaCl for nucleosome arrays without H1.8. Mean and S.E.M of three independent experiments are shown (bottom). **C**) Same as B, except that nucleosome array binding of condensin II is shown. Mean and S.E.M (wildtype)/range (Q-loop mutant) of three (wildtype) or two (Q-loop mutant) independent experiments are shown. **D**) Same as B, except that nucleosome array binding of TOP2A in buffer containing 1 mM MgCl₂ is shown. Mean and range of two independent experiments are shown. The p-value in D was calculated using an unpaired Student's t-test.

Figure S2- Recombinant human condensin I and *X. laevis* TOP2A are functional

Figure S3- Condensin I binding is inhibited by magnesium

Figure S4- H1.8 inhibits condensin I binding to mononucleosomes

(Eltsov et al., 2008; Finch and Klug, 1976). Indeed, both EDTA and ATP, which chelate magnesium, increased binding of the Q loop mutant of condensin I to the nucleosome array (**Figure S3C**), suggesting that condensin binding to the nucleosome array is sensitive to high magnesium concentration and not due to ATP binding of condensin I. In contrast, ATP did not stimulate condensin I binding to mononucleosomes (**Figure S3D**), indicating that excess magnesium may limit condensin binding through compaction of the nucleosome array.

In the buffer condition where excess ATP was present over magnesium, both wild-type and the Q-loop mutant of condensin I showed reduced binding to nucleosome arrays loaded with H1.8 at physiological salt concentrations (50 – 150 mM NaCl) (**Figure 2B**). H1.8 also suppressed binding of condensin II and its Q-loop mutant to the nucleosome array (**Figure 2C**), though, as expected (Kong et al., 2020), condensin II showed higher affinity to the nucleosome array than condensin I. This reduced binding of condensins in the presence of H1.8 can be explained by either direct competition between H1.8 and condensins for the linker DNA or by H1.8-mediated formation of a higher order structure of the nucleosome array (Song et al., 2014). The direct competition model can be tested by using mononucleosomes with linker DNAs instead of the nucleosome array, since H1.8 binding to mononucleosomes does not promote higher order structure or aggregation (White et al., 2016). Supporting the direct competition model, H1.8 reduced binding of condensin I to mononucleosomes (**Figure S4**).

We also examined if H1.8 interferes with binding of the recombinant *Xenopus* TOP2A to nucleosome arrays. As compared to condensins, TOP2A showed more stable binding to the nucleosome array, and H1.8 had no effect on TOP2A binding at low salt concentrations. However, H1.8 did reduce nucleosome array binding of recombinant *Xenopus* TOP2A at 120 mM NaCl (**Figure 2D**). Altogether, these data demonstrate that preloaded H1.8 on nucleosomes can directly interfere with binding of condensins and TOP2A to chromatin.

Chromosome elongation by H1.8 depletion is due to increased chromatin bound condensin I

If the increased amount of condensin I on chromatin is responsible for the chromosome elongation phenotype observed in Δ H1 extracts, reducing condensin I levels should reverse this phenotype. To measure lengths of mitotic chromosomes formed in *Xenopus* egg extracts, we assembled replicated metaphase chromosomes in extracts depleted with mock IgG, H1.8, CAP-G (condensin I) or CAP-D3 (condensin II) antibodies. These extracts were then diluted to disperse individualized chromosomes (**Figure 3A, 3B**) (Funabiki and Murray, 2000). As reported previously (Maresca et al., 2005), average chromosome length increased by ~50% by H1.8 depletion (**Figure 3C, 3D**). Supporting our hypothesis, when condensin I was co-depleted (Δ H1 Δ CAP-G), chromosomes became even shorter than chromosomes in mock depleted extracts (Δ IgG) (**Figure 3C, 3D**). In contrast, condensin II co-depletion (Δ H1 Δ CAP-D3) did not change the chromosome length (**Figure 3D**). Chromosome length in condensin II-depleted extracts (Δ CAP-D3) was also indistinguishable from mock depleted chromosomes (**Figure S5A**), consistent with the negligible effect of condensin II depletion in the presence of normal levels of condensin I in *Xenopus* egg extracts (Shintomi and Hirano, 2011).

If H1.8 regulates chromosome length by a mechanism independent of its role in condensin I inhibition, we expect that the chromosome lengths in Δ CAP-G extracts and Δ H1 Δ CAP-G extracts would be different. Unfortunately, since severe defects in chromosome individualization in Δ CAP-G extracts prevented us from measuring chromosome lengths, this comparison was not possible (see **Figure 5B**). To circumvent this issue, we asked if similar condensin I levels on both Δ IgG and Δ H1 chromatin would result in similar chromosome lengths. Since condensin I binding to chromosomes is suppressed by H1.8, to load condensin I to chromosomes in Δ H1.8 extracts at the levels seen in control (Δ IgG) extracts, we assembled chromosomes in extracts where condensin I was partially depleted (**Figure S5B**). When mitotic chromosomes were assembled in Δ H1.8 extracts that contain 40 % (of Δ IgG) condensin I (CAP-G), condensin I level on these chromosomes was equivalent to that on chromosomes in control (Δ IgG) extracts (**Figure 3E**). Consistent with the hypothesis that H1.8 does not

Figure 3. Chromosome elongation by H1.8 depletion is due to enhanced condensin I loading on chromatin

A) Schematic of extract dilution to disperse individualized chromosomes. **B)** Western blots of total egg extracts showing depletions of indicated proteins. **C)** Representative images of mitotic chromosomes after dilution of indicated extracts. Bar, 10 μ m. **D)** Quantification of the chromosome length. Data distribution of the length of individual chromosomes from three independent experiments (green, purple, grey) is shown. Each black dot represents the median length of chromosomes from a single experiment. Bar represents mean and S.E.M of three independent experiments. **E)** Quantification of CAP-G levels normalized to DNA signal (Cy3-dUTP) in the indicated conditions by immunofluorescence. Each dot represents the mean of CAP-G intensity normalized to DNA intensity of a single chromosome cluster (from one nucleus). The data plotted is median $\pm$ 95% C.I. **F)** Chromosome lengths in the indicated condition. Each dot represents length of a single chromosome. Bars represent median $\pm$ 95% C.I. The p-values in D compare the median chromosome lengths in each condition and were calculated using an unpaired Student's t-test and those in E, F compare the median values in a single experiment and were calculated using a two-tailed Mann-Whitney *U* test. The number of nuclei (E) or chromosomes (D, F) imaged in each condition for each experiment are indicated above the figure.

Figure S5. Condensin II depletion does not affect chromosome length

contribute to chromosome length regulation independently of condensin I, average chromosome length was essentially identical between these two conditions ($\Delta H1$ 40 % CAP-G and ΔIgG) (**Figure 3F**). In summary, these results demonstrate that H1.8 controls mitotic chromosome lengths primarily through limiting the chromosome binding of condensin I.

H1.8 increases condensin I driven mitotic loop layer organization

Experimental and simulation studies have shown that mitotic chromosome length and width are sensitive to the number of condensin I molecules on a chromosome, through affecting the loop size (Fitz-James et al., 2020; Gibcus et al., 2018; Goloborodko et al., 2016b, 2016a). Low levels of condensin I are predicted to result in fewer and larger loops, while higher levels of condensin I will lead to a larger number of smaller loops. Therefore, if linker histone H1.8 decreases chromosome length through limiting condensin I association with chromatin, we expect that the average size of mitotic condensin loops decreases upon H1.8 depletion. As an alternative way to quantitatively assess the effect of H1.8 and condensins on mitotic chromosome organization, we used the chromosome conformation capture assay Hi-C (Lieberman-Aiden et al., 2009).

Hi-C contact probability maps were generated from replicated metaphase *Xenopus laevis* sperm chromosomes 60 min after cycling back into mitosis in the presence of nocodazole. As seen in mitotic chromosomes in somatic cells (Gibcus et al., 2018; Naumova et al., 2013) and in early stages of mouse development (Du et al., 2017), all the Hi-C contact maps showed no checkerboard pattern commonly associated with interphase chromosome compartments, and lacked any sign of topologically associating domains (TADs) (Szabo et al., 2019) (**Figure 4A**- Data available at GEO under accession no. GSE164434). Hi-C interaction maps are characterized by the decay in contact probability, P , as a function of the genomic distance, s . To derive quantitative information about the polymer structure of the mitotic chromosomes, we plotted the genome-wide average $P(s)$ (**Figure 4B**). $P(s)$ plots were consistent among two biological replicates (**Figure S6A**) and among different chromosomes (**Figure S6B**). The interaction decay profile of the control (ΔIgG) chromosomes was also qualitatively similar to mitotic chromosomes in both DT40 and human

cells (Elbatsh et al., 2019; Gibcus et al., 2018; Naumova et al., 2013). Condensin I depletion ($\Delta CAP-G$) caused a major change in the Hi-C map (**Figure 4A**), reflecting its severe morphological defects in mitotic chromosomes in *Xenopus* egg extracts (Hirano et al., 1997). Unlike in DT40 cells, where depletion of condensin I and condensin II reduces interactions at shorter (< 6 Mb) and longer (> 6 Mb) distances respectively (Gibcus et al., 2018), condensin I depletion affected interactions at longer distances (~ 10 Mb), whereas condensin II depletion ($\Delta CAP-D3$) did not cause recognizable changes in interactions at both long and short distances (**Figure S6C**). A second diagonal band, which is indicative of a strong helical organization in the chromosome axis (Gibcus et al., 2018), was not seen in *Xenopus* egg extracts (**Figure 4A, S6C**), likely reflecting the minor contribution of condensin II in this system or perhaps due to a prolonged arrest in mitosis (Gibcus et al., 2018). In H1.8-depleted extracts ($\Delta H1$), a dramatic decrease in interactions at long genomic distances (1-10 Mb) was observed (**Figure 4A, 4B**), as expected from the thinner chromosomes (**Figure 3**).

$P(s)$ plots are useful to determine the underlying polymer structure of the chromosome such as average loop size (Gibcus et al., 2018). Specifically, the first derivative (slope) of the $P(s)$ plots can reveal both the average loop size and the amount of DNA per layer of the rod-shaped mitotic chromosome (layer size) (Abramo et al., 2019; Gassler et al., 2017; Gibcus et al., 2018) (**Figure 4C**). Average loop size can be estimated from the peak value in the derivative plot (Gassler et al., 2017; Patel et al., 2019). As the derivative plot in our data showed a peak in the 10 kb-1 Mb range, we estimated that this peak location reflects the average loop size. Unlike mitotic chromosomes in DT40 or HeLa cell lines (Abramo et al., 2019; Gibcus et al., 2018), metaphase chromosomes from *Xenopus* extracts had a flattened peak (**Figure S6D**), possibly due to a larger variation in loop sizes. Although it is difficult to estimate the exact loop size from these plots, chromosomes from $\Delta H1$ extracts showed a reproducible shift of the peak towards smaller genomic distances (**Figure S6D**). This is consistent with a decrease in loop size due to increased condensin I accumulation on chromatin upon H1.8 depletion. Derivative plots generated from Hi-C maps of dispersed chromosomes in diluted extracts (**Figure 3A**) showed a similar shift in peak towards smaller genomic distances upon

Choppakatla et al., April 2021-preprint copy- BioRxiv

Figure 4. Effects of H1 and/or condensin I, II depletion on mitotic genome folding

A) Hi-C maps of metaphase *X. laevis* chromosome 3S, binned to 250 kb, in the indicated condition. **B)** Genome wide average contact probability decay curves for the indicated conditions showing the changes in longer distance interactions. **C)** Derivative plots of the average contact probability decay curves for dispersed chromosomes from mock (Δ IgG) and H1.8 depleted extracts (Δ H1) showing the change in estimated loop size. The solid and dotted lines are from two independent biological replicates. **D)** Derivative plots of the genome wide average contact probability decay curves in the indicated conditions. The dotted lines indicate the layer sizes for each plotted condition. **E)** Estimates of layer size from derivatives of genome wide probability decay curves upon depletion of H1.8 and CAP-G (condensin I) or CAP-D3 (condensin II). The mean and range of two biological replicates is shown.

Figure S6- Effects of H1, condensin I and condensin II depletion on mitotic genome folding

Figure S7- H3-H4 depletion leads to even smaller loop sizes

H1.8 depletion (**Figure 4C**). The derivative plots from these dispersed chromosomes showed a better-defined peak, allowing a more precise loop size estimate of 140 kb in Δ IgG extracts and 110 kb in Δ H1 extracts. These loop sizes are comparable to the estimated size of condensin I-driven loops in DT40 and HeLa cells (Gibcus et al., 2018; Naumova et al., 2013).

$P(s)$ plots for mitotic chromosomes display three regimes that are typical for relatively stiff rod-shaped conformation (Gibcus et al., 2018; Naumova et al., 2013): for small genomic distance the contact probability is dominated by interactions between pairs of loci located within loops (up to 1-200 kb). For loci separated by up to a few megabases, the contact probability decays slowly with genomic distance. These interactions mostly reflect contacts between loci located in different loops which are relatively closely packed as a radial layer of loops around the central axis (intra-layer regime). The third regime is characterized by a steep decay in contact probability at several megabases. This represents pairs of loci separated by a relatively large distance along the axis of the rod-shaped chromosome so that they very rarely interact (inter-layer regime). The size of these layers can be obtained from the derivative of $P(s)$, where the steep drop in contact probability curve is marked by sharp drop in the derivative value. For control chromosomes the average amount of DNA in each layer of loops (layer size) was approximately 5 Mb, given the steep decay in contact probability observed for loci separated by more than 5 Mb for both replicates (**Figure 4B, 4D**). For H1.8 depleted chromosomes we observed a smaller layer size of around 3.5 Mb (**Figure 4B, 4D**). This 1.5-fold reduction of DNA content of each layer would explain the observed 1.5-fold increase in chromosome length (**Figure 3D**). Further, we can derive the number of loops per layer by dividing the layer size by the loop size in the corresponding condition. In both control (Δ IgG) and H1.8 depleted extracts (Δ H1), the number of loops per layer was ~ 40 . This indicates that the change in the layer size upon H1.8 depletion is a result of decreased loop size, while the number of loops per layer is not affected. The layer size in H1.8/condensin I co-depleted extracts (Δ H1 Δ CAP-G) was larger than those in control extracts (Δ IgG) (**Figure 4D, 4E**). This is consistent with the shorter chromosomes in Δ H1 Δ CAP-G extracts than in control Δ IgG extracts (**Figure 3D, 3F**). Condensin II co-depletion (Δ H1 Δ CAP-D3) did not affect the layer size (3.5 Mb) and also the chromosome length (**Figure 4D, 4E**,

3D). Condensin II depletion alone (Δ CAP-D3) also did not affect the layer size, consistent with the observed lack of change in the chromosome length (**Figure 4D, 4E, Figure S5A**). Taken together, these data support the hypothesis that H1.8 limits the condensin I level on chromatin and shortens chromosome lengths, allowing each condensin I to form a longer loop, and consequentially tuning the amount of DNA present in each layer. The data also suggest that unlike in chicken DT 40 cells, condensin I and not condensin II plays the dominant role in the organization of loop layers (Gibcus et al., 2018).

Since condensin I binding to DNA is also limited by nucleosomes (Kong et al., 2020; Shintomi et al., 2017; Zierhut et al., 2014), we expected that loss of nucleosomes would similarly reduce the loop and layer sizes. To test this, we depleted H3-H4 tetramers using an antibody to acetylated lysine 12 of histone H4 (H4K12ac)(Zierhut et al., 2014), and generated metaphase chromosomes for Hi-C (**Figure S7A**). Since *X. laevis* sperm contains preloaded paternal H3-H4 (Shechter et al., 2009), the number of nucleosomes in our metaphase chromosomes was expected to be reduced by at most 50 %. As nucleosomes occupy a large majority of the genomic DNA (Chereji et al., 2019; Lee et al., 2007), even a partial histone depletion (Δ H3-H4) increased chromatin bound condensin I beyond that of H1.8 depletion (Δ H1) (**Figure S7B**). Consequentially, the layer size in Δ H3-H4 extracts became much smaller (around 500 kb) than in Δ H1 extracts (**Figure S7C**). Assuming that the number of loops per layer is similar in these chromosomes (~ 40), the loop size estimate is 12 kb, which is much shorter than in H1.8 depletion. Altogether these results suggest that global occupancy of nucleosomes and linker histones can affect DNA loop size and chromosome through controlling the number of condensin molecules on the chromatin fiber.

Choppakatla et al., April 2021-preprint copy- BioRxiv

Figure 5. H1.8 suppresses condensin to limit chromosome individualization

A) Western blots of total egg extracts showing depletion levels in extract of condensin I and condensin II using the CAP-G and CAP-D3 antibodies respectively. * represents non-specific band **B**) Representative images of chromosomes after extract dilution, which disperses individualized chromosomes. DNA and centromere-associated CENP-A immunofluorescence are shown. Bar, 20 μ m. **C**) Percent frequency of individualized chromosomes (chromosomes in DNA masses with <4 CENP-A foci) in the indicated conditions. A large majority of DNA masses with no CENP-A foci are derived from Δ CAP-D3 extracts, where CENP-A loading is compromised (Bernad et al., 2011). DNA masses and CENP-A foci were identified using Otsu's thresholding algorithm and CENP-A foci in a binarized DNA mask were counted. The numbers of DNA masses counted in each condition were as follows: Δ IgG (502, 643), Δ H1 (1279, 839), Δ CAP-G (447, 170), Δ CAP-D3 (937), Δ H1 Δ CAP-G (1565, 301), Δ H1 Δ CAP-D3 (1536, 300), Δ H1 Δ CAP-G Δ CAP-D3 (300, 156). **D**) Quantification of CAP-G (condensin I) immunofluorescence normalized to the DNA signal for the indicated conditions. Each grey or orange dot represents the average signal intensity of a single chromosome cluster (from one nucleus). Each black dot represents the median signal intensity from a single experiment. Bars represent mean and range of the medians of two independent experiments. **E**) Quantification of CAP-G2 (condensin II) immunofluorescence intensity, normalized to the DNA signal for the indicated conditions. Each grey or magenta dot represents the average signal intensity of a single chromosome cluster (from one nucleus). Each black dot represents the median signal intensity from a single experiment. Bars represent mean and range of the medians of two independent experiments. The p-values in C, D and E were calculated by an unpaired Student's *t*-test. The number of nuclei imaged in each condition in D and E in each experiment are indicated above the figures.

Figure S8- Regulation of chromosome individualization by topo II, condensins and H1.8 in *Xenopus* egg extracts

H1.8 suppresses condensin-driven mitotic chromosome individualization.

Condensins and topo II act in concert to generate mitotic chromosomes from decondensed interphase nuclei (Cuvier and Hirano, 2003). Both experimental observations and *in silico* experiments also suggest that condensin can drive complete decatenation of sister-chromatids (Goloborodko et al., 2016a; Marko, 2009; Nagasaka et al., 2016). Condensin is also required to keep decatenating sister chromatids in metaphase (Piskadlo et al., 2017). Although it has been suggested that condensin-mediated chromosome compaction promotes chromosome individualization (Brahmachari and Marko, 2019; Sun et al., 2018), it remains to be established if different linear chromosomes (non-sisters) are catenated with each other even after completion of mitotic compaction since Ki-67 on chromosome surfaces may act as a barrier to prevent interchromosomal DNA interaction during mitosis (Cuylen et al., 2016). To investigate if substantial interchromosomal entanglements exist in metaphase *Xenopus* egg extracts, we examined if the topo II inhibitor ICRF-193 blocks chromosome individualization after mitotic chromosome compaction (**Figure S8A**). While chromosomes clustered on the metaphase spindle in DMSO-treated control extracts were effectively dispersed into individualized chromosomes after diluting extracts, most chromosomes remained clustered when ICRF-193 was added at the beginning of mitotic induction and incubated for 50 min (**Figure S8B, C**, ICRF-50 min). Even when ICRF-193 was added to metaphase egg extracts after completion of metaphase spindle formation, but 2 min before extract dilution (**Figure S8A**, bottom), efficiency of chromosome individualization decreased (**Figure S8B, C**, ICRF-2 min). These results suggest that substantial interchromosomal topological catenations remain unresolved in metaphase.

We then asked if H1.8-mediated suppression of condensins limits chromosome individualization and resolution of interchromosomal linkages. If so, H1.8 depletion may reduce the minimum required level of condensin activities to support chromosome individualization. In metaphase mock-depleted (Δ IgG) and H1.8-depleted (Δ H1) extracts with replicated chromosomes, the dilution procedure (**Figure 3A**) resulted mostly in physically separated single chromosomes but also a small number of clumped chromosomes (**Figure 5B**, **Figure S9B**). To better quantify chromosome individualization, we

measured the number of CENP-A foci in each distinct chromatin mass (**Figure S8D**). Note that each centromere pair in a chromosome is counted as one focus, and we defined individualized chromosomes as chromosomes present in DNA masses containing less than 4 CENP-A foci (**Figure 5C**; this threshold was to include largely individualized chromosomes that stochastically come into contact- counting chromosomes in DNA masses having <2 foci yielded a similar result- **Fig S8E**). Condensin I depletion (Δ CAP-G) resulted in defective chromosome individualization, suggesting that condensin I activity drives resolution of interchromosomal entanglements, and that condensin II is not sufficient to resolve these interchromosomal links in this background. Strikingly, co-depletion of H1.8 and condensin I (Δ H1 Δ CAP-G) almost completely rescued chromosome individualization without detectable CAP-G on chromatin (**Figure 5A-D**). This apparent bypass of condensin I requirement in chromosome individualization required condensin II as chromosome individualization failed in the triple-depleted extracts (Δ H1 Δ CAP-G Δ CAP-D3) (**Figure 5B, C**). The increased condensin II in Δ H1 Δ CAP-G extracts may replace the function of condensin I by condensin II in the absence of H1.8 (**Figure 5E**). These results demonstrate that H1.8-mediated suppression of condensin enrichment on chromatin limits chromosome individualization during mitosis.

H1.8 prevents chromosomes from hyper-individualization by suppressing condensins and topo II

We next examined the functional significance of H1.8-mediated suppression of TOP2A enrichment on mitotic chromosomes (**Figure 1C-E**), asking if H1.8 depletion could reduce the required TOP2A level in extracts for chromosome individualization. Complete loss of TOP2A inhibits decompaction of sperm nuclei, a process associated with replacement of protamines with histones (Adachi et al., 1991), so we addressed this question using extracts partially depleted of TOP2A (**Figure 6A**). We first generated nuclei with replicated chromosomes in extracts containing the normal level of TOP2A, and then the extracts were diluted with Δ TOP2A extracts to reduce the total TOP2A level to 25 % (**Figure S9A**). Under this condition, the level of chromosome-associated TOP2A also reduced to 25 % (**Figure 6B**).

Choppakatla et al., April 2021-preprint copy- BioRxiv

Figure 6. H1.8 suppresses hyper-individualization through condensins and topo II

A) Schematic of partial TOP2A depletion to test sensitivity of chromosome individualization to TOP2A levels. **B**) Quantification of chromosome-associated TOP2A upon partial TOP2A depletion. Each dot represents the mean of TOP2A intensity normalized to DNA intensity of a single chromosome cluster (from one nucleus). The data plotted is median $\pm$ 95 % C.I. **C**) Percent frequency of DNA clusters categorized as unindividualized nuclei (**Figure S9B**) upon partial TOP2A depletion. Mean and S.E.M from three independent experiments. Each dot represents the percentage of unindividualized chromosome clusters in the indicated condition in an independent biological replicate. **D**) Representative images of nuclei using DNA (Cy5-dUTP) showing the clustering phenotype in the indicated conditions. Scale bar, 10 μ m. **E**) Quantification of the three-dimensional surface area of the chromosome clusters in **D** normalized to DNA (Cy5-dUTP) signal. Each grey, magenta, green or orange dot represents the normalized surface area of single nucleus or chromosome cluster and each black square represents the median surface area of a single experiment. Data plotted is mean and S.E.M of four independent experiments. **F**) Quantification of TOP2A immunofluorescence intensity normalized to the DNA signal for the indicated conditions. Each grey or magenta dot represents the average signal intensity of a single chromosome cluster (from one nucleus). Each grey, magenta, orange or grey dot represents the median signal intensity from a single experiment. Mean and S.E.M of the median of four independent experiments are also shown. **G**) Representative images of nuclei using DNA (Cy3-dUTP) showing the clustering phenotype in the indicated conditions. Scale bar, 10 μ m. **H**) Quantification of the three-dimensional surface area of the chromosome clusters in **G** normalized to DNA (Cy3-dUTP) signal. Each black open circle represents the normalized surface area of single nucleus or chromosome cluster. Data plotted is median and 95 % C.I. The p-values in C and F were calculated by an unpaired Student's t-test and the p-values in H were calculated by a two-tailed Mann-Whitney U test. The number of nuclei imaged in each condition for each experiment in E, F and H are indicated above the figure.

Figure S9. H1.8 suppresses over-individualization through condensins and topo II

Upon reduction of TOP2A to 25 %, the frequency of unindividualized chromosome clusters increased about 3-fold in Δ IgG background (**Figure 6C**). However, in the absence of H1.8, extracts with 25% levels of TOP2A were still able to support maximum level of chromosome individualization as chromosome-associated levels of TOP2A became equivalent to untreated control extracts (**Figure 6C**). These data suggest that chromosome individualization is sensitive to TOP2A levels on chromatin, and that H1.8 suppression of TOP2A plays a role in suppressing chromosome individualization.

Since the purpose of mitosis is to equally segregate chromosomes to dividing cells, it may seem odd that H1.8 suppresses TOP2A and condensins to limit chromosome individualization. In many cells, the spindle gathers the individualized mitotic chromosomes at the metaphase plate before chromosomes are ready for segregation. In *Xenopus* egg extracts, however, even when spindle assembly is inhibited using nocodazole, all chromosomes from one nucleus tightly cluster into a single mass (**Figures 1C, 6D**). In contrast, individual chromosomes were more readily distinguished by DNA staining and spread to larger area in Δ H1 extracts, suggesting that the H1.8 mediated suppression of chromosome individualization is responsible for the chromosome clustering when spindle assembly is compromised. To quantify this phenotype, we measured the three-dimensional surface area of these chromosomes. The severe chromosome clustering defect in H1.8 depleted (Δ H1) extracts resulted in higher surface area than chromosomes in control (Δ IgG) extracts (**Figure 6D, 6E**). While condensin I depletion alone (Δ CAP-G) showed no significant differences to control (Δ IgG) extracts, condensin I co-depletion partially rescued the chromosome clustering defect of H1.8 depletion (Δ H1 Δ CAP-G). Notably, chromosomes were more spread in Δ H1 Δ CAP-G extracts than in Δ IgG extracts, despite the shorter average chromosome length in Δ H1 Δ CAP-G extracts (**Figure 3D, 3F**), suggesting that chromosome spreading upon H1.8 depletion is not primarily driven by chromosome elongation but by chromosome hyper-individualization. Since TOP2A is enriched on chromosomes upon H1.8 depletion in a condensin-independent manner (**Figure 6F, S9C**), we then asked if this increased TOP2A plays a role in this increased chromosome spreading. Partial TOP2A depletion had no effect on clustering in control (Δ IgG) background since these nuclei were already tightly packed together (**Figure**

6G, 6H- 25 % TOP2A). TOP2A depletion in H1.8 depleted extracts also did not revert the chromosome spreading despite that TOP2A levels on these chromosomes became comparable to control (Δ IgG-100 %) levels (**Figure 6G, 6H**), perhaps because increased condensin loading can compensate for reduced TOP2A activity. However, partial TOP2A depletion reduced chromosome spreading in H1.8/condensin I co-depleted (Δ H1 Δ CAP-G) extracts, suggesting that suppression of both condensin and topo II by H1.8 keeps egg extract chromosomes together during mitosis even in the absence of spindle microtubules.

Discussion

It has been thought that the linker histone H1 promotes local chromatin compaction through stabilizing linker DNA and facilitating nucleosome-nucleosome interaction (Li et al., 2016; Song et al., 2014; White et al., 2016). Here we demonstrated that H1.8 also plays a major role in regulation of long-range DNA interaction through controlling chromatin loading of condensins and topo II. Although condensins and topo II are activated in mitosis to drive chromosome segregation (Hirano and Mitchison, 1991; Kimura et al., 1998), we showed that their functionalities are antagonized by H1.8 to tune chromosome length and prevent hyper-individualization, which increases surface area (Figure 7). Thus, the linker histone H1, which can promote local chromatin compaction through promoting nucleosome-nucleosome interaction, can also suppress chromosome dispersion that is driven by condensins and topo II that organize the long-range DNA folding.

Nucleosomes reduce binding of condensins to DNA *in vitro* (Kong et al., 2020), *in vivo* (Toselli-Mollereau et al., 2016) and in *Xenopus* egg extracts (Shintomi et al., 2017; Zierhut et al., 2014). Now we showed that the linker histone H1.8 limits binding of condensin I and II to chromatin both *in vitro* and in *Xenopus* egg extracts. H1.8 suppressed condensin binding on both mononucleosomes and nucleosome arrays, suggesting that H1.8 is able to compete out condensins for the same linker DNA targets, though the capacity of linker histones to promote higher order structures or phase separation may also limit the access of condensin (Gibson et al., 2019; Song et al., 2014). Since chromosome lengths can be dictated by the amount of condensins (Figure 3F), we propose that regulating the linker histone that limits

chromatin-loading of condensins could serve as a rheostat to control chromosome length (**Figure 7**).

Since the average loop size is an aggregate result of loop extrusion rate, processivity and number of loop extruders (Alipour and Marko, 2012; Goloborodko et al., 2016b), it is unclear from our data whether H1.8 affects the loop extrusion kinetics of a single condensin molecule. However, our observation is consistent with the *in silico* simulation showing that increasing the number of loop extruders makes chromosomes thinner and longer by reducing average loop size (Goloborodko et al., 2016a). Similar DNA loop shortening accompanied with increased condensin I loading was also reported on integrated fission yeast genome DNA segments in mouse and human chromosomes (Fitz-James et al., 2020). Reduced condensin loading due to mutations in condensin I or expression of phosphomimetic H3 mutants in human cells also lead to reduced chromosome length (Elbatsh et al., 2019). Condensin binding is similarly suppressed by nucleosomes, but loop extrusion proceeds unhindered through sparsely distributed nucleosomes (Kong et al., 2020). Since the length of chromosomes with similar condensin I levels is unaffected by the presence of H1.8 (Figure 3E, 3F), H1.8 may increase the loop size by simply reducing the number of condensin molecules on chromatin but may not necessarily inhibit the loop extrusion rate or processivity.

The regulatory mechanisms of TOP2A recruitment to mitotic chromatin are less clear than those of condensins. TOP2A is a DNA binding protein that may also bind histone tails (Lane et al., 2013). Similar to condensin, both TOP2A and TOP2B preferentially localize at active and highly transcribed chromatin indicating a possible preference for nucleosome free regions (Canela et al., 2017; Thakurela et al., 2013; Yu et al., 2017). We observe that TOP2A levels on chromatin increase upon linker histone depletion in egg extracts and that linker histone inhibits TOP2A binding to nucleosome arrays *in vitro*, suggesting that a similar rheostat like mechanism for controlling TOP2A levels on mitotic chromatin by controlling H1 stoichiometry is possible. The preference of topo II for binding linker DNA may also explain the observation of well-spaced TOP2B binding peaks around the well-spaced nucleosomes around the CTCF- binding sites (Canela et al., 2017, 2019). However, in contrast to our finding that more TOP2A associates with elongated chromosomes in Δ H1 extracts, depletion of topo II leads to elongated

chromosomes in vertebrate somatic cell lines and in early embryos of *Caenorhabditis elegans* (Farr et al., 2014; Ladouceur et al., 2017; Nielsen et al., 2020; Samejima et al., 2012). It is unclear whether these observations suggest that increased topo II activity would lead to shorter chromosomes. One possibility is that impaired decatenation due to reduced topo II may interfere with condensin-mediated loop formation, leading to shorter loop size and thinner chromosomes as reported. On the other hand, excess topo II on chromatin, as seen in Δ H1 extracts, may not affect chromosome shape.

Our data also suggest that condensins and topo II combine to resolve extensive and persistent interchromosomal topological entanglement during mitotic compaction in *Xenopus* egg extracts (**Figure S8**), but these activities are counterbalanced by H1.8. In HeLa cells, some variants of H1 are phosphorylated and evicted along the inter-chromatid axis. This partial H1 eviction in prophase seems to be required for complete decatenation along the chromosome arms (Krishnan et al., 2017). This suggests that local enrichment of condensin II upon H1 eviction may play a role in sister chromatid decatenation. In contrast, H1.8 binding to nucleosomes is enhanced upon transition from interphase to M phase in *Xenopus* egg extracts (Arimura et al., 2020). It might be counterintuitive that activities of condensins and topo II are actively antagonized by H1.8 to suppress chromosome individualization. Since spindle assembly in oocytes relies on chromatin-induced microtubule nucleation (Heald et al., 1996), keeping chromosomes at metaphase plate might be important for maintaining robust bipolar spindle. As we showed that H1.8 is important for chromosome clustering when spindle assembly is inhibited (Figure 6), this microtubule-independent chromosome clustering may be particularly important for the large oocyte and early embryonic cells since it would be difficult to assemble a bipolar spindle onto all the chromosomes once they disperse into the large space of the cytoplasm. Clustering chromosomes through incomplete individualization may also avoid generation of chromosomes that do not associate with the spindle. Such a mechanism may be particularly important during early embryonic cell divisions when the spindle checkpoint cannot be activated by unattached chromosomes (Gerhart et al., 1984; Hara et al., 1980; Mara et al., 2019).

Figure 7. A graphical model of how H1.8 controls mitotic chromosome length

In the absence of H1.8, more condensins and topo II bind to more DNA loops of shorter length, resulting in longer and more individualized chromosomes (top). H1.8 limits chromatin levels of condensins and topo II to generate longer and thus fewer DNA loops, resulting in shorter and less individualized chromosomes (bottom).

Another possible reason for the suppressed individualization is related to the fact that oocyte chromosomes completely lose cohesion from arms at the end of meiosis I, while maintaining sister chromatid cohesion only at the centromeres (Lister et al., 2010). Normally, this centromeric cohesion is critical for supporting the kinetochore tension to establish bipolar attachment. During long natural arrest at meiotic metaphase II, these centromeres undergo cohesion fatigue, where centromeres prematurely separate. However, proper segregation may still be accomplished due to apparent inter-chromatids DNA linkages (Gruhn et al., 2019). Resolution of these DNA linkages may be prevented by H1.8-mediated suppression of condensin and TOP2A. However, our data do not eliminate the possibility that H1.8 plays a condensin and TOP2A independent role in regulating chromosome individualization. Regulation of Ki-67, which coats the surface of chromosomes, may be a good candidate of H1-mediated regulation (Cuylen et al., 2016; Gibson et al., 2019; White et al., 2016).

Linker histones are a dynamic component of chromatin (Misteli et al., 2000). Linker histone occupancy varies widely (Woodcock et al., 2006),

and can be controlled by both the linker histone variant and their posttranslational modifications (Christophorou et al., 2014; Hergeth and Schneider, 2015; Th'ng et al., 2005). We expect that changing the H1 stoichiometry on chromatin by changing the amount, subtype and/or affinity (e.g., through posttranslational modifications) of H1 affects the length and individualization of chromosomes through regulating the levels of condensins and topo II. In *Xenopus* embryos, exogenously added somatic linker histone variants, but not H1.8, are evicted from mitotic chromatin (Freedman and Heald, 2010). H1.8 is also more preferentially enriched on mitotic chromatin than on interphase chromatin in *Xenopus* egg extracts (Arimura et al., 2020), suggesting that loading of condensins and topo II to chromatin is actively limited by modulating mitotic H1 loading in oocytes and early embryos. Shortening chromosome length would reduce time and distance to complete chromosome segregation in rapidly dividing early embryonic cells. In addition, reducing the chromosome surface by shortening chromosomes may contribute to rapid completion of nuclear envelope formation. This is particularly important during early embryonic cell cycle in *Xenopus*, since

the nuclear envelope encapsulates each individual chromosome in late anaphase to form the karyomere, in which DNA replication can be accomplished (Lemaitre et al., 1998).

Linker histones serve important interphase roles through regulation of transcription (Izzo et al., 2008), and the epigenetic landscape of the chromatin determines the linker histone variant on chromatin (Izzo et al., 2013; Parseghian et al., 2001; Th'ng et al., 2005). Since H1.8 competitively inhibits

condensin binding in mitosis, it is tempting to speculate that linker histones also inhibit condensin II and cohesin binding in interphase. We suggest that local and global regulation of chromatin structure and function can be regulated by controlling differential expression of linker histone H1 variants and their modifications not just through promoting inter-nucleosomal interactions but also by controlling accessibility of SMC proteins and topo II.

Data availability

The Hi-C datasets used in the paper are available in the GEO under accession no. GSE164434.

Acknowledgments

We thank C. Zierhut for providing Δ H3-H4 depleted extracts for Hi-C analysis; S. Rankin, T. Hirano, Y. Azuma for sharing reagents; C. Jenness for anti-H1.8 antibodies; J. F. Martinez and M. P. Rout for their help with purifying TOP2A; A. North, C. Rico and K. Cialowicz at Bioimaging resource center (BIRC) for help with imaging; Y. Arimura, R. Heald, L. Mirny, A. Vannini and C. Zierhut and members of the Funabiki lab for helpful discussions. This work was supported by NIH grant R35 GM132111 to H.F. and NIH grant HG003143 to J.D. J.D. is an investigator of the Howard Hughes Medical Institute. A.V. is supported by a Cancer Research UK Programme Foundation (CR-UK C47547/A21536) and a Wellcome Trust Investigator Award (200818/Z/16/Z).

Author Contributions

P.C. conducted all experiments except for Hi-C library preparation, which was done by B.D. P.C., B.D., J.D. and H.F. analyzed Hi-C data. E.E.C. prepared human condensin I and II complexes. P.C., E.E.C., A.V. and H.F. analyzed *in vitro* condensin binding data. P.C. and H.F. designed the experiments and wrote the paper. B.D., E.E.C. and J.D. edited the paper. A.V. commented on the paper. H.F. oversaw the entire project.

Experimental Model and Subject Details

Xenopus laevis frogs

Mature female pigmented *X. laevis* frogs (NASCO-LM00535MX) were maintained in a temperature-controlled room (16-18°C) using a recirculating water system at The Rockefeller Comparative BioScience Center (CBC). Frogs were temporarily moved to a satellite facility for ovulation and egg-laying according to protocols approved by the IACUC (Protocol number 20031).

Method Details

Antibodies

H3 was detected with ab1791 (Abcam; 1 µg/ml for western blots). H2B was detected with ab1790 (Abcam; 1 µg/ml for western blots). α -tubulin was detected with T9026 (Sigma; 1:10000 for western blots).

H1.8 was detected using anti-H1M (H1.8) antibody (Jenness et al. 2018; 1 µg/ml for western blots). Anti-TOP2A was a gift from Y. Azuma (Ryu et al. 2010; 1 µg/ml for western blots and IF). Anti-CAPD2 was a gift from T. Hirano (Hirano, Kobayashi, and Hirano 1997; 2 µg/ml for western blots). Anti-CAPG2 was a gift from S. Rankin (4 µg/ml for IF, 2 µg/ml for western blots). xCAP-G (Zierhut et al., 2014) and xCAP-D3 custom antibodies were used at 2 µg/ml for western blots and 1 µg/ml for IF. CENP-A was detected using an antibody against N-terminal 50 amino acids of CENP-A (Wynne and Funabiki, 2015) and used at 4 µg/ml for IF.

xCAP-G antibody was also conjugated to Alexa488 using the Alexa488-NHS ester (Thermo-Fisher Scientific) using the manufacturer's instructions. The conjugated antibody was purified using the Sephadex G-25 in PD-10 desalting column (Cytiva). The

conjugated antibody was dialyzed into PBS+50% glycerol and stored in aliquots at -80°C after freezing with liquid nitrogen. The labelled antibody was used at 4 µg/ml for IF.

IRDye 680LT Goat anti-Mouse IgG (H+L), IRDye 680LT Goat anti-Rabbit IgG (H+L), IRDye 800CW Goat anti-Mouse IgG (H+L) and IRDye 800CW Goat anti-Rabbit IgG (H+L) were used at 1:15000 (LI-COR Biosciences) dilution for western blots. Alexa 488, Alexa 555 and Alexa 647 conjugated secondary antibodies (Jackson ImmunoResearch) were used for IF.

Antibody production

xCAP-D3 C-terminal peptide (CRQRISGKAPLKPSN) was synthesized at The Rockefeller University Proteomics Resource Center). The peptide was then coupled to the keyhole limpet hemocyanin according to the manufacturer's protocol (Thermo-Fisher Scientific) and used to immunize rabbits (Cocalico Biologicals). Antibody was purified from the immunized rabbit sera using affinity purification against the same peptide coupled to SulfoLink resin (Thermo-Fisher Scientific). The antibody was dialyzed into PBS+ 50% glycerol and stored with the addition of 0.05% sodium azide.

Xenopus egg extracts and Immunodepletion

Cytostatic Factor (CSF) arrested *X. laevis* egg extracts were generated as previously described (Murray, 1991). To generate replicated mitotic chromosomes, 0.3 mM CaCl₂ was added to CSF arrested extracts containing *X. laevis* sperm to cycle the extracts into interphase at 20°C. Ninety minutes after adding CaCl₂, half the volume of fresh CSF extract and 40 nM of the non-degradable cyclin BΔ90 fragment were added to interphase extracts to induce mitotic entry (Glotzer et al., 1991; Holloway et al., 1993). After 60 min of incubation, extracts were processed for morphological and biochemical assessments. For all experiments involving immunofluorescence, 10 nM nocodazole was added along with the cyclin BΔ90.

For immunodepletions of 50-100 µl extracts, 250 µg/ml antibodies were conjugated to Protein-A coupled Dynabeads (Thermo Fisher Scientific) either at room temperature for 60 min or overnight at 4 °C. Mock (IgG) and H1.8 (H1) antibody beads were crosslinked using 4 mM BS₃ (Thermo Fisher Scientific) at room temperature for 45 min and quenched using 10 mM Tris-HCl (Sigma). All antibody beads were washed extensively using Sperm Dilution Buffer (SDB; 10 mM HEPES, 1mM

MgCl₂, 100 mM KCl, 150 mM Sucrose) and separated from the buffer using a magnet before addition of extract. H1.8 depletions (ΔH1) were performed with two 45 min rounds of depletion at 4 °C using 2 volumes of antibody-coupled beads for each round. For condensin I and condensin II depletions, 1.5-2 volumes of xCAP-G or xCAP-D3 antibody-coupled beads were used in a single round for depletion for 60 min at 4 °C. For double depletion of condensin I and II, a single round of depletion using 1.5 volume each of xCAP-G and xCAP-D3 antibody-coupled beads was performed. For TopoII depletions (ΔTOP2A), a single round of depletion was performed using 1.2 volume of anti-TopoIIα coupled antibody beads for 60 min at 4 °C. After the incubations, the beads were separated using a magnet.

Western blots

For total egg extract samples, 1 µl sample was added to 25 µl 1x sample buffer (50 mM Tris-HCl pH 6.8, 2 % SDS, 10% Glycerol, 2.5 % β-mercaptoethanol) and boiled for 10 min. Samples were spun at 8000 rpm for 3 min before gel electrophoresis and overnight transfer at 4 °C. Blotting membranes were blocked with 4 % powdered skim-milk (Difco). Primary and secondary antibodies were diluted in LICOR Odyssey blocking buffer-PBS (LI-COR Biotechnology). Western blots were imaged on a LICOR Odyssey. Quantifications were done using Image-J.

Hi-C

Standard samples

10⁶ *X. laevis* sperm nuclei were added to 150 µl interphase extract and allowed to replicate at 21 °C for 90 min. The extracts were cycled back into mitosis by adding 100 µl CSF extract, 40 nM of the non-degradable cyclin BΔ90 and 10 µM nocodazole (Sigma). After 60 min at metaphase, the samples were diluted into 12 ml of fixing solution (80 mM K-PIPES pH 6.8, 1 mM MgCl₂, 1 mM EGTA, 30% glycerol, 0.1% Triton X-100, 1% formaldehyde) and incubated at room temperature with rocking for 10 min. The samples were then quenched with 690 µl 2.5 M glycine for 5 min at room temperature. The samples were then placed on ice for 15 min and then centrifuged at 6000 g at 4 °C for 20 min. The pellet was then resuspended in 1 ml ice-cold DPBS. The tube was then centrifuged again at 13000 g for 20 min at 4 °C. The buffer was aspirated, and the pellet was frozen in liquid nitrogen and then stored at -80 °C.

Dispersed chromosome samples

The metaphase chromosome samples were prepared as above, but nocodazole was omitted. The metaphase extracts were diluted by adding 1.2 ml chromosome dilution buffer (10 mM K-HEPES pH 8, 200 mM KCl, 0.5 mM EGTA, 0.5 mM MgCl₂, 250 mM Sucrose) and incubated at room temperature for 8 min. 6 ml fixation buffer (5 mM K-HEPES pH 8, 0.1 mM EDTA, 100 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 0.5% Triton X-100, 20% glycerol, 1% formaldehyde) was added to the tube, mixed by rotation 10 min at room temperature. 420 μ l 2.5 M glycine was added to quench the formaldehyde and the mixture was incubated for 5 min at room temperature. The samples were then placed on ice for 15 min and then centrifuged at 6500 g at 4 °C for 20 min. The pellet was then resuspended in 1 ml ice-cold DPBS. The tube was then centrifuged again at 13000 g for 20 min at 4 °C. The buffer was aspirated, and the pellet was frozen in liquid nitrogen and then stored at -80 °C.

Two biological replicates were performed for each sample, and they confirmed similar behavior among the replicates.

Library prep and sequencing

Hi-C protocol was performed as previously described (Belaghzal et al., 2017), with exception that cell disruption by douncing was omitted. Briefly, pellets were digested by DpnII overnight at 37 °C prior to biotin fill-in with biotin-14-dATP for 4 h at 23 °C. After ligation at 16 °C for 4 h, crosslinking was reversed by proteinase K at 65°C overnight. Purified ligation products were sonicated with 200 bp average size, followed by 100-350 bp size selection. End repair was performed on size selected ligation products, prior to purifying biotin tagged DNA fragments with streptavidin beads. A-tailing was done on the purified DNA fragments followed by Illumina Truseq adapter ligation. Hi-C library was finished by PCR amplification and purification to remove PCR primers. Final library was sequenced on Illumina HiSeq 4000 with PE50.

Hi-C Data Processing

Hi-C fastq files were mapped to the *Xenopus laevis* 9.2 genome with the distiller-nf pipeline (<https://github.com/open2c/distiller-nf>). The reads were aligned with bwa-mem, afterwards duplicate reads were filtered out. These valid pair reads were aggregated in genomic bins of 10, 25, 50, 100, 250, 500kb using the cooler format (Abdennur and Mirny,

2019). Cooler files were balanced using Iterative balancing correction (Imakaev et al., 2012), ignoring first two diagonals to avoid artifacts within the first bin such as re-ligation products. Contact heatmaps from balanced cooler files were viewed and exported with Hiclass (Kerpedjiev et al., 2018).

Contact Probability ($P(s)$) and derivatives

Contacts probability were calculated by contact frequency (P) as function of genomic distance (s). Interaction pairs were selected for genomic distance from 1kb till 100mb binned at log-scale. Within each genomic bin observed number of interactions were divided by total possible number of interactions within the bin. Distance decay plots were normalized by total number interactions, derivative plots were made from corresponding $P(s)$. The derivative plots plotted in **Figure 4C, 4D, Figure S6D** were drawn using LOESS smoothing.

Immunofluorescence

Immunofluorescence was performed according to previously published protocols (Desai et al., 1998). 10 μ l metaphase extracts containing chromosomes were diluted into 2 ml of fixing solution (80 mM K-PIPES pH 6.8, 1 mM MgCl₂, 1 mM EGTA, 30% glycerol, 0.1% Triton X-100, 2% formaldehyde) and incubated at room temperature for 7 min. The fixed chromosomes were then laid onto a cushion (80 mM K-PIPES pH 6.8, 1 mM MgCl₂, 1 mM EGTA, 50% glycerol) with a coverslip placed at the bottom of the tube and centrifuged at 5000g for 15 min at 18 °C in a swinging bucket rotor. The coverslips were recovered and fixed with methanol (-20 °C) for 4 min. The coverslips were then blocked overnight with antibody dilution buffer (50 mM Tris-Cl pH 7.5, 150 mM NaCl, 2% BSA). Primary and secondary antibodies were diluted in antibody dilution buffer and sealed in Prolong Gold AntiFade mounting media (Thermo-Fisher Scientific).

For coverslips stained with Alexa488-anti-CAP-G antibody (**Figure 1C, 3E, 5D, 5E**), coverslips stained with primary and secondary antibodies were washed three times with PBS-T (1x PBS +0.5% Tween-20). Then, they were blocked with 100 μ g/ml rabbit IgG or 30 min and were incubated with Alexa488-anti-xCAP-G antibody without any washing steps in between. The coverslips were then washed three times with PBS-T and then sealed in Prolong Gold AntiFade mounting media (Thermo-Fisher Scientific).

Chromosome individualization

Chromosome dilution was performed as before with some modifications (Funabiki and Murray, 2000). 40 μ l Chromosome Dilution Buffer (10 mM K-HEPES pH 8, 200 mM KCl, 0.5 mM EGTA, 0.5 mM $MgCl_2$, 250 mM Sucrose) was added to 10 μ l metaphase extract containing chromosomes and incubated at room temperature for 8 min. 200 μ l fixation buffer (5 mM K-HEPES pH 8, 0.1 mM EDTA, 100 mM NaCl, 2 mM KCl, 1 mM $MgCl_2$, 2 mM $CaCl_2$, 0.5% Triton X-100, 20% glycerol, 2% formaldehyde) was added to the tube and incubated for 10 min at room temperature. The samples were laid over a cushion (5 mM K-HEPES pH 8, 0.1 mM EDTA, 100 mM NaCl, 2 mM KCl, 1 mM $MgCl_2$, 2 mM $CaCl_2$, 50 % glycerol) with a coverslip placed under the cushion and centrifuged at 7000g for 20 min at 18 °C in a swinging bucket rotor. The coverslips were recovered and fixed with ice-cold methanol for 4 min, washed extensively and blocked overnight with antibody dilution buffer (50 mM Tris-Cl pH 7.5, 150 mM NaCl, 2% BSA). CENP-A immunofluorescence was performed on these coverslips for **Figure 5B, 5C, S8D, E**.

Chromosome purification

One volume of metaphase extracts with ~3000/ μ l sperm nuclei was diluted into 3 volumes of DB2 (10 mM K-HEPES, 50 mM β -glycerophosphate, 50 mM NaF, 20 mM EGTA, 2 mM EDTA, 0.5 mM spermine, 1 mM phenylmethylsulfonyl fluoride, 200 mM sucrose) and laid over 1 ml cushion (DB2 with 50% sucrose). The tube was centrifuged in a swinging bucket rotor at 10,000g for 30 min at 4 °C. Most of the cushion was aspirated and the pellet was resuspended in the remaining solution and transferred to a fresh tube. The sample was centrifuged again at 13000g for 15 min at 4 °C. The pellet was then resuspended in 1x sample buffer and boiled for 10 min before being subject to gel electrophoresis.

Image acquisition and analysis

All the quantitative immunofluorescence imaging and some of the spindle imaging was performed on a DeltaVision Image Restoration microscope (Applied Precision) which is a wide-field inverted microscope equipped with a pco.edge sCMOS camera (pco). The immunofluorescence and surface area measurement samples were imaged with z-sections of 200 nm width with a 100x (1.4 NA) objective and were processed with a iterative processive deconvolution algorithm using the SoftWoRx (Applied Precision). The dispersed chromosomes imaged for length measurements and chromosome individualization

were imaged in five 1 μ m z-sections with a 63x (1.33 NA) silicone oil objective.

More than twenty nuclei imaged for immunofluorescence and three-dimensional surface area quantification for most of the experiments for each condition. The differences in each experiment were analyzed using a two-tailed Mann-Whitney *U* test. The data in Figures **1D, 1G, 3D, 5D, 5E, 6E, 6F, Figure S9C** were combined data from multiple experiments, where the data were normalized to the control (Δ IgG) levels, whose medians were normalized to 1 for all conditions (Lord et al., 2020). The aggregate data were then analyzed using an unpaired Student's *t*-test.

For all the immunofluorescence quantifications, the maximum intensity single slice was selected, background subtraction was performed, and average intensities were calculated on a mask generated using the DNA signal. The analysis was performed using custom MATLAB (Mathworks) code available at https://github.com/pavancss/PC2021_microscopy. For surface area measurements, images were interpolated into stacks of 67 nm width. A surface mask was built in three-dimensional space and surface area and DNA signal was calculated using the regionprops3 MATLAB function. Only large objects (>10000 pixel³) were analyzed to compare significant fractions of each nucleus. The analysis was done automatically using custom MATLAB (Mathworks) code available at https://github.com/pavancss/PC2021_microscopy.

For the CENP-A foci counting in **Figure 5C and Figure S8D, E**, DNA and CENP-A were segmented by Otsu's thresholding algorithm. Each independent object in a binarized DNA image was treated as a single chromosomal mass and CENP-A foci were counted in each mass. This was done using custom MATLAB code available at https://github.com/pavancss/PC2021_microscopy.

For the categorization of unindividualized chromosomes in **Figure 6C**, a large area of coverslip was imaged in panels and all the observed DNA masses were counted and categorized as in **Figure S9B** in an unblinded fashion. The number of DNA masses counted in each condition in **Figure 6C** were: Δ IgG-100% TOP2A (201, 128, 146), Δ IgG-25% TOP2A (155, 177, 144), Δ H1-100% TOP2A (447, 135, 208), Δ H1-25% TOP2A (232, 187, 171).

For chromosome length measurements, > 54 chromosomes were measured in each experiment to ensure that a relatively even sampling of the 18 different chromosomes of each sperm nucleus was possible. Chromosome length measurements were done by manually tracing the chromosomes on a single maximum intensity slice in ImageJ 1.52p.

Mononucleosomes and nucleosome arrays

Nucleosome arrays were prepared as previously noted (Guse et al., 2011; Zierhut et al., 2014). The plasmid pAS696 which contains 19 repeats of the Widom 601 nucleosome position sequence (Lowary and Widom, 1998) was digested with EcoRI, XbaI, HaeII and DraI. The fragment containing the array was isolated using polyethylene glycol-based precipitation. The ends of the DNA fragment were filled in with dATP, dGTP, dCTP and Bio-16-dUTP (Chemcyte) using Klenow DNA polymerase (NEB) and purified using Sephadex G-50 Nick columns (Cytiva Biosciences).

Mononucleosomal DNA were prepared by digesting pAS696 using AvaI. The 196 bp fragment was isolated using polyethylene glycol-based precipitation. The ends of the fragment were filled in with dATP, dGTP, dTTP and Alexa647-aha-dCTP (Thermo-Fisher Scientific) using Klenow DNA polymerase (NEB) and purified using Sephadex G-50 Nick columns (Cytiva Biosciences).

For nucleosome deposition, 10 µg of DNA arrays or mononucleosomal DNA was mixed with equimolar amount of *X. laevis* H3-H4 tetramer and twice equimolar amount of *X. laevis* H2A-H2B dimers in 1x TE with 2 M NaCl. The mixture was added into in a Slide-A-Lyzer dialysis cassette (Thermo-Fisher Scientific) and placed into 500 ml High salt buffer (10 mM Tris-Cl pH 7.5 @ 4 °C, 2M NaCl, 1 mM EDTA, 5 mM β-mercaptoethanol, 0.01 % Triton X-100). Salt was reduced in a gradient by pumping in 2 L of Low salt buffer (10 mM Tris-Cl pH 7.5 at 4 °C, 100 mM NaCl, 1 mM EDTA, 5 mM β-mercaptoethanol, 0.01 % Triton X-100) at constant volume at 1 ml/min. The quality of the nucleosome arrays was ascertained by digesting the nucleosome arrays with AvaI overnight in low magnesium buffer (5 mM potassium acetate, 2 mM Tris-acetate, 0.5 mM magnesium acetate, 1 mM DTT, pH 7.9) and electrophoresed in a 5% polyacrylamide gel made in 0.5x TBE (45 mM Tris-borate, 1 mM EDTA). The mononucleosomes were assayed by direct electrophoresis.

Nucleosome binding assays

Nucleosome arrays were bound to M280 Streptavidin DynaBeads (Thermo-Fisher Scientific) in chromatin bead binding buffer (50 mM Tris-Cl pH 8, 150 mM NaCl, 0.25 mM EDTA, 0.05% Triton X-100, 2.5 % polyvinylalcohol) by shaking at 1300 rpm for 3.5 h. To block the Step tagged condensin complexes from binding the unconjugated streptavidin on the beads during the condensin pull downs, the beads were washed once in chromatin binding buffer and then incubated in 1 mM Biotin in chromatin binding buffer by shaking at 1300 rpm for 1 hour. The beads were then washed with chromatin binding buffer (50 mM Tris-Cl pH 8, 150 mM NaCl, 0.25 mM EDTA, 0.05% Triton X-100) three times, moved to a new tube, washed twice with SDB (10 mM HEPES, 1mM MgCl₂, 100 mM KCl, 150 mM Sucrose) and split into two tubes. SDB with 0.0008% poly-glutamic acid (Sigma)(Stein and Künzler, 1983) was mixed with 400 nM recombinant xH1.8 (buffer for control) and incubated for 5 min at room temperature. This mixture was incubated with the beads (half with buffer, half with xH1.8) with rotation at 16 °C. The beads were then washed 1x with SDB and 1x with binding buffer (10 mM HEPES pH 8, 40 mM NaCl, 2.5 mM MgCl₂, 0.5 mM DTT, 0.05% Triton X-100). Beads were washed 2x with binding buffer with indicated assay salt concentration and resuspended in binding buffer with 100 nM recombinant TOP2A, 380 nM human condensin I, condensin I Q loop mutant or 320 nM condensin II or condensin II Q loop mutant. The beads were rotated at room temperature for 30 min. Total reaction samples were taken and the beads were washed three times on a magnet in binding buffer and moved to a new tube. The beads were collected on a magnet and resuspended in 1x sample buffer (50 mM Tris-HCl pH 6.8, 2 % SDS, 10% Glycerol, 2.5 % β-mercaptoethanol) and boiled for 5 min. Gel electrophoresis was performed and the gels were stained with GelCode Blue Stain reagent (Thermo-Fisher Scientific).

Condensin Gel Shift assays

200 nM Alexa 647 labelled 196 bp mononucleosomes were mixed with 0.0008% poly-glutamic acid (Sigma)(Stein and Künzler, 1983) and half was mixed with 400 nM recombinant xH1.8 in 1x binding buffer (10 mM HEPES pH 8, 50 mM NaCl, 2.5 mM MgCl₂, 5 mM ATP, 0.5 mM DTT, 0.05% Triton X-100) and incubated for 30 min at room temperature. 100 nM of the mononucleosomes with or H1.8 were mixed with the indicated concentration of condensin I in 1x binding buffer at

4 °C for 30 min and subject to electrophoresis onto a 5 % polyacrylamide gel in 0.5x TBE at room temperature. The gels were imaged on a LICOR Odyssey (LI-COR Biotechnology). The binding curves were fitted using Graphpad Prism 8.4.3 using the sigmoidal binding curve option of the non-linear curve fitting.

Protein purification

H1.8

A pET51b vector expressing *Xenopus laevis* H1.8 with an N-terminal Strep-Tag II and C-terminal 6x Histidine-tag was a gift from Rebecca Heald (UC Berkeley). *E. coli* Rosetta2 (DE3 pLysS) cells containing expression plasmids were grown in TBG-M9 media (15 g/l tryptone, 7.5 g/l yeast extract, 5 g/l NaCl, 0.15 g/l MgSO₄, 1.5 g/l NH₄Cl, 3 g/l KH₂PO₄, 6 g/l Na₂HPO₄; 0.4% glucose) at 37 °C until they reach OD~0.6 and were supplemented with 1 mM isopropylthio-β-galactoside (IPTG) and grown at 18 °C for 14 h. Cells were collected and resuspend in lysis buffer (1x PBS, 500 mM NaCl, 10% glycerol, 20 mM Imidazole, 0.1% Triton X-100, 10 mM β-mercaptoethanol, 1 mM phenylmethylsulfonyl fluoride, 10 μg/ml leupeptin, 10 μg/ml pepstatin, 10 μg/ml chymostatin). All subsequent steps were carried out at 4 °C. After 30 min incubation, the cell suspension was sonicated and centrifuged at 45000g for 45 min at 4 °C. The supernatant was added to Ni-NTA beads (BioRad) and rotated for 60 min. The beads were then washed with Wash Buffer 1 (1x PBS, 20 mM imidazole, 500 mM NaCl, 4 mM β-mercaptoethanol, 10 mM ATP, 2.5 mM MgCl₂, cOmplete EDTA-free protease inhibitor cocktail-Roche). The beads were eluted with NTA elution buffer (1x PBS, 400 mM Imidazole, 500 mM NaCl). The correct fractions were collected and dialyzed into PBS supplement with 500 mM NaCl, concentrated using Amicon Ultra centrifugal filters (10k cutoff), flash frozen, aliquoted and stored at -80 °C.

TopoIIa

X. laevis TOP2A tagged with calmodulin binding protein (CBP) was purified from *P. pastoris* yeast as reported (Ryu et al., 2010) with some modifications. *Pichia pastoris* integrated with a CBP tagged TOP2A cassette under the influence of an alcohol oxidase (AOX) promoter (a gift from Yoshiaki Azuma) were grown in BMGY media (1% Yeast Extract, 2% Peptone, 100 mM potassium phosphate pH 6, 1.34% Yeast Nitrogen Base, 4x10⁻⁵% biotin, 1% glycerol) containing 50 μg/ml G418 (Thermo-Fisher

Scientific) at 30 °C until OD~4.0. The cells were collected by centrifugation and split into BMMY media (1% Yeast Extract, 2% Peptone, 100 mM potassium phosphate pH 6, 1.34% Yeast Nitrogen Base, 4x10⁻⁵% biotin, 0.5% methanol) and grown at 22 °C for 14 h. The cells were collected, packed into a syringe and extruded into liquid nitrogen in the form of noodles. These frozen noodles were lysed using a Retsch PM100 cryomill (Retsch) with continuous liquid nitrogen cooling. The cryomilled cells were then resuspended in Lysis Buffer (150mM NaCl, 18mM β-glycerophosphate, 1mM MgCl₂, 40mM HEPES (pH 7.8), 5% glycerol, 0.1% Triton X-100, 1mM DTT, cOmplete EDTA-free protease inhibitor tablet) and sonicated on ice. The cells were centrifuged at 35,000g for 45 min at 4 °C. 2 mM CaCl₂ was added to the supernatant along with Calmodulin-sepharose beads (Stratagene) and the mixture was incubated at 4 °C for 120 min. The beads were then washed with ATP-Wash Buffer (Lysis Buffer+ 5mM MgCl₂, 2mM CaCl₂, 1mM ATP), Wash Buffer 1 (Lysis Buffer + 2 mM CaCl₂), Wash Buffer 2 (300mM NaCl, 1mM MgCl₂, 2mM CaCl₂, 20mM HEPES (pH 7.8), 5% Glycerol, 1mM DTT) and then eluted into Elution Buffer (300mM NaCl, 1mM MgCl₂, 5mM EGTA, 20mM HEPES (pH 7.8), 5% glycerol, 1mM DTT).

The eluted protein was then passed through a MonoQ anion exchange column (Cytiva) on an AKTA-FPLC (Cytiva) to separate co-purified DNA. The flowthrough was then digested with TEV protease to cleave the CBP tag and then loaded on a HiTrap Heparin HP column (Cytiva) on an AKTA-FPLC and eluted using a salt gradient of 150 mM NaCl to 1 M NaCl. The selected fractions were then loaded on a Superose 6 gel filtration column (Cytiva) and eluted in Freezing buffer (250mM NaCl, 1mM MgCl₂, 20mM HEPES pH 7.8, 5% glycerol, 1mM DTT). The protein was then concentrated and frozen in aliquots at -80 °C.

Condensins

Human condensin complexes were purified as described previously (Kong et al., 2020). Briefly, the five subunits of human condensin I and II, sub-complexes and Q-loop mutations and were assembled into biGBac vectors (Weissmann et al., 2016) to create baculovirus for protein expression in HighFive insect cells. Cell were lysed in condensin purification buffer (20 mM HEPES [pH 8], 300 mM KCl, 5 mM MgCl₂, 1 mM DTT, 10% glycerol) supplemented with Pierce protease inhibitor EDTA-free tablet (Thermo Scientific) and Benzonase

(Sigma). Cleared lysate was loaded on to a StrepTrap HP (GE), washed with condensin purification buffer and eluted with condensin purification buffer supplemented with 5 mM Desthiobiotin (Sigma). Protein containing fractions were pooled, diluted 2-fold with Buffer A (20 mM HEPES [pH 8], 5 mM MgCl₂, 5% glycerol, 1 mM DTT), loaded on to HiTrap Heparin HP column (GE), washed with Buffer A with 250 mM NaCl, then eluted with buffer A with 500 mM NaCl. Finally, size exclusion chromatography was performed using Condensin purification buffer and a Superose 6 16/70 or increase 10/300 column (GE).

Mass spectrometry

Sperm chromosomes were purified as previously described (Funabiki and Murray, 2000). Briefly, extracts containing 8000/μl sperm nuclei were replicated along with 5 mM biotin-dUTP for 90 min and cycled back into metaphase with the addition of 1 volume of fresh CSF depleted (correspondingly

ΔIgG or ΔH1). After 60 min in metaphase, these chromosomes were diluted in 3 volumes of DB (10 mM K-HEPES [pH 7.6], 100 mM KCl, 2 mM EDTA, 0.5 mM EGTA, 0.5 mM spermine, 250 mM Sucrose, 1 mM PMSF, and 10 μg/ml each of leupeptin, pepstatin, and chymostatin) and centrifuged through a 60DB cushion (DB with 60 % (w/v) sucrose). The collected chromosome enriched pellet was then incubated with 15 μl streptavidin coupled DynaBeads (M280) and rotated at 4 °C for 2 h. The beads were then collected, washed and boiled in sample buffer before running on a 6 % polyacrylamide gel for 10 min. The gel was stained with commassie blue and the protein containing gel fragments were cut out and processed for mass spectrometry. The mass spectrometry was performed at the Rockefeller University Proteomics Resource Center as previously described (Zierhut et al., 2014), but the peptides were queried against the *X. laevis* database (Wühr et al., 2014) using the MaxQuant software (Max-Planck Institute).

References

- Abdennur, N., and Mirny, L.A. (2019). Cooler: scalable storage for Hi-C data and other genomically labeled arrays. *Bioinformatics* 36, 311–316.
- Abramo, K., Valton, A., Venev, S. V, Ozadam, H., Fox, A.N., and Dekker, J. (2019). A chromosome folding intermediate at the condensin-to-cohesin transition during telophase. *Nat. Cell Biol.* 21, 1393–1402.
- Adachi, Y., Luke, M., and Laemmli, U.K. (1991). Chromosome assembly in vitro: topoisomerase II is required for condensation. *Cell* 64, 137–148.
- Alipour, E., and Marko, J.F. (2012). Self-organization of domain structures by DNA-loop-extruding enzymes. *Nucleic Acids Res.* 40, 11202–11212.
- Arimura, Y., Shih, R.M., Froom, R., and Funabiki, H. (2020). Nucleosome structural variations in interphase and metaphase chromosomes. *BioRxiv* 2020.11.12.380386.
- Bakhrebah, M., Zhang, T., Mann, J.R., Kalitsis, P., and Hudson, D.F. (2015). Disruption of a conserved CAP-D3 threonine alters condensin loading on mitotic chromosomes leading to chromosome hypercondensation. *J. Biol. Chem.* 290, 6156–6167.
- Banigan, E.J., and Mirny, L.A. (2020). Loop extrusion: theory meets single-molecule experiments. *Curr. Opin. Cell Biol.* 64, 124–138.
- Bauer, C.R., Hartl, T.A., and Bosco, G. (2012). ΔIgG or ΔH1). After 60 min in metaphase, these chromosomes were diluted in 3 volumes of DB (10 mM K-HEPES [pH 7.6], 100 mM KCl, 2 mM EDTA, 0.5 mM EGTA, 0.5 mM spermine, 250 mM Sucrose, 1 mM PMSF, and 10 μg/ml each of leupeptin, pepstatin, and chymostatin) and centrifuged through a 60DB cushion (DB with 60 % (w/v) sucrose). The collected chromosome enriched pellet was then incubated with 15 μl streptavidin coupled DynaBeads (M280) and rotated at 4 °C for 2 h. The beads were then collected, washed and boiled in sample buffer before running on a 6 % polyacrylamide gel for 10 min. The gel was stained with commassie blue and the protein containing gel fragments were cut out and processed for mass spectrometry. The mass spectrometry was performed at the Rockefeller University Proteomics Resource Center as previously described (Zierhut et al., 2014), but the peptides were queried against the *X. laevis* database (Wühr et al., 2014) using the MaxQuant software (Max-Planck Institute).
- Condensin II Promotes the Formation of Chromosome Territories by Inducing Axial Compaction of Polyploid Interphase Chromosomes. *PLoS Genet.* 8, 1–12.
- Bednar, J., Garcia-Saez, I., Boopathi, R., Cutter, A.R., Papai, G., Reymer, A., Syed, S.H., Lone, I.N., Tonchev, O., Crucifix, C., et al. (2017). Structure and Dynamics of a 197 bp Nucleosome in Complex with Linker Histone H1. *Mol. Cell* 66, 384-397.e8.
- Belaghzal, H., Dekker, J., and Gibcus, J.H. (2017). Hi-C 2.0: An optimized Hi-C procedure for high-resolution genome-wide mapping of chromosome conformation. *Methods* 123, 56–65.
- Bernad, R., Sánchez, P., Rivera, T., Rodríguez-corsino, M., Boyarchuk, E., Vassias, I., Ray-gallet, D., Arnautov, A., Dasso, M., Almouzni, G., et al. (2011). HJURP and condensin II are required for CENP-A assembly. *192*, 569–582.
- Brahmachari, S., and Marko, J.F. (2019). Chromosome disentanglement driven via optimal compaction of loop-extruded brush structures. *Proc. Natl. Acad. Sci. U. S. A.* 116, 24956–24965.
- Brandão, H.B., Paul, P., van den Berg, A.A., Rudner, D.Z., Wang, X., and Mirny, L.A. (2019). RNA polymerases as moving barriers to condensin loop extrusion. *Proc. Natl. Acad. Sci.* 116, 20489–20499.
- Canela, A., Maman, Y., Jung, S., Wong, N., Callen, E., Day, A., Kieffer-Kwon, K.R., Pekowska, A., Zhang, H., Rao, S.S.P., et al. (2017). Genome Organization Drives Chromosome Fragility. *Cell*

170, 507–521.e18.

- Canela, A., Maman, Y., Huang, S.N., Wutz, G., Tang, W., Zagnoli-Vieira, G., Callen, E., Wong, N., Day, A., Peters, J.-M., et al. (2019). Topoisomerase II-Induced Chromosome Breakage and Translocation Is Determined by Chromosome Architecture and Transcriptional Activity. *Mol. Cell* 1–15.
- Chereji, R. V., Bryson, T.D., and Henikoff, S. (2019). Quantitative MNase-seq accurately maps nucleosome occupancy levels. *Genome Biol.* 20, 198.
- Christophorou, M.A., Castelo-Branco, G., Halley-Stott, R.P., Oliveira, C.S., Loos, R., Radzisheuskaya, A., Mowen, K.A., Bertone, P., Silva, J.C.R., Zernicka-Goetz, M., et al. (2014). Citrullination regulates pluripotency and histone H1 binding to chromatin. *Nature* 507, 104–108.
- Cremer, T., and Cremer, M. (2010). Chromosome territories. *Cold Spring Harb. Perspect. Biol.* 2, 1–23.
- Cutts, E.E., and Vannini, A. (2020). Condensin complexes: understanding loop extrusion one conformational change at a time. *Biochem. Soc. Trans.* 48, 2089–2100.
- Cuvier, O., and Hirano, T. (2003). A role of topoisomerase II in linking DNA replication to chromosome condensation. *J. Cell Biol.* 160, 645–655.
- Cuylen, S., Blaukopf, C., Politi, A.Z., Müller-Reichert, T., Neumann, B., Poser, I., Ellenberg, J., Hyman, A.A., and Gerlich, D.W. (2016). Ki-67 acts as a biological surfactant to disperse mitotic chromosomes. *Nature* 535, 308–312.
- Dasso, M., Dimitrov, S., and Wolffe, A.P. (1994). Nuclear assembly is independent of linker histones. *Proc. Natl. Acad. Sci. U. S. A.* 91, 12477–12481.
- Datta, S., Lecomte, L., and Haering, C.H. (2020). Structural insights into DNA loop extrusion by SMC protein complexes. *Curr. Opin. Struct. Biol.* 65, 102–109.
- Desai, A., Murray, A.W., Mitchison, T., and Walczak, C.E. (1998). Chapter 20 The Use of *Xenopus* Egg Extracts to Study Mitotic Spindle Assembly and Function in Vitro. *Methods Cell Biol.* 61, 385–412.
- Du, Z., Zheng, H., Huang, B., Ma, R., Wu, J., Zhang, X., He, J., Xiang, Y., Wang, Q., Li, Y., et al. (2017). Allelic reprogramming of 3D chromatin architecture during early mammalian development. *Nature* 547, 232–235.
- Dworkin-Rastl, E., Kandolf, H., and Smith, R.C. (1994). The maternal histone H1 variant, H1M (B4 protein), is the predominant H1 histone in *Xenopus* pregastrula embryos. *Dev. Biol.* 161, 425–439.
- Earnshaw, W.C., and Laemmli, U.K. (1983). Architecture of metaphase chromosomes and chromosome scaffolds. *J. Cell Biol.* 96, 84–93.
- Elbatsh, A.M.O., Kim, E., Eeftens, J.M., Raaijmakers, J.A., van der Weide, R.H., García-Nieto, A., Bravo, S., Ganji, M., uit de Bos, J., Teunissen, H., et al. (2019). Distinct Roles for Condensin's Two ATPase Sites in Chromosome Condensation. *Mol. Cell* 76, 724–737.e5.
- Eltsov, M., MacLellan, K.M., Maeshima, K., Frangakis, A.S., and Dubochet, J. (2008). Analysis of cryo-electron microscopy images does not support the existence of 30-nm chromatin fibers in mitotic chromosomes in situ. *Proc. Natl. Acad. Sci.* 105, 19732–19737.
- Farcas, A., Uluocak, P., Helmhart, W., and Nasmyth, K.A. (2011). Cohesin's concatenation of sister DNAs maintains their intertwining. *Mol. Cell* 44, 97–107.
- Farr, C.J., Antoniou-Kourounioti, M., Mimmack, M.L., Volkov, A., and Porter, A.C.G. (2014). The α isoform of topoisomerase II is required for hypercompaction of mitotic chromosomes in human cells. *Nucleic Acids Res.* 42, 4414–4426.
- Finch, J.T., and Klug, A. (1976). Solenoidal model for superstructure in chromatin. *Proc. Natl. Acad. Sci.* 73, 1897–1901.
- Fitz-James, M.H., Tong, P., Pidoux, A.L., Ozadam, H., Yang, L., White, S.A., Dekker, J., and Allshire, R.C. (2020). Large domains of heterochromatin direct the formation of short mitotic chromosome loops. *Elife* 9.
- Freedman, B.S., and Heald, R. (2010). Functional comparison of H1 histones in *Xenopus* reveals isoform-specific regulation by Cdk1 and RanGTP. *Curr. Biol.* 20, 1048–1052.
- Funabiki, H., and Murray, A.W. (2000). The *Xenopus* chromokinesin Xkid is essential for metaphase chromosome alignment and must be degraded to allow anaphase chromosome movement. *Cell* 102, 411–424.
- Gandhi, R., Gillespie, P.J., and Hirano, T. (2006). Human Wapl Is a Cohesin-Binding Protein that Promotes Sister-Chromatid Resolution in Mitotic Prophase. *Curr. Biol.* 16, 2406–2417.
- Ganji, M., Shaltiel, I.A., Bisht, S., Kim, E., Kalichava, A., Haering, C.H., and Dekker, C. (2018). Real-time imaging of DNA loop extrusion by condensin. *Science* (80-.). 360, 102–105.
- Gassler, J., Brandão, H.B., Imakaev, M., Flyamer, I.M., Ladstätter, S., Bickmore, W.A., Peters, J.-M.,

- Mirny, L.A., and Tachibana, K. (2017). A mechanism of cohesin-dependent loop extrusion organizes zygotic genome architecture. *EMBO J.* 36, 3600–3618.
- Gerhart, J., Wu, M., and Kirschner, M. (1984). Cell cycle dynamics of an M-phase-specific cytoplasmic factor in *Xenopus laevis* oocytes and eggs. *J. Cell Biol.* 98, 1247–1255.
- Gibcus, J.H., Samejima, K., Goloborodko, A., Samejima, I., Naumova, N., Nuebler, J., Kanemaki, M.T., Xie, L., Paulson, J.R., Earnshaw, W.C., et al. (2018). A pathway for mitotic chromosome formation. *Science* (80-.). 359.
- Gibson, B.A., Doolittle, L.K., Schneider, M.W.G., Jensen, L.E., Gamarra, N., Henry, L., Gerlich, D.W., Redding, S., and Rosen, M.K. (2019). Organization of Chromatin by Intrinsic and Regulated Phase Separation. *Cell* 1–15.
- Glutzer, M., Murray, A.W., and Kirschner, M.W. (1991). Cyclin is degraded by the ubiquitin pathway. 349.
- Golfier, S., Quail, T., Kimura, H., and Brugués, J. (2020). Cohesin and condensin extrude DNA loops in a cell-cycle dependent manner. *Elife* 9.
- Goloborodko, A., Imakaev, M. V., Marko, J.F., and Mirny, L.A. (2016a). Compaction and segregation of sister chromatids via active loop extrusion. *Elife* 5, 1–16.
- Goloborodko, A., Marko, J.F., and Mirny, L.A. (2016b). Chromosome Compaction by Active Loop Extrusion. *Biophys. J.* 110, 2162–2168.
- Goundaroulis, D., Lieberman Aiden, E., and Stasiak, A. (2020). Chromatin Is Frequently Unknotted at the Megabase Scale. *Biophys. J.* 118, 2268–2279.
- Green, L.C., Kalitsis, P., Chang, T.M., Cipetic, M., Kim, J.H., Marshall, O., Turnbull, L., Whitchurch, C.B., Vagnarelli, P., Samejima, K., et al. (2012). Contrasting roles of condensin I and condensin II in mitotic chromosome formation. *J. Cell Sci.* 125, 1591–1604.
- Grigoryev, S.A., Bascom, G., Buckwalter, J.M., Schubert, M.B., Woodcock, C.L., and Schlick, T. (2016). Hierarchical looping of zigzag nucleosome chains in metaphase chromosomes. *Proc. Natl. Acad. Sci.* 113, 1238–1243.
- Gruhn, J.R., Zielinska, A.P., Shukla, V., Blanshard, R., Capalbo, A., Cimadomo, D., Nikiforov, D., Chan, A.C.-H., Newnham, L.J., Vogel, I., et al. (2019). Chromosome errors in human eggs shape natural fertility over reproductive life span. *Science* (80-.). 365, 1466–1469.
- Guacci, V., Hogan, E., and Koshland, D. (1994). Chromosome condensation and sister chromatid pairing in budding yeast. *J. Cell Biol.* 125, 517–530.
- Guse, A., Carroll, C.W., Moree, B., Fuller, C.J., and Straight, A.F. (2011). In vitro centromere and kinetochore assembly on defined chromatin templates. *Nature* 477, 354–358.
- Hara, K., Tydeman, P., and Kirschner, M. (1980). A cytoplasmic clock with the same period as the division cycle in *Xenopus* eggs. *Proc. Natl. Acad. Sci. U. S. A.* 77, 462–466.
- Hassler, M., Shaltiel, I.A., Kschonsak, M., Simon, B., Merkel, F., Thärichen, L., Bailey, H.J., Macošek, J., Bravo, S., Metz, J., et al. (2019). Structural Basis of an Asymmetric Condensin ATPase Cycle. *Mol. Cell* 74, 1175–1188.e9.
- Heald, R., and Gibeaux, R. (2018). ScienceDirect Subcellular scaling : does size matter for cell division ? *Curr. Opin. Cell Biol.* 52, 88–95.
- Hergeth, S.P., and Schneider, R. (2015). The H1 linker histones: multifunctional proteins beyond the nucleosomal core particle. *EMBO Rep.* 16, 1439–1453.
- Hirano, T., and Mitchison, T.J. (1991). Cell cycle control of higher-order chromatin assembly around naked DNA in vitro. *J. Cell Biol.* 115, 1479–1489.
- Hirano, T., and Mitchison, T.J. (1994). A heterodimeric coiled-coil protein required for mitotic chromosome condensation in vitro. *Cell* 79, 449–458.
- Hirano, T., Kobayashi, R., and Hirano, M. (1997). Condensins, Chromosome Condensation Protein Complexes Containing XCAP-C, XCAP-E and a *Xenopus* Homolog of the *Drosophila* Barren Protein. *Cell* 89, 511–521.
- Hirota, T., Gerlich, D.W., Koch, B., Ellenberg, J., and Peters, J.-M. (2004). Distinct functions of condensin I and II in mitotic chromosome assembly. *J. Cell Sci.* 117, 6435–6445.
- Holloway, S.L., Glutzer, M., King, R.W., and Murray, A.W. (1993). Anaphase is initiated by proteolysis rather than by the inactivation of maturation-promoting factor. *Cell* 73, 1393–1402.
- Hopfner, K.P., Karcher, A., Shin, D.S., Craig, L., Arthur, L.M., Carney, J.P., and Tainer, J.A. (2000). Structural biology of Rad50 ATPase: ATP-driven conformational control in DNA double-strand break repair and the ABC-ATPase superfamily. *Cell* 101, 789–800.
- Hsieh, T.-H.S., Cattoglio, C., Slobodyanyuk, E., Hansen, A.S., Rando, O.J., Tjian, R., and Darzacq, X. (2020). Resolving the 3D Landscape of Transcription-Linked Mammalian Chromatin Folding. *Mol. Cell* 78, 539–553.e8.

- Imakaev, M., Fudenberg, G., McCord, R.P., Naumova, N., Goloborodko, A., Lajoie, B.R., Dekker, J., and Mirny, L.A. (2012). Iterative correction of Hi-C data reveals hallmarks of chromosome organization. *Nat. Methods* 9, 999–1003.
- Izzo, A., Kamieniarz, K., and Schneider, R. (2008). The histone H1 family: Specific members, specific functions? *Biol. Chem.* 389, 333–343.
- Izzo, A., Kamieniarz-Gdula, K., Ramírez, F., Noreen, N., Kind, J., Manke, T., vanSteensel, B., and Schneider, R. (2013). The Genomic Landscape of the Somatic Linker Histone Subtypes H1.1 to H1.5 in Human Cells. *Cell Rep.* 3, 2142–2154.
- Jenness, C., Giunta, S., Müller, M.M., Kimura, H., Muir, T.W., and Funabiki, H. (2018). HELLS and CDCA7 comprise a bipartite nucleosome remodeling complex defective in ICF syndrome. *Proc. Natl. Acad. Sci.* 201717509.
- Kerpedjiev, P., Abdennur, N., Lekschas, F., McCallum, C., Dinkla, K., Strobelt, H., Lubert, J.M., Ouellette, S.B., Azhir, A., Kumar, N., et al. (2018). HiGlass: web-based visual exploration and analysis of genome interaction maps. *Genome Biol.* 19, 125.
- Kimura, K., Hirano, M., Kobayashi, R., and Hirano, T. (1998). Phosphorylation and activation of 13S condensin by Cdc2 in vitro. *Science* 282, 487–490.
- Kimura, K., Cuvier, O., and Hirano, T. (2001). Chromosome condensation by a human condensin complex in *Xenopus* egg extracts. *J. Biol. Chem.* 276, 5417–5420.
- Kinoshita, K., and Hirano, T. (2017). Dynamic organization of mitotic chromosomes. *Curr. Opin. Cell Biol.* 46, 46–53.
- Kong, M., Cutts, E.E., Pan, D., Beuron, F., Kaliyappan, T., Xue, C., Morris, E.P., Musacchio, A., Vannini, A., and Greene, E.C. (2020). Human Condensin I and II Drive Extensive ATP-Dependent Compaction of Nucleosome-Bound DNA. *Mol. Cell* 683540.
- Krietenstein, N., Abraham, S., Venev, S. V., Abdennur, N., Gibcus, J.H., Hsieh, T.-H.S., Parsi, K.M., Yang, L., Maehr, R., Mirny, L.A., et al. (2020). Ultrastructural Details of Mammalian Chromosome Architecture. *Mol. Cell* 78, 554–565.e7.
- Krishnan, S., Smits, A.H., Vermeulen, M., and Reinberg, D. (2017). Phospho-H1 Decorates the Inter-chromatid Axis and Is Evicted along with Shugoshin by SET during Mitosis. *Mol. Cell* 67, 579–593.e6.
- Kschonsak, M., and Haering, C.H. (2015). Shaping mitotic chromosomes: From classical concepts to molecular mechanisms. *BioEssays* 37, 755–766.
- Ladouceur, A.-M., Dorn, J.F., and Maddox, P.S. (2015). Mitotic chromosome length scales in response to both cell and nuclear size. *J. Cell Biol.* 209, 645–652.
- Ladouceur, A.-M., Ranjan, R., Smith, L., Fadero, T., Heppert, J., Goldstein, B., Maddox, A.S., and Maddox, P.S. (2017). CENP-A and topoisomerase-II antagonistically affect chromosome length. *J. Cell Biol.* 216, jcb.201608084.
- Lane, A.B., Giménez-Abián, J.F., and Clarke, D.J. (2013). A novel chromatin tether domain controls topoisomerase IIα dynamics and mitotic chromosome formation. *J. Cell Biol.* 203, 471–486.
- Lee, W., Tillo, D., Bray, N., Morse, R.H., Davis, R.W., Hughes, T.R., and Nislow, C. (2007). A high-resolution atlas of nucleosome occupancy in yeast. *Nat. Genet.* 39, 1235–1244.
- Lemaitre, J.M., Géraud, G., and Méchali, M. (1998). Dynamics of the genome during early *Xenopus laevis* development: karyomeres as independent units of replication. *J. Cell Biol.* 142, 1159–1166.
- Li, W., Chen, P., Yu, J., Dong, L., Liang, D., Feng, J., Yan, J., Wang, P.Y., Li, Q., Zhang, Z., et al. (2016). FACT Remodels the Tetranucleosomal Unit of Chromatin Fibers for Gene Transcription. *Mol. Cell* 64, 120–133.
- Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009). Comprehensive Mapping of Long-Range Interactions Reveals Folding Principles of the Human Genome. *Science* (80-.). 326, 289–293.
- Lister, L.M., Kouznetsova, A., Hyslop, L.A., Kalleas, D., Pace, S.L., Barel, J.C., Nathan, A., Floros, V., Adelfalk, C., Watanabe, Y., et al. (2010). Age-related meiotic segregation errors in mammalian oocytes are preceded by depletion of cohesin and Sgo2. *Curr. Biol.* 20, 1511–1521.
- Lord, S.J., Velle, K.B., Mullins, R.D., and Fritz-Laylin, L.K. (2020). SuperPlots: Communicating reproducibility and variability in cell biology. *J. Cell Biol.* 219.
- Lowary, P., and Widom, J. (1998). New DNA sequence rules for high affinity binding to histone octamer and sequence-directed nucleosome positioning. *J. Mol. Biol.* 276, 19–42.
- Löwe, J., Cordell, S.C., and Van Den Ent, F. (2001). Crystal structure of the SMC head domain: An ABC ATPase with 900 residues antiparallel coiled-coil inserted. *J. Mol. Biol.* 306, 25–35.
- Mara, L., Paim, G., and Fitzharris, G. (2019). Cell-

Size-Independent Spindle Checkpoint Failure Underlies Chromosome Segregation Error in Mouse Report Cell-Size-Independent Spindle Checkpoint Failure Underlies Chromosome Segregation Error. 865–873.

Maresca, T.J., Freedman, B.S., and Heald, R. (2005). Histone H1 is essential for mitotic chromosome architecture and segregation in *Xenopus laevis* egg extracts. *J. Cell Biol.* *169*, 859–869.

Marko, J.F. (2008). Micromechanical studies of mitotic chromosomes. *Chromosom. Res.* *16*, 469–497.

Marko, J.F. (2009). Linking topology of tethered polymer rings with applications to chromosome segregation and estimation of the knotting length. *Phys. Rev. E - Stat. Nonlinear, Soft Matter Phys.* *79*, 1–16.

Micheli, G., Luzzatto, A.R., Carrì, M.T., de Capoa, A., and Pelliccia, F. (1993). Chromosome length and DNA loop size during early embryonic development of *Xenopus laevis*. *Chromosoma* *102*, 478–483.

Misteli, T., Gunjan, a, Hock, R., Bustin, M., and Brown, D.T. (2000). Dynamic binding of histone H1 to chromatin in living cells. *Nature* *408*, 877–881.

Murray, A.W. (1991). Chapter 30 Cell Cycle Extracts. In *Methods in Cell Biology*, pp. 581–605. Nagasaka, K., Hossain, M.J., Roberti, M.J., Ellenberg, J., and Hirota, T. (2016). Sister chromatid resolution is an intrinsic part of chromosome organization in prophase. *Nat. Cell Biol.* *18*, 692–699.

Nasmyth, K.A. (2001). Disseminating the Genome: Joining, Resolving, and Separating Sister Chromatids During Mitosis and Meiosis. *Annu. Rev. Genet.* *35*, 673–745.

Naumova, N., Imakaev, M., Fudenberg, G., Zhan, Y.Y.Y., Lajoie, B.R., Mirny, L.A., and Dekker, J. (2013). Organization of the Mitotic Chromosome. *Science* (80-.). *342*, 948–953.

Nielsen, C.F., Zhang, T., Barisic, M., Kalitsis, P., and Hudson, D.F. (2020). Topoisomerase IIa is essential for maintenance of mitotic chromosome structure. *Proc. Natl. Acad. Sci. U. S. A.* *117*.

Ohsumi, K., Katagiri, C., and Kishimoto, T. (1993). Chromosome condensation in *Xenopus* mitotic extracts without histone H1. *Science* (80-.). *262*, 2033–2035.

Ono, T., Losada, A., Hirano, M., Myers, M.P., Neuwald, A.F., and Hirano, T. (2003). Differential contributions of condensin I and condensin II to

mitotic chromosome architecture in vertebrate cells. *Cell* *115*, 109–121.

Ono, T., Fang, Y., Spector, D.L., and Hirano, T. (2004). Spatial and temporal regulation of condensins I and II in mitotic chromosome assembly in human cells. *Mol. Biol. Cell* *15*, 3296–3308.

Ou, H.D., Phan, S., Deerinck, T.J., Thor, A., Ellisman, M.H., and O'Shea, C.C. (2017).

ChromEMT: Visualizing 3D chromatin structure and compaction in interphase and mitotic cells. *Science* (80-.). *357*, eaag0025.

Parseghian, M.H., Newcomb, R.L., and Hamkalo, B.A. (2001). Distribution of somatic H1 subtypes is non-random on active vs. inactive chromatin II: Distribution in human adult fibroblasts. *J. Cell. Biochem.* *83*, 643–659.

Patel, L., Kang, R., Rosenberg, S.C., Qiu, Y., Raviram, R., Chee, S., Hu, R., Ren, B., Cole, F., and Corbett, K.D. (2019). Dynamic reorganization of the genome shapes the recombination landscape in meiotic prophase. *Nat. Struct. Mol. Biol.* *26*.

Paulson, J.R., and Laemmli, U.K. (1977). The structure of histone-depleted metaphase chromosomes. *Cell* *12*, 817–828.

Piskadlo, E., Tavares, A., and Oliveira, R.A. (2017). Metaphase chromosome structure is dynamically maintained by condensin I-directed DNA (de)catenation. *Elife* *6*, 1–22.

Potapova, T.A., Unruh, J.R., Yu, Z., Rancati, G., Li, H., Stampfer, M.R., and Gerton, J.L. (2019). Superresolution microscopy reveals linkages between ribosomal DNA on heterologous chromosomes. *J. Cell Biol.* *218*, 2492–2513.

Riggs, A.D. (1990). DNA methylation and late replication probably aid cell memory, and type I DNA reeling could aid chromosome folding and enhancer function. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* *326*, 285–297.

Rosin, L.F., Nguyen, S.C., and Joyce, E.F. (2018). Condensin II drives large-scale folding and spatial partitioning of interphase chromosomes in *Drosophila* nuclei. *PLoS Genet.* *14*, 1–26.

Ryu, H., Furuta, M., Kirkpatrick, D., Gygi, S.P., and Azuma, Y. (2010). PIASy-dependent SUMOylation regulates DNA topoisomerase IIa activity. *J. Cell Biol.* *191*, 783–784.

Samejima, K., Samejima, I., Vagnarelli, P., Ogawa, H., Vargiu, G., Kelly, D.A., Alves, F. de L., Kerr, A., Green, L.C., Hudson, D.F., et al. (2012). Mitotic chromosomes are compacted laterally by KIF4 and condensin and axially by topoisomerase IIa. *J. Cell Biol.* *199*, 755–770.

- Shakya, A., Park, S., Rana, N., and King, J.T. (2020). Liquid-Liquid Phase Separation of Histone Proteins in Cells: Role in Chromatin Organization. *Biophys. J.* 118, 753–764.
- Shamu, C.E., and Murray, A.W. (1992). Sister Chromatid Separation in Frog Egg Extracts Requires DNA Topoisomerase II Activity during Anaphase. 117, 921–934.
- Shechter, D., Nicklay, J.J., Chitta, R.K., Shabanowitz, J., Hunt, D.F., and David Allis, C. (2009). Analysis of histones in *Xenopus laevis* I. a distinct index of enriched variants and modifications exists in each cell type and is remodeled during developmental transitions. *J. Biol. Chem.* 284, 1064–1074.
- Shintomi, K., and Hirano, T. (2011). The relative ratio of condensin I to II determines chromosome shapes. *Genes Dev.* 25, 1464–1469.
- Shintomi, K., Inoue, F., Watanabe, H., Ohsumi, K., Ohsugi, M., and Hirano, T. (2017). Mitotic chromosome assembly despite nucleosome depletion in *Xenopus* egg extracts. *Science* (80-). 356, 1284–1287.
- Song, F., Chen, P., Sun, D., Wang, M., Dong, L., Liang, D., Xu, R.-M., Zhu, P., and Li, G. (2014). Cryo-EM Study of the Chromatin Fiber Reveals a Double Helix Twisted by Tetranucleosomal Units. *Science* (80-). 344, 376–380.
- Stein, A., and Künzler, P. (1983). Histone H5 can correctly align randomly arranged nucleosomes in a defined in vitro system. *Nature* 302, 548–550.
- Stigler, J., Çamdere, G., Koshland, D.E., and Greene, E.C. (2016). Single-Molecule Imaging Reveals a Collapsed Conformational State for DNA-Bound Cohesin. *Cell Rep.* 15, 988–998.
- Sun, M., Biggs, R., Hornick, J., and Marko, J.F. (2018). Condensin controls mitotic chromosome stiffness and stability without forming a structurally contiguous scaffold. *Chromosom. Res.* 26, 277–295.
- Sundin, O., and Varshavsky, A. (1981). Arrest of Segregation Leads to Accumulation of Highly Intertwined Catenated Dimers : Dissection of the Final Stages of SV40 DNA Replication. *Cell* 25, 659–669.
- Szabo, Q., Bantignies, F., and Cavalli, G. (2019). Principles of genome folding into topologically associating domains. *Sci. Adv.* 5.
- Tavares-Cadete, F., Norouzi, D., Dekker, B., Liu, Y., and Dekker, J. (2020). Multi-contact 3C reveals that the human genome during interphase is largely not entangled. *Nat. Struct. Mol. Biol.* 27, 1105–1114.
- Terakawa, T., Bisht, S., Eeftens, J.M., Dekker, C., Haering, C.H., and Greene, E.C. (2017). The condensin complex is a mechanochemical motor that translocates along DNA. *Science* 358, 672–676.
- Th'ng, J.P.H., Sung, R., Ye, M., and Hendzel, M.J. (2005). H1 Family Histones in the Nucleus. *J. Biol. Chem.* 280, 27809–27814.
- Thakurela, S., Garding, A., Jung, J., Schübeler, D., Burger, L., and Tiwari, V.K. (2013). Gene regulation and priming by topoisomerase II α in embryonic stem cells. *Nat. Commun.* 4.
- Toselli-Mollereau, E., Robellet, X., Fauque, L., Lemaire, S., Schiklenk, C., Klein, C., Hocquet, C., Legros, P., N'Guyen, L., Mouillard, L., et al. (2016). Nucleosome eviction in mitosis assists condensin loading and chromosome condensation. *EMBO J.* 35, 1565–1581.
- Umesono, K., Hiraoka, Y., Toda, T., and Yanagida, M. (1983). Visualization of chromosomes in mitotically arrested cells of the fission yeast *Schizosaccharomyces pombe*. *Curr. Genet.* 7, 123–128.
- Walther, N., Hossain, M.J., Politi, A.Z., Koch, B., Kueblbeck, M., Ødegård-Fougner, Ø., Lampe, M., and Ellenberg, J. (2018). A quantitative map of human Condensins provides new insights into mitotic chromosome architecture. *J. Cell Biol.* 217, 2309–2328.
- Weissmann, F., Petzold, G., VanderLinden, R., Huis in 't Veld, P.J., Brown, N.G., Lampert, F., Westermann, S., Stark, H., Schulman, B.A., and Peters, J.-M. (2016). biGBac enables rapid gene assembly for the expression of large multisubunit protein complexes. *Proc. Natl. Acad. Sci.* 113, E2564–E2569.
- White, A.E., Hieb, A.R., and Luger, K. (2016). A quantitative investigation of linker histone interactions with nucleosomes and chromatin. *Sci. Rep.* 6, 1–14.
- Woodcock, C.L., Skoultchi, A.I., and Fan, Y. (2006). Role of linker histone in chromatin structure and function: H1 stoichiometry and nucleosome repeat length. *Chromosom. Res.* 14, 17–25.
- Wühr, M., Freeman, R.M., Presler, M., Horb, M.E., Peshkin, L., Gygi, S., and Kirschner, M.W. (2014). Deep proteomics of the *Xenopus laevis* egg using an mRNA-derived reference database. *Curr. Biol.* 24, 1467–1475.
- Wynne, D.J., and Funabiki, H. (2015). Kinetochore function is controlled by a phosphodependent coexpansion of inner and outer components. *J. Cell Biol.* 210, 899–916.
- Yu, X., Davenport, J.W., Urtishak, K.A., Carillo, M.L., Gosai, S.J., Kolaris, C.P., Byl, J.A.W.,

Choppakatla et al., April 2021-preprint copy- BioRxiv

Rappaport, E.F., Osheroff, N., Gregory, B.D., et al. (2017). Genome-wide TOP2A DNA cleavage is biased toward translocated and highly transcribed loci. *Genome Res.* 27, 1238–1249.

Zhou, B., Jiang, J., Feng, H., Ghirlando, R., Xiao, S., Bai, Y., Zhou, B., Jiang, J., Feng, H., Ghirlando, R., et al. (2015). Structural Mechanisms of Nucleosome Recognition by Linker Histones
Article Structural Mechanisms of Nucleosome Recognition by Linker Histones. *Mol. Cell* 59, 628–638.

Zhou, B., Feng, H., Kale, S., Fox, T., Khant, H., de Val, N., Ghirlando, R., Panchenko, A.R., and Bai, Y. (2021). Distinct Structures and Dynamics of Chromatosomes with Different Human Linker Histone Isoforms. *Mol. Cell* 81, 166-182.e6.

Zierhut, C., Jenness, C., Kimura, H., and Funabiki, H. (2014). Nucleosomal regulation of chromatin composition and nuclear assembly revealed by histone depletion. *Nat. Struct. Mol. Biol.* 21, 617–625.

Figure S1. Linker histone H1.8 depletion does not lead to global accumulation of DNA-binding proteins (related to Figure 1)

A) Experimental scheme to incorporate Cy3-labelled nucleotides to use normalization of immunofluorescence signals on chromosomes. **B**) Quantification of Cy3-dUTP signals normalized to Hoechst 33342 signals, showing uniform normalization across two coverslips used for quantification of condensin I and TOP2A. The result also indicates no detectable defect in DNA replication in $\Delta H1$ extracts. **C**) CAP-G (condensin I) and TOP2A immunofluorescence signal levels on chromosomes normalized with Hoechst and incorporated Cy3-dUTP. Normalization using Hoechst 33342 and Cy3-dUTP/Cy3-dUTP signals were shown to be consistent across many experiments. **D**) Quantification of immunofluorescence signal levels of two proteins that prefer nucleosomes (RCC1 and Dasra A) and two that prefer to bind nucleosome-free DNA (Dppa2 and Xkid) (Zierhut et al., 2014). In B-D, distribution of signal intensity per chromosome cluster (dots), and median and SEM from one experiment are shown. This analysis was performed twice and the similar results were obtained. **E**) Protein abundance on metaphase sperm chromosomes purified from mock and H1.8 depleted extracts. Subunits from the SMC family complexes are labelled in pink. Core histones and their variants are in blue and linker histone variants are shown in red. The lines for 0.25x, 0.5x, 1, 2x and 4x abundance on $\Delta H1$ compared to ΔIgG chromosomes are shown. The mass spectrometry analysis was performed once. The number of nuclei imaged in each condition for each experiment in B, C and D are indicated above the figure.

Description	Gene ID	$\Delta H1/\Delta IgG$
Structural maintenance of chromosomes protein 4	SMC4	1.83
Structural maintenance of chromosomes protein 2	SMC2	1.87
DNA topoisomerase 2-alpha	TOP2A	3.13
Condensin complex subunit 3	NCAPG	2.54
Condensin complex subunit 1	NCAPD2	3.65
SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 5	SMARCA5	2.75
Condensin complex subunit 2	NCAPH	2.22
DNA replication licensing factor MCM7	MCM7	2.17
DNA replication licensing factor MCM5	MCM5	2.64
FACT complex subunit SPT16	SUPT16H	1.80
FACT complex subunit SSRP1	SSRP1	1.74
Shugoshin-like 1	SGOL1	1.79
Mediator of DNA damage checkpoint protein 1	MDC1	2.50
Fanconi anemia group I protein	FANCI	2.52
Histone H1.3	HIST1H1D	2.20

Table S1- High abundance proteins with the highest increases in $\Delta H1$ depleted chromosomes

High abundance proteins ($>10^7$ ion current) showing the highest increase in $\Delta H1$ depleted chromosomes.

Choppakatla et al., April 2021-preprint copy- BioRxiv

Description	Gene ID	$\Delta H1/\Delta IgG$
DNA damage-binding protein 1	DDB1	0.19
Histone H1oo	H1FOO	0.15
Bromodomain and WD repeat-containing protein 3	BRWD3	0.35
GTP-binding nuclear protein Ran	RAN	0.50
Lymphoid-specific helicase	HELLS	0.38
DNA topoisomerase 2-binding protein 1	TOPBP1	0.49
Telomere-associated protein RIF1	RIF1	0.01
Exportin-2	CSE1L	0.22
Probable ATP-dependent RNA helicase DDX6	DDX6	0.35
Polyadenylate-binding protein 1	PABPC1	0.25
Prohibitin	PHB	0.34
Importin subunit beta-1	KPNB1	0.10
RuvB-like 1	RUVBL1	0.02
Importin alpha 5.1	Imp 5.1	0.25
Nuclear pore complex protein Nup98-Nup96	NUP98	0.24

Table S2- High abundance proteins with the highest decreases in $\Delta H1$ depleted chromosomes

High abundance proteins ($>10^7$ ion current) showing the highest increase in $\Delta H1$ depleted chromosomes.

Figure S2- Recombinant human condensin I and *X. laevis* TOP2A are functional

A) Native PAGE gel analysis of nucleosome array beads loaded with or without H1.8 after digestion of the array with *Ava*I, which released monomers of nucleosome positioning sequence. A complete shift of monomer bands by H1.8 addition indicates the saturated occupancy of nucleosome and H1.8. **B)** Representative Hoechst (DNA) and CENP-A immunofluorescence images of chromosomes in indicated metaphase egg extracts after dilution, which disperses individualized chromosomes (left). Extracts depleted of condensin I (Δ CAP-G) were complemented with recombinant human condensin I and condensin I Q loop mutant protein complexes. Total egg extract western blots of depletion and the rescue conditions (right). **C)** Recombinant *X. laevis* TOP2A is functional in *Xenopus* egg extracts. Sperm nuclei were added to undepleted, TOP2A-depleted CSF extracts (Δ TOP2A), or Δ TOP2A extracts with purified recombinant TOP2A. Representative Hoechst (DNA) images of sperm are shown. In undepleted or Δ TOP2A extracts with supplemented TOP2A, proper sperm remodeling led to mitotic chromosomes formation. In Δ TOP2A extracts, sperm remodeling failed and sperm nuclei remained compact crescent-like shape. Bar, 20 μ m. **D)** Recombinant *X. laevis* TOP2A possesses decatenating activity in vitro. Agarose gel of kinetoplast decatenation assay. Recombinant TOP2A promoted ATP-dependent decatenation of kinetoplast DNA.

Choppakatla et al., April 2021-preprint copy- BioRxiv

Choppakatla et al., April 2021-preprint copy- BioRxiv

Figure S4- H1.8 inhibits condensin binding to mononucleosomes

Alexa647-labelled 196 bp mononucleosomes with or without H1.8 were incubated with indicated concentrations of condensin I and electrophoresed on a 5 % native PAGE. Alexa647-labeled DNAs are shown. Condensin I binding curves with mononucleosomes with and without H1.8, showing the large increase in binding constant in the presence of H1.8. The binding constants derived from fitting the data to a sigmoidal binding curve are shown. The data reproducibility was confirmed by another biological replicate.

Figure S5- Condensin II depletion does not affect chromosome length

A) Length of mitotic chromosomes in the indicated conditions after extract dilution showing no effect of CAP-D3 (condensin II) depletion in both mock and H1.8 depletion background. Each dot represents length of a single chromosome. Bars represent median $\pm$ 95% C.I. The length of >50 chromosomes were measured in each condition for every experiment. **B)** Total egg extract blots showing the partial depletions of condensin in the indicated conditions. The p-value in A was calculated by a two-tailed Mann-Whitney *U* test. The number of chromosomes imaged in each condition for each experiment in A are indicated above the figure.

Figure S6- Effects of H1, condensin I and condensin II depletion on mitotic genome folding

A) Contact probability decay curves of mitotic chromosomes in mock (ΔIgG) and H1-depleted (ΔH1) extracts from two different experiments showing the replicability of the Hi-C features. The solid and dashed lines are from biological replicates. **B)** Hi-C maps (left) binned to 500 kb, and contact probability curves (right) of single chromosomes showing the uniform effects of H1.8 depletion genome wide. **C)** Genome average contact probability decay curves upon H1.8, CAP-G and CAP-D3 depletions. **D)** Genome average contact probability derivative curves of metaphase chromosomes in undiluted mock (ΔIgG) and H1.8 depleted (ΔH1) extracts. The layer size estimates from the valley in the derivative plots are indicated. The dotted lines are a biological replicate.

Figure S7- H3-H4 depletion leads to even smaller loop sizes

A) Western blotting showing the depletion of H3-H4 in total egg extract using the H4K12ac antibody (Zierhut et al., 2014). **B)** Quantification of condensin I (CAP-G) immunofluorescence levels on chromosomes normalized to DNA in ΔH1 and ΔH3-H4 extracts. Each dot represents the average signal intensity of a single chromosome cluster (from one nucleus). Data plotted is median and 95 % C.I. >40 nuclei were quantified for each condition plotted. The statistical significance between ΔH1 and ΔH3-H4 was analyzed by a two-tailed Mann-Whitney U-test. **C)** Hi-C probability decay derivative plots estimating the loop sizes in ΔH1 and ΔH3-H4 extracts. The dashed lines indicate the loop sizes in the corresponding conditions and the dotted lines indicated the layer size. Two biological replicates for Hi-C on chromosomes from ΔH3-H4 extracts were performed and similar results were obtained from both experiments. The number of nuclei imaged in B are indicated above the figure.

Choppakatla et al., April 2021-preprint copy- BioRxiv

Figure S8. Regulation of chromosome individualization by topo II, condensins and H1.8 in *Xenopus* egg extracts

A) Schematic of ICRF-193 addition to check for requirement of topo II activity in individualizing chromosomes. Interphase nuclei were first formed in Δ IgG or Δ H1 extracts, topo II inhibitor ICRF-193 (50 μ M) was added to egg extracts, either together with corresponding depleted egg extracts and incubated for 50 min (ICRF-50 min), or 48 min after adding the depleted extracts, followed by 2 min incubation with ICRF-193 (ICRF-2 min). **B)** Metaphase extracts processed as A were diluted to disperse individualized chromosomes. Representative Hoechst images of chromosomes are shown. Bar, 20 μ m. **C)** Quantification of the Hoechst-stained area of chromosomes in B. Each dot represents the area of a single chromosome or a chromosome cluster. Large values indicate the extent of chromosome clusters. The box shows the 10th-90th percentile limits of the sample values. The number of DNA masses imaged in each condition are indicated above the figure. **D)** Distribution of CENP-A foci per DNA mass in the experiment shown in **Figure 5A-C**. Clusters of unresolved chromosomes are represented by higher numbers (>2) of CENP-A foci per DNA mass indicates clusters. No detectable CENP-A focus in Δ CAP-D3 (Δ condensin II) extracts is due to low CENP-A signal, reflecting the reported role of condensin II in CENP-A loading (Bernad et al., 2011). The number noted in each sample label is the number of chromosomes/clusters counted in each condition. **E)** Percent frequency of single chromosomes (chromosomes in DNA masses with <2 CENP-A foci) in the indicated conditions showing that the cutoff for number of CENP-A foci does not change the relative individualization measured. The number of DNA masses imaged in each condition are the same as in **Figure 5C**.

Figure S9. H1.8 suppresses over-individualization through condensins and topo II

A) Western blots of total egg extracts showing partial depletion of TOP2A. Left three lanes; dilution series of total mock-depleted extracts (Δ IgG) for signal quantitation. Middle Δ IgG lane; extracts with 25% level of TOP2A. right two Δ H1 lanes; extracts depleted of H1.8 with either 25% or 100% levels of TOP2A. **B)** Categories of chromosome clusters observed upon performing chromosome individualization assay. **C)** Quantification of TOP2A signal normalized to the DNA signal in the indicated conditions. Each gray, orange or violet dot represents the mean normalized TOP2A level in a single nucleus. The black squares represent the median of the TOP2A values of all the nuclei in a single experiment. Data plotted is mean and S.E.M. The p-value in C was calculated using an unpaired Student's t-test. The number of nuclei imaged in C in each condition are indicated above the figure.