Integrative Clinical and Molecular Characterization of Translocation Renal Cell Carcinoma

Ziad Bakouny^{1,2,3}, Ananthan Sadagopan¹, Praful Ravi¹, Nebiyu Y. Metaferia¹, Jiao Li¹, Shatha AbuHammad¹, Stephen Tang¹, Thomas Denize^{3,4}, Emma R. Garner¹, Xin Gao^{3,5}, David A. Braun^{1,2,3}, Laure Hirsch^{1,3}, John A. Steinharter¹, Gabrielle Bouchard¹, Emily Walton⁴, Destiny West⁴, Chris Labaki¹, Shaan Dudani⁶, Chun-Loo Gan⁷, Vidyalakshmi Sethunath¹, Filipe LF. Carvalho⁸, Alma Imamovic¹, Cora Ricker¹, Natalie I. Vokes⁹, Jackson Nyman¹, Jihye Park¹, Michelle S. Hirsch^{3,4}, Rizwan Haq¹, Gwo-Shu Mary Lee¹, Bradley A. McGregor¹, Steven L. Chang⁸, Adam S. Feldman¹⁰, Catherine J. Wu^{1,2,11}, David F. McDermott¹², Daniel Y.C. Heng⁷, Sabina Signoretti^{4,13}, Eliezer M. Van Allen^{1,2,3*}, Toni K. Choueiri^{1,3,11*#}, Srinivas R. Viswanathan^{1,2,3*#}

*Senior authors

#Corresponding authors

Affiliations:

- 1) Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA
- 2) Broad Institute of MIT and Harvard, Cambridge, MA, USA
- 3) Harvard Medical School, Boston, MA, USA
- 4) Department of Pathology, Brigham and Women's Hospital, Boston, MA, USA
- 5) Department of Medicine, Massachusetts General Hospital Cancer Center, Boston, MA, USA
- 6) Division of Medical Oncology/Hematology, William Osler Health System, Brampton, ON, Canada
- 7) Division of Medical Oncology, Tom Baker Cancer Centre, University of Calgary, AB, Canada
- 8) Division of Urology, Brigham and Women's Hospital, Boston, MA, USA
- 9) Department of Thoracic/Head and Neck Medical Oncology; Department of Genomic Medicine, MD Anderson Cancer Center, Houston, TX, USA
- 10) Department of Urology, Massachusetts General Hospital, Boston, MA, USA
- 11) Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA
- 12) Beth Israel Deaconess Medical Center, Boston, MA, USA
- 13) Department of Oncologic Pathology, Dana-Farber Cancer Institute, Boston, MA, USA

Corresponding authors:

Toni K. Choueiri, MD, Department of Medical Oncology, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, Massachusetts, 02215 (toni_choueiri@dfci.harvard.edu). Tel: +1 617-632-5456

Srinivas R. Viswanathan, MD, PhD, Department of Medical Oncology, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, Massachusetts, 02215 (srinivas.viswanathan@dfci.harvard.edu). Tel: +1 617-632-2429

Abstract: 150 words

Main text: 4,439 words

Methods: 4,067 words

Figures: 6

Supplementary figures: 6

Supplementary tables: 6

46 ABSTRACT

47 Translocation renal cell carcinoma (tRCC) is an aggressive and poorly-characterized subtype of
 48 kidney cancer driven by *Mit/TFE* gene fusions. Here, we define the landmarks of tRCC through an
 49 integrative analysis of 152 tRCC patients identified across multiple genomic, clinical trial, and
 50 retrospective cohorts. Most tRCCs harbor few somatic alterations apart from *Mit/TFE* fusions and
 51 homozygous deletions at chromosome 9p21.3 (19.2% of cases). Transcriptionally, tRCCs display a
 52 heightened NRF2-driven antioxidant response that is associated with resistance to many targeted
 53 therapies. Consistently, we find that outcomes for tRCC patients treated with vascular endothelial
 54 growth factor receptor inhibitors (VEGFR-TKI) are worse than those treated with immune checkpoint
 55 inhibition (ICI). Multiparametric immunofluorescence confirmed the presence of CD8⁺ tumor-
 56 infiltrating T cells compatible with a clinical benefit from ICI and revealed an exhaustion
 57 immunophenotype distinct from clear cell RCC. Our findings comprehensively define the clinical and
 58 molecular features of tRCC and may inspire new therapeutic hypotheses.

59

60 KEYWORDS

61 Translocation renal cell carcinoma; genomics; TFE3; TFEB; MITF; NRF2; VEGFR; immune
 62 checkpoint inhibition; immunotherapy; oxidative stress

63

64

65 INTRODUCTION

66 Translocation renal cell carcinoma (tRCC) is an aggressive subtype of non-clear cell kidney cancer that
 67 comprises up to 5% of all RCCs in adults and up to 50% of RCCs in children^{1,2}. Prior case series have
 68 suggested that tRCC has a demographic profile that is distinct from more common subtypes of kidney
 69 cancer, with a younger age at diagnosis, advanced stage at presentation, and a female predominance^{3–}
 70 ⁵. Biologically, tRCCs are driven by activating gene fusions involving transcription factors in the *MIT/TFE*
 71 gene family^{6–12}. There are currently no molecularly-targeted therapies specific to tRCC and effective
 72 treatments for this aggressive cancer remain a major unmet medical need.

73 A significant barrier to the development of mechanism-inspired therapeutics for tRCC is an incomplete
 74 understanding of the molecular landscape and clinical features of the disease. Owing to the rarity of
 75 tRCC, prior genomic profiling studies have been limited in scope. While *MIT/TFE* fusions are universal in
 76 tRCC, it remains unclear whether there are co-occurring genetic alterations or transcriptional programs
 77 that represent additional defining features of the disease^{13–15}. Like the molecular landscape, the clinical
 78 treatment landscape in tRCC is also largely undefined, with no established standard of care. As a result,
 79 tRCC patients are typically treated with therapies originally developed for clear cell RCC (ccRCC)¹⁶,
 80 including vascular endothelial growth factor receptor inhibitors (VEGFR-TKI), multikinase inhibitors
 81 (cabozantinib), mTOR inhibitors, or immune checkpoint inhibitors (ICIs). Although some responses to
 82 each of these classes of agents have been reported in tRCC, outcomes have been variable between
 83 series, and it remains unclear which class(es) of therapeutics are best suited to the biology of tRCC^{17–23}.

84 An intriguing feature of tRCC is that it can exhibit diverse histologic features that may mimic almost all
 85 other subtypes of RCC^{24,25}. As a result, tRCC cases have been retrospectively identified within ccRCC
 86 and papillary RCC (pRCC) sequencing cohorts^{7,26,27}. In this study, we leveraged this “histologic overlap”
 87 between tRCC and other RCC subtypes to identify tRCC cases from across multiple genomic, clinical
 88 trial, and retrospective datasets. We combined these cases with profiling of prospectively identified
 89 patients with tRCC to comprehensively characterize the molecular landscape, clinical features, and
 90 treatment outcomes for this disease.

91 RESULTS

92 Identification of tRCC Cases in Large-scale Clinical and Genomic Datasets

93 To comprehensively characterize both the molecular and clinical features of tRCC, we interrogated RCC
 94 cases across multiple large-scale datasets. In a retrospective analysis of metastatic RCC patients from
 95 the Dana-Farber/Harvard Cancer Center (Harvard cohort), we identified 734 patients with ccRCC, 97
 96 patients with pRCC, 23 patients with chromophobe RCC (chRCC), and 19 patients with tRCC. tRCC
 97 patients were identified on the basis of positive *TFE3* fluorescence *in situ* hybridization (FISH) or strongly
 98 positive TFE3 immunohistochemistry with FISH not available. Among this cohort, we observed that tRCC
 99 patients had significantly worse outcomes than did patients with the other major histologies of RCC (**Fig.**
 100 **1a**), a trend that held in an independent metastatic RCC dataset (International Metastatic RCC Database
 101 Consortium, IMDC; **Fig. S1a**). Similarly, patients with localized tRCC trended towards the shortest
 102 progression-free interval after nephrectomy (**Fig. 1a**). Consistent with smaller case series^{3,5}, we used
 103 data from three large independent cohorts (Harvard, IMDC, TCGA) to confirm that tRCCs are female-
 104 predominant (**Fig S1b**), present at a younger age (**Fig S1c**), higher stage (**Fig S1d**), and are associated
 105 with worse clinical prognostic groups in metastatic disease (**Fig S1e**) as compared with the other major
 106 histologies of RCC. Collectively, these data establish tRCC as a disease that predominantly impacts
 107 young female patients and is more aggressive than other forms of RCC in both the localized and
 108 metastatic settings.

109 To aggregate tRCC cases for genomic analysis, we leveraged the fact that tRCCs have been reported
 110 to share overlapping histologic features with the most frequent histologic subtypes of kidney cancer
 111 (ccRCC and pRCC)²⁸. As a result, a small number of tRCC cases – harboring defining *MiT/TFE* fusions
 112 – have been inadvertently included in several RCC genomic datasets^{26,29–32}. As an example, tRCC cases
 113 with histopathologic features indistinguishable from ccRCC and pRCC were included in the Cancer
 114 Genome Atlas (TCGA) effort^{26,29} (**Fig. 1b** and **Supplementary Table 1**). Building on this observation, we
 115 interrogated fusion calls and/or FISH results for 2818 RCCs across 9 independent datasets profiled by
 116 DNA sequencing (exome, genome, or panel sequencing) and/or RNA sequencing (**Fig. 1c**). We identified

117 a total of 90 tRCCs with genomic (DNA) or transcriptomic (RNA) profiling data (42 with only genomic
118 data, 16 with only transcriptomic data, 32 with both, **Fig. S1f**).

119 **Somatic Mutational and Copy Number Alterations in tRCC**

120 We analyzed the 74 tRCC cases on which DNA profiling data were available to elucidate the genomic
121 landscape of tRCC. Among these cases, 36 were profiled via WES, 3 via WGS, and 35 via panel
122 sequencing (**Methods**). tRCC cases showed few mutations overall, with a median (interquartile range)
123 tumor mutational burden of 0.82 (0.43 - 1.28) per megabase (on WES), a rate significantly lower than
124 ccRCC and pRCC and comparable to chRCC (**Fig. S2a**), with similar trends for all (**Fig. S2b**) and
125 frameshift (**Fig. S2c**) indels. Of the most frequently mutated genes in tRCC, none exceeded a frequency
126 of 10% (**Fig. 2a**). These included genes involved in the DNA Damage response (*ATM* (8.1%), *BRCA2*
127 (8.1%), and *WRN* (4.4%)), genes involved in ATP-dependent chromatin remodeling via the
128 SWItch/Sucrose Non-Fermentable (SWI/SNF) complex (*ARID1A* (5.4%), *SMARCA4* (5.4%)), and
129 mutations in *TERT* (6.8%; primarily non-coding mutations in the *TERT* promoter)³³. Among the 52 cases
130 with gene-level copy number profiling data available, the only recurrent focal alteration in tRCC was
131 homozygous deletion at the *CDKN2A/2B* locus (9p21.3), found in 19.2% of cases. Notably, 50.0% (37/74)
132 of cases in our cohort showed no detectable somatic alterations in either the most frequently mutated
133 tRCC genes or genes that are significantly mutated in clear cell, papillary, or chromophobe RCC (**Fig.**
134 **2a**)²⁷. Analysis of arm-level copy number alterations among 17 tRCC cases in the TCGA cohort³⁴
135 revealed the most frequent alterations to be hemizygous loss of chromosome 3p (28.6%; though
136 markedly less frequent versus ccRCC 86.8%; $p < 0.001$), chromosome 9p (23.5%), chromosome 18
137 (29.4%), and chromosome 22q (18.8%), as well as gain of 17q (20.0%) (**Fig. S2e**). Several of these
138 alterations are defining features of other tumor types of neural/neuroendocrine origin, including
139 monosomy 18 in small intestinal neuroendocrine tumors³⁵, 17q gain in neuroblastoma³⁶, and 22q loss in
140 pediatric ependymoma³⁷.

141 We next conducted an enrichment analysis of driver gene alteration frequencies between tRCC and other
142 RCC subtypes. We computed pairwise enrichment (tRCC versus ccRCC, pRCC, and chRCC separately)

for each locus within each dataset, then used a random-effects meta-analysis to obtain a pooled estimate of gene alteration enrichment or depletion in tRCC versus comparator RCC histologies across datasets (see **Methods**). We found that the genes most frequently altered in tRCC – most notably *CDKN2A/2B* locus (9p21.3) deletions – are highly enriched in tRCC versus other RCC histologies. In contrast, mutations in genes that are significantly mutated in ccRCC, pRCC, and chRCC tended to be depleted in tRCC (**Fig. 2b**). Thus, while tRCCs are genomically quiet overall (with a lower mutational and copy number alteration burden than other RCC histologies), a subset harbor recurrent alterations -- distinct in profile from those seen in other RCCs -- that may cooperate with the *Mit/TFE* fusion to drive cancer.

Structure of *Mit/TFE* fusions in tRCC

We next turned our attention to further analysis of the *Mit/TFE* fusion, the defining genetic lesion in tRCC. Across the combined tRCC cohort, we found that the vast majority of cases (78 cases; 88.6%) harbored *TFE3* fusions, while the remainder harbored *TFEB* (8 cases; 9.1%) or *MITF* (2 cases; 2.3%) fusions (**Fig. 3a**). Seventeen different *Mit/TFE* fusion partners were observed across the cohort and the spectrum of fusion partners was largely distinct between *TFE3*, *TFEB*, and *MITF* (**Fig. S3a**). The most common *TFE3* fusion partners were *ASPSR1*, *SFPQ*, *PRCC*, and *NONO*. Interestingly several chromosomes harbored multiple potential *Mit/TFE* fusion partners (chr1, chr17, chrX) (**Fig. 3b**). *Mit/TFE* fusion partners showed an enrichment for ontology terms involving RNA processing and RNA splicing, and this was driven predominantly by *TFE3* fusion partners (**Fig. 3c** and **Fig. S3b-c**). Analysis of fusion breakpoints revealed that all fusions preserved the C-terminal helix-loop-helix/leucine zipper domain (HLH-LZ) of the *Mit/TFE* transcription factor, the region of the protein critical for dimerization and DNA binding³⁸; the activation domain was variably preserved in the fusion product (**Fig. 3d** and **Supplementary Table 2**). Interestingly, large N-terminal portions of most *TFE3* fusion partners were included in the fusion, including, domains with RNA-binding potential in cases where the fusion partner was an RNA binding protein. In contrast, *TFEB* and *MITF* fusion partners tended to preserve less of the N-terminal fusion partner in the fusion product (**Fig. 3e**). Overall, our results point to a coherent logic to the structure of *Mit/TFE* fusions despite great diversity in fusion partners and breakpoints.

Distinctive transcriptional features of tRCC

Given our observation that most tRCCs harbor few genomic alterations aside from the *MIT/TFE* fusion, we next sought to determine whether the transcriptional program of tRCC is largely driven by the fusion. We ectopically expressed either wild type (WT) *TFE3* or four of the most common *TFE3* fusions (*ASPCR1-TFE3*, *NONO-TFE3*, *PRCC-TFE3*, *SFPQ-TFE3*) in 293T cells and performed RNA-Seq (**Fig. 4a** and **Supplementary Table 3**). We derived a 139-gene transcriptional signature based on genes differentially expressed upon *TFE3* fusion, but not WT *TFE3*, expression (**Fig. S4a**, **Supplementary Table 4** and **Methods**). Subsequently, we performed unsupervised hierarchical clustering using this fusion-specific signature. We observed that tRCC samples clustered tightly together across four independent datasets^{30,39–41} (**Fig. 4b** and **Fig. S4b**). Clustering based on our fusion-derived signature resulted in superior grouping of tRCCs than did clustering based on the 1000 most variable genes in each dataset (**Fig. S4c**). We then performed differential expression analysis to identify a consensus set of genes overexpressed in tRCC as compared with all comparator tumor types. In each dataset, we performed pairwise comparisons between tRCC and each comparator tumor type to identify genes selectively overexpressed in tRCC (q-value <0.05; **Fig. S4d-e**). We identified a consensus list of 76 genes that were selectively overexpressed in tRCC (q-value <0.05) in 9/13 or more pairwise comparisons (**Fig. 4c** and **Fig. S4e**). Notably, several of these have been previously annotated as MITF target genes^{42,43} on the basis of prior ChIP-Seq studies and include genes involved in neuronal development (*SNCB*, *TRIM67*, *IRX6*)^{44–46}, ion flux and the antioxidant stress response (*SQSTM1*, *TMEM64*, *SLC39A1*)^{46–48}, and lysosomal function/mTORC1 signaling (*RAB7A*, *RHEB*, *RRAGC*, *ATP6V1C1*)^{49–51}. We performed gene set enrichment analysis (GSEA)⁵² using hallmark gene sets⁵³ to identify pathways selectively activated in tRCC. This revealed a strong enrichment for gene sets pertaining to reactive oxidative species (ROS) sensing and the response to oxidative stress and xenobiotics (top tRCC-enriched gene sets shown in **Fig. 4d**). In sum, the transcriptional program of tRCC appeared to be driven by the MIT/TFE fusion and resulted in overexpression of genes implicated in mTORC1 signaling, antioxidant stress response, ROS sensing, and the response to oxidative stress and xenobiotics.

195 **An antioxidant response signature associated with resistance to targeted therapies in tRCC**

196 The transcription factor NRF2 (nuclear factor erythroid-derived-2-like 2, *NFE2L2*) is a master regulator of
 197 the cellular antioxidant response and controls the expression of genes involved in the response to
 198 xenobiotics and oxidative stress⁵⁴. Notably, activation of the NRF2 pathway has been reported in certain
 199 subsets of RCC via diverse mechanisms that include somatic alteration or hypermethylation of NRF2
 200 pathway members⁵⁵ and the production of oncometabolites that modify and inhibit KEAP1, a negative
 201 regulator of NRF2^{7,10,27}. Given evidence of activated ROS-sensing in tRCC (**Fig. 4c-d**), we derived an
 202 NRF2 activity score using single sample GSEA (ssGSEA)⁵⁶ (based on a 55-gene NRF2 signature⁵⁷)
 203 across all RCC samples with available transcriptome profiling data (46 total tRCC samples across 4
 204 datasets; NRF2 activity calculated and Z-scored separately within each individual dataset). We observed
 205 that NRF2 activity was universally high amongst tRCC samples as compared with other RCC types and
 206 normal kidney tissue (**Fig. 5a**).

207 We next investigated whether high NRF2 activity in tRCC was attributable to somatic alterations in this
 208 pathway. We observed that somatic alterations in the NRF2 pathway (most commonly *KEAP1* or *NFE2L2*
 209 alteration) were associated with an increased NRF2 activity score in ccRCC and pRCC, as was a CpG
 210 island methylator phenotype (CIMP), consistent with prior reports (**Fig. 5b** and **Fig. S5a**)²⁷. Interestingly,
 211 however, tRCC samples showed uniformly elevated NRF2 activity, comparable to ccRCC/pRCC samples
 212 with somatic alterations in the NRF2 pathway (**Fig. 5b**), despite having no detectable NRF2 pathway
 213 alterations. The expression of strong oncogenes has been linked to NRF2 pathway activation⁵⁸ and our
 214 transcriptomic analyses revealed overlapping targets between NRF2 and MITF (**Fig. 4c**, hypergeometric
 215 one-tailed p-value < 0.001). Consistently, we observed that the NRF2 gene signature was enriched upon
 216 ectopic expression of all *TFE3* fusions in 293T cells as compared to the mock treatment condition,
 217 suggesting that expression of the *TFE3* fusion may be directly linked to activation of the NRF2 pathway
 218 (**Fig. 5c**).

219 Activation of the NRF2 pathway has been associated with resistance to a number of ROS-producing
 220 drugs, including inducers of ferroptosis, a regulated form of iron-dependent oxidative cell death^{57,59,60}. We

calculated a correlation between NRF2 activity score and drug sensitivity across 593 cell lines and 481 compounds assayed in the Cancer Therapeutics Response Portal⁶¹. Strikingly, high NRF2 activity was associated with relative resistance to almost all agents assayed, including several targeted therapies used in the treatment of RCC (e.g. sunitinib, axitinib, lenvatinib, temsirolimus), and most notably, to multiple compounds known to induce electrophilic stress and oxidative cell death (e.g. PRIMA-1, PX-12, piperlongumine, ML-210, RSL-3) (**Fig. 5d**)⁶². In order to uncover potential vulnerabilities of this otherwise drug-resistant state, we next surveyed pooled genetic (shRNA and CRISPR) screening data generated as part of the Cancer Dependency Map effort^{63,64}. In both the CRISPR and shRNA datasets, we found that the outlier dependency of NRF2-high cells is *NFE2L2* itself (**Fig. S5b**). Although tRCC cell lines are not currently included among those assayed in the Cancer Dependency Map effort, we separately validated that three tRCC cell lines all demonstrated variable levels of dependency on *NFE2L2* knockdown, consistent with the notion that direct inhibition of NRF2 is a vulnerability of the NRF2-high state observed in tRCC (**Fig. S5c**).

Next, to determine whether elevated NRF2 activity might be associated with resistance to targeted therapies in patients, we evaluated molecular data from the IMmotion151 trial (NCT02420821), a Phase III trial of 915 RCC patients with clear cell or sarcomatoid histology who were randomized to either sunitinib (multitargeted kinase inhibitor against VEGFRs and PDGFRs) or the combination of atezolizumab (monoclonal antibody targeting PD-L1) and bevacizumab (monoclonal antibody targeted VEGF-A)⁶⁵. RNA-Seq performed on tumor biopsies from patients enrolled on this trial revealed 15 patients with *TFEB/TFE3* translocations among 822 with available RNA-seq data (**Fig. 1c**), of which 6 were treated on the sunitinib arm and 9 were treated on the atezolizumab + bevacizumab (AtezoBev) arm³⁰. While AtezoBev showed a modest benefit over sunitinib in progression-free survival (PFS) in the overall study and amongst ccRCC patients, we observed that tRCC patients receiving sunitinib did dramatically worse than those receiving AtezoBev (median PFS 3.5 months with sunitinib vs. 15.8 months with AtezoBev; log-rank $p=0.004$). Consistent with this observation, the extent of benefit derived from AtezoBev as compared with sunitinib, in patients with tRCC vs. ccRCC, was significantly greater

(histology-by-treatment arm interaction Cox p-value=0.008) (**Fig. 5e**). When ccRCC patients treated with sunitinib were dichotomized based on NRF2 activity score, those with high-NRF2 scores had shorter PFS compared to with low-NRF2 scores (median PFS 7.1 months for high-NRF2 vs. 11.1 months for low-NRF2; log-rank p=0.002). In contrast, NRF2 activity score was not associated with a significant difference in outcome in ccRCC patients treated on the AtezoBev arm (**Fig. 5f**). In the CheckMate cohort including 311 patients with ccRCC with available RNA-seq data (pooled analysis of the CheckMate 009 [NCT01358721], 010 [NCT01354431], and 025 [NCT01668784] clinical trials)⁶⁶, a similar signal was observed whereby ccRCC patients with a high NRF2 activity score experienced shorter PFS than did those with a low NRF2 activity score (**Fig. S5d**), on the everolimus arm (median PFS 9.7 months for high-NRF2 vs. 14.3 months for low-NRF2; log-rank p= 0.031), but not the nivolumab arm⁶⁷. Together, these results indicate that high NRF2 activity – a defining feature of tRCC – is associated with resistance to targeted agents used in the treatment of RCC, but may not preclude responses to ICI.

Response to immune checkpoint inhibition in tRCC

We sought to further explore the possibility that tRCC may be responsive to ICI. Analysis of responses from the IMmotion151 study showed that tRCC patients derived significantly greater clinical benefit (CB) on AtezoBev than on sunitinib (77.8% with AtezoBev vs. 16.7% with sunitinib; Fisher p-value= 0.041). However, tRCC patients tended to not derive clinical benefit (no clinical benefit; NCB) from sunitinib as compared with AtezoBev (11.1% with AtezoBev vs. 50.0% with sunitinib; Fisher p-value= 0.235). In contrast, ccRCC patients tended to have similar CB (65.1% with AtezoBev vs. 64.0% with sunitinib; Fisher p-value= 0.767) and NCB (15.6% with AtezoBev vs. 16.0% with sunitinib; Fisher p-value= 0.923) rates whether they received AtezoBev or sunitinib (**Fig. 6a**).

In a combined analysis of the IMDC and Harvard datasets, we identified 12 metastatic tRCC patients who had received ICI in any line of therapy as well as 10 tRCCs that had been treated by TKIs (n= 8 sunitinib; n= 2 pazopanib). Among this cohort, 5 achieved either partial response (n= 3) or stable disease (n= 2) on an ICI-containing regimen, with several ongoing responses (**Fig. 6b** and **Fig. S6a-b**). Overall, in this retrospective combined cohort of tRCC patients, the response rate (25.0% with ICI and 0% with

TKI; Fisher p-value= 0.220) and overall survival (OS; median OS 62.4 months with ICI and median OS 10.3 months with TKI; log-rank p-value= 0.267) tended to be increased on ICI-based regimens compared to TKIs (**Fig. S6c-d**), corroborating the result that tRCC patients may derive greater benefit from ICI-based therapies than VEGF-targeted therapies.

We next examined whether immunogenomic features of tRCC could explain responses to ICI in this RCC subtype, despite a low burden of mutations and CNAs (**Fig. 2a** and **Fig. S2**). In the TCGA cohort, tumor purity (which is inversely correlated to immune cell infiltration), was lower in tRCC than chRCC (a classically ICI-resistant subtype^{68,69}) (Wilcoxon p-value< 0.001), similar to pRCC (Wilcoxon p-value= 0.160), and higher than ccRCC (Wilcoxon p-value= 0.005) (**Fig. 6c**). Consistently, immune deconvolution analyses (CIBERSORTx⁷⁰) showed that the inferred percentage of cluster of differentiation 8 (CD8)⁺ T cells was higher in tRCC than in chRCC (Wilcoxon p-value< 0.001), and comparable to that seen in ccRCC (Wilcoxon p-value= 0.190) and pRCC (Wilcoxon p-value= 0.150) (**Fig. 6d**). Additionally, PD-L1 protein levels on tumor-infiltrating immune cells, as assessed by IHC, in patients on the IMmotion151 trial, were comparable between tRCC and ccRCC patients (41.8% with ccRCC vs. 33.3% with tRCC; Fisher p-value= 0.604) (**Fig. S6e**).

Finally, we sought to more carefully characterize the CD8⁺ tumor-infiltrating T cells in tRCC via multiparametric immunofluorescence^{71,72}. We examined 11 ccRCC cases (including 10 with adjacent normal tissue) and 11 tRCC cases for T cells expressing CD8 or the immune checkpoint markers PD1, T-cell immunoglobulin and mucin-domain containing-3 (TIM3), and lymphocyte activation protein-3 (LAG3). While the overall CD8⁺ T cell density tended to be lower in tRCC samples than in ccRCC samples (Wilcoxon p-value = 0.065) (**Fig. 6e-f**), the percentage of CD8⁺PD1⁺TIM3⁻LAG3⁻ cells (the subset predictive of a response to PD1/PD-L1-based ICI^{71,72}) was not significantly different between tRCC and ccRCC (**Fig. S6f**). Moreover, the profile of immune checkpoint markers differed significantly between ccRCC and tRCC; tRCC cases displayed a higher percentage of CD8⁺PD1⁺TIM3⁻LAG3⁺ T cells (Wilcoxon p-value = 0.009) whereas ccRCC cases displayed a higher percentage of CD8⁺PD1⁺TIM3⁺LAG3⁻ T cells (Wilcoxon p-value = 0.040). Altogether, our results are consistent with the notion that tRCCs may benefit

from ICI as a result of a permissive immune microenvironment characterized by a tumor-infiltrating T cell profile distinct from that observed in ccRCC.

301

302 DISCUSSION

303 We performed a comprehensive and multicenter characterization of the molecular and clinical features
304 of 152 tRCCs. While prior studies have identified some genomic and transcriptional features of tRCC,
305 the broader extensibility of these findings, their clinical actionability, as well as an understanding of how
306 they compare to other subtypes of RCC have remained unclear^{13–15}. Our integrative analysis spans
307 genomic and transcriptomic data, immunophenotypic analysis, functional validation, and clinical outcome
308 data from both retrospective cohorts and randomized clinical trials. From these efforts, an increasingly
309 well-defined landscape of tRCC emerges.

310 The defining – and often singular – genomic alteration in tRCC is the *MiT/TFE* fusion. Our results show
311 that *TFE3* is by far the most frequently involved *MiT/TFE* gene. While there exists a great diversity of
312 *MiT/TFE* fusion partners, these partners are highly enriched on certain chromosomes (chr1, chr17, chrX),
313 raising intriguing questions about whether patterns of spatial genome organization underlie these
314 recurrent translocations^{73–75}. Moreover, our analysis of breakpoint locations across fusions highlights that
315 the vast majority of *TFE3* fusions arise via in-frame events that preserve functional domains from both
316 *TFE3* and its partner protein (most of which are RNA binding proteins); this opens the possibility that
317 *TFE3* fusion partners may confer neomorphic activity to the fusion product. In contrast, much smaller
318 regions of *TFEB* and *MiTF* partner genes appear to be involved in the fusion product. Whether differences
319 in fusion structure translate to histologic and/or phenotypic differences between *TFE3*-, *TFEB*-, and *MiTF*-
320 translocation RCC warrants further investigation^{1,76,77}.

321 Overall, tRCCs are genomically quiet tumors with a low mutational and copy number alteration burden,
322 a reduced frequency of alterations in genes known to be significantly mutated in other RCC subtypes,
323 and few recurrent alterations aside from the *MiT/TFE* fusion. A notable exception is homozygous loss at

chromosome 9p21.3, which harbors the *CDKN2A/2B* genes, and is found in 19.2% of tRCC cases. Loss of CDKN2 proteins may be associated with high CDK4/6 activity and may sensitize to CDK4/6 inhibitors⁷⁸. Co-deletion of *MTAP*, which is located in close proximity to *CDKN2A*, may sensitize to PRMT5 inhibitors^{79,80}. Mutations in *TERT* (primarily in the promoter region) were also found in 6.8% of cases. Notably, both *CDKN2A/B* loss and *TERT* promoter mutations are defining genetic features of malignant melanoma, a cancer type driven by activated MITF signaling^{33,81–83}. Less frequent alterations in the cohort included multiple genes involved in the DNA damage response (*ATM*, *BRCA2*, *WRN*), though the lack of specific variant information, the absence of matched normal-based filtering of mutation calls for some samples, and low alteration frequency preclude drawing strong conclusions about this class of mutations.

We identified a heightened response to oxidative stress as a transcriptional hallmark of tRCC. Activated NRF2 signaling has been linked to oncogenesis and resistance to chemotherapies in various contexts⁸⁴. Prior studies have indicated that small subsets of both ccRCC and pRCC display heightened NRF2 signaling, generally linked to somatic alterations or DNA methylation in the NRF2 pathway^{7,27,85}. Interestingly, our results suggest that NRF2 signaling is uniformly activated in tRCC in the absence of detectable somatic alterations in the NRF2 pathway. Notably, multiple NRF2 target genes are also annotated as MIT/TFE targets (**Fig. 4c**), suggesting a direct link between MIT/TFE fusions and the NRF2 pathway in tRCC. Our results may explain why tRCCs (and ccRCCs with elevated NRF2 signaling) display worse outcomes with sunitinib than with ICI in clinical datasets, and are consistent with *in vitro* data suggesting that NRF2 confers resistance to sunitinib and other TKIs^{55,86,87}. Whether this signal holds for extended spectrum kinase inhibitors such as cabozantinib and lenvatinib remains to be determined, as patients receiving these therapies were not represented in our retrospective cohort. We validate that *NFE2L2* represents a clear genetic dependency of the NRF2-high state, and suggest that specific NRF2 pathway inhibitors, if developed, may be effective in tRCC^{54,88}.

Responses to ICI in tRCC are notable given the apparent lack of potential sources of tumor-associated antigens (i.e. low burden of mutations and indels). Our immune deconvolution analyses and immunofluorescence studies both support the notion that tRCCs do contain an appreciable density of

tumor-infiltrating CD8⁺ T cells. The tumor neoantigens recruiting T cells in tRCC may be derived from the fusion junction, as has also been reported for other fusion-driven malignancies^{15,89}. Interestingly, there is no significant difference in the percentage of CD8⁺PD1⁺TIM3⁺LAG3⁺ T cells – the activated non-exhausted T-cell subset that is implicated in an effective antitumor response – between ccRCC (a classically ICI-responsive tumor) and tRCC^{90–92}. The immunophenotype of exhausted T cells also appears to differ between ccRCC and tRCC: CD8⁺PD1⁺TIM3⁺LAG3⁺ T cells are predominant in tRCC while CD8⁺PD1⁺TIM3⁺LAG3⁺ T cells are predominant in ccRCC. Both TIM3 and LAG3 have been proposed as immune checkpoints that can be targeted in combination with PD-1/PD-L1. Notably, several trials combining LAG3 blockade with PD1 blockade are currently underway (and include patients with RCC)⁹⁰ and this combination has recently shown to have efficacy in patients with previously untreated metastatic melanoma⁹³. Our immunophenotypic data provide rationale for the development of this therapeutic combination in tRCC. Our findings are also consistent with those of a prior study that showed, using a lung adenocarcinoma mouse model, that activated NRF2 and PI3K/mTOR signaling can lead to changes in the immune microenvironment that are permissive to ICI response⁹⁴. In tRCC, our results suggest that both the PI3K/AKT/mTOR pathway and NRF2 may be activated downstream of MiT/TFE fusions (**Fig. 4c**)²¹.

Our study does have several limitations. First, the cohort is heterogeneous in terms of stage of disease (localized and metastatic), sequencing platform used, and data types available for analysis. While the heterogeneity of the cohort is inevitable given the rarity of the disease, the analysis methods we apply account for dataset-specific biases (**Methods**) and the scale of this study has enabled us to make multiple novel insights. Second, tRCCs are themselves a heterogeneous group of tumors with respect to fusion partners, biology, and prognosis⁹⁵. Larger studies or more homogeneous cohorts comprised of prospectively collected samples will be required to draw strong conclusions about how the specific *MiT/TFE* gene or its fusion partner influence disease biology. Third, some of our clinical data are retrospective, which has inherent limitations. Nonetheless, we suggest that the signals observed from misclassified tRCC patients enrolled on randomized clinical trials for ccRCC, and the corroboration of

376 these signals by translational and retrospective clinical data, may have important implications for the
377 treatment of tRCC.

378 Altogether, we demonstrate the power of integrative clinico-genomic analysis to illuminate the molecular
379 underpinnings and clinical features of tRCC. Our work inspires multiple hypotheses that can be pursued
380 in future studies to further dissect the biology of this rare cancer. These data also lay the framework for
381 the development and testing of mechanism-driven therapeutic regimens in tRCC.

382

383 METHODS

384 Clinical tRCC cohorts

385 The comparison of baseline characteristics and clinical outcomes was done using data from patients
 386 included in two retrospective cohorts of consecutive patients: (1) Harvard cohort (n= 734 ccRCC, n= 97
 387 pRCC, n= 23 chRCC, n= 19 tRCC), a retrospective cohort from the Dana-Farber/Harvard Cancer
 388 Center including patients from Dana-Farber Cancer Institute, Beth Israel Deaconess Medical Center,
 389 and Massachusetts General Hospital and (2) IMDC cohort (n= 6107 ccRCC, n= 396 pRCC, n= 107
 390 chRCC, n= 40 tRCC): a retrospective multi-center cohort of metastatic RCC that includes more than 40
 391 international cancer centers and more than 10,000 patients with metastatic RCC⁹⁶. All patients
 392 consented to an institutional review board (IRB) approved protocol to have their clinical data
 393 retrospectively collected for research purposes and the analysis was performed under a secondary use
 394 protocol, approved by the Dana-Farber Cancer Institute IRB. For the Harvard cohort, tRCC patients
 395 were defined as: (1) positive *TFE3* FISH test or (2) positive *TFE3* test by IHC along with a strongly
 396 suggestive clinico-pathologic history and no FISH testing results available (missing). For the IMDC
 397 cohort, patients were included as tRCCs if they (1) had a positive *TFE3* FISH test, (2) had a positive
 398 *TFE3* IHC test and suggestive clinico-pathologic history and no FISH testing data available (missing),
 399 or (3) no *TFE3* FISH or *TFE3* IHC test results available but suggestive clinico-pathologic history.
 400 Clinico-pathologic diagnoses were used to define comparator RCC histologies (ccRCC, pRCC, and
 401 chRCC). For the IMDC cohort, comparator histologies (controls) were only used from clinical sites that
 402 contributed tRCC cases.

403 Genomic tRCC cohorts

404 For genomic datasets, tRCCs were identified based on RNA-seq-based fusion calls, a positive *TFE3*
 405 FISH test, or DNA-based fusion calls derived from panel data (MSK-IMPACT or OncoPanel). Clinico-
 406 pathologic diagnoses were used to define the cases of other RCC histologies (ccRCC, pRCC, chRCC,
 407 normal kidney, or other). Data for the Memorial-Sloan Kettering (MSK) cohort was obtained from the

study by Marcon et al.¹⁵ and Zehir et al.⁹⁷. Fusion calls for the TCGA cohort were obtained from the study by Gao et al.²⁹, clinico-pathologic data was obtained from Genomic Data Commons (<https://gdc.cancer.gov/about-data/publications/pancanatlas>), and the pathology slides used in **Fig. 1b** were obtained from <https://portal.gdc.cancer.gov/>. Data for the PCAWG⁹⁸ cohort were obtained from the ICGC data portal (<https://dcc.icgc.org/releases/PCAWG>). Data for the IMmotion151 (NCT02420821, Motzer et al.)³⁰, Wang et al.⁴⁰, Durinck et al.³², Malouf et al.⁹⁹, and Sato et al.³¹ cohorts were obtained from the corresponding studies. For the OncoPanel cohort, DNA extraction, sequencing, and mutation and copy number calling were performed as previously described for the OncoPanel gene panel assay¹⁰⁰. The OncoPanel assay is an institutional analytic platform that is certified for clinical use and patient reporting under the Clinical Laboratory Improvement Amendments (CLIA) Act. The panel includes 275 to 447 cancer genes (versions 1 to 3 of the panel). Sample-level data for the OncoPanel cohort (mutations, gene-level CNA, and clinical metadata) are provided in **Supplementary Table 5**. The data types available for each dataset are illustrated in **Fig. 1C**, but not all data types were available for all samples in each cohort. The full list of samples used (including the data types available) and sequencing platform used for DNA-sequencing (WGS, WES, or panel) are provided in **Supplementary Table 1**.

Analysis of mutation and copy number variants in genomic tRCC cohorts

Mutation calls (all aligned to human genome reference build hg19) were obtained as detailed above. Specifically, for the MSK cohort^{15,97}, WES-based calls were used where available and panel-based data were otherwise used for tRCC samples. For the TCGA cohort, the mc3 MAF calls¹⁰¹ (<https://gdc.cancer.gov/about-data/publications/pancanatlas>) were used. For the Durinck et al.³² and Malouf et al. cohort⁹⁹, only samples from patients that had mutation calling based on matched normal sequencing were included. For the Sato et al. cohort³¹, only the WES calls were used. All mutations were annotated uniformly using Oncotator¹⁰² (except for the IMmotion151 cohort, for which a MAF was not available). In order to filter out potential germline mutations in the OncoPanel cohort, mutations present at an allelic frequency of 0.5% in one of the superpopulations from the 1000 Genomes

Project¹⁰³ (<https://www.internationalgenome.org/data>) were excluded from all downstream analyses. For the enrichment analyses, mutations were included if they were truncating (nonsense or splice site), insertions-deletions (indels), missense mutations, or *TERT* promoter mutations. For the IMmotion151 cohort, mutations were included if they were short-variants or truncating. The mutation load was computed as the number of all non-synonymous mutations per sample. The indel load was computed as the number of all indels per sample (either all indels or only frameshift indels). For the OncoPanel and MSK-IMPACT samples, the mutation and indel loads were normalized to the bait sets of the version of the panel used. The bait sets¹⁰⁴ for OncoPanel were: v1, 0.753334 Megabases [Mb]; v2, 0.826167 Mb; and v3, 1.315078 Mb. For MSK-IMPACT, the bait sets were: IMPACT341, 0.896665; IMPACT410, 1.016478; and IMPACT468, 1.139322 Mb.

Gene-level copy number data calls were available for the MSK cohort⁹⁷, IMmotion151 cohort³⁰, OncoPanel cohort (**Supplementary Table 5b**), PCAWG (https://dcc.icgc.org/releases/PCAWG/consensus_cnv/GISTIC_analysis/all_thresholded.by_genes.rmcnv.pt_170207.txt), and TCGA (<http://firebrowse.org/>; KIPAN dataset). For all gene-level analyses only focal events (deep deletions and high amplifications) were considered. As measures of the copy number alteration burden, the aneuploidy score and fraction genome altered were obtained for the TCGA¹⁰⁵ and MSK⁹⁷ cohorts, respectively. Arm-level calls were obtained for the TCGA cohort¹⁰⁵.

Genomic enrichment analyses

In order to account for the inherent differences between the included cohorts and to maximize the power of the study to detect differences in mutations and copy number alterations in tRCC versus other RCC histologies, a meta-analytic approach was adopted for all gene-level enrichment analyses, as has been done in prior studies^{106,107}. First, Fisher's exact tests were used to evaluate the enrichment of gene alterations (mutations and copy number alterations separately) within each cohort (combined WES cohort, IMmotion151, PCAWG, OncoPanel, and MSK-IMPACT). For panel-based cohorts, this enrichment took into account the bait set of each version of the panel used for sequencing (i.e. a gene

was counted as missing, and not non-mutated, if not included in the bait set of a version of the panel). The conditional maximal likelihood estimate of the odds ratio and its 95% confidence interval were computed using the `fisher.test()` function from the stats package in R. For each gene, we then obtained pooled estimates of the odds ratio and its 95% confidence interval using a random-effects model with the Paule-Mandel estimator for tau, with treatment arm continuity correction and Knapp-Hartung adjustment. The meta-analysis was performed using the `metabin()` function from the meta package in R^{108–110}. The enrichment analysis was performed pairwise between tRCC and each comparator RCC histology separately (ccRCC, pRCC, and chRCC). Genes were included in the enrichment analysis if: (1) they were altered in at least two different cohorts; (2) alteration frequency in tRCC was 3% or more; and (3) were Tier 1 cancer genes as defined in the Cancer Gene Census (accessed on February 17 2021)¹¹¹. Genes that had been previously reported to be significantly mutated in ccRCC, pRCC, and chRCC²⁷ were also included in the analysis. For all analyses, samples that were originally part of the TCGA and PCAWG cohorts were only included in one of the two cohorts as part of the enrichment analyses (cohort assignment reported in **Supplementary Table 1**). The CoMut plot was generated using the CoMut package in Python¹¹² and genes that were not assessed in specific samples (i.e. not included in the bait sets of the gene panel used) are shown as gray boxes; the corresponding alteration frequency (bar graph at the right-hand side of the CoMut) was adjusted accordingly and reflects only samples in which a particular gene was assessable for alteration. Arm-level comparisons (TCGA cohort) were performed pairwise with RCC histologies using Fisher’s exact tests. The mutation and indel loads, as well as the aneuploidy score and fraction genome altered, were compared pairwise with each RCC histology (ccRCC, pRCC, and chRCC) using Wilcoxon rank-sum tests.

***MiT/TFE* fusion identification and characterization**

Fusion calls were obtained as detailed under “Genomic tRCC cohorts” above. In particular, for the MSK cohort, determination of fusion partners was based on MSK-IMPACT and/or RNA-seq^{15,97} and fusion breakpoints were based on MSK-IMPACT and available for a subset of samples⁹⁷. For the OncoPanel cohort, fusion partners and breakpoints were based on an in-house fusion calling pipeline and were

available for a subset of samples. For the TCGA, PCAWG, Wang et al., Sato et al., Durinck et al., and Malouf et al. cohorts, fusion partners were based on RNA-seq. Of those, the fusion breakpoints were available for the TCGA, PCAWG, Sato et al., and Durinck et al. cohorts. For the Malouf et al. cohort, fusion breakpoint locations were inferred based on the reported fusion breakpoint sequences using BLAT (<https://genome.ucsc.edu/cgi-bin/hgBlat>). All breakpoint locations were aligned to human genome reference build hg19, except for the TCGA breakpoints which had been originally mapped to hg38 and were converted to hg19, for the purposes of this analysis, using liftOver (<https://genome.ucsc.edu/cgi-bin/hgLiftOver>). The Circos Perl package¹¹³ was used to represent the chromosomal locations of fusions in a circos plot. The enrichr¹¹⁴ tool was used to evaluate enrichment of Gene Ontology (GO) terms among the *Mit/TFE* partner genes. In order to annotate the fusion protein products based on the breakpoints, breakpoints were first aligned to human genome GRCH37.p13 on NCBI Genome Data viewer. Functional domains were then annotated using UniPort Protein knowledgebase UniProtKB/Swiss-Prot And NCBI Conserved Domain Database¹¹⁵ (CDD v3.19). The presence of Prion-Like domains (PLD) was analyzed using Prion-Like Amino Acid Composition (PLAAC) web-based program¹¹⁶. Illustrations were made using Illustrator for Biological Sequences (IBS)¹¹⁷ version 1.0. Annotated functional domains with abbreviations are provided in **Supplementary table 2**.

Cell lines

293T cells were obtained from the American Type Culture Collection. UOK109 and UOK146 cells were a kind gift of Dr. Marston Linehan (National Cancer Institute). FU-UR-1 cells were a kind gift of Dr. Masako Ishiguro (Fukuoka University School of Medicine). Cell lines were grown in base media of DMEM (293T, UOK109, UOK146) or DMEM/F12 (FU-UR-1), supplemented with 10% FBS, 100 U mL⁻¹ penicillin, 100 µg mL⁻¹ streptomycin, 2 mM L-glutamine, and 100 µg mL⁻¹ Normocin (Invivogen).

TFE3 fusion-specific signature

For *TFE3* fusion overexpression experiments, 293T cells were seeded in 6-well plates at 2 x 10⁵ cells per well and after 24 hours were transfected with 500 ng of plasmids encoding *ASPSCR1-TFE3*, *NONO-*

511 *TFE3*, *PRCC-TFE3*, *SFPQ-TFE3*, wild type (WT) *TFE3*, or an empty vector control (all in pLX313). All
 512 transfections were performed in three biological replicates. Cells were harvested 48 hours after
 513 transfection and total RNA was collected using the RNeasy Plus Mini Kit (QIAGEN, #74136). Sample
 514 concentrations were measured using a NanoDrop 8000 Spectrophotometer (Thermo Fisher Scientific)
 515 and sequencing libraries were prepared with poly(A) selection. Libraries were pooled and paired-end 150
 516 bp RNA-sequencing was performed on an Illumina HiSeq. Paired-end sequencing reads were aligned to
 517 the human genome reference build hg38 using STAR v2.7.2¹¹⁸ and quantified using RSEM v1.3.2¹¹⁹.
 518 Transcripts were filtered based on read support (sum of expected read counts across three biological
 519 replicates > 30) prior to gene-level differential expression analysis using the voom transformation in limma
 520 v3.40.6¹²⁰. Transcripts-per-million (TPMs) were used for visualization and clustering. Expected count and
 521 TPM matrices are provided in **Supplementary Table 3**.

522 In order to derive a transcriptional signature that is specific to the *TFE3* fusion, we performed differential
 523 gene expression of each of the fusion conditions (*ASPCR1-TFE3*, *NONO-TFE3*, *PRCC-TFE3*, *SFPQ-*
 524 *TFE3*) versus the WT *TFE3* condition. Genes that were significantly upregulated ($q < 0.05$ and $\log_2(\text{fold-}$
 525 $\text{change}) > 0$) or significantly downregulated ($q < 0.05$ and $\log_2(\text{fold-change}) > 0$) across all four comparisons
 526 defined a *TFE3* fusion-specific signature (**Supplementary Table 4**). In order to evaluate the relevance
 527 of the *in vitro*-derived signature to tRCC tumor samples, we performed clustering on 4 independent RNA-
 528 seq datasets that included tRCC samples. The normalized expression matrices used for clustering were
 529 those obtained from TCGA (<https://gdc.cancer.gov/about-data/publications/pancanatlas>), PCAWG
 530 (<https://dcc.icgc.org/releases/PCAWG>), IMmotion151, and Wang et al. as described under “Genomic
 531 tRCC cohorts” above. Clustering was performed in each dataset independently using the Heatmap
 532 function from the ComplexHeatmap¹²¹ package in R, using hierarchical clustering with ward.D2 as the
 533 clustering method and the Kendall correlation distance metric. The average intra-tRCC distance was
 534 used as a metric for density of clustering of tRCCs and was compared to the distance obtained from
 535 clustering using the 1000 most variable genes within each dataset (**Fig. S4**).

536 Differential gene expression analysis

Pairwise differential gene expression analysis was performed between tRCC and each other sample type, within each dataset independently (TCGA, PCAWG, IMmotion151, Wang et al., and 293T cell line experiment). Differential gene analysis for the cell line experiment was performed as described above using the limma package. For the tumor datasets, differential gene expression was performed using pairwise Wilcoxon rank-sum tests. For all tests, the Benjamini-Hochberg correction was used to compute q-values and a q-value<0.05 was taken as statistically significant. In order to define a transcriptional signature specific to tRCC, an Upset plot was computed using the UpsetR package¹²². The 76 genes that were found to be significantly upregulated in 9 or more of the 14 pairwise comparisons were plotted in a heatmap (**Fig. 4c**), which included tRCC samples and comparator samples. Gene pathway annotations were obtained from enrichr¹¹⁴. Overlap between the NRF2 and MITF target genes was evaluated using a one-tailed hypergeometric test. In order to adjust for potential RNA-seq batch effects between datasets in visualization, gene expression was Z-scored within each dataset independently. For volcano plots, log₂(fold-change) of the mean expression of genes in each group was used.

Gene set enrichment analysis

Pre-ranked gene set enrichment analysis (GSEA) was performed pairwise between tRCC and each comparator, within each dataset independently, using the using $-\log_{10}(\text{q-value})$ signed by the sign of the log₂(fold-change) of mean gene expression. GSEA was used on the Hallmark gene sets v7.1 from the Molecular Signatures Database (MSigDB)¹²³ and a previously defined 55-gene NRF2 signature¹²⁴. For the Hallmark analysis, the gene sets were ranked by the number of pairwise comparisons that had a normalized enrichment score (NES)>1 in tRCC vs the other comparators (with the top gene sets visualized as a dot plot) (**Fig. 4d**).

In addition, single sample GSEA (ssGSEA) scores were computed for the 55-gene NRF2 signature using the GSVA package⁵⁶ in R to infer the level of activity of NRF2 in each sample. In order to adjust for potential RNA-seq batch effects in visualization, NRF2 signature scores were Z-scored within dataset prior to visualization as a waterfall plot (**Fig. 5a**). Comparison of ssGSEA scores between tumor

types in the TCGA cohort was performed using Wilcoxon rank-sum tests. To examine the relationship of the NRF2 signature with survival outcomes, the NRF2 score was dichotomized at the median in each treatment arm of each cohort.

Analysis of CTRP and DepMap datasets

RNAi genetic dependence scores were obtained from the DEMETER2 Data v6 dataset¹²⁵, CRISPR genetic dependence scores were obtained from the CRISPR (Avana) Public 21Q1 dataset^{126,127} and drug area under the curve (AUC) values were obtained from the CTRP v2.0 2015 CTD² dataset^{61,128}. Cell lines were excluded if they had multiple AUC values for each drug. All datasets were downloaded from the DepMap Data Download Portal (<https://depmap.org/portal/download/>). NRF2 ssGSEA scores were calculated from the Broad Institute CCLE RNA-seq dataset. Expression values were upper quartile normalized prior to analysis. For each drug (or gene), drug AUCs (or gene dependence scores) were Z-scored and the NRF2 ssGSEA scores were Z-scored, amongst samples having data for both data types. Pearson's correlation coefficient was used to assess the association between drug AUC Z-score and NRF2 ssGSEA Z-score as well as between gene dependence Z-score and NRF2 ssGSEA Z-score. For each correlation, t-statistics were computed ($t = r \cdot ((n-2)/(1-r^2))^{0.5}$), a two-tailed Student's t-distribution was used to determine p-values, and q-values were computed using a Benjamini-Hochberg correction.

Colony forming assays

shRNAs were cloned into a doxycycline-inducible lentiviral vector as previously described¹²⁹. The indicated cell lines were transduced with lentivirus expressing doxycycline-inducible shRNA (shRNA target sequence: CCGGCATTTCACTAAACACAA) and selected with 500 µg/mL of G418 prior to seeding at equal densities with or without the addition of 1 µg/mL doxycycline. Cell densities ranged from 500-1500 cells per well of 12-well plate depending on the cell line. Fresh complete culture media with/without doxycycline was replaced every two days prior to fixation and staining with crystal violet after 12-20 days. Colony areas were quantified using Image J v1.53.

588 Multiplex immunofluorescence and image analysis

589 Cluster of differentiation (CD8), programmed death 1 (PD1), T cell immunoglobulin and mucin domain-3
590 (TIM3), and Lymphocyte-activation gene 3 (LAG3) multiplex immunofluorescence (IF) was performed as
591 previously described⁶⁶. Briefly, we used the Perkin Elmer Opal tyramide signal system on a Bond RX
592 Autostainer (Leica Biosystems). The anti-CD8 antibody (1:5,000, C8/144B, mouse monoclonal antibody,
593 Agilent) was detected using the Opal 520 fluorophore (1:150, FITC); the anti-TIM3 antibody (1:1,000,
594 AF2365 goat monoclonal antibody, R&D Systems) was detected using the Opal 540 fluorophore (1:50,
595 Cy3); the anti-LAG3 antibody (1/10,000, 17B4 mouse monoclonal antibody, LifeSpan Biosciences) was
596 detected using the Opal 560 fluorophore (1:150, Texas Red); the validated anti-PD-1 antibody (1:5,000,
597 EH33 mouse monoclonal antibody, Dr. Freeman laboratory, Dana-Farber Cancer Institute, Boston, MA)
598 was detected using the Opal 690 fluorophore (1:50, Cy5). Whole slide images were acquired at 10x using
599 the Vectra 3 automated quantitative pathology imaging system (PerkinElmer). Subsequently, at least 5
600 stamps of 931x698 um were selected per slide in areas of high immune infiltration (hotspots) using Perkin
601 Elmer Phenochart v 1.0 software. Each stamp was then acquired at 20x using the Vectra 3. Inform 2.2
602 software was then used in order to deconvolute the multispectral images, as previously described⁷¹.
603 Hotspot deconvoluted images in .tiff format were uploaded into Indica Lab HALO platform version 3.0.
604 For each hotspot, the tumor area was manually annotated by a pathologist (TD). CD8 cells were
605 phenotyped according to the expression of PD1, TIM3 and LAG3 using the Indica Lab High-Plex FL v2.0
606 module, using DAPI-based nuclear segmentation and detection of FITC (CD8), Cy3 (TIM3), Texas Red
607 (LAG3), Cy5 (PD1) positive cells by adapting a dye cytoplasm positive threshold for each slide. A unique
608 algorithm was created for each whole slide, and each group of hotspots and its accuracy was validated
609 through visual inspection by two pathologists (TD, SS). Sample-level results of the multiplex
610 immunofluorescence analysis are provided in **Supplementary Table 6**. Comparisons between tRCC (n=
611 11), ccRCC (n= 11), and normal (ccRCC adjacent, (n= 10)) were performed using Wilcoxon rank-sum
612 tests. All tRCC samples were either (1) *TFE3* FISH positive or (2) positive *TFE3* test by IHC along with a
613 strongly suggestive clinico-pathologic history and no FISH testing results available (missing). For each T

cell subset, T cell subset density was calculated as the number of T cells per mm². Percentage of a T cell subset was defined as the density of the T cell subset divided by the density of CD8⁺ T cells in the sample.

Immune deconvolution and immune analyses

CIBERSORTx⁷⁰ (Job type: “Impute cell fractions”), in absolute mode, with B mode batch correction, with quantile normalization disabled, and in 1000 permutations was used on the LM22 signature in order to infer the immune cell composition of samples from RNA-seq in the TCGA cohort. All samples which had a p-value for deconvolution >0.05 were considered to have failed deconvolution and were therefore discarded from all downstream analyses. Relative cell proportions were obtained by normalizing the CIBERSORTx output to the sample-level sum of cell counts (in order to obtain percentages of immune infiltration). Purity estimates for the TCGA cohort were obtained for the TCGA cohort from the Taylor et al. study¹⁰⁵. CD8⁺ T cell density and purity were compared pairwise between tRCC and each other RCC histology (ccRCC, pRCC, and chRCC) using Wilcoxon rank-sum tests. Sample-level PD-L1 protein expression by IHC on tumor-infiltrating immune cells (PD-L1 ≥ 1%) for the IMmotion151 trial were obtained from the Motzer et al. study³⁰ and compared using a Fisher’s exact test between tRCC and ccRCC.

Clinical and survival analyses

Tumor stage was obtained from Genomic Data Commons (<https://gdc.cancer.gov/about-data/publications/pancanatlas>) for the TCGA cohort and was defined using American Joint Committee on Cancer (AJCC) 8th edition for the IMDC and Harvard cohorts. IMDC risk groups (a previously validated prognostic model for patients with metastatic RCC) were defined as previously described¹³⁰. Tumor stage (I/II vs III/IV), IMDC risk groups (favorable, intermediate, poor), and sex were compared pairwise between tRCC and each other RCC histology (ccRCC, pRCC, and chRCC) using Fisher’s exact test. Age at initial RCC diagnosis was compared between tRCC and each other RCC histology (ccRCC, pRCC, and chRCC) using Wilcoxon rank-sum tests. Sankey diagrams for the Harvard and IMDC cohorts were computed using the ggalluvial package in R.

For all survival endpoints, the Kaplan-Meier method was used to summarize survival distributions. For the TCGA cohort, progression-free interval (PFI) was defined as the period from the date of diagnosis until the date of the first occurrence of a new tumor event (includes disease progression, locoregional recurrence, distant metastasis, new primary tumor, or death with tumor). Patients were censored if they were alive without any of these events at last follow-up or had died without tumor¹³¹. Overall survival (OS) was defined as the period from the start of systemic therapy until death. Patients were censored if they were alive at last follow-up. Time-to-treatment failure (TTF) was defined from the start of the line of systemic therapy to the end of that line of therapy or death from any cause. Since assessment of responses in retrospective cohorts (Harvard and IMDC cohorts) was not subject to radiological review specifically for the purpose of this study, responses were defined based on RECIST v1.1 criteria¹³² as available by retrospective review. Patients were censored if they were alive and still on the line of therapy at last follow-up. Progression-free survival (PFS) was defined (in the CheckMate and IMmotion151 cohorts) from the time of randomization or start of first dose until disease progression or death. Patients were censored if they were alive at last follow-up. For all survival endpoints, pairwise comparisons were performed using log-rank tests. In the IMmotion151 cohort, a Cox model that included an interaction term (histology-by-treatment arm) was used to evaluate the difference in the extent of benefit derived with atezolizumab + bevacizumab versus sunitinib in tRCC versus ccRCC. In the Harvard/IMDC pooled cohort, all patients who got ICI-based therapies were included in the ICI group. If patients received multiple lines of ICI-based therapies, the first ICI-based regimen was used for the analysis of clinical outcomes. All other patients had received TKIs and were assigned to the TKI group. If patients received multiple lines of TKIs, the first TKI regimen was used for the analysis of clinical outcomes.

Clinical benefit (CB) was defined as an objective response (complete response or partial response) or stable disease with PFS of at least 6 months. No clinical benefit (NCB) was defined as progressive disease with PFS less than 3 months. All other patients (not meeting criteria for CB or NCB) were classified as having intermediate clinical benefit (ICB). In the IMmotion151 cohort, rates of CB and NCB

were compared between the atezolizumab + bevacizumab and sunitinib arms, in patients with tRCC and ccRCC separately, using Fisher's exact test.

Statistics

All downstream analyses were done using R v3.6.1, Python v3.8.5 (on Spyder v4.1.5), Circos v0.69.9, or GraphPad PRISM 9. For boxplots, the upper and lower hinges represent the 75th and 25th percentiles, respectively. The whiskers extend in both directions until the largest or lowest value not further than 1.5 times the interquartile range from the corresponding hinge. All tests were two-tailed (unless otherwise specified) and considered statistically significant if $p < 0.05$ or $q < 0.05$.

Data availability statement

All relevant data are available from the authors and/or are included with the manuscript. The list of samples used (including the data types available) and sequencing platform used for DNA-sequencing (WGS, WES, or panel) are provided in **Supplementary Table 1**. The expression matrix (RSEM expected counts and TPMs) derived from the RNA-sequencing of the cell lines in the *in vitro* experiment represented in **Figure 4a** is provided in **Supplementary Table 3**. For the OncoPanel cohort, sample-level data (mutation, copy number, and clinical metadata) are provided in **Supplementary Table 5**. Sample-level data from the multiparametric immunofluorescence cohort are provided in **Supplementary Table 6**.

Code availability statement

Algorithms used for data analysis are all publicly available from the indicated references in the **Methods** section. Any other queries about the custom code used in this study should be directed to the corresponding authors of this study.

687 **Acknowledgements**

688 We thank the OncoPanel study team and the patients who contributed their data to research and
 689 participated in clinical trials. We thank all contributors to The International Metastatic Renal-Cell
 690 Carcinoma Database Consortium for their data contributions. This work was supported in part by The
 691 Friends of Dana-Farber (S.R.V.), Claudia Adams Barr Program for Innovative Cancer Research
 692 (S.R.V.), Clinician-Scientist Development Award from the Doris Duke Charitable Foundation (S.R.V.),
 693 Department of Defense Kidney Cancer Research Program (W81XWH-19-1-0815) (S.R.V.), an
 694 Independent Investigator Grant from the Rally Foundation for Childhood Cancer Research (S.R.V.), and
 695 the Dana-Farber / Harvard Cancer Center Kidney Cancer SPORE (P50-CA101942) (D.A.B., D.F.M.,
 696 S.S., T.K.C., S.R.V.). T.K.C. is supported by the Kohlberg Chair at Harvard Medical School and the
 697 Trust Family, Michael Brigham, and Loker Pinard Funds for Kidney Cancer Research at DFCI. J.N.
 698 acknowledges support by NIH F31 CA250136. N.I.V. is supported by Damon Runyon Physician-
 699 Scientist Training Award, Conquer Cancer Foundation YIA, SITC Genentech Award. D.A.B. is
 700 supported by the Dana-Farber / Harvard Cancer Center Kidney Cancer SPORE Career Enhancement
 701 Program (NCI P50CA101942-15), DoD CDMRP (KC170216, KC190130), and the DoD Academy of
 702 Kidney Cancer Investigators (KC190128).

703 **Author contributions**

704 Conception and Design: Ziad Bakouny, Eliezer M. Van Allen, Toni K. Choueiri, Srinivas R. Viswanathan
 705 Provision of study material or patients: Ziad Bakouny, Praful Ravi, Xin Gao, David A. Braun, Eliezer M.
 706 Van Allen, Toni K. Choueiri, Srinivas R. Viswanathan
 707 Collection and assembly of data: Ziad Bakouny, Praful Ravi, Jiao Li, Stephen Tang, Thomas Denize,
 708 Emma R. Garner, Laure Hirsch, John A. Steinharter, Gabrielle Bouchard, Emily Walton, Destiny West,
 709 Chris Labaki, Shaan Dudani, Chun-Loo Gan, Vidyalakshmi Sethunath, Filipe LF. Carvalho, Michelle S.
 710 Hirsch, Gwo-Shu Mary Lee, Bradley A. McGregor, Steven L. Chang, Adam S. Feldman, Catherine J.

711 Wu, David F. McDermott, Daniel Y.C. Heng, Eliezer M. Van Allen, Toni K. Choueiri, Srinivas R.
712 Viswanathan

713 Data Analysis and Interpretation: Ziad Bakouny, Ananthan Sadagopan, Nebiyu Y. Metaferia, Shatha
714 AbuHammad, Stephen Tang, Thomas Denize, David A. Braun, Alma Imamovic, Cora Ricker, Natalie I.
715 Vokes, Jackson Nyman, Jihye Park, Rizwan Haq, Eliezer M. Van Allen, Toni K. Choueiri, Srinivas R.
716 Viswanathan

717 Manuscript writing and revision: All authors

718 Final approval of manuscript: All authors

719 Accountable for all aspects of work: All authors

720 Competing Interests statement

721 The following authors report competing interests (all outside of the submitted work):

722 Z.B.: Research funding from Bristol-Myers Squibb & Genentech/imCORE; Honoraria from UpToDate.

723 X.G.: Advisory board for Exelixis.

724 D.A.B.: Nonfinancial support from Bristol Myers Squibb, honoraria from LM Education/Exchange
725 Services, advisory board fees from Exelixis, and consulting/personal fees from Octane Global, Defined
726 Health, Dedham Group, Adept Field Solutions, Slingshot Insights, Blueprint Partnerships, Charles River
727 Associates, Trinity Group, and Insight Strategy, outside of the submitted work.

728 N.I.V.: Advisory board to Sanofi/Genzyme.

729 M.S.H.: Consultant, Janssen Pharmaceuticals; UpToDate.

730 R.H.: Research funding from Novartis.

731 B.A.M.: Consultant for Bayer, Astellas, AstraZeneca, Seattle Genetics, Exelixis, Nektar, Pfizer,
732 Janssen, Genentech, Eisai, Dendreon, Bristol Myers Squibb, Calithera, and EMD Serono. Research
733 funding to the institution from Bristol Myers Squibb, Calithera, Exelixis, and Seattle Genetics.

734 A.S.F.: Olympus America, Inc – Consultant; Roche, Janssen – Honoraria; Vessi Medical - Advisory
735 Board.

736 C.J.W.: Equity holder of BioNTech, Inc; Research funding from Pharmacyclics.

737 D.F.M.: Honoraria from BMS, Pfizer, Merck, Alkermes, Inc., EMD Serono, Eli Lilly and Company,
738 lovance, Eisai Inc., Werewolf Therapeutics, and Calithera Biosciences; Research Support from BMS,
739 Merck, Genentech, Pfizer, Exelixis, X4 Pharma, and Alkermes, Inc.

740 D.Y.C.H.: Consultancies and research funding from Pfizer, Novartis, BMS, Merck, Eisai, Ipsen, Exelixis.

741 S.S.: Grants from Exelixis, grants from Bristol-Myers Squibb, personal fees from Merck, grants and
742 personal fees from AstraZeneca, personal fees from CRISPR Therapeutics, personal fees from NCI,
743 and personal fees from AACR; a patent for Biogenex with royalties paid.

744 E.M.V.A.: Advisory/Consulting: Tango Therapeutics, Genome Medical, Invitae, Enara Bio, Janssen,
745 Manifold Bio, Monte Rosa; Research support: Novartis, BMS; Equity: Tango Therapeutics, Genome
746 Medical, Syapse, Enara Bio, Manifold Bio, Microsoft, Monte Rosa; Patents: Institutional patents filed on
747 chromatin mutations and immunotherapy response, and methods for clinical interpretation.

748 T.K.C.: Research (Institutional and personal): Alexion, Analysis Group, AstraZeneca, Aveo, Bayer,
749 Bristol Myers-Squibb/ER Squibb and sons LLC, Calithera, Cerulean, Corvus, Eisai, Exelixis, F.
750 Hoffmann-La Roche, Foundation Medicine Inc., Genentech, GlaxoSmithKline, Ipsen, Lilly, Merck,
751 Novartis, Peloton, Pfizer, Prometheus Labs, Roche, Roche Products Limited, Sanofi/Aventis, Takeda,
752 Tracon; Consulting/honoraria or Advisory Role: Alexion, Analysis Group, AstraZeneca, Aveo, Bayer,
753 Bristol Myers-Squibb/ER Squibb and sons LLC, Cerulean, Corvus, Eisai, EMD Serono, Exelixis,
754 Foundation Medicine Inc., Genentech, GlaxoSmithKline, Heron Therapeutics, Infinity Pharma, Ipsen,
755 Jansen Oncology, IQVIA, Lilly, Merck, NCCN, Novartis, Peloton, Pfizer, Pionyr, Prometheus Labs,
756 Roche, Sanofi/Aventis, Surface Oncology, Tempest, Up-to-Date. CME-related events (e.g.: OncLive,
757 PVI, MJH Life Sciences). NCI GU Steering Committee; Stock ownership: Pionyr, Tempest; Patents
758 filed, royalties or other intellectual properties: related to biomarkers of immune checkpoint blockers and

759 ctDNA; Travel, accommodations, expenses, medical writing in relation to consulting, advisory roles, or
 760 honoraria; No speaker's bureau.

761 S.R.V.: Consulting, MPM Capital and Vida Ventures. Spouse is an employee of and holds equity in
 762 Kojin Therapeutics. J.L. & S.R.V: institutional patent application on targets in tRCC.

763 All other authors report no competing interests.

REFERENCES

1. Kauffman, E. C. *et al.* Molecular genetics and cellular features of TFE3 and TFEB fusion kidney cancers. *Nat Rev Urol* **11**, 465–475 (2014).
2. Moch, H., Cubilla, A. L., Humphrey, P. A., Reuter, V. E. & Ulbright, T. M. The 2016 WHO Classification of Tumours of the Urinary System and Male Genital Organs—Part A: Renal, Penile, and Testicular Tumours. *European Urology* **70**, 93–105 (2016).
3. Argani, P. *et al.* Xp11 Translocation Renal Cell Carcinoma in Adults: Expanded Clinical, Pathologic, and Genetic Spectrum. *The American Journal of Surgical Pathology* **31**, 1149–1160 (2007).
4. Calìò, A., Segala, D., Munari, E., Brunelli, M. & Martignoni, G. MiT Family Translocation Renal Cell Carcinoma: from the Early Descriptions to the Current Knowledge. *Cancers (Basel)* **11**, (2019).
5. Xu, L. *et al.* Xp11.2 translocation renal cell carcinomas in young adults. *BMC Urology* **15**, 57 (2015).
6. Baba, M. *et al.* TFE3 Xp11.2 Translocation Renal Cell Carcinoma Mouse Model Reveals Novel Therapeutic Targets and Identifies GPNMB as a Diagnostic Marker for Human Disease. *Mol Cancer Res* **17**, 1613–1626 (2019).
7. Comprehensive Molecular Characterization of Papillary Renal-Cell Carcinoma. *New England Journal of Medicine* **374**, 135–145 (2016).
8. Clark, J. *et al.* Fusion of splicing factor genes PSF and NonO (p54nrb) to the TFE3 gene in papillary renal cell carcinoma. *Oncogene* **15**, 2233–2239 (1997).
9. Linehan, W. M. Genetic basis of kidney cancer: role of genomics for the development of disease-based therapeutics. *Genome Res* **22**, 2089–2100 (2012).
10. Linehan, W. M. *et al.* The Metabolic Basis of Kidney Cancer. *Cancer Discov* **9**, 1006–1021 (2019).
11. Sidhar, S. K. *et al.* The t(X;1)(p11.2;q21.2) translocation in papillary renal cell carcinoma fuses a novel gene PRCC to the TFE3 transcription factor gene. *Hum Mol Genet* **5**, 1333–1338 (1996).

- 787 12. Zhong, M. *et al.* Dual-color, Break-apart FISH Assay on Paraffin-embedded Tissues as an Adjunct to Diagnosis
788 of Xp11 Translocation Renal Cell Carcinoma and Alveolar Soft Part Sarcoma. *The American Journal of*
789 *Surgical Pathology* **34**, 757–766 (2010).
- 790 13. Malouf, G. G. *et al.* Genomic heterogeneity of translocation renal cell carcinoma. *Clin. Cancer Res.* **19**, 4673–
791 4684 (2013).
- 792 14. Malouf, G. G. *et al.* Next-Generation Sequencing of Translocation Renal Cell Carcinoma Reveals Novel RNA
793 Splicing Partners and Frequent Mutations of Chromatin-Remodeling Genes. *Clin Cancer Res* **20**, 4129–4140
794 (2014).
- 795 15. Marcon, J. *et al.* Comprehensive Genomic Analysis of Translocation Renal Cell Carcinoma Reveals Copy-
796 Number Variations as Drivers of Disease Progression. *Clin. Cancer Res.* (2020) doi:10.1158/1078-0432.CCR-
797 19-3283.
- 798 16. Choueiri, T. K. & Motzer, R. J. Systemic Therapy for Metastatic Renal-Cell Carcinoma. *New England Journal of*
799 *Medicine* **376**, 354–366 (2017).
- 800 17. Ambalavanan, M. & Geller, J. I. Treatment of advanced pediatric renal cell carcinoma. *Pediatr Blood Cancer*
801 **66**, e27766 (2019).
- 802 18. Boilève, A. *et al.* Immune checkpoint inhibitors in MITF family translocation renal cell carcinomas and
803 genetic correlates of exceptional responders. *J. immunotherapy cancer* **6**, 159 (2018).
- 804 19. Chanzá, N. M. *et al.* Cabozantinib in advanced non-clear-cell renal cell carcinoma: a multicentre,
805 retrospective, cohort study. *Lancet Oncol* **20**, 581–590 (2019).
- 806 20. Choueiri, T. K. *et al.* Vascular endothelial growth factor-targeted therapy for the treatment of adult
807 metastatic Xp11.2 translocation renal cell carcinoma. *Cancer* **116**, 5219–5225 (2010).
- 808 21. Damayanti, N. P. *et al.* Therapeutic targeting of TFE3/IRS-1/PI3K/mTOR axis in translocation renal cell
809 carcinoma. *Clinical Cancer Research* clincanres.0269.2018 (2018) doi:10.1158/1078-0432.CCR-18-0269.
- 810 22. Malouf, G. G. *et al.* Targeted agents in metastatic Xp11 translocation/TFE3 gene fusion renal cell carcinoma
811 (RCC): a report from the Juvenile RCC Network. *Ann. Oncol.* **21**, 1834–1838 (2010).

- 812 23. Rua Fernández, O. R. *et al.* Renal Cell Carcinoma Associated With Xp11.2 Translocation/TFE3 Gene-fusion: A
813 Long Response to mammalian target of rapamycin (mTOR) Inhibitors. *Urology* **117**, 41–43 (2018).
- 814 24. Hayes, M. *et al.* Molecular-genetic analysis is essential for accurate classification of renal carcinoma
815 resembling Xp11.2 translocation carcinoma. *Virchows Arch* **466**, 313–322 (2015).
- 816 25. Wang, X.-T. *et al.* RNA sequencing of Xp11 translocation-associated cancers reveals novel gene fusions and
817 distinctive clinicopathologic correlations. *Mod. Pathol.* (2018).
- 818 26. Chen, F. *et al.* Multilevel Genomics-Based Taxonomy of Renal Cell Carcinoma. *Cell Reports* **14**, 2476–2489
819 (2016).
- 820 27. Ricketts, C. J. *et al.* The Cancer Genome Atlas Comprehensive Molecular Characterization of Renal Cell
821 Carcinoma. *Cell Reports* **23**, 313–326.e5 (2018).
- 822 28. Sukov, W. R. *et al.* TFE3 rearrangements in adult renal cell carcinoma: clinical and pathologic features with
823 outcome in a large series of consecutively treated patients. *Am J Surg Pathol* **36**, 663–670 (2012).
- 824 29. Gao, Q. *et al.* Driver Fusions and Their Implications in the Development and Treatment of Human Cancers.
825 *Cell Reports* **23**, 227–238.e3 (2018).
- 826 30. Motzer, R. J. *et al.* Molecular Subsets in Renal Cancer Determine Outcome to Checkpoint and Angiogenesis
827 Blockade. *Cancer Cell* **38**, 803–817.e4 (2020).
- 828 31. Sato, Y. *et al.* Integrated molecular analysis of clear-cell renal cell carcinoma. *Nature Genetics* **45**, 860–867
829 (2013).
- 830 32. Durinck, S. *et al.* Spectrum of diverse genomic alterations define non-clear cell renal carcinoma subtypes.
831 *Nat Genet* **47**, 13–21 (2015).
- 832 33. Huang, F. W. *et al.* Highly recurrent TERT promoter mutations in human melanoma. *Science* **339**, 957–959
833 (2013).
- 834 34. Taylor, A. M. *et al.* Genomic and Functional Approaches to Understanding Cancer Aneuploidy. *Cancer cell*
835 (2018) doi:10.1016/j.ccell.2018.03.007.

- 836 35. Cunningham, J. L. *et al.* Common pathogenetic mechanism involving human chromosome 18 in familial and
837 sporadic ileal carcinoid tumors. *Genes Chromosomes Cancer* **50**, 82–94 (2011).
- 838 36. Brady, S. W. *et al.* Pan-neuroblastoma analysis reveals age- and signature-associated driver alterations. *Nat*
839 *Commun* **11**, (2020).
- 840 37. Suarez-Merino, B. *et al.* Microarray analysis of pediatric ependymoma identifies a cluster of 112 candidate
841 genes including four transcripts at 22q12.1-q13.3,. *Neuro-oncol* **7**, 20–31 (2005).
- 842 38. Hershey, C. L. & Fisher, D. E. Mitf and Tfe3: members of a b-HLH-ZIP transcription factor family essential for
843 osteoclast development and function. *Bone* **34**, 689–696 (2004).
- 844 39. Hoadley, K. A. *et al.* Cell-of-Origin Patterns Dominate the Molecular Classification of 10,000 Tumors from 33
845 Types of Cancer. *Cell* **173**, 291-304.e6 (2018).
- 846 40. Wang, X. *et al.* Malignant melanotic Xp11 neoplasms exhibit a clinicopathologic spectrum and gene
847 expression profiling akin to alveolar soft part sarcoma: a proposal for reclassification. *J. Pathol.* **251**, 365–
848 377 (2020).
- 849 41. Pan-cancer analysis of whole genomes | Nature. <https://www.nature.com/articles/s41586-020-1969-6>.
- 850 42. Hoek, K. S. *et al.* Novel MITF targets identified using a two-step DNA microarray strategy. *Pigment Cell &*
851 *Melanoma Research* **21**, 665–676 (2008).
- 852 43. Lachmann, A. *et al.* ChEA: transcription factor regulation inferred from integrating genome-wide ChIP-X
853 experiments. *Bioinformatics* **26**, 2438–2444 (2010).
- 854 44. Canela, L. *et al.* Transcriptional profiling of striatal neurons in response to single or concurrent activation of
855 dopamine D2, adenosine A2A and metabotropic glutamate type 5 receptors: Focus on beta-synuclein
856 expression. *Gene* **508**, 199–205 (2012).
- 857 45. Boyer, N. P., Monkiewicz, C., Menon, S., Moy, S. S. & Gupton, S. L. Mammalian TRIM67 Functions in Brain
858 Development and Behavior. *eNeuro* **5**, (2018).
- 859 46. Star, E. N. *et al.* Regulation of retinal interneuron subtype identity by the Iroquois homeobox gene *lrx6*.
860 *Development* **139**, 4644–4655 (2012).

861 47. Rezaei, K. A., Chen, Y., Cai, J. & Sternberg, P. Modulation of Nrf2-Dependent Antioxidant Functions in the
862 RPE by Zip2, a Zinc Transporter Protein. *Invest. Ophthalmol. Vis. Sci.* **49**, 1665–1670 (2008).

863 48. Kim, H. *et al.* Tmem64 modulates calcium signaling during RANKL-mediated osteoclast differentiation. *Cell*
864 *Metab* **17**, 249–260 (2013).

865 49. Sun, M., Luong, G., Plastikwala, F. & Sun, Y. Control of Rab7a activity and localization through endosomal
866 type Igamma PIP 5-kinase is required for endosome maturation and lysosome function. *FASEB j.* **34**, 2730–
867 2748 (2020).

868 50. Groenewoud, M. J. & Zwartkuis, F. J. T. Rheb and Rags come together at the lysosome to activate mTORC1.
869 *Biochemical Society Transactions* **41**, 951–955 (2013).

870 51. McConnell, M. *et al.* Atp6v1c1 Facilitates Breast Cancer Growth and Bone Metastasis through the mTORC1
871 Pathway. *The FASEB Journal* **29**, 284.8 (2015).

872 52. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-
873 wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 15545–15550 (2005).

874 53. Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Systems* **1**, 417–425
875 (2015).

876 54. Menegon, S., Columbano, A. & Giordano, S. The Dual Roles of NRF2 in Cancer. *Trends in Molecular Medicine*
877 **22**, 578–593 (2016).

878 55. Clerici, S. & Boletta, A. Role of the KEAP1-NRF2 Axis in Renal Cell Carcinoma. *Cancers* **12**, 3458 (2020).

879 56. Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-Seq
880 data. *BMC Bioinformatics* **14**, 7 (2013).

881 57. Takahashi, N. *et al.* 3D Culture Models with CRISPR Screens Reveal Hyperactive NRF2 as a Prerequisite for
882 Spheroid Formation via Regulation of Proliferation and Ferroptosis. *Mol Cell* **80**, 828-844.e6 (2020).

883 58. DeNicola, G. M. *et al.* Oncogene-induced Nrf2 transcription promotes ROS detoxification and tumorigenesis.
884 *Nature* **475**, 106–109 (2011).

- 885 59. Harris, I. S. *et al.* Deubiquitinases Maintain Protein Homeostasis and Survival of Cancer Cells upon
886 Glutathione Depletion. *Cell Metab* **29**, 1166-1181.e6 (2019).
- 887 60. Xue, D., Zhou, X. & Qiu, J. Emerging role of NRF2 in ROS-mediated tumor chemoresistance. *Biomed*
888 *Pharmacother* **131**, 110676 (2020).
- 889 61. Rees, M. G. *et al.* Correlating chemical sensitivity and basal gene expression reveals mechanism of action.
890 *Nat Chem Biol* **12**, 109–116 (2016).
- 891 62. Viswanathan, V. S. *et al.* Dependency of a therapy-resistant state of cancer cells on a lipid peroxidase
892 pathway. *Nature* **547**, 453–457 (2017).
- 893 63. Ghandi, M. *et al.* Next-generation characterization of the Cancer Cell Line Encyclopedia. *Nature* **569**, 503–
894 508 (2019).
- 895 64. Tsherniak, A. *et al.* Defining a Cancer Dependency Map. *Cell* **170**, 564-576.e16 (2017).
- 896 65. Rini, B. I. *et al.* Atezolizumab plus bevacizumab versus sunitinib in patients with previously untreated
897 metastatic renal cell carcinoma (IMmotion151): a multicentre, open-label, phase 3, randomised controlled
898 trial. *The Lancet* **393**, 2404–2415 (2019).
- 899 66. Braun, D. A. *et al.* Interplay of somatic alterations and immune infiltration modulates response to PD-1
900 blockade in advanced clear cell renal cell carcinoma. *Nat Med* **26**, 909–918 (2020).
- 901 67. Motzer, R. J. *et al.* Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. *New England Journal of*
902 *Medicine* **373**, 1803–1813 (2015).
- 903 68. McDermott, D. F. *et al.* Open-Label, Single-Arm, Phase II Study of Pembrolizumab Monotherapy as First-Line
904 Therapy in Patients With Advanced Non-Clear Cell Renal Cell Carcinoma. *J Clin Oncol* **39**, 1029–1039 (2021).
- 905 69. McGregor, B. A. *et al.* Results of a Multicenter Phase II Study of Atezolizumab and Bevacizumab for Patients
906 With Metastatic Renal Cell Carcinoma With Variant Histology and/or Sarcomatoid Features. *J Clin Oncol* **38**,
907 63–70 (2020).
- 908 70. Newman, A. M. *et al.* Determining cell type abundance and expression from bulk tissues with digital
909 cytometry. *Nature Biotechnology* **37**, 773–782 (2019).

- 910 71. Pignon, J.-C. *et al.* irRECIST for the Evaluation of Candidate Biomarkers of Response to Nivolumab in
911 Metastatic Clear Cell Renal Cell Carcinoma: Analysis of a Phase II Prospective Clinical Trial. *Clin Cancer Res*
912 **25**, 2174–2184 (2019).
- 913 72. Ficial, M. *et al.* Expression of T-Cell Exhaustion Molecules and Human Endogenous Retroviruses as Predictive
914 Biomarkers for Response to Nivolumab in Metastatic Clear Cell Renal Cell Carcinoma. *Clin Cancer Res* **27**,
915 1371–1380 (2021).
- 916 73. Engreitz, J. M., Agarwala, V. & Mirny, L. A. Three-Dimensional Genome Architecture Influences Partner
917 Selection for Chromosomal Translocations in Human Disease. *PLOS ONE* **7**, e44196 (2012).
- 918 74. Lin, C. *et al.* Nuclear Receptor-Induced Chromosomal Proximity and DNA Breaks Underlie Specific
919 Translocations in Cancer. *Cell* **139**, 1069–1083 (2009).
- 920 75. Roix, J. J., McQueen, P. G., Munson, P. J., Parada, L. A. & Misteli, T. Spatial proximity of translocation-prone
921 gene loci in human lymphomas. *Nature Genetics* **34**, 287–291 (2003).
- 922 76. Inamura, K. *et al.* Diverse Fusion Patterns and Heterogeneous Clinicopathologic Features of Renal Cell
923 Carcinoma With t(6;11) Translocation. *The American Journal of Surgical Pathology* **36**, 35–42 (2012).
- 924 77. Xia, Q.-Y. *et al.* Novel gene fusion of PRCC–MITF defines a new member of MiT family translocation renal cell
925 carcinoma: clinicopathological analysis and detection of the gene fusion by RNA sequencing and FISH.
926 *Histopathology* **72**, 786–794 (2018).
- 927 78. Álvarez-Fernández, M. & Malumbres, M. Mechanisms of Sensitivity and Resistance to CDK4/6 Inhibition.
928 *Cancer Cell* **37**, 514–529 (2020).
- 929 79. Kryukov, G. V. *et al.* MTAP deletion confers enhanced dependency on the PRMT5 arginine
930 methyltransferase in cancer cells. *Science* **351**, 1214–1218 (2016).
- 931 80. Mavrakis, K. J. *et al.* Disordered methionine metabolism in MTAP/CDKN2A-deleted cancers leads to
932 dependence on PRMT5. *Science* **351**, 1208–1213 (2016).
- 933 81. Garraway, L. A. *et al.* Integrative genomic analyses identify MITF as a lineage survival oncogene amplified in
934 malignant melanoma. *Nature* **436**, 117–122 (2005).

- 935 82. Liu, L. *et al.* Mutation of the CDKN2A 5' UTR creates an aberrant initiation codon and predisposes to
936 melanoma. *Nature Genetics* **21**, 128–132 (1999).
- 937 83. Walker, G. J. *et al.* Virtually 100% of melanoma cell lines harbor alterations at the DNA level within CDKN2A,
938 CDKN2B, or one of their downstream targets. *Genes, Chromosomes and Cancer* **22**, 157–163 (1998).
- 939 84. Sporn, M. & Liby, K. NRF2 and cancer: the good, the bad and the importance of context. *Nature reviews.*
940 *Cancer* **12**, (2012).
- 941 85. Li, L. *et al.* SQSTM1 Is a Pathogenic Target of 5q Copy Number Gains in Kidney Cancer. *Cancer Cell* **24**, 738–
942 750 (2013).
- 943 86. Huang, H. *et al.* Downregulation of Keap1 contributes to poor prognosis and Axitinib resistance of renal cell
944 carcinoma via upregulation of Nrf2 expression. *Int J Mol Med* (2019) doi:10.3892/ijmm.2019.4134.
- 945 87. Ji, S., Xiong, Y., Zhao, X., Liu, Y. & Yu, L. Q. Effect of the Nrf2-ARE signaling pathway on biological
946 characteristics and sensitivity to sunitinib in renal cell carcinoma. *Oncology Letters* **17**, 5175–5186 (2019).
- 947 88. Romero, R. *et al.* Keap1 mutation renders lung adenocarcinomas dependent on Slc33a1. *Nature Cancer* **1**,
948 589–602 (2020).
- 949 89. Yang, W. *et al.* Immunogenic neoantigens derived from gene fusions stimulate T cell responses. *Nature*
950 *Medicine* **25**, 767–775 (2019).
- 951 90. Braun, D. A. *et al.* Beyond conventional immune-checkpoint inhibition — novel immunotherapies for renal
952 cell carcinoma. *Nat Rev Clin Oncol* **18**, 199–214 (2021).
- 953 91. Jansen, C. S. *et al.* An intra-tumoral niche maintains and differentiates stem-like CD8 T cells. *Nature* **576**,
954 465–470 (2019).
- 955 92. Pignon, J.-C. *et al.* irRECIST for the Evaluation of Candidate Biomarkers of Response to Nivolumab in
956 Metastatic Clear Cell Renal Cell Carcinoma: Analysis of a Phase II Prospective Clinical Trial. *Clin Cancer Res*
957 **25**, 2174–2184 (2019).

958 93. Press Release. Bristol Myers Squibb Announces RELATIVITY-047, a Trial Evaluating Anti-LAG-3 Antibody
959 Relatlimab and Opdivo (nivolumab) in Patients with Previously Untreated Metastatic or Unresectable
960 Melanoma, Meets Primary Endpoint of Progression-Free Survival.

961 94. Best, S. A. *et al.* Synergy between the KEAP1/NRF2 and PI3K Pathways Drives Non-Small-Cell Lung Cancer
962 with an Altered Immune Microenvironment. *Cell Metabolism* **27**, 935-943.e4 (2018).

963 95. Zhong, M. *et al.* Translocation renal cell carcinomas in adults: a single-institution experience. *Am J Surg*
964 *Pathol* **36**, 654–662 (2012).

965 96. Heng, D. Y. C. *et al.* External validation and comparison with other models of the International Metastatic
966 Renal-Cell Carcinoma Database Consortium prognostic model: a population-based study. *Lancet Oncol* **14**,
967 141–148 (2013).

968 97. Zehir, A. *et al.* Mutational landscape of metastatic cancer revealed from prospective clinical sequencing of
969 10,000 patients. *Nature Medicine* **23**, 703–713 (2017).

970 98. Campbell, P. J. *et al.* Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2020).

971 99. Malouf, G. G. *et al.* Next-Generation Sequencing of Translocation Renal Cell Carcinoma Reveals Novel RNA
972 Splicing Partners and Frequent Mutations of Chromatin-Remodeling Genes. *Clin Cancer Res* **20**, 4129–4140
973 (2014).

974 100. Garcia, E. P. *et al.* Validation of OncoPanel: A Targeted Next-Generation Sequencing Assay for the
975 Detection of Somatic Variants in Cancer. *Arch Pathol Lab Med* **141**, 751–758 (2017).

976 101. Ellrott, K. *et al.* Scalable Open Science Approach for Mutation Calling of Tumor Exomes Using Multiple
977 Genomic Pipelines. *cells* **6**, 271-281.e7 (2018).

978 102. Ramos, A. H. *et al.* Oncotator: cancer variant annotation tool. *Hum Mutat* **36**, E2423-2429 (2015).

979 103. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation. *Nature* **526**,
980 68–74 (2015).

981 104. Vokes, N. I. *et al.* Harmonization of Tumor Mutational Burden Quantification and Association With
982 Response to Immune Checkpoint Blockade in Non–Small-Cell Lung Cancer. *JCO Precision Oncology* 1–12
983 (2019) doi:10.1200/PO.19.00171.

984 105. Taylor, A. M. *et al.* Genomic and Functional Approaches to Understanding Cancer Aneuploidy. *Cancer*
985 *Cell* **33**, 676–689.e3 (2018).

986 106. Turajlic, S. *et al.* Tracking Cancer Evolution Reveals Constrained Routes to Metastases: TRACERx Renal.
987 *Cell* **173**, 581–594.e12 (2018).

988 107. Bakouny, Z. *et al.* Integrative molecular characterization of sarcomatoid and rhabdoid renal cell
989 carcinoma. *Nature Communications* **12**, 808 (2021).

990 108. Harrer, M., Cuijpers, P., Furukawa, T. & Ebert, D. Doing meta-analysis in R: A hands-on guide. *PROTECT*
991 *Lab Erlangen* (2019).

992 109. Efthimiou, O. Practical guide to the meta-analysis of rare events. *Evid Based Ment Health* **21**, 72–76
993 (2018).

994 110. Veroniki, A. A. *et al.* Methods to estimate the between-study variance and its uncertainty in meta-
995 analysis. *Res Synth Methods* **7**, 55–79 (2016).

996 111. Sondka, Z. *et al.* The COSMIC Cancer Gene Census: describing genetic dysfunction across all human
997 cancers. *Nature Reviews Cancer* **18**, 696–705 (2018).

998 112. Crowdis, J., He, M. X., Reardon, B. & Van Allen, E. M. CoMut: visualizing integrated molecular
999 information with comutation plots. *Bioinformatics* **36**, 4348–4349 (2020).

1000 113. Krzywinski, M. *et al.* Circos: An information aesthetic for comparative genomics. *Genome Res.* **19**, 1639–
1001 1645 (2009).

1002 114. Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Current Protocols* **1**, e90 (2021).

1003 115. Marchler-Bauer, A. *et al.* CDD: NCBI’s conserved domain database. *Nucleic Acids Res* **43**, D222–226
1004 (2015).

1005 116. Lancaster, A. K., Nutter-Upham, A., Lindquist, S. & King, O. D. PLAAC: a web and command-line
1006 application to identify proteins with prion-like amino acid composition. *Bioinformatics* **30**, 2501–2502
1007 (2014).

1008 117. Liu, W. *et al.* IBS: an illustrator for the presentation and visualization of biological sequences.
1009 *Bioinformatics* **31**, 3359–3361 (2015).

1010 118. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

1011 119. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a
1012 reference genome. *BMC Bioinformatics* **12**, 323 (2011).

1013 120. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray
1014 studies. *Nucleic Acids Res* **43**, e47 (2015).

1015 121. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional
1016 genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

1017 122. Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of intersecting sets
1018 and their properties. *Bioinformatics* **33**, 2938–2940 (2017).

1019 123. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**,
1020 417–425 (2015).

1021 124. Takahashi, N. *et al.* 3D Culture Models with CRISPR Screens Reveal Hyperactive NRF2 as a Prerequisite
1022 for Spheroid Formation via Regulation of Proliferation and Ferroptosis. *Molecular Cell* **80**, 828-844.e6
1023 (2020).

1024 125. McFarland, J. M. *et al.* Improved estimation of cancer dependencies from large-scale RNAi screens using
1025 model-based normalization and data integration. *Nat Commun* **9**, 4610 (2018).

1026 126. Meyers, R. M. *et al.* Computational correction of copy number effect improves specificity of CRISPR-Cas9
1027 essentiality screens in cancer cells. *Nat Genet* **49**, 1779–1784 (2017).

1028 127. Dempster, J. M. *et al.* Extracting Biological Insights from the Project Achilles Genome-Scale CRISPR
1029 Screens in Cancer Cell Lines. *bioRxiv* 720243 (2019) doi:10.1101/720243.

1030 128. Seashore-Ludlow, B. *et al.* Harnessing Connectivity in a Large-Scale Small-Molecule Sensitivity Dataset.
1031 *Cancer Discov* **5**, 1210–1223 (2015).

1032 129. Viswanathan, S. R. *et al.* Genome-scale analysis identifies paralog lethality as a vulnerability of
1033 chromosome 1p loss in cancer. *Nat Genet* **50**, 937–943 (2018).

1034 130. Heng, D. Y. C. *et al.* Prognostic factors for overall survival in patients with metastatic renal cell carcinoma
1035 treated with vascular endothelial growth factor-targeted agents: results from a large, multicenter study. *J*
1036 *Clin Oncol* **27**, 5794–5799 (2009).

1037 131. Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival
1038 Outcome Analytics. *Cell* **173**, 400-416.e11 (2018).

1039 132. Eisenhauer, E. A. *et al.* New response evaluation criteria in solid tumours: Revised RECIST guideline
1040 (version 1.1). *European Journal of Cancer* **45**, 228–247 (2009).

1041 133. Rees, M. G. *et al.* Correlating chemical sensitivity and basal gene expression reveals mechanism of
1042 action. *Nat Chem Biol* **12**, 109–116 (2016).

1043 134. Wang, X. *et al.* Malignant melanotic Xp11 neoplasms exhibit a clinicopathological spectrum and gene
1044 expression profiling akin to alveolar soft part sarcoma: a proposal for reclassification. *The Journal of*
1045 *Pathology* **n/a**,.

1046

1047

FIGURES

Fig. 1 | Identification of tRCC cases in multiple clinical and molecular datasets. **a**, *Top*, Kaplan-Meier curves for time-to-treatment failure in metastatic ccRCC, pRCC, chrRCC, or tRCC (Harvard cohort). *Bottom*, Kaplan-Meier curves for progression-free interval for localized ccRCC, pRCC, chrRCC, or tRCC (TCGA cohort). P-values were calculated by pairwise log-rank test. **b**, Representative H&E micrographs (x10) of cases originally included in the TCGA ccRCC or pRCC sequencing cohorts. The right case in each pair was subsequently found to have a *TFE3* gene fusion on RNA-Seq. **c**, Aggregation of tRCC cases from across 9 independent NGS datasets. The data type(s) analyzed are indicated for each dataset. tRCC cases were identified based on the presence of a fusion involving an *MIT/TFE* family member (see **Methods**). The number and proportion of tRCC samples as well as number of total RCC samples is indicated for each dataset.

Fig. 2 | Landscape of genomic alterations in tRCC. a, CoMut plot of mutational and copy number alterations in tRCC across all datasets. Genes listed include those found to be most frequently altered in tRCC across all datasets as well as previously reported significantly mutated genes in ccRCC, pRCC, and chrRCC²⁷ (indicated in the left track). **b**, Pairwise enrichment analysis for genomic alteration frequencies in tRCC versus other RCC histologies for the indicated genes, presented as the pooled odds ratio and 95% confidence interval from the random-effects meta-analysis in tRCC versus comparator histology. From left to right, genes listed on plots indicate: most frequently altered genes in tRCC, significantly mutated genes in ccRCC, significantly mutated genes in pRCC, and significantly mutated genes in chRCC. Pairwise enrichment between tRCC and comparator was calculated individually for each locus or gene within each dataset and pooled estimates across datasets were obtained as detailed in **Methods**.

Fig. 3 | Structure of *Mit/TFE* fusions in tRCC. a, Number and percentage of tRCC cases displaying gene fusions involving *TFE3*, *TFEB*, or *MITF* across all datasets analyzed. **b**, Genomic location of *Mit/TFE* fusion partners. Stroke thickness is proportional to the number of times a given gene was observed to be an *Mit/TFE* fusion partner across all datasets analyzed. **c**, Gene ontology terms (GO Biological Process) enriched amongst *Mit/TFE* fusion partners. **d**, Breakpoints observed within *TFE3*, *TFEB*, or *MITF* across all samples analyzed. Solid portion represents the portion of the *Mit/TFE* gene retained within the oncogenic fusion product. Fusion partner genes observed to join at a given breakpoint are listed. Functional domains within each *Mit/TFE* gene are indicated (legend in **Supplementary Table 2**). **e**, Breakpoints observed within *Mit/TFE* partner genes. Solid portion represents the portion of each partner gene retained within the oncogenic fusion product. Genes are grouped by whether they were observed to fuse with *TFE3* (top), *TFEB* (middle), or *MITF* (bottom). Functional domains within each *Mit/TFE* partner gene are indicated (legend in **Supplementary Table 2**).

Fig. 4 | Distinctive transcriptional features of tRCC. **a**, Schematic of *in vitro* experiment used to derive *TFE3*-fusion-specific transcriptional signature. **b**, Transcriptome sequencing data from three independent datasets (TCGA, PCAWG, IMmotion151) were subjected to unsupervised hierarchical clustering using the fusion-specific signature derived in (a). Blue bars indicate *Mit/TFE*-fusion-positive cases within each dataset. Gray bars indicate other RCC histologic subtypes or normal kidney. **c**, Heatmap of genes overexpressed in tRCC as compared with other RCC subtypes or normal kidney, across all datasets (see **Fig.S4**). Membership of genes in key pathways related to tRCC pathogenesis is indicated in the track at left. **d**, Gene set enrichment analysis showing top enriched Hallmark pathways in tRCC samples versus comparators across all datasets analyzed. Dataset and pairwise comparison across which the GSEA was performed is indicated in the track at the top of each column. Dot size is proportional to normalized enrichment score (NES) in tRCC versus comparator; dot color reflects $-\log_{10}(q\text{-value})$ for the enrichment.

1099 **Fig. 5 | tRCC displays activated NRF2 pathway signaling and a relative resistance to targeted therapies. a,**
1100 Waterfall plot showing NRF2 signature score for all RCC samples across all datasets analyzed. tRCC samples
1101 are depicted in blue (n=46); other samples (ccRCC, pRCC, chRCC, normal kidney, or other tumors) are shown in
1102 gray (n=1999). **b,** NRF2 signature score for TCGA RCC samples of the indicated histologies. For each histology,
1103 samples with somatic alterations in the NRF2 pathway are shown separately. No chRCC or tRCC samples
1104 displayed somatic alterations in the NRF2 pathway. **c,** Gene set enrichment analysis showing enrichment of
1105 NRF2 gene signature in 293T cells expressing *TFE3* fusions versus mock (untransfected) control condition. **d,**
1106 Volcano plot showing correlation of NRF2 signature score with drug sensitivity in the Broad Institute Cancer
1107 Therapeutics Response Portal dataset¹³³. A high NRF2 signature score is significantly associated with resistance
1108 to the agents shown in red. Agents annotated to act through the induction of oxidative stress or ferroptosis are
1109 colored in purple. Selected targeted agents used in the treatment of kidney cancer are labeled. **e,** Progression-
1110 free survival curves for tRCC (dark and light orange) or ccRCC (dark and light purple) patients treated with either
1111 atezolizumab and bevacizumab (AtezoBev) or sunitinib in the randomized Phase III IMmotion151 trial. **f,**
1112 Progression-free survival curves for ccRCC patients with high (red) or low (blue) NRF2 signature score treated
1113 with either sunitinib (top) atezolizumab + bevacizumab (bottom) on the IMmotion151 trial. For **e-f**, NRF2 signature
1114 score was dichotomized at the median in each arm.

1115

Fig. 6 | Immunogenomic features of tRCC associated with responses to immune checkpoint inhibition. a, Percentage of tRCC patients showing clinical benefit (CB), intermediate clinical benefit (ICB), or no clinical benefit (NCB) to either AtezoBev or sunitinib on the IMmotion151 trial. **b,** Swimmer plot showing response types and response times to immune checkpoint inhibitor-based regimens in tRCC patients in the combined IMDC and Harvard retrospective cohort. Line (L) in which ICI was received as well as specific ICI regimen is indicated to the right of each patient. **c,** Sample purity in tRCC, ccRCC, chRCC, and pRCC in the TCGA cohort. **d,** CD8+ T cell infiltration imputed from gene expression (CIBERSORTx) in tRCC, ccRCC, chRCC, and pRCC in the TCGA cohort. **e,** Multiparametric immunofluorescence for CD8, TIM3, LAG3, and PD1 in representative tRCC cases (top two rows) and ccRCC cases (bottom two rows). Red arrows indicate CD8+PD1+LAG3+TIM3+ tumor-infiltrating T cells in tRCC cases. Yellow arrows indicate CD8+PD1+LAG3+TIM3+ tumor-infiltrating T cells in ccRCC cases. **f,** Quantification of CD8+ T-cell density (top), percentage of CD8+PD1+TIM3+LAG3+ T cells (middle), and percentage of CD8+PD1+TIM3+LAG3+ T cells (bottom) in tRCC (n= 11), ccRCC (n= 11), and adjacent normal tissue (from ccRCC cases, n= 10).

1131 SUPPLEMENTARY FIGURES

1133 **Fig. S1 | Clinical features of tRCC.** **a**, Kaplan-Meier curves for time-to treatment-failure in metastatic ccRCC,
1134 pRCC, chrRCC, and tRCC from patients in the IMDC cohort. **b**, Proportion of male and female ccRCC, pRCC,
1135 chrRCC, and tRCC cases in the Harvard, IMDC, and TCGA cohorts. **c**, Age distribution of tRCC, ccRCC,
1136 chrRCC, and pRCC cases in the Harvard, IMDC, and TCGA datasets. **d** Distribution stage at diagnosis
1137 among ccRCC, pRCC, chrRCC, and tRCC patients in the Harvard and IMDC cohorts. **e**, Distribution of IMDC
1138 risk group at start of first-line of systemic therapy among ccRCC, pRCC, chrRCC, and tRCC patients in the
1139 Harvard and IMDC cohorts. **f**, Number of tRCC samples with DNA sequencing (WGS, WES, or Panel
1140 sequencing), RNA sequencing, or both data types, available for analysis across all NGS data sets.

1141

Fig. S2 | Genomic features of tRCC as compared with other RCC subtypes. **a**, Number of mutations per sample in tRCC versus other RCC histologies in the TCGA, MSK, and OncoPanel cohorts. **b**, Number of indels per sample in tRCC versus other RCC histologies in the TCGA, MSK, and OncoPanel cohorts. **c**, Number of frameshift indels per sample in tRCC versus other RCC histologies in the TCGA, MSK, and OncoPanel cohorts. In **a-c**, for the OncoPanel and MSK cohorts, the numbers of mutations and indels were normalized to the bait set of each version of each panel (**Methods**). **d**, *Left*, Aneuploidy score³⁴ in tRCC versus other RCC histologies in the TCGA cohort. *Right*, Fraction of genome altered in tRCC versus other

1150 RCC histologies in the MSK cohort. **e**, Frequency of arm-level copy number alterations among tRCC
 1151 samples in the TCGA cohort³⁴.
 1152

a

b

c

Fig. S3 | Characterization of *MIT/TFE* fusion partners. a, Frequency of various partner genes observed to fuse with *TFE3*, *TFEB*, or *MITF* across all datasets. b-c, Terms enriched amongst *MIT/TFE* fusion partner genes using either the GO Molecular Function (b) or WikiPathways (c) databases.

1159 **Fig. S4 | Transcriptional features of tRCC.** **a**, Expression of genes included in the *in vitro*-derived *TFE3*
1160 fusion-specific gene signature. **b**, Hierarchical clustering of RNA-Seq data¹³⁴ based on fusion-specific gene
1161 signature. **c**, Quality of clustering (based on $-\log_2(\text{intra-tRCC distance})$) in the TCGA, PCAWG, Wang et al.,
1162 or IMmotion151 datasets using either 1000 most variable genes (grey) or the fusion-specific gene signature
1163 (orange). **d**, Upset plot showing overlap of upregulated ($q < 0.05$) genes in tRCC versus other sample types
1164 in each of the datasets analyzed. **e**, Volcano plots showing differentially expressed genes in tRCC samples
1165 versus normal kidney, ccRCC, pRCC, and chrRCC in the TCGA cohort. Labels indicate selected genes that
1166 emerged as commonly upregulated in tRCC versus other sample types (**Figures 4c and S4c**) across all
1167 datasets analyzed.

1168

Fig. S5 | Activation of the NRF2 pathway in tRCC. **a**, NRF2 signature score in tRCC samples compared with ccRCC, pRCC, or chrRCC samples from the TCGA effort. Papillary RCC subtypes are annotated as previously described^{26,27}. Somatic alterations in the NRF2 pathway genes are indicated on the bottom track. **b**, Volcano plot displaying gene dependencies correlated to high NRF2 score in the DepMap RNAi (top) and CRISPR (bottom) datasets. **c**, Colony-forming assay in three tRCC cell lines (FU-UR1, UOK109, UOK146) transduced with a lentiviral doxycycline-inducible shRNA targeting *NRF2*. Quantification represents mean $\pm$ s.d. for $n=3$ independent replicates. **d**, Progression-free survival curves for ccRCC patients with high (red) or low (blue) NRF2 signature score treated with either everolimus (top) or nivolumab (bottom) in the CheckMate cohort. NRF2 signature score was dichotomized at the median in each arm.

1180 **Fig. S6 | Immunogenomic features and treatment patterns in tRCC.** **a**, Sankey flow diagram showing
1181 lines of systemic treatment received by patients with metastatic tRCC in the retrospective IMDC cohort
1182 (n=40). **b**, Sankey flow diagram showing lines of systemic treatment received by patients with metastatic
1183 tRCC in the retrospective Harvard cohort (n=17). **c**, Kaplan-Meier curves for overall survival in metastatic tRCC
1184 patients who received ICI-based (n=12) or tyrosine kinase inhibitor (TKI, n=10) regimens in the combined Harvard
1185 + IMDC retrospective cohort. **d**, Percentage of tRCC patients showing a response to either immune
1186 checkpoint inhibitor (ICI-based) or tyrosine kinase inhibitor (sunitinib and pazopanib) in the combined IMDC
1187 and Harvard retrospective cohorts. **e**, PD-L1 protein expression on infiltrating immune cells (PD-L1 $\geq$ 1%) in
1188 tRCC (n=15) and ccRCC (n=797) in the IMmotion151 cohort. **f**, Quantification of percentage of CD8⁺PD1⁺ T-
1189 cells (left), percentage of CD8⁺PD1⁺TIM3⁺LAG3⁺ T cells (middle), and percentage of CD8⁺PD1⁺TIM3⁺LAG3⁺ T
1190 cells (right) in tRCC (n=11), ccRCC (n=11), and adjacent normal tissue (from ccRCC cases, n= 10) analyzed by
1191 multiparametric immunofluorescence.

1192

1193 SUPPLEMENTARY TABLE LEGENDS

1194 **Supplementary Table 1:** List of samples in the NGS datasets included in the analysis.

1195 **Supplementary Table 2:** List and legend of functional domains used in the annotation of *MIT/TFE* and partners
1196 genes in **Figures 3d-e**.

1197 **Supplementary Table 3:** RSEM expected counts (**Supplementary Table 3a**) and transcript-per-million (TPM;
1198 **Supplementary Table 3b**) derived from the RNA-sequencing of the cell lines in the *in vitro* experiment
1199 represented in **Figure 4a**.

1200 **Supplementary Table 4:** List of genes that are in the *TFE3*-fusion-specific transcriptional signature developed in
1201 **Figure 4a**.

1202 **Supplementary Table 5:** Sample-level MAF (**Supplementary Table 5a**) and gene-level copy number
1203 (**Supplementary Table 5b**) data for the OncoPanel cohort.

1204 **Supplementary Table 6:** Sample-level data for the multiparametric immunofluorescence cohort.

1205