Development and comparison of Loop-Mediated Isothermal Amplification with Quantitative PCR for the specific detection of *Saprolegnia* spp.

Satyaki Ghosh^{1, #}, David L. Straus², Christopher Good³ and Vipaporn Phuntumart^{1, *}

¹ Department of Biological Sciences, Bowling Green State University, Bowling Green, Ohio, United States of America

² United States Department of Agriculture, Agricultural Research Service, Harry K. Dupree-Stuttgart National Aquaculture Research Center, Stuttgart, Arkansas, United States of America

³ The Conservation Fund's Freshwater Institute, Shepherdstown, West Virginia, United States of America

*Corresponding author

E-mail: vphuntu@bgsu.edu

Current Address: AgriPlex Genomics, 11000 Cedar Ave., Suite 250, Cleveland, Ohio 44106.

Abstract

Saprolegniasis is an important disease in freshwater aquaculture, and is associated with oomycete pathogens in the genus *Saprolegnia*. Early detection of significant levels of *Saprolegnia* spp. pathogens would allow informed decisions for treatment which could significantly reduce losses. This study is the first to report the development of loop-mediated isothermal amplification (LAMP) for the detection of *Saprolegnia* spp. and compares it with quantitative PCR (qPCR). The developed protocols targeted the internal transcribed spacer (ITS) region of ribosomal DNA and the cytochrome C oxidase subunit 1 (CoxI) gene and was shown to be specific only to *Saprolegnia* genus. This LAMP method can detect as low as 10 fg of *S. salmonis* DNA while the qPCR method has a detection limit of 2 pg of *S. salmonis* DNA, indicating the superior sensitivity of LAMP compared to qPCR. When applied to detect the pathogen in water samples, both methods could detect the pathogen when only one zoospore of *Saprolegnia* was present. We propose LAMP as a quick (about 20-60 minutes) and sensitive molecular diagnostic tool for the detection of *Saprolegnia* spp. suitable for on-site applications.

Keywords: *Saprolegnia*, pathogen, diagnostic procedure, LAMP, qPCR

Introduction

Oomycetes represent a diverse group of eukaryotic pathogens that can infect a wide range of hosts [1, 2, 3]. Pathogens in the *Saprolegnia* genus can infect and kill crustaceans and fish in freshwater, especially in aquaculture where fish are raised at high densities [4]. The most widely studied members of this genus include *S. diclina*, *S. ferax* and *S. parasitica* [5].

Saprolegniasis has been reported in channel catfish (*Ictalurus punctatus*), rainbow trout (*Oncorhynchus mykiss*), redear sunfish (*Lepomis microlophus*), Indian major carps (including *Labeo rohita*, *Catla catla*, *L. calbasu*) and Atlantic salmon (*Salmo salar*) [6, 7, 8]. The majority saprolegniasis occurrence has been reported in temperate environments and commonly found associated with salmonid farming. *Saprolegnia* can infect all stages of the fish life cycle including the egg stage [9, 10, 11]. In the U.S., economic losses due to saprolegniasis in commercial aquaculture is estimated to be approximately 40 million dollars [12, 13]. Almost 50% of the commercial losses to farmed channel catfish is attributed to saprolegniasis [14]. It is estimated that 10% of all hatched salmon succumb to saprolegniasis, which represents a major financial loss given that this industry accounts for approximately 30% of the global fish production for consumption [4, 14, 15, 16]. The incidence of saprolegniasis extends to Asian tropical aquaculture systems where over 80% of fish is produced through aquaculture [17]. Therefore, saprolegniasis is among the most impactful diseases in commercial aquaculture.

Conventional culture-based strategies coupled with conventional PCR-based approaches for pathogen identification usually take several days [18]. Currently, quantitative-PCR (qPCR) is one of the methods of choice for molecular detection and quantification of target organisms [19]. Detection of pathogens based on qPCR has been reported in several studies including

Trypanosoma cruzi, *Fusarium* spp., Japanese encephalitis virus and bacteria such as *Listeria monocytogenes*, *Francisella tularensis*, and *Mycobacterium avium* [20-23].

Loop-Mediated Isothermal Amplification (LAMP) has emerged as a novel tool for the sensitive and rapid detection of nucleic acids and is suitable for field applications. Since the first publication of LAMP [24], there has been a progressive increase in the number of publications based on this technique to diagnose and detect a variety of organisms including bacteria, viruses and eukaryotic pathogens indicating the success of LAMP as a reliable tool for detection [25, 26, 27]. The present focuses on development of a LAMP and a qPCR method for specific detection of *Saprolegnia* species, and compared these two tests based on sensitivity and specificity. These two molecular approaches targeted the internal transcribed spacer (ITS) region of ribosomal DNA and the mitochondrial cytochrome c oxidase subunit I (C0xI) gene. We first report the standard protocols for specific detection of the pathogens, followed by testing our methods under various scenarios for detection of *Saprolegnia* species including water samples, infected fish tissue, mycelium and zoospores. Our results show that both of the methods are reliable, specific and sensitive to *Saprolegnia*, and have the potential to be applied for the detection of other pathogens.

Materials and Methods

Isolation and maintenance of oomycete cultures

All *Saprolegnia* spp. cultures used in this study were grown and maintained on yeast peptone sucrose (YPS) agar (20 g/L D-glucose, 1 g/L KH₂PO₄, 0.5 mg/L MgSO₄, and 1 g/L yeast extract) supplemented with 68 µg/ml chloramphenicol and 68 µg/ml streptomycin. The cultures were

90 incubated at 24°C in the dark [28]. Unless otherwise noted, all laboratory grade chemicals used
91 in this study were purchased from Sigma Aldrich, St. Louis, MO.

92 *Saprolegnia* spp. isolates used in this study (Table 1) were isolated from various sources
93 including fish, eggs, and waters from commercial aquaculture ponds and from recirculating
94 aquaculture systems (RASs). For isolation of *Saprolegnia* spp. from water, water samples from
95 each source were collected and filtered through two layers of miracloth (Calbiochem, La Jolla,
96 CA). One ml of the filtrate was spread aseptically onto YPS media and incubated in the dark at
97 24°C. The resultant mycelial growths were subcultured for several generations to ensure the
98 purity of the culture.

99 **Table 1. *Saprolegnia* spp. isolates used in the present study.**

Organism/Sample ID	Isolation Source
<i>S. salmonis</i>	Sunshine bass eggs from Keo Fish Farm (Keo, AR).
3, 9, 6A, 11A, 1B	Pond water from J.M. Malone and Sons fish farm (Lonoke, AR)
F1, F2, JMM Shad 3, JMM Shad 72	American Shad (<i>A. sapidissima</i>) from ponds at J. M. Malone and Son fish farm (Lonoke, AR)
RAS1, RAS2, RAS3, RAS4, RAS5	RAS water from The Freshwater Institute (Shepherdstown, WV)

For isolation of *Saprolegnia* spp. from infected tissues, adult infected American shad (*Alosa sapidissima*) with lengths ranging between 48.3cm to 49.5cm and exhibiting visible symptoms of saprolegniasis were used for isolation of *Saprolegnia* spp following previously published protocols [29]. Briefly, areas of the fish exhibiting visible symptoms were dissected using sterilized scalpels; individual pieces were surface sterilized by treatment with 2.5% bleach for 45 s, followed by submersion in distilled water for 1 min, then in 70% ethanol for 45 s, and finally in distilled water for 1 min. These pieces were then aseptically placed onto YPS agar plates and incubated in the dark at 24°C and observed daily.

DNA isolation

Mycelial mats (90-100 mg) from 4-day old *Saprolegnia* cultures were collected and were crushed using a mortar and pestle in liquid nitrogen. DNA was isolated using the method of Zelaya-Molina et al. [30] with some modifications. Briefly, the crushed mycelia were treated with lysis buffer (10 mM Tris, 50 mM EDTA, 0.5% SDS, 0.5% Tween-20, and 0.5% Triton X-100), 20 mg/ml Proteinase K and 100 mg/ml RNase A. The tubes were incubated at 37°C for 30 min in a shaking incubator at 150 rpm, vortexed for 30 s and then incubated at 55°C for 30 min with a gentle inversion at 10-min intervals. Next, an equal volume of 25:24:1 phenol-chloroform-isoamyl alcohol (VWR, Radnor, PA) was added and the tubes were vortexed for 30 s followed by centrifugation at 10,000g for 10 min. The supernatant was transferred into a new tube, an equal volume of 24:1 chloroform-isoamyl alcohol was added, vortexed for 30 s and centrifuged at 10,000g for 10 min. The supernatant was transferred into a new tube, DNA was precipitated by adding an equal volume of isopropanol (VWR, Radnor, PA) and incubated at 20°C for 15 min. The tubes were then centrifuged at 10,000g for 10 min and the supernatant was discarded. The resultant pellet was washed with 70% ethanol (Pharmco-AAPER, Shelbyville,

KY), dried and resuspended in molecular grade water. The isolated DNA was quantified spectrophotometrically using a Nanodrop 2000c (Thermo Scientific, Wilmington, DE). Assessment of the quality of the isolated DNA was performed by electrophoresis on 0.8% agarose gels stained with SYBRTM Green I Nucleic Acid Gel stain (Invitrogen, Carlsbad, CA) and visualized using a UV transilluminator (Ultra-Lum Inc, Claremont, CA).

Sequence analysis and phylogenetic analysis

Identification of *Saprolegnia* spp. isolates was performed by PCR of the internal transcribed spacer (ITS) region and a portion of the ribosomal large subunit [16, 31, 32] and of cytochrome C oxidase subunit 1 (CoxI) gene [33]. Amplification products were analyzed by electrophoresis and were purified using the QIAquick PCR purification kit (Qiagen, Germantown, MD) according to the manufacturer's protocol. The purified amplicons were sequenced at the University of Chicago Comprehensive Cancer Centre, DNA sequencing and Genotyping Facility (Chicago, IL). The resultant sequences were subjected to bioinformatic analysis via BLAST [34] and compared to sequences on both the NCBI [35] and the FungiDB [36] databases.

Multiple sequence alignments were generated using Clustal Omega [37]. The maximum-likelihood phylogenetic trees of CoxI and ITS markers were constructed using Mega7 software [38] based on the Tamura-Nei Model [39] with 1,000 bootstrap replicates and default parameters. Reference ITS and CoxI sequences were obtained from NCBI GenBank database; *Phytophthora sojae* voucher P6497 and *Aphanomyces euteiches* voucher BR694 were used as the outgroups for both the trees. Concatenated phylogenetic trees of ITS and CoxI regions were constructed using SequenceMatrix version 1.8 (<http://www.ggvaidya.com/taxondna/>) [40] with 1,000 bootstrap replicates and default parameters.

qPCR for specific detection of *Saprolegnia* spp.

Primers specific to the ITS region and the CoxI gene of the *Saprolegnia* genus were designed using Primer3 software with manual modifications (<https://bioinfo.ut.ee/primer3-0.4.0/>) [41]. Quantitative PCR reactions were carried out using the iTaq™ Universal SYBR® Green Supermix (Bio-Rad, Hercules, CA) according to the manufacturer's instructions in a CFX Connect Real-Time PCR Detection system (Bio-Rad, Hercules, CA). To generate the standard-curve for absolute quantification, reactions were carried out using a 10-fold dilution series of *S. salmonis* DNA (ranging from 2 ng/μl to 20 ng/μl, measured spectrophotometrically). To assess the specificity of the developed qPCR, 20 ng/μl of DNA purified from all the *Saprolegnia* isolates described in Table 1 were used with both the ITS and CoxI primers. The reactions were carried out in triplicate.

LAMP for specific detection of *Saprolegnia* spp.

To design primers for LAMP against the conserved ITS region specific for *Saprolegnia* spp., the ITS sequences for *S. parasitica* CBS 223.65 (Accession No. - PRJNA280969), *S. diclina* VS20 (Accession No. - PRJNA255245), *S. ferax* (Accession No. - PRJNA12392), *S. delica* clone 130 (Accession No. - JX212906.1) and *S. salmonis* from Argentina (Accession No. - EU551153.1) were retrieved from the NCBI sequence database [35] and aligned using Clustal Omega [37]. Inner and outer primers were generated based on this alignment, using the Primer Explorer v5 (<https://primerexplorer.jp/e/>, Eiken Chemical Co. Ltd., Tokyo, Japan) tool, with default parameters.

To optimize the LAMP reaction, each LAMP reaction (25 μl) contained 20 ng/μl of *S. salmonis* gDNA, 1.4 mM dNTPs, 0.32 U/ μl Bst 3.0 (New England Biolabs, Ipswich, MA), 8 mM MgSO₄,

1.6 μ M inner primers, 0.2 μ M outer primers, 0.4 μ M loop primers and molecular grade water (Invitrogen, Carlsbad, CA). The reactions were incubated at 65°C for 60 min, followed by inactivation of the enzyme to terminate the reaction at 80°C for 5 min. No template controls (NTC) using molecular grade water served as negative controls. To each reaction, 1 μ l of 1,000X SYBR Green I (AMRESCO Inc., Solon, OH) was added. Visually, a green color formation was indicative of a positive reaction whereas a golden-brown color indicated a negative reaction. The amplification products were also verified visually by 2% agarose gel electrophoresis.

Comparison of qPCR and LAMP

A sensitivity assay was performed using a 10-fold dilution of purified *S. salmonis* DNA at 100 ng, 10 ng, 1 ng, 100 pg, 10 pg, 1 pg, 100 fg, 10 fg and 1 fg. Protocols for qPCR and LAMP assay were as described above. For detection of zoospores, one and ten zoospores were used in each reaction for both LAMP and qPCR. The reactions were performed in triplicate.

Results

Pathogen Identification and phylogenetic analysis

Fifteen cultures of the oomycete were isolated from water samples as well as from infected fish tissues (Table 1). To identify the isolates, PCR was performed using both the oomycete-specific CoxI primers and universal ITS primers. The PCR products were visualized on 0.8% agarose gel and were purified, sequenced, and analyzed by BLAST [34] with an E-value cutoff of 0. All of the isolates were identified as *Saprolegnia*. To further validate our results and to track their genetic diversity, phylogenetic trees were constructed. The maximum-likelihood tree of the ITS sequences showed that all the isolated species grouped within the *Saprolegnia* genus, in agreement with BLAST results (Fig 1). The *S. salmonis* isolate grouped closely with *S.*

parasitica with a bootstrap value of 68%. For cultures isolated from the ponds of J.M. Malone and Sons fish farm (Lonoke, AR), isolates 6A and 11A were closest to *S. diclina* with a bootstrap value of 90%. Isolate 1B was grouped with *S. parasitica* with a bootstrap value of 98%. Isolate 9 grouped with *S. ferax* with 80% bootstrap support. Isolate 3 was taxonomically closest to *S. diclina* with a bootstrap value of 63%. For cultures isolated from infected fish (J.M. Malone and Sons fish farm, Lonoke, AR), isolates F1 and F2 grouped together with *S. parasitica* with 67% bootstrap support. The JMM Shad 3 and JMM Shad 72 also grouped together and appeared to have *S. parasitica* as the closest phylogenetic neighbor with 68% bootstrap support. Isolates RAS1, RAS2 and RAS3 grouped closest to *S. parasitica*, while RAS4 grouped closest to *S. ferax* with 80% bootstrap support; RAS5 had the closest phylogenetic relationship with *S. monoica* with 89% bootstrap support. A second maximum-likelihood tree based on the CoxI sequences showed the same phylogenetic groupings with the ITS based tree with slightly different bootstrap values (Fig 2). To increase discriminatory power and the robustness for identification of species within the genus, the sequences of both ITS and CoxI were used to constitute the concatenated phylogenetic tree. Fig 3 showed the same phylogenetic groupings with both the ITS- and CoxI-based trees with slightly different bootstrap values. Overall, the results of the three phylogenetic trees confirmed that the isolates in Table 1 belonged to the *Saprolegnia* genus.

qPCR for specific detection

The qPCR reaction for CoXI gene using qCOX R1 (5'CTGAAGGACCGAGTGHGCTTG 3') and qCOX F1 (5'GGDGCTCCWGATAGGCTTTNCC 3') was optimized by gradient qPCR. Results showed that annealing temperature at 59°C gave the best specificity and efficiency. We then generated a standard curve using a 10-fold dilution of purified gDNA from *S. salmonis* (20 ng/μl to 0.2 pg/μl). The C_q values of these standards were plotted against the logarithm of their

concentrations. The assay efficiency was calculated to be at 91.4% using the equation $y = (-3.546)x + 54.138$ with the slope values of -3.546 and R^2 values of 0.990 (Fig 4). The melt curve analysis gave rise to a single distinct peak indicating that the qPCR product was a pure, single amplicon and was specific to *S. salmonis*.

Development of LAMP

Primers for the specific detection of *Saprolegnia* genus were designed targeting a conserved 201 bp section of the ITS region (Fig 5A). The inner and outer primers were generated based on sequence alignment using Clustal Omega alignment followed by the Primer Explorer v5 (<https://primerexplorer.jp/e/>, Eiken Chemical Co. Ltd., Tokyo, Japan) tool. The outer and inner primer sets with the highest dG value for dimerization and with acceptable room for generating loop primers, were chosen to ensure that the free energies and the 5' end of F1c and B1c, and the 3' end of F2 and B2, were less than or equal to -4.0 kcal/mol (Fig 5B, Table 2).

Table 2. LAMP primers used in the present study.

Primer	Abbreviation	Sequence
Forward Inner Primer	FIP	5' CACTTACATGAGAAATCTCCGAA-TAGCCGAAGAACGCTTTGGAAGC 3'
Backward Inner Primer	BIP	5' AATTCAGTGAGTCATCTAAATA-AACATACTCCCAGGACTAACCCGC 3'
Forward Outer Primer	FwdOut	5' GGTAATGGTGTGGTTTTTTGTGG 3'

Backward Outer Primer	RevOut	5' TGAAAGAAGTTTGTGTTG 3'
Loop Forward Primer	LF	5' GCGACGGGAACACCGT 3'
Loop Backward Primer	LB	5' AGTGCAATATGCGTT 3'

226

227 For optimization of the LAMP reaction, the first factor that was considered was the Mg^{2+}
228 concentration. We tested the reactions using 2 ng of DNA/reaction with eight different
229 concentrations of Mg^{2+} ranging from 2 mM-10 mM Mg^{2+} . The experiments were performed in
230 triplicate and repeated three times. Our results showed that concentration of Mg^{2+} at and above 4
231 mM showed positive reactions. Interestingly, in no template controls, when concentrations of
232 Mg^{2+} was at 5 mM and above, the colors of the reactions showed false positives (Fig 6).
233 Therefore, to avoid false positive reaction, the optimal concentration of Mg^{2+} at 4mM was
234 chosen. Optimization of the other factors including various temperatures, reaction times and
235 primer ratios did not have any impact on the integrity of the reaction. Hence, the optimal
236 condition for the LAMP reaction was selected as 4mM Mg^{2+} , 1:8 outer to inner primer ratio and
237 a reaction temperature of 65°C.

238 To establish the sensitivity of the LAMP reaction, the experiments were carried out with a 10-
239 fold dilution series of purified *S. salmomonis* DNA ranging from 1 fg to 100 ng. Results showed
240 that LAMP successfully detected as low as 10 fg of *S. salmomonis* DNA (Fig 7). This detection
241 sensitivity was higher than that of qPCR which was at 2 pg (Fig 4). Further testing to determine
242 whether our method can be used to directly detect zoospore from water samples without DNA

purification, we subjected one or ten zoospores of *S. salmonis* to LAMP and qPCR methods. Ten ng of *S. salmonis* genomic DNA was used as the positive control and NTC was used as negative controls. The results showed that both the qPCR and LAMP procedures successfully detected one zoospore in the water samples (Fig 8), indicating that our developed methods were highly specific and highly sensitive.

The specificity of the LAMP assay was tested using purified DNA from different isolates of *Saprolegnia* spp., infected fish tissues other non-*Saprolegnia* oomycetes (*A. astaci* and *P. sojiae*), a fungus (*Fusarium* spp.), *Daphnia magna* and a bacterium (*Escherichia coli*). Results showed that the LAMP reaction was specific only to the *Saprolegnia* genus (Table 3), demonstrating that our developed LAMP method was specific to pathogens in the genus *Saprolegnia* and is highly sensitive.

Table 3. Specificity of qPCR and LAMP.

Species	Isolation Source	qPCR result	LAMP result
<i>S. salmonis</i> (+ control)	Pond water from J.M. Malone and Sons fish farm (Lonoke, AR)	+	+
<i>S. ferax</i>	Infected crayfish tissue (Bowling Green, OH)	+	+
<i>S. parasitica</i>	RAS water from The Freshwater Institute (Shepherdstown, WV)	+	+
<i>S. diclina</i>	Pond water from J.M. Malone and Sons fish farm (Lonoke, AR)	+	+

<i>S. delica</i>	Pond water from J.M. Malone and Sons fish farm (Lonoke, AR)	+	+
<i>A. astaci</i>	Infected crayfish tissue (Bowling Green, OH)	-	-
<i>P. sojae</i>	Infected soybean (Bowling Green, OH)	-	-
<i>Fusarium</i> spp.	Water from cichlid aquaria (Bowling Green, OH)	-	-
<i>D. magna</i>	Water from BGSU Greenhouse reservoir (Bowling Green, OH)	-	-
<i>E. coli</i>	TOP10 (Invitrogen, CA)	-	-

Discussion

The pathogens in the genus *Saprolegnia* cause significant economic losses in aquaculture; therefore, accurate identification and early detection of pathologically relevant levels is critical. Morphological identification is challenging and less reliable compared to molecular studies. To ensure reliable identification of oomycetes, a combination of molecular markers based on ITS regions along with CoxI and/or CoxII are recommended [33]. The present study develops a standard protocol for identification of pathogens in the genus *Saprolegnia* using both ITS and CoxI. To determine genetic relationships among *Saprolegnia* spp. used in this study, phylogenetic trees with a single locus of ITS or CoxI marker as well as a concatenated phylogeny were constructed. These three phylogenetic trees confirmed that the isolates were related and belonged to the *Saprolegnia* genus. Due to low bootstrap values at some nodes, additional markers are required to conclusively determine the phylogenetic positioning of the isolates within the *Saprolegnia* genus. For instance, in all three phylogenetic trees, *S. salmonis*

clustered with *S. parasitica* but bootstrap values were below 68%. Multilocus phylogenetic analysis should, in theory, increase phylogenetic resolution and improve the analysis. Such multi-gene analyses has been conducted by Göker et al. [42] using α -tubulin, NADH1 and LSU rDNA sequences for several oomycetes. Phylogenetic resolution of the *Phytophthora* genus was accomplished in this manner by Martin et al. [43]. Richter and Rósselló-Mora [44] suggested that measurement of average nucleotide identity between two genomes could yield more accurate results of species resolution. Finally, it has been suggested that sampling more individuals per species can help resolve incongruences amongst single gene trees [45]. An interesting observation in our phylogenetic studies was that the variation at the species level for the *Saprolegnia* isolates from RAS, even though the water samples from these systems was collected from the same tank at the same time. However, similar species level spatio-temporal variations have been reported for *Phytophthora*, *Pythium* and *Phytopythium* species in nursery irrigation systems [46].

In this study, we were able to isolate and purify *Saprolegnia* from environmental sources (water samples and infected tissues), using a similar method for the isolation of *Saprolegnia* spp. from infected rainbow trout [8], infected crayfish [47], salmon hatcheries [11] and amphibians [48]. Current protocols for detecting and identifying *Saprolegnia* from environmental sources involve lab culture-based approaches followed by PCR-based molecular identification. Another method of oomycete detection which has been used with some success is baiting [49] using rice seeds [50], hemp seeds [28] and sesame seeds [51], followed by PCR for accurate identification. Finally, immunological detection of *Saprolegnia* spp. using monoclonal antibodies has also been reported [15].

This study is the first to develop a rapid and sensitive molecular technique for the detection of *Saprolegnia* spp. using the novel LAMP technique. Previously, LAMP has been employed for the detection of other oomycetes such as *Plasmopara viticola* [52], *P. sojae* [53] and *Pythium* spp. [54]. As an on-site detection tool, LAMP has several advantages compared to conventional PCR and qPCR. It has been applied successfully for on-site detections of the bacteria responsible for foot rot in sheep [55].

A major advantage of LAMP is that it is highly sensitive and does not require major equipment. Our developed LAMP assay successfully detected all 15 isolates in the present study and offered positive results with as low as 10 fg DNA and with only one zoospore of *S. salmonis*. It is noteworthy that our method can directly detect an organism from a sample without DNA purification, which refers to a one-step LAMP. This method has been reported in several studies such as detection of *E. coli*, *Mycobacterium smegmatis* [56] and *Mycoplasma ovipneumoniae* [57]. Similarly, detection by LAMP at femtogram level has been reported in numerous studies such as acute viral necrobiosis virus in scallop [58], classical swine flu virus [59] and *Enterococcus hirae* [60]. When our developed LAMP assay from this study was tested on several members of the oomycetes, it was found to be specific for all 15 *Saprolegnia* isolates used in this study. The sensitivity and specificity of LAMP makes it a suitable candidate for the detection of target organisms from environmental samples, especially when present in extremely low amounts. Although false positives are one of the major drawbacks associated with LAMP, much of the false positive amplification can be eliminated by using six primers and careful optimization of the reaction [61]. We did not experience any false positives in our experiments. While both LAMP and qPCR have their own advantages and drawbacks depending on the circumstance, they can both be used to specifically detect the members of the genus *Saprolegnia*.

The LAMP method is inexpensive and more rapid and sensitive compared to qPCR; therefore, we propose the use of LAMP for sensitive detection of pathologically relevant levels of *Saprolegnia* for on-site applications. However, to quantify *Saprolegnia* loads, qPCR is more useful, especially when the epidemic threshold of the disease is known. Incorporation of the LAMP method and qPCR into real-time LAMP would provide the best outcome for detection and quantification. This would enable monitoring the dynamics of *Saprolegnia* spp. in the water, and at what points their level becomes an issue. Real-time LAMP has been reported for the detection and quantification of *Enterocytozoon hepatopenaei* [62] and *Ustilago maydis* [63] and future research should investigate its development use for to detect *Saprolegnia* and other oomycete pathogens.

Acknowledgements

This research was funded by Ohio Sea Grant (10000960) and Bowling Green State University. We thank Gayathri Beligala for her contribution to the experiments. We thank Dr. Paul F. Morris for his technical suggestions regarding experimental design and for his comments on the manuscript. Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by Bowling Green State University, The Conservation Fund's Freshwater Institute or the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

References

1. Densmore CL, Green DE. Diseases of amphibians. ILAR Journal. 2007;48(3):235-54.
doi: 10.1093/ilar.48.3.235.
2. Ruthig GR. Water molds of the genera *Saprolegnia* and *Leptolegnia* are pathogenic to the North American frogs *Rana catesbeiana* and *Pseudacris crucifer*, respectively. Diseases of Aquatic Organisms. 2009;84(3):173-8. doi: 10.3354/dao02042.
3. Banfield MJ, Kamoun S. Hooked and Cooked: A Fish Killer Genome Exposed. PLOS Genetics. 2013;9(6):e1003590. doi: 10.1371/journal.pgen.1003590.
4. van West P. *Saprolegnia parasitica*, an oomycete pathogen with a fishy appetite: new challenges for an old problem. Mycologist. 2006;20(3):99-104. doi: <https://doi.org/10.1016/j.mycol.2006.06.004>.
5. Diéguez-Uribeondo J, Cerenius L, Söderhäll K. Repeated zoospore emergence in *Saprolegnia parasitica*. 1994;98(7):810-5. doi: 10.1016/s0953-7562(09)81060-5.
6. Bly JE, Lawson LA, Szalai AJ, Clem LW. Environmental factors affecting outbreaks of winter saprolegniosis in channel catfish, *Ictalurus punctatus* (Rafinesque). Journal of Fish Diseases. 1993;16(6):541-9. doi: <https://doi.org/10.1111/j.1365-2761.1993.tb00890.x>.
7. Das SK, Murmu K, Das A, Shakuntala I, Das RK, Ngachan SV, Majhi SK. Studies on the identification and control of pathogen *Saprolegnia* in selected Indian major carp fingerlings at mid hill altitude. J Environ Biol. 2012;33(3):545-9. Epub 2012/10/04. PubMed PMID: 23029901.
8. Shin S, Kulatunga DCM, Dananjaya SHS, Nikapitiya C, Lee J, De Zoysa M. *Saprolegnia parasitica* isolated from rainbow trout in Korea: Characterization, anti-*Saprolegnia* activity and host pathogen interaction in zebrafish disease model. Mycobiology.

2017;45(4):297-311. Epub 2017/12/31. doi: 10.5941/MYCO.2017.45.4.297. PubMed

PMID: 29371797.

9. Straus DL, Mitchell AJ, Carter RR, Steeby JA. Optimizing copper sulfate treatments for fungus control on channel catfish eggs. Journal of Aquatic Animal Health. 2009;21(2):91-7. doi: <https://doi.org/10.1577/H07-057.1>.

10. Straus DL, Farmer BD, Ledbetter CK, Beck BH, Williams RS, Clark ML, Freeze MT. Use of copper sulfate to control egg saprolegniasis at a commercial sunshine bass hatchery. North American Journal of Aquaculture. 2016;78(3):243-50. Doi: <https://doi.org/10.1080/15222055.2016.1146183>.

11. Thoen E, Evensen Ø, Skaar I. Factors influencing *Saprolegnia* spp. spore numbers in Norwegian salmon hatcheries. Journal of Fish Diseases. 2016;39(6):657-65. doi: 10.1111/jfd.12392.

12. Torto-Alalibo T, Tian M, Gajendran K, Waugh ME, van West P, Kamoun S. Expressed sequence tags from the oomycete fish pathogen *Saprolegnia parasitica* reveal putative virulence factors. BMC Microbiol. 2005;5:46-. doi: 10.1186/1471-2180-5-46. PubMed PMID: 16076392.

13. van den Berg AH, McLaggan D, Diéguez-Urbeondo J, van West P. The impact of the water moulds *Saprolegnia diclina* and *Saprolegnia parasitica* on natural ecosystems and the aquaculture industry. Fungal Biology Reviews. 2013;27(2):33-42. doi: <https://doi.org/10.1016/j.fbr.2013.05.001>.

14. Phillips AJ, Anderson VL, Robertson EJ, Secombes CJ, van West P. New insights into animal pathogenic oomycetes. Trends in Microbiology. 2008;16(1):13-9. Doi: <https://doi.org/10.1016/j.tim.2007.10.013>.

- 379 **15.** Fregeneda-Grandes JM, Rodríguez-Cadenas F, Aller-Gancedo JM. Fungi isolated from
380 cultured eggs, alevins and broodfish of brown trout in a hatchery affected by
381 saprolegniosis. *Journal of Fish Biology*. 2007;71(2):510-8. doi: 10.1111/j.1095-
382 8649.2007.01510.x.
- 383 **16.** Sandoval-Sierra JV, Martín MP, Diéguez-Uribeondo J. Species identification in the
384 genus *Saprolegnia* (Oomycetes): Defining DNA-based molecular operational taxonomic
385 units. *Fungal Biology*. 2014;118(7):559-78. doi:
386 <https://doi.org/10.1016/j.funbio.2013.10.005>.
- 387 **17.** Karunasagar I, Karunasagar I, Otta SK. disease problems affecting fish in tropical
388 environments. *Journal of Applied Aquaculture*. 2003;13(3-4):231-49. doi:
389 10.1300/J028v13n03_03.
- 390 **18.** Leung MH, Bryson K, Freystatter K, Pichon B, Edwards G, Charalambous BM, Gillespie
391 SH. Sequotyping: serotyping *Streptococcus pneumoniae* by a single PCR sequencing
392 strategy. *J Clin Microbiol*. 2012;50(7):2419-27. Epub 2012/05/02. doi:
393 10.1128/JCM.06384-11. PubMed PMID: 22553238.
- 394 **19.** Kralik P, Ricchi M. A Basic Guide to Real Time PCR in Microbial Diagnostics:
395 Definitions, parameters, and everything. *Front Microbiol*. 2017;8:108-. doi:
396 10.3389/fmicb.2017.00108. PubMed PMID: 28210243.
- 397 **20.** de Souza Godoi PA, Piechnik CA, de Oliveira AC, Sfeir MZ, de Souza EM, Rogez H,
398 Soccol VH. qPCR for the detection of foodborne *Trypanosoma cruzi*. *Parasitology*
399 International. 2017;66(5):563-6. doi: <https://doi.org/10.1016/j.parint.2017.06.001>.

21. Zitnick-Anderson K, Simons K, Pasche JS. Detection and qPCR quantification of seven *Fusarium* species associated with the root rot complex in field pea. Canadian Journal of Plant Pathology. 2018;40(2):261-71. doi: 10.1080/07060661.2018.1429494.
22. Bharucha T, Sengvilaipaseuth O, Vongsouvath M, Vongsouvath M, Davong V, Panyanouvong P, Piorkowski G, Garson JA, Newton PN, de Lamballerie X, Dubot-Pères A. Development of an improved RT-qPCR Assay for detection of Japanese encephalitis virus (JEV) RNA including a systematic review and comprehensive comparison with published methods. PloS one. 2018;13(3):e0194412-e. doi: 10.1371/journal.pone.0194412. PubMed PMID: 29570739.
23. Ricchi M, Bertasio C, Boniotti MB, Vicari N, Russo S, Tilola M, Bellotti MA, Bertasi B. Comparison among the quantification of bacterial pathogens by qPCR, dPCR, and cultural methods. Front Microbiol. 2017;8:1174. Epub 2017/07/14. doi: 10.3389/fmicb.2017.01174. PubMed PMID: 28702010; PubMed Central PMCID: PMC5487435.
24. Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, Hase T. Loop-mediated isothermal amplification of DNA. Nucleic Acids Res. 2000;28(12):E63-E. doi: 10.1093/nar/28.12.e63. PubMed PMID: 10871386.
25. Seki M, Kilgore PE, Kim EJ, Ohnishi M, Hayakawa S, Kim DW. Loop-mediated isothermal amplification methods for diagnosis of bacterial meningitis. Front Pediatr. 2018;6:57-. doi: 10.3389/fped.2018.00057. PubMed PMID: 29594087.
26. Niihara H, Kohno K, Tobita R, Ishikawa N, Morita E. Rapid diagnosis of herpes zoster by loop-mediated isothermal amplification in a pregnant woman showing folliculitis-like

eruption without vesicles. The Journal of Dermatology. 2017;44(7):e174-e5. doi:

10.1111/1346-8138.13861.

27. Nakayama T, Yamazaki T, Yo A, Tone K, Mahdi Alshahni M, Fujisaki R, Makimura K.

Detection of fungi from an indoor environment using loop-mediated isothermal

amplification (LAMP) method. Biocontrol Science. 2017;22(2):97-104. doi:

10.4265/bio.22.97.

28. Stueland S, Hatai K, Skaar I. Morphological and physiological characteristics of

Saprolegnia spp. strains pathogenic to Atlantic salmon, *Salmo salar* L. Journal of Fish

Diseases. 2005;28(8):445-53. doi: 10.1111/j.1365-2761.2005.00635.x.

29. Wolf K, Quimby MC. 5 Fish cell and tissue culture. In: Hoar WS, Randall DJ, editors.

Fish Physiology. 3: Academic Press; 1969. p. 253-305.

30. Zelaya-Molina LX, Ortega MA, Dorrance AE. Easy and efficient protocol for oomycete

DNA extraction suitable for population genetic analysis. Biotechnol Lett.

2011;33(4):715-20. Epub 2010/11/24. doi: 10.1007/s10529-010-0478-3. PubMed PMID:

21107649.

31. Bakkeren G, Kronstad JW, Lévesque CA. Comparison of AFLP fingerprints and ITS

sequences as phylogenetic markers in Ustilaginomycetes. Mycologia. 2000;92(3):510-21.

doi: 10.1080/00275514.2000.12061187.

32. Bala K, Robideau GP, Désaulniers N, de Cock AWAM, Lévesque CA. Taxonomy, DNA

barcoding and phylogeny of three new species of *Pythium* from Canada. Persoonia.

2010;25:22-31. Epub 2010/07/30. doi: 10.3767/003158510X524754. PubMed PMID:

21339964.

- 444 **33.** Robideau GP, De Cock AWAM, Coffey MD, Voglmayr H, Brouwer H, Bala K, Chitty
445 DW, Désaulniers N, Eggertson QA, Gachon CM, Hu CH, Küpper FC, Rintoul TL,
446 Sarhan E, Verstappen EC, Zhang Y, Bonants PJ, Ristaino JB, Lévesque CA. DNA
447 barcoding of oomycetes with cytochrome c oxidase subunit I and internal transcribed
448 spacer. *Molecular Ecology Resources*. 2011;11(6):1002-11. doi: 10.1111/j.1755-
449 0998.2011.03041.x.
- 450 **34.** Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search
451 tool. *Journal of Molecular Biology*. 1990;215(3):403-10. doi:
452 [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
- 453 **35.** NCBI Resource Coordinators. Database resources of the National Center for
454 Biotechnology Information. *Nucleic Acids Res*. 2016 Jan 4;44(D1):D7-19. doi:
455 10.1093/nar/gkv1290.
- 456 **36.** Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D,
457 Warrenfeltz S, Aurrecoechea C, Stoeckert CJ Jr, Kissinger JC, Roos DS, Hertz-Fowler C.
458 FungiDB: An integrated bioinformatic resource for fungi and oomycetes. *J Fungi (Basel)*.
459 2018;4(1):39. doi: 10.3390/jof4010039. PubMed PMID: 30152809.
- 460 **37.** Madeira F, Park Ym, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN,
461 Potter SC, Finn RD, Lopez R. The EMBL-EBI search and sequence analysis tools APIs
462 in 2019. *Nucleic Acids Res*. 2019;47(W1):W636-W41. doi: 10.1093/nar/gkz268.
- 463 **38.** Kumar S, Stecher G, Tamura K. MEGA7: Molecular evolutionary genetics analysis
464 Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*. 2016;33(7):1870-4.
465 doi: 10.1093/molbev/msw054.

39. Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*. 1993;10(3):512-26. doi: 10.1093/oxfordjournals.molbev.a040023.
40. Vaidya G, Lohman DJ, Meier R. SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics*. 2011;27(2):171-80. doi: 10.1111/j.1096-0031.2010.00329.x.
41. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. In: Misener S, Krawetz SA, editors. *Bioinformatics Methods and Protocols*. Totowa, NJ: Humana Press; 1999. p. 365-86.
42. Göker M, Voglmayr H, Riethmüller A, Oberwinkler F. How do obligate parasites evolve? A multi-gene phylogenetic analysis of downy mildews. *Fungal Genetics and Biology*. 2007;44(2):105-22. doi: <https://doi.org/10.1016/j.fgb.2006.07.005>.
43. Martin FN, Abad ZG, Balci Y, Ivors K. Identification and detection of *Phytophthora*: reviewing our progress, identifying our needs. *Plant Disease*. 2012;96(8):1080-103. doi: 10.1094/pdis-12-11-1036-fe. PubMed PMID: 30727075.
44. Richter M, Rosselló-Móra R. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A*. 2009;106(45):19126-31. Epub 2009/10/23. doi: 10.1073/pnas.0906412106. PubMed PMID: 19855009.
45. Maddison WP, Knowles LL. Inferring phylogeny despite incomplete lineage sorting. *Systematic Biology*. 2006;55(1):21-30. doi: 10.1080/10635150500354928.
46. Redekar NR, Eberhart JL, Parke JL. Diversity of *Phytophthora*, *Pythium*, and *Phytopythium* Species in Recycled Irrigation Water in a Container Nursery. *Phytobiomes Journal*. 2019;3(1):31-45. doi: 10.1094/pbiomes-10-18-0043-r.

47. Söderhäll K, Dick MW, Clark G, Fürst M, Constantinescu O. Isolation of *Saprolegnia parasitica* from the crayfish *Astacus leptodactylus*. Aquaculture. 1991;92:121-5. doi: [https://doi.org/10.1016/0044-8486\(91\)90014-X](https://doi.org/10.1016/0044-8486(91)90014-X).
48. Fernández-Benítez MJ, Ortiz-Santaliestra ME, Lizana M, Diéguez-Urbeondo J. *Saprolegnia diclina*: another species responsible for the emergent disease ‘Saprolegnia infections’ in amphibians. FEMS Microbiology Letters. 2008;279(1):23-9. doi: 10.1111/j.1574-6968.2007.01002.x.
49. Nechwatal J, Wielgoss A, Mendgen K. Diversity, host, and habitat specificity of oomycete communities in declining reed stands (*Phragmites australis*) of a large freshwater lake. Mycological Research. 2008;112(6):689-96. doi: <https://doi.org/10.1016/j.mycres.2007.11.015>.
50. Sandoval-Sierra JV, Diéguez-Urbeondo J. A Comprehensive Protocol for Improving the Description of Saprolegniales (Oomycota): Two Practical Examples (*Saprolegnia aenigmatica* sp. nov. and *Saprolegnia racemosa* sp. nov.). PloS ONE. 2015;10(7):e0132999. doi: 10.1371/journal.pone.0132999.
51. Ali SE, Thoen E, Vrålstad T, Kristensen R, Evensen Ø, Skaar I. Development and reproduction of *Saprolegnia* species in biofilms. Veterinary Microbiology. 2013;163(1):133-41. doi: <https://doi.org/10.1016/j.vetmic.2012.12.012>.
52. Kong X, Qin W, Huang X, Kong F, Schoen CD, Feng J, Wang Z, Zhang H. Development and application of loop-mediated isothermal amplification (LAMP) for detection of *Plasmopara viticola*. Scientific reports. 2016;6:28935-. doi: 10.1038/srep28935. PubMed PMID: 27363943.

- 511 **53.** Dai T-T, Lu C-C, Lu J, Dong S, Ye W, Wang Y, Zheng X. Development of a loop-
512 mediated isothermal amplification assay for detection of *Phytophthora sojae*. FEMS
513 Microbiology Letters. 2012;334(1):27-34. doi: 10.1111/j.1574-6968.2012.02619.x.
- 514 **54.** Feng W, Hieno A, Kusunoki M, Suga H, Kageyama K. LAMP Detection of four plant-
515 pathogenic oomycetes and its application in lettuce fields. Plant Disease.
516 2019;103(2):298-307. Doi: 10.1094/pdis-05-18-0858-re. PubMed PMID: 30608214.
- 517 **55.** Best N, Rawlin G, Suter R, Rodoni B, Beddoe T. Optimization of a loop-mediated
518 isothermal amplification (LAMP) assay for in-field detection of *Dichelobacter nodosus*
519 with aprV2 (VDN LAMP) in Victorian sheep flocks. Frontiers in Veterinary Science.
520 2019;6(67). doi: 10.3389/fvets.2019.00067.
- 521 **56.** Naik P, Jaitpal S, Shetty P, Paul D. An integrated one-step assay combining thermal lysis
522 and loop-mediated isothermal DNA amplification (LAMP) in 30 min from *E. coli* and *M.*
523 *smegmatis* cells on a paper substrate. bioRxiv. 2019:594374. doi: 10.1101/594374.
- 524 **57.** Zhang J, Cao J, Zhu M, Xu M, Shi F. Loop-mediated isothermal amplification-lateral-
525 flow dipstick (LAMP-LFD) to detect *Mycoplasma ovipneumoniae*. World J Microbiol
526 Biotechnol. 2019;35(2):31-. doi: 10.1007/s11274-019-2601-5. PubMed PMID:
527 30701329.
- 528 **58.** Zhang H, Zeng L, Fan Y, Zhou Y, Xu J, Ma J. A loop-mediated isothermal amplification
529 assay for rapid detection of cyprinid herpesvirus 2 in Gibel carp (*Carassius auratus*
530 *gibelio*). ScientificWorld Journal. 2014;2014:716413-. doi: 10.1155/2014/716413.
531 PubMed PMID: 24574914.
- 532 **59.** Parida M, Shukla J, Sharma S, Ranghia Santhosh S, Ravi V, Mani R, Thomas M, Khare
533 S, Rai A, Kant Ratho R, Pujari S, Mishra B, Lakshmana Rao PV. Development and

evaluation of reverse transcription loop-mediated isothermal amplification assay for rapid and real-time detection of the swine-origin influenza A H1N1 virus. J Mol Diagn. 2011;13(1):100-7. Epub 2011/01/14. doi: 10.1016/j.jmoldx.2010.11.003. PubMed PMID: 21227400; PubMed Central PMCID: PMC3069812.

60. Dolka B, Cisek AA, Szeleszczuk P. The application of the loop-mediated isothermal amplification (LAMP) method for diagnosing *Enterococcus hirae*-associated endocarditis outbreaks in chickens. BMC Microbiol. 2019;19(1):48-. doi: 10.1186/s12866-019-1420-z. PubMed PMID: 30791887.

61. Hardinge P, Murray JAH. Reduced false positives and improved reporting of loop-mediated isothermal amplification using quenched fluorescent primers. Scientific Reports. 2019;9(1):7400. Doi: 10.1038/s41598-019-43817-z.

62. Cai S-X, Kong F-D, Xu S-F, Yao C-L. Real-time loop-mediated isothermal amplification for rapid detection of *Enterocytozoon hepatopenaei*. PeerJ. 2018;6:e5993-e. doi: 10.7717/peerj.5993. PubMed PMID: 30533299.

63. Cao Y, Wang L, Duan L, Li J, Ma J, Xie S, Shi L, Li H. Development of a real-time fluorescence loop-mediated isothermal amplification assay for rapid and quantitative detection of *Ustilago maydis*. Scientific reports. 2017;7(1):13394-. doi: 10.1038/s41598-017-13881-4. PubMed PMID: 29042629.

Figure Legends:

Figure 1. Maximum-likelihood phylogenetic tree showing evolutionary relationships of 53

***Saprolegnia* isolates based on the ITS region.** Isolates examined in this study are shown in boldface. Bootstrap values $\geq 50\%$ (1000 replicates) are given at the branchpoints. The scale bar indicates the number of substitutions per site. *P. sojae* voucher P6497 and *A. euteiches* BR694 were used as outgroups.

Figure 2. Maximum-likelihood phylogenetic tree showing evolutionary relationships of 53

***Saprolegnia* isolates based on the CoxI gene.** Isolates examined in this study are shown in boldface. Bootstrap values $\geq 50\%$ (1000 replicates) are given at the branchpoints. The scale bar indicates the number of substitutions per site. *P. sojae* voucher P6497 and *A. euteiches* BR694 were used as outgroups.

Figure 3. Concatenation phylogenetic tree based on the sequences of ITS and CoxI. Isolates

examined in this study are shown in boldface. Bootstrap values $\geq 50\%$ (1000 replicates) are given at the branchpoints. The scale bar indicates the number of substitutions per site. *P. sojae* voucher P6497 and *A. euteiches* BR694 were used as outgroups.

Figure 4. Standard curve of CoxI gene. Ten-fold dilutions of *S. salmonis* DNA ranging from 1

fg to 100 ng were amplified in triplicate. The iTaq™ Universal SYBR® Green Supermix and qCOXF1+R1 primers were used. The reactions were repeated three times.

576

577 **Figure 5. Lamp primer design.** A. Schematic representation of the ITS region, showing the
578 positioning of the LAMP primers used in the present study. Scale bar represents 100bp.
579 FwdOut=Forward Outer Primer; RevOut=Backward Outer Primer; F2+F1c=Forward Inner
580 Primer; B2c+B1=Backward Inner Primer; LF=Loop Forward Primer; LB=Loop Backward
581 Primer. B. Positioning and orientation of LAMP primers developed in the present study.
582 Representative ITS sequences from *Saprolegnia* and non-*Saprolegnia* oomycetes were aligned
583 and the LAMP primers were positioned on the alignment manually. The LAMP primers
584 amplified the ITS1 regions of the target sequences.

585

586 **Figure 6. Optimization of Mg²⁺ concentration for LAMP.** The concentrations of Mg²⁺
587 ranging from 2 mM-10 mM were tested. Top panel represents positive controls using 2ng/μl *S.*
588 *salmonis* gDNA. The bottom panel, No Template Control (NTC) represents negative controls.
589 Color change to green indicates positive reaction, while golden brown indicates negative
590 reaction. The experiments were performed in triplicate and repeated three times.

591

592 **Figure 7: The sensitivity of LAMP.** Ten-fold dilution *S. salmonis* DNA ranging from 1 fg to
593 100 ng were used in the presence of 1000X SYBR Green I. Color changes from golden brown to
594 green indicates a positive reaction. The concentrations of template DNA are indicated, NTC= no
595 template control, represents a negative control. The reactions were performed in triplicate and
596 were repeated three times.

597

598 **Figure 8. Direct detection of zoospores.** A. qPCR and B. LAMP. One and ten zoospores were
599 used in each reaction, *S. salmonis* DNA (10 ng/μl) was used as the positive control while no
600 template control (NTC) was used as the negative control. Each sample was performed in
601 triplicate and was repeated three times.

602

Figure 1

bioRxiv preprint doi: <https://doi.org/10.1101/2021.04.15.439949>; this version posted April 15, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

Figure 2

bioRxiv preprint doi: <https://doi.org/10.1101/2021.04.15.439949>; this version posted April 15, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

Figure 3

bioRxiv preprint doi: <https://doi.org/10.1101/2021.04.15.439949>; this version posted April 15, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

Figure 4

bioRxiv preprint doi: <https://doi.org/10.1101/2021.04.15.439949>; this version posted April 15, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a U.S. Government work. It is not subject to copyright under 17 USC 105 and is also made available under a CC-BY 4.0 International license.

Figure 5

bioRxiv preprint doi: <https://doi.org/10.1101/2021.04.15.439949>; this version posted April 15, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

Figure 6

bioRxiv preprint doi: <https://doi.org/10.1101/2021.04.15.439949>; this version posted April 15, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. This article is a US Government work. It is not subject to copyright under 17 USC 105 and is also made available for use under a CC0 license.

Figure 7

Figure 8