Contrasting processing tomato cultivars unlink yield and pollen viability under heat stress

Running title: The response of processing tomato cultivars to heat stress

4

5 Golan Miller¹, Avital Beery¹, Prashant Kumar Singh^{1,2}, Fengde Wang^{1,3}, Rotem
6 Zelingher^{1,4}, Etel Motenko¹, Michal Lieberman-Lazarovich¹

7 ¹ Institute of plant sciences, Agricultural Research organization – Volcani Center,
8 Rishon LeZion, Israel.

9 ² Department of Biotechnology, Mizoram University (A Central University),
10 Pachhunga University College Campus, Aizawl-796005, Mizoram, India.

11 ³ Institute of vegetables and flowers, Shandong Academy of Agricultural Sciences,
12 Jinan, China.

13 ⁴ Université Paris-Saclay, INRAE, AgroParisTech, Economie Publique, 78850,
14 Thiverval-Grignon, France

15

16

Abstract:

The occurring climate change is causing temperature increment in crop production areas worldwide, generating conditions of heat stress that negatively affect crop productivity. Tomato (*Solanum lycopersicum*), a major vegetable crop, is highly susceptible to conditions of heat stress. When tomato plants are exposed to ambient day/night temperatures that exceed 32°C/20°C respectively during the reproductive phase, fruit set and fruit weight are reduced, leading to a significant decrease in yield. Processing tomato cultivars are cultivated in open fields, where environmental conditions are not controlled, therefore plants are exposed to multiple abiotic stresses, including heat stress. Understanding the physiological response of modern processing tomato cultivars to heat stress may facilitate the development of thermotolerant cultivars. Here, we compared two tomato processing cultivars, H4107 and H9780, that we found to be constantly differing in yield performance. Using field and temperature-controlled greenhouse experiments, we show that the observed difference in yield is attributed to the occurrence of heat stress conditions. In addition, fruit-set and seed production were significantly improved in the thermotolerant cultivar H4107, compared with H9780. Despite the general acceptance of pollen viability as a measure of thermotolerance, there was no difference in the percentage of viable pollen between H4107 and H9780 under either of the conditions tested. Therefore, processing tomato cultivars may present a particular case, in which other factors are central for heat stress tolerance. Our results also demonstrate the value of combining controlled with uncontrolled experimental settings, in order to identify heat stress related responses and facilitate the development of thermotolerant processing tomato cultivars.

Keywords: Heat Stress, Tomato, Yield, Processing cultivars, Pollen viability

Introduction

Plant physiology and development are prominently affected by changes in ambient temperatures. With the current global climate change, temperatures are gradually shifting and temperature extremes occur more frequently. Predictions of the effect of temperature increment on major crops yield show that each degree-Celsius increase in global mean temperature would cause yield reduction by 3.1-7.4% on average (Zhao et al. 2017). Recent IPCC reports estimate global warming is likely to reach a 1.5°C increase in average surface temperature between 2030 and 2052 if it continues to

increase at the current rate, and reach a 2–4°C increase by the end of the twenty-first century (IPCC, 2018), thus challenging crop productivity and food security. High temperature is a major abiotic stress that disturbs basic molecular processes, such as protein folding, photosynthesis and assimilates metabolism (Bokszczanin et al. 2013). These effects cause morphological and physiological changes, negatively affecting plant growth and development (Wahid et al. 2007; Bitu and Gerats 2013). Yield reduction due to heat stress was documented in various crops such as cereals (wheat, rice, barley, sorghum and maize), pulses (chickpea) and oil yielding crops (mustard, canola) fruits and vegetables (potato, eggplant, cabbage, cauliflower, lettuce, onion, cucumber, musk melon, watermelon and pumpkin) (Hasanuzzaman et al. 2013). When heat stress occurs during the reproductive phase of plant development, the observed consequences include morphological alterations of anthers, style elongation, bud abscission and reduced fruit number, size and seed set. The development of pollen is considered the most heat-sensitive stage (Lohani et al. 2020) as it was shown to be more sensitive than both the sporophyte and female gametophyte tissues (Peet et al. 1998; Young et al. 2004; Wang et al. 2019). Heat stress disrupts of meiotic cell division, abnormal pollen morphology and size, and reduced grain number, viability, and germination capacity (Endo et al. 2009; M. M. Peet et al. 1998; Djanaguiraman et al. 2013; Giorno et al. 2013; Pressman et al. 2002; Firon et al. 2006; Begcy et al. 2019; Prasad et al. 2006). Specifically, pollen viability is considered a central element for heat stress tolerance as high temperatures were shown to impair pollen viability in numerous crop species such as wheat (Begcy et al. 2018), rice (Jagadish et al. 2007), sorghum (Djanaguiraman et al. 2018), soybean (Djanaguiraman et al. 2013), and tomato (Firon et al. 2006), leading to male sterility and reduced fruit/grain production.

Tomato (*Solanum lycopersicum*), an important vegetable crop worldwide, cultivated in a wide range of agro-climatic regions, is very sensitive to heat stress. The tomato fruit set is optimal when the average day and night temperatures range between 21°C - 29°C and 18°C - 21°C, respectively (Pelzer 2008). Prolonged stress of day temperatures exceeding 32°C with night temperature above 20°C cause reduced fruit set, fruit weight, total yield and seed production (El Ahmadi and Stevens 1979; Peet et al. 1998, Sato 2000; Firon 2006). In tomato, pollen heat stress related damage, exhibited by morphological alterations and reduced pollen viability and germination rates, was observed after short episodes of high temperatures at 40°C, or after chronic exposure to milder heat stress of 31-32°C/25-28°C day/night for several months (Firon et al.

2006; Iwahori 1966; Giorno et al. 2013). The decrease in pollen viability and/or germination was shown to cause a significant decrease in fruit set (Iwahori 1965; Rudich et al. 1977; Abdul-Baki 1992; Sato et al. 2000), therefore pollen viability was used as a screening approach to identify heat stress tolerant tomato genotypes. Consequently, several tomato genotypes were identified, that maintain a higher level of pollen viability under heat stress conditions (Dane et al. 1991; Paupière et al. 2017; Driedonks et al. 2018). Pollen viability is therefore often used as a measure of thermotolerance, establishing the correlation between pollen viability and fruit (Pressman et al. 2002; Xu et al. 2017; Pham et al. 2020; Rutley et al. 2021; Firon et al. 2006).

In contrast to the wealth of data demonstrating the correlation between pollen heat stress damage and fruit set, examples of heat stress tolerance/sensitivity not correlated with pollen viability are very scarce. To the best of our knowledge, only two such cases were described. Gonzalo *et al.* (2020) performed a population screen of introgression lines from the wild species *Solanum pimpinellifolium* for reproductive traits under controlled heat stress conditions, and no correlation was found between pollen viability and fruit set (Gonzalo et al. 2020). In a more recent study, Ayenan et al (2021) screened a collection of 42 cultivated and wild tomato genotypes with good yield components under long term mild heat stress and did not find association between the proportion of viable pollen and fruit set percentage (Ayenan et al., 2021). In this paper, we present yet another example for heat stress tolerance that is not correlated with pollen viability, in a processing cultivar of tomato.

Tomato processing cultivars are used by the food industry to produce tomato paste and sauces, canned crushed, diced, or peeled tomatoes and various juices and soups. For these purposes, breeding companies developed cultivars suited for mechanical harvesting and canning processes. These cultivars are characterized by a determinate growth habit, synchronized fruit set and firm flesh (Hanna 1971; Gould et al. 1992), unlike the indeterminate fresh market cultivars, grown primarily in greenhouses or other covered facilities. Processing tomato plants are cultivated only in open fields, where heat stress conditions are prevalent. Particularly in the Mediterranean basin, including the major tomato producers Italy and Spain, the growing season starts in March–April, when the probability of high temperatures during the sensitive reproductive stage is very high (<http://www.wptc.to>). However, information regarding the response of processing cultivars to heat stress is very limited.

Here, we characterized the heat stress response of two processing tomato cultivars, which are usually grown in open field conditions therefore exposed to a combination of stress factors, including heat stress, during the reproductive stage. We show that the constant difference in yield between these cultivars is attributed to high temperature conditions. In order to gain information specifically for the response to heat-stress, the same cultivars were tested in a controlled greenhouse, under heat stress and control conditions in a parallel setup. This setup allows the identification of specific heat stress related traits, which is not possible under the uncontrolled, multi-stress field conditions. Our results demonstrate a clear difference in performance under heat stress, which is, unexpectedly, not related to pollen viability.

Materials and methods

Plant material and growth conditions

Two tomato (*Solanum lycopersicum*) commercial processing cultivars H4107 and H9780 (Green Seeds Ltd.), were grown during 2018 in three different experimental fields, in different locations as follows: 1. ‘Upper Galilee’ site, at the Northern part of Israel (33°10'50.6"N latitude 35°34'49.6"E longitude; Field size 100 plants), 2. ‘Eden’ site (32°27'58.2"N latitude 35°29'12.2"E longitude; Field size 80 plants) and 3. ‘Volcani’ site at a central region of Israel (31°59'34.6"N latitude 34°49'01.8"E longitude; field size 40 plants). The two cultivars were grown in a completely randomized design in 3-5 replicas (plots). Seeds were sown in germination trays and transplanted in open fields after three weeks. Mature plants were maintained under standard horticultural practices. During the whole growing period climatic data were recorded using the weather stations ‘Khavat Eden’, ‘Beit Dagan’ and ‘Mop Tzafon’ located in Eden, Volcani and Upper Galilee fields, respectively. In addition, the two cultivars were grown in climate controlled greenhouses at the Naan site of Evogene LTD company. In this controlled experiment, four plants from each cultivar were grown under moderate chronic heat stress (MCHS) conditions (32°C-22°C day-night, starting at flowering) and control conditions (25°C-18°C day-night), in a randomized setup, identical between the two rooms. The seeds were sown in germination trays and transplanted into 10L pots filled with soil 21 days after sowing.

Reproductive traits evaluation

Fruit set and fruit production were evaluated in all three experimental fields and in the controlled experiment. Fruit production (FW – fruit weight) was evaluated by weighing total red-ripe fruits per repeat (plot or plant in the field or controlled experiments, respectively). Fruit set ratio (FS) was evaluated from 10 randomly selected inflorescences from each plot in the field experiments. In the controlled experiment, FS was evaluated from three randomly selected inflorescences in 4 different plants (a total of 12 inflorescences per cultivar). Seed number per fruit (SN) was examined by seeds extraction using three fruits from five plants (Volcani field) or three fruits from five plots (Upper Galilee field). In the controlled experiment, 5-25 fruits from all four plants were sampled. Seeds were extracted using the sulfuric acid method; the locular gel containing the seeds was extracted and soaked in 2% sulfuric acid solution. After 3 hours, the seeds were transferred into a net bag and rinsed under tap water. Seeds were then thoroughly dried in the open air for few days. Seed number was calculated using the weighing method: a small portion was manually counted and weighed, and then the total amount of seeds was estimated by weighing.

Pollen viability analysis

For pollen viability analysis, flowers at anthesis were collected in the morning (7 to 10 am). In total, three flowers per plant were collected and three plants were used per cultivar. Each anther was cut into two pieces and put in a 1.5 mL tube filled with 0.5mL germination solution [1 mM KNO₃, 3 mM Ca (NO₃)₂·4H₂O, 0.8 mM MgSO₄·7 H₂O, 1.6 mM H₃BO₃; (Pressman et al. 2002)], followed by 20 µl of Alexander dye. The Alexander dye consisted of 20 ml of ethanol, 20 mg of malachite green, 50 ml of distilled water, 40 ml of glycerol, 100mg of Acid fuchsin, 2 gr Phenol, and 2 ml of Lactic acid for a 100 ml solution (Alexander 1980). Samples were observed under Leica DMLB epi-fluorescence microscope (Germany) using BF filter, magnified by 10-20. Three fields containing representative pollen pattern were captured with DS-Fi1 digital camera using NIS-Elements BR3.0 software (Nikon). Viable (purple) and non-viable (blue-green) pollen grains were counted manually in ImageJ version 1.43 software using the 'Cell counter' plugin (Schneider et al. 2012).

Statistical analysis

One-way ANOVA was employed to identify significant differences ($p < 0.05$) between the cultivars for each trait. When ANOVA identified significant differences among genotypes, we used the student t-test method as an exact test for all differences

between means. These conservative procedures limited the probability of rejecting a true null hypothesis to the desired ($p < 0.05$) level. All statistical analyses were performed using JMP Version 3.2.2 (SAS Institute, Inc., Cary, N.C.).

Results

Consistent difference in yield between H4107 and H9780 across multiple years and locations.

Following a survey of processing tomato field-testing data from 15 years (between 2005 and 2019) across 17 different locations (Table S1), we detected a consistent difference between two cultivars, i.e., H4107 and H9780. While the yield of H4107 was always above the test average, the yield of H9780 was always lower than the test average (Table S1). When we compared the results of specific years and locations where both cultivars were tested simultaneously, the average yield was 12.4 and 10.8 kg/m^2 for H4107 and H9780, respectively, providing a significant difference (Figure 1a, b). We aimed to understand the source of this difference in order to promote breeding efforts for high yield in field-grown processing tomato. Since the field environment imposes various stresses to the plants, and tomato being particularly sensitive to elevated temperatures, we set to test the possibility that the high temperature conditions usually prevalent in those regions are causing the difference in yield.

The difference in yield between H4107 and H9780 is associated with high temperature conditions.

To test whether the observed difference in yield between H4107 and H9780 is due to their differential response to high temperature, we set field experiments in two locations that are routinely used for processing tomato cultivation, however, differing by their environmental conditions. The 'Upper Galilee' field is located in a region that is characterized by hot days and cooler nights during the processing tomato season (May-July), whereas the "Eden" field is located in the Jordan Valley which is characterized by high day and night temperatures, and high humidity. For this reason, planting in "Eden" starts earlier (February until May), to avoid extreme heat stress and yield losses. In addition, we set a small experimental field at the Volcani Center, located in a more temperate region. Overall, we tested the plants under field conditions in three different environments. Environmental data were obtained for each field from a local meteorological station, enabling recording temperature every 3 hours, hence we

calculated day and night average and maximum temperatures. Considering that tomato plants experience heat stress when day temperature exceeds 32°C and night temperature exceeds 20°C, our analysis shows that heat stress conditions were indeed prevalent in all three locations, though with some differences (Figure 2a-d). In the Eden field, due to the early planting, heat stress conditions developed around 50 days after flowering. Nonetheless, day and night maximal temperatures surpassed threshold values already 5 days after flowering, generating heat stress conditions throughout the entire reproductive period. In the Upper Galilee field, daily average temperatures were around 32°C, reaching a maximum of approximately 35°C in most days, including three incidences of above 40°C. Night temperatures in the Upper Galilee field were higher than 20°C throughout the period, reaching a maximum of over 30°C on several occasions, presenting more severe heat stress than in the Eden field. Lower temperatures were observed in the Volcani field, where the daily average was usually under 32°C, with four exceptional heat waves. Night temperatures were still high averaging around 25°C throughout the tested period, thus the plants in the Volcani field also experienced heat stress conditions (Figure 2a-d). Under the above-described conditions, we found that the yield of H4107 was significantly higher than that of H9780 in all fields (Figure 2e), in agreement with our analysis of multiple years and locations data (Figure 1). While H4107 produced 9.0, 6.9, and 11.0Kg fruit/m² in Upper Galilee, Volcani, and Eden, respectively, H9780 produced 5.1, 3.3, and 8.0Kg fruit/m² in the same respective fields. Moreover, yield levels in both cultivars were higher in Eden than in the Upper Galilee and Volcani fields that experienced a more substantial heat stress, suggesting that yield levels are indeed affected by the high temperatures in these locations. The reproductive difference between H4107 and H9780 was further demonstrated by testing fruit set ratio and seed production in the Upper Galilee and Volcani fields (Figure 3). In these locations, H4107 reached 28% and 35% fruit set, respectively, while H9780 had 17% fruit set in both locations (Figure 3a). Similarly, H4107 produced a higher number of seeds per fruit versus H9780, reaching 244 and 96, respectively, in the Upper Galilee field. In the Volcani field, H4107 had on average 61 seeds per fruit, and H9780 produced only 21 seeds per fruit on average, maintaining a significant difference (Figure 3b).

In order to validate the effect of heat stress on the productivity of H4107 and H9780, we set a controlled experiment in which the same cultivars were grown under

either MCHS (32°C/22°C day/night), or control conditions (25°C/18°C day/night) in separate rooms. At the beginning of the experiment, both rooms were maintained under control conditions. Once plants started to flower, MCHS was initiated in one room while the other room was kept at control conditions throughout the rest of the plants growth (Figure 4a). Fruit set rate and seed production were analyzed under both conditions. We found no significant difference between H4107 and H9780 in both parameters measured (i.e. 64-68% fruit set and 52-92 seeds per fruit) under control conditions. However, under MCHS conditions, H4107 performed better than H9780, as the fruit set was 36% versus 19% in H9780. Seed number per fruit was 71 and 23 for H4107 and H9780, respectively (Figure 4b-c). Markedly, fruit set ratios were very similar between field and controlled heat stress for both cultivars, supporting the occurrence of heat stress conditions in the field experiments. Importantly, these results confirm that the observed difference in yield and other reproductive traits under open field conditions are due to high temperatures, and suggest that H4107 is more tolerant than H9780 to heat stress.

The difference in heat tolerance between H4107 and H9780 is not related to pollen viability.

Since pollen viability is widely recognized as a main parameter determining plant heat stress tolerance (Dane et al. 1991; Paupière et al. 2017; Driedonks et al. 2018), we aimed to test whether the heat stress tolerance of H4107 can be at least partially explained by higher degree of pollen viability under heat stress conditions. To address that, we analyzed pollen viability percentage in field and controlled conditions. In the Upper Galilee field, we found no significant difference between H4107 and H9780, as both showed 60-70% viable pollen out of total pollen grains (Figure 5a). Pollen viability was lower in the Volcani field (30-45%), yet still similar between the cultivars (Figure 5b). In the controlled experiment, pollen viability reached 90-100%, even under MCHS conditions, and again, similarly between H4107 and H9780. Interestingly, the same levels were found under control conditions (Figure 5c), meaning that pollen viability was not affected by heat stress in these cultivars and is not linked with the heat stress tolerance of H4107. Our results also suggest that the low rates of pollen viability in field conditions is not due to the high temperatures, but rather to another environmental factor.

Discussion

Current literature on processing tomatoes in general and on their response to heat stress in particular is very limited. We identified a consistent difference in yield between H4107 and H9780 across multiple years and locations. This difference is manifested by higher fruit set rate and total fruit weight of H4107. We found this difference to be associated with the response to heat stress, meaning that H4107 is more heat stress tolerant than H9780, presenting better reproductive performance in terms of fruit set and seed production under high temperature conditions. H4107 was bred and adapted for humid and arid environments by the Heinz company (https://d36rz30b5p7lzd.cloudfront.net/372/studio/assets/v1611911409263_1054604699/2021%20HeinzSeed%20International%20Brochure.pdf), but heat stress tolerance was not reported so far. Interestingly, the heat stress tolerance we observed was not correlated with better pollen viability, implying that other factors mediate the tolerance in this system. In one of the earliest studies on heat stress response in tomato, Levy *et al.* (1978) showed that the characters contributing to low fruit set under heat stress were bud drop and style exertion which were more pronounced in susceptible cultivars. Actually, no fruit set was ever observed when the style protruded out of the antheridial cone (Levy *et al.* 1978). Fruit setting was correlated with bud abscission and style elongation under field conditions as well (Singh *et al.* 2015; Kugblenu *et al.* 2013). Considering this aspect, we tested bud abscission and style elongation ratios in field and greenhouse but no significant difference was found between H4107 and H9780 (data not shown). Alternatively, ovule development and post-pollination interactions were also demonstrated to negatively influence fruit set, by applying pollen from control condition flowers onto freshly open flowers grown under heat stress conditions (Peet *et al.*, 1997; Xu *et al.*, 2017).

Ayenan *et. al* (2021) showed recently that in some tomato genotypes grown in the greenhouse, pollen viability was not correlated with fruit set and yield (Ayenan *et al.*, 2021). On the same hand, our results suggest that while pollen viability is a valid trait demonstrating heat stress tolerance in various tomato genotypes, it may not be the case in open field processing cultivars. If this is due to their genetic structure or the complex environment they were bred in, or a combination of both, is yet to be determined following a comprehensive follow-up study.

Generally, in plant science research, field and greenhouse data are inconsistent, explained by the big difference in environmental conditions between the two experimental systems. We found that fruit set is highly similar between the controlled experiment (36% and 19% for H4107 and H9780, respectively) and the field experiments (28-36% and 17% for H4107 and H9780, respectively). Thus, our results demonstrate consistency in regard to a complex trait (yield), suggesting that in our system, controlled greenhouse experiments are highly relevant for agricultural conditions, facilitating translating research from lab to practice. Moreover, our results demonstrate the importance of temperature-controlled experimental systems in isolating specific heat-stress related phenomena.

In order to address the challenge of maintaining crop productivity in areas of temperature increment, the development of thermo-tolerant cultivars is needed. To achieve that, a comprehensive understanding of the agronomical, physiological and molecular responses of crop plants to heat stress is vital (Berry and Bjorkman 1980; Brestic et al. 2018). In light of the research presented here, which demonstrates a unique feature of specific cultivars, emphasis should be put on local and relevant cultivars that may offer different attributes in terms of response to the environment.

Acknowledgements

(removed)

Authors declare no conflict of interests.

References

- Abdul-Baki, A A. 2019. "Determination of Pollen Viability in Tomatoes." *Journal of the American Society for Horticultural Science* 117 (3): 473–76. <https://doi.org/10.21273/jashs.117.3.473>.
- Abdul-Baki, A A, and J R Stommel. 1995. "Pollen Viability and Fruit Set of Tomato Genotypes under Optimum- and High-Temperature Regimes." *HortScience* 30 (1): 115–17. <https://doi.org/10.21273/hortsci.30.1.115>.
- Akhoundnejad, Yelderem, Yildiz H. Dasgan, and Senay Karabiyik. 2020. "Pollen Quality, Pollen Production and Yield of Some Tomato (*Solanum Lycopersicum*) Genotypes under High Temperature Stress in Eastern Mediterranean." *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 48 (2): 893–905. <https://doi.org/10.15835/nbha48211896>.

- 343 Alexander, M P. 1980. "A Versatile Stain for Pollen Fungi, Yeast and
344 Bacteria." *Stain Technology* 55 (1): 13–18.
- 345 Ayenan, Mathieu Anatole Tele, Agyemang Danquah, Peter Hanson, Isaac K
346 Asante, and Eric Y Danquah. 2021. "Identification of New Sources of Heat
347 Tolerance in Cultivated and Wild Tomatoes." *Euphytica* 217 (3): 1–16.
- 348 Begcy, Kevin, Tetyana Nosenko, Liang Zi Zhou, Lena Fragner, Wolfram
349 Weckwerth, and Thomas Dresselhaus. 2019. "Male Sterility in Maize after
350 Transient Heat Stress during the Tetrad Stage of Pollen Development." *Plant
351 Physiology* 181 (2): 683–700. <https://doi.org/10.1104/pp.19.00707>.
- 352 Begcy, Kevin, Anna Weigert, Andrew Ogolla Egesa, and Thomas
353 Dresselhaus. 2018. "Compared to Australian Cultivars, European Summer Wheat
354 (Triticum Aestivum) Overreacts When Moderate Heat Stress Is Applied at the
355 Pollen Development Stage." *Agronomy* 8 (7): 99.
356 <https://doi.org/10.3390/agronomy8070099>.
- 357 Berry, J, and O Bjorkman. 1980. "Photosynthetic Response and Adaptation
358 to Temperature in Higher Plants." *Annual Review of Plant Physiology* 31 (1):
359 491–543. <https://doi.org/10.1146/annurev.pp.31.060180.002423>.
- 360 Bitá, Craita E, and Tom Gerats. 2013. "Plant Tolerance to High Temperature
361 in a Changing Environment: Scientific Fundamentals and Production of Heat
362 Stress-Tolerant Crops." *FRONTIERS IN PLANT SCIENCE* 4.
363 <https://doi.org/10.3389/fpls.2013.00273>.
- 364 Bokszezanin, Kamila L, Sotirios Fragkostefanakis, and Solanaceae Pollen
365 Thermotolerance. 2013. "Perspectives on Deciphering Mechanisms Underlying
366 Plant Heat Stress Response and Thermotolerance." *FRONTIERS IN PLANT
367 SCIENCE* 4. <https://doi.org/10.3389/fpls.2013.00315>.
- 368 Brestic, Marian, Marek Zivcak, Pavol Hauptvogel, Svetlana Misheva,
369 Konstantina Kocheva, Xinghong Yang, Xiangnan Li, and Suleyman I.
370 Allakhverdiev. 2018. "Wheat Plant Selection for High Yields Entailed
371 Improvement of Leaf Anatomical and Biochemical Traits Including Tolerance to
372 Non-Optimal Temperature Conditions." *Photosynthesis Research* 136 (2): 245–
373 55. <https://doi.org/10.1007/s11120-018-0486-z>.
- 374 Buckle, Simon. 2009. "Mitigation of Climate Change." *Weather* 64 (6):
375 165–66. <https://doi.org/10.1002/wea.422>.
- 376 Dane, Fenny, A Gene Hunter, and Oyetete L Chambliss. 2019. "Fruit Set,
377 Pollen Fertility, and Combining Ability of Selected Tomato Genotypes under
378 High-Temperature Field Conditions." *Journal of the American Society for
379 Horticultural Science* 116 (5): 906–10. <https://doi.org/10.21273/jashs.116.5.906>.
- 380 Djanaguiraman, M., R. Perumal, S. V.K. Jagadish, I. A. Ciampitti, R. Welti,
381 and P. V.V. Prasad. 2018. "Sensitivity of Sorghum Pollen and Pistil to High-

382 Temperature Stress.” *Plant Cell and Environment* 41 (5): 1065–82.
383 <https://doi.org/10.1111/pce.13089>.

384 Djanaguiraman, M., P. V.V. Prasad, D. L. Boyle, and W. T. Schapaugh.
385 2013. “Soybean Pollen Anatomy, Viability and Pod Set under High Temperature
386 Stress.” *Journal of Agronomy and Crop Science* 199 (3): 171–77.
387 <https://doi.org/10.1111/jac.12005>.

388 Driedonks, Nicky, Mieke Wolters-Arts, Heidrun Huber, Gert-Jan de Boer,
389 Wim Vriezen, Celestina Mariani, and Ivo Rieu. 2018. “Exploring the Natural
390 Variation for Reproductive Thermotolerance in Wild Tomato Species.”
391 *Euphytica* 214 (4): 1–12.

392 Ahmadi, A Beshir El, and M Allen Stevens. 1979. “Reproductive Responses
393 of Heat-Tolerant Tomatoes to High Temperatures.” *Journal of the American*
394 *Society for Horticultural Science* 104: 686–91.

395 Endo, Makoto, Tohru Tsuchiya, Kazuki Hamada, Shingo Kawamura,
396 Kentaro Yano, Masahiro Ohshima, Atsushi Higashitani, Masao Watanabe, and
397 Makiko Kawagishi-Kobayashi. 2009. “High Temperatures Cause Male Sterility
398 in Rice Plants with Transcriptional Alterations during Pollen Development.”
399 *Plant and Cell Physiology* 50 (11): 1911–22. <https://doi.org/10.1093/pcp/pcp135>.

400 Firon, N., R. Shaked, M. M. Peet, D. M. Pharr, E. Zamski, K. Rosenfeld, L.
401 Althan, and E. Pressman. 2006. “Pollen Grains of Heat Tolerant Tomato
402 Cultivars Retain Higher Carbohydrate Concentration under Heat Stress
403 Conditions.” *Scientia Horticulturae* 109 (3): 212–17.
404 <https://doi.org/10.1016/j.scienta.2006.03.007>.

405 Giorno, Filomena, Mieke Wolters-Arts, Celestina Mariani, and Ivo Rieu.
406 2013. “Ensuring Reproduction at High Temperatures: The Heat Stress Response
407 during Anther and Pollen Development.” *Plants* 2 (3): 489–506.
408 <https://doi.org/10.3390/plants2030489>.

409 Gonzalo, Maria José, Yi Cheng Li, Kai Yi Chen, David Gil, Teresa
410 Montoro, Inmaculada Nájera, Carlos Baixauli, Antonio Granell, and Antonio
411 José Monforte. 2020. “Genetic Control of Reproductive Traits in Tomatoes
412 Under High Temperature.” *Frontiers in Plant Science* 11 (April): 1–15.
413 <https://doi.org/10.3389/fpls.2020.00326>.

414 Gould, Wilbur A, Wilbur A Gould, and others. 1992. “Tomato Production,
415 Processing & Technology.”

416 Grandillo, Silvana, Daniel Zamir, and Steven D. Tanksley. 1999. “Genetic
417 Improvement of Processing Tomatoes: A 20 Years Perspective.” *Euphytica* 110
418 (2): 85–97. <https://doi.org/10.1023/A:1003760015485>.

419 Hanna, G C. 1971. “Breeding Tomatoes for Mechanical Harvesting in
420 California.” *Genet, Agt* 25: 379–90.

- 421 Hasanuzzaman, Mirza, Kamrun Nahar, Md Mahabub Alam, Rajib
422 Roychowdhury, and Masayuki Fujita. 2013. "Physiological, Biochemical, and
423 Molecular Mechanisms of Heat Stress Tolerance in Plants." *International*
424 *Journal of Molecular Sciences* 14 (5): 9643–84.
425 <https://doi.org/10.3390/ijms14059643>.
- 426 Iwahori, Shuichi. 1965. "High Temperature Injuries in Tomato. IV.
427 Development of Normal Flower Buds and Morphological Abnormalities of
428 Flower Buds Treated with High Temperature." *Journal of the Japanese Society*
429 *for Horticultural Science* 34 (1): 33–41.
- 430 Iwahori, Shuichi. 1966. "High Temperature Injuries in Tomato." *Engei*
431 *Gakkai Zasshi* 35 (4): 379–86. <https://doi.org/10.2503/jjshs.35.379>.
- 432 Jagadish, S. V.K., P. Q. Craufurd, and T. R. Wheeler. 2007. "High
433 Temperature Stress and Spikelet Fertility in Rice (*Oryza Sativa* L.)." *Journal of*
434 *Experimental Botany* 58 (7): 1627–35. <https://doi.org/10.1093/jxb/erm003>.
- 435 Kugblenu, Yvonne O, Eric Oppong Danso, Kwadjo Ofori, Mathias N
436 Andersen, Stephen Abenney-Mickson, Edward B Sabi, Finn Plauborg, et al.
437 2013. "Screening Tomato Genotypes for Adaptation to High Temperature in
438 West Africa." *Acta Agriculturae Scandinavica, Section B-Soil & Plant Science*
439 63 (6): 516–22.
- 440 Levy, A., H. D. Rabinowitch, and N. Kedar. 1978. "Morphological and
441 Physiological Characters Affecting Flower Drop and Fruit Set of Tomatoes at
442 High Temperatures." *Euphytica* 27 (1): 211–18.
443 <https://doi.org/10.1007/BF00039137>.
- 444 Lohani, Neeta, Mohan B Singh, and Prem L Bhalla. 2020. "High
445 Temperature Susceptibility of Sexual Reproduction in Crop Plants." *Journal of*
446 *Experimental Botany*. Oxford University Press.
447 <https://doi.org/10.1093/jxb/erz426>.
- 448 Masson-Delmotte, V, P Zhai, H.-O. Pörtner, D Roberts, J Skea, P R Shukla,
449 A Pirani, et al. 2018. "IPCC, 2018: Summary for Policymakers. In: Global
450 Warming of 1.5°C. An IPCC Special Report on the Impacts of Global Warming
451 of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas
452 Emission Pathways, in the Context of Strengthening the Global." *IPCC Special*
453 *Report* 1 (3): 374–81.
- 454 Müller, Florian, and Ivo Rieu. 2016. "Acclimation to High Temperature
455 during Pollen Development." *Plant Reproduction* 29 (1–2): 107–18.
456 <https://doi.org/10.1007/s00497-016-0282-x>.
- 457 Parlevliet, J E. 1994. "Breeding for Abiotic Stress Tolerance." *Plant*
458 *Production on the Threshold of a New Century*. [https://doi.org/10.1007/978-94-](https://doi.org/10.1007/978-94-011-1158-4_27)
459 [011-1158-4_27](https://doi.org/10.1007/978-94-011-1158-4_27).

- 460 Paupière, Marine J., Pauline van Haperen, Ivo Rieu, Richard G.F. Visser,
461 Yury M. Tikunov, and Arnaud G. Bovy. 2017. "Screening for Pollen Tolerance
462 to High Temperatures in Tomato." *Euphytica* 213 (6).
463 <https://doi.org/10.1007/s10681-017-1927-z>.
- 464 Peet, M. M., S. Sato, and R. G. Gardner. 1998. "Comparing Heat Stress
465 Effects on Male-Fertile and Male-Sterile Tomatoes." *Plant, Cell and*
466 *Environment* 21 (2): 225–31. <https://doi.org/10.1046/j.1365-3040.1998.00281.x>.
- 467 Peet, Mary M, D H Willits, and R Gardner. 1997. "Response of Ovule
468 Development and Post-Pollen Production Processes in Male-Sterile Tomatoes to
469 Chronic, Sub-Acute High Temperature Stress." *Journal of Experimental Botany*
470 48 (1): 101–11.
- 471 Pelzer, Nancy L. 2008. "A Review of Tomato Plant Culture: In the Field,
472 Greenhouse, and Home Garden." *Journal of Agricultural & Food Information*.
473 Taylor & Francis. <https://doi.org/10.1080/10496500802286343>.
- 474 Pham, Dung, Ken Hoshikawa, Satoshi Fujita, Shoma Fukumoto, Tadayoshi
475 Hirai, Yoshihito Shinozaki, and Hiroshi Ezura. 2020. "A Tomato Heat-Tolerant
476 Mutant Shows Improved Pollen Fertility and Fruit-Setting under Long-Term
477 Ambient High Temperature." *Environmental and Experimental Botany* 178:
478 104150. <https://doi.org/10.1016/j.envexpbot.2020.104150>.
- 479 Prasad, P V.Vara, Kenneth J Boote, and L Hartwell Allen. 2006. "Adverse
480 High Temperature Effects on Pollen Viability, Seed-Set, Seed Yield and Harvest
481 Index of Grain-Sorghum [Sorghum Bicolor (L.) Moench] Are More Severe at
482 Elevated Carbon Dioxide Due to Higher Tissue Temperatures." *Agricultural and*
483 *Forest Meteorology* 139 (3–4): 237–51.
484 <https://doi.org/10.1016/j.agrformet.2006.07.003>.
- 485 Pressman, Etan, Mary M. Peet, and D. Mason Pharr. 2002. "The Effect of
486 Heat Stress on Tomato Pollen Characteristics Is Associated with Changes in
487 Carbohydrate Concentration in the Developing Anthers." *Annals of Botany* 90
488 (5): 631–36. <https://doi.org/10.1093/aob/mcf240>.
- 489 Rieu, Ivo, David Twell, and Nurit Firon. 2017. "Pollen Development at High
490 Temperature: From Acclimation to Collapse." *Plant Physiology* 173 (4): 1967–
491 76. <https://doi.org/10.1104/pp.16.01644>.
- 492 Rudich, J, E Zamski, and Yael Regev. 1977. "Genotypic Variation for
493 Sensitivity to High Temperature in the Tomato: Pollination and Fruit Set."
494 *Botanical Gazette* 138 (4): 448–52. <http://www.jstor.org/stable/2473878>.
- 495 Rutley, Nicolas, Golan Miller, Fengde Wang, Jeffrey F Harper, Gad Miller,
496 and Michal Lieberman Lazarovich. 2020. "Enhanced Reproductive
497 Thermotolerance Is Associated with Increased Accumulation of Flavonols in
498 Pollen of the Tomato High-Pigment 2 Mutant." *BioRxiv*.

499 Sato, S., M. M. Peet, and J. F. Thomas. 2000. "Physiological Factors Limit
500 Fruit Set of Tomato (*Lycopersicon Esculentum* Mill.) under Chronic, Mild Heat
501 Stress." *Plant, Cell and Environment* 23 (7): 719–26.
502 <https://doi.org/10.1046/j.1365-3040.2000.00589.x>.

503 Sato, S, M Kamiyama, T Iwata, N Makita, H Furukawa, and H Ikeda. 2006.
504 "Moderate Increase of Mean Daily Temperature Adversely Affects Fruit Set of
505 *Lycopersicon Esculentum* by Disrupting Specific Physiological Processes in
506 Male Reproductive Development." *Annals of Botany* 97 (5): 731–38.
507 <https://doi.org/10.1093/aob/mcl037>.

508 Schneider, Caroline A, Wayne S Rasband, and Kevin W Eliceiri. 2012.
509 "NIH Image to ImageJ: 25 Years of Image Analysis." *Nature Methods* 9 (7):
510 671–75.

511 Singh, Umesh, Pradeep Kumar Patel, Amit Kumar Singh, Vivek Tiwari,
512 Rajesh Kumar, N Rai, Anant Bahadur, Shailesh K Tiwari, Major Singh, and B
513 Singh. 2015. "Screening of Tomato Genotypes Underhigh Temperature Stress
514 for Reproductive Traits." *Vegetable Science* 42 (2): 52–55.

515 Wahid, Abdul, Saddia Gelani, M Ashraf, and Majid R Foolad. 2007. "Heat
516 Tolerance in Plants: An Overview." *Environmental and Experimental Botany* 61
517 (3): 199–223.

518 Wang, Yuanyuan, Hongbin Tao, Beijing Tian, Dechang Sheng, Chenchen
519 Xu, Heming Zhou, Shoubing Huang, and Pu Wang. 2019. "Flowering Dynamics,
520 Pollen, and Pistil Contribution to Grain Yield in Response to High Temperature
521 during Maize Flowering." *Environmental and Experimental Botany* 158: 80–88.
522 <https://doi.org/10.1016/j.envexpbot.2018.11.007>.

523 Xu, Jiemeng, Mieke Wolters-Arts, Celestina Mariani, Heidrun Huber, and
524 Ivo Rieu. 2017. "Heat Stress Affects Vegetative and Reproductive Performance
525 and Trait Correlations in Tomato (*Solanum Lycopersicum*)." *Euphytica* 213 (7):
526 156. <https://doi.org/10.1007/s10681-017-1949-6>.

527 Young, Lester W, Ron W Wilen, and Peta C Bonham-Smith. 2004. "High
528 Temperature Stress of Brassica Napus during Flowering Reduces Micro-and
529 Megagametophyte Fertility, Induces Fruit Abortion, and Disrupts Seed
530 Production." *Journal of Experimental Botany* 55 (396): 485–495.
531 <https://academic.oup.com/jxb/article-abstract/55/396/485/489044>.

532 Zhao, Chuang, Bing Liu, Shilong Piao, Xuhui Wang, David B Lobell, Yao
533 Huang, Mengtian Huang, et al. 2017. "Temperature Increase Reduces Global
534 Yields of Major Crops in Four Independent Estimates." *Proceedings of the
535 National Academy of Sciences of the United States of America* 114 (35): 9326–
536 31. <https://doi.org/10.1073/pnas.1701762114>.

537

538

Figure 1. Consistent difference in yield between H4107 and H9780 across years and locations. (A) Average yield of H4107 and H9780 in years and locations testing both cultivars. The test average obtained by yield measurements of multiple cultivars is presented as well. (B) Average yield of H4107 and H9780 across years and locations presented in A. *, statistically significant difference (P-value < 0.05).

Figure 2. Field experiments conditions and yield. Temperatures were recorded constantly in the three experimental sites: Upper Galilee, Volcani and Eden. Daily average (A), daily maximum (B), night average (C) and night maximum (D) were calculated for the reproductive period and are presented from the first day of flowering until the end of the experiment (88 days after flowering). (E) Yield performance for H4107 and H9780 in the Volcani (left), Upper Galilee (middle) and Eden (right) fields. **, statistically significant difference (P-value < 0.01).

Figure 3. Fruit set and seed number measurements in the field experiments. (A) Fruit set rates of H4107 and H9780 in Upper Galilee (left) and Volcani (right) fields. (B) Seeds number per fruit for H4107 and H9780 in Upper Galilee (left) and Volcani (right) fields. *, statistically significant difference (P-value < 0.05).

Figure 4. Controlled experiment conditions and reproductive measurements. (A) Temperatures measured every five minutes in both control (blue) and MCHS (brown) greenhouses. Black arrows denote day of flowering and day of stress initiation. Threshold temperatures for heat stress conditions in tomato are marked by dotted lines. (B) Fruit set ratio for H4107 and H9780 under control (white bars) and MCHS (grey bars) conditions. (C) Seeds number per fruit in H4107 and H9780 under control (white bars) and MCHS (grey bars) conditions. MCHS, moderate chronic heat stress. *, statistically significant difference (P-value < 0.05). ns, not significant.

Figure 5. Pollen viability in field and controlled experiments. Percentage of viable pollen from post-anthesis flowers of H4107 and H9780 at the (A) Upper Galilee field, (B) Volcani field and (C) controlled greenhouses, under control (white bars) and MCHS (grey bars) conditions. MCHS, moderate chronic heat stress. ns, not significant.

Table S1. Yield measurements in field trials of processing tomatoes between 2005 and 2019 in different locations. Presented here are yield values for H4107 and H9780 as well as the whole test average. na, not applicable – the cultivar was not tested

Year	Location	Yield (Kg/m ²)		
		H4107	Test Average	H9780
2005	Akko	na	13.1	14.2
2005	Upper Galilee	na	12.3	14.2
2005	Yifat	na	12.1	10.8
2006	Akko	na	13.4	12.9
2006	Beit HaShita	na	9.9	10.4
2006	Eden	na	11.9	11.8
2006	Kfar Hahores	na	11.2	9.9
2006	Yaen	na	7.6	5.1
2007	Geva	na	11.5	11.5
2007	Megido	na	10.8	10.8
2007	Upper Galilee	na	12.6	13.7
2008	Akko	na	11.6	10.7
2008	Geva	na	12.2	12.4
2008	Megido	na	10.7	8.8
2008	Upper Galilee	na	8.1	7.9
2009	Eden	na	12.7	13.4
2009	Geva	na	13.4	14.5
2009	Megido	na	11.8	11.8
2009	Upper Galilee	na	12.8	13.9
2010	Akko	na	13.8	13.3
2010	Eden	na	13.0	13.0
2010	Ramat David	na	10.8	10.8
2010	Geva	na	12.5	12.7
2010	Mesilot	na	8.4	7.3
2010	Upper Galilee	na	5.8	5.8
2011	Akko	15.1	14.2	13.7
2011	Eden	10.6	9.3	9.1
2011	Geva	13.1	9.8	9.3
2011	Gilad	8.9	8.1	8.0
2011	Upper Galilee	12.9	10.9	11.1
2012	Akko	15.2	13.0	na
2012	Eden	11.7	10.7	na
2012	Gadash haemek	10.8	10.7	na
2012	Geva	16.5	14.7	na
2012	Neve Eitan	10.7	9.8	na
2012	Upper Galilee	11.9	10.2	na
2013	Eden	11.2	11.3	na
2013	Gadash haemek	na	13.7	12.9
2013	Geva	na	9.9	10.0
2013	Upper Galilee	na	12.3	12.2
2014	Akko	16.0	13.5	13.0
2014	Gadash haemek	14.5	12.6	na
2014	Geva	15.8	13.2	na
2014	Upper Galilee	13.9	9.4	na
2015	Akko	15.7	14.4	na
2015	Gadash haemek	13.2	11.3	na
2015	Upper Galilee	14.7	13.3	na

2016	Akko	12.1	10.1	na
2016	Eden	16.4	14.5	na
2016	Gadash haemek	7.8	7.3	na
2016	Geva	9.8	10.0	na
2016	Upper Galilee	7.8	6.5	na
2017	Akko	17.7	14.8	na
2017	Midrach Oz	10.3	9.7	na
2017	Geva	14.3	14.0	na
2017	Upper Galilee	13.1	11.1	na
2018	Upper Galilee	9.4	8.4	6.1
2018	Geva	10.8	9.4	na
2018	Midrach Oz	9.4	8.9	na
2018	Akko	13.5	12.5	na
2018	Upper Galilee	10.3	7.6	na
2019	Akko	13.7	12.8	na
2019	Upper Galilee	6.9	4.7	3.3

581

582

583