Integrative transcriptomic analysis of tissue-specific metabolic crosstalk after myocardial infarction

Muhammad Arif^{1, #}, Martina Klevstig^{2, #}, Rui Benfeitas³, Stephen Doran⁴, Hasan Turkez⁵, Mathias Uhlén¹, Maryam Clausen⁶, Johannes Wikström⁷, Damla Etal⁶, Cheng Zhang¹, Malin Levin², Adil Mardinoglu^{1, 4, *}, Jan Boren^{2, *}

¹Science for Life Laboratory, KTH - Royal Institute of Technology, Stockholm, SE-17121, Sweden.

²Department of Molecular and Clinical Medicine, University of Gothenburg, The Wallenberg Laboratory, Sahlgrenska University Hospital, Gothenburg, Sweden.

³National Bioinformatics Infrastructure Sweden (NBIS), Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, S-10691 Stockholm, Sweden

⁴Centre for Host-Microbiome Interactions, Faculty of Dentistry, Oral & Craniofacial Sciences, King's College London, London, SE1 9RT, United Kingdom.

⁵Department of Medical Biology, Faculty of Medicine, Atatürk University, Erzurum, 25240, Turkey.

⁶Translational Genomics, BioPharmaceuticals R&D, Discovery Sciences, AstraZeneca, Gothenburg, Sweden.

⁷Bioscience Cardiovascular, Research and Early Development, Cardiovascular, Renal and Metabolism (CVRM), BioPharmaceuticals R&D, AstraZeneca, Gothenburg, Sweden

[#]Contributed equally.

*Corresponding authors: Adil Mardinoglu and Jan Boren

Email: adilm@scilifelab.se; Jan.Boren@wlab.gu.se

33 **Abstract/Summary**

34 Myocardial infarction (MI) promotes a range of systemic effects, many of which are
 35 unknown. Here, we investigated the alterations associated with MI progression in heart and
 36 other metabolically active tissues (liver, skeletal muscle, and adipose) in a mouse model of
 37 MI (induced by ligating the left ascending coronary artery) and sham-operated mice. We
 38 performed a genome-wide transcriptomic analysis on tissue samples obtained 6- and 24-hours
 39 post MI or sham operation. By generating tissue-specific biological networks, we observed:
 40 (1) dysregulation in multiple biological processes (including immune system, mitochondrial
 41 dysfunction, fatty-acid beta-oxidation, and RNA and protein processing) across multiple
 42 tissues post MI; and (2) tissue-specific dysregulation in biological processes in liver and heart
 43 post MI. Finally, we validated our findings in two independent MI cohorts. Overall, our
 44 integrative analysis highlighted both common and specific biological responses to MI across a
 45 range of metabolically active tissues.

46

47 **Keywords**

48 Systems biology; network analysis; whole-body modelling; cardiovascular disease;
 49 Myocardial infarction; multi-tissue; metabolically active tissues; liver; adipose; muscle

50

51 Introduction

52 Cardiovascular disease (CVD) is the leading cause of death worldwide, accounting for more
53 than 17 million deaths globally in 2016 ¹. Myocardial infarction (MI) is one of the most
54 common causes of CVD-related death, and is the result of severe coronary artery disease that
55 develops from tapered arteries or chronic blockage of the arteries caused by accumulation of
56 cholesterol or plaque (atherosclerosis). Many behavioral risk factors (including unhealthy
57 diet, physical inactivity, excessive use of alcohol, and tobacco consumption), which are
58 responsible for hypertension, obesity, diabetes, and hyperlipidemia by significantly altering
59 metabolism, are also implicated in MI. These abnormalities are known as the high-risk factors
60 of MI and CVDs in general.

61
62 Systems biology has been used in many studies to reveal the underlying molecular
63 mechanisms of complex human diseases and to answer important biological questions related
64 to the progression, diagnosis and treatment of the diseases. The use of systems biology has
65 aided the discovery of new therapeutic approaches in multiple diseases ¹⁻³ by identifying
66 novel therapeutic agents and repositioning of existing drugs ⁴. Systems biology has also been
67 employed in the identification of novel biomarkers, characterization of patients and
68 stratification of heterogenous cancer patients ⁵⁻⁷. Specifically, integrated networks (INs) ⁷ and
69 co-expression networks (CNs) ⁸ have been proven to be robust methods for revealing the key
70 driver of metabolic abnormalities, discovering new therapy strategies, as well as gaining
71 systematic understanding of diseases ^{9,10}.

72
73 Previously, multiple studies in individual tissues have been performed and provided new
74 insights into the underlying mechanisms of diseases ¹¹⁻¹⁴. However, the crosstalk between
75 different tissues and their dysregulation has not been examined in MI and other CVD-related
76 complications ¹⁵. Here we performed an integrated analysis of heart and other metabolically
77 active tissues (liver, skeletal muscle and adipose tissue) using a mouse model of MI. We used
78 several systems biology approaches to obtain a systematic picture of the metabolic alterations
79 that occur after an MI (**Figure 1A**), and validated our findings in two independent datasets.

80 Results

81 *Differential expression analysis shows a pronounced effect on gene expression 24 h post*

82 MI

83 To study global biological alterations and systemic whole-body effects associated with MI,
84 we obtained heart, liver, skeletal muscle, and white adipose tissue from mice 6 h and 24 h
85 after either an MI (induced by ligating the left ascending coronary artery) or a sham operation
86 (as control). Total of 20 mice were used in this study (5 mice in each time and condition
87 combination) (**Figure 1A**). We generated transcriptomics data and identified differentially
88 expressed genes (DEGs) 6 and 24 h post MI and sham operation in all tissues, with the most
89 significant differences occurring after 24 h (**Supplementary File 1, Figure 1B**). Principal
90 component analysis (PCA) showed a close clustering between the control (for both time
91 points) and MI (6 h and 24 h separately) samples for heart tissue but clustering by extraction
92 time points (6 h and 24 h clusters) for the other tissues (**Figure 1 - Figure Supplement 1**).
93 We present the transcriptional changes associated with MI in **Supplementary File 1** and the
94 DEGs (FDR < 5%) using an UpSet plot¹⁶ in **Figure 1C**.

95

96 All tissues showed a more pronounced effect in terms of the number of DEGs 24 h post MI
97 (**Figure 1C**). As expected, the most affected tissue was the heart (393 DEGs at 6 h, 3318
98 DEGs at 24 h, and 318 DEGs were the same at both time points). By contrast, 136, 641 and
99 374 genes were significantly changed in liver, skeletal muscle and adipose tissues 24 h post
100 MI compared to control, respectively. More than 33% of the DEGs that significantly changed
101 in the other tissues also changed in the heart (**Figure 1C**). Interestingly, more than 97% of the
102 shared DEGs between heart and skeletal muscle changed in the same direction, with
103 corresponding numbers of 88% and 64% in adipose and liver, respectively.

104

Functional analysis reveals widespread alterations of mitochondrial, fatty acid, immune, and protein and RNA-related biological processes post MI with liver shows contrasting trend

We performed gene-set enrichment analysis (GSEA) with KEGG pathways (**Supplementary File 2, Figure 1D**) and gene ontology (GO) biological processes (BPs) (**Supplementary File 3, Figure 2A**) to identify altered biological functions and pathways 24 h after an MI.

Mitochondrial functions (specifically, mitochondrial translation, respiratory chain and oxidative phosphorylation) were significantly downregulated in the heart, muscle and adipose tissues but not in the liver. Processes related to oxidative stress were upregulated in the heart and skeletal muscle. Fatty acid beta-oxidation was downregulated in the heart and adipose but upregulated in the liver. Processes and pathways related to immune systems were significantly upregulated in the heart and skeletal muscle but significantly downregulated in liver. Processes associated with protein and RNA processing, ribosome biogenesis and protein targeting endoplasmic reticulum were upregulated in all tissues except liver whereas protein processing in endoplasmic reticulum and RNA transport pathways were upregulated in all tissues.

We also observed that liver was showing opposite trends compared to the other tissues in other important functions, such as fatty acid metabolism and immune response. By checking regulation at the gene level, we observed that only 16 DEGs in liver showed opposite regulation compared to the other tissues whereas 97 out of the 136 DEGs in liver were not DEGs in any other tissues (**Supplementary File 4**). Therefore, the differences we observed in liver were mainly due to different DEGs rather than opposite regulation compared to other tissues.

Tissue-specific altered biological functions point to specificity of metabolic and signaling responses to MI

The functional analysis also indicated that several metabolic pathways (including cholesterol, ascorbate and aldarate, linoleic acid, and sphingolipid metabolism pathways) and signaling pathways (including GnRH, FoxO, cAMP and prolactin signaling pathways) were significantly upregulated in heart 6 h after an MI (**Supplementary File 2, Figure 1 - Figure Supplement 2**). We also observed significant down regulation of tryptophan metabolism and upregulation of glycosaminoglycan biosynthesis in heart 24 h after an MI (**Supplementary File 2, Figure 1 - Figure Supplement 2**). Processes related to retinol metabolism were upregulated in heart at both timepoints. Pathways that were previously associated with cardiac hypertrophy and cardiac remodeling (e.g. JAK-STAT, MAPK, estrogen, and TNF signaling pathways, and ECM-receptor interaction) were significantly upregulated in heart 6 and 24 h after an MI (**Figure 1 - Figure Supplement 4**).

Our analysis also indicated significant metabolic differences in adipose tissue 24 h after an MI (**Figure 1 - Figure Supplement 3**). Fructose and mannose metabolism, glyoxylate and dicarboxylate metabolism, glycolysis/gluconeogenesis, and pentose phosphate pathways, glycine, serine and threonine metabolism and pyrimidine metabolism, as well as endocrine systems (e.g. insulin signaling pathway and regulation of lipolysis in adipocytes) were downregulated in adipose tissue.

We observed that the PPAR signaling pathway was upregulated whereas glutathione was downregulated in liver 24 h post-infarction (**Figure 1 - Figure Supplement 3**). We found that sphingolipid metabolism and immune-related pathways were upregulated in skeletal muscle 24 h post-infarction (**Figure 1 - Figure Supplement 3**).

Reporter metabolite analyses show significant alterations in fatty acid, amino acid, retinol, and estrogen metabolism post MI

To predict the effect of the transcriptional changes on metabolism, we performed reporter metabolite analyses (**Supplementary File 5**) using the gene-to-metabolites mapping from the Mouse Metabolic Reaction database¹⁷; results in each tissue 24 h after MI are shown in **Figure 2B**. In agreement with our analyses above, reporter metabolites related to oxidative phosphorylation, such as *ubiquinol*, *ubiquinone*, *NADH* and *NAD+*, were downregulated in all tissues except liver. Moreover, *linolenoyl-CoA*, *acetyl CoA*, and several other fatty acyl-CoA-related metabolites were downregulated in heart and adipose tissue but upregulated in liver. We also found that several *5-S-glutathionyl* metabolite forms, known to be related to phenylalanine, tyrosine and tryptophan biosynthesis, were downregulated in heart, liver and skeletal muscle. The same pattern of downregulation was also observed for metabolites related to estrogen metabolism, specifically metabolites related to oestrone and its glutathione conjugate derivative. Moreover, *12-keto-LTB4* and *12-oxo-c-LTB3*, related to leukotriene metabolism, and *hepoxilin A3*, an arachidonic acid, were also found to be downregulated in heart, liver, and skeletal muscle.

The liver showed the highest alteration in reporter metabolites, which is attributed to its role as one of the most metabolically active tissues. We found that several reporter metabolites related to retinol metabolism, namely *retinal*, *retinol*, *retinoate*, and *all-trans-18-hydroxyretinoic acid*, were significantly downregulated only in liver tissue. Retinol metabolism has been previously associated with MI^{18,19}.

Network analyses unveil universal and tissue-specific clusters and mechanisms post MI

The use of co-expression network (CN) analyses can assist in elucidating the functional relationships between genes in a specific cell and tissue⁸. Here, we performed CN analysis to reveal the functional relationship between the DEGs by generating tissue-specific CNs and selected highly connected genes (the top 5% positively correlated genes that fulfilled FDR < 0.05) (**Table 1**). To better define the structure of the networks, we used the Leiden clustering algorithm²⁰ by maximizing the modularity scores (**Figure 3A-D**) and selected the clusters that include more than 30 genes. Next, we superimposed DEGs 24 h post-infarction onto the network (**Supplementary File 1**) and identified the components of the clusters that were

affected by an MI. We also used functional analysis with GO BP and KEGG pathways to understand the specific functions associated with each cluster by using the Enrichr algorithm (FDR < 0.05)^{21,22}. We summarized the GO BP terms with Revigo (**Supplementary File 6**)²³ and checked the average clustering coefficient to define the centrality of each cluster (**Supplementary File 6**)⁸. Among the clusters, we identified the key clusters as those with the highest average clustering coefficient, allowing us to identify sets of genes whose time-dependent coordinated changes showed the strongest relationships.

Interestingly, key clusters contained genes with similar functionalities including RNA processing, transports, and RNA metabolic processes in all tissue-specific CNs (**Supplementary File 6**). In addition, we found that the majority of the DEGs associated with those clusters were significantly upregulated. These observations strengthen the findings of the functional analysis above (**Figure 2A**) and further highlight how embryonically distinct tissues display similar functional responses to MI, with the most highly connected groups of genes preserved between different tissues (**Supplementary File 6, Figure 3E**).

Community detection reveals tissue-specific clusters post MI

We investigated the tissue specificity of each cluster by performing enrichment analysis with data from the Mouse Gene Atlas²⁴, which involved counting the number of tissue-specific genes.

The heart network showed the highest number of tissue-specific genes in cluster Heart-3 (302 genes). Based on DEG analysis, we found that 522 genes were downregulated and 192 genes were upregulated in the cluster. The enriched GO BP terms in the cluster were mitochondrial transport, protein processing and respiratory chain, cardiac muscle cell action potential, response to muscle stretch, and heart contraction (**Figure 3F**). We observed that the results of the KEGG pathway enrichment analysis were consistent with those obtained from GO BP analysis (**Supplementary File 6**).

In the liver network, cluster Liver-2 showed the highest tissue specificity (479 genes). In this cluster, we found that 15 genes were significantly downregulated and 17 genes were significantly upregulated. Based on GO BP enrichment analysis, the genes in this cluster were associated with cholesterol metabolism and homeostasis, lipid transport, glutathione

metabolism, lipoprotein metabolism, and glucose 6-phosphate metabolism (**Supplementary File 6**). KEGG enrichment analysis also showed that the genes in the cluster were related to retinol, carbohydrate, lipid and amino-acid metabolism (**Supplementary File 6**).

The muscle network had two clusters with high tissue specificity: cluster Muscle-4 (276 genes) and Muscle-5 (143 genes). Muscle-4 showed association with GO BP terms such as mitochondrial transport, protein processing and respiratory chain, response to muscle stretch, and muscle contraction (**Supplementary File 6**). In contrast, the KEGG pathway in this cluster showed relation to glycolysis/glucogenesis, propanoate metabolism, glyoxylate and dicarboxylate metabolism, and several signaling pathways (e.g. oxytocin, glucagon, cGMP-PKG and HIF-1) (**Supplementary File 6**). Muscle-5 was enriched in GO BP terms associated with protein dephosphorylation, muscle contraction and intracellular protein transport (**Supplementary File 6**). We also found that insulin, MAPK and Wnt signaling pathways were associated to Muscle-5 from the KEGG enrichment analysis (**Supplementary File 6**).

The adipose tissue network showed tissue specificity in cluster Adipose-2 (33 genes), which is associated with GO BP processes including mRNA processing, regulation of mitotic cell cycle phase, ribosome biogenesis, and viral processes (**Supplementary File 6**). We observed that the results of the KEGG pathway enrichment analysis were consistent with those obtained from GO BP analysis, with additional associations with multiple signaling and regulatory pathways (**Supplementary File 6**).

Tissue-specific clusters show important tissue-specific changes post MI

To understand the specific behavior of each tissue, we further studied the tissue-specific clusters in the CNs (**Figure 4A**). Heart specific cluster, Heart-3, was driven by several central genes including *Pln*, *Pde4b*, and *Atp2a2* (related to regulation of cardiac muscle contraction) and *Pdha1* and *Vdac1* (related to mitochondrial functions). These genes were also found to be significantly differentially expressed in heart 24 hours post MI (**Supplementary File 1**). Genes in the heart-specific cluster were related to multiple other processes/pathways, e.g. oxytocin signaling pathway, and several metabolic pathways (glycogen, inositol phosphate and purine) (**Supplementary File 6**).

Mitochondrial dysfunction in the heart leads to disturbance of energy (ATP) production^{25,26} and, in the presence of oxygen, to accumulation of reactive oxygen species (ROS), which can cause oxidative stress. *Vdac1*, a key gene for regulation of mitochondria function and one of the central genes in the heart-specific cluster (see above), is significantly downregulated in MI²⁷. *Vdac1* is located in the outer mitochondrial membrane and is involved directly in cardioprotection²⁹ within the cGMP/PKG pathway (**Figure 4 - Figure Supplement 1**). In the same pathway, we also observed down-regulation of the reporter metabolite hydrogen peroxide (**Supplementary File 5**), a ROS that is related to cardioprotection^{28,29}. We also observed downregulation of *Pdha1*, which is known to have a substantial role in both the HIF-1 signaling pathway and the pyruvate metabolism pathway that converts pyruvate to acetyl-CoA in the mitochondria (**Figure 4 - Figure Supplement 2**). Acetyl-CoA is used in the TCA cycle to produce NADH and FADH₂, which are both needed for ATP production and were downregulated in our reporter metabolite analysis of the heart. Our findings are thus consistent with dysfunctional mitochondria and ATP production in the heart in response to an MI. *Pdha1* has been also been linked to the heart sensitivity during to ischemic stress, where its deficiency can compromise AMP-activated protein kinase activation³⁰.

In skeletal muscle and adipose tissue, we found that central genes in their respective tissue-specific clusters related to fatty acid metabolism and lipid metabolism were significantly altered (**Supplementary File 6, Figure 5**). In liver-specific cluster, we found that their central genes were related to fatty-acid beta oxidation (*Cyp4a31*, *Cyp4a32*) and glutathione metabolism (*Gstm3*) (**Supplementary File 6, Figure 5A**). Alterations of fatty acid beta-oxidation and glutathione metabolism have previously been reported in non-alcoholic fatty liver disease, a known risk factor of CVD^{31,32}. Moreover, in liver, we also found that retinol metabolism was uniquely related to genes in the liver-specific cluster, mainly driven by four significantly differentially expressed central genes of the clusters, i.e. *Cyp26a1*, *Cyp4a31*, *Cyp4a32*, and *Hsd17b6* (**Supplementary File 6**). A previous study showed that mortality from CVD in older individuals was accompanied by impaired liver ability to store retinol¹⁸.

Multi-tissue modeling reveals key metabolic pathways affected post MI

To investigate the metabolic responses to MI in and across tissues in the mice, we constructed a multi-tissue genome-scale metabolic model. The model consisted of five tissue-specific genome scale metabolic models, namely heart, liver, skeletal muscle, adipose, and small

intestine. The small intestine model (for which we do not have transcriptomic data) was added to include ingestion and conversion of dietary nutrients into chylomicrons, which are directly secreted into blood and transport lipids to other tissues¹⁷. The final mouse multi-tissue model included 19,859 reactions, 13,284 metabolites, 7,116 genes and 41 compartments. We predicted the metabolic fluxes in mice 24 h after an MI or sham operation by integrating the dietary input, tissue-specific resting energy expenditure and transcriptomics data.

The modeling showed that oxygen uptake, carbon dioxide production and the oxidative phosphorylation pathway in heart, adipose and skeletal muscle were decreased in MI mice, in agreement with the downregulation of oxidative phosphorylation we observed in these tissues (**Supplementary File 7**). By contrast, liver showed slightly increased oxygen uptake, which might be due to the slightly (not statistically significant) upregulated oxidative phosphorylation (**Supplementary File 7**). These findings indicate that the changes in oxygen and carbon dioxide fluxes and the oxidative phosphorylation pathway could serve as a positive control for predicting the changes due to MI in the fluxes.

Next, we investigated the tissue-specific metabolic flux changes in the same model (**Supplementary File 7**). We found that the pentose phosphate pathway was upregulated in heart 24 hours post MI, consistent with upregulated glucose metabolism after an MI. Elevated glycolysis could allow the heart to rapidly generate energy under stress conditions, and the enhanced pentose phosphate pathway could increase the NADPH level, which could help maintain the level of reduced glutathione in heart³³. In addition, we observed an increase uptake of alpha-ketoglutarate (AKG) of heart 24h after MI. It has been reported that supplementation of AKG could prevent heart from ischaemic injury³⁴, and the increased uptake of AKG we observed after MI might be a natural protective metabolic response to MI. Moreover, we found there is a net lactate metabolic flux coming from liver to heart in the MI group. The influx of lactate has been reported to be positively correlated with the fraction of regional ejection of heart³⁵ and this net flux not only agrees well with the previous report but also additionally suggested the source of the lactate. We also found that adipose tissue secreted more ketone bodies, including acetoacetate and butyrate, into plasma; the plasma level of ketone bodies has been reported as a stress marker in acute MI³⁶. Notably, relatively small metabolic changes were found in liver and skeletal muscle, which is probably due to the small number of transcriptomic changes in metabolic pathways in these tissues.

Validating our findings with publicly available datasets

We validated our observations in heart tissue in two independent cohorts of bulk RNA-seq data from mouse heart (**Supplementary File 8**). We filtered both validation cohorts to get and analyzed only 24 hours post-MI data. We found that there were 2169 DEGs from our heart 24 h post MI data were validated in at least one of the independent cohorts (959 DEGs validated in both) (**Figure 6A**). We also found that 109 out of the 123 most connected genes in our heart-specific cluster were also significantly differentially expressed in at least one of the independent cohorts (81 in both). By performing functional analysis of the validation cohorts, we found that ~61% of GO BP and 84% of KEGG pathways identified in our analysis of the heart were also present in at least one of the validation cohorts 24 h after infarction (**Figure 6B-C**). In both cohorts, we observed downregulation of mitochondrial functions and fatty acid metabolism processes. We also observed upregulation of processes and pathways related to retinol metabolism and inflammatory response in both validation cohorts.

Identification of driver genes in MI

We observed that *Flnc*, *Lgals3*, *Prkaca* and *Pprc1* showed important role to MI. These genes were 4 of 16 genes that were DEGs in at least three tissues and validated in both validation cohorts (**Supplementary File 9**). *Flnc*, *Lgals3* and *Pprc1* were upregulated in heart, skeletal muscle, and adipose, whereas *Prkaca* was downregulated in these three tissues. We further retrieved their neighbors at each tissue specific CNs, showed their regulations from differential expression results, and performed functional analysis in **Supplementary File 9**.

Flnc, which encodes filamin-C, was part of heart and skeletal muscle-specific CN cluster (**Figure S4**). Its neighbor genes were found to be significantly (FDR < 0.05) associated to several functions, including TCA cycle, pyruvate metabolism, glycolysis pathway, and involved in mitochondrial functions. Specifically, they were related to heart-specific processes in heart, VEGF signaling pathway in muscle, carbohydrate metabolism in adipose, and to MAPK signaling pathway and muscle contraction in heart and muscle.

Lgals3 (encodes galectin-3) and *Prkaca* were among the most central genes in central clusters (**Supplementary File 6**). The neighbors of *Lgals3* were significantly related to cell cycle and protein digestion and absorption pathway in all tissues, and to RNA and mRNA related-processes in muscle and adipose tissue. The neighbors of *Prkaca* were related to insulin

351 signaling pathway in heart and adipose, and several mitochondrial functions in adipose. *Pprc1*
 352 was part of most central clusters in heart and adipose tissue CN, and its neighbors were
 353 related to ribosomal RNA processing and ribosome biogenesis.
 354

Discussion

CVD has a complex etiology and is responsible for a range of systemic effects, hindering our understanding of its consequences on different tissues. Here, we took advantage of the technological advances in high-throughput RNA-seq and applied integrative network analyses to comprehensively explore the underlying biological effects of MI. Specifically, we generated RNA-seq data from heart, liver, skeletal muscle and adipose tissue obtained from mice 6 and 24 h after an MI or sham operation. We used transcriptomics data analyses (differential expression, functional analysis, and reporter metabolites analysis) to determine the systemic effects of the MI across multiple tissues. Moreover, we performed CN analyses to pinpoint important key and tissue-specific clusters in each tissue, and identified the key genes in each cluster. Finally, we used a whole-body modelling approach to identify the crosstalk between tissues and reveal the global metabolic alterations, before finally validating our findings with publicly available independent MI cohorts.

Based on our analyses, we observed downregulation of heart-specific functions and upregulation of lipid metabolism and inflammatory response in heart, muscle, and adipose tissue after an MI (**Figure 4B**). Liver showed a distinct response with respect to the other three tissues, including downregulation of inflammatory response. We observed that fatty acid metabolism was downregulated in heart and adipose tissue, whereas fatty acid beta-oxidation was upregulated and glutathione metabolism was downregulated in liver. We also observed upregulation of oxidative stress in heart and skeletal muscle. We also observed downregulation of mitochondrial functions in heart, muscle, and adipose tissue. Furthermore, we found upregulation of retinol metabolism in heart and downregulation of retinol metabolites in liver (**Figure 4B**).

We hypothesized that downregulation of fatty acid metabolism from adipose tissue was due to exchange of fatty acids with other tissues (liver and muscle) (**Figure 4B**). We also observed the flow of retinol from liver to heart during MI, consistent with previous reports¹⁹. These MI-associated alterations lead to dysfunctional mitochondria and decreased energy production, especially in heart and skeletal muscle.

We also validated our results with publicly available MI datasets generated in separate independent studies. The validation results strengthened our findings on the altered functions/pathways and the important heart-specific genes after an MI.

Importantly, our analyses of gene clusters highlighted multiple key genes in the response to MI in different tissues. Specifically, we observed that *Flnc*, *Prkaca*, *Lgals3*, and *Pprc1* showed important responses in heart, skeletal muscle, and adipose tissue. *Flnc* is involved in actin cytoskeleton organization in heart and skeletal muscle, and previous studies have shown that this gene has critical role in CVD^{37,38}. Similarly, *Prkaca*, an important metabolic gene, has also been shown to play an important function during CVD³⁹⁻⁴¹. *Lgals3*, related to acute inflammation response, has been studied intensively in recent years as a key gene in CVD, and as a potential CVD therapy target^{42,43}. Lastly, *Pprc1*, as important regulator of mitochondrial biogenesis, has not been explored for its direct relationship with CVD; however, mitochondrial biogenesis appears to be an important response to CVD⁴⁴⁻⁴⁶.

We recognized several limitations to be noted on this research. First, only transcriptomic data was analyzed in this research, hence the sensitivity might be limited especially for short timepoint, e.g. 6 hours after MI. Second, we focused our analysis in this research only on protein-coding genes. Third, to explore more about the shift in metabolism due to MI, longer timepoints needs to be explored. This opens new opportunities for future research, including analyzing the non-protein-coding gene signatures and longer timepoints.

In summary, we systematically unveiled the deregulation of biological processes and pathways that resulted from MI in heart, liver, muscle, and adipose tissue by integrating transcriptomic data and the use of biological networks. We also identified the key clusters and central genes using generated tissue-specific CNs. In this study, we demonstrated a strategy to utilize multi-tissue transcriptomic data to identify alteration of biological processes and pathways to systemically explore the effect of a disease.

Author Contribution

MK performed the animal experiments, MA performed the computational analysis and analyzed the clinical data together with RB, SD, HT, MU, MC, JW, DE, CZ, AM, and JB

coordinated the generation of the clinical data. MA, MK, AM and JB wrote the paper and all authors were involved in editing the paper.

Acknowledgements

This work was financially supported by the Knut and Alice Wallenberg Foundation, Swedish Research Foundation and Swedish Heart-Lung Foundation.

Conflict of Interest

JW, MC, DE are employees at AstraZeneca. The other authors declare no conflict of interest.

428 Material and Methods

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
commercial assay or kit	RNeasy Fibrous Tissue Mini Kit	Qiagen		Heart and Skeletal Muscle Tissue
commercial assay or kit	RNeasy Mini Kit	Qiagen		Liver Tissue
commercial assay or kit	RNeasy Lipid Tissue Mini Kit	Qiagen		Adipose Tissue
commercial assay or kit	cDNA Reverse Transcription Kit	Applied Biosystems		
commercial assay or kit	TaqMan real-time PCR in a ViiA™ 7 system	Applied Biosystems		
commercial assay or kit	NovaSeq6000	Illumina		
software, algorithm	NovaSeq Control Software 1.6.0/RNA v3.4.4	Illumina		
software, algorithm	CASAVA Software Suite	Illumina		
software, algorithm	Kallisto		RRID:SCR_016582	
software,	Python 3.7	Python	RRID:SCR_008394	

algorithm		Programmin g Language		
software, algorithm	sklearn	Python Package	RRID:SCR_019053	
software, algorithm	R	R Project for Statistical Computing	RRID:SCR_001905	
software, algorithm	Rpy2	Python Package	https://rpy2.github.io/	
software, algorithm	DESeq2	R Package	RRID:SCR_015687	
software, algorithm	PIANO	R Package	RRID:SCR_003200	
software, algorithm	SciPy	Python Package	RRID:SCR_008058	
software, algorithm	Statsmodel	Python Package	RRID:SCR_016074	
software, algorithm	iGraph	Python Package	RRID:SCR_019225	
software, algorithm	Leiden Clustering	Python Package	https://github.com/vtraag/leidenalg	
software, algorithm	Matlab	Mathworks	RRID:SCR_001622	

Induction of MI

10-week-old male C57Bl/6N mice were fasted for 4 h before induction of myocardial infarction. The mice were then anesthetized with isoflurane, orally intubated, and connected to a small-animal ventilator (SAR-830, Geneq, Montreal, Canada) distributing a mixture of oxygen, air and 2–3% isoflurane. ECG electrodes were placed on the extremities, and cardiac rhythm was monitored during surgery. An incision was made between the 4th and 5th ribs to reveal the upper part of the anterior left ventricle (LV) wall and the lower part of the left atrium. Myocardial infarction was induced by ligating the left anterior descending (LAD) coronary artery immediately after the bifurcation of the left coronary artery 1. The efficacy of the procedure was immediately verified by characteristic ECG changes, and akinesis of the LV anterior wall. After verification of the infarction, the lungs were hyperinflated, positive

end-expiratory pressure was applied, and the chest was closed. Sham mice were handled identically (fasted, anesthetized, intubated, and connected to ventilator, and subsequently incised between 4th and 5th ribs), but no ligation of the LAD coronary artery was performed (and thus, no ischemia was induced in these mice). The mice received an intraperitoneal injection of 0.1 ml buprenorphine to relieve postoperative pain and were allowed to recover spontaneously after stopping isoflurane administration. Mice were killed with an overdose of isoflurane 6 h or 24 h after occlusion or sham operation. We collected the left ventricle (the whole left ventricle containing mainly infarcted tissue) of the heart, whereas white adipose tissue (WAT) was collected from the abdomen and musculus soleus was taken as the muscle tissue. Mouse hearts and biopsies from the liver, muscle and WAT were snap-frozen in liquid nitrogen and stored at -80°C until analysis. All mice studies were approved by the local animal ethics committee and conform to the guidelines from Directive 2010/63/EU of the European Parliament on the protection of animals used for scientific purposes.

Echocardiography in mice

Echocardiographic examination, using VisualSonics VEVO 2100 system (VisualSonics Inc, Ontario, Canada), which includes an integrated rail system for consistent positioning of the ultrasound probe was performed 6 and 24 h after an MI to determine the size of the MI. We calculated infarct size based on wall motion score index (WMSI) 24 h after myocardial infarction by a 16-segments model on 3 short axis images, as 0 for normal, ½ for reduced wall thickening and excursion in a segment and 1 for no wall thickening and excursion in a segment. WMSI was calculated as the sum of scores divided by the total number of segments. Hair removal gel was applied to isoflurane-anesthetized (1.2%) mice chest to minimize resistance to ultrasonic beam transmission. The mice were then placed on a heating pad and extremities were connected to an ECG. A 55 MHz linear transducer (MS550D) was used for imaging. An optimal parasternal long axis (LAX) cine loop of >1000 frames/s was acquired using the ECG-gated kilohertz visualization technique. Parasternal short axis cine-loops were acquired at 1, 3, and 5 mm below the mitral annulus. Infarct size was calculated based on wall motion score index 6 and 24 hours after myocardial infarction by a 16-segments model on LAX and 3 short axis images view, as 0 for normal, ½ for reduced wall thickening and excursion in a segment and 1 for no wall thickening and excursion in a segment. The data were evaluated using VevoStrain™ software system (VisualSonics Inc, Ontario, Canada).

RNA extraction and sequencing

Total RNA was isolated from snap-frozen tissues using RNeasy Fibrous Tissue Mini Kit (Qiagen) for heart and skeletal muscle, RNeasy Mini Kit (Qiagen) for liver, or RNeasy Lipid Tissue Mini Kit (Qiagen) for adipose tissue. cDNA was synthesized with the high-capacity cDNA Reverse Transcription Kit (Applied Biosystems) and random primers. mRNA expression of genes of interest was analyzed with TaqMan real-time PCR in a ViiA™ 7 system (Applied Biosystems). RNA sequencing library were prepared with Illumina RNA-Seq with Poly-A selections. Subsequently, the libraries were sequenced on NovaSeq6000 (NovaSeq Control Software 1.6.0/RNA v3.4.4) with a 2x51 setup using ‘NovaSeqXp’ workflow in ‘S1’ mode flow cell. The Bcl was converted to FastQ by bcl2fastq_v2.19.1.403 from CASAVA software suite (Sanger/phred33/Illumina 1.8+ quality scale).

RNA-sequencing data analysis

The raw RNA-sequencing results were processed using Kallisto⁴⁷ with index file generated from the Ensembl mouse reference genome (Release-96)⁴⁸. The output from Kallisto, both estimated count and TPM (Transcript per kilobase million), were subsequently mapped to gene using the mapping file retrieved from Ensembl BioMart website, by filtering only protein coding genes and transcripts. Genes with mean expression less than 1 TPM in each condition were filtered. For data exploration, we used PCA from sklearn package⁴⁹ in Python 3.7 and used TPM values as the input.

Subsequently, we performed differential gene expression analysis using DESeq2⁵⁰ package in R. We utilized the capabilities from DESeq2 to normalize the rounded estimated count data and to correct for confounding factors (such as time). To define a gene as differentially expressed (DEGs), a gene has to fulfill a criterion of FDR < 5%. The results of differential expression analysis were then used for functional analysis.

We checked the tissue specificity of the DEGs in each tissue with the data from Mouse Gene Atlas²⁴. For all the tissue-specific genes, we also checked their human-homolog genes in the human secretome database⁵¹.

Functional analysis

We performed functional analysis using the R package PIANO⁵². As the input, we used the fold changes and p-values from the DESeq2, and also GO BP and KEGG pathways gene-set collections from Enrichr^{21,22}, and metabolites from Mouse Metabolic Reaction database¹⁷. To define a process or pathway as significant, we used a cut off of FDR < 5% for the distinct direction of PIANO (both up and down).

Co-expression network generation

We generated the co-expression network by generating gene-gene Spearman correlation ranks within a tissue type, using *spearmanr* function from SciPy⁵³ in Python 3.7. Using the same environment, we performed multiple hypothesis testing using Benjamini-Hochberg method from *statsmodels*⁵⁴. Correlation data were filtered with criterion of adjusted p-value < 5%.

The top 5% of filtered correlation results were then loaded into iGraph module⁵⁵ in Python 3.7 as an unweighted network. To find the subnetworks, we employed the Leiden clustering algorithm²⁰ with *ModularityVertexPartition* method. Each cluster was analyzed by using Enrichr^{21,22} to get the enriched GO BP and KEGG pathways. Criterion FDR < 0.05 were used to find the significantly enriched terms. Clusters with less than 30 genes were discarded, to be able to get significant functional analysis results. Since GO BP was relatively sparse, we used Revigo²³ to summarize the GO BP into a higher level. Revigo was further employed to build a GO BP network. Clustering coefficient was calculated based on the average local clustering coefficient function within iGraph.

Multi-tissue metabolic modeling

We combined tissue-specific models (of heart, liver, muscle, adipose and small intestine) constructed previously¹⁷ in a multi-tissue model by adding an additional compartment representing the plasma, which allows the exchange of metabolites among different tissues. Blocked reactions that could not carry fluxes (and the unused metabolites and genes linked to these reactions) were removed from the models. In addition, the dietary input reactions and constraints were added to the small intestine model to simulate the food intake (**Supplementary File 7**). Specifically, we assumed that the mice weighed 30 g and consumed 4.5 g chow diet per day (15 g/100 g body weight) based on a previous study⁵⁶. We also

calculated the tissue-specific resting energy expenditures and set them as mandatory metabolic constraints based on previous studies and resting energy expenditure for other tissues was incorporated by including a mandatory glucose secretion flux out from the system with the lower bound calculated based on ATP (**Supplementary File 7**)⁵⁶.

To simulate the metabolic flux distribution in the sham-operated mice, we set the lipid droplet accumulation reaction in adipose tissue (m3_Adipose_LD_pool) as the objective function as we assume the energy additional to the resting energy expenditure will be mostly stored as fat rather than used by the muscle for physical activities because mice raised in the cages might have very little exercise. Then, we used parsimonious FBA to calculate the flux distribution. To simulate the flux distribution after an MI, we calculated an expected flux fold change of each reaction based on the FDR and expression fold changes of all genes associated with the reaction, and obtains a flux distribution that is closest to this expected flux distribution while satisfying the stoichiometric balance and flux constraints of the model. The mathematical formulation of the method is described as below,

$$\begin{aligned} \text{minimize } Z &= \sum_i |v_i - v_i^{exp}| \\ \text{s. t. } S * v &= 0 \\ lb &\leq v \leq ub \end{aligned}$$

where S , v , lb , ub represent the stoichiometric matrix, flux distribution, lower bound and upper bound of all reactions, respectively. The v_i^{exp} represents the expected flux of i^{th} reaction which is calculated as follows,

$$v_i^{exp} = v_i^{ref} * \sqrt[m]{\prod_{j=1}^m FC_j}$$

where n is the number of gene sets that could independently catalyze the corresponding reaction, and FC_j represents the expected expression changes of j^{th} gene set which is calculated below,

$$FC_j = (1 - P_1) * fc_1 + P_1(1 - P_2) * fc_2 + \dots + \prod_{k=1}^{m-1} P_k(1 - P_n) * fc_n$$

where m is the number of genes in the i^{th} gene sets, and P_j and fc_j respectively represents the FDR and fold change of gene expression with j^{th} smallest fold change in this gene set. In this way, genes with lowest fold change will have a dominating effect within a gene set encoding

a protein complex, while the geometric mean of expected fold changes of gene sets encoding different isozymes of this reaction will be used as the final expected flux fold change of this reaction.

Validation of the results

We validated our findings by performing similar steps of RNA sequencing and functional analysis for the publicly available mouse MI datasets GSE104187 and GSE52313^{13,14}.

Data and code availability

All raw RNA-sequencing data generated from this study can be accessed through accession number GSE153485. Codes used during the analysis are available on https://github.com/sysmedicine/ArifEtAl_2020_MultiTissueMI

References

- 1 Mardinoglu, A., Boren, J., Smith, U., Uhlen, M. & Nielsen, J. The employment of systems biology in gastroenterology and hepatology. *Nat. Rev. Gastroenterol. Hepatol* (2017).
- 2 Mardinoglu, A. & Nielsen, J. New paradigms for metabolic modeling of human cells. *Current Opinion in Biotechnology* **34**, 91-97 (2015).
- 3 Nielsen, J. Systems biology of metabolism: a driver for developing personalized and precision medicine. *Cell metabolism* **25**, 572-579 (2017).
- 4 Turanli, B. *et al.* Discovery of therapeutic agents for prostate cancer using genome-scale metabolic modeling and drug repositioning. *EBioMedicine* **42**, 386-396 (2019).
- 5 Benfeitas, R. *et al.* Characterization of heterogeneous redox responses in hepatocellular carcinoma patients using network analysis. *EBioMedicine* **40**, 471-487 (2019).
- 6 Bidkhori, G. *et al.* Metabolic network-based stratification of hepatocellular carcinoma reveals three distinct tumor subtypes. *Proceedings of the National Academy of Sciences* **115**, E11874-E11883 (2018).
- 7 Lee, S. *et al.* Integrated network analysis reveals an association between plasma mannose levels and insulin resistance. *Cell metabolism* **24**, 172-184 (2016).
- 8 Lee, S. *et al.* Network analyses identify liver-specific targets for treating liver diseases. *Molecular systems biology* **13** (2017).
- 9 Bakhtiarzadeh, M. R., Hosseinpour, B., Shahhoseini, M., Korte, A. & Gifani, P. Weighted Gene Co-expression Network Analysis of Endometriosis and Identification of Functional Modules Associated With Its Main Hallmarks. *Frontiers in Genetics* **9**, doi:10.3389/fgene.2018.00453 (2018).
- 10 Mukund, K. & Subramaniam, S. Co-expression network approach reveals functional similarities among diseases affecting human skeletal muscle. *Frontiers in physiology* **8**, 980 (2017).
- 11 Pedrotty, D. M., Morley, M. P. & Cappola, T. P. Transcriptomic biomarkers of cardiovascular disease. *Progress in cardiovascular diseases* **55**, 64-69 (2012).
- 12 Das, S. *et al.* Transcriptomics of cardiac biopsies reveals differences in patients with or without diagnostic parameters for heart failure with preserved ejection fraction. *Scientific reports* **9**, 3179 (2019).
- 13 Ounzain, S. *et al.* Genome-wide profiling of the cardiac transcriptome after myocardial infarction identifies novel heart-specific long non-coding RNAs. *European heart journal* **36**, 353-368 (2014).
- 14 Williams, A. L. *et al.* HIF1 mediates a switch in pyruvate kinase isoforms after myocardial infarction. *Physiological genomics* **50**, 479-494 (2018).
- 15 Priest, C. & Tontonoz, P. Inter-organ cross-talk in metabolic syndrome. *Nature Metabolism*, doi:10.1038/s42255-019-0145-5 (2019).
- 16 Lex, A., Gehlenborg, N., Strobel, H., Vuilleumot, R. & Pfister, H. UpSet: visualization of intersecting sets. *IEEE transactions on visualization and computer graphics* **20**, 1983-1992 (2014).

614 17 Mardinoglu, A. *et al.* The gut microbiota modulates host amino acid and glutathione
615 metabolism in mice. *Molecular Systems Biology* **11**, 834, doi:10.15252/msb.20156487
616 (2015).

617 18 Lima, I., Peres, W., Cruz, S. & Ramalho, A. Association of Ischemic Cardiovascular
618 Disease with Inadequacy of Liver Store of Retinol in Elderly Individuals. *Oxidative*
619 *medicine and cellular longevity* **2018** (2018).

620 19 Palace, V. P., Hill, M. F., Khaper, N. & Singal, P. K. Metabolism of vitamin A in the
621 heart increases after a myocardial infarction. *Free Radical Biology and Medicine* **26**,
622 1501-1507 (1999).

623 20 Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing
624 well-connected communities. *Scientific reports* **9** (2019).

625 21 Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment
626 analysis tool. *BMC Bioinformatics* **14**, 128, doi:10.1186/1471-2105-14-128 (2013).

627 22 Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web
628 server 2016 update. *Nucleic Acids Research* **44**, W90-W97, doi:10.1093/nar/gkw377
629 (2016).

630 23 Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and visualizes
631 long lists of gene ontology terms. *PloS one* **6**, e21800 (2011).

632 24 Su, A. I. *et al.* A gene atlas of the mouse and human protein-encoding transcriptomes.
633 *Proceedings of the National Academy of Sciences of the United States of America* **101**,
634 6062-6067, doi:10.1073/pnas.0400782101 (2004).

635 25 Kiyuna, L. A., e Albuquerque, R. P., Chen, C.-H., Mochly-Rosen, D. & Ferreira, J. C.
636 B. Targeting mitochondrial dysfunction and oxidative stress in heart failure:
637 challenges and opportunities. *Free Radical Biology and Medicine* **129**, 155-168
638 (2018).

639 26 Palaniyandi, S. S., Qi, X., Yogalingam, G., Ferreira, J. C. B. & Mochly-Rosen, D.
640 Regulation of mitochondrial processes: a target for heart failure. *Drug Discovery*
641 *Today: Disease Mechanisms* **7**, e95-e102 (2010).

642 27 Camara, A. K., Zhou, Y., Wen, P.-C., Tajkhorshid, E. & Kwok, W.-M. Mitochondrial
643 VDAC1: a key gatekeeper as potential therapeutic target. *Frontiers in physiology* **8**,
644 460 (2017).

645 28 Schwartz, H. *et al.* Myocardial ischemia/reperfusion causes VDAC phosphorylation
646 which is reduced by cardioprotection with a p38 MAP kinase inhibitor.
647 *PROTEOMICS* **7**, 4579-4588, doi:10.1002/pmic.200700734 (2007).

648 29 Yada, T. *et al.* Cardioprotective role of endogenous hydrogen peroxide during
649 ischemia-reperfusion injury in canine coronary microcirculation in vivo. *American*
650 *Journal of Physiology-Heart and Circulatory Physiology* **291**, H1138-H1146 (2006).

651 30 Sun, W. *et al.* Cardiac-specific deletion of the Pdha1 gene sensitizes heart to
652 toxicological actions of ischemic stress. *Toxicological Sciences* **151**, 193-203 (2016).

653 31 Mardinoglu, A. *et al.* Personal model-assisted identification of NAD⁺ and
654 glutathione metabolism as intervention target in NAFLD. *Molecular systems biology*
655 **13** (2017).

656 32 Alexander, M. *et al.* Non-alcoholic fatty liver disease and risk of incident acute
657 myocardial infarction and stroke: findings from matched cohort study of 18 million
658 European adults. *bmj* **367**, 15367 (2019).

659 33 Tran, D. H. & Wang, Z. V. Glucose Metabolism in Cardiac Hypertrophy and Heart
660 Failure. *Journal of the American Heart Association* **8**, e012673 (2019).

661 34 Kjellman, U. *et al.* Alpha-ketoglutarate for myocardial protection in heart surgery.
662 *Lancet* **345**, 552-553, doi:10.1016/s0140-6736(95)90466-2 (1995).

663 35 Hattori, R. *et al.* Lactate metabolism in acute myocardial infarction and its relation to
664 regional ventricular performance. *J Am Coll Cardiol* **5**, 1283-1291,
665 doi:10.1016/s0735-1097(85)80338-7 (1985).

666 36 Miyamoto, T., Katayama, Y. & Harano, Y. Blood ketone body as a stress marker in
667 acute myocardial infarction. *Nihon Kyokyu Igakukai Zasshi* **10**, 621-622 (1999).

668 37 Zhou, Y. *et al.* Loss of Filamin C Is Catastrophic for Heart Function. *Circulation* **141**,
669 869-871 (2020).

670 38 Hall, C. L. *et al.* RNA sequencing-based transcriptome profiling of cardiac tissue
671 implicates novel putative disease mechanisms in FLNC-associated arrhythmogenic
672 cardiomyopathy. *International journal of cardiology* **302**, 124-130 (2020).

673 39 Diviani, D., Dodge-Kafka, K. L., Li, J. & Kapiloff, M. S. A-kinase anchoring proteins:
674 scaffolding proteins in the heart. *American Journal of Physiology-Heart and*
675 *Circulatory Physiology* **301**, H1742-H1753 (2011).

676 40 Turnham, R. E. & Scott, J. D. Protein kinase A catalytic subunit isoform PRKACA;
677 History, function and physiology. *Gene* **577**, 101-108 (2016).

678 41 Bers, D. M. Calcium cycling and signaling in cardiac myocytes. *Annu. Rev. Physiol.*
679 **70**, 23-49 (2008).

680 42 Zhong, X., Qian, X., Chen, G. & Song, X. The role of galectin-3 in heart failure and
681 cardiovascular disease. *Clinical and Experimental Pharmacology and Physiology* **46**,
682 197-203 (2019).

683 43 Suthahar, N. *et al.* Galectin-3 activation and inhibition in heart failure and
684 cardiovascular disease: an update. *Theranostics* **8**, 593 (2018).

685 44 Ren, J., Pulakat, L., Whaley-Connell, A. & Sowers, J. R. Mitochondrial biogenesis in
686 the metabolic syndrome and cardiovascular disease. *Journal of molecular medicine*
687 **88**, 993-1001 (2010).

688 45 Siasos, G. *et al.* Mitochondria and cardiovascular diseases—from pathophysiology to
689 treatment. *Annals of translational medicine* **6** (2018).

690 46 Piantadosi, C. A. & Suliman, H. B. Transcriptional control of mitochondrial
691 biogenesis and its interface with inflammatory processes. *Biochimica et Biophysica*
692 *Acta (BBA)-General Subjects* **1820**, 532-541 (2012).

693 47 Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-
694 seq quantification. *Nature biotechnology* **34**, 525 (2016).

695 48 Zerbino, D. R. *et al.* Ensembl 2018. *Nucleic Acids Research* **46**, D754-D761,
696 doi:10.1093/nar/gkx1098 (2017).

697 49 Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *Journal of machine*
698 *learning research* **12**, 2825-2830 (2011).

699 50 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
700 dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550,
701 doi:10.1186/s13059-014-0550-8 (2014).

702 51 Uhlén, M. *et al.* The human secretome. *Science Signaling* **12**, eaaz0274,
703 doi:10.1126/scisignal.aaz0274 (2019).

704 52 Våremo, L., Nielsen, J. & Nookaew, I. Enriching the gene set analysis of genome-
705 wide data by incorporating directionality of gene expression and combining statistical
706 hypotheses and methods. *Nucleic Acids Research* **41**, 4378-4391,
707 doi:10.1093/nar/gkt111 (2013).

708 53 Jones, E., Oliphant, T. & Peterson, P. SciPy: Open source scientific tools for Python.
709 (2001).

710 54 Perktold, J., Seabold, S. & Taylor, J. (Oct, 2017).

711 55 Csardi, G. & Nepusz, T. The igraph software package for complex network research.
712 *InterJournal, Complex Systems* **1695**, 1-9, doi:10.5281/zenodo.3630268 (2006).

713 56 Kummitha, C. M., Kalhan, S. C., Saidel, G. M. & Lai, N. Relating tissue/organ energy
714 expenditure to metabolic fluxes in mouse and human: experimental data integrated
715 with mathematical modeling. *Physiological reports* **2** (2014).

716

717

Table

Table 1 Properties of the co-expression network

Tissue	# of Genes	# of Edges	# of Clusters	Modularity Scores
Heart	8793	1570898	7	0.540
Liver	7760	1103589	6	0.577
Muscle	8834	1660603	7	0.521
Adipose	10790	2636378	8	0.495

Figure Supplements

Figure 1 - Figure Supplement 1 Data Exploration of the Samples

Figure 1 - Figure Supplement 2 KEGG pathway analysis results for Heart 6- and 24- hours post MI

Figure 1 - Figure Supplement 3 KEGG pathway analysis results for each tissue Liver, Muscle, and Adipose tissue 24 hours post MI.

Figure 1 - Figure Supplement 4 KEGG pathways related to cardiac problems show activation after an MI.

Figure 4 - Figure Supplement 1 cGMP-PKG with overlay data from differential expression and reporter metabolites analysis.

Figure 4 - Figure Supplement 2 HIF-1 signaling pathway with overlay data from differential expression and reporter metabolites analysis.

Supplementary Files

Supplementary File 1 Differential Expression Analysis Results

- 735 **Supplementary File 2** KEGG Pathways
- 736 **Supplementary File 3** Gene Ontology Biological Processes
- 737 **Supplementary File 4** DEG comparison between Liver and other tissues
- 738 **Supplementary File 5** Reporter Metabolite Analysis
- 739 **Supplementary File 6** Enrichment Analyses of Clusters, Clusters properties
- 740 **Supplementary File 7** Food Intake, Energy Expenditure, and Flux Balance Analysis (FBA)
- 741 of Whole-Body Modeling
- 742 **Supplementary File 8** Validation Result (Differential Expression and Functional Analysis)
- 743 **Supplementary File 9** Detailed Information of 16 Key Genes that are DEGs in at least 3
- 744 tissues and Neighbors and Functional Analysis Results of The Neighbors of 4 key genes

Figure 1 (A) Overview of this study (B) Number of differentially expressed genes for each tissue at each time point. Effect of MI shown to be more pronounced after 24 h. (C) UpSet plot to show intersection between differentially expressed genes (FDR < 5%) in different tissues. The plot showed that each tissue has its specific set of genes that were affected by MI. (D) KEGG pathway analysis (FDR < 0.05 in at least 3 tissues) for 24 hours post MI compared to its control for each tissue. We observed that 141 (5 upregulated) and 125 (14 upregulated) pathways are significantly altered in heart 6 and 24 h after infarction, respectively. For other tissues, we found that 24 (9 upregulated), 61 (54 upregulated) and 48 (15 upregulated) pathways are altered in liver, muscle, and adipose, respectively.

E Reg. of transcription from RNAP-II mRNA Processing Regulation of Mitotic Cell Cycle Phase Ubiquitin-dependent Protein Catabolic Process

Figure 3 Network analyses. (A) Heart co-expression network clusters with superimposed DEGs 24 h post-infarction (Blue = down-regulated, Red = up-regulated) marked with the cluster numbers. The edges between the clusters were aggregation of the inter-cluster edges (B) Liver. (C) Muscle. (D) Adipose. (E) Intersection of the most central clusters in all tissues shows that the central architecture of the network was conserved in all tissues. We found 4 sub-clusters within the network intersection. Top 10 most connected genes are marked in black. (F) Enriched GO BP in heart-specific cluster generated by Revigo.

A

B

Figure 4 (A) Similarity of functions in the most central cluster and specific functions of each tissue-specific cluster. (B) Functional analysis for each tissue and hypothesized flow of metabolites

A

bioRxiv preprint doi: <https://doi.org/10.1101/2020.10.28.358556>; this version posted April 18, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

B

Figure 5 (A) Significantly differentially expressed central genes of each tissue-specific cluster to fatty acid metabolism, as one of the most affected metabolic process. (B) Lipid metabolism. Red = upregulated, blue = downregulated.

Figure 6 (A) DEGs intersection of our data and validation cohort (B) & (C) Intersection of functional analysis results (GO BP and KEGG Pathways) of our data and validation cohort

Figure 1 - Figure Supplement 1
Data Exploration of the Samples

PCA plots of each tissue showing data from mice 6 and 24 h after an MI or sham operation. The plot showed that heart was affected the most by the change in conditions and the rest were most affected by time shifts..

Figure 1 - Figure Supplement 3

KEGG pathway analysis results for each tissue

Liver, Muscle, and Adipose tissue 24 hours post MI.

Figure 1 - Figure Supplement 4

KEGG pathways related to cardiac problems show activation after an MI.

Figure 4 - Figure Supplement 1

cGMP-PKG with overlay data from differential expression and reporter metabolites analysis.

Figure 4 - Figure Supplement 2

HIF-1 signaling pathway with overlay data from differential expression and reporter metabolites analysis.