Spatially resolved analysis of *Pseudomonas aeruginosa* biofilm proteomes measured by laser ablation sample transfer

Aruni Chathurya Pulukkody,¹ Yeni P. Yung,¹ Fabrizio Donnarumma,² Kermit K. Murray²,
Ross P. Carlson,³ and Luke Hanley^{1,*}

¹Department of Chemistry, University of Illinois at Chicago, Chicago, IL 60607

²Department of Chemistry, Louisiana State University, 232 Choppin Hall, Baton Rouge, LA 70803

³Department of Chemical and Biological Engineering, Center for Biofilm Engineering, Montana State University, Bozeman, MT 59717

Abstract

Heterogeneity in the distribution of nutrients and O₂ gradients during biofilm growth gives rise to changes in phenotype. There has been long term interest in identifying spatial differences during biofilm development including clues that identify chemical heterogeneity. Laser ablation sample transfer (LAST) allows site-specific sampling combined with label free proteomics to distinguish radially and axially resolved proteomes for *Pseudomonas aeruginosa* biofilms. Specifically, differential protein abundances on oxic vs. anoxic regions of a biofilm was observed by combining LAST with bottom up proteomics. This study reveals active metabolism in the anoxic region of the biofilm with respect to the oxic region in *P. aeruginosa*, an aerobe by nature. Protein abundance data related to cellular acclimations to chemical gradients include identification of glucose catabolizing proteins, high abundance of proteins from arginine and polyamine metabolism, and proteins that could also support virulence and environmental stress mediation on the anoxic region. Finally, this methodology requires only a few mm² of biofilm area to identify hundreds of proteins.

Emails: apuluk2@uic.edu, yenipyung@gmail.com, fabrizio@lsu.edu, kkmurray@lsu.edu, rossc@montana.edu, and LHanley@uic.edu

I. Introduction

Pseudomonas aeruginosa is a metabolically versatile bacteria that synchronizes expression of genes in a controlled framework such that survival in different environments is matched to manage efficiency and fitness during its growth stages.¹ *P. aeruginosa* can generate bacterial virulence factors that amplify antibiotic resistance, making it an opportunistic pathogen that takes advantage of immunocompromised hosts. Thus, *P. aeruginosa* is prominent within infections that occur in patients with cystic fibrosis, diabetic foot ulcers, burn wounds, and implanted biomaterials.² Such nosocomial infections can transition from acute to chronic stages, making them challenging to treat and in some cases, life threatening.³ The severity of such infections arises in part from antibiotic resistance which causes morbidity and mortality in patients, with over two million of the American population suffering from them per year.^{4,5}

The sessile aggregates or biofilms of *P. aeruginosa* have been widely studied and chemical and physical heterogeneity along radial and axial (vertical) axes are common.⁶ Chemical heterogeneity in *P. aeruginosa* biofilms has been measured axially for compounds like O₂ using microelectrode technology,^{7,8} while metabolic activity has been measured using reporter proteins like green fluorescent protein^{6,9,10} and transcriptome.^{6,9} Proteomic analyses of whole biofilms have also been performed previously.^{8, 11-13} However, there have been relatively few studies focused on spatially resolved *P. aeruginosa* biofilm proteomics.^{6,14,15}

Mass spectrometry (MS) imaging studies the spatial distribution of multiple analytes ranging from small molecules to lipids, peptides and proteins by creating a two-dimensional heat map based on the abundance and distribution of representative ions.¹⁵ Matrix assisted laser desorption ionization (MALDI) is the most popular ionization mode for MS imaging and most commonly employs nanosecond (ns), ultraviolet laser pulses to desorb biomolecules from a

chemical matrix-coated sample.^{16,17} Improvements in spatial resolution and sensitivity in MALDI-MS imaging have been achieved using high mass resolution/accuracy MS instrumentation, methodologies to spray thin and even coatings of the matrix, new matrices, and better laser spot focusing.^{15,17-19} Another popular method in MS imaging crosses the ablation plume formed by a mid-infrared ns laser pulse with the output from an electrospray ionization source (sometimes referred to as laser ablation electrospray ionization or LAESI).²⁰ Laser wavelengths near $\sim 3 \mu\text{m}$ are resonant with the OH stretch vibrations of water, facilitating explosive plume-like ablation from biofilms and other hydrated biological samples.²⁰ MALDI, LAESI, and other popular methods of MS imaging like desorption electrospray ionization (DESI)²¹ typically lack the chromatographic separation that is so effective for MS analysis of chemically complex samples such as bacterial biofilms.

Localized sampling of biological surfaces can be accomplished in several ways beyond the use of focused probes. Manual dissection²²⁻²⁴ can remove a portion of the surface from the bulk material, while laser capture microdissection²⁵⁻²⁷ allows more precise localized sampling. Liquid microjunctions²⁸ have also been used to achieve extraction of specific analytes from a biological surface through tuning the composition of the extraction mixture. However, it is either difficult or impossible to perform a layered extraction or measure a depth profile using these methods.

Laser ablation sample transfer (LAST) also utilizes a mid-infrared ns-pulsed laser to ablate hydrated biological samples,^{29,30} like LAESI. Murray and co-workers have been able to demonstrate multiple uses of this method on a diversity of samples including mouse brain, mouse lung tissue, peptide and protein standards, fingerprints, ink, enzymes and lipids on bacterial colonies of *E. coli* and *B. cereus* grown liquid media.²⁹⁻³³ The LAST method can be used as a field sampling or offline technique,³⁴ where collected material can be further processed and analyzed

separately. LAST also permits spatially resolved sampling and can be sequentially combined with liquid chromatographic (LC) separation and tandem mass spectrometry (MS/MS) for bottom-up proteomic analysis of biomolecules in complex biological samples. Selection of the laser fluence and irradiation geometry enables different ablation modalities. Higher laser fluence irradiation in reflection geometry (front side) can ablate the entire irradiated biofilm volume. Lower laser fluence irradiation in transmission geometry (back side of the glass slide, through translucent biofilm) induces laser spallation^{35,36} that leads to ablation of only the outmost layer of the biofilm (see Figure S1b in Supplementary Information). Spallation is thought to proceed via a pressure wave propagating from the point of irradiation along the longitudinal axis of the biofilm.^{35,36}

LAST is demonstrated here to be effective for region-specific proteomics analysis of bacterial biofilms by coupling it with bottom-up proteomics via LC and high-resolution Orbitrap MS/MS. The method is found here to be particularly sensitive in that it requires only a few mm² area of a biofilm to identify hundreds of proteins. This work combines current advances in bottom-up proteomics to spatially resolve bacterial proteomes within *P. aeruginosa* biofilms, further revealing the remarkable molecular machinery involved in their acclimation to environmental conditions.

II. Experimental Section

A. LAST. The laser ablation sample transfer setup was constructed in-house based upon a published design of Murray and Donnarumma³² (see Supplementary Information, Figure S1a). 2.94 μm , ~7 ns laser pulses emitted at a 20 Hz repetition rate from an Nd:YAG-pumped optical parametric oscillator (IR Opolette, Opotek, Carlsbad, CA) were transmitted through an iris onto a 45° angle gold mirror. The reflected laser beam was focused normal to the sample surface through a 50 mm CaF₂ lens in transmission geometry onto biofilm samples adhered to glass microscope

slides. Inverted sample slides were placed on a xyz translation stage operated using 25 mm motorized linear actuators (Model 462, CMA-25CCCL, Newport, Irvine, CA, USA), run using a previously described software.³⁷ Biofilm samples were ablated while the sample stage moved at a speed of 0.2 mm/s with the laser irradiation perpendicular to the sample surface. The spot size of the focused beam was 260 μm with a laser fluence of $4.6 \pm 2.1 \text{ J/cm}^2$. The laser spot size was determined by ablation of agar gels, followed by examination of the gels using a digital camera (DFC 495, Leica Microsystems Inc., Buffalo Grove, IL), where $\sim 50 \mu\text{m}$ thick agar gels were prepared from 1.5 g of agar powder dissolved in 5.0 mL of water, coated on glass microscope slides and air-dried. Ablation followed the perimeter of 4 mm square of a biofilm, requiring a total of ~ 1500 laser shots. A microcentrifuge tube with buffered solution (10 mM NH_4HCO_3 , pH 8.0) was used to collect the ablated material.

B. Growth and ablation of biofilms. A glycerol stock of *P. aeruginosa* chronic wound clinical isolate strain 215 was used to inoculate a CSP-agar (chemically defined nutrient media for bacterial growth) streak plate, as described previously.⁸ The streak plate inoculum was incubated for 16 hours at 37 °C, then used to establish fresh colonies of *P. aeruginosa* in planktonic culture (10 mL of CSP liquid media). This planktonic culture was grown until it attained optical density of 0.1 at 600 nm absorbance (OD_{600}).

Planktonic cultures of *P. aeruginosa* at $\text{OD}_{600} = 0.1$ (20 μL) were inoculated onto sterile polycarbonate membranes (25 mm diam, 0.2 μm pore size from GE Life Sciences, Marlborough, MA), then were grown for three days on chemically-defined CSP medium introduced into solid agar. Biofilms supported on membranes were transferred onto fresh media plates every 24 hours.⁸ These polycarbonate membrane biofilms were used to develop and evaluate the analytical methodology and to measure the radially resolved proteome.

Biofilms were also grown from $OD_{600} = 0.1$ aliquots of planktonic culture which were first diluted to $OD_{600} \approx 0.03$. Transwell cell culture plate inserts (0.4 μm pore size, Corning Inc. Tewksbury, PA) were placed in six well plates with 2.5 mL of CSP media. Membranes were inoculated with 100 μL of diluted *P. aeruginosa* culture and grown for three days with agitation. The transwell inserts were transferred every 12 hours to fresh CSP media in new cell culture plates. A scalpel blade was used to separate the transwell membrane from the plastic insert that was then inverted onto a microscope slide. Next, the glass slide then placed in a secondary container and stored in dry ice for 15 min, then the membrane was peeled off while still frozen to expose its anoxic side (see Figure S2), and finally thawed prior to laser irradiation for examination of the axially resolved proteome. A similar process was used to expose the oxic surface via a second membrane, as shown in Figure 1.

C. Proteomic analyses. The basic LC-MS/MS analysis strategy as well as the qualitative and quantitative proteomics were adapted from a method that was described in detail previously⁸ and is only summarized here. Laser ablated material from the polycarbonate membrane or transwell membrane insert grown biofilms was captured in a buffered solution then concentrated by vacuum centrifugation (Labconco Corp., Kansas City, MO) at 30°C and 1725 rpm. The residues were reconstituted in 20 μL of Laemmli buffer with 5% β -mercaptoethanol to reduce disulfide bonds, then heated to 90°C for complete denaturation of proteins for sodium dodecyl sulphate - polyacrylamide gel electrophoresis (SDS-PAGE) analysis. The bands on each well represents the separation of proteins from the ablated material.

Each lane in the SDS gel was excised and processed with an in-gel digestion protocol.³⁸ The processing of entire lanes was done to maintain consistency with the SDS-PAGE results. The resultant tryptic digested peptides were reconstituted in 10 μL of 5% acetonitrile, 0.1% formic acid

buffer for LC-MS/MS analysis. Difficulties arose in quantifying the relatively low levels of protein by a Bradford assay.^{43,44} Peptide concentrations for LC-MS/MS injection from the transwell membranes were instead measured using 1 μ L aliquots of each sample on a micro-volume UV-visible spectrophotometer (Nano Drop One, Thermo Fisher Scientific, Wilmington, DE),³⁹ where the absorbance ratios of 280 nm to 205 nm (due to amide bonds of proteins) were used to determine the volume of injection corresponding to an equivalent amount of protein for normalization across biological replicates. This strategy reduced interference in UV absorbance from detergents and also reduced bias arising from consideration of only aromatic peptides.

Approximately 200 ng of digested peptides resuspensions were separated through nano-LC (1260 Infinity LC System, Agilent, Santa Clara, CA) with a C18 column (Agilent, Zorbax 300SB, and column dimensions of 150 mm length $\times$ 75 μ m internal diameter, with 3.5 μ m pore size particles) and a 90-min increasing organic gradient from 7% to 95%. The aqueous phase was 0.1% formic acid in water and the organic phase was 0.1% formic acid and 80% acetonitrile in water, leading to a total run time of 120 min at a flow rate of 300 nL/min.

The chromatographically separated peptides were then electrosprayed at 1.90 kV spray voltage and 275 °C capillary temperature. Peptide ions were analyzed in a high mass resolution tandem mass spectrometer (Orbitrap Velos Pro, Thermo Fisher Scientific, Waltham, MA) with automatic gain control set at 10^6 ions and an injection time between 1 and 200 ms. Full-scan mass spectra were collected in data-dependent acquisition mode from m/z 400 - 2000 at 30,000 mass resolution. Ten precursor ions were selected from each full mass spectrum for MS/MS analyses in high energy collision-induced dissociation mode using 30% energy for fragmentation. Label free quantitation was employed by the use of ion intensities as a direct (albeit semi-quantitative) measure of protein abundance.

The Perseus⁴⁰ and MaxQuant⁴¹ computational platforms were used for protein identification using the UniProt Knowledgebase⁴² and statistical analysis via label free quantification (LFQ), generally following the previously described procedure which is summarized here.⁸ Mass spectra data (.raw) files were directly uploaded and processed using a FASTA file of reviewed *P. aeruginosa* proteins (taxonomy identification 287) downloaded from the UniProt Knowledgebase database source and used for protein identification via MaxQuant. The main search parameters for identification were set at peptide tolerance of 4.5 ppm, MS/MS match tolerance of 20 ppm, MS/MS *de novo* tolerance of 10 ppm, minimum peptide length of 7, false discovery rate of 0.01, carbamidomethyl fixed modification, oxidation and acetylation variable modifications, and enabled contaminant search. An unpaired Students' t-test was used, with permutation based false discovery rate accepting values $\leq 5\%$, in order to reduce identifications of false positives. The final set of identified peptides was further reduced in a spreadsheet by application of two additional selection criteria: $\geq 10\%$ sequence coverage and at least two unique peptides.

The ablated material from three biological replicates of transwell membrane biofilms were analyzed separately by LC-MS/MS and the output averaged to represent the axially-resolved proteome. The proteins from oxic vs. anoxic regions were compared to identify statistically significant ($q\text{-value} \leq 0.05$, and a corresponding adjusted $p\text{-value}$) high and low abundance proteins with a ≥ 1.5 -fold change, referred to as up- or downregulated, respectively. However, no statistics are provided for the one replicate analysis for each region by LC-MS/MS of polycarbonate membrane grown biofilms. The resultant radially-resolved proteome is only preliminary, but is reported here nonetheless in case it might be of use to future studies.

III. Results and Discussion

A. Verification of proteins ablated from polycarbonate membrane biofilms. Preliminary studies required identification of the presence of ablated protein in the material laser ablated from the polycarbonate membrane-grown biofilms. SDS-PAGE was performed on ablated cell lysates from total areas of 6, 12, 18 and 24 mm². These areas correspond the perimeters of 1 - 4 mm² squares ablated from the same position on distinct biofilms, then multiplied by six to account for the pooling of sample collected from six separate biofilms. The images of the protein bands on a polyacrylamide gel from each lane from the pooled samples are shown in Figure 2a and confirm the ability of LAST to sample small areas of *P. aeruginosa* biofilms. Clearly distinguishable bands were observed on lanes resulting from at least 12 mm² of total biofilm area. These 12 mm² lanes subsequently yielded a considerable number of protein identifications via LC-MS/MS, as discuss below. However, 24 mm² of ablated material was designated as the minimum required for the oxic vs. anoxic analyses by LC-MS/MS to increase protein coverage.

B. Preliminary radially-resolved proteome from *P. aeruginosa* biofilms. The biofilms evaluated for their radially-resolved proteome were grown for three days on polycarbonate membranes placed on CSP containing agar plates. This biofilm cultivation method establishes opposing concentration gradients of O₂ and medium components similar to chronic wounds and was methodologically convenient for generating the multiple biofilms necessary for development of the methods described herein.⁴⁵⁻⁴⁷ Furthermore, growth on polycarbonate membranes on agar plates was analogous to the biofilm growth on membranes in tissue culture wells.^{8,14}

Radially-resolved proteomic analyses were performed by collecting ablated material from two radially distinct regions three day old polycarbonate membrane-grown biofilms: a central biofilm region which corresponded with older cells (brown square on Figure 3a) and outer region

of the biofilm which corresponded with younger cells (green square on Figure 3b) as biofilms grew radially outward from the central inoculation site. Figure 3a shows schematically that ablation was actually performed in the shape of concentric squares, a pattern that was readily available to the motion stage while minimizing the age distribution of cells sampled. Figure 3b shows an optical image of post-ablation markings on an actual biofilm.

LC-MS/MS analysis of the ablated material from these preliminary samples resulted in identification of ~400 proteins, but only ~200 of them satisfied the more stringent criteria of more than two unique peptides and $\geq 10\%$ sequence coverage. Analysis of single samples grouped from six biofilms found 86 proteins unique to the newer cell region (24 mm² total area) and fewer proteins unique to the older cell region, as expected given that the 12 mm² was only half the total area. The analyses carried out on radial ablations from biofilms on polycarbonate membranes were performed mainly to support method validation, but these preliminary protein identities are documented in Supplementary Information (see Table S1). All further work was focused on the younger cell region of the transwell membrane-grown biofilms.

C. Oxic and anoxic regions of *P. aeruginosa* biofilms have major differences in proteome.

Examination of Figure 2b indicates that laser ablation of anoxic vs. oxic regions of biofilms led to no visually apparent differences in the gel bands from SDS-PAGE, whereas bottom-up proteomics by LC-MS/MS showed clear differences. 24 mm² square perimeters were laser ablated from three anoxic or oxic regions of six biofilms grown for three days on transwell membranes from the same culture. Proteins from each sample was separated via SDS-PAGE, then the proteins were digested on the gel and all bands of the lane were extracted and the resultant peptides injected into the LC-MS/MS. All proteins identified by this procedure are tabulated in Table S2 of the Supplementary Information along with their three or four-letter codes, full protein names, the number of unique

peptides identified, the sequence coverage, the differential abundance for anoxic vs. oxic regions, and statistical significance.

LAST combined with LC-MS/MS revealed proteins with statistically significant different abundances between anoxic and oxic regions of these transwell biofilms. The volcano plot in Figure 4 includes 250 proteins identified that satisfied the stringency conditions of at least two peptides and $\geq 10\%$ sequence coverage, out of a total of 430 proteins identified. An oxygen-selective electrode distinguished the oxic from anoxic regions in biofilms of this same strain grown in the same medium and cultivation system (albeit for a shorter growth period),⁸ but the proteins observed therein were averaged across the entire biofilm. The results presented here show that the bottom, anoxic region of the biofilm had the majority of statistically significant increases in protein abundance which was unexpected as *P. aeruginosa* relies largely on respiration for energy generation. The CSP medium did not contain alternative electron acceptors like nitrate constraining respiration in the lower regions of the biofilm. The bottom, anoxic portion of the *P. aeruginosa* biofilm had 47 higher abundance proteins relative to the top of the biofilm, while the oxic region had only 14 higher abundance proteins relative to the bottom. The observed bottom vs. top differences in the proteins support the argument that LAST distinguishes anoxic from oxic regions. However, these experiments cannot rule out the possibility that cells from the two regions are at least partially mixed in the native biofilms or as a result the ablation process. This was in stark contrast to previous analyses of vertical differences in transcriptome of *P. aeruginosa* biofilms which indicated essentially all transcripts were higher in the upper portion of the biofilm.^{6,14}

Ribosomal proteins (RpsS, RpsK, RpmG RpsM, RpsH and RplR) were down regulated by two to threefold in the anoxic zone, consistent with the widely held hypothesis of minimal

growth in anoxic zones. However, up regulation of many metabolic proteins suggested the cells were generating cellular energy and acclimating to stressful chemical environments.⁵⁰ The anoxic region of the biofilm had a roughly twofold up regulation of glucose catabolizing proteins (Edd, PckA, and Pck) as compared to the oxic region. *P. aeruginosa* catabolizes glucose using the Entner-Doudoroff glycolysis pathway (Edd, Eda) and the data suggest cells in the anoxic regions were catabolizing the sugar and likely producing byproducts like acetate. Cells in the oxic region of the biofilm had an almost fourfold up regulation for the key acetate metabolic protein, AceA, from the glyoxylate shunt, suggesting either acetate from the anoxic region may be metabolized in the oxic zone or that there is mixing of cells from the two regions. Up regulation of AceA implies that the oxic zone might play a dominant role in the carbon catabolite strategy known as reverse diauxie which prefers organic acids to glucose and which was observed previously in *P. aeruginosa* biofilms.⁵¹

The cells in the anoxic region also displayed a fivefold up regulation of proteins associated with adenosine triphosphate (ATP) synthase proteins (AtpD and AtpG). The data suggest there may be more than substrate level phosphorylation occurring as operating an oxidative tricarboxylic acid (TCA) cycle and ATP synthase requires external electron acceptors. The soluble transhydrogenase protein (SthA) was also up regulated by twofold in the anoxic region potentially linking the conversion of anabolic and catabolic electrons. The exchange of redox active metabolites could enable respiration at a distance using small molecules like phenazines, although proteins associated with these processes were not identified.^{52,53}

Cells in the anoxic portion of the biofilm displayed up regulation of proteins from arginine and polyamine metabolism. CSP medium contained arginine and it would be preferentially available to cells on the bottom of the biofilm. Arginine is an important amino acid for *P.*

aeruginosa acclimation to biofilm conditions and can be used for substrate level energy generation, pH modulation via deamination (ArcA) and decarboxylase reactions (SpeA).^{54,55} Arginine also serves as a precursor for synthesis of putrescine and spermidine which stabilize cellular membranes to stress.^{54,55} Multiple arginine proteins were associated with the arginine succinyltransferase (AST), arginine decarboxylase/agmatine deiminase pathway, and arginine biosynthesis including the AnsB and ArcA proteins up regulated by almost eightfold in the anoxic region; ArgC, ArgG, ArgJ, and SpuD, by fivefold; AruC and GdhB by fourfold; and ArgS and ArcB by twofold. Additionally, cytosolic aminopeptidase protein (PepA) was found to be up regulated by threefold in the anoxic regions and could be degrading cellular proteins to repurpose the amino acids for new proteins, stress responses like pH mediation, or cellular energy synthesis.

A large number of higher abundance proteins in the anoxic region could also support virulence and environmental stress mediation. Some of these proteins include large-conductance mechanosensitive ion channels (MscL, up regulated fivefold in the anoxic region) a type of protein transport system which is sensitive to mechanical perturbations and changes in osmotic pressure.⁵⁶ Multidrug resistance protein MexA was up regulated by roughly twofold in the anoxic region consistent with previous transcript analyses of anoxic cultivation which indicate *P. aeruginosa* is more tolerant to antibiotics in hypoxic environments.⁵⁷ Membrane protein insertase (YidC) is a ubiquitous protein from the Sec, general secretory pathway and aids in the translocation of proteins across the cytoplasmic membrane. It functions together with other proteins to promote insertion of membrane proteins.⁵⁸ It is also responsible for modulating the expression of hypoxic genes and when present can aid in the assembly of ATP synthases (AtpD, AtpG, and AtpA, all identified in this study). YidC also causes a roughly tenfold down regulation of some ribosomal proteins (RplY and RplV), in the oxic region of the biofilm.⁵⁹ DNA-binding protein (HupB) was up regulated

eightfold in the anoxic region. HupB are histone-like small proteins - also known as heat unstable (HU) proteins - that prevent DNA damage caused by oxidative stress and enable adaptation to stress response.⁶⁰ HU proteins can also control the transcription of a gene required for anoxic respiration.⁶¹ The chaperone protein ClpB was upregulated by twofold in the anoxic region and can rescue stress damaged proteins in an ATP consuming process.⁶² The energy requirement to maintain protein function under environmental stress may partially explain the apparent substantial energy metabolism occurring in the anoxic region of the biofilm. Finally, the extracellular polymeric substance (EPS) producing protein, AlgC, was up regulated threefold in the anoxic zone, possibly driving resource flux toward additional EPS synthesis.⁶³

IV. Conclusions

This study demonstrated several important advantages of laser ablation sample transfer for regio-specific determination of a biofilm proteome. Prior work by some of the authors required the harvest of entire *P. aeruginosa* bacterial biofilms of several cm² area to achieve an average characterization of phenotype via proteomics.⁸ However, that prior work did not distinguish the changes in the proteome that occurred as O₂ abundance varied with depth.⁸ LAST is shown here to distinguish protein response in the anoxic vs. oxic regions selected by the flipping of the biofilm as depicted in Figures 1 and S2, determining a phenotypic difference between the two regions. LAST also allows regio-specific sampling across the surface of biofilm by the use of an ~260 μm laser irradiation spot diameter.

There are other experimental advantages of the LAST strategy. Ablated material can be concentrated via vacuum centrifugation, then stored at -80°C to prevent deterioration until analysis can be conveniently performed. Sample processing for analysis is carried out off-line by standard LC-MS/MS-based bottom-up proteomics. Furthermore, the version of the LAST setup deployed

here is on wheels, allowing it to be moved into a bio-safety cabinet when used with pathogenic bacteria. Furthermore, the setup is remotely operated, avoiding exposure of the experimenter to ablated pathogenic aerosols.

Rapid dehydration and flaking of the biofilm can occur when the laser beam directly impacts a surface in reflection geometry, leading to a disruption of the water-based laser ablation process.⁶⁴ This problem was addressed here by inverting the microscope glass slide on which the sample rests so that the laser radiation penetrated through the glass. Transmission geometry is also advantageous for heat diffusion,²⁹ minimizing the potential for thermal degradation of the biofilm. Furthermore, the sticky and mucous-like texture of the biofilm can minimize the collection of ablated material via reflection geometry, while the transmission geometry assists collection of ablated material via spallation.

There are considerable advantages to LAST compared to direct MS imaging techniques such as MALDI^{15,17-19,65} or DESI,²¹ despite LAST's lower lateral resolution. The delocalization introduced by the matrix⁶⁶ or the solvent spray⁶⁷ and the less efficient desorption of large biomolecules⁶⁸ in direct MS imaging methods are eliminated using LAST. Direct MS imaging techniques have been combined with advanced instrumentation such as ion mobility and Fourier transform ion cyclotron resonance MS^{15,69,70} to simultaneously achieve high protein coverage and lateral resolution, but such strategies are still not widely available. By contrast, a general LC-MS/MS workflow combined with LAST provides a significantly high number of protein identities using relatively low cost and widely accessible instrumentation, albeit at a relatively low lateral resolution.

In comparison to previous work that required several entire biofilms for proteomic analyses,⁸ only a small region of the biofilm was needed for LAST. About 430 proteins were

identified here by laser ablation of the surface region of a 24 mm² area of biofilm. This is comparable to the ~600 proteins identified from analysis of the cell lysate from three entire biofilms (of several cm² total area, where the entire thickness of the biofilm was required).⁸ Loosening of the protein identification criteria (i.e., the 10% sequence matching) would have considerably increased the number of protein identifications.

There has been long term interest in identifying spatial differences during biofilm development including clues that identify chemical heterogeneity therein.⁹ The novel results presented here develop the radial and axial resolved proteome in important biofilm-forming bacteria, *P. aeruginosa*, which provides key insight to metabolic activity in the slow growing anoxic zone that was not apparent from previous transcript studies. An extension of this work to multi-species biofilms will provide insights to synchronize between the omics data to highlight the synergy in biofilms that are ubiquitous in the environment.

Data Availability Statement

The proteomic data that support the findings of this study are available on the Mass Spectrometry Interactive Virtual Environment (MassIVE) repository at <ftp://massive.ucsd.edu/MSV000086320/>.

Acknowledgements

This work was supported by the National Institute of Biomedical Imaging and Bioengineering under grant 1 U01 EB019416. Purchase of the liquid chromatograph and Orbitrap mass spectrometer was funded by the National Center for Research Resources under grant 1 S10 RR025653-01A1.

Figure Captions

Figure 1. Technique of exposing the oxic surface of a biofilm for LAST by using a second membrane to aid in the inversion of the biofilm.

Figure 2. a) Preliminary identification of the presence of proteins observed by SDS-PAGE performed on cell lysates for a range of laser ablated areas from biofilms grown on polycarbonate membranes. b) SDS-PAGE of cell lysates from 24 mm² area regions that were laser ablated from anoxic and oxic regions of biofilms grown on transwell membranes.

Figure 3. a) Schematic of concentric ablation squares 1 mm (older cell region, in brown) and 2 mm (younger cell region, in green) from the center of the biofilm, ablation of polycarbonate biofilms carried used to evaluate regio-specific methodology, b) optical image of post-ablation marks on biofilm.

Figure 4. Differential protein abundances for anoxic vs. oxic regions of *P. aeruginosa* three day grown transwell biofilms. Proteins with statistically significant ($-\log_{10}(\text{p-value}) > 1.3$) changes in abundance are highlighted above horizontal black line. Of those, proteins with $|\log_2(X_{\text{anoxic}}/X_{\text{oxic}})| \geq 0.58$, where $X_{(\text{an})\text{oxic}}$ is protein abundance, are deemed as significantly up or down regulated and are annotated as red or blue points, respectively. The ten proteins that displayed the greatest fold changes in up or down regulation are labelled with their three or four-letter protein codes (see Table S2 of Supplementary Information).

References

- (1) Rojo, F. Carbon catabolite repression in *pseudomonas* : Optimizing metabolic versatility and interactions with the environment. *FEMS Microbiology Reviews* **2010**, *34*, 658-684.
- (2) Ma, Y.-X.; Wang, C.-Y.; Li, Y.-Y.; Li, J.; Wan, Q.-Q.; Chen, J.-H.; Tay, F. R.; Niu, L.-N. Considerations and caveats in combating escape pathogens against nosocomial infections. *Advanced Science* **2020**, *7*, 1901872.
- (3) Obritsch, M. D.; Fish, D. N.; MacLaren, R.; Jung, R. Nosocomial infections due to multidrug-resistant *Pseudomonas aeruginosa*: Epidemiology and treatment options. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* **2005**, *25*, 1353-1364.
- (4) Mulcahy, L. R.; Isabella, V. M.; Lewis, K. *Pseudomonas aeruginosa* biofilms in disease. *Microbial Ecology* **2014**, *68*, 1-12.
- (5) Center for Disease Control, Antibiotic resistance threats in the United States, U.S. Department of Health and Human Services: Atlanta, GA, 2019, <https://www.cdc.gov/drugresistance/pdf/threats-report/2019-ar-threats-report-508.pdf>
- (6) Williamson, K. S.; Richards, L. A.; Perez-Osorio, A. C.; Pitts, B.; McInnerney, K.; Stewart, P. S.; Franklin, M. J. Heterogeneity in *Pseudomonas aeruginosa* biofilms includes expression of ribosome hibernation factors in the antibiotic-tolerant subpopulation and hypoxia-induced stress response in the metabolically active population. *Journal of Bacteriology* **2012**, *194*, 2062-2073.
- (7) Rani, S. A.; Pitts, B.; Beyenal, H.; Veluchamy, R. A.; Lewandowski, Z.; Davison, W. M.; Buckingham-Meyer, K.; Stewart, P. S. Spatial patterns of DNA replication, protein synthesis, and oxygen concentration within bacterial biofilms reveal diverse physiological states. *Journal of Bacteriology* **2007**, *189*, 4223-4233.
- (8) Yung, Y. P.; McGill, S. L.; Chen, H.; Park, H.; Carlson, R. P.; Hanley, L. Reverse diauxie phenotype in *Pseudomonas aeruginosa* biofilm revealed by exometabolomics and label-free proteomics. *npj Biofilms and Microbiomes* **2019**, *5*, 31.
- (9) Stewart, P. S.; Franklin, M. J. Physiological heterogeneity in biofilms. *Nature Reviews Microbiology* **2008**, *6*, 199-210.
- (10) Folsom, J. P.; Richards, L.; Pitts, B.; Roe, F.; Ehrlich, G. D.; Parker, A.; Mazurie, A.; Stewart, P. S. Physiology of *Pseudomonas aeruginosa* in biofilms as revealed by transcriptome analysis. *BMC Microbiology* **2010**, *10*, 294.
- (11) Yoon, S. S.; Hennigan, R. F.; Hilliard, G. M.; Ochsner, U. A.; Parvatiyar, K.; Kamani, M. C.; Allen, H. L.; DeKievit, T. R.; Gardner, P. R.; Schwab, U.; Rowe, J. J.; Iglewski, B. H.; McDermott, T. R.; Mason, R. P.; Wozniak, D. J.; Hancock, R. E. W.; Parsek, M. R.; Noah, T. L.; Boucher, R. C.; Hassett, D. J. *Pseudomonas aeruginosa* anaerobic

- respiration in biofilms: Relationships to cystic fibrosis pathogenesis. *Developmental Cell* **2002**, 3, 593-603.
- (12) Guina, T.; Purvine, S. O.; Yi, E. C.; Eng, J.; Goodlett, D. R.; Aebersold, R.; Miller, S. I. Quantitative proteomic analysis indicates increased synthesis of a quinolone by *Pseudomonas aeruginosa* isolates from cystic fibrosis airways. *Proceedings of the National Academy of Sciences U.S.A.* **2003**, 100, 2771-2776.
- (13) Patrauchan, M. A.; Sarkisova, S. A.; Franklin, M. J. Strain-specific proteome responses of *Pseudomonas aeruginosa* to biofilm-associated growth and to calcium. *Microbiology* **2007**, 153, 3838-3851.
- (14) Stewart, P. S. A review of experimental measurements of effective diffusive permeabilities and effective diffusion coefficients in biofilms. *Biotechnology and Bioengineering* **1998**, 59, 261-272.
- (15) Wakeman, C. A.; Moore, J. L.; Noto, M. J.; Zhang, Y.; Singleton, M. D.; Prentice, B. M.; Gilston, B. A.; Doster, R. S.; Gaddy, J. A.; Chazin, W. J.; Caprioli, R. M.; Skaar, E. P. The innate immune protein calprotectin promotes *Pseudomonas aeruginosa* and *Staphylococcus aureus* interaction. *Nature Communications* **2016**, 7, 11951.
- (16) Palmer, A.; Trede, D.; Alexandrov, T. Where imaging mass spectrometry stands: Here are the numbers. *Metabolomics* **2016**, 12, 107.
- (17) Murray, K. K.; Seneviratne, C. A.; Ghorai, S. High resolution laser mass spectrometry bioimaging. *Methods* **2016**, 104, 118-126.
- (18) Kompauer, M.; Heiles, S.; Spengler, B. Autofocusing MALDI mass spectrometry imaging of tissue sections and 3D chemical topography of nonflat surfaces. *Nature Methods* **2017**, 14, 1156-1158.
- (19) Bednařík, A.; Machálková, M.; Moskovets, E.; Coufalíková, K.; Krásenský, P.; Houška, P.; Kroupa, J.; Navrátilová, J.; Šmarda, J.; Preisler, J. MALDI MS imaging at acquisition rates exceeding 100 pixels per second. *Journal of the American Society for Mass Spectrometry* **2019**, 30, 289-298.
- (20) Flanigan, P.; Levis, R. Ambient femtosecond laser vaporization and nanosecond laser desorption electrospray ionization mass spectrometry. *Annual Review of Analytical Chemistry* **2014**, 7, 229-256.
- (21) a) Wu, C.; Dill, A. L.; Eberlin, L. S.; Cooks, R. G.; Ifa, D. R. Mass spectrometry imaging under ambient conditions. *Mass Spectrometry Reviews* **2013**, 32, 218-243. b) Honarvar, E.; Venter, A. R. Comparing the effects of additives on protein analysis between desorption electrospray (DESI) and electrospray ionization (ESI), *Journal of the American Society for Mass Spectrometry* **2018**, 29, 2443-2455.

- (22) Franck, J.; Quanico, J.; Wisztorski, M.; Day, R.; Salzet, M.; Fournier, I. Quantification-based mass spectrometry imaging of proteins by parafilm assisted microdissection. *Analytical Chemistry* **2013**, *85*, 8127-8134.
- (23) Quanico, J.; Franck, J.; Gimeno, J. P.; Sabbagh, R.; Salzet, M.; Day, R.; Fournier, I. Parafilm-assisted microdissection: A sampling method for mass spectrometry-based identification of differentially expressed prostate cancer protein biomarkers. *Chemical Communications* **2015**, *51*, 4564-4567.
- (24) Quanico, J.; Franck, J.; Wisztorski, M.; Salzet, M.; Fournier, I. Integrated mass spectrometry imaging and omics workflows on the same tissue section using grid-aided, parafilm-assisted microdissection. *Biochimica et Biophysica Acta (BBA) - General Subjects* **2017**, *1861*, 1702-1714.
- (25) Simone, N. L.; Bonner, R. F.; Gillespie, J. W.; Emmert-Buck, M. R.; Liotta, L. A. Laser-capture microdissection: Opening the microscopic frontier to molecular analysis. *Trends in Genetics* **1998**, *14*, 272-276.
- (26) Legres, L. G.; Janin, A.; Masselon, C.; Bertheau, P. Beyond laser microdissection technology: Follow the yellow brick road for cancer research. *American journal of cancer research* **2014**, *4*, 1-28.
- (27) Mahalingam, M. Laser capture microdissection: Insights into methods and applications in *Laser capture microdissection: Methods and protocols*, Murray, G. I., Ed.; Springer New York: New York, NY, 2018, pp 1-17.
- (28) Quanico, J.; Franck, J.; Dauly, C.; Strupat, K.; Dupuy, J.; Day, R.; Salzet, M.; Fournier, I.; Wisztorski, M. Development of liquid microjunction extraction strategy for improving protein identification from tissue sections. *Journal of Proteomics* **2013**, *79*, 200-218.
- (29) Park, S.-G.; Murray, K. K. Infrared laser ablation sample transfer for MALDI imaging. *Analytical Chemistry* **2012**, *84*, 3240-3245.
- (30) Park, S.-G.; Murray, K. K. Infrared laser ablation sample transfer for on-line liquid chromatography electrospray ionization mass spectrometry. *Journal of Mass Spectrometry* **2012**, *47*, 1322-1326.
- (31) Donnarumma, F.; Cao, F.; Murray, K. Laser ablation with vacuum capture for MALDI mass spectrometry of tissue. *Journal of the American Society for Mass Spectrometry* **2016**, *27*, 108-116.
- (32) Donnarumma, F.; Murray, K. K. Laser ablation sample transfer for localized LC-MS/MS proteomic analysis of tissue. *Journal of Mass Spectrometry* **2016**, *51*, 261-268.
- (33) Wang, K.; Donnarumma, F.; Baldone, M. D.; Murray, K. K. Infrared laser ablation and capture of enzymes with conserved activity. *Analytica Chimica Acta* **2018**, *1027*, 41-46.

- (34) Hayes, J. M.; Murray, K. K. Ambient laser ablation sample transfer with nanostructure-assisted laser desorption ionization mass spectrometry for bacteria analysis. *Rapid Communications in Mass Spectrometry* **2014**, *28*, 2382-2384.
- (35) Vogel, A.; Venugopalan, V. Mechanisms of pulsed laser ablation of biological tissues. *Chemical Reviews* **2003**, *103*, 577-644.
- (36) Zhidkov, A. G.; Zhigilei, L. V.; Sasaki, A.; Tajima, T. Short-laser-pulse-driven emission of energetic ions into a solid target from a surface layer spalled by a laser prepulse. *Applied Physics A* **2001**, *73*, 741-747.
- (37) Cui, Y.; Hanley, L. ChiMS: Open-source instrument control software platform on labview for imaging/depth profiling mass spectrometers. *Review of Scientific Instruments* **2015**, *86*, 065106.
- (38) Shevchenko, A.; Tomas, H.; Havli, J.; Olsen, J. V.; Mann, M. In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nature Protocols* **2006**, *1*, 2856-2860.
- (39) Anthis, N. J.; Clore, G. M. Sequence-specific determination of protein and peptide concentrations by absorbance at 205 nm. *Protein Science* **2013**, *22*, 851-858.
- (40) Tyanova, S.; Temu, T.; Sinitcyn, P.; Carlson, A.; Hein, M. Y.; Geiger, T.; Mann, M.; Cox, J. The perseus computational platform for comprehensive analysis of (prote)omics data. *Nature Methods* **2016**, *13*, 731-740.
- (41) Tyanova, S.; Temu, T.; Cox, J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nature Protocols* **2016**, *11*, 2301-2319.
- (42) Consortium, T. U. UniProt: A worldwide hub of protein knowledge. *Nucleic Acids Research* **2018**, *47*, D506-D515.
- (43) Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical and Bioanalytical Chemistry* **1976**, *72*, 248-254.
- (44) ThermoScientific. Pierce detergent compatible Bradford assay kit, MAN0016018, Rev A.0 **2016**. https://assets.thermofisher.com/TFS-Assets/LSG/manuals/23246_23246S_deter_compat_bradford_UG.pdf
- (45) Anderl, J. N.; Franklin, M. J.; Stewart, P. S. Role of antibiotic penetration limitation in *Klebsiella pneumoniae* biofilm resistance to ampicillin and ciprofloxacin. *Antimicrobial Agents and Chemotherapy* **2000**, *44*, 1818-1824.
- (46) Batchelor, S. E.; Cooper, M.; Chhabra, S. R.; Glover, L. A.; Stewart, G. S. A. B.; Williams, P.; Prosser, J. I. Cell density-regulated recovery of starved biofilm populations of ammonia-oxidizing bacteria. *Applied and Environmental Microbiology* **1997**, *63*, 2281-2286.

- (47) Zuroff, T. R.; Bernstein, H.; Lloyd-Randolfi, J.; Jimenez-Taracido, L.; Stewart, P. S.; Carlson, R. P. Robustness analysis of culturing perturbations on *Escherichia coli* colony biofilm beta-lactam and aminoglycoside antibiotic tolerance *BMC Microbiology* **2010**, *10*, 185.
- (48) Vogel, C.; Marcotte, E. M. Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nature Reviews Genetics* **2012**, *13*, 227-232.
- (49) Corona, F.; Reales-Calderón, J. A.; Gil, C.; Martínez, J. L. The development of a new parameter for tracking post-transcriptional regulation allows the detailed map of the *Pseudomonas aeruginosa* CRC regulon. *Scientific Reports* **2018**, *8*, 16793.
- (50) Aseev, L. V.; Boni, I. V. Extraribosomal functions of bacterial ribosomal proteins. *Molecular Biology* **2011**, *45*, 739.
- (51) Park, H.; McGill, S. L.; Arnold, A. D.; Carlson, R. P. Pseudomonad reverse carbon catabolite repression, interspecies metabolite exchange, and consortial division of labor. *Cellular and Molecular Life Sciences* **2020**, *77*, 395-413.
- (52) Ciemniecki, J. A.; Newman, D. K. The potential for redox-active metabolites (RAMs) to enhance or unlock anaerobic survival metabolisms in aerobes. *Journal of Bacteriology* **2020**, JB.00797-00719.
- (53) Schiessl, K. T.; Hu, F.; Jo, J.; Nazia, S. Z.; Wang, B.; Price-Whelan, A.; Min, W.; Dietrich, L. E. P. Phenazine production promotes antibiotic tolerance and metabolic heterogeneity in *Pseudomonas aeruginosa* biofilms. *Nature Communications* **2019**, *10*, 762.
- (54) Gogoi, M.; Datey, A.; Wilson, K. T.; Chakravorty, D. Dual role of arginine metabolism in establishing pathogenesis. *Current Opinion in Microbiology* **2016**, *29*, 43-48.
- (55) Barrientos-Moreno, L.; Molina-Henares, M. A.; Ramos-González, M. I.; Espinosa-Urgel, M. Arginine as an environmental and metabolic cue for cyclic diguanylate signalling and biofilm formation in *Pseudomonas putida*. *Scientific Reports* **2020**, *10*, 13623.
- (56) Pivetti, C. D.; Yen, M.-R.; Miller, S.; Busch, W.; Tseng, Y.-H.; Booth, I. R.; Saier, Milton H. Two families of mechanosensitive channel proteins. *Microbiology and Molecular Biology Reviews* **2003**, *67*, 66-85.
- (57) Schaible, B.; Taylor, C. T.; Schaffer, K. Hypoxia increases antibiotic resistance in *Pseudomonas aeruginosa* through altering the composition of multidrug efflux pumps. *Antimicrobial Agents and Chemotherapy* **2012**, *56*, 2114-2118.
- (58) Ma, Q.; Zhai, Y.; Schneider, J. C.; Ramseier, T. M.; Saier, M. H. Protein secretion systems of *Pseudomonas aeruginosa* and *P. fluorescens*. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **2003**, *1611*, 223-233.

- (59) Thakur, P.; Gantasala, N. P.; Choudhary, E.; Singh, N.; Abdin, M. Z.; Agarwal, N. The preprotein translocase YidC controls respiratory metabolism in *Mycobacterium tuberculosis*. *Scientific Reports* **2016**, *6*, 24998.
- (60) Stojkova, P.; Spidlova, P.; Stulik, J. Nucleoid-associated protein HU: A lilliputian in gene regulation of bacterial virulence. *Frontiers in Cellular and Infection Microbiology* **2019**, *9*, 159.
- (61) Kugelberg, E.; Löfmark, S.; Wretling, B.; Andersson, D. I. Reduction of the fitness burden of quinolone resistance in *Pseudomonas aeruginosa*. *Journal of Antimicrobial Chemotherapy* **2005**, *55*, 22-30.
- (62) Lee, S.; Sowa, M. E.; Watanabe, Y.-h.; Sigler, P. B.; Chiu, W.; Yoshida, M.; Tsai, F. T. F. The structure of ClpB: A molecular chaperone that rescues proteins from an aggregated state. *Cell* **2003**, *115*, 229-240.
- (63) Davies, D. G.; Geesey, G. G. Regulation of the alginate biosynthesis gene *algC* in *Pseudomonas aeruginosa* during biofilm development in continuous culture. *Applied and Environmental Microbiology* **1995**, *61*, 860-867.
- (64) Park, S.-G.; Murray, K. K. Infrared laser ablation sample transfer for MALDI and electrospray. *Journal of the American Society for Mass Spectrometry* **2011**, *22*, 1352-1362.
- (65) Caprioli, R. M. Imaging mass spectrometry: Enabling a new age of discovery in biology and medicine through molecular microscopy. *Journal of the American Society for Mass Spectrometry* **2015**, *26*, 850-852.
- (66) Wang, X.; Han, J.; Hardie, D. B.; Yang, J.; Borchers, C. H. The use of matrix coating assisted by an electric field (MCAEF) to enhance mass spectrometric imaging of human prostate cancer biomarkers. *Journal of Mass Spectrometry* **2016**, *51*, 86-95.
- (67) Tillner, J.; Wu, V.; Jones, E. A.; Pringle, S. D.; Karancsi, T.; Dannhorn, A.; Veselkov, K.; McKenzie, J. S.; Takats, Z. Faster, more reproducible DESI-MS for biological tissue imaging. *Journal of the American Society for Mass Spectrometry* **2017**, *28*, 2090-2098.
- (68) Garza, K. Y.; Feider, C. L.; Klein, D. R.; Rosenberg, J. A.; Brodbelt, J. S.; Eberlin, L. S. Desorption electrospray ionization mass spectrometry imaging of proteins directly from biological tissue sections. *Analytical Chemistry* **2018**, *90*, 7785-7789.
- (69) Cole, L. M.; Mahmoud, K.; Haywood-Small, S.; Tozer, G. M.; Smith, D. P.; Clench, M. R. Recombinant "IMS TAG" proteins – a new method for validating bottom-up matrix-assisted laser desorption/ionisation ion mobility separation mass spectrometry imaging. *Rapid Communications in Mass Spectrometry* **2013**, *27*, 2355-2362.
- (70) Benigni, P.; Porter, J.; Ridgeway, M. E.; Park, M. A.; Fernandez-Lima, F. Increasing analytical separation and duty cycle with nonlinear analytical mobility scan functions in tims-FT-ICR MS. *Analytical Chemistry* **2018**, *90*, 2446-2450.

Figure 1

Figure 2

a

b

Figure 3

Figure 4