Positive modulation of cerebellar $\alpha 6$ GABA_A receptors for treating essential tremor: a proof-of-concept study in harmaline-treated mice

Ya-Hsien Huang,¹ Ming Tatt Lee,^{1,6} Werner Sieghart,³ Daniel E. Knutson,⁴ Laurin R. Wimmer,⁵
Dishary Sharmin,⁴ James Cook,⁴ Marko D. Mihovilovic,⁵ and Lih-Chu Chiou,^{1,2,7*}

Affiliations:

¹Department and Graduate Institute of Pharmacology, ²Graduate Institute of Brain and Mind Sciences, College of Medicine, National Taiwan University, Taipei 10051, Taiwan.

³Center for Brain Research, Department of Molecular Neurosciences, Medical University Vienna, Vienna, 1090, Wien, Austria.

⁴Department of Chemistry and Biochemistry, Milwaukee Institute for Drug Discovery, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin, WI, 53211, USA.

⁵Institute of Applied Synthetic Chemistry, TU Wien, 1060 Vienna, Austria.

⁶Faculty of Pharmaceutical Sciences, UCSI University, 56000 Kuala Lumpur, Malaysia.

⁷Graduate Institute of Acupuncture Science, China Medical University, Taichung 40402, Taiwan.

Short title: Cerebellar $\alpha 6$ GABA_ARs and essential tremor

***To whom correspondence should be addressed:**

Professor Lih-Chu Chiou, Department of Pharmacology, College of Medicine, National Taiwan University, No. 1, Jen-Ai Rd., Section 1, Taipei 100, Taiwan. Phone: 886-2-2312-3456 ext 88323; FAX: 886-2-2341-4788; Email: lcchiou@ntu.edu.tw

Abstract

Background: The etiology of essential tremor (ET) remains unclear but may involve abnormal firing of Purkinje cells, which receive excitatory inputs from granule cells in the cerebellum. Since $\alpha 6$ subunit-containing GABA_A receptors ($\alpha 6$ GABA_ARs) are abundantly expressed in granule cells, we validated a hypothesis that $\alpha 6$ GABA_AR-selective positive allosteric modulators (PAMs) are promising pharmacological interventions for ET therapy.

Methods: Employing the harmaline-induced ET model in male ICR mice, we evaluated the possible anti-tremor effects of four $\alpha 6$ GABA_AR-selective PAMs, the pyrazoloquinolinones Compound 6 and LAU-463 and their respective deuterated derivatives. Propranolol, a clinical anti-tremor agent, was employed as positive control. To investigate the involvement of cerebellar $\alpha 6$ GABA_ARs in the anti-tremor effect of intraperitoneal (*i.p.*) Compound 6, furosemide, an $\alpha 6$ GABA_AR antagonist, was intracerebellarly (*i.cb.*) co-administered with Compound 6. The burrowing activity, an indicator of well-being in rodents, was measured concurrently.

Results: Harmaline (10-30 mg/kg, *s.c.*) induced action tremor in ICR mice dose-dependently and markedly reduced their burrowing activity. Compound 6 (3 and 10 mg/kg, *i.p.*) significantly attenuated harmaline (20 mg/kg)-induced action tremor and burrowing activity impairment. Propranolol (20 mg/kg, *i.p.*) diminished tremor but failed to restore the burrowing activity in harmaline-treated mice. Importantly, both anti-tremor and burrowing activity restorative effects of Compound 6 (10 mg/kg, *i.p.*) was significantly reversed by co-administration of *i.cb.* furosemide at a dose (10 nmol/0.5 μ l) having no effect *per se*. All four $\alpha 6$ GABA_AR PAMs exhibited a similar therapeutic efficacy.

Conclusion: $\alpha 6$ GABA_AR-selective PAMs significantly attenuated action tremor and restored physical well-being in a mouse model mimicking ET by acting in the cerebellum. Thus, $\alpha 6$ GABA_AR-selective PAMs may be potential therapeutic agents for ET.

Keywords:

Essential tremor, $\alpha 6$ subunit-containing GABA_A receptor, cerebellum, harmaline, burrowing activity

Introduction

Essential tremor (ET) is a neurological disorder with symptoms characterized by uncontrollable rhythmic shaking of one or more body parts (Deuschl *et al.*, 1998; Haubenberger and Hallett, 2018). It is one of the most common movement disorders, especially in the elderly (Louis and Ferreira, 2010; Sepúlveda Soto and Fasano, 2020), and a potential risk factor for other neuropsychiatric conditions, such as depression and anxiety (Huang *et al.*, 2019; Louis, 2010). The socio-economic burden inflicted by ET is insurmountable, as it negatively affects the well-being and productivity of the patients and their caregivers (Monin *et al.*, 2017). Current therapeutic agents for ET are limited and often hampered by either inadequate efficacy or intolerable side effects (Shanker, 2019). Therefore, it is an urgent need to develop novel anti-tremor agents with a promising efficacy and minimal unwanted side effects.

The pathogenesis of ET remains not fully understood. Several human studies suggested a general hypo-function of GABAergic transmission in the brains of ET patients. ET patients had lower GABA, but higher glutamate concentrations in the cerebrospinal fluid (CSF) than normal controls (Mally *et al.*, 1996). Positron emission tomography (PET) studies demonstrated that ET patients had a higher binding potential of ^{11}C -flumazenil at the benzodiazepine (BDZ) site of GABA_ARs than normal subjects in brain regions involved in the pathogenesis of ET, including the deep cerebellar nuclei (DCN), cerebellar vermis, and ventral intermediate nucleus of the thalamus, and these changes correlated with their ET severity (Boecker *et al.*, 2010; Gironell *et al.*, 2012). An increased binding of ^{11}C -flumazenil suggests an increased expression of GABA_A receptors that contain BDZ binding sites in ET patients and might be a compensatory change in these brain regions due to the reduced GABA levels.

Consistent with a reduced GABAergic inhibition, ET patients have been reported to have excessive cerebellar activity revealed by neuroimaging studies (Sharifi *et al.*, 2014). Postmortem studies in ET patients revealed a significant loss of cerebellar Purkinje cells (PCs) (Louis *et al.*,

2013; Louis *et al.*, 2007) and their dendritic arborization (Kuo *et al.*, 2017; Louis *et al.*, 2007; Louis *et al.*, 2014). Other abnormalities in the cerebellum affected basket cells and climbing fibers (Louis and Faust, 2020). Such changes could have been the cause or the consequence of the excessive cerebellar activity.

In any case, Purkinje cells (PCs) are at the center of the repeating microcircuits that form the cerebellar cortex. They integrate information from five major classes of excitatory and inhibitory interneurons. The cerebellar cortex has two excitatory inputs, the mossy fibers and the climbing fibers. Mossy fibers provide the sensory and contextual information to the excitatory granule cells that elicit the coordinated motor programs via PCs. Climbing fibers provide error signals directly to the PCs, leading to an improved motor program and final timing for motor activity (D'Angelo *et al.*, 2013). Mossy fibers, in addition to granule cells, also excite the inhibitory Golgi cells. Golgi cells directly and indirectly inhibit granule cell excitation via feed forward and feedback loops, respectively, and sharpen their excitatory signal (Mapelli *et al.*, 2014; Nieuws *et al.*, 2014). In addition, Golgi cells cause a broad lateral inhibition of granule cells that generates dense clusters of granule cell activity organized in center-surround structures, implementing combinatorial operations of multiple mossy fiber inputs (D'Angelo *et al.*, 2013). The axons of granule cells are ascending to the molecular layer where they originate the parallel fibers that form excitatory synapses on the inhibitory PCs, Golgi cells, basket cells and stellate cells. In contrast to the granule cell-inhibiting Golgi cells, basket cells and stellate cells inhibit PCs, thus increasing the precision of PC firing in response to mossy fiber and climbing fiber excitation (Person and Raman, 2012). PCs are the major output neurons of the cerebellar cortex, providing a precise inhibitory timing control on the neuronal excitability in the downstream deep cerebellar nuclei (DCN) (Hirano, 2019; Pugh and Raman, 2009).

In agreement with their central role in the function of the cerebellum, PCs were recently reported to be significantly involved in the excessive cerebellar activity and play an important role in the pathogenesis of ET. In an ET model in which mice were treated with harmaline, a β -carboline

neurotoxin (Lavita *et al.*, 2016), it was demonstrated that the onset of tremor coincides with abnormal rhythmic firing of cerebellar PCs, which alters the downstream firing regularity of DCN neurons, and that a genetic reduction of PC-DCN transmission reduces tremor activity (Brown *et al.*, 2020). This conclusion was also supported by another study indicating that misfiring of cerebellar PCs elicited by synaptic pruning deficits of climbing fiber-to-PC synapses caused by PC-specific glutamate receptor subunit insufficiency, cause excessive cerebellar oscillations and might be responsible for tremor generation (Pan *et al.*, 2020).

Therefore, an enhancement of GABAergic transmission by positive allosteric modulators (PAMs) of GABA_ARs, like benzodiazepines (BDZs), may be a therapeutic strategy for ET patients. BDZs indeed are used clinically as a second-line treatment for ET (Shanker, 2019). However, the use of BDZs in ET treatment is limited by their unwanted side effects, like sedation, muscle weakness, addiction, confusion and amnesia (Bruno *et al.*, 2015). Conversely, PAMs that selectively modulate the GABA_ARs in the cerebellum would be an ideal option for ET treatment.

GABA_ARs are ligand-gated chloride channels consisting of 5 subunits. A total of 6 α , 3 β , 3 γ , δ , ϵ , π , θ and 3 ρ subunits have been identified in the mammalian nervous system (Olsen and Sieghart, 2008). The majority of GABA_ARs consists of two α , two β and one γ or δ subunits (Sieghart and Savic, 2018). The $\alpha 6$ subunit-containing GABA_A receptors ($\alpha 6$ GABA_ARs) are ideal targets for modulating the activity of PCs and a substantial diversity of heterocyclic scaffolds have been employed as allosteric modulators (Vega Alanis *et al.*, 2020). They are insensitive to classical benzodiazepines, like diazepam, (Luddens *et al.*, 1990; Sieghart, 1995) and are massively expressed in cerebellar granule cells (Gutierrez *et al.*, 1996; Pirker *et al.*, 2000). They form $\alpha 6\beta\gamma 2$ and $\alpha 1\alpha 6\beta\gamma 2$ GABA_ARs (Jechlinger *et al.*, 1998), mediating GABAergic inhibition at the Golgi cell-granule cell synapses (Nusser *et al.*, 1998). In addition, they form $\alpha 6\beta\delta$ and $\alpha 1\alpha 6\beta\delta$ GABA_ARs (Jechlinger *et al.*, 1998; Olsen and Sieghart, 2008; Scholze *et al.*, 2020) that are present at extrasynaptic sites at the Golgi cell-granule cell synapse and all over granule cells, their cell bodies, axons and their parallel

fibers, mediating tonic inhibition of granule cells (Lee *et al.*, 2010; Nusser *et al.*, 1998). These $\alpha 6$ GABA_ARs are much less expressed in other brain regions (Gutierrez *et al.*, 1996). Since $\alpha 6$ GABA_ARs regulate the excitability of granule cells, which represent one of the two main excitatory inputs of the cerebellum, enhancement of their inhibitory efficacy by $\alpha 6$ GABA_AR-selective PAMs seems to be one possible way to reduce the excessive activity of PCs. Here, we confirmed this concept using the harmaline-induced tremor model and four pyrazoloquinolinone compounds (PQs) (Zhang *et al.*, 1995) that we have previously identified to be PAMs highly selective to $\alpha 6$ GABA_ARs (Knutson *et al.*, 2018; Varagic *et al.*, 2013).

Harmaline can induce action tremor, involving the limbs, tail, trunk, head, and whiskers-, with a frequency range, 10-16 Hz in mice (Martin *et al.*, 2005), 8-12 Hz in rats (Martin *et al.*, 2005), 8-12 Hz in cats (Lamarre and Mercier, 1971) and 6-8 Hz in monkeys (Ohye *et al.*, 1970), resembling the action tremor in ET patients at the frequency of 4-12 Hz (Kuo *et al.*, 2019). Thus, the harmaline-induced tremor model is recognized as an animal model for screening potential therapeutic agents for ET treatment (Martin *et al.*, 2005). The PQ compounds tested include Compound 6, originally coded as PZ-II-029 (**Fig. 1A**), and its structural analogue LAU463 (**Fig. 1B**), as well as their respective deuterated analogues, DK-I-56-1 (**Fig. 1C**) and DK-I-58-1 (**Fig. 1D**), that have a better pharmacokinetic profile with longer half-lives (Knutson *et al.*, 2018). We first established the dose- and time-dependence of harmaline-induced tremor, as well as that of harmaline-induced disruption in the burrowing activity of mice, which is one of the “Activities of Daily Living” in laboratory rodents and has been employed as an indicator of well-being in rodents (Jirkof, 2014). Then, we investigated whether the PQ Compound 6 would inhibit harmaline-induced tremor and restore the burrowing activity disrupted by harmaline. To identify the site-of-action of Compound 6, we subsequently examined whether intracerebellar (*i.cb.*) microinjection of furosemide, a selective $\alpha 6$ GABA_AR antagonist, would reverse the effects of Compound 6. Lastly, we compared effects of the four PQ compounds on tremor and burrowing activities in harmaline-treated mice.

Materials and Methods

Animals

The animal care and experimental procedures reported in this study were approved by the Institutional Animal Care and Use Committee of National Taiwan University, College of Medicine. Male adult mice (ICR strain) were purchased from BioLASCO (Taiwan Co., Ltd) were housed in groups of 5 in a cage in a holding room with a 12 h light-dark reversed cycle (lights on at 20:00 and off at 08:00) and access to food and water *ad libitum*. On the experimental day, mice (6-10 weeks) were moved with their home cages to a behavior room and acclimated there for at least one hour before testing.

Intra-cerebellar (i.cb), intraperitoneal (i.p.) and subcutaneous (s.c.) injections

The *i.cb.* cannulation procedure was performed as reported in our previous studies (Chiou *et al.*, 2018; Liao *et al.*, 2016) with modifications. Briefly, mice were anaesthetized with sodium pentobarbital (60 mg/kg, *i.p.*) and placed in a stereotaxic frame keeping the bregma-lambda axis horizontal. After shaving the hair and exposing the skull surface, each mouse was implanted with a 24-gauge stainless-steel guide cannula directing toward the vermis (-7.0 mm caudal, -0.4 mm ventral from bregma) according to the stereotaxic coordinate of the mouse (Franklin and Paxinos, 1997). Each mouse was allowed to recover for at least 1 week. On the day for behavioral tests, a 30-gauge injection cannula connected to a 1- μ l Hamilton syringe via a 50 cm polyethylene tube was inserted into the guide cannula for drug injection. The drug solution of 0.5 μ l was slowly infused with a microinfusion pump (KDS311, KD Scientific Inc.) for 3 min with a further “hold” time for 2 min, while the mouse was allowed to freely move in an open field arena. The microinjection site was confirmed by the positive staining of trypan blue, which was injected through the cannula after the behavioral tests. Data from mice with offsite injections were discarded. For *i.p.* and *s.c.* injection, the drug solution was injected at a volume of 10 ml/kg.

Tremor induction and measurement

Action tremor was induced in mice by harmaline in accordance with the protocol described previously with minor modifications (Handforth *et al.*, 2018; Martin *et al.*, 2005). The frequency and intensity of tremor were measured and analyzed using a Tremor Monitor (San Diego Instruments, SDI, CA, USA) as reported previously (White *et al.*, 2016), which can accurately differentiate tremor activity from ambulatory/stereotyped movements and global motion activity (Paterson *et al.*, 2009). Each mouse was placed in the chamber for acclimatization, and then the baseline motion power was recorded for 10 min. Tested drugs were administered by *i.p.* or *i.cb.* injection 5 min before harmaline injection. Each raw trace of the movement activity recorded by Tremor Monitor was converted by a fast Fourier transform to generate a frequency domain-based motion power spectrum with the bin size at 1 Hz. The tremor activity in each mouse was measured as the ratio of the motion power at 10-16 Hz over the tremor frequency (0-34 Hz) of the global motion power over a 10-min duration (**Fig. 2A**). The overall tremor activity was expressed by the area under curve (AUC) derived from time-dependent tremor responses. To ensure that the harmaline-induced tremor detected in our setups was indeed action tremor, we videotaped the harmaline-treated mice's locomotor activity in the tremor chamber and subsequently analyzed the action/immobile motion of the mice using the SMART Video Tracking System (Harvard Apparatus, MA, USA) (**Fig.2B**).

Burrowing activity assessment

The burrowing activity of a mouse was measured as described previously (Ballon Romero *et al.*, 2020; Deacon, 2012) with modifications. For acclimatization, an empty tube was placed into the home cage, such that all five mice in the cage were exposed to the tube. On the day of testing, a separate cage was prepared for each mouse to be tested individually. The burrowing tube was filled with 200 g of food pellets. The burrowing activity was calculated as the weight of the expelled food pellet by subtracting the weight of the food pellet left at the end of the experiment from the initial weight (200

g).

Experimental protocol

As depicted in **Fig. 3A**, 10 min after harmaline (*s.c.*) injection, the mouse was placed in the tremor chamber and its tremor activity was measured for 10 min (pink shade). Right after the 10-min tremor measurement, the mouse was placed in a designated cage with the burrowing tube filled with 200 g of food pellets, and its burrowing activity was measured for another 10 min (grey shade). Then, the mouse was returned to the tremor chamber for tremor measurement. This tremor-burrowing measurement alternation was repeated for several cycles as indicated in each experiment. As harmaline-induced tremor is an action tremor, alternations were designed to refresh the alertness of the mouse once back in the tremor chamber, enhancing tremor consistency by replacing the rest period described in previous studies (Handforth *et al.*, 2018; Martin *et al.*, 2005) with the burrowing activity measurement.

Drugs and chemicals

Compound 6, LAU 463, DK-I-56-1 and DK-I-58-1 were synthesized as previously described (Knutson *et al.*, 2018). Harmaline and propranolol were purchased from Sigma-Aldrich (St. Louis, MO, USA) and furosemide from Tocris Bioscience (Bristol, UK). Compound 6, LAU463, DK-I-56-1, DK-I-58-1 (**Fig. 1, A-D**, respectively) and propranolol were dissolved in a vehicle containing 20% DMSO, 20% Cremophor® EL (polyoxyethylene castor; Sigma-Aldrich) and 60% normal saline. Harmaline was dissolved in normal saline. Furosemide administered by *i.cb.* microinjection was dissolved in DMSO as reported previously (Chiou *et al.*, 2018; Liao *et al.*, 2016).

Statistical analysis

Data were expressed as the mean $\pm$ S.E.M and the n number indicates the number of animals used. The two-way ANOVA followed by Holm-Sidak's *post hoc* test was employed to examine the difference in the tremor intensity among different treatment groups across time. The one-way ANOVA followed by Holm-Sidak's *post hoc* test was employed to examine the differences among treatment groups. However, if inhomogeneity was found in one-way ANOVA, as demonstrated in a significant Brown-Forsythe test, non-parametric Kruskal-Wallis followed by Dunn's *post hoc* test was employed. Statistical differences were considered significant if $P < 0.05$.

Results

Harmaline consistently induced tremor and disrupted burrowing activity in mice

As depicted in **Fig. 3A**, harmaline at doses of 10, 20 and 30 mg/kg (*s.c.*), significantly induced tremor activity in mice as demonstrated by two-way ANOVA with repeated measures over time, which showed main effects of time [$F(6,114) = 12.89, P < 0.001$] and treatment [$F(3,19) = 12.73, P < 0.001$], and a significant interaction between time and treatment [$F(18,114) = 2.778, P < 0.001$]. The tremor activity peaked at 10-16 Hz as demonstrated in the motion power-frequency distribution plot and heatmap spectrogram (**Fig. 2A**). As shown in **Fig 2B**, tremor activity was detected during mobile (red) periods, but not during the immobile phase (black periods) of the harmaline-treated mice, suggesting that harmaline-induced tremor is an action tremor.

Harmaline-induced tremor activity reached peak at the first assessing time interval, 10-20 min after harmaline injection, and was similar among various dosage groups, but the time-dependent analysis with one-way ANOVA showed a significant difference among treatment groups [$F(3,19) = 6.472, P = 0.003$] (**Fig. 3A**). The tremor activity declined gradually and dose-dependently. The peak tremor activity lasted for at least 50, 75 and 115 min, respectively, induced by 10, 20 and 30 mg/kg harmaline. The area under curves (AUC) of the tremor activity over the 120 min-measuring period showed that the tremor-inducing effect of harmaline was dose-dependent (**Fig. 3B**). Besides inducing tremor, harmaline at doses of 10 and 20 mg/kg, significantly suppressed the burrowing activity in mice (**Fig. 3C**), but not at 30 mg/kg. The latter negative result might have been due to the lower number of mice used in this experiment, or to unwanted detrimental effects induced by a high dose of harmaline since it is a neurotoxin that can act at many ion channels and neurotransmitter receptors (Du *et al.*, 1997; Jiang *et al.*, 2019; Zhan and Graf, 2012). Thus, harmaline at the dose of 20 mg/kg (*s.c.*) was chosen to induce tremor and burrowing activity deficit for subsequent pharmacological studies, with the measuring period limited to the first 75 mins after harmaline injection.

Compound 6 (i.p.) suppressed tremor and restored burrowing activity in harmaline-treated mice

Similar to the protocol described in **Fig. 3A**, vehicle, Compound 6 (3 or 10 mg/kg) or propranolol (20 mg/kg) were *i.p.* administered 5 min prior to harmaline (20 mg/kg, *s.c.*) injection (**Fig. 4A**). Two-way ANOVA with repeated measurements over time showed main effects of time [$F(4,164) = 41.59$, $P < 0.001$] and treatment [$F(3,41) = 12.57$, $P < 0.001$], and a significant interaction between time and treatment [$F(12,164) = 8.327$, $P < 0.001$]. Both doses (3 and 10 mg/kg, *i.p.*) of Compound 6 and propranolol (20 mg/kg) significantly reduced tremor activity, as compared with the vehicle-treated group (**Fig. 4A**). The AUC derived from the time-dependent analysis showed that both doses of Compound 6 exerted significant anti-tremor effect, but propranolol showed the highest suppressive effect (**Fig. 4B**). One-way ANOVA showed a significant difference among treatment groups [$F(3,41) = 12.85$, $P < 0.001$]. However, one-way ANOVA followed by Holm-Sidak *post hoc* test demonstrated that the tremor-suppressant dose of propranolol failed to significantly restore the burrowing activity in harmaline-induced mice ($P = 0.18$), in contrast to Compound 6 at 3 mg/kg ($P = 0.05$) and 10 mg/kg ($P = 0.05$).

Furosemide (i.cb.) antagonized effects of Compound 6 (i.p.) on tremor and burrowing activities in harmaline-treated mice

In order to discern whether cerebellar $\alpha 6\text{GABA}_A\text{Rs}$ are involved in the anti-tremor effect of Compound 6, we treated mice with co-administration of *i.cb.* furosemide (10 nmol) and *i.p.* Compound 6 (10 mg/kg), 5 min before harmaline injection (**Fig. 5A**). Two-way ANOVA with repeated measures over time showed main effects of time [$F(4,164) = 41.59$, $P < 0.001$] and treatment [$F(3,41) = 12.57$, $P < 0.001$], and a significant interaction between time and treatment [$F(12,164) = 8.327$, $P < 0.001$] (**Fig. 5A**). In both time-dependent analyses (yellow diamonds, **Fig. 5A**) and the derived AUC of motion power (yellow bar, **Fig. 5B**) in harmaline-treated mice, *i.cb.* furosemide alone had no effect on the tremor activity. Co-administered of Compound 6 with *i.cb.*

microinjection of furosemide (green bar, **Fig. 5B**), but not its vehicle (blue bar, **Fig. 5A**), did not display significant anti-tremor effect, suggesting *i.cb.* furosemide, at the dose without affecting tremor activity *per se*, significantly antagonizes the anti-tremor effect of Compound 6. Interestingly, the restorative effect of Compound 6 on burrowing-activity in harmaline-treated mice was also completely reversed by *i.cb.* furosemide (green bar, **Fig. 5C**) while *i.cb.* furosemide *per se* did not affect the burrowing activity (yellow bar, **Fig. 5C**).

Compound 6 analogues and deuterated derivatives displayed similar anti-tremor and burrowing-restorative effects in harmaline-treated mice

Compound 6 and its structural analogue, LAU463, as well as their respective deuterated derivatives, DK-I-56-1 and DK-I-58-1, were previously demonstrated to possess similar selectivity towards $\alpha 6$ GABA_AR as PAMs. In the present study, administration of DK-I-56-1 (half-filled square symbols, **Fig. 6A**), LAU-463 (filled diamonds, **Fig. 6B**) and DK-I-58-1 (half-filled diamonds, **Fig. 6C**) at the of 3 and 10 mg/kg (*i.p.*) demonstrated time and treatment-dependent suppression of harmaline-induced tremor in mice, lasting for almost 1 hr. By comparing the AUC derived from time-dependent analysis showed that each drug group significantly suppressed tremor activity as compared with vehicle group, except DK-I-58-1 at the dose of 3 mg/kg. Interestingly, as compared with the vehicle group within each drug group, all treatment groups significantly restored burrowing activity in harmaline-treated mice (**Fig. 6E**), similar to the effect of Compound 6, but not of propranolol (**Fig. 4C**),

Discussion

$\alpha 6$ GABA_AR-selective PQ compounds suppressed tremor and restored burrowing activity in harmaline-treated mice.

In this study, we found that harmaline at 10-30 mg/kg (*s.c.*) induced significant action tremor at 10-16 Hz, with a similar maximal tremor activity but a dose-dependent action duration in ICR mice (**Fig. 3**), consistent with a previous report (Martin *et al.*, 2005). The predictive validity of harmaline-induced tremor in modeling ET in patients was strongly supported by the finding that it responded to clinically used ET-relieving agents, such as propranolol, primidone, alcohol, benzodiazepines, gabapentin, gammahydroxybutyrate, 1-octanol, and zonisamide, and was exacerbated by drugs that worsen ET, such as tricyclics and caffeine (Handforth, 2012). In this action tremor model, we provided a proof-of-concept pre-clinical study indicating that $\alpha 6$ GABA_AR-selective PAMs are a novel pharmacotherapy for ET.

Compound 6 as well as its three structural analogues effectively suppressed tremor activity and restored burrowing activity in harmaline-treated mice. Whereas Compound 6 and LAU-463 are structural analogues with different functional groups on the A-ring, DK-I-56-1 and DK-I-58-1 are their respective derivatives with a deuterated methoxy group on the D-ring (**Fig. 1**). These deuterated derivatives, compared with their parent compounds, had a similar potency and the same $\alpha 6$ GABA_AR-selectivity in enhancing GABA currents of recombinant $\alpha 6\beta\gamma$ GABA_ARs expressed in *Xenopus* oocytes (Knutson *et al.*, 2018; Treven *et al.*, 2018) but displayed longer half-lives in rodents (Knutson *et al.*, 2018). Although the merit of longer half-lives of deuterated compounds, cannot be observed in these behavioral assays that only lasted for 90 minutes, for future research and a possible clinical development of $\alpha 6$ GABA_AR-selective PQ compounds, it was important to also investigate the activity of these deuterated compounds in these assays.

The burrowing activity in laboratory rodents is one of their “Activities of Daily Living” (ADL) (Deacon, 2012), and has been used as an indicator of well-being (Jirkof, 2014), because it is

negatively associated with stress (Van Loo *et al.*, 2007) or chronic pain (Huang *et al.*, 2013) in rodents. In addition, a deficit of the burrowing activity has also been reported in mouse models of neurodegenerative diseases, such as Alzheimer's disease (Geiszler *et al.*, 2016) and Parkinson's disease (Baumann *et al.*, 2016). The finding that harmaline markedly disrupted the burrowing activity in mice (**Fig. 3C**) suggests that this ET animal model mimics not only the action tremor but also the reduced ADL score manifested in ET patients because of the motor disorder-induced distress that severely affects their well-being. Interestingly, $\alpha 6$ GABA_AR-selective PQ compounds also restored the burrowing activity in harmaline-treated mice (**Fig. 4C** and **Fig. 6E**) in addition to attenuating their tremor activity. In contrast, propranolol, the first-line medication for essential tremor, did not restore the burrowing activity in mice (**Fig. 2D**). This is probably due to the reduction of the locomotor activity induced by propranolol (Poli and Palermo-Neto, 1985), which also causes muscle weakness in humans (Deuschl *et al.*, 2011). Since $\alpha 6$ GABA_AR PAMs can both reduce tremor activity and restore burrowing activity in harmaline-treated mice, they may be viable candidates to complement current anti-tremor drugs.

Cerebellar $\alpha 6$ GABA_ARs are the site of action of anti-tremor effects of $\alpha 6$ GABA_AR-selective PQ compounds

The finding that *i.cb.* microinjection of furosemide, an $\alpha 6$ GABA_AR-selective antagonist (Korpi and Luddens, 1997), antagonized the anti-tremor effect of Compound 6 (**Fig. 3, B & C**) suggests that the $\alpha 6$ GABA_ARs in the cerebellum are the site of action of Compound 6. A previous study indicated that gaboxadol, a selective agonist of the δ -subunit containing GABA_ARs (δ GABA_ARs), including the $\alpha 6\beta\delta$ subtype of GABA_ARs, significantly suppressed harmaline-induced tremor in wild type, but not in *Gabra6*^{-/-} or *Gabrd*^{-/-} mice (Handforth *et al.*, 2018), suggesting that the $\alpha 6\beta\delta$ GABA_AR subtype is involved in harmaline-induced tremor. Our previous electrophysiological study on oocyte expressing various recombinant GABA_AR subtypes demonstrated that Compound 6

exerted its PAM activity at both $\alpha 6\beta\gamma 2$ and $\alpha 6\beta\delta$ containing $\alpha 6$ GABA_ARs although having a lower efficacy for the $\alpha 6\beta\delta$ subtype (Chiou *et al.*, 2018). Even though, it enhanced the GABA current of $\alpha 6\beta\delta$ GABA_ARs to 270% at 3 μ M (Chiou *et al.*, 2018). Thus, it is likely that cerebellar $\alpha 6\beta\delta$ GABA_ARs are the action target of Compound 6. On the other hand, a contribution of the $\alpha 6\beta\gamma 2$ GABA_AR subtype in the anti-tremor effect of Compound 6 cannot be excluded. Compound 6 could reduce the neuronal excitability of granule cells by enhancing GABAergic transmission at extrasynaptic $\alpha 6\beta\delta$ GABA_ARs as well as at $\alpha 6\beta\gamma 2$ GABA_ARs at Golgi cell-granule cell synapses (Nusser *et al.*, 1998), thus ultimately reducing the excitatory input of granule cells and hence the mis-firing of PCs.

Current essential tremor therapy is insufficient

Currently, essential tremor is symptomatically treated with medicines that can suppress tremor activity, like β -blockers, anti-epileptics, neuron stabilizers and GABA-related drugs (Gironell, 2014; Shanker, 2019). Among these, propranolol and primidone remain the first line medications for relieving tremor in patients with essential tremor, even decades after their initial applications (Shanker, 2019). A combination of these two drugs yields a better response than using either of them alone (Zesiewicz *et al.*, 2005). However, one third of patients developed drug resistance (Deuschl *et al.*, 2011), and a significant proportion of the patients are even refractory to both medications (Louis *et al.*, 2010). The reported adverse effects differ between these two drugs with acute reactions more common with primidone and chronic reactions with propranolol (Bain, 1997). The acute side effects associated with primidone are sedation, nausea and vertigo. They are manifest after the first dose and tend to subside on chronic administrations (Koller and Royse, 1986; Sasso *et al.*, 1990). On the other hand, chronic propranolol may lead to bradycardia, hypotension and breathlessness, especially when higher doses are used. Therefore, propranolol is contraindicated in

conditions such as chronic obstructive pulmonary disease, asthma, severe peripheral vascular disease and diabetes (Winkler and Young, 1974).

Limited data from randomized controlled trials are available to support the use of other medications in essential tremor, including topiramate, alprazolam, gabapentin, and other beta-blockers besides propranolol (e.g., atenolol, nadolol, and sotalol) (Deuschl *et al.*, 2011; Haubenberger and Hallett, 2018; Schaefer *et al.*, 2018; Zesiewicz *et al.*, 2011). Randomized controlled trials have shown no significant benefit for several other drugs for essential tremor, including levetiracetam, amifampridine, flunarizine, trazodone, pindolol, acetazolamide, mirtazapine, nifedipine, and verapamil (Zesiewicz *et al.*, 2011). As a result, the need for an effective anti-tremor drug that is well-tolerated is urgent and obvious.

$\alpha 6$ GABA_AR-selective PQ compounds as a novel therapy for essential tremor

We have previously demonstrated that Compound 6, one of the first $\alpha 6$ GABA_AR-selective PAMs identified (Knutson *et al.*, 2018; Varagic *et al.*, 2013), via acting at cerebellar $\alpha 6$ GABA_ARs, ameliorated disrupted prepulse inhibition (PPI) of the startle response (Chiou *et al.*, 2018), social withdrawal and cognitive impairment (Chiou *et al.*, 2019) in animal models mimicking schizophrenia. By acting via $\alpha 6$ GABA_ARs at trigeminal ganglia, it inhibited nociceptive activation of the trigeminovascular system and migraine-like grimaces in animal models of migraine (Fan *et al.*, 2018; Tzeng *et al.*, 2020). And, a deuterated Compound 6 analogue, DK-I-56-1, prevented and reduced trigeminal neuropathic pain in a rat model (Vasovic *et al.*, 2019). Here, we further substantiated the effectiveness of $\alpha 6$ GABA_AR-selective PQ compounds in an animal model of essential tremor. The effective dose range (3-10 mg/kg, *i.p.*) of Compound 6 in suppressing tremor was similar to that in mouse models of schizophrenia (Chiou *et al.*, 2018; Chiou *et al.*, 2019) and migraine (Tzeng *et al.*, 2020). At these doses, Compound 6 did not impair motor functions in rats (Knutson *et al.*, 2018), nor display sedative (Wu and Chiou, 2016) or hypolocomotor activity (Chiou

et al., 2019) in mice. In contrast to the GABA binding site agonist, gaboxadol (THIP), which directly activates all $\alpha 4\beta\delta$ and $\alpha 6\beta\delta$ GABA_ARs in nervous tissues (Brown *et al.*, 2020; Meera *et al.*, 2011) and exhibits sedation and motor-incoordination (Wafford and Ebert, 2006), $\alpha 6$ GABA_AR-selective PAMs can only allosterically modulate the activity of those $\alpha 6\beta\delta$ and $\alpha 6\beta\gamma 2$ GABA_ARs, which are activated by GABA in a certain task, explaining the absence of such side effects. In contrast to BDZs and gaboxadol, $\alpha 6$ GABA_AR-selective PQ compounds may thus exert therapeutic effects without sedative or motor-impairing activity.

In addition, the investigated $\alpha 6$ GABA_AR-selective PQ compounds have a satisfactory safety pharmacology profile. In a panel of radioligand binding assays targeting 46 CNS receptors/channels/enzymes, these PQ compounds displayed an affinity for GABA_AR binding that was more than 1000-fold higher than that for 45 other off targets, including the hERG channel, an arrhythmia risk potassium channel (Knutson *et al.*, 2018). In cytotoxicity tests on HEK293 (kidney) and HEPGS (liver) cells, PQ compounds at the concentration up to 400 μ M did not display significant cytotoxicity (Knutson *et al.*, 2018). However, due to their $\alpha 6$ GABA_AR selectivity, these PQ compounds only modulated the excitatory input of mossy fibers, but not that of climbing fibers in cerebellum, possibly explaining their incomplete suppression of the tremor activity. Nevertheless, in view of their novel target-acting property, satisfactory safety pharmacology profile, and absence of side effects, the investigated $\alpha 6$ GABA_AR-selective PAMs can be developed as an alternative monotherapy or a non-toxic add-on therapy for the treatment of ET.

Acknowledgement

This study was supported by grants from the Ministry of Science and Technology, Taiwan (MOST 104-2923-B-002-006-MY3, MOST 108-2320-B-002-029-MY3 and 109-2320-B-002-042-MY3 to LCC), National Health Research Institutes, Taiwan (NHRI-EX107-10733NI to LCC), Ministry of Education, Taiwan (107M4022-3 to LCC), UCSI University Research Excellence and Innovation Grant (REIG-FPS-2020/065 to MTL) National Institutes of Health, USA (R01AA029023 and R01MH096463 to JC), and the National Science Foundation, USA, Division of Chemistry (CHE-1625735 to JC). We thank the Milwaukee Institute of Drug Discovery for the mass spectroscopy. LRW was supported by the Austrian Science Fund FWF grant W1232 (Graduate School Program MolTag).

Authors' Contributions

YHH conducted experiments, analyzed data, and wrote the paper; MTL and WS contributed to data interpretation, data analysis, result discussion, and paper writing; DEK, LRW, DS, MDM and JC synthesized and provided compounds; LCC provided supervision, designed experiments, analyzed data, and wrote the paper.

References

- Bain PG (1997). The Effectiveness of Treatments for Essential Tremor. *Neurologist* 3: 305-321.
- Ballon Romero SS, Lee YC, Fuh LJ, Chung HY, Hung SY, Chen YH (2020). Analgesic and Neuroprotective Effects of Electroacupuncture in a Dental Pulp Injury Model-A Basic Research. *Int J Mol Sci* 21.
- Baumann A, Moreira CG, Morawska MM, Masneuf S, Baumann CR, Noain D (2016). Preliminary Evidence of Apathetic-Like Behavior in Aged Vesicular Monoamine Transporter 2 Deficient Mice. *Front Hum Neurosci* 10: 587.
- Boecker H, Weindl A, Brooks DJ, Ceballos-Baumann AO, Liedtke C, Miederer M, *et al.* (2010). GABAergic dysfunction in essential tremor: an 11C-flumazenil PET study. *J Nucl Med* 51: 1030-1035.
- Brown AM, White JJ, van der Heijden ME, Zhou J, Lin T, Sillitoe RV (2020). Purkinje cell misfiring generates high-amplitude action tremors that are corrected by cerebellar deep brain stimulation. *Elife* 9.
- Bruno E, Nicoletti A, Quattrocchi G, Filippini G, Zappia M, Colosimo C (2015). Alprazolam for essential tremor. *Cochrane Database Syst Rev*: CD009681.
- Chiou LC, Lee HJ, Ernst M, Huang WJ, Chou JF, Chen HL, *et al.* (2018). Cerebellar alpha6 - subunit-containing GABAA receptors: a novel therapeutic target for disrupted prepulse inhibition in neuropsychiatric disorders. *Br J Pharmacol* 175: 2414-2427.
- Chiou LC, Lee HJ, Lee MT, Tzeng HR, Fan PC, Kubota H, *et al.* (2019). The alpha-6 subunit-containing GABAA receptor is a novel therapeutic target for schizophrenia and migraine. *2019 Neuroscience Meeting Planner*; Chicago, IL, USA. Society for Neuroscience.
- D'Angelo E, Solinas S, Mapelli J, Gandolfi D, Mapelli L, Prestori F (2013). The cerebellar Golgi cell and spatiotemporal organization of granular layer activity. *Front Neural Circuits* 7: 93.
- Deacon R (2012). Assessing burrowing, nest construction, and hoarding in mice. *J Vis Exp*: e2607.

- Deuschl G, Bain P, Brin M (1998). Consensus statement of the Movement Disorder Society on Tremor. Ad Hoc Scientific Committee. *Mov Disord* 13 Suppl 3: 2-23.
- Deuschl G, Raethjen J, Hellriegel H, Elble R (2011). Treatment of patients with essential tremor. *Lancet Neurol* 10: 148-161.
- Du W, Aloyo VJ, Harvey JA (1997). Harmaline competitively inhibits [3H]MK-801 binding to the NMDA receptor in rabbit brain. *Brain Res* 770: 26-29.
- Fan PC, Lai TH, Hor CC, Lee MT, Huang P, Sieghart W, *et al.* (2018). The alpha6 subunit-containing GABAA receptor: A novel drug target for inhibition of trigeminal activation. *Neuropharmacology* 140: 1-13.
- Franklin K, Paxinos G (1997). The mouse brain in stereotaxic coordinates, 1997. Academic, San Diego.
- Geiszler PC, Barron MR, Pardon MC (2016). Impaired burrowing is the most prominent behavioral deficit of aging htau mice. *Neuroscience* 329: 98-111.
- Gironell A (2014). The GABA Hypothesis in Essential Tremor: Lights and Shadows. *Tremor Other Hyperkinet Mov (N Y)* 4: 254.
- Gironell A, Figueiras FP, Pagonabarraga J, Herance JR, Pascual-Sedano B, Trampal C, *et al.* (2012). Gaba and serotonin molecular neuroimaging in essential tremor: a clinical correlation study. *Parkinsonism Relat Disord* 18: 876-880.
- Gutierrez A, Khan ZU, De Blas AL (1996). Immunocytochemical localization of the alpha 6 subunit of the gamma-aminobutyric acidA receptor in the rat nervous system. *J Comp Neurol* 365: 504-510.
- Handforth A (2012). Harmaline tremor: underlying mechanisms in a potential animal model of essential tremor. *Tremor Other Hyperkinet Mov (N Y)* 2.

- Handforth A, Kadam PA, Kosoyan HP, Eslami P (2018). Suppression of Harmaline Tremor by Activation of an Extrasynaptic GABAA Receptor: Implications for Essential Tremor. *Tremor Other Hyperkinet Mov* (N Y) 8: 546.
- Haubenberger D, Hallett M (2018). Essential Tremor. *N Engl J Med* 379: 596-597.
- Hirano T (2019). GABA and Synaptic Transmission in the Cerebellum. In: Manto M, Gruol D, Schmähmann J, Koibuchi N, Sillitoe R (ed)^(eds). *Handbook of the Cerebellum and Cerebellar Disorders*, edn. Cham: Springer International Publishing. p[^]pp 1-14.
- Huang H, Yang X, Zhao Q, Chen Y, Ning P, Shen Q, *et al.* (2019). Prevalence and Risk Factors of Depression and Anxiety in Essential Tremor Patients: A Cross-Sectional Study in Southwest China. *Front Neurol* 10: 1194.
- Huang W, Calvo M, Karu K, Olausen HR, Bathgate G, Okuse K, *et al.* (2013). A clinically relevant rodent model of the HIV antiretroviral drug stavudine induced painful peripheral neuropathy. *Pain* 154: 560-575.
- Jechlinger M, Pelz R, Tretter V, Klausberger T, Sieghart W (1998). Subunit composition and quantitative importance of hetero-oligomeric receptors: GABAA receptors containing alpha6 subunits. *J Neurosci* 18: 2449-2457.
- Jiang B, Meng L, Zou N, Wang H, Li S, Huang L, *et al.* (2019). Mechanism-based pharmacokinetics-pharmacodynamics studies of harmine and harmaline on neurotransmitters regulatory effects in healthy rats: Challenge on monoamine oxidase and acetylcholinesterase inhibition. *Phytomedicine* 62: 152967.
- Jirkof P (2014). Burrowing and nest building behavior as indicators of well-being in mice. *J Neurosci Methods* 234: 139-146.
- Knutson DE, Kodali R, Divovic B, Treven M, Stephen MR, Zahn NM, *et al.* (2018). Design and Synthesis of Novel Deuterated Ligands Functionally Selective for the gamma-Aminobutyric

- Acid Type A Receptor (GABAAR) alpha6 Subtype with Improved Metabolic Stability and Enhanced Bioavailability. *J Med Chem* 61: 2422-2446.
- Koller WC, Royse VL (1986). Efficacy of primidone in essential tremor. *Neurology* 36: 121-124.
- Korpi ER, Luddens H (1997). Furosemide interactions with brain GABA_A receptors. *Br J Pharmacol* 120: 741-748.
- Kuo SH, Louis ED, Faust PL, Handforth A, Chang SY, Avlar B, *et al.* (2019). Current Opinions and Consensus for Studying Tremor in Animal Models. *Cerebellum* 18: 1036-1063.
- Kuo SH, Wang J, Tate WJ, Pan MK, Kelly GC, Gutierrez J, *et al.* (2017). Cerebellar Pathology in Early Onset and Late Onset Essential Tremor. *Cerebellum* 16: 473-482.
- Lamarre Y, Mercier LA (1971). Neurophysiological studies of harmaline-induced tremor in the cat. *Can J Physiol Pharmacol* 49: 1049-1058.
- Lavita SI, Aro R, Kiss B, Manto M, Duez P (2016). The Role of beta-Carboline Alkaloids in the Pathogenesis of Essential Tremor. *Cerebellum* 15: 276-284.
- Lee S, Yoon BE, Berglund K, Oh SJ, Park H, Shin HS, *et al.* (2010). Channel-mediated tonic GABA release from glia. *Science* 330: 790-796.
- Liao YH, Lee HJ, Huang WJ, Fan PC, Chiou LC (2016). Hispidulin alleviated methamphetamine-induced hyperlocomotion by acting at alpha6 subunit-containing GABA_A receptors in the cerebellum. *Psychopharmacology (Berl)* 233: 3187-3199.
- Louis ED (2010). Essential tremor as a neuropsychiatric disorder. *J Neurol Sci* 289: 144-148.
- Louis ED, Babij R, Lee M, Cortes E, Vonsattel JP (2013). Quantification of cerebellar hemispheric purkinje cell linear density: 32 ET cases versus 16 controls. *Mov Disord* 28: 1854-1859.
- Louis ED, Faust PL (2020). Essential tremor pathology: neurodegeneration and reorganization of neuronal connections. *Nat Rev Neurol* 16: 69-83.

- Louis ED, Faust PL, Vonsattel JP, Honig LS, Rajput A, Robinson CA, *et al.* (2007). Neuropathological changes in essential tremor: 33 cases compared with 21 controls. *Brain* 130: 3297-3307.
- Louis ED, Ferreira JJ (2010). How common is the most common adult movement disorder? Update on the worldwide prevalence of essential tremor. *Mov Disord* 25: 534-541.
- Louis ED, Lee M, Babij R, Ma K, Cortes E, Vonsattel JP, *et al.* (2014). Reduced Purkinje cell dendritic arborization and loss of dendritic spines in essential tremor. *Brain* 137: 3142-3148.
- Louis ED, Rios E, Hencheliffe C (2010). How are we doing with the treatment of essential tremor (ET)? Persistence of patients with ET on medication: data from 528 patients in three settings. *Eur J Neurol* 17: 882-884.
- Luddens H, Pritchett DB, Kohler M, Killisch I, Keinanen K, Monyer H, *et al.* (1990). Cerebellar GABAA receptor selective for a behavioural alcohol antagonist. *Nature* 346: 648-651.
- Mally J, Baranyi M, Vizi ES (1996). Change in the concentrations of amino acids in CSF and serum of patients with essential tremor. *J Neural Transm (Vienna)* 103: 555-560.
- Mapelli L, Solinas S, D'Angelo E (2014). Integration and regulation of glomerular inhibition in the cerebellar granular layer circuit. *Front Cell Neurosci* 8: 55.
- Martin FC, Thu Le A, Handforth A (2005). Harmaline-induced tremor as a potential preclinical screening method for essential tremor medications. *Mov Disord* 20: 298-305.
- Meera P, Wallner M, Otis TS (2011). Molecular basis for the high THIP/gaboxadol sensitivity of extrasynaptic GABA(A) receptors. *J Neurophysiol* 106: 2057-2064.
- Monin JK, Gutierrez J, Kellner S, Morgan S, Collins K, Rohl B, *et al.* (2017). Psychological Suffering in Essential Tremor: A Study of Patients and Those Who Are Close to Them. *Tremor Other Hyperkinet Mov (N Y)* 7: 526.
- Nieus TR, Mapelli L, D'Angelo E (2014). Regulation of output spike patterns by phasic inhibition in cerebellar granule cells. *Front Cell Neurosci* 8: 246.

- Nusser Z, Sieghart W, Somogyi P (1998). Segregation of different GABAA receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci* 18: 1693-1703.
- Ohye C, Bouchard R, Larochelle L, Bedard P, Boucher R, Raphy B, *et al.* (1970). Effect of dorsal rhizotomy on postural tremor in the monkey. *Exp Brain Res* 10: 140-150.
- Olsen RW, Sieghart W (2008). International Union of Pharmacology. LXX. Subtypes of gamma-aminobutyric acid(A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 60: 243-260.
- Pan MK, Li YS, Wong SB, Ni CL, Wang YM, Liu WC, *et al.* (2020). Cerebellar oscillations driven by synaptic pruning deficits of cerebellar climbing fibers contribute to tremor pathophysiology. *Sci Transl Med* 12.
- Paterson NE, Malekiani SA, Foreman MM, Olivier B, Hanania T (2009). Pharmacological characterization of harmaline-induced tremor activity in mice. *Eur J Pharmacol* 616: 73-80.
- Person AL, Raman IM (2012). Synchrony and neural coding in cerebellar circuits. *Front Neural Circuits* 6: 97.
- Poli A, Palermo-Neto J (1985). Effects of d,l-propranolol on open field behavior of rats. *Psychopharmacology (Berl)* 86: 153-155.
- Pugh JR, Raman IM (2009). Nothing can be coincidence: synaptic inhibition and plasticity in the cerebellar nuclei. *Trends in neurosciences* 32: 170-177.
- Sasso E, Perucca E, Fava R, Calzetti S (1990). Primidone in the long-term treatment of essential tremor: a prospective study with computerized quantitative analysis. *Clin Neuropharmacol* 13: 67-76.
- Schaefer SM, Vives Rodriguez A, Louis ED (2018). Brain circuits and neurochemical systems in essential tremor: insights into current and future pharmacotherapeutic approaches. *Expert Rev Neurother* 18: 101-110.

- Scholze P, Pokl M, Langle S, Steudle F, Fabjan J, Ernst M (2020). Two Distinct Populations of $\alpha_1\alpha_6$ -Containing GABAA-Receptors in Rat Cerebellum. *Front Synaptic Neurosci* 12: 591129.
- Sepúlveda Soto MC, Fasano A (2020). Essential tremor: New advances. *Clinical Parkinsonism & Related Disorders* 3: 100031.
- Shanker V (2019). Essential tremor: diagnosis and management. *BMJ* 366: 14485.
- Sharifi S, Nederveen AJ, Booij J, van Rootselaar AF (2014). Neuroimaging essentials in essential tremor: a systematic review. *Neuroimage Clin* 5: 217-231.
- Sieghart W (1995). Structure and pharmacology of gamma-aminobutyric acidA receptor subtypes. *Pharmacol Rev* 47: 181-234.
- Sieghart W, Savic MM (2018). International Union of Basic and Clinical Pharmacology. CVI: GABAA Receptor Subtype- and Function-selective Ligands: Key Issues in Translation to Humans. *Pharmacol Rev* 70: 836-878.
- Treven M, Siebert DCB, Holzinger R, Bampali K, Fabjan J, Varagic Z, *et al.* (2018). Towards functional selectivity for $\alpha_6\beta_3\gamma_2$ GABAA receptors: a series of novel pyrazoloquinolinones. *Br J Pharmacol* 175: 419-428.
- Tzeng HR, Lee MT, Fan PC, Knutson DE, Lai TH, Sieghart W, *et al.* (2020). α_6 GABAA Receptor Positive Modulators Alleviate Migraine-like Grimaces in Mice via Compensating GABAergic Deficits in Trigeminal Ganglia. *Neurotherapeutics*.
- Van Loo PL, Kuin N, Sommer R, Avsaroglu H, Pham T, Baumans V (2007). Impact of 'living apart together' on postoperative recovery of mice compared with social and individual housing. *Lab Anim* 41: 441-455.
- Varagic Z, Ramerstorfer J, Huang S, Rallapalli S, Sarto-Jackson I, Cook J, *et al.* (2013). Subtype selectivity of α + β - site ligands of GABAA receptors: identification of the first highly

specific positive modulators at $\alpha 6\beta 2/3\gamma 2$ receptors. *Br J Pharmacol* 169: 384-399.

Vasovic D, Divovic B, Treven M, Knutson DE, Steudle F, Scholze P, *et al.* (2019). Trigeminal neuropathic pain development and maintenance in rats are suppressed by a positive modulator of $\alpha 6$ GABAA receptors. *Eur J Pain* 23: 973-984.

Vega Alanis BA, Iorio MT, Silva LL, Bampali K, Ernst M, Schnurch M, *et al.* (2020). Allosteric GABAA Receptor Modulators-A Review on the Most Recent Heterocyclic Chemotypes and Their Synthetic Accessibility. *Molecules* 25.

Wafford KA, Ebert B (2006). Gaboxadol--a new awakening in sleep. *Curr Opin Pharmacol* 6: 30-36.

White JJ, Arancillo M, King A, Lin T, Miterko LN, Gebre SA, *et al.* (2016). Pathogenesis of severe ataxia and tremor without the typical signs of neurodegeneration. *Neurobiol Dis* 86: 86-98.

Winkler GF, Young RR (1974). Efficacy of chronic propranolol therapy in action tremors of the familial, senile or essential varieties. *N Engl J Med* 290: 984-988.

Wu CY, Chiou LC (2016). A novel therapeutic target for hyper-dopaminergic neuropsychiatric disorders with stereotypy behaviors: Cerebellar $\alpha 6$ subunit-containing GABAA receptors. Master, National Taiwan University, Taipei, Taiwan, 2016.

Zesiewicz TA, Elble R, Louis ED, Hauser RA, Sullivan KL, Dewey RB, Jr., *et al.* (2005). Practice parameter: therapies for essential tremor: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 64: 2008-2020.

Zesiewicz TA, Elble RJ, Louis ED, Gronseth GS, Ondo WG, Dewey RB, Jr., *et al.* (2011). Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards subcommittee of the American Academy of Neurology. *Neurology* 77: 1752-1755.

Zhan X, Graf WM (2012). Harmaline attenuates voltage--sensitive Ca^{2+} currents in neurons of the inferior olive. *J Pharm Pharm Sci* 15: 657-668.

Zhang P, Zhang W, Liu R, Harris B, Skolnick P, Cook JM (1995). Synthesis of novel imidazobenzodiazepines as probes of the pharmacophore for "diazepam-insensitive" GABAA receptors. *J Med Chem* 38: 1679-1688.

Figures

Fig 1. Chemical structures of pyrazoloquinolinone compounds, the $\alpha 6$ subunit-containing GABA_A receptor ($\alpha 6$ GABA_AR)-selective positive allosteric modulators (PAMs). (A) Compound 6, originally encoded as PZ-II-029, was the first synthesized $\alpha 6$ GABA_AR-selective PAMs. (B) LAU463 is a structural analogue of Compound 6. (C) DK-I-56-1 and (D) DK-I-58-1 are deuterated derivatives of Compound 6 and LAU463, respectively.

Fig 2. Representative figures of frequency domain-based motion power spectra and motion power spectrograms in mice treated with saline, harmaline and Compound 6 plus harmaline.

(A) The motion waves in mice measured by a Tremor Monitor (San Diego Instruments, CA, USA) were converted into frequency domain motion power spectra (upper panels) and motion power heatmap spectrograms (lower panels) by a fast Fourier transformer. Note that harmaline induced tremor response at the frequency of 10-16 Hz in ICR mice, and Compound 6 attenuated harmaline-induced tremor activity. **(B)** A sample of 10-min motion power spectrograms (lower panel) and concurrent motor activity detected by SMART motion detector showing that the tremor activity occurs during movement (red region), but not immobile (black region) periods, suggesting that harmaline induces action tremor, but not rest tremor, in mice.

Fig. 3. Dose-dependent effects of harmaline on tremor activity and burrowing activity in mice.

(A) Time courses of the tremor activity induced by saline and 10, 20 and 30 mg/kg (*s.c.*) of harmaline. The tremor activity was measured for 10 min (pink shade), 15 min before, 5 min after, and then every 20 min after harmaline injection. Immediately after the tremor activity measurement, the mouse was placed in a chamber and its burrowing activity (grey shade) was measured for 10 min. Tremor activity and burrowing activity were measured alternatively for 120 min. The means of tremor activity for each 10-min period were plotted against time and compared among treatment groups. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. Saline group, two-way ANOVA with repeated measures over time followed by Holm-Sidak's *post hoc* test. (B) The total tremor activity, as measured by the area under curve (AUC) of the tremor activity against time, and (C) the burrowing activity, as measured by total displaced pellets, during 120 min-recording period in saline- or harmaline-treated mice. * $P < 0.05$, ** $P < 0.01$ vs. Saline group, Kruskal-Wallis test followed by

Dunn's *post hoc* test. The numbers in the parentheses denoted the n number of mice tested in each group. Data are expressed as mean $\pm$ S.E.M.

Fig. 4. Effects of Compound 6, an $\alpha 6$ GABA_AR-selective PAM, and propranolol on harmaline-induced tremor and burrowing impairment in mice. (A) Time courses of effects of 3 and 10 mg/kg (*i.p.*) of Compound 6 and its vehicle, and propranolol (20 mg/kg) on harmaline (20 mg/kg, *s.c.*)-induced tremor. The tremor activity and burrowing activity were measured alternatively for 80 min as described in Fig. 3A. Compound 6 and vehicle were injected (*i.p.*) 5 min before harmaline treatment. * $P < 0.05$, *** $P < 0.001$ vs. Saline group, two-way ANOVA with repeated measures over time followed by Holm-Sidak's *post hoc* test. The total tremor activity (B) and burrowing activity (C) were measured as described in Figure 3 in harmaline-treated mice in various treatment groups. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. Vehicle group, one-way ANOVA followed by Holm-Sidak's *post hoc* test. The numbers in the parentheses denoted the n number of mice tested in each group. Data are expressed as mean $\pm$ S.E.M. Veh: vehicle, C6: Compound 6.

Fig. 5. Effects of intra-cerebellar microinjection (*i.cb.*) of furosemide, an $\alpha 6$ GABA_AR antagonist, on anti-tremor and burrow-restorative effects of Compound 6 in harmaline-treated mice. (A) Time courses of effects of 10 mg/kg (*i.p.*) of Compound 6 and its vehicle without and with *i.cb.* furosemide (10 nmol) co-treatment on the tremor activity of harmaline-treated mice. Tremor activity and burrowing activity were measured alternatively for 80 min as described in Fig. 4A. * $P < 0.05$, *** $P < 0.001$ vs. Saline group, two-way ANOVA with repeated measures over time followed by Holm-Sidak's *post hoc* test. The total tremor activity (**B**) and burrowing activity (**C**) were measured as described in Figure 3 in harmaline-treated mice pretreated with Compound 6 alone or in combination with *i.cb.* furosemide. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. Vehicle group, # $P < 0.05$ vs. Veh (*i.cb.*)+C6 group, one-way ANOVA followed by Holm-Sidak's *post hoc* test. Furo: furosemide, Veh: vehicle, C6: Compound 6.

Fig. 6. Effects of DK-I-56-1, LAU463 and DK-I-58-1, structural analogues of Compound 6, on harmaline-treated mice. LAU463 is a structural analogue of Compound 6. DK-I-58-1 and DK-I-56-1 are their respective deuterated derivatives. These compounds are all $\alpha 6\text{GABA}_A\text{R}$ PAMs with similar $\alpha 6\text{GABA}_A\text{R}$ -selectivity and efficacy while deuterated derivatives have longer half-lives. Similar procedure as described in Fig. 4A were performed with Compound 6 replaced with (A) DK-I-56-1 (half-filled square symbols), (B) LAU463 (diamond symbols) or (C) DK-I-58-1 (half-filled diamond symbols), at doses of 3 and 10 mg/kg (*i.p.*). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. Saline group, two-way ANOVA with repeated measures over time followed by Holm-Sidak's *post hoc* test. The total tremor activity (D) and burrowing activity (E) were measured alternatively for 80 min as described in Figure 3 in various treatment groups. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. Vehicle group, one-way ANOVA followed by Holm-Sidak's *post hoc* test. Veh: vehicle, PAM: positive allosteric modulator.