1 Genomic diversification of dehydrin gene family in 2 vascular plants: three distinctive orthologue groups and a 3 novel KS-dehydrin conserved protein motif.

4 Alejandra E. Melgar^{1,2} and Alicia M. Zelada^{1,2*}

5 ¹Laboratorio de Agrobiotecnología, Departamento de Fisiología, Biología Molecular y Celular,
6 Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

7 ²Instituto de Biodiversidad y Biología Experimental y Aplicada, Consejo Nacional de
8 Investigaciones Científicas y Técnicas-Universidad de Buenos Aires (IBBEA, CONICET-UBA),
9 Buenos Aires, Argentina

10 *Author for correspondence: azelada@fbmc.fcen.uba.ar

12 Abstract

13 Dehydrins (DHNs) are a family of plant proteins that play important roles on abiotic stress tolerance
14 and seed development. They are classified into five structural subgroups: K-, SK-, YK-, YSK-, and
15 KS-DHNs, according to the presence of conserved motifs named K-, Y- and S- segments. We
16 carried out a comparative structural and phylogenetic analysis of these proteins, focusing on the
17 less-studied KS-type DHNs. A search for conserved motifs in DHNs from 56 plant genomes
18 revealed that KS-DHNs possess a unique and highly conserved N-terminal, 15-residue amino acid
19 motif not previously described. This novel motif, that we named H-segment, is present in DHNs of
20 angiosperms, gymnosperms and lycopphytes, suggesting that HKS-DHNs were present in the first
21 vascular plants. Phylogenetic and microsynteny analyses indicate that the five structural subgroups
22 of angiosperm DHNs can be assigned to three groups of orthologue genes, characterized by the
23 presence of the H-, F- or Y- segments. Importantly, the hydrophilin character of DHNs correlate
24 with the phylogenetic origin of the DHNs rather than to the traditional structural subgroups. We
25 propose that angiosperm DHNs can be ultimately subdivided into three orthologous groups, a
26 phylogenetic framework that should help future studies on the evolution and function of this protein
27 family.

29 Introduction

30 Plants have to deal with different environmental stresses that can negatively affect their growth and
31 development. Loss of intracellular water in response to abiotic stresses like drought, salinity and
32 low temperature results in the accumulation of Late Embryogenesis Abundant (LEA) proteins in
33 different vegetative tissues. These proteins, which belong to several different families, were first
34 identified in cotton seeds as proteins upregulated during a programmed maturation drying event
35 during seed development^{1,2}. LEA proteins belong to a large group of proteins known as
36 "hydrophilins" characterized by glycine-rich, highly hydrophilic disordered amino acid sequences³.
37 Based on sequence similarity, LEAs are classified into 7 families distinguished by the presence of
38 different conserved motifs^{4,5}.

Dehydrins (DHNs) constitute a biochemically and evolutionarily distinct group of LEAs with a highly modular structure consisting of a combination of different conserved motifs, variable in number and type, interspersed within weakly conserved amino acid segments. The presence of at least one conserved lysine rich-motif, named the K-segment, is usually used as a *sine qua non* condition to define a protein as a dehydrin⁶. Two other conserved motifs have been described, the Y- and S-segments, that in conjunction with the K-segment are the basis for the general classification of DHNs into 5 structural subgroups: KnS, SKn, YnK, YnSKn and Kn-DHNs, where n refers to the number of repetitions of a given motif⁷.

The Y-segment, whose conserved consensus sequence is [V/T]D[E/Q]YGNP, is usually located at the N-terminus of the protein in one or several tandem copies, while the S-segment, a tract of Ser combined with Asp and Glu residues, is always found in one copy per protein⁷. Recently, Strimbeck⁸ described a new conserved motif present in a subgroup of SK-DHNs that consists of a 11-residue amino acid consensus sequence (DRGLDFDLGKK), named the F-segment. These conserved motifs are usually surrounded by less conserved sequences denoted Phi-segments, characterized by a higher proportion of Gly, Thr, and Glu residues.

Several studies have identified, classified and determined the role of DHNs in different plants. A positive relationship between the level of DHN transcripts and/or protein accumulation and plant stress tolerance has been reported⁹⁻¹¹. Furthermore, it has been observed that DHN overexpression in transgenic lines increases resistance to unfavourable environmental conditions, such as cold, drought and salinity¹²⁻¹⁴. In vitro experimental evidence from biochemical assays and localization experiments suggests multiple roles for dehydrins, including membrane protection, cryoprotection of enzymes, interaction with DNA and protection from reactive oxygen species¹⁵⁻¹⁷.

In most of these studies, the biochemical and functional characteristics of these proteins were analysed within the framework of conserved structural domains, but a comparative analysis taking an evolutionary point of view has not been fully explored. The phylogenetic relationships of DHNs have been studied in many different plants, but most of these studies are limited to one genus or species¹⁸⁻²⁰. Only recently, a comprehensive understanding of the evolutionary history of DHNs has been attempted. A phylogenetic and structural analysis of a large number of plant DHNs by Riley et al (2019) suggests that the ancestral DHN belonged to a Kn or SKn group, and that YSKn and YKn-DHNs first arose in angiosperms²¹. On the other hand, Artur (2019) showed that angiosperm DHNs with Y- and F- segments belong to two different orthologue groups that can be distinguished by synteny conservation across angiosperms²². The evolutionary origin of KS-DHNs is still elusive, since previous works have neglected this group.

Here, we present a thorough phylogenetic and structural analysis of DHNs obtained from a wide spectrum of plant genomes. Even though KS-DHNs have previously been described only in a handful of species, we show that this DHN group is actually present in all angiosperms as well as in gymnosperms and lycophytes, indicating its ancient origin in vascular plant evolution. We show that KS-dehydrin genes share a conserved synteny neighbourhood in angiosperm genomes and possess a conserved N-terminal domain, that we named H-segment, and propose that all angiosperm DHNs belong to one of three orthologue groups, the H, F and Y groups. We also carried out a comparative analysis of the different domain structures and biochemical characteristics inherent to the hydrophilin quality of DHNS to investigate how they correlate with their evolutionary origin.

Methods

Dehydrin protein sequences database construction

Initially, DHN proteins were obtained by searching plant genomes or transcriptomes with the Hidden Markov Model (HMM) profile assigned to the DHN protein family (PF0027), downloaded from the Pfam database (<http://pfam.xfam.org/>), using the HMMER 3.1 software (<http://hmmerr.org/>). The HMM profile was used to search the Phytozome v13 database (<https://phytozome-next.jgi.doe.gov/>) which harbours 56 genomes from species spanning the whole viridiplantae clade, including one rodophyte, nine chlorophytes, two briophytes (*Ceratodon purpureus* and *Physcomitrella patens*), the lychophyte *Selaginella moellendorffii*, the angiosperm basal species *Amborella trichopoda* and *Nymphaea colorata* and a subset of 9 monocots and 28 eudicots representing different families. To include gymnosperm species in our search, we employed the Gymno PLAZA 1.0 database²³ and the ConGeniE database (<http://congenie.org/>) which contain the transcriptomes of *Ginkgo biloba*, *Picea abies* and *Picea glauca*.

To identify the conserved motif structures of DHN proteins, we used the MEME software (<http://meme-suite.org/>)²⁴. Since we noticed that KS-type DHNs were underrepresented in this preliminary DHN database, we searched the National Center for Biotechnology Information (NCBI) database to retrieve homologues of *Arabidopsis thaliana* HIRD11 using the Blastp algorithm, and the new KS-DHN sequences were used to construct a HMM profile specific for this DHN group. In parallel, HMM profiles were also constructed for F- and Y-DHNs. Finally, the three HMM profiles were used to reanalyse the databases. All DHN sequences identified in this work are shown in Supplementary Table S1.

MEME searching conserved motif in DHNs database.

The conserved motif structures of DHN proteins were identified using MEME software to find recurrent ungapped motifs assuming that each sequence may contain any number of non-overlapping motifs. The results presented correspond to an analysis made with the following parameters: number of motifs = 8, motif width = 6 to 20, and number of sites for each motif = 2 to 600 (Supplementary Figure S1). The E-values of the different motifs predicted by MEME for our DHN database were compared to E-values calculated from the same sequenced randomly shuffled using the same MEME run parameters to confirm the significance of the discovered motifs.

Multiple sequence alignments and phylogenetic tree construction.

In order to establish orthology/paralogy relationships among the sequences, phylogenetic relationships within each DHN family were estimated. The DHN protein sequences were aligned using Clustal Omega²⁵ or T-coffee²⁶ with default parameters, and multiple sequence alignments (MSA) were visualized using Jalview²⁷. The phylogenetic tree was constructed using an MSA that included only angiosperm DHNs, in order to prevent very divergent sequences from reducing the quality of the alignment. Phylogenetic trees were estimated by the Maximum Likelihood (ML) method as implemented in the NGPhylogeny website (<https://ngphylogeny.fr/>)²⁸ using FastTree²⁹ with the LG amino acid substitution model³⁰ and the GAMMA model with invariant sites for rate heterogeneity. A total of one thousand bootstrap samplings were run. The resulting tree was visualized using iTOL³¹.

Microsynteny analysis.

For microsynteny analysis of selected DHN genes, the corresponding proteins were identified in the NCBI database by pairwise BLASTP searches. Annotations with 100% identity were selected and the genomic context analysed using the NCBI Genome Data Viewer (GDV). Protein sequences of ten to twenty genes flanking both sides of DHN genes were compared between species, using the loci of *A. trichopoda* DHN genes as references. Reciprocal BLASTP analysis were used to confirm homology, with sequences that matched with an E-value of $<10^{-5}$ being considered homologous to each other.

Estimation of physicochemical properties of DHNs proteins.

The theoretical physicochemical properties of DHNs such as grand average hydropathicity index (GRAVY), molecular weight (MW), isoelectric point (pI) and glycine percentage were calculated with the ProtParam tool of Expasy (<https://web.expasy.org/protparam/>). The GRAVY index indicates the hydrophobicity of the protein and was calculated as the sum of the hydropathy values (Kyte and Doolittle parameters) of all amino acids divided by the sequence length. Proteins with positive GRAVY scores are hydrophobic whereas proteins with negative GRAVY scores are hydrophilic. The fold index of proteins was estimated using the FoldIndex© software (<https://fold.weizmann.ac.il/fldbin/findex>).

Results

Unbiased genome-wide identification of dehydrins in Viridiplantae genomes

As a first step to understand the evolutionary history of KS-DHNs and their relationship to the other structural subgroups (YnSKn-, YnKn- SKn- and Kn-DHNs), we performed a genome-wide sequence homology search to identify the complete repertoires of DHNs across 56 genomes of species belonging to the Viridiplantae clade, including representative members of chlorophytes (green algae) and streptophytes (see Materials and Methods). The initial screening was made using a Hidden Markov Model (HMM) profile defined for dehydrin family proteins (Pfam2057) obtained from the Pfam 33.1 database³². Surprisingly, when we analysed the sequences retrieved, we noticed that well known KS-DHNs, such as the HIRD11 dehydrin from the dicot *Arabidopsis thaliana* (At1g54410)³³ and the ZmDHN13 from the monocot *Zea mays*¹³ were not detected by the algorithm. That prompted us to hypothesize that a Pfam00257-based HMM is not sensitive enough to recognize KS-DHNs as members of the dehydrin family. To overcome this limitation we built three different HMM profiles: one (KS-HMM) using KS-DHN sequences from angiosperm genomes identified by Blastp searches using *A. thaliana* HIRD11, and two other profiles (F-HMM and Y-HMM) based on angiosperm proteins belonging to the F- and Y-DHNs orthologous groups recently described²².

After searching with Pfam2057 and the three DHN group-specific HMM profiles, we recovered a total of 305 non-redundant DHN sequences from genomes of representative species of briophytes (4), lycophytes (1), gymnosperms (3) and angiosperms (36) (Supplementary Table 1). No sequences were retrieved from the 9 chlorophyta green algae genomes analysed, neither from the genome of the streptophyte alga *Chara brunii*, that belongs to a sister group to embryophytes³⁴, confirming that the DHNs family emerged in land plants³⁵.

Remarkably, the KS-HMM profile displayed an increased sensitivity in recognising DHN homologues, since it was able to identify 92.2% of DHN proteins, while the Pfam00257-based HMM identified 81.7% and the other two HMM profiles only 83% of DHN sequences (Fig. 1). Among a total of 62 DHNs exhibiting the KS-architecture, only 17 could be retrieved using the Pfam00257 profile, confirming its poor performance in recognizing KS-DHNs. While F-HMM has a better performance and could recognize 26 KS-DHNs, only the KS-HMM profile was able to

Figure 1. DHNs identified in Viridiplantae genomes by different HMM profiles. Sequences retrieved from species genomes using F-HMM, Y-HMM, KS-HMM profiles constructed in this study and PF00257 the Pfam profile for dehydrin family proteins are displayed as a Venn diagram. White numbers indicate the number of KS-DHNs present in each subset. Note that most KS-DHNs are not recovered using Pfam00257.

retrieve all KS-DHNs. Indeed, 32 KS-DHNs could only be retrieved using the KS-HMM profile. Conversely, the KS-HMM profile failed to recognize 22 DHN sequences that were identified by the other HMM profiles. No DHNs were retrieved solely by Pfam00257, indicating that group specific-HMM profiles are necessary and sufficient for a thorough search of DHN proteins in angiosperm genomes.

174 Analysis of conserved protein motif and classification of the dehydrin database.

175 To classify the DHNs of our unbiased database into the structural subgroups, we used the MEME
176 program to check for the presence of known dehydrin motifs (K- Y- F- and S-segments) and to
177 discover putative novel motifs (Supplementary Figure S1). The LOGO representations of the
178 conserved motifs detected and the distribution of DHNs sequences in the different structural
179 subgroups are shown in Figure 2.

180 We confirmed the presence of the K-segment in 302 of the 307 dehydrins identified by homology
181 searches based on HMM profiles. The MEME program failed to recognize a sequence similar to K-
182 segment in a few proteins, all of which from non-angiosperm species. However, these proteins all

Figure 2. Identification of conserved protein motifs and structural classification of DHNs. (a) LOGO representation of the different conserved motifs detected by MEME in the set of DHNs of the unbiased database. (b) Number of members of each angiosperm DHN structural subgroup identified in the database. We distinguished FSK2 and FSK3 structural subgroups in accordance to Strimbeck (Strimbeck, 2017). All classified DHNs are listed in Supplementary Table 1. The dotted pattern indicates monocots, while the filled pattern indicates eudicots.

183 possess a degenerate, less conserved K-segment, as well as other DHN motifs, indicating that they
184 are bona fide DHNs. This is the case, for instance, for DHNs from the lycophyte *S. moellendorffii*
185 and the gymnosperm *Ginkgo biloba* (see Fig. 3).

186 We identified 75 DHNs bearing a unique F-segment located in the N-terminal region of the
187 proteins, that we classified within the FSKn-DHN structural subgroup. The F-segment predicted
188 with our protein database is similar to the one described by Strimbeck (2017) (Fig. 2A).
189 Importantly, even a search with a specific F-HMM profile failed to identify FSKn-DHNs in the
190 genomes of four bryophyte and one lycophyte species, but we did find them in the three
191 gymnosperms included in this study, *Picea abies*, *Picea glauca* and *Ginkgo biloba*, confirming that
192 this subtype of DHN probably arose in seed plants. We observed an expansion of the FSKn-DHN
193 gene family in the Pinaceae clade, in accordance with previous observations³⁶, but that was not a
194 general feature of gymnosperm species. Only three DHNs were identified in the gymnosperm
195 *Ginkgo biloba*, two of them with a F-segment (FSK2 and FK2) and a third harbouring a novel

identified in the genomes of the mosses *P. patens* (six proteins), *Ceratodon purpureus* (six), *Sphagnum fallax* (three) and the liverwort *Marchantia polymorpha* (three) belong to the Kn-structural subgroup. The exception is an atypical DHN containing a series of repetitive motifs resembling the Y-segment in the N-terminus that is present in *P. patens* (PpDHNA)³⁷ and *C. purpureus* (Supplementary Figure S10). A phylogenetic analysis indicates the presence of five DHN orthologue groups in *P. patens* and *C. purpureus* (Supplementary Figure S9), which reflects the phylogenetic proximity of the Funariidae and Dicranidae clades³⁸. The DHNs from the more distantly related *S. fallax* (Sphagnophytina) did not cluster with the DHNs of the other mosses, but multiple sequence alignments and reciprocal Blastp analyses suggest that two of the *S. fallax* DHNs (Sphfalx0010s0103.1 and Sphfalx0064s0013.1) are related to groups III and V of *P. patens* and *C. purpureus* (Supplementary Figure S10). The three DHN proteins of the liverwort *M. polymorpha* do not display any obvious homology to moss DHN groups outside the K-segment.

The H-segment is a novel conserved motif present in all KS-dehydrins.

Our MEME analysis identified a highly conserved motif, not previously described, at the N-terminal region of all angiosperm KS-DHNs analysed. This 15-residue segment is characterized by a combination of hydrophobic amino acids Ile and Leu with amphipathic amino acids Lys and Glu, framed by two Gly at positions 3 and 15 conserved in 91% and 87% KS-DHNs (Fig. 2). In addition, the high percentage of conservation of the Lys (97%) and Ile (97%) located in the central positions 7 and 8 strongly suggest an important function in KS-DHNs. Ile residues at positions 4 and 5 are less conserved, and are often replaced by other hydrofobic amino acids like Phe, Val or Met. KS-DHNs are characterized by sequences enriched in His amino acids, as reflected in the name HIRD11 for the *A. thaliana* KS-DHN, which stands for Histidine-Rich Domain 11 kDa protein³⁹. Two His residues are found in positions 6 and 13 in 56% and 77% of KS-DHNs, respectively. Since this novel motif seems to be a signature of KS-DHNs, we propose to name it the H-segment, reflecting the particular feature of these kind of proteins, even when histidines are not the most prevalent aminoacids in the motif.

A structural prediction of representative angiosperms KS-DHNs, obtained by Phyre2⁴⁰, indicates with high confidence the presence of a helical α -helix spanning the H-motif in all proteins analysed (Fig. 4). This helical wheel projection is conformed by the alternation of hydrophobic and hydrophilic amino acids and is surrounded by highly conserved Gly amino acids that could function as a helix breaker due to their high conformational flexibility, which makes it entropically expensive to adopt the relatively constrained α -helical structure. A very similar structure is predicted for the K-segment, suggesting that the H-segment could also have amphiphilic membrane or protein binding properties as described for the K-segment^{15,41}.

The K- and S-segments of KS-DHNs present some particular characteristics compared to FSK and YSK-DHNs. The prevalence of amino acids at positions 6, 16 and 17 differs in the K-segments of KS-DHNs (Fig. 2 and Supplementary Figure S7). Thus, position 6 is occupied by an Asp in all KS-DHNs instead of the Lys that it is typically present in DHNs, with the exception of three FSK-dehydrins of the gymnosperm *P. abies*. Concerning position 16, KS-DHNs usually have an Ile instead of a Leu. It is notable that in the species with more KS-DHN genes, proteins with one or the other amino acid in this position can be found (see Supplementary Table 1; *Solanum tuberosum*; *Populus trichocarpa*, *Phaseolus vulgaris*). Even though Ile and Leu amino acids are generally considered conservative, there is evidence that these amino acids are not always interchangeable, affecting the affinity and specificity of protein-protein and protein-membrane interactions⁴², which might potentially lead to functional diversification of the KS-DHNs by modulating K-segment

behaviour. In contrast to other DHNs, KS-dehydrins do not show a clear prevalence of Pro at position 17; instead, His is the most frequent amino acid at this position, while Pro is only found in the DHNs of Rutaceae and in a subgroup of Malpighiales species, and Thr is prevalent in monocot KS-DHNs at this position. The capacity to tolerate different kinds of amino acids at that position could indicate that it is not essential for K-segment functionality. In spite of these differences, the predicted α -helix structure delimited by conserved Gly amino acids of the K-segment is conserved in KS-DHNs (Fig. 4).

Figure 4. Predicted H-DHN structures. The secondary structure of H-DHN proteins from *A. trichopoda*, *S. bicolor* and *A. thaliana* (HIRD11) were predicted by the Phyre2 program. Note that a helical α -helix is predicted to be present near the N-terminus of the proteins, spanning the H-segment. Red colour indicates high confidence in the prediction.

As for S-segments, which are characterised by a stretch of Ser residues, there are differences in the length of the Ser-amino acid stretch and neighbouring amino acids between KS-DHNs and other DHNs. We observed that the core of 6 to 9 Ser residues usually ended with negatively-charged Asp or Glu amino acids in all structural subgroups of DHNs. On the other hand, the triad Leu-His-Arg that precedes the Ser stretch, which is highly conserved in all FSK-DHNs and in the majority of YSK-DHNs (segment S1) and the one found in KS-DHNs (segment S2, see also the alignments in Supplementary Figure S2 and Figure S7). The S-segment of all types of DHNs has been shown to

be a hotspot for phosphorylation by kinases^{13,43,44}, and the differences between the S1- and S2-segment could result in different kinase specificities. For instance, the triad Leu-His-Arg constitutes part of the recognition sequence for SnRK2 kinases⁴⁵, which have been recently demonstrated to phosphorylate *A. thaliana* dehydrins ERD4 and ERD10 in response to osmotic stress⁴⁶.

A 11-residue Lys-rich motif has been consistently detected in all KS-DHNs as well as in all FSK2 and the majority of FSK3-DHNs (Supplementary Figure S1 to Figure S3; Fig. 5). The whole motif comprises 9-11 Lys residues preceded by Gly and Asp amino acids in positions 2 and 3 (Fig. 2). In KS- and FSK-DHNs, the Lys-rich motif is located between the S-segment and the K-segment while, at the same position, YK- and YSK-DHNs usually have a RRKK or RRKKK sequence framed by Gly residues, a motif that resembles monopartite nuclear localization signals^{47,48}. It has been demonstrated that monopartite NLS require specific residues flanking the core basic cluster for their complete activity, and in particular the inclusion of Asp- and Glu-aminoacid seems to be detrimental for its activity⁴⁸. Some Lys-rich motifs could constitute a NLS sequence, but the presence of Asp or Glu amino acids at positions 2 and 8 in KS- and FSK-DHNs suggests that the conservation of the Lys-rich motif could fulfill a distinct function. In conclusion, the KS-DHNs can be better described as having a H-K-S structural organization, with H being a newly described segment, exclusively present in this group of DHNs.

Phylogenetic analysis reveals three basic groups of DHNs in angiosperms

Having identified the motifs that characterize the KS-structural subgroup, we sought to infer the phylogenetic relationships between the angiosperm DHN protein sequences that we identified. We decided to use only angiosperm DHNs to build a phylogenetic tree due to the sparse taxonomic sampling of other land plant DHNs. We used the approximately-maximum-likelihood principle as implemented in FastTree 2⁴⁹ and estimated statistical robustness with the bootstrap method.

The resulting tree is roughly organised in three branches or groups (Fig. 5). All KS-DHNs, characterised by the presence of the H-segment (H-DHNs), are grouped together in a branch with high bootstrap support (96%). The other DHNs are separated into two branches, one harbouring most DHNs that contain the F-segment (FSK), while the other contains DHNs carrying the Y-segment (YnSK, YnKn). Interestingly, the few DHNs that contain only the K-segment or a combination of K- and S-segments are placed within the F or Y branches, indicating that these DHNs actually belong to either of these groups. Overall, the phylogenetic tree suggests that each angiosperm DHN belongs to one of three phylogenetic families, basically distinguished by the presence of the H-, F- or Y-segments. This conclusion is reinforced by the observation of the DHNs of plants at key phylogenetic positions. Thus, the basal angiosperm *Amborella trichopoda* possesses three DHNs, each one belonging to the H, F or Y groups (Fig. 5). Similarly, the three DHNs from the basal dicot *Nelumbo nucifera* are also each one placed into the H, F and Y groups. Overall, the phylogenetic results suggest that these three groups of DHNs were present since the beginning of angiosperm evolution.

H-DHNs belong to a separate synteny community in angiosperms

Even though the phylogenetic tree described above separates angiosperm DHNs into three groups, the large number of different motifs and their divergent arrangement in DHNs makes the sequences difficult to align and reduces the certainty of the phylogenetic reconstruction. Since synteny analyses of orthologue genes can give important hints about the evolution of genomes and gene families⁵⁰, we performed an analysis of the genomic neighbourhood (microsynteny) of DHN genes in order to reinforce our phylogenetic results.

Figure 5. Phylogenetic tree of DHN proteins. Amino acid sequences of DHN proteins from angiosperms were aligned and an approximately maximum-likelihood reconstruction of the phylogenetic relationships was generated using FastTree 2 (Price et al. 2010). Nodes with bootstrap support over 95% are indicated by a violet dot. The motifs of each protein are indicated by coloured boxes, as indicated. Note that all H-DHNs are grouped together in a branch with high support, and most F- and Y-DHNs are also grouped together, forming three groups. DHN proteins from the basal angiosperm *A. trichopoda* (red) and the basal eudicot *N. nucifera* (blue) are indicated to show that they possess one DHN in each group.

328 Recently, Artur et al. (2018) analysed DHN genes plant genomes and identified two main synteny
329 blocks (or communities) among angiosperms, corresponding to DHNs containing the F (community
330 1) and Y (community 2) motifs²². Their analysis, however, did not include DHNs of the KS group,
331 presumably due to the difficulty of retrieving these sequences using the Pfam motif PF00257. To
332 verify whether DHNs containing the H-segment would also be part of a syntenic community, we
333 compared 40 genes surrounding the unique H-DHN of the basal angiosperm, *Amborella trichopoda*
334 to the genomic neighbourhoods of H-DHN loci of the waterlily *Nymphaea colorata*⁵¹, the basal
335 eudicot sacred locus, *Nelumbo nucifera*⁵², the legume *Medicago truncatula*⁵³, the model plant *A.*

336 *thaliana* (HIRD11)⁵⁴ and the monocot grass *Sorghum bicolor*⁵⁵. All of these species possess only
 337 one H-dehydrin paralogue except for *M. truncatula*, which has two (Supplementary Table 1). We
 338 chose *A. trichopoda* as the basis for synteny comparison since this flowering plant belongs to a
 339 sister lineage to all other angiosperms (Amborellales), did not undergo the whole genome
 340 duplications that affected other lineages and its genome exhibits conserved synteny with other
 341 angiosperms, features that facilitate the study of gene family evolution in plants^{56,57}. As shown in
 342 Figure 6, 17 genes that surround the H-DHN of *A. trichopoda* (LOC18421535) are also present
 343 around the H-DHN gene of *N. colorata*, which belongs to a group, the Nymphaeales, that is a sister
 344 lineage to all angiosperms except for Amborellales⁵⁸. A smaller number of conserved genes are
 345 present around the H-DHN genes from the eudicots *N. nucifera*, *A. thaliana* (HIRD11) and *M.*
 346 *truncatula* and the monocot *S. bicolor* (Fig. 6). The microsynteny of H-DHN genes of other
 347 angiosperms is likewise conserved (not shown). H-DHNs possess two exons, with the whole coding
 348 region contained within the first exon and the second exon being no-coding, while F- and Y-DHNs
 349 usually have two coding exons (not shown). The conserved exon-intron structure also points to a
 350 common origin of H-DHNs. In conclusion, the microsynteny of H-DHN genes is conserved in
 351 angiosperms, indicating their true orthologous status and common evolutionary origin.

Figure 6. Microsynteny analysis of angiosperm H-DHNs genes. The genomic neighbourhood of the H-DHN gene of *A. trichopoda* (LOC18421535) is compared to that of other angiosperms. H-DHN genes are indicated as black dots, and a colour code indicates homologous genes present in the other species. Grey dots indicate genes only present in *A. trichopoda*. Some intervening genes in species other than *A. trichopoda* are not shown for clarity.

352 Along with one H-DHN gene, the genome of *A. trichopoda* contains two other DHN genes
 353 (LOC18424350 and LOC18426770). As mentioned above, in our phylogenetic tree, LOC18424350
 354 is grouped together with F-DHNs and LOC18424350 with Y-DHNs (Fig. 5). Curiously, the F and Y
 355 motifs of these proteins are quite degenerated and are not readily recognised by the MEME
 356 program. A comparison of the genomic neighbourhoods of LOC18424350 and LOC18426770 of *A.*
 357 *trichopoda* with F- and Y-DHNs of *N. colorata* and *N. nucifera*, which likewise have only three
 358 DHN genes, reveals microsynteny conservation around these genes (Supplementary Figure S11),
 359 confirming that LOC18424350 and LOC18426770 belong to the F- and Y-DHN syntenic
 360 communities, respectively.

361 In summary, it is apparent that DHN genes of angiosperms can be generally divided into three
 362 syntenic communities, each one characterised, among other features, by the presence of the H, F or
 363 the Y motif. We propose that these orthologous groups be called F-dehydrins (community 1), Y-
 364 dehydrins (community 2) and H-dehydrins (community 3). The presence of only three dehydrin

genes in the basal angiosperm *Amborella*, as well as in the early diverging Nymphaeales and the basal eudicot *N. nucifera*, suggests that the genomes of the first flowering plants had one H-, F- and Y-dehydrin gene each. Subsequent whole genome duplications in eudicots and monocots greatly increased the repertoire of these genes, specially those encoding F- and Y-dehydrins.

Each DHN orthologous group presents distinctive hydrophilin biochemical properties.

To analyse if the existence of three DHNs orthologous groups could result in proteins with distinctive characteristics, we compared the biochemical and biophysical properties of angiosperm DHNs from the H-, F- and Y-orthologous groups. Specifically, we determined general biochemical features such as molecular weight (MW) and isoelectric point (pI), as well as parameters related to the hydrophilin character of the proteins (Supplementary Table 1). We observed that each DHN orthologous group has a different MW distribution, with a characteristic statistical median (Fig.7A). H-DHNs are the smallest DHNs with the narrowest range of MW (10-16 kDa), reflecting that the number of residues and domain structure of the members of this DHN group are relatively constant. F-DHNs also have a compact MW distribution that ranges from 18 kDa to 35 kDa. Y-DHNs, on the other hand, present a main subgroup of proteins ranging from 10 kDa to 25 kDa and a number of DHNs with MW over 30 kDa that belong exclusively to monocot species. The high MW of this latter subgroup is not due to an increased number of conserved Y or K domains, but to the presence of long Gly-rich regions separating these domains (Supplementary Figure S4 to Figure S6). As for the isoelectric point, most H-DHNs present acidic pI values, with neutral and basic isoforms being found in some species (Fig 7B). F-DHNs have a very homogeneous acidic pI profile, with a unimodal distribution between 5 and 6. In contrast, Y-DHNs display a bimodal distribution consisting of two main subgroups of DHNs with basic and acid pI values, and a smaller subgroup with pI values close to neutrality. Interestingly, we were able to determine that almost all plant species have at least one basic and one acidic Y-DHN isoform, suggesting that a functional specialization of both types of proteins may have occurred during evolution. In monocots, the number of basic Y-DHNs is always greater than the acidic ones, and the opposite occurs in dicots. It should be noted that the early-diverging angiosperms *A. thrichopoda* and *N. colorata* encode a single basic Y-DHN in their genomes, which may represent the original pI character of these proteins in angiosperms (Supplementary Table 1).

When analysing the pI distribution in the five traditional DHN structural subgroups, it can be noted that the bimodal character observed in SK- and K-type DHNs strongly correlates with their evolutionary origin (Fig. 7C). For example, five of the K-type DHNs that display acidic pI, which corresponds to the pI of F-DHNs, belong to this orthologous group. Similarly, all members of the SK- and K-DHNs structural subgroups with high pI values belong to the Y-DHN orthologous group (Fig. 7B-7C). This suggests that DHN orthologous groups are better indicators of the pI character of DHNs compared to the traditional structural classification.

As for glycine content, both F- and H-DHNs present a compact and homogeneous distribution of percentage of glycine residues (Fig 7D). The DHNs with the lowest percentage of glycine (around 10%) are the F-DHNs. Remarkably, DHNs of the FSK3 structural subgroup are characterized by the presence of many proline stretches, which might play an equivalent role to that of glycine in terms of the disruption of the protein structure (Supplementary Figure S2 and Figure S3). Notably, a larger dispersion in the percentage of glycine residues is observed in Y-DHNs (range: 5% to 35%).

408 All DHNs display a negative GRAVY index, showing the characteristic hydrophilicity of this type
 409 of proteins (Fig. 7E). The scores for F-DHNs are distributed in the range of -1 to -1.7, overlapping
 410 with the other two groups. The Y-DHNs include the least hydrophilic proteins, with scores in the
 411 range of -0.5 to -1.5. H-DHNs, in contrast, are the most hydrophilic proteins, with GRAVY indexes
 412 ranging from -2.8 to -1.3. The atypical H-DHNs from the Malpighiales species *S. purpurea* and *P.*
 413 *trichocarpa* are the least hydrophilic dehydrins in this group, with a GRAVY index around -1.3. We
 414 also evaluated the Fold Index of DHNs using the FoldIndex algorithm, which estimates the mean
 415 net charge and hydrophobicity of a given protein sequence to predict if it is intrinsically unfolded³⁹.
 416 The fold index of DHNs shows a similar distribution to that of the GRAVY index, with H-DHNs
 417 being the most intrinsically unfolded, while F- and Y-DHNs have a less unfolded character
 418 (Fig.7F).

419 In summary, in general terms, the biochemical and biophysical characteristics of DHNs correlate
 420 well with the three orthologous groups (Fig. 7G). Since these features are likely related to the
 421 function of DHNs, this suggests that functional studies of these proteins should take into
 422 consideration the phylogenetic framework proposed here.

Figure 7. Distribution of biochemical and biophysical properties of angiosperm DHNs. Scatter plots show the distribution of molecular weight (a), isoelectric point (b and c), glycine content (d), GRAVY index (e), Fold Index (f) or glycine content, GRAVY and Fold index simultaneously (g) in orthologous or structural subgroups of DHNs. Members of the three orthologous groups of DHNs are colour-coded: Y- (green), F- (orange) and H-DHNs (violet).

Discussion

Dehydrins are characterised by a great diversity of structural domains, arranged in various ways, which constitute the basis for the current classification into six structural subgroups, namely Kn-, SKn-, YnKn-, YnSKn-, KS-DHNs and the recently proposed FSKn-DHNs. However, the underlying evolutionary relationships between these DHNs in angiosperms and other plant groups have been unclear. In this work, we present a phylogenetic framework for DHNs that sheds light on the relationships between these proteins, specially in angiosperms. The main points of our work are: i) searches of DHN in plant genomic databases need to be done with a combination of HMM profiles to retrieve all types of DHN proteins; ii) KS-DHNs possess a new, conserved structural domain present at the N-terminus, which we named the H-domain; iii) phylogenetic and synteny analyses show that all angiosperm DHNs can be subdivided into three DHN orthologous groups, distinguished by the presence of the H-, F- or Y-domains, and iv) the physicochemical characteristics that are typical for DHNs correlate with each orthologous group, indicating that the evolutionary origin of DHNs should be taken into consideration when studying their function.

The reconstruction of the evolutionary history of DHNs is a complex task, due to the modular nature of these proteins, which are characterized by the presence of various small conserved segments surrounded by less conserved sequences of various lengths. Thus, the coupling of phylogenetic reconstruction with microsynteny analyses was crucial for the determination of the evolutionary relationships between DHNs. Angiosperm DHNs can be divided into three orthologous groups, H-DHNs, F-DHNs and Y-DHNs, which can, in most cases, be readily recognised by the presence of the H-, F- or Y-segments. All angiosperms analysed by us possess at least one DHN member of each homologous group, including the basal angiosperms *A. trichopoda* and *N. colorata*, indicating that the first angiosperms had genes encoding the three types of DHNs. Synteny analyses could not be extended to non-angiosperm species due to the fast rate of synteny loss that is typical for plants^{50,60}.

Our analysis indicated that H-, F- and Y-DHNs are clearly distinguished from each other in features that characterize hydrophilins and intrinsically-disordered proteins (IDPs). Since all dehydrins of K and SK-structural subgroups actually belong to the F- and Y- syntenic groups, we were able to show that the classification based on the structural subgroups ends up putting in the same category DHNs with very different physicochemical properties. It has been observed that the cryoprotective capacity of DHNs depends on the size (hydrodynamic radius) and the intrinsic disorder, highlighting the importance of the composition and size of the Phi segments, which are generally less conserved than the structural motifs⁶¹. It has been also demonstrated that it is the size and sequence composition of DHNs that is the most important for preventing aggregation, while for freeze damage it is the sequence composition that is most significant⁶². Thus, it seems that the simple presence of K and S segments would not be necessarily good predictors of the functional characteristics of DHNs.

It should be noted that the diversity of DHNs is not encompassed by the HMM model that is usually employed to search for DHN genes in the scientific literature, namely Pfam00257. Indeed, we show here that most H-DHNs (which belong to the KS-DHN structural subgroup) are not recognized by this model, which might be the reason that genomic-wide analyses of DHNs usually failed to retrieve many members of this orthologous group^{18,20,22}. In view of this, we propose that studies aimed at identifying DHNs should use HMM profiles based on H-, F- and Y-DHNs separately in order to pinpoint all members of this protein family.

Importantly, we describe that KS-DHNs possess a new motif that we named the H-segment, due to the presence of two conserved His residues. This segment is always located at the N-terminus of the proteins and is predicted to have an α -helical structure. Thus, KS-DHNs can be better described as bearing a H-K-S organization of motifs. As mentioned above, phylogenetic and synteny analyses indicate that angiosperm DHNs are all evolutionarily related. The presence of DHNs with a distinct H-K-S organization in the lycophyte *S. moellendorffii* and the gymnosperm *Ginkgo biloba* strongly suggests that H-DHNs appeared in the early evolution of vascular plants. Although some KS-DHNs have been described before in the scientific literature, our work is the first, to our knowledge, to provide a thorough description of this group of DHNs.

The best studied member of the H-DHN group is the HIRD11 protein from *A. thaliana*. AtHIRD11 is expressed ubiquitously, with somewhat higher levels in flowers⁵⁴. Functional studies showed that HIRD11 binds to metal ions and can protect proteins from heavy metal damage^{54,63} and can also reduce free radical generation⁶⁴. Interestingly, both the binding to metals and the inhibition properties of HIRD11 depend on His residues, which are present in the H-segment. Importantly, Yokoyama et al (2020) have recently showed that both the K- and the H-segments (which the authors called K and NK1, respectively) of AtHIRD11 can protect proteins from freezing damage with similar efficiencies. Structurally, the presence of the K- and H-segments were needed for AtHIRD11 to transition from a disordered to an ordered state⁶⁵. Overall, the functional results by Yokoyama et al (2020) show that the H-segment is an important component of H-DHNs, as implied by its high degree of phylogenetic conservation, and suggests that K- and H-segments might play overlapping roles in the activity of H-DHNs.

In conclusion, we consider that the classification of angiosperm DHNs into three homologous groups, as proposed here, better reflects the diversity of DHNs and should complement the traditional classification into six structural subgroups in the study of the function of these proteins.

References

1. Dure, L., Greenway, S. C. & Galau, G. A. Developmental biochemistry of cottonseed embryogenesis and germination: changing messenger ribonucleic acid populations as shown by in vitro and in vivo protein synthesis. *Biochemistry* **20**, 4162–4168 (1981).
2. Galau, G. A., Bijaisoradat, N. & Hughes, D. W. Accumulation kinetics of cotton late embryogenesis-abundant mRNAs and storage protein mRNAs: coordinate regulation during embryogenesis and the role of abscisic acid. *Dev. Biol.* **123**, 198–212 (1987).
3. Garay-Arroyo, A., Colmenero-Flores, J. M., Garciarrubio, A. & Covarrubias, A. A. Highly Hydrophilic Proteins in Prokaryotes and Eukaryotes Are Common during Conditions of Water Deficit. *J. Biol. Chem.* **275**, 5668–5674 (2000).
4. Dure, L. A repeating 11-mer amino acid motif and plant desiccation. *Plant J. Cell Mol. Biol.* **3**, 363–369 (1993).

505 5. Battaglia, M., Olvera-Carrillo, Y., Garcarrubio, A., Campos, F. & Covarrubias, A. A. The
506 enigmatic LEA proteins and other hydrophilins. *Plant Physiol.* **148**, 6–24 (2008).

507 6. Close, T. J., Kortt, A. A. & Chandler, P. M. A cDNA-based comparison of dehydration-induced
508 proteins (dehydrins) in barley and corn. *Plant Mol. Biol.* **13**, 95–108 (1989).

509 7. Campbell, S. A. & Close, T. J. Dehydrins: genes, proteins, and associations with phenotypic
510 traits. *New Phytol.* **137**, 61–74 (1997).

511 8. Richard Strimbeck, G. Hiding in plain sight: the F segment and other conserved features of
512 seed plant SKn dehydrins. *Planta* **245**, 1061–1066 (2017).

513 9. Singh, J., Reddy, P. S., Reddy, C. S. & Reddy, M. K. Molecular cloning and characterization of
514 salt inducible dehydrin gene from the C4 plant Pennisetum glaucum. *Plant Gene* **4**, 55–63
515 (2015).

516 10. Cao, Y., Xiang, X., Geng, M., You, Q. & Huang, X. Effect of HbDHN1 and HbDHN2 Genes on
517 Abiotic Stress Responses in Arabidopsis. *Front. Plant Sci.* **8**, (2017).

518 11. Zhang, H. *et al.* Molecular Cloning and Functional Characterization of the Dehydrin (IpDHN)
519 Gene From Ipomoea pes-caprae. *Front. Plant Sci.* **9**, 1454 (2018).

520 12. Hill, W., Jin, X.-L. & Zhang, X.-H. Expression of an arctic chickweed dehydrin, CarDHN,
521 enhances tolerance to abiotic stress in tobacco plants. *Plant Growth Regul.* **80**, 323–334 (2016).

522 13. Liu, Y., Wang, L., Zhang, T., Yang, X. & Li, D. Functional characterization of KS-type
523 dehydrin ZmDHN13 and its related conserved domains under oxidative stress. *Sci. Rep.* **7**, 7361
524 (2017).

525 14. Luo, D. *et al.* CaDHN5, a Dehydrin Gene from Pepper, Plays an Important Role in Salt and
526 Osmotic Stress Responses. *Int. J. Mol. Sci.* **20**, (2019).

527 15. Hughes, S. & Graether, S. P. Cryoprotective mechanism of a small intrinsically disordered
528 dehydrin protein. *Protein Sci. Publ. Protein Soc.* **20**, 42–50 (2011).

529 16. Eriksson, S., Eremina, N., Barth, A., Danielsson, J. & Harryson, P. Membrane-Induced Folding
530 of the Plant Stress Dehydrin Lti30. *Plant Physiol.* **171**, 932–943 (2016).

531 17. Boddington, K. F. & Graether, S. P. Binding of a Vitis riparia dehydrin to DNA. *Plant Sci.* **287**,
532 110172 (2019).

- 533 18. Jing, H. *et al.* Genome-Wide Identification, Expression Diversification of Dehydrin Gene Family
534 and Characterization of CaDHN3 in Pepper (*Capsicum annuum* L.). *PLoS One* **11**, e0161073
535 (2016).
- 536 19. Abedini, R. *et al.* Plant dehydrins: shedding light on structure and expression patterns of
537 dehydrin gene family in barley. *J. Plant Res.* **130**, 747–763 (2017).
- 538 20. Nagaraju, M. *et al.* Genome-wide in silico analysis of dehydrins in *Sorghum bicolor*, *Setaria*
539 *italica* and *Zea mays* and quantitative analysis of dehydrin gene expressions under abiotic
540 stresses in *Sorghum bicolor*. *Plant Gene* **13**, 64–75 (2018).
- 541 21. Riley, A. C., Ashlock, D. A. & Graether, S. P. Evolution of the modular, disordered stress
542 proteins known as dehydrins. *PLoS One* **14**, e0211813 (2019).
- 543 22. Artur, M. A. S., Zhao, T., Ligterink, W., Schranz, E. & Hilhorst, H. W. M. Dissecting the
544 Genomic Diversification of Late Embryogenesis Abundant (LEA) Protein Gene Families in
545 Plants. *Genome Biol. Evol.* **11**, 459–471 (2019).
- 546 23. Proost, S. *et al.* PLAZA 3.0: an access point for plant comparative genomics. *Nucleic Acids*
547 *Res.* **43**, D974–D981 (2015).
- 548 24. Bailey, T. L. *et al.* MEME Suite: tools for motif discovery and searching. *Nucleic Acids Res.* **37**,
549 W202–W208 (2009).
- 550 25. Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple sequence alignments
551 using Clustal Omega. *Mol. Syst. Biol.* **7**, 539 (2011).
- 552 26. Notredame, C., Higgins, D. G. & Heringa, J. T-Coffee: A novel method for fast and accurate
553 multiple sequence alignment. *J. Mol. Biol.* **302**, 205–217 (2000).
- 554 27. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview Version
555 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–
556 1191 (2009).
- 557 28. Lemoine, F. *et al.* NGPhylogeny.fr: new generation phylogenetic services for non-specialists.
558 *Nucleic Acids Res.* **47**, W260–W265 (2019).
- 559 29. Lemoine, F. *et al.* Renewing Felsenstein’s phylogenetic bootstrap in the era of big data. *Nature*
560 **556**, 452–456 (2018).
- 561 30. Le, S. Q. & Gascuel, O. An improved general amino acid replacement matrix. *Mol. Biol. Evol.*
562 **25**, 1307–1320 (2008).

563 31. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
564 developments. *Nucleic Acids Res.* **47**, W256–W259 (2019).

565 32. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–
566 D419 (2021).

567 33. Hundertmark, M. & Hinch, D. K. LEA (Late Embryogenesis Abundant) proteins and their
568 encoding genes in *Arabidopsis thaliana*. *BMC Genomics* **9**, 118 (2008).

569 34. Nishiyama, T. *et al.* The Chara Genome: Secondary Complexity and Implications for Plant
570 Terrestrialization. *Cell* **174**, 448–464.e24 (2018).

571 35. Malik, A. A., Veltri, M., Boddington, K. F., Singh, K. K. & Graether, S. P. Genome Analysis of
572 Conserved Dehydrin Motifs in Vascular Plants. *Front. Plant Sci.* **8**, 709 (2017).

573 36. Stival Sena, J., Giguère, I., Rigault, P., Bousquet, J. & Mackay, J. Expansion of the dehydrin
574 gene family in the Pinaceae is associated with considerable structural diversity and drought-
575 responsive expression. *Tree Physiol.* **38**, 442–456 (2018).

576 37. Saavedra, L. *et al.* A dehydrin gene in *Physcomitrella patens* is required for salt and osmotic
577 stress tolerance. *Plant J. Cell Mol. Biol.* **45**, 237–249 (2006).

578 38. Chang, Y. & Graham, S. W. Patterns of clade support across the major lineages of moss
579 phylogeny. *Cladistics* **30**, 590–606 (2014).

580 39. Hara, M., Fujinaga, M. & Kuboi, T. Metal binding by citrus dehydrin with histidine-rich
581 domains. *J. Exp. Bot.* **56**, 2695–2703 (2005).

582 40. Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E. The Phyre2 web
583 portal for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845–858 (2015).

584 41. Hara, M., Endo, T., Kamiya, K. & Kameyama, A. The role of hydrophobic amino acids of K-
585 segments in the cryoprotection of lactate dehydrogenase by dehydrins. *J. Plant Physiol.* **210**,
586 18–23 (2017).

587 42. Brosnan, J. T. & Brosnan, M. E. Branched-Chain Amino Acids: Enzyme and Substrate
588 Regulation. *J. Nutr.* **136**, 207S–211S (2006).

589 43. Alsheikh, M. K., Heyen, B. J. & Randall, S. K. Ion Binding Properties of the Dehydrin ERD14
590 Are Dependent upon Phosphorylation*. *J. Biol. Chem.* **278**, 40882–40889 (2003).

591 44. Rahman, L. N. *et al.* Phosphorylation of Thellungiella salsuginea dehydrins TsDHN-1 and
592 TsDHN-2 facilitates cation-induced conformational changes and actin assembly. *Biochemistry*
593 **50**, 9587–9604 (2011).

594 45. Vlad, F., Turk, B. E., Peynot, P., Leung, J. & Merlot, S. A versatile strategy to define the
595 phosphorylation preferences of plant protein kinases and screen for putative substrates. *Plant J.*
596 **55**, 104–117 (2008).

597 46. Maszkowska, J. *et al.* Phosphoproteomic analysis reveals that dehydrins ERD10 and ERD14
598 are phosphorylated by SNF1-related protein kinase 2.10 in response to osmotic stress. *Plant*
599 *Cell Environ.* **42**, 931–946 (2019).

600 47. Lange, A. *et al.* Classical nuclear localization signals: definition, function, and interaction with
601 importin alpha. *J. Biol. Chem.* **282**, 5101–5105 (2007).

602 48. Kosugi, S. *et al.* Six classes of nuclear localization signals specific to different binding grooves
603 of importin alpha. *J. Biol. Chem.* **284**, 478–485 (2009).

604 49. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree: Computing Large Minimum Evolution Trees
605 with Profiles instead of a Distance Matrix. *Mol. Biol. Evol.* **26**, 1641–1650 (2009).

606 50. Jiao, Y., Li, J., Tang, H. & Paterson, A. H. Integrated syntenic and phylogenomic analyses
607 reveal an ancient genome duplication in monocots. *Plant Cell* **26**, 2792–2802 (2014).

608 51. Zhang, L. *et al.* The water lily genome and the early evolution of flowering plants. *Nature* **577**,
609 79–84 (2020).

610 52. Ming, R. *et al.* Genome of the long-living sacred lotus (*Nelumbo nucifera* Gaertn.). *Genome*
611 *Biol.* **14**, R41 (2013).

612 53. Burks, D., Azad, R., Wen, J. & Dickstein, R. The Medicago truncatula Genome: Genomic Data
613 Availability. *Methods Mol. Biol. Clifton NJ* **1822**, 39–59 (2018).

614 54. Hara, M. *et al.* Biochemical characterization of the Arabidopsis KS-type dehydrin protein,
615 whose gene expression is constitutively abundant rather than stress dependent. *Acta Physiol.*
616 *Plant.* **33**, 2103–2116 (2011).

617 55. McCormick, R. F. *et al.* The Sorghum bicolor reference genome: improved assembly, gene
618 annotations, a transcriptome atlas, and signatures of genome organization. *Plant J. Cell Mol.*
619 *Biol.* **93**, 338–354 (2018).

620 56. Amborella Genome Project. The Amborella genome and the evolution of flowering plants.
621 *Science* **342**, 1241089 (2013).

622 57. Seader, V. H., Thornsberry, J. M. & Carey, R. E. Utility of the Amborella trichopoda expansin
623 superfamily in elucidating the history of angiosperm expansins. *J. Plant Res.* **129**, 199–207
624 (2016).

625 58. Soltis, P. S. *et al.* Floral variation and floral genetics in basal angiosperms. *Am. J. Bot.* **96**, 110–
626 128 (2009).

627 59. Prilusky, J. *et al.* FoldIndex©: a simple tool to predict whether a given protein sequence is
628 intrinsically unfolded. *Bioinformatics* **21**, 3435–3438 (2005).

629 60. Zhao, T. & Schranz, M. E. Network-based microsynteny analysis identifies major differences
630 and genomic outliers in mammalian and angiosperm genomes. *Proc. Natl. Acad. Sci. U. S. A.*
631 **116**, 2165–2174 (2019).

632 61. Hughes, S. L. *et al.* The importance of size and disorder in the cryoprotective effects of
633 dehydrins. *Plant Physiol.* **163**, 1376–1386 (2013).

634 62. Palmer, S. R., De Villa, R. & Graether, S. P. Sequence composition versus sequence order in the
635 cryoprotective function of an intrinsically disordered stress-response protein. *Protein Sci. Publ.*
636 *Protein Soc.* **28**, 1448–1459 (2019).

637 63. Hara, M. *et al.* The Arabidopsis KS-type dehydrin recovers lactate dehydrogenase activity
638 inhibited by copper with the contribution of His residues. *Plant Sci. Int. J. Exp. Plant Biol.* **245**,
639 135–142 (2016).

640 64. Hara, M., Kondo, M. & Kato, T. A KS-type dehydrin and its related domains reduce Cu-
641 promoted radical generation and the histidine residues contribute to the radical-reducing
642 activities. *J. Exp. Bot.* **64**, 1615–1624 (2013).

643 65. Yokoyama, T., Ohkubo, T., Kamiya, K. & Hara, M. Cryoprotective activity of Arabidopsis KS-
644 type dehydrin depends on the hydrophobic amino acids of two active segments. *Arch. Biochem.*
645 *Biophys.* **691**, 108510 (2020).

646

647 **Competing interests**

648 The authors declare no competing interests.

649

Acknowledgements

This work was supported by grant from the National Agency for the Promotion of Science and Technology of Argentina (ANPCyT, PICT 2015-3527) and National Scientific and Technical Research Council (CONICET, PIP 11220150100584). AMZ is a career research scientist from CONICET and AEM is a CONICET fellow.

Author contributions

AMZ planned and designed the research. AEM and AMZ performed the research and analysed the data. AMZ wrote the article with contribution of AEM. Both authors read and approved the manuscript.

Supplemental material

Table S1: List of all DHNs analysed in this study. It includes sequence name, taxonomic data, accession number, syntenic/homologous group (H-, F- or Y-DHN), segmental structure and physicochemical characteristics.

Fig. S1: MEME analysis of unbiased DHN database.

Fig. S2: Multiple sequence alignment of FSK2 dehydrins. Protein sequences of FSK2 from eudicot species were aligned with Clustal Omega and visualized with Jalview. Structural segments are indicated and a consensus sequence is shown below the alignment.

Fig. S3: Multiple sequences alignment of FSK3 dehydrins. Protein sequences of FSK3-DHNs from angiosperms were aligned with Clustal Omega and visualized with Jalview. Structural segments are indicated and a consensus sequence is shown below the alignment. Note that there is a lysine-rich region adjacent to the S-segment but it is not as conserved as the B-segment found in FSK2-DHNs (compare to Fig. S2).

Fig. S4: Multiple sequences alignment of YSKn dehydrins. Protein sequences of YSKn-DHNs from angiosperms were aligned with T-Coffee and visualized with Jalview. Structural segments are indicated and a consensus sequence is shown below the alignment.

Fig. S5: Multiple sequence alignment of dehydrins Y2SKn. Protein sequences of Y2SKn-DHNs from angiosperms were aligned with T-Coffee and visualized with Jalview. Structural segments are indicated and a consensus sequence is shown below the alignment. Y2SK2 and Y2SK3-DHNs are present in eudicots and the grass *Brachypodium distachyon*, while other Poaceae only have Y2SK2-DHNs.

Fig. S6: Multiple sequences alignment of Y3SKn dehydrins. Protein sequences of Y3SKn-DHNs from angiosperms were aligned with T-Coffee and visualized with Jalview. Structural segments are indicated and a consensus sequence is shown below the alignment.

Fig. S7: Multiple sequence alignment of HSK-dehydrins. Protein sequences of H-DHNs from vascular plants were aligned with Clustal Omega and visualized with Jalview. Structural segments

are indicated and a consensus sequence is shown below the alignment. DHNs come from angiosperms except for proteins from *Selaginella moellendorffii* (Smo) and *Ginkgo biloba* (Gbi).

Fig. S8: Multiple sequence alignment of atypical H-DHNs from Malpighiales. (A) Alignment of HKS-DHNs from *P. trichocarpa* and *S. purpurea* and a HS-DHN from *P. trichocarpa*. (B) Atypical H-DHNs with multiple K segments interspersed with Phi-segments. Segments are indicated by a colour code: H (purple), K (red), S2 (blue) and Phi (green). Sequences were aligned with Clustal Omega and visualized with Jalview.

Fig. S9: Evolutionary relationships of bryophyte dehydrins. (A) Maximum-likelihood phylogenetic tree constructed with PhyML 3.0. Branches with bootstrap values over 90 are indicated with a circle. Note that DHN sequences from *P. patens* and *C. purpureus* form five homologous groups, while *S. fallax* DHNs are not grouped with the other sequences. (B) DHN sequences and homologous groups of *P. patens* and *C. purpureus*.

Fig. S10: Multiple sequence alignment of bryophyte DHNs. Segments are indicated by a colour code: K (red), Y (green) and S (blue). Note that Group I has a Y8K structure; the Y-segments with an asterisk (*) have a sequence identical to the Y-segments of angiosperms (DEYGNP), while the others have a modified Y-segment (DNYGN/QP). Group II has a KS-structure, Group III and V have a K2-structure and Group IV a K-structure. Sequences were aligned with T-Coffee and visualized with Jalview.

Fig. S11: Scatter plots of physicochemical features of angiosperm DHN-structural groups: Glycine content, GRAVY index and Fold index. Homologous groups are colour-coded: H-DHNs (purple), F-DHNs (orange) and Y-DHNs (green).