1 Title

2 The gene expression profile of the song control nucleus HVC shows sex specificity,
3 hormone responsiveness, and species specificity among songbirds

4 Author affiliation

5 Meng-Ching Ko* (ORCID 0000-0002-2234-9380)

6 Carolina Frankl-Vilches*

7 Antje Bakker*

8 Manfred Gahr*

9 *Department of Behavioural Neurobiology, Max Planck Institute for Ornithology,

10 Eberhard-Gwinner Straße 6a, 82319 Seewiesen, Germany

11 Corresponding author

12 Meng-Ching Ko

13 E-mail: mcko@orn.mpg.de

14 Keywords

15 Sex differences; Songbirds; Gene expression; Brain; HVC; Testosterone

16

17

18

19 Please format this manuscript in British English.

20

Abstract

Singing occurs in songbirds of both sexes, but some species show typical degrees of sex-specific performance. We studied the transcriptional sex differences in the HVC, a brain nucleus critical for song pattern generation, of the forest weaver (*Ploceus bicolor*), the blue-capped cordon-bleu (*Uraeginthus cyanocephalus*), and the canary (*Serinus canaria*), which are species that show low, medium, and high levels of sex-specific singing, respectively. We observed persistent sex differences in gene expression levels regardless of the species-specific sexual singing phenotypes. We further studied the HVC transcriptomes of defined phenotypes of canary, known for its testosterone-sensitive seasonal singing. By studying both sexes of canaries during both breeding and nonbreeding seasons, nonbreeding canaries treated with testosterone, and spontaneously singing females, we found that the circulating androgen levels and sex were the predominant variables associated with the variations in the HVC transcriptomes. The comparison of natural singing with testosterone-induced singing in canaries of the same sex revealed considerable differences in the HVC transcriptomes. Strong transcriptional changes in the HVC were detected during the transition from nonsinging to singing in canaries of both sexes. Although the sex-specific genes of singing females shared little resemblance with those of males, our analysis showed potential functional convergences. Thus, male and female songbirds achieve comparable singing behaviours with sex-specific transcriptomes.

Introduction

Most of the genomes of male and female individuals of the same species are the same but stark sex differences in physiological, phenotypical, or behavioural traits between the sexes are common and widespread in the animal kingdom. The songbird clade (a suborder of the perching birds) consists of more than 4000 extant avian species, and these exhibit a great diversity of sex differences in singing behaviour. Among domesticated canaries (*Serinus canaria*), males are known for their singing behaviour and have been selected for their sophisticated songs for centuries, whereas female canaries seldom sing (Hartley et al., 1997; Herrick and Harris, 1957; Ko et al., 2020; Pesch and Güttinger, 1985; Shoemaker, 1939; Vallet et al., 1996). Such substantial sex differences in singing behaviour are commonly found in the majority of northern temperate songbird species, even though females of many tropical and southern temperate species sing regularly, and their songs play an important role in inter-sexual communication (Hall et al., 2015; Price, 2019; Price et al., 2009; Slater and Mann, 2004). For example, female blue-capped cordon-bleus (*Uraeginthus cyanocephalus*) use their song as advertising signals and address their songs to their mates (Immelmann, 1968; Ota et al., 2018), although the female songs appear shorter and less complex than those of males (Geberzahn and Gahr, 2011). In another tropical songbird species, forest weavers (*Ploceus bicolor*), males and females develop their songs during pair binding and eventually learn to sing identical duets, which they use to defend their territories (Wickler and Seibt, 1980).

Although the extent of sex differences in singing behaviour varies greatly in the songbird clade (Ball, 2016), the song quality and occurrence in males and females of many songbird species are dependent on testosterone (Dittrich et al., 2014; Fusani et al., 2003; Ko et al., 2020; Leitner et al., 2001b; Nottebohm et al., 1987; Voigt and Leitner, 2008). In canaries that breed seasonally, the optimal breeding conditions are tightly associated with an increase in the day length, which initiates gonadal growth and testosterone production as well as various types of breeding activity, including singing (Leitner et al., 2001b; Nottebohm et al., 1987; Voigt and Leitner, 2008). The length and syllable repetition rate of songs during the breeding season are greater than those of songs in nonbreeding seasons (Leitner et al., 2001a; Leitner et al., 2001b; Voigt and Leitner, 2008). Castrated male canaries sing shorter songs than sham-operated males, and the subcutaneous implantation of testosterone results in the recovery of singing performance (Heid et al., 1985). The local administration of testosterone into the preoptic brain region increases the male canary singing rate by increasing motivation (Alward et al., 2013). Similarly, the systemic administration of testosterone reliably and repeatedly induces

singing behaviour in otherwise nonsinging female canaries (Fusani et al., 2003; Herrick and Harris, 1957; Leonard, 1939; Madison et al., 2015; Nottebohm, 1980; Shoemaker, 1939; Vellema et al., 2019b). Although female canaries rarely exhibit spontaneous singing (Hartley et al., 1997; Herrick and Harris, 1957; Ko et al., 2020; Pesch and Güttinger, 1985; Shoemaker, 1939; Vallet et al., 1996), its occurrence appears to be associated with the plasma androgen levels (Ko et al., 2020).

A set of interconnected neural circuits called the song control system controls the production and learning of singing behaviour (Nottebohm et al., 1976; Wild, 2004). The premotor nucleus HVC (used as the proper name), which is a sensorimotor integration centre in the song control system, is involved in the frequency and temporal modulation of songs in male and female songbirds (Hahnloser et al., 2002; Hoffmann et al., 2019) and in the sexual preferences regarding conspecific song displays (Brenowitz, 1991; Del Negro et al., 1998). The HVC is the only brain nucleus within the song control system that expresses receptors for both androgens and estrogens (Frankl-Vilches and Gahr, 2018; Gahr, 2001). Intriguingly, the anatomical properties of the HVC, such as volume (Figure 1), neuron number, and dendrite complexity, are male-biased (greater, higher and more complex, respectively, in males than in females) in all songbird species that have been examined, irrespective of the existence of sex differences in singing behaviour (Brenowitz et al., 1985; Gahr et al., 2008; Gahr et al., 1998; Gurney and Konishi, 1980; Hall et al., 2010; Lobato et al., 2015; MacDougall-Shackleton and Ball, 1999; Nixdorf et al., 1989; Nottebohm and Arnold, 1976; Schwabl et al., 2015). Testosterone treatment increases the delineable volume of the HVC in both male and female canaries (Fusani et al., 2003; Madison et al., 2015; Nottebohm, 1980) and many other species (Bernard and Ball, 1997; Dittrich et al., 2014; Gulledge and Deviche, 1999; Smith et al., 1997; Van Meir et al., 2004). Nevertheless, the HVC volume of female canaries implanted with testosterone remains markedly smaller than that of males (Nottebohm, 1980) (Figure 1). Testosterone clearly regulates singing behaviour and the anatomy of the HVC in both male and female songbirds. However, an intrinsic limit to the alterations induced by testosterone appears to exist, and this limit prevents female songbirds from reaching the same levels of “maleness” in terms of song characteristics and HVC anatomy.

How are such sex-specific differences in singing and HVC anatomy achieved in different species? We hypothesize the existence of a fundamental difference in gene regulation and expression between male and female conspecific songbirds. Based on this hypothesis, sex differences in the HVC transcriptomes of male and female conspecifics should always be observable, even though both sexes share an almost identical genome.

The female-specific W chromosome harbours 30-50 genes, whereas one and two copies of the Z chromosome (harbouring approximately 1,080 genes) are present in female and male birds, respectively (Frankl-Vilches et al., 2015; Sayers et al., 2020; Smeds et al., 2015). We further hypothesize that such intrinsic differences cannot be overwritten by manipulating the testosterone level in females. To test our hypotheses, we compared the transcriptomes of HVCs microdissected from three songbird species—the forest weaver, the blue-capped cordon-bleu, and the canary. These species represent three categories of sex differences in singing behaviour (low, medium, and high). Furthermore, we compared male and female canaries with and without testosterone implantation to evaluate the effects of testosterone (Table 1). The results revealed sex differences in the HVC transcriptome between- and within-species comparisons. Our results indicate that although the extent of sex-biased gene expression is context-dependent, it is persistent regardless of hormonal manipulation, behavioural phenotypes, and distinct genetic backgrounds.

Materials and methods

Animals

Forest weaver (*Ploceus bicolor*) pairs were observed in their breeding territories in eastern South Africa; the songs of the pairs were recorded to ensure that both mates were singing. Subsequently, the animals were caught and sacrificed in accordance with permits issued by the local authorities (Chief Professional Officer for Research at the Natal Parks, Game and Fish Preservation Board, P. O. B. 662, Pietermaritzburg 3200). Blue-capped cordon-bleus (*Uraeginthus cyanocephalus*) and canaries (*Serinus canaria*) were bred at the animal facilities at the Max Planck Institute for Ornithology in Seewiesen, Germany. The procedures used for animal housing and welfare complied with the European directives for the protection of animals used for scientific purposes (2010/63/EU), and the protocols were approved by the Government of Upper Bavaria (AZ No. 55.2-1-54-2532-181-12). Adult canaries and blue-capped cordon-bleus (aged at least 1 year) were housed in pairs under long-day conditions (light:dark = 14:10 hours), and their breeding activities were monitored. The birds (male canaries and cordon-bleus) were sacrificed after singing activity was observed during the breeding season. Nonbreeding canaries were housed pairwise with a 9-hour light:15-hour dark schedule for at least 8 weeks before their singing activity was monitored (generally starting in late September). Nonbreeding male and female canaries were maintained alone in sound-attenuated boxes (70 × 50 × 50 cm), recorded continuously during the song monitoring phase (two weeks)

and sacrificed after confirmation of no singing activity. Additional groups of male and female nonbreeding and nonsinging canaries were implanted with testosterone for 2 weeks and sacrificed after observation of singing activity. Water and food were available *ad libitum*. The sex was confirmed by PCR using the P2 and P8 primers for CHD genes (Griffiths et al., 1998) and by visual inspection of the reproduction system after sacrifice. We previously found that six nonbreeding female canaries exhibited singing behaviour during long-term monitoring, and their songs have been described in detail (Ko et al., 2020). In this study, we included the transcriptomes of these birds. All the birds were sacrificed via an overdose of isoflurane followed by decapitation, the body weights were recorded, and the brains were dissected, weighed, snap-frozen on dry ice and stored at -80°C until further use. Table 1 summarizes the information of the experimental groups used in this study.

Testosterone implantation

A Silastic™ tube (Dow Corning; 1.47-mm inner diameter, 1.96-mm outer diameter, 0.23-mm thickness) was cut to a length of 7 mm and loaded with testosterone (86500, Fluka) as densely as possible. The two ends of the Silastic™ tube were sealed with silicone elastomer (3140, Dow Corning). After closure, the implants were cleaned with 100% ethanol to remove testosterone particles and then immersed in ethanol overnight in a hood to ensure no leakage at either end. Implants with apparent dampness were discarded. One day prior to the implantation, the implants were incubated in 0.1 M phosphate buffered saline (PBS) overnight to enable the immediate release of testosterone upon implantation (Rasika et al., 1994). We started implantation at approximately 8:30 am (immediately after the light was turned on in the morning), which resulted in a 20-minute interval between birds based on the scarification time. A small incision was made on the back of the bird over the pectoral musculature, and one testosterone implant was placed subcutaneously. The skin was closed by the application of tissue glue. After 2 weeks, the animals were sacrificed. The testosterone implants were checked, and this inspection revealed that the implants were all in place and were not empty at the end of the experiments.

Radioimmunoassay of plasma testosterone

Blood was sampled (< 150 µl) at the time sacrifice. All blood samples were taken between 8 and 11 am and were taken within 3 min after caught to avoid the effect of handling (Wingfield et al., 1982). Blood samples were centrifuged (5,000 rpm, 10 min) to separate

the plasma from blood cells. Testosterone metabolites were measured with a radioimmunoassay using a commercial antiserum against testosterone (T3-125, Endocrine Sciences, Tarzana, USA) as previously described (Goymann et al., 2002). Standard curves and sample concentrations were calculated with Immunofit 3.0 (Beckman Inc. Fullerton, CA) using a four-parameter logistic curve fit and corrected for individual recoveries.

The testosterone concentrations were assayed in duplicate in five separate assays. The mean extraction efficiency for plasma testosterone was $85.0 \pm 3.9\%$ (mean $\pm$ SD, N = 42). The lower detection limits of the testosterone assays were 0.34, 0.35, 0.36, 0.38 and 0.35 pg per tube, and all the samples were above the detection limit. The intra-assay coefficients of variation of a chicken plasma pool were 8.7%, 3.4%, 12.8%, 1.9%, and 4.4%. The interassay coefficient of variation as determined by the variation in the chicken plasma pool between all the assays was 5.1%. Because the testosterone antibody used shows significant cross-reactions with 5 α -dihydrotestosterone (44%), our measurement might include a fraction of 5 α -DHT.

Brain sectioning

The birds were killed by an overdose of isoflurane, and their brains were snap-frozen on dry ice. The brains were sectioned sagittally into four series of 40- μ m sections and two series of 20- μ m sections with a cryostat (Jung CM3000 Leica). The 40- μ m sections were mounted on glass slides for subsequent tissue dissection for total RNA extraction, whereas the thin sections were mounted on RNase-free Superfrost slides for Nissl staining and measurement of the HVC volume. All sections were stored at -80°C until further processing.

Measurement of the HVC volume

One series of 20- μ m sections mounted on RNase-free Superfrost slides was subjected to Nissl staining with 0.1% thionin (Sigma-Aldrich), dehydrated, immersed in xylene and cover-slipped with Roti-Histokitt II mounting medium (Roth). The HVC areas (typically 8-10 slices) were measured with a Leica DM6000 B microscope connected to a computer-based image-analysis system (IMAtec). All brains were coded such that the observers were blind towards any additional information about the sections they measured during the delineations. The volumes were derived from the summed area measurements multiplied by the section thickness and the intersection distance.

210 **Microarray procedures and annotation**

211 For total RNA extraction, the song control nucleus HVC and the visual area of the
 212 entopallium were dissected from the abovementioned 40- μ m sections under a
 213 stereomicroscope (typically 24-32 slices for the HVC and 16-20 slices for the entopallium)
 214 and transferred into an Eppendorf tube containing 340 μ l of RLT buffer mixture
 215 (containing DTT, Qiagen). This dissection procedure using rather thin sections reduces
 216 the contamination of HVC tissue with surrounding tissue. RNA was then extracted using
 217 the RNeasy® Micro Kit (Qiagen) with the optional DNase digest step. The RNA quality
 218 was assessed using the Agilent Model 2100 Bioanalyzer (Agilent Technologies), and the
 219 RNA concentrations were assessed using a Nanodrop 1000 spectrometer (Thermo Fisher
 220 Scientific). The RNA quality of all samples was good (RIN > 7). The purified total RNA
 221 samples (at least 100 ng per sample) were subsequently processed and hybridized using
 222 the Ambion WT Expression Kit and the Affymetrix WT Terminal Labelling and Controls
 223 Kit. The resulting cDNA was hybridized to the Custom Affymetrix Gene Chip® MPIO-
 224 ZF1s520811 Exon Array, which has been used successfully and validated in cross-species
 225 hybridization studies (Dittrich et al., 2014; Frankl-Vilches et al., 2015). The 5.7 million
 226 male zebra finch-specific probes spotted on this array correspond to approximately
 227 4,711,133 probe sets and hence to 25,816 transcripts published in public databases (NCBI
 228 and Ensembl) (Warren et al., 2010). We annotated more than 90% of the transcripts to
 229 12,729 human orthologous genes using several publicly available databases (Ensembl,
 230 GenBank, UniProt, and DAVID (Benson et al., 2005; Consortium, 2015; Flicek et al., 2014;
 231 Huang et al., 2008, 2009; Yates et al., 2016)) and commercial databases (El Dorado,
 232 Genomatix, Precigen Bioinformatics Germany GmbH (PBG), Munich, Germany).
 233 Hybridization was performed for 16 hours at 45°C and 60 rpm in a GeneChip
 234 Hybridization Oven 640 (Affymetrix). The arrays were washed, stained, and scanned
 235 using the Affymetrix GeneChip Fluidics Station 450 and the Affymetrix GeneChip scanner
 236 3000 7G. The CEL files were generated using Affymetrix® GeneChip® Command
 237 Console® Software (AGCC), and for quality control, the success of individual
 238 hybridizations was assessed using Affymetrix® Expression Console™ software.

239 Differential expression was calculated using ChipInspector software version 21 (El
 240 Dorado Database version E28R1306 (Genomatix)). ChipInspector is a single probe-based
 241 analysis tool for microarray data that can show increased sensitivity compared with that
 242 obtained with conventional probe set-based analyses, such as robust multiarray analysis.
 243 ChipInspector consists of four steps: single probe-transcript annotation (ensuring up-to-
 244 date annotation), total intensity normalization, SAM (significance analysis of microarrays,

adapted to single-probe handling) analysis (Tusher et al., 2001), and transcript identification based on significantly changed probes (Cohen et al., 2008). We set the delta threshold to 0 (to control the false positive rate) and used the groupwise exhaustive comparison tool in ChipInspector. The minimum coverage for each transcript was set to 10 significant probes, and the minimum expression difference was $|\log_2(\text{fold change})| \geq 0.5$. The significantly differentially expressed transcripts obtained were annotated to human orthologous genes as described above. For transcripts belonging to the same genes, the average expression was calculated if all transcripts were regulated in the same manner (e.g., all upregulated or all downregulated). If a gene contains both up- or downregulated transcripts, the transcripts that showed changes in expression in the minority direction were discarded (<40% of the total transcripts), and the average expression was instead calculated from the remaining transcripts (60% or more). Transcripts without human orthologous gene annotation were removed prior to subsequent analyses. We were cautious about possible cross-species bias in hybridization, and differential expression analyses were only performed between two conspecific groups. The microarray data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Edgar et al., 2002) and are accessible through the GEO Series accession number [GSE83674](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83674).

Hierarchical clustering

The normalized gene expression levels (across all transcriptome samples) were calculated using the “justRMA” function of the R package “affy” v1.66.0 (Gautier et al., 2004), and the expression levels were further collapsed to gene levels using the “collapseRows” functions of the R package “WGCNA” v1.69 (Langfelder and Horvath, 2008, 2012). The group-level expression levels were calculated by averaging the gene expression levels. Spearman's ρ was calculated between samples (or groups) using the “cor” function, and a Euclidean distance matrix was calculated ($\text{dist}(1-\text{cor})$) and used for hierarchical clustering analysis (method: complete) with the “hclust” function. This computation was performed with R v4.0.2 (R_Core_Team, 2020) using the “dendextend” v1.14.0 package for visualization (Galili, 2015).

Principal component analysis (PCA)

PCA (Ringner, 2008) was performed using the normalized and collapsed HVC gene expression data (described in the hierarchical clustering section) from 41 canaries (Table 1). The computation was performed using the “pca” function of the R package

“pcaMethods” (v1.60.0 (Stacklies et al., 2007), method = svd). The data were centred but not scaled because the expression data had already been normalized.

Fisher’s exact testing for chromosomal enrichment

Fisher’s exact test was used to test whether sex-biased genes were enriched on a chromosome by comparing the gene lists of interest to the zebra finch annotation (<https://www.ncbi.nlm.nih.gov/genome/?term=zebra+finch>). We used the “fisher.test” function in R (R_Core_Team, 2020) with the alternative set to “greater” for enrichment. The male- and female-biased genes identified from each comparison as well as the male-specific, female-specific, and sex-shared genes were tested separately. The P values were adjusted using the Bonferroni correction to account for multiple comparisons using “p.adjust” in R (R_Core_Team, 2020).

Gene Ontology (GO) term enrichment analysis

We used ClueGO v2.2.4, an application built in the Cytoscape environment v3.3.0 (Bindea et al., 2009; Shannon et al., 2003), to predict the putative biological functions of the genes of interest. This software performs GO term enrichment hierarchical analyses and fuses GO terms with similar functions. The enrichment was determined by the right-sided test and corrected using the Bonferroni step-down method considering multiple comparisons.

Results and discussion

The species identity distinguishes its HVC transcriptomes

We quantified the total mRNA from the HVC of male and female birds belonging to three songbird species (forest weavers, blue-capped cordon-bleus, and canaries) during the breeding season (Table 1). For intraspecies sex comparisons, we included male and female canaries during the nonbreeding seasons (both nonsinging), and subgroups were treated with testosterone to induce singing behaviour in both sexes (Supplementary Figure 1). In addition, we obtained a rare group of nonbreeding female canaries that sang spontaneously without exogenous testosterone manipulation. However, their plasma androgen levels were intrinsically higher than those of nonbreeding and nonsinging females (Ko et al., 2020). To visualize the similarities in the gene expression patterns among and within the three species, we calculated Spearman’s rank correlation coefficient σ for 12,360 gene expression levels and calculated the distance matrix based

on the coefficient σ . The resulting cladogram indicated that the HVC transcriptomes of the eleven groups were clustered primarily by phylogenetic relatedness (Figure 1B and Supplementary Figure 2). We obtained a similar result with the transcriptomes of another tissue, the entopallium (Supplementary Figure 3), which is an area of the avian visual system that is functionally equivalent to the primary visual cortex of mammals. The within-species comparison of the canary HVC showed that the individual transcriptomes clustered well by sex, with the exception of the nonbreeding canaries implanted with testosterone (Figure 1B and Supplementary Figure 2). The subclusters of forest weaver and cordon-bleu HVC transcriptomes were less sexually differential than those of canaries, i.e., male and female birds were intermittent within the subcluster of the two species (Supplementary Figure 2 and Supplementary Figure 3).

320

321 **Figure 1. Species, sex, and plasma androgen level are the major determinants of the**
322 **HVC gene expression patterns in songbirds.**

A, HVC volume of birds used in this study. Forest weavers: males 0.528 mm³ (mean), females 0.357 mm³. Mann-Whitney Test, U = 7, P value = 0.0931. Cordon-bleus: males 0.245 mm³, females 0.136

mm³. Mann-Whitney Test, U = 0, P value = 0.00217. Breeding canaries: singing males 0.439 mm³, nonsinging females 0.116 mm³. Mann-Whitney Test, U = 0, P value = 0.00217. Nonbreeding canaries: nonsinging males 0.228 mm³, nonsinging females 0.103 mm³, singing females 0.176 mm³. Mann-Whitney Test (nonsinging males vs. nonsinging females), U = 0, P value = 0.00492. Mann-Whitney Test (nonsinging males vs. singing females), U = 8, P value = 0.127. Testosterone-implanted canaries: males 0.478 mm³, females 0.253 mm³. Mann-Whitney Test, U = 0, P value = 0.0238. * P value < 0.05; ** P value < 0.01. Each colour-coded dot indicates the measurement from one bird. The boxes indicate the 25th/50th/75th percentiles (bottom/middle/top bar), and the extent of the whiskers indicates the most extreme values that are within 1.5 times the IQR (interquartile range) of the hinge. T: testosterone; music note: presence of singing behaviour. **B**, Hierarchical clustering showing that phylogenetic relatedness accounts for the most variation in the HVC transcriptomes of females and males of the forest weaver, cordon-bleu, and canary. Among the seven canary groups, the testosterone-treated animals were the least similar to the untreated canaries. **C**, A PCA of the HVC transcriptomes of the seven canary groups distinguished male birds (PC2 < 0) from female birds (PC2 > 0). Each point is colour-coded by group (see B) and represents an HVC sample.

Sex-biased gene expression in the HVC of songbirds

We defined the degree of sex differences in HVC gene expression levels as the number of sex-biased genes whose expression levels showed significant differences between males and females of the same species (male-biased genes showed higher expression levels in males than in females, $\log_2$ (fold change) ≥ 0.5 ; female-biased genes presented higher expression levels in females than in males, $\log_2$ (fold change) ≤ -0.5). For canary, we quantified the sex differences from five distinct male-to-female comparisons (Figure 2A, comparisons 3 to 7).

Interestingly, we found persistent sex differences in HVC gene expression levels from all the comparisons, regardless of the degree of sex differences in singing behaviour (Figure 2A). The majority (> 90%) of sex-biased genes were lowly biased ($0.5 \leq |\log_2$ (fold change)| < 1, Supplementary Figure 4). The comparison of forest weaver (the female birds can sing identical songs to the males; least differences in the HVC volumes between the sexes; Figure 1A) showed the least prominent sex differences in the HVC transcriptomes across all the comparisons performed in this study. Nevertheless, the extent of sex differences in gene expression levels was extensive (> 1,300 sex-biased genes, Figure 2A, comparison 1). The comparison showing the second-lowest degree of sex-differences was the comparison of nonbreeding male and nonbreeding female canaries (> 1,500 sex-biased genes, Figure 2A, comparison 7), both lacking singing behaviour, but HVC volume was male biased (Figure 1A). Testosterone implantation induced singing behaviour and increased the HVC volume of both male and female canaries, although a markedly smaller

344 HVC volume continued to be observed in these females (Figure 1A and Supplementary
345 Figure 1C). Such processes in females are referred to as “masculinization” of the female
346 brain and behaviour (Arnold and Gorski, 1984; Wade, 2001). However, the gene
347 expression levels of the testosterone-stimulated singing female canaries were very
348 different from those of the male canaries administered the same treatment (> 5,000 sex-
349 biased genes, Figure 2A, comparison 4). The comparisons showing the most striking sex
350 differences were the two that included spontaneously singing female canaries. As
351 revealed in these review comparisons, many genes showed markedly different regulation in
352 spontaneously singing female canaries, as reflected by expression levels, compared with
353 breeding singing male canaries (> 8,600 sex-biased genes, comparison 5) and
354 nonbreeding nonsinging male canaries (> 7,500 sex-biased genes, comparison 6, Figure
355 2A).

356

Figure 2. Female- and male-biased sex differences in HVC gene expression levels in songbirds.

A, The bar graph summarizes the number of sex-biased genes in the HVC transcriptome identified from each male-to-female comparisons. The phenotypes of the groups being compared in each male-to-female comparison are listed at the bottom of the graph. **B**, The bar graph summarizes the number of autosomal and Z chromosomal genes classified as male- and female-biased genes in the HVC transcriptome based on each male-to-female comparison. Abbreviations: B, breeding; NB, nonbreeding; T, testosterone implantation. The music note indicates the presence or absence of singing behaviour; the asterisk (*) indicates enrichment for Z chromosomal genes (adjusted $p < 0.05$, Fisher's exact test followed by the Bonferroni correction for multiple testing).

For each comparison, we decomposed the sex-biased genes into two classes, autosomal genes and Z sex chromosomal genes, to assess whether the sex-biased genes were mainly concentrated on the sex chromosomes. Annotation for W chromosomal genes was unfortunately not available in our study. We used Fisher's exact test to evaluate whether a particular chromosome was enriched in sex-biased genes. We observed enrichments of Z chromosomal genes among the male-biased genes identified in most of the comparisons (Figure 2B and Supplementary Table 1). This observation was not surprising because males are the homogametic sex (have two copies of Z), whereas females are the heterogametic sex (have one copy of Z and one copy of W) in birds, and the dosage compensation for Z-linked genes is less complete in birds than that in mammals (Itoh et al., 2007; Mank, 2013; Nätt et al., 2014; Uebbing et al., 2015; Wolf and Bryk, 2011). However, one canary comparison (breeding singing males and nonbreeding singing females, Figure 2B, comparison 5) was an exception (Supplementary Table 1). In this comparison, the spontaneously singing female canaries expressed a high number of Z chromosomal genes (260 genes, approximately 24% of all Z chromosomal genes) at higher levels than the breeding singing male canaries, which eliminated the male enrichment of Z chromosomal genes.

Sex-biased genes were not only Z chromosomal genes; in fact, sizeable numbers of the autosomal genes were found to be sex-biased genes, particularly female-biased genes (male-biased genes: >40%; female-biased genes: >96%, Figure 2B). Many studies have reported sex differences in the gene expression levels in multiple tissues and animal species (Frésard et al., 2013; Itoh et al., 2007; Nätt et al., 2014; Uebbing et al., 2015; Wolf and Bryk, 2011; Yang et al., 2006). Gene expression levels are generally under tight regulation, and perturbed expression levels might result in functional consequences. For example, several genes, including transcription factors, modulate distinct gene sets

384 depending on their expression levels (Birchler et al., 2001; Doghman et al., 2013; Schulz,
385 2017).

386 We compared the male-biased genes in a pairwise manner across the
387 comparisons and found that the male-biased autosomal genes showed low similarity to
388 each other (Supplementary Figure 5A), whereas the male-biased Z chromosomal genes
389 showed some similarity across the groups (Supplementary Figure 5B). Pairwise
390 comparison of the female-biased genes across groups showed low similarity, and this
391 finding was obtained for both autosomal genes (Supplementary Figure 5C) and Z
392 chromosomal genes (Supplementary Figure 5D). Fisher's exact test showed that
393 chromosome 2 (comparisons 1, 4, and 5) and chromosome 3 (comparisons 1, 2, and 4)
394 are hotspots for female-biased genes; the enrichments of these chromosomes were
395 observed in four out of the seven pairwise comparisons (Supplementary Table 1).

396

397 **Figure 3. Cross-species comparisons of sex-biased genes expressed in the HVC of**
398 **birds with singing phenotypes shows high species specificity.**

Venn diagrams of male-biased (A) and female-biased (B) autosomal and Z chromosomal genes. The numbers of sex-biased genes are indicated in parentheses. C, GO term enrichment analysis predicting the biological functions of sex-biased (autosomal and Z chromosomal) genes. The GO terms were categorized and are colour coded. Bonferroni-adjusted P values are shown by colour scales in the heatmap. Only five GO terms with the lowest Bonferroni-adjusted P values are shown for each set (see Supplementary Table 2 for the complete results). The phenotypes of the groups used for comparisons 1,

2, and 5 are listed at the bottom. Abbreviations: B, breeding; NB, nonbreeding. The music note indicates the presence or absence of singing behaviour

Further examination of the sex-biased genes (comparisons 1, 2, and 5) identified in the groups of naturally singing birds revealed that almost none of the male-biased autosomal genes (Figure 3A) and none of the female-biased Z chromosomal genes (Figure 3B) were shared across the three songbird species. In addition, relatively low numbers of male-biased Z chromosomal genes (Figure 3A) and female-biased autosomal genes (Figure 3B) were shared across the three songbird species. This observation indicates that only a small set of genes were regulated in the same manner (male- or female-biased) between species even though all birds exhibit singing behaviours. Moreover, a Gene Ontology (GO) enrichment analysis suggested that male- and female-biased genes did not functionally converge to similar pathways in the three studied species (Figure 3C).

The plasma androgen levels substantially alter the HVC transcriptomes of canaries

With the aim of understanding the features that distinguish the HVC transcriptomes in within-species contexts, we performed a principal component analysis (PCA) of the HVC transcriptomic data from the seven canary groups to identify variables that would explain the most variation in the data. By calculating the correlation coefficient between the PCs and the variables (plasma testosterone levels (Supplementary Figure 1), HVC volume (Figure 1A), sex, and singing), we identified the variables that were highly correlated with the most important PCs. PC1 explained 32% of the data variance (Figure 1C and Supplementary Figure 6) and was strongly correlated with the blood plasma androgen concentrations and the presence of singing activity (plasma androgen level: Pearson's $r = -0.57$, Bonferroni-adjusted $p = 1.0 \times 10^{-4}$; singing: Pearson's $r = -0.51$, Bonferroni-adjusted $p = 0.001$, Supplementary Table 3). Sex and HVC volume were strongly correlated with PC2 (sex: Pearson's $r = -0.91$, Bonferroni-adjusted $p = 4.4 \times 10^{-16}$; HVC volume: Pearson's $r = -0.73$, Bonferroni-adjusted $p = 2.1 \times 10^{-7}$), which explained 19% of the variance in the data (Supplementary Table 3). Taken together, the hierarchical clustering and PCA results suggest that although phylogenetic relationships dominate the variation between songbird species, the circulating testosterone levels, the presence of singing activity, and sex identity dominate the HVC gene expression patterns within a single species.

The application of testosterone does not yield an HVC transcriptome that mimics that of natural singing canaries

To investigate the molecular mechanisms underlying the singing phenotype, we performed a differential gene expression analysis of the HVC transcriptomes of the two groups of singing canary females (nonbreeding spontaneously singing females and nonbreeding testosterone-stimulated singing females) against the nonsinging nonbreeding female canaries. Similarly, a differential gene expression analysis of the HVC transcriptomes of the two groups of singing male canaries (breeding singing males and nonbreeding testosterone-stimulated singing males) with nonsinging nonbreeding male canaries was performed. A substantial number of genes were differentially expressed between singing and nonsinging birds (Figure 4A, nonbreeding singing females: 4,125 genes; nonbreeding testosterone-stimulated singing females: 7,702 genes; breeding singing males: 3,359 genes; nonbreeding testosterone-stimulated singing males: 5,106 genes). Approximately 65% of the differentially expressed genes found in the naturally singing birds overlapped with those found in the testosterone-treated birds of the same sex (male: 65%; female: 62%). However, both testosterone-treated groups had markedly higher numbers of differentially expressed genes than the naturally singing groups of the same sex (Figure 4A). Thus, although testosterone implantation induced singing in females and males, most testosterone-responsive genes (female: 67%; male: 57%) might not be necessary for singing behaviour *per se* but rather a response to nonphysiological levels of testosterone (Supplementary Figure 1). Alternatively, mechanisms that lead to the first song in life might be very different from those that reinduce singing in animals that sang before (Vellema et al., 2019a). Because male canaries sing regularly starting from approximately 50 days of age (Nottebohm et al., 1986) while most female canaries never sing (Ko et al., 2020), the induction of singing in females by testosterone likely activated genes related to first-time singing, whereas in males, this treatment activated genes related to reinduced singing.

The majority of differentially expressed genes between singing and nonsinging canaries are sex-specific

We examined whether the overlapping gene sets of the female canary signing groups were similar to the overlapping gene sets of the singing males (Figure 4C). The Venn diagram indicated that only approximately 25% (642 of 2569 genes, Supplementary Table 4) of the female-specific expressed genes and 29% (642 of 2210 genes,

Supplementary Table 5) of the male-specific expressed genes were shared by both sexes of singing canaries (Supplementary Table 6).

462

463 **Figure 4. HVC transcriptomes of naturally singing canaries are dissimilar from**
464 **testosterone-stimulated singing canaries.**

Venn diagrams comparing (A) the two female singing groups, (B) the two male singing groups, and (C) the female- and male-specific expressed genes resulting from comparisons A and B. Note that only approximately 20% of the genes were shared among the singing females and approximately 26% were shared among the singing males. The majority of these overlapping genes were not sex-shared;

approximately 75% of female intersections were specific to female groups, whereas 71% of male intersections were specific to male groups. **D**, GO term enrichment analysis of female-specific, male-specific, and sex-shared expressed genes derived from comparisons A to C. Because many putative functions show similarities between male- and female-specific expressed genes, the results suggest functional intraspecies convergence based on sex-specific gene expression in the HVC. See Supplementary Table 7 for the full results of the GO term enrichment analysis.

The sex-specific genes show functional overlap

To understand the putative biological functions of the female-specific, male-specific, and sex-shared expressed genes in the HVC, we performed a GO term enrichment analysis. Interestingly, the results suggested that the female- and male-specific expressed genes largely overlapped at the functional level (Figure 4D and Supplementary Table 7). GO terms such as nervous system development (GO:0007399), neuron development (GO:0048666), and intracellular signal transduction (GO:0035556) were shared among the sexes. Female-specific GO terms were mainly related to cellular maintenance, such as organonitrogen compound metabolic process (GO:1901564) and phospholipid metabolic process (GO:0006644). In contrast, the male-specific GO terms were axon development (GO:0061564), DNA replication (GO:0006260), cell migration (GO:0016477), and blood vessel development (GO:0001568). In summary, although only approximately one-fourth to one-third of the sex-specific expressed genes are shared among the sexes in terms of their identities, most of the predicted functions of the sex-specific expressed genes are nevertheless sex-shared in singing canaries.

All four groups of singing canaries had elevated plasma androgen levels compared with nonbreeding nonsinging canaries of the same sex (Supplementary Figure 1 and (Ko et al., 2020)). The activation of singing is likely testosterone-dependent in females as in males (Hartley and Suthers, 1989; Heid et al., 1985; Leitner et al., 2001a; Nottebohm et al., 1987). Thus, the potential master regulator for inducing singing behaviour might be testosterone-sensitive and Z-linked. One such candidate is *DMRT1* (doublesex and mab-3 related transcription factor 1), which is present in the sex-shared expressed gene list. The Z-linked gene *DMRT1* is needed for male sex determination in birds and other animal species (Herpin and Scharf, 2015; Lambeth et al., 2014; Smith et al., 2009). The overexpression of *DMRT1* in female chicken embryos reduces aromatase expression in the gonads and triggers development of the testis (Lambeth et al., 2014). The role of *DMRT1* in adult avian tissues in general and in the brain in particular is unknown. Thus, whether *DMRT1* affects steroid metabolism in the HVC, such as converting testosterone to more active metabolites (see below), or whether it regulates other mechanisms that

direct the HVC into a configuration that enables singing needs to be validated by future experiments.

Testosterone can be converted to 5 α -dihydrotestosterone (5 α -DHT) and 17 β -estradiol, which activate androgen receptor (AR) and estrogen receptors (ER α and ER β), respectively. Both AR (encoded by *AR*) and ERs (ER α encoded by *ESR1* and ER β encoded by *ESR2*) are transcription factors that play important roles in the transcription of numerous genes (Bourdeau et al., 2004; Pihlajamaa et al., 2015; Takayama et al., 2007; Wilson et al., 2016). Interestingly, *ESR1* was specifically expressed in males, whereas *ESR2* was female-specifically expressed in the HVC of canaries, which suggests that estrogen receptor paralogues could provide finer-tuned mechanisms for sex-specific regulation. Empirical results have shown that ER α and ER β bind to the same estrogen response element motifs (Zhao et al., 2010) and might functionally overlap in some tissues. Moreover, ER paralogues show sex differences in expression levels and tissue specificity (Zhang et al., 2017). In quail, the administration of an agonist specific to ER β on embryonic day 7 demasculinizes male sexual behaviour and midbrain nuclei characteristics in Japanese quails, whereas an agonist specific to ER α does not exert this effect (Court et al., 2020). The specific roles of each ER paralogue in the adult HVC of male and female songbirds warrant further investigation. Until now, ER α but not ER β was expected to regulate the function of HVC neurons of adult songbirds in addition to the AR (Frankl-Vilches and Gahr, 2017).

Conclusion

In this study, we investigated the sex differences in the gene expression patterns in the HVC of three songbird species with different levels of sex-specific singing and several different song-related phenotypes between male and female canaries. Our inter- and intraspecies comparisons yielded large-scale transcriptional sex differences regardless of singing behaviour. Instead, fundamental sex differences in gene expression levels were found in the HVC, and these differences were highly species-specific. By leveraging several experimental groups of canaries, we found that the plasma androgen levels and sex were the major contributors to the variations in the HVC transcriptome. Although testosterone reliably induced singing in both female and male canaries, testosterone treatment did not alter the transcriptome to imitate that of natural singing birds. Our results suggest that female and male canaries rely on different gene networks for singing behaviour, but the sex-specific gene networks might show functional convergence.

Data accessibility

The microarray CEL files (GEO Series accession number [GSE83674](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE83674)) are available on NCBI's Gene Expression Omnibus. The processed microarray data, plasma androgen levels, HVC volume measures, and all scripts used for analysis and visualization are available in GitHub (<https://github.com/maggieMCKO/SongbirdSexDiff>).

Authors' contributions

M-CK collected the nonbreeding testosterone-implanted canaries, analysed and visualized the transcriptomic data and was a major contributor to the writing of the manuscript. AB and M-CK performed the RNA extractions and microarray hybridizations. MG and M-CK isolated the tissues for microarray. MG collected the forest weavers and cordon-bleus. MG conceived the study and obtained financial support for the study. M-CK, CF-V and MG drafted the manuscript. All the authors read and approved the final manuscript.

Acknowledgements

We thank Dieter Schmidl for catching the forest weavers, Stefan Leitner for providing the breeding canaries, Roswitha Brighton and David Witkowski for maintaining the breeding colonies of the blue-capped cordon-bleus and the canaries, Doris Walcher for providing data from the breeding male and female canaries as well as the nonbreeding nonsinging male and female canaries, and Wolfgang Goymann and Monika Trappschuh for the RIA of the plasma testosterone. Furthermore, we express our appreciation to Maude Baldwin, Glenn Cockburn, Falk Dittrich, Lisa Gill, Vincent Van Meir, Keren Sadanandan, and Michiel Vellema for their comments on previous versions of this manuscript and to the International Max Planck Research School for Organismal Biology for the training and support provided.

Competing interests

The authors declare no competing financial interests.

Funding

This research was supported by Max-Planck-Gesellschaft to MG. The funder had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Alward, B.A., Balthazart, J., and Ball, G.F. (2013). Differential effects of global versus local testosterone on singing behavior and its underlying neural substrate. *Proc Natl Acad Sci U S A* 110, 19573-19578.
- Arnold, A.P., and Gorski, R.A. (1984). Gonadal steroid induction of structural sex differences in the central nervous system. *Annu Rev Neurosci* 7, 413-442.
- Ball, G.F. (2016). Species variation in the degree of sex differences in brain and behaviour related to birdsong: adaptations and constraints. *Philos Trans R Soc Lond B Biol Sci* 371.
- Benson, D.A., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., and Wheeler, D.L. (2005). GenBank. *Nucleic Acids Research* 33, D34-D38.
- Bernard, D.J., and Ball, G.F. (1997). Photoperiodic Condition Modulates the Effects of Testosterone on Song Control Nuclei Volumes in Male European Starlings. *General and Comparative Endocrinology* 105, 276-283.
- Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.-H., Pagès, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* 25, 1091-1093.
- Birchler, J.A., Bhadra, U., Bhadra, M.P., and Auger, D.L. (2001). Dosage-Dependent Gene Regulation in Multicellular Eukaryotes: Implications for Dosage Compensation, Aneuploid Syndromes, and Quantitative Traits. *Developmental Biology* 234, 275-288.
- Bourdeau, V.r., Deschênes, J., Métivier, R.I., Nagai, Y., Nguyen, D., Bretschneider, N., Gannon, F., White, J.H., and Mader, S. (2004). Genome-Wide Identification of High-Affinity Estrogen Response Elements in Human and Mouse. *Molecular Endocrinology* 18, 1411-1427.
- Brenowitz, E. (1991). Altered perception of species-specific song by female birds after lesions of a forebrain nucleus. *Science* 251, 303 - 305.
- Brenowitz, E.A., Arnold, A.P., and Levin, R.N. (1985). Neural correlates of female song in tropical duetting birds. *Brain Res* 343, 104-112.
- Cohen, C.D., Lindenmeyer, M.T., Eichinger, F., Hahn, A., Seifert, M., Moll, A.G., Schmid, H., Kiss, E., Grone, E., Grone, H.J., *et al.* (2008). Improved elucidation of biological processes linked to diabetic nephropathy by single probe-based microarray data analysis. *PLoS One* 3, e2937.
- Consortium, T.U. (2015). UniProt: a hub for protein information. *Nucleic Acids Research* 43, D204-D212.
- Court, L., Vandries, L., Balthazart, J., and Cornil, C.A. (2020). Key role of estrogen receptor beta in the organization of brain and behavior of the Japanese quail. *Horm Behav* 125, 104827.
- Del Negro, C., Gahr, M., Leboucher, G., and Kreutzer, M. (1998). The selectivity of sexual responses to song displays: effects of partial chemical lesion of the HVC in female canaries. *Behavioural Brain Research* 96, 151-159.
- Dittrich, F., Ramenda, C., Grillitsch, D., Frankl-Vilches, C., Ko, M.C., Hertel, M., Goymann, W., Ter Maat, A., and Gahr, M. (2014). Regulatory mechanisms of testosterone-stimulated song in the sensorimotor nucleus HVC of female songbirds. *BMC Neurosci* 15, 128.

602 Doghman, M., Figueiredo, B.C., Volante, M., Papotti, M., and Lalli, E. (2013). Integrative
603 analysis of SF-1 transcription factor dosage impact on genome-wide binding and gene
604 expression regulation. *Nucleic Acids Research* 41, 8896-8907.

605 Edgar, R., Domrachev, M., and Lash, A. (2002). Gene Expression Omnibus: NCBI gene
606 expression and hybridization array data repository. *Nucleic Acids Res* 30, 207 - 210.

607 Flicek, P., Amode, M.R., Barrell, D., Beal, K., Billis, K., Brent, S., Carvalho-Silva, D.,
608 Clapham, P., Coates, G., Fitzgerald, S., *et al.* (2014). Ensembl 2014. *Nucleic Acids*
609 *Research* 42, D749-D755.

610 Frankl-Vilches, C., and Gahr, M. (2017). Androgen and estrogen sensitivity of bird song: a
611 comparative view on gene regulatory levels. *Journal of Comparative Physiology A*.

612 Frankl-Vilches, C., and Gahr, M. (2018). Androgen and estrogen sensitivity of bird song: a
613 comparative view on gene regulatory levels. *Journal of Comparative Physiology A* 204,
614 113-126.

615 Frankl-Vilches, C., Kuhl, H., Werber, M., Klages, S., Kerick, M., Bakker, A., de Oliveira, E.H.,
616 Reusch, C., Capuano, F., Vowinckel, J., *et al.* (2015). Using the canary genome to decipher
617 the evolution of hormone-sensitive gene regulation in seasonal singing birds. *Genome*
618 *Biol* 16, 19.

619 Frésard, L., Morisson, M., Brun, J.-M., Collin, A., Pain, B., Minvielle, F., and Pitel, F. (2013).
620 Epigenetics and phenotypic variability: some interesting insights from birds. *Genetics*
621 *Selection Evolution* 45, 16.

622 Fusani, L., Metzdorf, R., Hutchison, J., and Gahr, M. (2003). Aromatase inhibition affects
623 testosterone-induced masculinization of song and the neural song system in female
624 canaries. *J Neurobiol* 54, 370 - 379.

625 Gahr, M. (2001). Distribution of sex steroid hormone receptors in the avian brain:
626 Functional implications for neural sex differences and sexual behaviors. *Microscopy*
627 *Research and Technique* 55, 1-11.

628 Gahr, M., Metzdorf, R., Schmidl, D., and Wickler, W. (2008). Bi-directional sexual
629 dimorphisms of the song control nucleus HVC in a songbird with unison song. *PLoS One*
630 3, e3073.

631 Gahr, M., Sonnenschein, E., and Wickler, W. (1998). Sex difference in the size of the
632 neural song control regions in a duetting songbird with similar song repertoire size of
633 males and females. *Journal of Neuroscience* 18, 1124-1131.

634 Galili, T. (2015). dendextend: an R package for visualizing, adjusting and comparing
635 trees of hierarchical clustering. *Bioinformatics* 31, 3718-3720.

636 Gautier, L., Cope, L., Bolstad, B.M., and Irizarry, R.A. (2004). affy--analysis of Affymetrix
637 GeneChip data at the probe level. *Bioinformatics* 20, 307-315.

638 Geberzahn, N., and Gahr, M. (2011). Undirected (solitary) birdsong in female and male
639 blue-capped cordon-bleus (*Uraeginthus cyanocephalus*) and its endocrine correlates.
640 *PLoS One* 6, e26485.

641 Genomatix (<http://www.genomatix.de>: Precigen Bioinformatics Germany GmbH
642 (PBG)).

643 Goymann, W., Möstl, E., and Gwinner, E. (2002). Non-invasive methods to measure
644 androgen metabolites in excrements of European stonechats, *Saxicola torquata rubicola*.
645 *General and Comparative Endocrinology* 129, 80-87.

646 Griffiths, R., Double, M.C., Orr, K., and Dawson, R.J.G. (1998). A DNA test to sex most
647 birds. *Molecular Ecology* 7, 1071-1075.

648 Gullledge, C.C., and Deviche, P. (1999). Photoperiod and testosterone independently
649 affect vocal control region volumes in adolescent male songbirds. *Journal of*
650 *Neurobiology* 36, 550-558.

651 Gurney, M.E., and Konishi, M. (1980). Hormone-induced sexual differentiation of brain
652 and behavior in zebra finches. *Science* 208, 1380-1383.

653 Hahnloser, R.H.R., Kozhevnikov, A.A., and Fee, M.S. (2002). An ultra-sparse code
654 underlies the generation of neural sequences in a songbird. *Nature* 419, 65-70.

655 Hall, M.L., Rittenbach, M.R.D., and Vehrencamp, S.L. (2015). Female song and vocal
656 interactions with males in a neotropical wren. *Frontiers in Ecology and Evolution* 3.

657 Hall, Z.J., MacDougall-Shackleton, S.A., Osorio-Beristain, M., and Murphy, T.G. (2010).
658 Male bias in the song control system despite female bias in song rate in streak-backed
659 orioles (*Icterus pustulatus*). *Brain Behav Evol* 76, 168-175.

660 Hartley, R.S., Chinn, M.S., and Ullrich, N.F.E. (1997). Left Syringeal Dominance in
661 Testosterone-Treated Female Canaries. *Neurobiology of Learning and Memory* 67, 248-
662 253.

663 Hartley, R.S., and Suthers, R.A. (1989). Airflow and pressure during canary song: direct
664 evidence for mini-breaths. *Journal of Comparative Physiology A* 165, 15-26.

665 Heid, P., Güttinger, H.R., and Pröve, E. (1985). The Influence of Castration and
666 Testosterone Replacement on the Song Architecture of Canaries (*Serinus canaria*).
667 *Zeitschrift für Tierpsychologie* 69, 224-236.

668 Herpin, A., and Scharf, M. (2015). Plasticity of gene-regulatory networks controlling sex
669 determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO*
670 *Rep* 16, 1260-1274.

671 Herrick, E.H., and Harris, J.O. (1957). Singing Female Canaries. *Science* 125, 1299-1300.

672 Hoffmann, S., Trost, L., Voigt, C., Leitner, S., Lemazina, A., Sagunsky, H., Abels, M.,
673 Kollmansperger, S., Maat, A.T., and Gahr, M. (2019). Duets recorded in the wild reveal
674 that interindividually coordinated motor control enables cooperative behavior. *Nat*
675 *Commun* 10, 2577.

676 Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2008). Systematic and integrative
677 analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 4,
678 44-57.

679 Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009). Bioinformatics enrichment tools:
680 paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids*
681 *Research* 37, 1-13.

682 Immelmann, K. (1968). *Zur biologischen Bedeutung des Estrildidengesanges*, Vol 109.

683 Itoh, Y., Melamed, E., Yang, X., Kampf, K., Wang, S., Yehya, N., Van Nas, A., Replogle, K.,
684 Band, M.R., Clayton, D.F., *et al.* (2007). Dosage compensation is less effective in birds
685 than in mammals. *Journal of Biology* 6, 1-15.

686 Ko, M.-C., Van Meir, V., Vellema, M., and Gahr, M. (2020). Characteristics of song, brain-
687 anatomy and blood androgen levels in spontaneously singing female canaries.
688 *Hormones and Behavior* 117, 104614.

689 Lambeth, L.S., Raymond, C.S., Roeszler, K.N., Kuroiwa, A., Nakata, T., Zarkower, D., and
690 Smith, C.A. (2014). Over-expression of DMRT1 induces the male pathway in embryonic
691 chicken gonads. *Developmental Biology* 389, 160-172.

692 Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation
693 network analysis. *BMC Bioinformatics* 9, 1-13.

Langfelder, P., and Horvath, S. (2012). Fast R Functions for Robust Correlations and Hierarchical Clustering. *Journal of statistical software* 46, i11.

Leitner, S., Voigt, C., and Gahr, M. (2001a). Seasonal Changes in the Song Pattern of the Non-Domesticated Island Canary (*Serinus Canaria*) a Field Study. *Behaviour* 138, 885-904.

Leitner, S., Voigt, C., Garcia-Segura, L., Van't Hof, T., and Gahr, M. (2001b). Seasonal activation and inactivation of song motor memories in wild canaries is not reflected in neuroanatomical changes of forebrain song areas. *Horm Behav* 40, 160 - 168.

Leonard, S.L. (1939). Induction of Singing in Female Canaries by Injections of Male Hormone. *Experimental Biology and Medicine* 41, 229-230.

Lobato, M., Vellema, M., Gahr, C., Leitão, A., de Lima, S.M.A., Geberzahn, N., and Gahr, M. (2015). Mismatch in sexual dimorphism of developing song and song control system in blue-capped cordon-bleus, a songbird species with singing females and males. *Frontiers in Ecology and Evolution* 3, 117.

MacDougall-Shackleton, S.A., and Ball, G.F. (1999). Comparative studies of sex differences in the song-control system of songbirds. *Trends in Neurosciences* 22, 432-436.

Madison, F.N., Rouse, M.L., Jr., Balthazart, J., and Ball, G.F. (2015). Reversing song behavior phenotype: Testosterone driven induction of singing and measures of song quality in adult male and female canaries (*Serinus canaria*). *Gen Comp Endocrinol* 215, 61-75.

Mank, J.E. (2013). Sex chromosome dosage compensation: definitely not for everyone. *Trends in Genetics* 29, 677-683.

Nätt, D., Agnvall, B., and Jensen, P. (2014). Large Sex Differences in Chicken Behavior and Brain Gene Expression Coincide with Few Differences in Promoter DNA-Methylation. *PLoS ONE* 9, e96376.

Nixdorf, B.E., Davis, S.S., and DeVoogd, T.J. (1989). Morphology of Golgi-impregnated neurons in hyperstriatum ventralis, pars caudalis in adult male and female canaries. *J Comp Neurol* 284, 337-349.

Nottebohm, F. (1980). Testosterone triggers growth of brain vocal control nuclei in adult female canaries. *Brain Res* 189, 429-436.

Nottebohm, F., and Arnold, A.P. (1976). Sexual dimorphism in vocal control areas of the songbird brain. *Science* 194, 211-213.

Nottebohm, F., Nottebohm, M., Crane, L., and Wingfield, J. (1987). Seasonal changes in gonadal hormone levels of adult male canaries and their relation to song. *Behav Neural Biol* 47, 197 - 211.

Nottebohm, F., Nottebohm, M.E., and Crane, L. (1986). Developmental and seasonal changes in canary song and their relation to changes in the anatomy of song-control nuclei. *Behavioral and neural biology* 46, 445-471.

Nottebohm, F., Stokes, T.M., and Leonard, C.M. (1976). Central control of song in the canary, *Serinus canarius*. *The Journal of Comparative Neurology* 165, 457-486.

Ota, N., Gahr, M., and Soma, M. (2018). Couples showing off: Audience promotes both male and female multimodal courtship display in a songbird. *Sci Adv* 4, eaat4779.

Pesch, A., and Güttinger, H.-R. (1985). Der Gesang des weiblichen Kanarienvogels. *Journal für Ornithologie* 126, 108-110.

739 Pihlajamaa, P., Sahu, B., and Jänne, O.A. (2015). Determinants of Receptor- and Tissue-
740 Specific Actions in Androgen Signaling. *Endocrine Reviews* 36, 357-384.

741 Price, J.J. (2019). Sex differences in song and plumage color do not evolve through sexual
742 selection alone: new insights from recent research. *Journal of Ornithology* 160, 1213-
743 1219.

744 Price, J.J., Lanyon, S.M., and Omland, K.E. (2009). Losses of female song with changes
745 from tropical to temperate breeding in the New World blackbirds. *Proceedings of the*
746 *Royal Society B: Biological Sciences* 276, 1971-1980.

747 R_Core_Team (2020). R: A language and environment for statistical computing
748 [Internet]. Vienna, Austria: R Foundation for Statistical Computing; (R Foundation for
749 Statistical Computing).

750 Rasika, S., Nottebohm, F., and Alvarez-Buylla, A. (1994). Testosterone increases the
751 recruitment and/or survival of new high vocal center neurons in adult female canaries.
752 *Proceedings of the National Academy of Sciences* 91, 7854-7858.

753 Ringner, M. (2008). What is principal component analysis? *Nat Biotechnol* 26, 303-304.

754 Sayers, E.W., Beck, J., Brister, J.R., Bolton, E.E., Canese, K., Comeau, D.C., Funk, K., Ketter,
755 A., Kim, S., Kimchi, A., *et al.* (2020). Database resources of the National Center for
756 Biotechnology Information. *Nucleic Acids Res* 48, D9-D16.

757 Schulz, E.G. (2017). X-chromosome dosage as a modulator of pluripotency, signalling
758 and differentiation? *Philosophical Transactions of the Royal Society B: Biological*
759 *Sciences* 372.

760 Schwabl, H., Dowling, J., Baldassarre, D.T., Gahr, M., Lindsay, W.R., and Webster, M.S.
761 (2015). Variation in song system anatomy and androgen levels does not correspond to
762 song characteristics in a tropical songbird. *Animal Behaviour* 104, 39-50.

763 Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
764 Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for
765 integrated models of biomolecular interaction networks. *Genome Res* 13, 2498-2504.

766 Shoemaker, H.H. (1939). Effect of Testosterone Propionate on Behavior of the Female
767 Canary. *Proceedings of the Society for Experimental Biology and Medicine* 41, 299-302.

768 Slater, P.J.B., and Mann, N.I. (2004). Why do the females of many bird species sing in the
769 tropics? *Journal of Avian Biology* 35, 289-294.

770 Smeds, L., Warmuth, V., Bolivar, P., Uebbing, S., Burri, R., Suh, A., Nater, A., Bures, S.,
771 Garamszegi, L.Z., Hogner, S., *et al.* (2015). Evolutionary analysis of the female-specific
772 avian W chromosome. *Nat Commun* 6, 7330.

773 Smith, C.A., Roeszler, K.N., Ohnesorg, T., Cummins, D.M., Farlie, P.G., Doran, T.J., and
774 Sinclair, A.H. (2009). The avian Z-linked gene DMRT1 is required for male sex
775 determination in the chicken. *Nature* 461, 267-271.

776 Smith, G.T., Brenowitz, E.A., and Wingfield, J.C. (1997). Roles of photoperiod and
777 testosterone in seasonal plasticity of the avian song control system. *J Neurobiol* 32, 426-
778 442.

779 Stacklies, W., Redestig, H., Scholz, M., Walther, D., and Selbig, J. (2007). pcaMethods—a
780 bioconductor package providing PCA methods for incomplete data. *Bioinformatics* 23,
781 1164-1167.

782 Takayama, K., Kaneshiro, K., Tsutsumi, S., Horie-Inoue, K., Ikeda, K., Urano, T., Ijichi, N.,
783 Ouchi, Y., Shirahige, K., Aburatani, H., *et al.* (2007). Identification of novel androgen

784 response genes in prostate cancer cells by coupling chromatin immunoprecipitation and
785 genomic microarray analysis. *Oncogene* 26, 4453.

786 Tusher, V.G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays
787 applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* 98, 5116-5121.

788 Uebbing, S., Konzer, A., Xu, L., Backström, N., Brunström, B., Bergquist, J., and Ellegren, H.
789 (2015). Quantitative Mass Spectrometry Reveals Partial Translational Regulation for
790 Dosage Compensation in Chicken. *Molecular Biology and Evolution*.

791 Vallet, E., Kreutzer, M., and Gahr, M. (1996). Testosterone Induces Sexual Release Quality
792 in the Song of Female Canaries. *Ethology* 102, 617-628.

793 Van Meir, V., Verhoye, M., Absil, P., Eens, M., Balthazart, J., and Van der Linden, A. (2004).
794 Differential effects of testosterone on neuronal populations and their connections in a
795 sensorimotor brain nucleus controlling song production in songbirds: a manganese
796 enhanced-magnetic resonance imaging study. *Neuroimage* 21, 914-923.

797 Vellema, M., Diales Rocha, M., Bascones, S., Zsebők, S., Dreier, J., Leitner, S., Van der
798 Linden, A., Brewer, J., and Gahr, M. (2019a). Accelerated redevelopment of vocal skills is
799 preceded by lasting reorganization of the song motor circuitry. *eLife* 8, e43194.

800 Vellema, M., Rocha, M., Bascones, S., Zsebők, S., Dreier, J., Leitner, S., Van der Linden, A.,
801 Brewer, J., and Gahr, M. (2019b). Accelerated redevelopment of vocal skills is preceded
802 by lasting reorganization of the song motor circuitry. *eLife* 8, e43194.

803 Voigt, C., and Leitner, S. (2008). Seasonality in song behaviour revisited: Seasonal and
804 annual variants and invariants in the song of the domesticated canary (*Serinus canaria*).
805 *Hormones and Behavior* 54, 373-378.

806 Wade, J. (2001). Zebra finch sexual differentiation: The aromatization hypothesis
807 revisited. *Microscopy Research and Technique* 54, 354-363.

808 Warren, W., Clayton, D., Ellegren, H., Arnold, A., Hillier, L., and Kunstner, A. (2010). The
809 genome of a songbird. *Nature* 464, 757-762.

810 Wickler, W., and Seibt, U. (1980). Vocal Dueting and the Pair Bond II. Uniseno dueting of
811 the African forest weaver, *Symplectes bicolor*. *Zeitschrift für Tierpsychologie* 52, 217-
812 226.

813 Wild, J.M. (2004). Functional Neuroanatomy of the Sensorimotor Control of Singing.
814 *Annals of the New York Academy of Sciences* 1016, 438-462.

815 Wilson, S., Qi, J., and Filipp, F.V. (2016). Refinement of the androgen response element
816 based on ChIP-Seq in androgen-insensitive and androgen-responsive prostate cancer
817 cell lines. *Scientific Reports* 6, 32611.

818 Wingfield, J.C., Smith, J.P., and Farner, D.S. (1982). Endocrine Responses of White-
819 Crowned Sparrows to Environmental Stress. *The Condor* 84, 399-409.

820 Wolf, J.B., and Bryk, J. (2011). General lack of global dosage compensation in ZZ/ZW
821 systems? Broadening the perspective with RNA-seq. *BMC Genomics* 12, 91.

822 Yang, X., Schadt, E.E., Wang, S., Wang, H., Arnold, A.P., Ingram-Drake, L., Drake, T.A., and
823 Lusis, A.J. (2006). Tissue-specific expression and regulation of sexually dimorphic genes
824 in mice. *Genome Res* 16, 995-1004.

825 Yates, A., Akanni, W., Amode, M.R., Barrell, D., Billis, K., Carvalho-Silva, D., Cummins, C.,
826 Clapham, P., Fitzgerald, S., Gil, L., *et al.* (2016). Ensembl 2016. *Nucleic Acids Research* 44,
827 D710-D716.

828 Zhang, R., Yin, Y., Sun, L., Yan, P., Zhou, Y., Wu, R., and Wu, X. (2017). Molecular cloning of
 829 ESR2 and gene expression analysis of ESR1 and ESR2 in the pituitary gland of the
 830 Chinese alligator (*Alligator sinensis*) during female reproductive cycle. *Gene* 623, 15-23.
 831 Zhao, C., Dahlman-Wright, K., and Gustafsson, J.-Å. (2010). Estrogen Signaling via
 832 Estrogen Receptor β . *Journal of Biological Chemistry* 285, 39575-39579.

833 Tables

834 **Table 1. Experimental groups and sample sizes.**

Species	Group	Sex	Singing	Tissue	Sample size
Forest weaver	Breeding	Female	Yes	HVC	6
		Male	Yes	HVC	6
Blue-capped cordon-bleu	Breeding	Female	Yes	HVC	6
		Male	Yes	HVC	6
	Breeding	Female	No	HVC	6
		Male	Yes	HVC	6
Canary	Nonbreeding	Female	Yes	HVC	6
		Female	No	HVC	5
		Male	No	HVC	6
	Nonbreeding + testosterone	Female	Yes	HVC	6
		Male	Yes	HVC	6

835

836 **Supplementary Table 1. Fisher's exact tests of sex-biased genes for chromosome**
837 **enrichment.**

838 **Supplementary Table 2. GO term enrichment analysis of sex-biased genes**
839 **identified from the forest weaver (comparison 1), cordon-bleu (comparison 2), and**
840 **canary (comparison 5) comparisons.**

Abbreviations: FWm, forest weaver male-biased genes; CBm, cordon-bleu male-biased genes; Cm, canary male-biased genes; FWf, forest weaver female-biased genes; CBf, cordon-bleu female-biased genes; CfS, canary female-biased genes.

841 **Supplementary Table 3. Pearson's correlation analysis of principal components**
842 **and variables (plasma androgen levels, HVC volume, sex, and singing).**

843 **Supplementary Table 4. Female-specific expressed genes.**

Abbreviations: SDfS: nonbreeding spontaneously singing female canaries; SDfT: nonbreeding testosterone-stimulated singing female canaries.

844 **Supplementary Table 5. Male-specific expressed genes.**

Abbreviations: LDm: breeding singing male canaries; SDmT: nonbreeding testosterone-stimulated singing male canaries.

845 **Supplementary Table 6. Sex-shared expressed genes.**

Abbreviations: LDm: breeding singing male canaries; SDfS: nonbreeding spontaneously singing female canaries; SDmT: nonbreeding testosterone-stimulated singing male canaries; SDfT: nonbreeding testosterone-stimulated singing female canaries.

846 **Supplementary Table 7. GO term enrichment analysis of female-specific, male-**
847 **specific and sex-shared genes.**

848

Supplementary Figures

Supplementary Figure 1. Plasma androgen levels of seven groups of canaries on the day of sacrifice.

A, Canaries in breeding conditions. Singing males 1.76 ng/ml (mean), nonsinging females 59.2 pg/ml. Mann-Whitney Test, $U = 6$, P value = 0.07. **B**, Canaries in nonbreeding conditions. Nonsinging males 66.7 pg/ml (mean), nonsinging females 47.2 pg/ml, singing females 52.6 pg/ml. Mann-Whitney Test (nonsinging males vs. nonsinging females), $U = 9$, P value = 0.172. Mann-Whitney Test (nonsinging males vs. singing females), $U = 24$, P value = 0.377. **C**, Testosterone-implanted nonbreeding canaries. Males 13.5 ng/ml, females 9.31 ng/ml. Mann-Whitney Test, $U = 13$, P value = 0.485. Testosterone implantation significantly increased the plasma androgen levels of both nonbreeding males and females Mann-Whitney Test (nonbreeding nonsinging males vs. testosterone-implanted singing males), $U = 0$, P value = 0.00492. Mann-Whitney Test (nonbreeding nonsinging females vs. testosterone-implanted singing females), $U = 0$, P value = 0.00217. The plasma androgen levels were higher in the breeding males than in the nonbreeding nonsinging males (Mann-Whitney Test, $U = 32$, P value = 0.0275). The boxes indicate the 25th/50th/75th percentiles (bottom/middle/top bar), and the extent of the whiskers indicates the most extreme values that are within 1.5 times the IQR (interquartile range) of the hinge. Each colour-coded dot indicates the measurement from one bird. The nonbreeding singing female canaries data were obtained from (Ko et al., 2020).

854

855 **Supplementary Figure 2. Hierarchical clustering of the HVC transcriptomes of 65**
 856 **birds used in this study.**

Hierarchical clustering showed that the HVC transcriptomes were first clustered based on the phylogenetic relationship; among canaries, nonbreeding females implanted with testosterone showed the most distinctive patterns. +T: testosterone implantation.

Supplementary Figure 3. Hierarchical clustering of the entopallium transcriptomes of 65 birds used in this study.

Hierarchical clustering showed that the entopallium transcriptomes were first clustered based on the phylogenetic relationship. The canaries were clustered by sex. +T: testosterone implantation.

860

861 **Supplementary Figure 4. The majority of sex-biased genes showed low sex bias.**

The bar graph summarizes the number of sex-biased genes in the HVC transcriptome identified from each male-to-female comparison. High bias: $|\log_2(\text{fold change})| \geq 2$; moderate bias: $1 \leq |\log_2(\text{fold change})| < 2$; low bias: $0.5 \leq |\log_2(\text{fold change})| < 1$. The phenotypes of the groups being compared in each male-to-female comparison are listed at the bottom of the graph.

Supplementary Figure 5. Pairwise comparisons between autosomal and Z chromosomal sex-biased genes.

A, Autosomal male-biased genes. **B**, Z chromosomal male-biased genes. **C**, Autosomal female-biased genes. **D**, Z chromosomal female-biased genes. The numbers indicated in the matrices are the proportions of identical genes identified from each pair. The phenotypes of the groups used for each comparison are listed at the bottom. B: breeding; NB: nonbreeding; T: testosterone implantation; the music note indicates the presence or absence of singing behaviour.

Supplementary Figure 6. PCA scree plot showing the percentage of variance explained by each principal component.

A **B** **C**

A

B

male	B ♪	B ♪	B ♪	NB + T ♪	B ♪	NB 🚫	NB 🚫
female	B ♪	B ♪	B 🚫	NB + T ♪	NB ♪	NB ♪	NB 🚫

A Male-biased genes

B Female-biased genes

C Species-specific putative functions

- GO:0072358: cardiovascular system development
- GO:0032963: collagen metabolic process
- GO:0048598: embryonic morphogenesis
- GO:0009887: organ morphogenesis
- GO:0048729: tissue morphogenesis
- GO:0030900: forebrain development
- GO:0007399: nervous system development
- GO:0061564: axon development
- GO:0007409: axonogenesis
- GO:0048667: cell morphogenesis involved in neuron differentiation
- GO:0048812: neuron projection morphogenesis
- GO:0048646: anatomical structure formation involved in morphogenesis
- GO:0000902: cell morphogenesis
- GO:0048858: cell projection morphogenesis
- GO:0032989: cellular component morphogenesis
- GO:0060271: cilium morphogenesis
- GO:0030574: collagen catabolic process
- GO:0051297: centrosome organization
- GO:0000280: nuclear division
- GO:0000819: sister chromatid segregation
- GO:0006352: DNA-templated transcription, initiation
- GO:0016071: mRNA metabolic process
- GO:0000398: mRNA splicing, via spliceosome
- GO:0008380: RNA splicing
- GO:0006367: transcription initiation from RNA polymerase II promoter
- GO:0032446: protein modification by small protein conjugation
- GO:0070647: protein modification by small protein conjugation or removal
- GO:0070646: protein modification by small protein removal
- GO:0016601: Rac protein signal transduction
- GO:1900746: regulation of vascular endothelial growth factor signaling pathway
- GO:0046486: glycerolipid metabolic process
- GO:0030258: lipid modification
- GO:0051270: regulation of cellular component movement
- GO:0070848: response to growth factor
- GO:0050954: sensory perception of mechanical stimulus
- GO:0007605: sensory perception of sound

Comparison

		1	2	5
M	B♂	B♂	B♂	
F	B♂	B♂	NB♂	
	Forest weaver	Cordon bleu	Canary

D Putative functions of the sex-specific singing-associated genes