1 *In vitro* efficacy of gentamicin against multidrug-resistant *Neisseria gonorrhoeae*:
2 synergy of three gentamicin antimicrobial combinations

3 Xuechun Li¹, Wenjing Le¹, Xiangdi Lou¹, Biwei Wang¹, Caroline A. Genco², Peter A.
 4 Rice³ and Xiaohong Su^{1#}

5 ¹Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union
 6 Medical College, Nanjing, China

7 ²Department of Immunology, School of Medicine, Tufts University, Boston, MA,
 8 USA.

9 ³Division of Infectious Diseases and Immunology, Department of Medicine,
 10 University of Massachusetts Medical School, Worcester, MA, United States

11

12 Running Head: gentamicin against multidrug-resistant gonococci

13 #Corresponding author:

14 Xiaohong Su, MD, PhD.

15 Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union

16 Medical College,

17 No. 12, Jiangwangmiao Road,

18 Nanjing 210042, Jiangsu Province, China

19 Tel: (86) 25-85478090

20 Fax: (86) 25-85414477

Email: suxh@ncstdlc.org

ABSTRACT

Objectives: To determine *in vitro* activities of gentamicin alone and in combination with ceftriaxone, ertapenem and azithromycin against multidrug-resistant (MDR) *N. gonorrhoeae* isolates.

Methods: 407 isolates from Nanjing, China, obtained in 2016 and 2017, had minimum inhibitory concentrations (MICs) determined for gentamicin using the agar dilution method. Antimicrobial combinations were also tested in 97 MDR strains using the antimicrobial gradient epsilometer test (Etest); results ranging from synergy to antagonism were interpreted using the fractional inhibitory concentration (FICI).

Results: All 407 gonococcal isolates were susceptible to gentamicin. MICs ranged from 2 mg/L to 16 mg/L. Synergy was demonstrated in 16.5% (16/97), 27.8% (27/97) and 8.2% (8/97) MDR strains when gentamicin was combined with ceftriaxone [geometric mean (GM) FICI; 0.747], ertapenem (GM FICI; 0.662) and azithromycin (GM FICI; 1.021), respectively. No antimicrobial antagonism was observed with any combination. The three antimicrobial combinations were indifferent overall. The overall GM MICs of gentamicin were reduced by 2.63-, 3.80- and 1.98-fold when tested in combination with ceftriaxone, ertapenem and azithromycin, respectively. The GM MICs of the three antimicrobials by themselves were reduced by 3-, 2.57-

and 1.98-fold respectively, when each was tested in combination with gentamicin. No antimicrobial antagonism was observed with any combination.

Conclusions: Gentamicin alone was effective *in vitro* against MDR *N. gonorrhoeae* and in combination with ceftriaxone, ertapenem or azithromycin. Combination testing of resistant strains, overall, showed lower effective MICs against gentamicin itself and each of the three antimicrobials when used in combination with gentamicin.

Keywords: *Neisseria gonorrhoeae*, multidrug resistance, combination, gentamicin

INTRODUCTION

Gonorrhea, caused by *Neisseria gonorrhoeae*, is a common bacterial sexually transmitted infection (STI), which is responsible for an increasing burden of disease globally. In 2016, the World Health Organization (WHO) estimated 87 million new cases of gonorrhea worldwide in adults, aged 15-49(1). Common features of gonococcal infection include urethritis and cervicitis; more serious manifestations include pelvic inflammatory disease, chronic pelvic pain, infertility, chorioamnionitis in women and epididymitis in men. Disseminated gonococcal infections (DGI) occurs occasionally in adults and neonates born to infected mothers; in the pre-antibiotic era endocarditis and meningitis sometimes resulted from gonococcal bacteremia but is rarely seen today. Concomitant gonococcal and HIV infections also increase the risk of HIV transmission (2).

61 The emergence of antibiotic resistance among *N. gonorrhoeae* is a global public
62 health threat. *N. gonorrhoeae* has developed resistance to all antimicrobials that have
63 been used for treatment, including penicillin, tetracycline, ciprofloxacin and
64 azithromycin, leading to the emergence of multidrug-resistant (MDR) isolates, which
65 are difficult to treat (3). Currently, dual antimicrobial therapy with an extended
66 spectrum cephalosporin (ESC), either ceftriaxone 250 mg intramuscularly or cefixime
67 400 mg orally, plus azithromycin 1g orally, is recommended as first-line treatment of
68 uncomplicated gonorrhea by the World Health Organization (WHO) (4). Increased
69 prevalence of azithromycin resistance globally prompted a revision of prior
70 recommendations in the United Kingdom (U.K.) (5) in 2018 and the United States
71 (U.S.) (6) in 2020 from dual therapy with ceftriaxone and azithromycin to
72 recommendations of higher doses of ceftriaxone: 1 gram (U.K.) and 500 mg (U.S.).
73 Unfortunately, resistance to ESCs (7-9), the last remaining option for empirical
74 first-line monotherapy, threatens future use of this class of antimicrobials. Treatment
75 failures with both mono- and dual-therapy (including azithromycin) have been
76 reported in recent years (10, 11).
77 Treatment options for gonorrhea including infections caused by MDR organisms are
78 diminishing; there is an urgent need to explore new or repurposed antimicrobial
79 agents and/or therapeutic strategies. Gentamicin, an aminoglycoside antibiotic that
80 inhibits protein synthesis by binding to the 30S ribosomal subunit, has been used as
81 first-line therapy for treatment of gonorrhea in several countries, including Malawi,

where it has been used officially for nearly 30 years (12). Numerous *in vitro* susceptibility studies have shown that gentamicin is active against *N. gonorrhoeae* (13-16), suggesting that gentamicin is appropriate for treatment of uncomplicated gonococcal infection. A single 240 mg dose of intramuscular gentamicin was 95% efficacious in a randomized controlled clinical trial to treat urethritis (17). In multicenter clinical trials, dual therapy with gentamicin plus azithromycin was 94%-100% effective compared to 98% for ceftriaxone plus azithromycin (18, 19). Little is known about *in vitro* susceptibility of gentamicin in Chinese isolates where this antimicrobial has not been used to treat gonorrhea. Use of antimicrobial combinations is a therapeutic strategy intended to increase efficacy and slow the development of resistance (20). Antimicrobial combinations may prove useful to successfully manage MDR *N. gonorrhoeae* infections and they are included in current WHO and CDC guidelines (4, 6). Initially, we evaluated gentamicin susceptibility of gonococcal strains isolated between 2016 and 2017 in Nanjing, China. Second, we carried out studies with antimicrobial combinations that included gentamicin to evaluate possible *in vitro* enhancement of gentamicin activity against MDR strains when tested in combination with either ceftriaxone (an ESC), ertapenem (a carbapenem) or azithromycin (a macrolide). Third, we determined the MICs of each of the three antimicrobials by themselves when tested in combination with gentamicin.

RESULTS

All 407 *N. gonorrhoeae* strains were sensitive to gentamicin by agar dilution; minimum inhibitory concentrations (MICs) ranged from 2mg/L to 16mg/L; the MIC₅₀ was 8 mg/L and the MIC₉₀ was 16mg/L. Among the 407 isolates, 34 (8.4%) were susceptible (MIC ≤ 4mg/L), 373 (91.6%) were intermediately susceptible (MIC range, 8-16 mg/L) and none was resistant (MIC ≥ 32 mg/L) (Fig.1). Among the 34 fully susceptible isolates, 6 (17.6%) had decreased susceptibility to ESCs i.e. ceftriaxone or cefixime or both and 1 (2.9%) was fully resistant to ceftriaxone (MIC; 1mg/L) and cefixime (MIC; 2mg/L). There was no change in intermediate susceptibility to gentamicin of isolates between 2016 and 2017 (89.3% vs. 93.4%; $\chi^2=2.225$, P=0.136) (Fig. 2).

The distribution of antimicrobial susceptibility patterns for the 97 MDR strains selected for antimicrobial combination testing showed 6 unique patterns of susceptibility (Table S1). 93 (95.9%) strains had decreased susceptibility to ceftriaxone or cefixime or both plus resistance to ciprofloxacin and penicillin, and 4 (4.1%) strains had decreased susceptibility to ceftriaxone or cefixime or both plus resistance to ciprofloxacin, penicillin and azithromycin. Resistance to gentamicin in MDR isolates was not detected by either the agar dilution or the Etest method. Agreement of MICs between agar dilution and Etest among MDR isolates is summarized in Table S2. Overall agreement of MICs (≤ 2-fold different) between the two methods was 93.8%. Etest always resulted in one to two dilutions lower MIC values than agar dilution.

Gentamicin agar dilution versus Etest MIC results (mg/L) were as follows: MIC₅₀ = 8

versus 6; MIC₉₀ =16 versus 8; geometric mean [GM] MIC =11.3 versus 5.84.

Additionally, there was a discrepancy in the full susceptibility category (29.9% fully susceptible by Etest versus 9.3% by agar dilution; $\chi^2=13.090$, $P<0.001$).

The three antimicrobial combinations used to test each of the 97 strains were examined for effects that were classified as: synergistic, indifferent or antagonistic-- summarized in Table 1. For example, the gentamicin GM MIC, when tested alone, was 5.840 mg/L; when combined with ceftriaxone, the GM MIC was reduced to 2.217 mg/L (2.63-fold reduction, $P<0.001$) (Table 2). Together with ceftriaxone, gentamicin exhibited synergy against 16.5% (16/97) MDR strains; overall, the combination was indifferent; FICI, 0.747. When tested alone, MICs of ceftriaxone ranged from 0.016-0.75 mg/L; in combination with gentamicin, the GM MIC against ceftriaxone decreased from 0.078 to 0.026 mg/L (3-fold reduction, $P<0.001$).

The gentamicin GM MIC when combined with ertapenem decreased to 1.536 mg/L (3.80-fold reduction, $P<0.001$) (Table 2). Gentamicin together with ertapenem was the most synergistic combination, displaying synergy against 27.8% (27/97) of MDR gonococcal isolates; overall, this combination was indifferent; FICI, 0.662. Ertapenem MICs of 97 MDR isolates, when tested alone, ranged from 0.006 to 0.064 mg/L (GM MIC, 0.018 mg/L); when combined with gentamicin, the GM MIC decreased to 0.007 mg/L (2.57-fold reduction, $P<0.001$).

The gentamicin GM MIC when combined with azithromycin decreased to 2.949 mg/L (1.98-fold reduction, $P<0.001$) (Table 2). Together with azithromycin, gentamicin

exhibited synergy against 8.2% (8/97) MDR strains; overall, the combination was indifferent; FICI, 1.021. When tested alone, MICs of azithromycin ranged from 0.047-8 mg/L; in combination with gentamicin, the GM MIC against azithromycin decreased from 0.347 to 0.175 mg/L (1.98-fold reduction, $P < 0.001$).

DISCUSSION

Our study provides data on gentamicin susceptibility against *N. gonorrhoeae* isolated in Nanjing (Jiangsu Province), China. On agar dilution testing, most strains displayed intermediate susceptibility to gentamicin (MIC 8-16 mg/L), similar to a study that examined gonococcal isolates from seven hospitals in a neighboring eastern Chinese province; in that study 97.8% (493/504) of strains possessed gentamicin MICs of 8-16 mg/L (21). European and U.S. studies have reported 82.7% (15) and 73% (18), intermediate susceptibility, respectively, of *N. gonorrhoeae* isolates to gentamicin. A recent report of gentamicin susceptibility of *N. gonorrhoeae* in which 86.0% of 470 isolates were fully susceptible (MICs ≤ 4 mg/L), examined isolates from seven geographically distributed Chinese provinces as part of the China Gonococcal Resistance Surveillance Programme (China-GRSP) (22) similar to an Indian study where 90.7% of isolates were reported as fully susceptible (15). Our study compared gentamicin MICs using agar dilution and Etest methods for 97 multidrug-resistant (MDR) *N. gonorrhoeae* isolates. Similar to previous studies (13, 23), we found that over 90% of gentamicin MICs determined by agar dilution and

Etest were ≤ 2 -fold different; typically, Etest resulted in lower MICs and identified a larger proportion of fully susceptible isolates. In particular, all MDR isolates were fully or intermediately susceptible to gentamicin.

Synergistic or additive effects of combining antimicrobials for treatment may slow the development of antimicrobial resistance of *N. gonorrhoeae* (24). We assessed the *in vitro* activity of gentamicin in combination with 3 antibiotics. Determining synergy has several challenges. Several test methods are available to evaluate synergistic effects of antimicrobial combinations, however, they are not well standardized. We chose Etest because it is practical and also correlates well with agar dilution, time–kill curves and checkerboard testing in demonstrating synergy for two-drug combination (25-27). Nonetheless, FICI interpretation depends on the criteria used. FICI values between 0.5 and 1 have been interpreted as additive in Indian (28) and Japanese (29) studies, differing from our criteria, which classifies FICI values in this range as indifferent.

Ceftriaxone, in higher doses, is now recommended as single therapy by the U.S. and the U.K. for treatment of uncomplicated gonorrhea (5, 6). Our study showed that the combination of ceftriaxone and gentamicin exhibited an indifferent effect overall in $>80\%$ of MDR strains ($< 20\%$ synergy). In the Indian study by Singh *et al* (28), 14.7% synergy and 6.3% antagonism were reported for this combination against 95 *N. gonorrhoeae* strains including 79 MDR and one extensively drug-resistant (XDR) strain. In a Canadian study, a mean FICI₅₀ value of 1.2 (0.8-2.0) was shown for nine

reference strains of *N. gonorrhoeae* (WHO F, G, K, L, M, N, O, P and ATCC 49226) with this combination (30). An American study reported a mean FICI of 1.25 (0.73-2) using gonococcal isolates that displayed different cefixime MICs (26). No synergistic/antagonistic effect (resulting in 100% indifference) was observed in either study (26, 30).

We chose ertapenem as a candidate for *in vitro* synergy testing because its mechanism of action differs from that of gentamicin and it has been used to treat infection with combined high-level azithromycin and ceftriaxone resistant *N. gonorrhoeae* (11). Ertapenem has demonstrated an advantage over ceftriaxone for MDR or ceftriaxone resistant isolates and has also been suggested for possible use in a dual antimicrobial regimen (31). We showed that gentamicin plus ertapenem in combination resulted in synergistic and indifferent effects with no antagonism demonstrated in any MDR strain. In the study by Singh *et al* (28), this combination displayed either synergy (31.6%) or indifference (68.4%) in 100% of strains; no antagonism was seen.

Gentamicin in combination with azithromycin is currently recommended as an option for retreatment by the WHO when dual therapy fails (4). Also, it is proposed as an alternative CDC recommendation when higher dose ceftriaxone therapy cannot be used (6). In our studies, this combination demonstrated synergy in fewer MDR isolates (<10%) than combinations with either ceftriaxone or ertapenem and exhibited the highest FICI value of the three combinations tested. Similar to our results, Sood *et al* (32) demonstrated synergistic effects in 22.9% of isolates displaying different

ceftriaxone MICs and no antagonism for this combination. All isolates with differing cefixime/ ceftriaxone MICs showed indifference with a mean FICI of 1.7/0.83 in studies from the UK (33) and Japan (29). The separate study by Singh *et al* (28) differed from these results, with 6.3% of strains exhibiting antagonism when this combination was used.

A summary of results from these previous studies is shown in Table S3. No synergistic/antagonistic effect (resulting in 100% indifference) was observed in studies from Canada (30), Japan (29), the US (26) and the UK (33). None of these studies incorporated isolates with multi-drug resistance. However, a certain proportion of synergistic effects was observed in the two Indian studies (28, 32). Antagonism was also observed in combinations of gentamicin with ceftriaxone and azithromycin in the study by Singh *et al* (28). Sood *et al* (32) used the same Etest that we used. In contrast, Singh *et al* (28) incubated the Etest strip of the first antimicrobial (antimicrobial A) for 1 h, and then replaced it with the Etest strip of the second antimicrobial (antimicrobial B) at the same location and looked for synergism. A mild degree of antagonism may have been missed in our tests when the zone of inhibition ran under the strips where they crossed and therefore was unreadable and interpreted as indifference (25).

From a pharmacokinetic perspective, all antimicrobials in the 3 combinations in our study result in peak levels of drug during the first 3 hours after administration (34-37). However, azithromycin has a longer half-life than gentamicin (approximately 68 and

2 hours, respectively) (34, 37). Gentamicin in combination with azithromycin also produced the lowest synergistic effects among the 3 combinations in our study so it may not be optimal for clinical use where synergy would not be prolonged.

In conclusion, resistance to gentamicin was not observed in gonococcal isolates examined in this study, including MDR isolates, indicating potential for its therapeutic use. Antimicrobial combinations of gentamicin plus ertapenem, ceftriaxone, and azithromycin showed no antagonistic effects; enhanced efficacy of individual antimicrobials in the presence of other antimicrobials was also demonstrated. Improved efficacy of antimicrobial combinations suggest that dual therapy might be a promising approach to manage MDR *N. gonorrhoeae*. Further studies to correlate *in vitro* results with clinical outcomes are warranted.

MATERIALS AND METHODS

Bacterial strains 407 gonococcal isolates were recovered from men with symptomatic urethritis (urethral discharge and/or dysuria) attending the STD clinic at the Institute of Dermatology, Chinese Academy of Medical Sciences in Nanjing, between January 2016 and December 2017. Urethral specimens were collected with cotton swabs and immediately streaked onto modified Thayer-Martin medium (Zhuhai DL Biotech Co. Ltd.) and cultured in candle jars at 36°C for 24–48 h. Gonococcal isolates were identified by colonial morphology, Gram's stain, and oxidase testing and sub-cultured onto GC chocolate agar base (Difco, Detroit, MI)

supplemented with 1% IsovitaleX™ (Oxoid, USA); pure cultures were swabbed, suspended in tryptone-based soy broth and frozen (−80°C) until used for antimicrobial testing.

Antimicrobial susceptibility testing The MICs of the 407 isolates were determined using the agar dilution method for gentamicin, penicillin, tetracycline, ciprofloxacin, azithromycin, spectinomycin, cefixime and ceftriaxone, used singly, according to the Clinical and Laboratory Standards Institute (CLSI) guidelines (38). *N. gonorrhoeae* ATCC 49226, WHO reference strains F, G, L, O and P were used as quality control strains in susceptibility tests. Although formal susceptibility criteria for gentamicin have not been established by CLSI, criteria described by CDC (16) have used MICs of ≤4 mg/L as fully susceptible, 8 to 16 mg/L as intermediately susceptible and ≥32 mg/L as resistant. Resistance to azithromycin (MIC ≥1 mg/L) was determined using EUCAST criteria (39). Susceptibilities to other antibiotics were assessed based on CLSI standards (38). Decreased susceptibility to cephalosporins was determined according to WHO standards (40). Based on criteria proposed by Tapsall *et al* in 2009 (41), MDR isolates were defined as those resistant or with decreased susceptibility to one or more widely used antimicrobials (ceftriaxone and cefixime; category I) and resistant to two or more antimicrobials, which are used less frequently (penicillin; ciprofloxacin and azithromycin; category II) (Table S4).

Synergy testing and interpretation Of 407 clinical isolates of *N. gonorrhoeae*, 97 MDR isolates (determined below) were selected for antimicrobial combination testing

(synergy/ antagonism) according to MDR criteria. WHO-P was used as a quality control strain. Dual antimicrobial testing was performed to evaluate the efficacy of gentamicin in combination with either ceftriaxone, ertapenem or azithromycin, using the Etest method, described previously (25). Briefly, MICs for individual antimicrobials (MIC A_{alone} and MIC B_{alone}) against *Neisseria gonorrhoeae* isolates were determined using Etest strips (Liofilchem, Italy); *in vitro* activity of each combination was determined by placing E-test strips of the two antimicrobials on the agar plates at a 90° angle, with intersections at the points of their individual MICs (Fig. 3). Agar plates were inverted during incubation at 36°C in 5% CO₂ for 16–18 hours, and the MIC of each antimicrobial in the combination (MIC A_{in combination} and MIC B_{in combination}) was read.

To determine whether each antimicrobial combination resulted in a synergistic, indifferent or antagonistic effect, the fractional inhibitory concentration index (FICI) was calculated using the following formula: $FICI = (MIC A_{in combination} / MIC A_{alone}) + (MIC B_{in combination} / MIC B_{alone})$ (25). FICI values were interpreted using the following criteria: synergy: ≤ 0.5 ; indifference: $FIC > 0.5$ to ≤ 4.0 and antagonism: FICI of > 4.0 (25).

Statistical Analysis The chi-square test was used to compare gentamicin susceptibility trend data and the categorical assignments of gentamicin susceptibility by the agar dilution or Etest method. Mean values of MICs and FICIs were calculated as geometric means (GM). The statistical significance of the difference between the MIC

of each of the three antimicrobials tested alone and in combination with gentamicin was determined using the nonparametric Mann-Whitney's U-test. P values less than 0.05 were considered statistically significant.

ACKNOWLEDGEMENTS

This work was supported by the grants from the Chinese Academy of Medical Sciences Initiative for Innovative Medicine (2016-I2M-3-021) and the U.S. National Institutes of Health (AI084048 and AI116969).

CONFLICTS OF INTEREST

No conflicts for all authors.

REFERENCES

1. World Health Organization. Report on global sexually transmitted infection surveillance -2018[M]. World Health Organization, 2018; Available from: <https://www.who.int/publications/i/item/9789241565691>.
2. Mlisana, K., N. Naicker, L. Werner, L. Roberts, F. van Loggerenberg, C. Baxter, J. A. Passmore, A. C. Grobler, A. W. Sturm, C. Williamson, K. Ronacher, G. Walzl, and K. S. Abdool. 2012. Symptomatic vaginal discharge is a poor predictor of sexually transmitted infections and genital tract inflammation in high-risk women in South Africa. *J Infect Dis* 206:6-14.

- 313 3. Unemo, M., C. S. Bradshaw, J. S. Hocking, H. de Vries, S. C. Francis, D. Mabey,
314 J. M. Marrazzo, G. Sonder, J. R. Schwebke, E. Hoornenborg, R. W. Peeling, S. S.
315 Philip, N. Low, and C. K. Fairley. 2017. Sexually transmitted infections: challenges
316 ahead. *Lancet Infect Dis* 17:e235-e279.
- 317 4. World Health Organization (WHO). 2016. WHO guidelines for the treatment of
318 *Neisseria gonorrhoeae*. WHO: Geneva, Switzerland, 2016; Available from:
319 [https://www.who.int/reproductivehealth/publications/rtis/gonorrhoea-treatment-guidel](https://www.who.int/reproductivehealth/publications/rtis/gonorrhoea-treatment-guidelines/en/)
320 [ines/en/](https://www.who.int/reproductivehealth/publications/rtis/gonorrhoea-treatment-guidelines/en/).
- 321 5. Fifer, H., J. Saunders, S. Soni, S. T. Sadiq, and M. FitzGerald. 2020. 2018 UK
322 national guideline for the management of infection with *Neisseria gonorrhoeae*. *Int J*
323 *STD AIDS* 31:4-15.
- 324 6. St, C. S., L. Barbee, K. A. Workowski, L. H. Bachmann, C. Pham, K. Schlanger,
325 E. Torrone, H. Weinstock, E. N. Kersh, and P. Thorpe. 2020. Update to CDC's
326 Treatment Guidelines for Gonococcal Infection, 2020. *MMWR Morb Mortal Wkly*
327 *Rep* 69:1911-1916.
- 328 7. Ohnishi, M., D. Golparian, K. Shimuta, T. Saika, S. Hoshina, K. Iwasaku, S.
329 Nakayama, J. Kitawaki, and M. Unemo. 2011. Is *Neisseria gonorrhoeae* initiating a
330 future era of untreatable gonorrhea?: detailed characterization of the first strain with
331 high-level resistance to ceftriaxone. *Antimicrob Agents Chemother* 55:3538-45.
- 332 8. Camara, J., J. Serra, J. Ayats, T. Bastida, D. Carnicer-Pont, A. Andreu, and C.
333 Ardanuy. 2012. Molecular characterization of two high-level ceftriaxone-resistant

334 *Neisseria gonorrhoeae* isolates detected in Catalonia, Spain. J Antimicrob Chemother
335 67:1858-60.

336 9. Unemo, M., D. Golparian, R. Nicholas, M. Ohnishi, A. Gallay, and P. Sednaoui.
337 2012. High-level cefixime- and ceftriaxone-resistant *Neisseria gonorrhoeae* in
338 France: novel penA mosaic allele in a successful international clone causes treatment
339 failure. Antimicrob Agents Chemother 56:1273-80.

340 10. Fifer, H., U. Natarajan, L. Jones, S. Alexander, G. Hughes, D. Golparian, and M.
341 Unemo. 2016. Failure of Dual Antimicrobial Therapy in Treatment of Gonorrhea. N
342 Engl J Med 374:2504-6.

343 11. Eyre, D. W., N. D. Sanderson, E. Lord, N. Regisford-Reimmer, K. Chau, L.
344 Barker, M. Morgan, R. Newnham, D. Golparian, M. Unemo, D. W. Crook, T. E. Peto,
345 G. Hughes, M. J. Cole, H. Fifer, A. Edwards, and M. I. Andersson. 2018. Gonorrhoea
346 treatment failure caused by a *Neisseria gonorrhoeae* strain with combined ceftriaxone
347 and high-level azithromycin resistance, England, February 2018. Euro Surveill 23.

348 12. Ross, J. D., and D. A. Lewis. 2012. Cephalosporin resistant *Neisseria*
349 *gonorrhoeae*: time to consider gentamicin? Sex Transm Infect 88:6-8.

350 13. Chisholm, S. A., N. Quaye, M. J. Cole, H. Fredlund, S. Hoffmann, J. S. Jensen,
351 M. J. van de Laar, M. Unemo, and C. A. Ison. 2011. An evaluation of gentamicin
352 susceptibility of *Neisseria gonorrhoeae* isolates in Europe. J Antimicrob Chemother
353 66:592-595.

354 14. Lagace-Wiens, P., H. J. Adam, N. M. Laing, M. R. Baxter, I. Martin, M. R.

355 Mulvey, J. A. Karlowsky, D. J. Hoban, and G. G. Zhanel. 2017. Antimicrobial
356 susceptibility of clinical isolates of *Neisseria gonorrhoeae* to alternative
357 antimicrobials with therapeutic potential. J Antimicrob Chemother 72:2273-2277.

358 15. Bala, M., V. Singh, A. Bhargava, M. Kakran, N. C. Joshi, and R. Bhatnagar.
359 2016. Gentamicin Susceptibility among a Sample of Multidrug-Resistant *Neisseria*
360 *gonorrhoeae* Isolates in India. Antimicrob Agents Chemother 60:7518-7521.

361 16. Mann, L. M., R. D. Kirkcaldy, J. R. Papp, and E. A. Torrone. 2018. Susceptibility
362 of *Neisseria gonorrhoeae* to Gentamicin—Gonococcal Isolate Surveillance Project,
363 2015-2016. Sex Trans Dis 45:96-98.

364 17. Lule, G., F. M. Behets, I. F. Hoffman, G. Dallabetta, H. A. Hamilton, S. Moeng,
365 G. Liomba, and M. S. Cohen. 1994. STD/HIV control in Malawi and the search for
366 affordable and effective urethritis therapy: a first field evaluation. Genitourin Med
367 70:384-388.

368 18. Kirkcaldy, R. D., H. S. Weinstock, P. C. Moore, S. S. Philip, H. C. Wiesenfeld, J.
369 R. Papp, P. R. Kerndt, S. Johnson, K. G. Ghanem, E. W. Hook, L. M. Newman, D.
370 Dowell, C. Deal, J. Glock, L. Venkatasubramanian, L. McNeil, C. Perlowski, J. Y.
371 Lee, S. Lensing, N. Trainor, S. Fuller, A. Herrera, J. S. Carlson, H. Harbison, C.
372 Lenderman, P. Dixon, A. Whittington, I. Macio, C. Priest, A. Jett, T. Campbell, A.
373 Uniyal, L. Royal, M. Mejia, J. Vonghack, S. Tobias, J. Zenilman, J. Long, A. Harvey,
374 K. Pettus, and S. Sharpe. 2014. The Efficacy and Safety of Gentamicin Plus
375 Azithromycin and Gemifloxacin Plus Azithromycin as Treatment of Uncomplicated

376 Gonorrhea. Clin Infect Dis 59:1083-1091.

377 19. Ross, J. D., J. Harding, L. Duley, A. A. Montgomery, T. Hepburn, W. Tan, C.

378 Brittain, G. Meakin, K. Sprange, S. Thandi, L. Jackson, T. Roberts, J. Wilson, J.

379 White, C. Dewsnap, M. Cole, and T. Lawrence. 2019. Gentamicin as an alternative to

380 ceftriaxone in the treatment of gonorrhoea: the G-TOG non-inferiority RCT. Health

381 Technol Assess 23:1-104.

382 20. Bollenbach, T. 2015. Antimicrobial interactions: mechanisms and implications

383 for drug discovery and resistance evolution. Curr Opin Microbiol 27:1-9.

384 21. Yang, F., J. Yan, J. Zhang, and S. van der Veen. 2020. Evaluation of alternative

385 antibiotics for susceptibility of gonococcal isolates from China. Int J Antimicrob

386 Agents 55:105846.

387 22. Liu, J. W., W. Q. Xu, X. Y. Zhu, X. Q. Dai, S. C. Chen, Y. Han, J. Liu, X. S.

388 Chen, and Y. P. Yin. 2019. Gentamicin susceptibility of *Neisseria gonorrhoeae*

389 isolates from 7 provinces in China. Infect Drug Resist 12:2471-2476.

390 23. Daly, C. C., I. Hoffman, M. Hobbs, M. Maida, D. Zimba, R. Davis, G.

391 Mughogho, and M. S. Cohen. 1997. Development of an antimicrobial susceptibility

392 surveillance system for *Neisseria gonorrhoeae* in Malawi: comparison of methods. J

393 Clin Microbiol 35:2985-8.

394 24. Lee, H., K. Lee, and Y. Chong. 2016. New treatment options for infections

395 caused by increasingly antimicrobial-resistant *Neisseria gonorrhoeae*. Expert Rev

396 Anti Infect ther 14:243.

- 397 25. White, R. L., D. S. Burgess, M. Manduru, and J. A. Bosso. 1996. Comparison of
398 three different in vitro methods of detecting synergy: time-kill, checkerboard, and E
399 test. *Antimicrob Agents Chemother* 40:1914-8.
- 400 26. Barbee, L. A., O. O. Soge, K. K. Holmes, and M. R. Golden. 2014. In vitro
401 synergy testing of novel antimicrobial combination therapies against *Neisseria*
402 *gonorrhoeae*. *J Antimicrob Chemother* 69:1572-1578.
- 403 27. Wind, C. M., H. J. de Vries, and A. P. van Dam. 2015. Determination of in vitro
404 synergy for dual antimicrobial therapy against resistant *Neisseria gonorrhoeae* using
405 Etest and agar dilution. *Int J Antimicrob Agents* 45:305-8.
- 406 28. Singh, V., M. Bala, A. Bhargava, M. Kakran, and R. Bhatnagar. 2018. In vitro
407 efficacy of 21 dual antimicrobial combinations comprising novel and currently
408 recommended combinations for treatment of drug resistant gonorrhoea in future era.
409 *PLoS One* 13:e0193678.
- 410 29. Furuya, R., Y. Koga, S. Irie, M. Tanaka, F. Ikeda, A. Kanayama, and I.
411 Kobayashi. 2013. In vitro activities of antimicrobial combinations against clinical
412 isolates of *Neisseria gonorrhoeae*. *J Infect Chemother* 19:1218-1220.
- 413 30. Bharat, A., I. Martin, G. G. Zhanel, and M. R. Mulvey. 2016. In vitro potency and
414 combination testing of antimicrobial agents against *Neisseria gonorrhoeae*. *J Infect*
415 *Chemother* 22:194-7.
- 416 31. Unemo, M., D. Golparian, A. Limnios, D. Whiley, M. Ohnishi, M. M. Lahra, and
417 J. W. Tapsall. 2012. In vitro activity of ertapenem versus ceftriaxone against

418 *Neisseria gonorrhoeae* isolates with highly diverse ceftriaxone MIC values and
 419 effects of ceftriaxone resistance determinants: ertapenem for treatment of gonorrhea?
 420 Antimicrob Agents Chemother 56:3603-9.

421 32. Sood, S., S. K. Agarwal, R. Singh, S. Gupta, and V. K. Sharma. 2019. In vitro
 422 assessment of gentamicin and azithromycin-based combination therapy against
 423 *Neisseria gonorrhoeae* isolates in India. J Med Microbiol 68:555-559.

424 33. Pereira, R., M. J. Cole, and C. A. Ison. 2013. Combination therapy for
 425 gonorrhoea: in vitro synergy testing. J Antimicrob Chemother 68:640-643.

426 34. Siber, G. R., P. Echeverria, A. L. Smith, J. W. Paisley, and D. H. Smith. 1975.
 427 Pharmacokinetics of gentamicin in children and adults. J Infect Dis 132:637-51.

428 35. Richards, D. M., R. C. Heel, R. N. Brogden, T. M. Speight, and G. S. Avery.
 429 1984. Ceftriaxone. A review of its antibacterial activity, pharmacological properties
 430 and therapeutic use. Drugs 27:469-527.

431 36. Nix, D. E., A. K. Majumdar, and M. J. DiNubile. 2004. Pharmacokinetics and
 432 pharmacodynamics of ertapenem: an overview for clinicians. J Antimicrob
 433 Chemother 53 Suppl 2:ii23-8.

434 37. Crokaert, F., A. Hubloux, and P. Cauchie. 1998. A Phase I Determination of
 435 Azithromycin in Plasma during a 6-Week Period in Normal Volunteers after a
 436 Standard Dose of 500mg Once Daily for 3 Days. Clin Drug Investig 16:161-6.

437 38. Clinical and Laboratory Standards Institute. 2018. Performance Standards for
 438 Antimicrobial Susceptibility Testing: Twenty-Eighth Informational Supplement M100.

439 CLSI, Wayne, PA, USA.

440 39. The European Committee on Antimicrobial Susceptibility Testing. 2015.

441 Breakpoint tables for interpretation of MICs and zone diameters. Version 5.0, 2015.

442 [https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/](https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_5.0_Breakpoint_Table_01.pdf)

443 [v_5.0_Breakpoint_Table_01.pdf](https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_5.0_Breakpoint_Table_01.pdf).

444 40. WHO. 2012. Global Action Plan to Control the Spread and Impact of

445 Antimicrobial Resistance in *Neisseria gonorrhoeae*. Available

446 from: http://whqlibdoc.who.int/publications/2012/9789241503501_eng.pdf?ua=1

447 41. Tapsall, J. W., F. Ndowa, D. A. Lewis, and M. Unemo. 2009. Meeting the public

448 health challenge of multidrug- and extensively drug-resistant *Neisseria gonorrhoeae*.

449 Expert Rev Anti Infect Ther 7:821-34.

450

451

452

453

454 Figure 1. Susceptibility of gentamicin against *N. gonorrhoeae* strains isolated in

455 Nanjing, China, 2016–2017 (n =407).

456

457

Figure 2. Distributions of MICs of gentamicin against *N. gonorrhoeae* isolates in 2016 (n=178) and 2017 (n=229).

Figure 3. Photograph of strip placement for E-test synergy method.

478

479 Table 1. Synergy test results for combinations of gentamicin plus azithromycin,

480 ceftriaxone, ertapenem against 97 MDR *N. gonorrhoeae* isolates

Effect n (%)	Antimicrobial combinations		
	GEN +CRO	GEN +ETP	GEN +AZM
Synergistic n (%)	16(16.5%)	27(27.8%)	8(8.2%)
Indifferent n (%)	81(83.5%)	70(72.2%)	89(91.8%)
Antagonistic n (%)	0	0	0
FICI (geometric mean)	0.747	0.662	1.021
Classification Overall	Indifferent	Indifferent	Indifferent

481 Abbreviations: n, number; GEN, gentamicin; CRO, ceftriaxone; ETP, ertapenem;

482 AZM, azithromycin.

483

484

485

486

487

488

489

490

491

492 Table 2. *In vitro* antibacterial activity of tested antibiotics combinations against 97

493 MDR *N. gonorrhoeae* isolates

MIC	Antimicrobial combinations		
	GEN +CRO	GEN +ETP	GEN +AZM
MIC A _{alone}	5.840(2-12)	5.840(2-12)	5.840(2-12)
MIC A _{in combination}	2.217(0.38-6)	1.536(0.25-6)	2.949(1-8)
MIC B _{alone}	0.078(0.016-0.75)	0.018(0.006-0.064)	0.347(0.047-8)
MIC B _{in combination}	0.026(0.004-0.25)	0.007(0.002-0.047)	0.175(0.023-2)

494 Abbreviations: GEN, gentamicin; CRO, ceftriaxone; ETP, ertapenem; AZM,

495 azithromycin; MIC, minimum inhibitory concentration.

496

497