1 Title: Mega- and meta-analyses of fecal metagenomic studies assessing response to immune checkpoint inhibitors

2
3 Authors: Alya Heirali^{1, *}, Bo Chen^{2, *}, Matthew Wong¹, Pierre HH Schneeberger¹, Victor Rey¹, Anna Spreafico^{2,3}, Wei Xu^{2,4, #}, Bryan
4 A. Coburn^{1,5, #}

5
6 Institutions:

- 7
8 1. Toronto General Hospital Research Institute, University Health Network, Toronto, Canada.
9 2. Princess Margaret Cancer Centre, Toronto, Canada.
10 3. Department of Medicine, Division of Medical Oncology and Hematology, University of Toronto, Toronto, Canada.
11 4. Dalla Lana School of Public Health, University of Toronto, Toronto, Canada.
12 5. Department of Medicine, Division of Infectious Diseases, University of Toronto, Toronto, Canada.
13

14 * Equally contributed

15 # Corresponding authors:

16 Bryan Coburn
17 bryan.coburn@utoronto.ca
18 University Health Network
19 101 College St., Rm 10-358
20 Toronto, Ontario, M5G 1L7
21

22 Wei Xu
23 wei.xu@uhnresearch.ca
24 Princess Margaret Cancer Centre
25 610 University Ave., Rm 10-511
26 Toronto, Ontario, M5G 2M9
27
28

29 Running title: Meta/mega analysis of fecal metagenomics and response to ICI

30
31 Keywords: Immune checkpoint inhibitors, programmed cell death protein 1, gut microbiome, RECIST response, biomarkers
32

Abstract

Purpose: Gut microbiota have been associated with response to immune checkpoint inhibitors (ICI) including anti-PD-1 and anti-CTLA-4 antibodies. However, inter-study difference in design, patient cohorts and data analysis pose challenges to identifying species consistently associated with response to ICI or lack thereof.

Experimental Design: We uniformly processed and analyzed data from three studies of microbial metagenomes in cancer immunotherapy response (four distinct data sets) to identify species consistently associated with response or non-response (n=190 patient samples). Metagenomic data were processed and analyzed using Metaphlan v2.0. Meta- and mega-analyses were performed using a two-part modelling approach of species present in at least 20% of samples to account for both prevalence and relative abundance differences between responders/non-responders.

Results: Meta- and mega-analyses identified five species that were concordantly significantly different between responders and non-responders. Amongst them, *Bacteroides thetaiotaomicron* and *Clostridium bolteae* relative abundance (RA) were independently predictive of non-response to immunotherapy when data sets were combined and analyzed using mega-analyses (AUC 0.59 95% CI 0.51-0.68 and AUC 0.61 95% CI 0.52-0.69, respectively).

Conclusions: Meta- and mega-analysis of published metagenomic studies identified bacterial species both positively and negatively associated with immunotherapy responsiveness across four published cohorts.

Introduction

A number of studies have demonstrated the gut microbiome is associated with response to immune checkpoint inhibitors (ICI)¹⁻⁵. Anti-programmed cell death protein 1 (PD-1) and anti-cytotoxic T lymphocyte-associated protein 4 (CTLA-4) targeting agents derepress anti-tumor T-cells. In the past five years, mouse models and human observational cohort studies have shown that the gut microbiome of responders to ICI is compositionally different from non-responders¹⁻⁶. However, inter-study differences in taxa associated with response to ICIs make it challenging to discern which organisms are consistently associated with response or non-response across studies and different cancers. This variability may in part be due to differences in sequence analytical pipelines. We conducted both meta- and mega-analyses of metagenomic studies of the gut microbiome in ICI recipients to determine species consistently enriched/depleted in responders compared to non-responders. Here, metagenomic data were selected to maximize the taxonomic resolution and delineate species-specific associations that may not be evident when taxa are annotated at the genus level.

Methods

Cohort Inclusion

We included three studies with publicly available metagenomic data and meta-data. Data was further divided into distinct data sets for the Routy *et al.* study to assess differences based on tumor type. These data sets include individuals with melanoma, renal cell carcinoma (RCC) and non-small cell lung cancer (NSCLC).

Definitions

Patients were classified as responders (R) or non-responders (NR) using the response evaluation criteria in solid tumors across all three studies (RECIST v1.1)^{3-5,7}. Data sets were analyzed in aggregate (meta/mega analysis) and separately. The primary outcome of

interest was to detect species consistently enriched/depleted in responders to ICI across data sets. This was achieved using a number of statistical methods described below. The secondary outcome was to identify predictors of response using receiver operator characteristics area under the curve (ROCAUC) analyses.

Microbiome Data processing

Raw sequencing data were obtained, and quality filtered using FastQC and MultiQC^{8,9}. Metaphlan v2.0 was used for its fast and robust species level annotation of microbial genomes¹⁰. Alpha-diversity measures were calculated using Phyloseq¹¹. Statistical analyses and data visualization were conducted in R and Prism^{12,13}.

Statistical Analyses

Two-part log-normal model

The raw sequencing data was observed as relative abundance (RA) of each species. Our target was to estimate the R/NR group differences of each species, where the differences consist of both species prevalence (presence/absence) and magnitude of abundance when species are present. We conducted two-part model analysis for species prevalence and relative abundance separately. Specifically, we used a logistic regression model to detect the association between species prevalence and R/NR. For RA data greater than zero, we observed log-normal distributions, and thus used linear regression model to find its association to R/NR group.

The logistic regression estimates (effect size) represent the log odds ratio of species prevalence between R and NR group, and linear regression estimates (effect size) represent the change of mean RA between R and NR group after excluding all zeros. We calculated 95% confidence intervals (CIs) for all estimates and tested if each estimate is significantly different from 0. A positive estimate means that R had a higher proportion of non-zeros among all samples, or R had a higher RA on average compared to NR. We noted that for

each species, estimates for prevalence and abundance may have different directions, which is described as “dissonant” effect¹⁴. In order not to let the opposite estimates reduce statistical test power, we tested each estimate by chi-squared test without considering its direction. We then combined these two test statistics to achieve the overall p-value, which tests the overall difference between R and NR for each species.

Meta-analysis

Meta-analysis is a two-stage approach¹⁵. In the initial stage data are analyzed separately for each study. In the second-stage test results for each study are aggregated to obtain summary data across all studies¹⁵. On the other hand, mega-analysis is a one-stage approach that pools and analyzes raw data from a number of studies to estimate the overall effect¹⁵. Both approaches have their strengths and limitations¹⁵. We conducted a meta-analysis by combining the test results for each of the four data sets in Table 1. Logistic regression and linear regression estimates are only accurate if the zero proportion in each species is sufficiently low. Accordingly, species were excluded if their prevalence was lower than 20%. As species prevalence differs by study, some species were filtered out only in a subset of the data sets. Assuming the independence of each data set, the overall meta-analysis p-values were computed for the combined test statistic, which is the sum of the test statistics from the four data sets.

Mega-analysis

For each species included in the meta-analysis, we also conducted a mega-analysis. The mega-analysis simply combined all samples from all four data sets, and two-part log-normal models were implemented similarly. We also calculated the CIs for all estimates and overall chi-squared test p-values combining species prevalence and abundance, which were then compared with the meta-analysis results.

Results

Characteristics of cohorts included in analyses

In this mega- and meta-analysis, we included three studies that assessed fecal metagenomics of R and NR to ICI in patients with varying tumor types (Table 1). Data from Routy *et al.*, was analyzed as subsets based on tumor type RCC vs advanced NSCLC for a total of 4 analyzed cohorts. DNA extraction, sequencing platform and sequencing depth varied by study and are summarized in Table S1. A total of 190 patients (n=103 R; n=87 NR) were included from the three studies.

A total of 469 species were identified across all four data sets of which 167 species met our inclusion criteria of being present in >20% of samples and were included in the two-part log-normal analysis (Table S2). A total of 34 species were differentially abundant based on response in at least one dataset or mega- and meta-analyses (Table S2).

114 **Table 1:** Characteristics of cohorts included in meta-analysis and corresponding taxa associated with response to ICI using
 115 metagenomic analyses.
 116

Patient population	N (= total number of patients)	Responders	non-Responders	Organisms enriched in RECIST responders	Organism Enriched in RECIST non-responders	Reference
Metastatic melanoma	N=25	N=14	N=11	<i>Faecalibacterium</i> spp.	<i>Bacteroides thetaiotaomicron</i> , <i>Escherichia coli</i> , <i>Anaerotruncus colihominis</i>	Gopalakrishnan <i>et al.</i> 2018
Metastatic melanoma	N=38	N=14	N=24	<i>Enterococcus faecalis</i> , <i>E. coli</i> , <i>Escherichia</i> spp., <i>Bacteroides ovatus</i> , <i>Turicibacter sanguinis</i> , <i>Clostridium nexile</i> , <i>Enterococcus faecium</i> , <i>Collinsella aerofaciens</i> , <i>Bifidobacterium adolescentis</i> , <i>Klebsiella oxytoca</i> , <i>Veillonella parvula</i> , <i>Parabacteroides merdae</i> , <i>Lactobacillus</i> species, and <i>Bifidobacterium longum</i> , <i>Campylobacter gracilis</i>	<i>Burkholderiales</i> <i>bacterium</i> 1147, <i>Holdemania filiformis</i> , <i>Coproccoccus comes</i>	Matson <i>et al.</i> 2018
Advanced NSCLC	N=65	N=33	N=32	<i>Akkermansia muciniphila</i> , <i>Ruminococcaceae</i> , <i>Faecalibacterium</i> , <i>Alistipes</i> spp., <i>Eubacterium</i> species, <i>Firmicutes</i> , <i>Cloacibacilius porcorum</i> , <i>Enterococcus faecium</i> , <i>Clostridiales</i> , <i>Firmicutes</i> , <i>Prevotella</i> ,	<i>Prevotella</i> spp., <i>Clostridiales</i> , <i>Blautia</i> , <i>Bacteroides calrus</i> , <i>Proteobacteria</i> , <i>Bacteroides nordii</i> , <i>Parabacteroides distasonis</i>	Routy <i>et al.</i> 2018
RCC	N=62	N=42	N=20			

Species consistently enriched or depleted in responders compared to non-responders across data sets

The primary goal for our study was to determine species consistently enriched or depleted in R compared to NR across data sets. While each data set had unique signatures of differentially abundant species between R/NR, no species were identified that were statistically significantly associated with response across all four data sets. A number of species were significantly differentially enriched/depleted in R/NR per data set; Matson et al., (n=7 species), Routy et al., (RCC) (n=6), Routy et al., (NSCLC) (n=10), Gopalakrishnan et al., (n=9) (Figures S1 and S2 and Table S2). A total of three species were significantly enriched or depleted in R in two data sets: *Clostridiaceae bacterium JC118* was enriched in R in the Routy et al., (NSCLC) and Matson et al., data sets; *Bacteroides thetaiotaomicron* RA was significantly depleted in R in both the Routy et al., (RCC) and Gopalakrishnan et al., data sets; and an increase in *Lachnospiraceae bacterium 5163FAA* abundance was associated with response in both the Gopalakrishnan et al., and Routy et al., (NSCLC) data sets. While not significantly different, *Streptococcus australis* trended towards a higher relative abundance in R across all cohorts. In contrast, *B. thetaiotaomicron* trended towards lower relative abundance in R across all data sets. In addition, we were interested in assessing species that are consistently enriched or depleted in R across data sets based on prevalence (Figure S2 and Table S2). *Clostridium bolteae*, *Escherichia coli*, *Flavonifractor plautii*, *Ruminococcus lactaris* and *Streptococcus australis* were consistently less prevalent in R. In contrast, *Bacteroides caccae*, *Barnesiella intestinihominis* and *Lachnospiraceae bacterium 8157FAA* were more prevalent in R. Only one species had statistically significant but opposing associations between cohorts; *Ruminococcus gnavus* was more abundant in R in the Routy et al., (RCC) cohort and less abundant in the cohort in the Gopalakrishnan et al., study.

Meta-analysis reveals species consistently associated with response to immunotherapy

A summary of a metagenomic meta-analysis of species level ICI response associations is shown in Figure 1. A total of thirteen species were significantly differentially abundant between R and NR in the meta-analysis. Of these thirteen species, twelve were identified as significantly different between R and NR in at least one data set on its own. *Clostridium hathewayi* was the only species identified uniquely in the meta-analysis. Five species which were significantly associated with R/NR in one dataset demonstrated a non-significant trend in the meta-analysis: *Coprobacillus spp*, *Parabacteroides spp.*, Lachnospiraceae bacterium 8157FAA, *Eubacterium ramulus*, and *Clostridium symbiosum*. Lastly, 16 species were not statistically significantly different nor trended towards response using the meta-analysis despite being significantly different in at least one dataset.

144 **Figure 1:** Taxa that are significantly different between responders and non-responders in at least one dataset and or mega/meta-
145 analyses using a two-part log-normal analysis accounting for both taxa abundance and prevalence.
146

Mega-analysis assessing differences in responders vs non-responders across all data sets

Mega-analysis allowed us to determine the effect size of the differences between R and NR for both prevalence and relative abundance of species across all studies (Figures S1 and S2, and Table S2). In the mega-analysis, thirteen species were statistically differentially abundant between R and NR (Figure 1). Of these thirteen species, eight were associated with response in at least one dataset. Five species were uniquely associated with response using the mega-analysis including *Anaerofustis stercorihominis*, *Flavonifractor plautti*, *Ruminococcus obeum*, *Rothia mucilaginosa* and *Barnesiella intestinitominis*. Of the 34 species associated with response in at least one data set or the meta-analysis, 19 species were not associated with response in the mega-analysis.

Identification of species consistently associated with response in meta- and mega-analyses and analysis of sensitivity and specificity to predict response

We next identified species that were concordantly associated with response in both the meta and mega-analyses. Using this criterion five species were identified including *B. thetaiotaomicron*, *Clostridium bolteae*, *H. filiformis*, *Clostridiaceae bacterium JC118* and *E. coli*. We tested the sensitivity and specificity of the relative abundance of these five organisms and alpha-diversity measures to predict response to ICI using a ROCAUC analyses (Table 2). Our findings demonstrate inter-study differences in the ability to discriminate R vs NR based on species RA. No species or alpha-diversity metric consistently demonstrated a significant AUC across studies. However, each dataset had at least two species or alpha-diversity measures with AUC's >0.60. In the Matson et al., dataset *H. filiformis* RA significantly discriminated between R/NR with an AUC of 0.72 [95% CI 0.55-0.88]. *C. bolteae* RA had a significant AUC of 0.73 [95% CI 0.60-0.87] in the Routy et al., (RCC) dataset suggesting that a higher RA of this species discriminates between

164 R and NR. In the Gopalakrishnan et al., dataset both *E. coli* and *B. thetaiotaomicron* RA had significant sensitivity and specificity to
165 detect response. When all four cohorts were combined sensitivity and specificity of predicting response were significant for the RA of
166 both *C. bolteae* and *B. thetaiotaomicron*. SDI did not discriminate between R/NR.

Table 2: Receiver operator characteristic curve area under the curve (AUC) analysis between responders and non-responders for core taxa and alpha-diversity measures.

Study				
Matson <i>et al.</i> , (n=38 samples)	AUC	SE [^]	95% CI ⁺	P-value
SDI [†]	0.52	0.10	0.33-0.72	0.81
SIM [‡]	0.55	0.10	0.35-0.74	0.63
<i>Escherichia coli</i>	0.64	0.10	0.44-0.84	0.15
<i>Clostridiaceae bacterium JC118</i>	0.63	0.10	0.43-0.82	0.19
<i>Holdemania filiformis</i>	0.72	0.085	0.55-0.88	0.027
<i>Clostridium bolteae</i>	0.57	0.097	0.39-0.76	0.45
<i>Bacteroides thetaiotaomicron</i>	0.61	0.092	0.43-0.79	0.25
Routy <i>et al.</i> , (RCC, n=62)				
SDI	0.51	0.080	0.35-0.67	0.88
SIM	0.53	0.079	0.37-0.68	0.74
<i>Escherichia coli</i>	0.54	0.075	0.40-0.69	0.58
<i>Clostridiaceae bacterium JC118</i>	0.63	0.077	0.47-0.78	0.11
<i>Holdemania filiformis</i>	0.53	0.086	0.36-0.69	0.76
<i>Clostridium bolteae</i>	0.73	0.070	0.60-0.87	0.0030
<i>Bacteroides thetaiotaomicron</i>	0.55	0.090	0.37-0.72	0.55
Routy <i>et al.</i> , (NSCLC, n=65)				
SDI	0.62	0.071	0.48-0.76	0.096
SIM	0.61	0.071	0.47-0.75	0.13
<i>Escherichia coli</i>	0.56	0.072	0.42-0.71	0.37
<i>Clostridiaceae bacterium JC118</i>	0.58	0.071	0.44-0.72	0.28
<i>Holdemania filiformis</i>	0.50	0.072	0.36-0.64	0.97
<i>Clostridium bolteae</i>	0.52	0.073	0.37-0.66	0.81
<i>Bacteroides thetaiotaomicron</i>	0.57	0.072	0.43-0.71	0.33

Gopalakrishnan et al., (n=25)				
SDI	0.55	0.13	0.30-0.79	0.70
SIM	0.64	0.12	0.41-0.88	0.25
<i>Escherichia coli</i>	0.73	0.11	0.52-0.93	0.055
<i>Clostridiaceae bacterium JC118</i>	0.50	0.12	0.27-0.74	0.98
<i>Holdemanian filiformis</i>	0.55	0.13	0.31-0.81	0.64
<i>Clostridium bolteae</i>	0.63	0.12	0.40-0.86	0.27
<i>Bacteroides thetaiotaomicron</i>	0.76	0.10	0.55-0.96	0.029
Mega-analysis (n=190)				
SDI	0.56	0.042	0.47-0.64	0.18
SIM	0.55	0.042	0.46-0.63	0.28
<i>Escherichia coli</i>	0.53	0.042	0.45-0.62	0.44
<i>Clostridiaceae bacterium JC118</i>	0.54	0.042	0.46-0.62	0.33
<i>Holdemanian filiformis</i>	0.54	0.043	0.46-0.63	0.27
<i>Clostridium bolteae</i>	0.61	0.041	0.52-0.69	0.013
<i>Bacteroides thetaiotaomicron</i>	0.59	0.043	0.51-0.68	0.027

† Shannon alpha-diversity index

‡ Simpson alpha-diversity index

Δ Standard Error

+ Confidence Interval

Discussion

In our study, we leveraged the additional statistical power of mega and meta-analyses and taxonomic resolution of metagenomic data to identify species consistently associated with R or NR across three studies (four cohorts) in which gut microbial community composition has been previously associated with response to ICI³⁻⁵. We conducted a two-part log-normal analysis that accounts for the skewness and non-Gaussian distribution of microbiome data^{14,16-18}. Compared to simply assessing changes in relative abundance of species, this approach has the advantage of accounting for both prevalence and relative abundance data^{14,16-18}. After uniform bioinformatic analysis, no species were identified that had a significant association with response across individual studies, but five were identified in combined analysis. Importantly, these associations were identified in spite of the inter-cohort heterogeneity in tumor type, treatment regimens, geographic location and sequencing methods, suggesting that they may be more generalizable. Our study confirms findings of some previously identified R/NR associated species^{2-5,19}. Notably, a higher abundance of *Bacteroides* has been associated with lack of response whereas a higher abundance of *Firmicutes* has been associated with response to ICI. Specifically, we observed R had a higher abundance of *Clostridiaceae bacterium JC118* and *E. coli* and a lower abundance of *B. thetaiotaomicron*, *H. filiformis* and *C. bolteae* compared to NR. In addition, we demonstrate that of the five organisms that were concordantly associated with response in the meta and mega-analyses, a higher relative abundance of *B. thetaiotaomicron* and *C. bolteae* were predictive of NR using a ROCAUC analyses. These data suggest that strategies that deplete non-response associated microbes (such as specific antibiotics or bacteriophages) may be viable therapeutic approaches. This is in contrasts to “pro-microbial”

191 approaches currently being investigated in many clinical trials such as the use of probiotics, fecal microbiome transplants and stool
 192 substituents to augment response to ICI²⁰.

193 We observed a number of discrepancies to previously published data. Of interest, *A. muciniphila*, *B. longum* and *F. prausnitzii* were
 194 associated with response in the original studies however we did not detect differences between R and NR in our analyses. While
 195 limitations in statistical power based on sample size may partly explain this, inter-cohort variability in treatment regimens and the
 196 biology of host-treatment-microbe interactions may be important additional factors. As the mechanisms conferring microbe-induced
 197 ICI-responsiveness are being elucidated, it is clear that some will be tumor-type, host or even tumor-specific, such as the molecular
 198 mimicry of tumor antigens in gut microbes²¹, which will only confer microbe-response associations in a defined number of patients
 199 with specific tumors. Other factors, such as diet-microbe-metabolite-immune interactions may also drive tumor, host or population-
 200 specific associations which do not generalize⁶.

201 We acknowledge a number of limitations to our study. Firstly, differences in sample collection, DNA extraction methods and
 202 sequencing platforms from each study may bias findings- making it challenging to detect universal signals associated with response to
 203 ICI. Implementing standardized methods for microbiome studies will reduce experimental and technical biases and improve our
 204 ability to detect differences between R and NR. Second, patient heterogeneity in disease stage, diet, sex, treatment, co-morbidities
 205 limit the ability of meta or mega-analytical approaches to identify associations that may exist in only defined subsets of patients. Our
 206 sample size of 190 metagenomic samples, while as large as some cohorts analyzed by lower taxonomic resolution methods such as

16S rRNA sequencing, was still limited. Additional multi-centre prospective studies are required to validate the associations identified in this analysis and their diagnostic or therapeutic utility.

Conclusion

Our study confirms previous findings suggesting that there are differences in the gut microbiome of R vs NR to ICI. Despite consistent bioinformatic analyses no species were found to be consistently differentially abundant between R/NR across data sets. However, using meta- and mega-analyses we identified five species that were concordantly differentially abundant between R and NR. Of these five organisms, *B. thetaiotaomicron* and *C. bolteae* were predictors of NR to ICI. These data suggest future clinical trials should assess the use of narrow spectrum anti-biotics targeting NR associated species.

Acknowledgments

The authors acknowledge and are grateful for the support of the Tomczyk AI and Microbiome Working Group. AH was supported by the Tomczyk AI and Microbiome Working Group and the Princess Margaret Cancer Foundation.

Conflicts of Interest of Statement

AAH, BC, PS, MW, VR, WX, BAC, have no conflicts to report. AS has received compensation from: Merck, Bristol-Myers Squibb, Novartis, Oncorus, Janssen; and grant/research support from: Novartis, Bristol-Myers Squibb, Symphogen AstraZeneca/Medimmune, Merck, Bayer, Surface Oncology, Northern Biologics, Janssen Oncology/Johnson & Johnson, Roche, Regeneron, Alkermes, Array Biopharma/Pfizer, GSK.

Literature cited

1. Vétizou, M. *et al.* Anticancer immunotherapy by CTLA-4 blockade relies on the gut microbiota. *Science* (80-.). **350**, 1079–1084 (2015).
2. Frankel, A. E. *et al.* Metagenomic Shotgun Sequencing and Unbiased Metabolomic Profiling Identify Specific Human Gut Microbiota and Metabolites Associated with Immune Checkpoint Therapy Efficacy in Melanoma Patients. *Neoplasia (United States)* **19**, (2017).
3. Matson, V. *et al.* The commensal microbiome is associated with anti-PD-1 efficacy in metastatic melanoma patients. *Science* (80-.). **359**, 104–108 (2018).
4. Gopalakrishnan, V. *et al.* Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* (80-.). **359**, 97–103 (2018).
5. Routy, B. *et al.* Gut microbiome influences efficacy of PD-1–based immunotherapy against epithelial tumors. *Science* (80-.). **359**, 91–97 (2018).
6. Mager, L. F. *et al.* Microbiome-derived inosine modulates response to checkpoint inhibitor immunotherapy. *Science* (80-.). **369**, (2020).
7. Schwartz, L. H. *et al.* RECIST 1.1 - Update and clarification: From the RECIST committee. *Eur. J. Cancer* **62**, 132–137 (2016).
8. Andrews, S. FASTQC A Quality Control tool for High Throughput Sequence Data. *Babraham Inst.* (2015).
9. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, (2016).
10. Segata, N. *et al.* Metagenomic microbial community profiling using unique clade-specific marker genes. *Nat. Methods* **9**, (2012).
11. McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).
12. Wickham H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New York* (2009). Available at: <https://cran.r-project.org/web/packages/ggplot2/citation.html>.
13. R Core Team. R: a language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria.* (2013).
14. Chen, B. & Xu, W. Generalized estimating equation modeling on correlated microbiome sequencing data with longitudinal measures. *PLoS Comput. Biol.* **16**, e1008108 (2020).
15. Burke, D. L., Ensor, J. & Riley, R. D. Meta-analysis using individual participant data: one-stage and two-stage approaches, and why they may differ. *Stat. Med.* **36**, (2017).
16. Wirbel, J. *et al.* Meta-analysis of fecal metagenomes reveals global microbial signatures that are specific for colorectal cancer. *Nat. Med.* **25**, (2019).
17. Ho, N. T., Li, F., Wang, S. & Kuhn, L. MetamicrobiomeR: An R package for analysis of microbiome relative abundance data using zero-inflated beta GAMLSS and meta-analysis across studies using random effects models. *BMC Bioinformatics* **20**, (2019).
18. Wagner, B. D., Robertson, C. E. & Harris, J. K. Application of two-part statistics for comparison of sequence variant counts. *PLoS One* **6**, (2011).
19. Derosa, L. *et al.* Gut Bacteria Composition Drives Primary Resistance to Cancer Immunotherapy in Renal Cell Carcinoma Patients. *Eur. Urol.* **78**, (2020).
20. Araujo, D. V *et al.* Bugs as Drugs: The Role of Microbiome in Cancer Focusing on Immunotherapeutics. *Cancer Treat. Rev.* (2020).
21. Fluckiger, A. *et al.* Cross-reactivity between tumor MHC class I–restricted antigens and an enterococcal bacteriophage. *Science* (80-.). **369**, (2020).

- 270 22. Suau, A. *et al.* Direct analysis of genes encoding 16S rRNA from complex communities
271 reveals many novel molecular species within the human gut. *Appl. Environ. Microbiol.* **65**,
272 (1999).
- 273 23. Godon, J. J., Zumstein, E., Dabert, P., Habouzit, F. & Moletta, R. Molecular microbial
274 diversity of an anaerobic digester as determined by small-subunit rDNA sequence
275 analysis. *Appl. Environ. Microbiol.* **63**, (1997).
276