Smoothened and ARL13B are critical in mouse for superior cerebellar peduncle targeting

Sarah K. Suci^{*,†,‡} Alyssa B. Long^{†,‡} and Tamara Caspary[†]

^{*}Genetics and Molecular Biology Graduate Program, [†]Department of Human Genetics,
Emory University, Atlanta, GA 30322, [‡]These authors contributed equally

ORCID ID:

Suci: 0000-0002-2416-9519

Long: 0000-0002-4467-4213

Caspary: 0000-0002-6579-7589

Summary statement

Joubert syndrome is diagnosed by the hindbrain “molar tooth sign” malformation. Using mouse models, we show loss of the ciliary GTPase ARL13B, mutations in which lead to Joubert syndrome, result in two features of the molar tooth sign: hypoplasia of the cerebellar vermis and inappropriate targeting of the superior cerebellar peduncles. Furthermore, we demonstrate that loss of vertebrate Hedgehog signaling may be the underlying disrupted mechanism as we extend its role in axon guidance to the superior cerebellar peduncles.

Running title (~35 characters, including spaces):

SMO and ARL13B in SCP targeting

Key words/phrases (up to ten):

molar tooth sign, Joubert syndrome, Smo, ARL13B, SCP targeting, cerebellar vermis

Corresponding author:

Tamara Caspary

Department of Human Genetics

615 Michael Street, Suite 301

Atlanta, GA 30322

404-727-9862

tcaspar@emory.edu

Abstract

Patients with the ciliopathy Joubert syndrome present with physical anomalies, intellectual disability, and a hindbrain malformation described as the “molar tooth sign” due to its appearance on an MRI. This radiological abnormality results from a combination of hypoplasia of the cerebellar vermis and inappropriate targeting of the white matter tracts of the superior cerebellar peduncles. *ARL13B* is a cilia-enriched regulatory GTPase established to regulate cell fate, cell proliferation and axon guidance through vertebrate Hedgehog signaling. In patients, mutations in *ARL13B* cause Joubert syndrome. In order to understand the etiology of the molar tooth sign, we used mouse models to investigate the role of *ARL13B* during cerebellar development. We found *ARL13B* regulates superior cerebellar peduncle targeting and these fiber tracts require Hedgehog signaling for proper guidance. However, in mouse the Joubert-causing R79Q mutation in *ARL13B* does not disrupt Hedgehog signaling nor does it impact tract targeting. We found a small cerebellar vermis in mice lacking *ARL13B* function but no cerebellar vermis hypoplasia in mice expressing the Joubert-causing R79Q mutation. Additionally, mice expressing a cilia-excluded variant of *ARL13B* that transduces Hedgehog normally, showed normal tract targeting and vermis width. Taken together, our data indicate that *ARL13B* is critical for control of cerebellar vermis width as well as superior cerebellar peduncle axon guidance, likely via Hedgehog signaling. Thus, our work highlights the complexity of *ARL13B* in molar tooth sign etiology.

INTRODUCTION

Joubert Syndrome and Related Disorders (JSRD) are recessive congenital disorders with a variety of symptoms including developmental delay, intellectual disability, abnormal respiratory rhythms, ataxia, oculomotor apraxia, polydactyly, craniofacial defects, retinal dystrophy, nephronophthisis, and hepatic fibrosis (Parisi *et al.* 2007, Bachmann-Gagescu *et al.* 2020). While the exact prevalence of JSRD is not known, published statistics range from 1:80,000 to 1:100,000, but these may be underestimates as suggested by a recent study (Brancati *et al.* 2010, Nuovo *et al.* 2020). The characteristic neuroanatomical feature of JSRD is the molar tooth sign (MTS), which is caused by hypoplasia of the cerebellar vermis and thickened, elongated superior cerebellar peduncles (SCPs) that fail to decussate (Maria *et al.* 1997; Yachnis and Rorke 1999; Poretti *et al.* 2007). However, little is known about the mechanistic etiology of this hindbrain malformation. This is especially significant as several symptoms of JSRD arise from defects in the hindbrain: cerebellar dysfunction commonly causes ataxia and hypotonia, while some patients manifest life-threatening breathing problems (Parisi 2019).

To date, mutations in over 35 genes cause JSRD, and their associated proteins almost always localize to the primary cilium or the centrosome (Parisi 2019). Thus, JSRD is classified as a ciliopathy, a category of human disease stemming from ciliary dysfunction. One of the genes implicated in JSRD is *ARL13B*, which encodes a regulatory GTPase highly enriched in cilia (Cantagrel *et al.* 2008; Bachmann-Gagescu *et al.* 2015; Thomas *et al.* 2015; Shaheen *et al.* 2016; Rafiullah *et al.* 2017). As a GTPase, *ARL13B* is expected to have multiple effector proteins which interact with

specific ARL13B residues. ARL13B can function as a guanine exchange factor (GEF) for ARL3, mutations in which also lead to JSRD (Gotthardt *et al.* 2015; Ivanova *et al.* 2017). JSRD-causing mutations in either ARL3 or ARL13B can disrupt their interaction or ARL13B's GEF activity, consistent with the notion that specific ARL13B function is affected by JS-causing point mutations (Gotthardt *et al.* 2015; Ivanova *et al.* 2017; Alkanderi *et al.* 2018). Most JSRD-causing ARL13B mutations cluster within the protein's GTPase domain, although two are located in the coiled coil domains in the C terminal half of the protein (Cantagrel *et al.* 2008; Bachmann-Gagescu *et al.* 2015; Thomas *et al.* 2015; Shaheen *et al.* 2016; Rafiullah *et al.* 2017). ARL13B complexes with the inositol phosphatase INPP5E, which is also implicated in causing JSRD (Bielas *et al.* 2009; Humbert *et al.* 2012). ARL13B is critical for targeting INPP5E to cilia and JSRD-causing ARL13B mutations disrupt INPP5E ciliary targeting (Humbert *et al.* 2012). INPP5E controls ciliary lipid composition through its phosphatase activity and most JSRD-causing mutations are within its phosphatase domain (Bielas *et al.* 2009; Chavez *et al.* 2015; Garcia-Gonzalo *et al.* 2015). Other proteins implicated in JSRD also affect ciliary targeting with many functioning at the transition zone, supporting the notion that abnormal ciliary traffic leading to defective signaling underlies JSRD (Arts *et al.* 2007; Delous *et al.* 2007; Garcia-Gonzalo *et al.* 2011; Hopp *et al.* 2011; Srour *et al.* 2012; Roberson *et al.* 2015).

The mechanistic connection between the cilia-related proteins implicated in JSRD and the MTS are elusive in part because distinct biological processes are at play. Abnormal proliferation may underlie the hypoplastic cerebellar vermis whereas defective axonal targeting is likely involved in the abnormal SCP tracts. One signaling pathway

potentially linked to both processes is vertebrate Hedgehog (Hh) which relies on cilia (Huangfu *et al.* 2003). Sonic hedgehog (Shh) is a mitogenic cue that controls proliferation in the developing cerebellum so its misregulation could underlie the cerebellar hypoplasia (Dahmane and Ruiz i Altaba 1999; Wechsler-Reya and Scott 1999; Kenney and Rowitch 2000). While the SCP tracts that normally project from the deep cerebellar nuclei to the contralateral thalamus are guided by unknown signals, Shh is a known commissural axon guidance cue (Charron *et al.* 2003). JSRD patients can also display axon guidance defects in decussation of the pyramidal tracts (Yachnis and Rorke 1999; Poretti *et al.* 1997).

ARL13B and *INPP5E*, encoding ciliary proteins linked to JSRD, are known to regulate vertebrate Hh signaling. In mouse models, *ARL13B* loss disrupts cell fate specification in the neural tube, proliferation of the cerebellar granule precursor cells in the cerebellum and Shh-directed guidance of commissural axons in the spinal cord (Caspary *et al.* 2007; Bay *et al.* 2018; Ferent *et al.* 2019). These data support a model whereby disruption of Shh signaling by *ARL13B* mutation could provide a single mechanism underlying the MTS. This model is bolstered by the fact that additional phenotypes exhibited by JSRD patients, such as craniofacial defects or polydactyly, can arise from aberrant Hh signaling (Valente *et al.* 2008; Lan and Jiang 2009; Lipinski *et al.* 2010).

As attractive as a Hh-based model for JSRD may be, not all the data support that JSRD phenotypes result from misregulation of Hh signaling. Some features of JSRD, such as the renal and liver anomalies, are not clearly due to misregulation of Hh signaling (Doherty 2009; Breslow *et al.* 2018). Additionally, of over 35 genes implicated

in JSRD, only 22 play some role in Hh pathway regulation (Davey *et al.* 2006; Reiter and Skarnes 2006; Caspary *et al.* 2007; Vierkotten *et al.* 2007; Huang *et al.* 2009; Weatherbee *et al.* 2009; Bimonte *et al.* 2011; Dowdle *et al.* 2011; Sang *et al.* 2011; Chih *et al.* 2011; Christopher *et al.* 2012; Thomas *et al.* 2012; Abdelhamed *et al.* 2013; Hynes *et al.* 2014; Wu *et al.* 2014; Chavez *et al.* 2015; Garcia-Gonzalo *et al.* 2015; Asadollahi *et al.* 2018; Frikstad *et al.* 2019; Munoz-Estrada and Ferland 2019). Some of these links are tenuous. For example, mouse *Arl3* mutants mislocalize the Hh transcription factor, GLI3, in their cilia yet do not exhibit any of the phenotypes normally displayed by mutants in the Hh pathway (Schrack *et al.* 2006, Lai *et al.* 2011; Schwarz *et al.* 2017). Additional signaling pathways are linked to cilia including others known to be important in cell proliferation and axon guidance. Loss of either of the JSRD-linked genes *Ahi1* or *Cep290* in mouse leads to a small cerebellar vermis due to aberrant Wnt signaling (Lancaster *et al.* 2011; Ramsbottom *et al.* 2020). Possible JSRD-causing mutations in *Znf423* are associated with defects in Wnt, BMP and retinoic acid signaling (Hata *et al.* 2000, Huang *et al.* 2009, Casoni *et al.* 2020, Deshpande *et al.* 2020). Conditional *Arl13b* or *Inpp5e* deletion in the SCPs results in their disorganization and thickening through misregulation of ciliary PI3 kinase and AKT (Guo *et al.* 2019).

Ciliopathies are well established to be genetically complex. JSRD patients with different ARL13B mutations can display distinct phenotypes, such as obesity in an individual expressing ARL13B^{Y86C} and occipital encephalocele in a patient expressing ARL13B^{R79Q} (Cantagrel *et al.* 2008; Thomas *et al.* 2015). This is further exemplified in cases of related individuals carrying the same mutation but exhibiting different phenotypes and even diagnoses. For example, JSRD-causing mutations in TMEM67

(R208X) and TMEM216 (R73H) can also cause the more severe disease Meckel syndrome (Consugar *et al.* 2007; Otto *et al.* 2009; Valente *et al.* 2010). Understanding the genetic modifiers and environmental contribution underlying the phenotypic variation will be key to understanding disease etiology as will understanding when and how relevant pathways interact. In mouse models of the JSRD- and Meckel Syndrome-linked gene *Tmem67*, two phenotypic categories emerged: one with cerebellar malformations resembling JSRD and another with more severe CNS defects reminiscent of Meckel syndrome (Abdelhamed *et al.* 2013). The two categories correlated with whether cilia were retained, with the severe Meckel-like phenotype observed in animals lacking cilia. Furthermore, the two phenotypic groups impacted Hh and Wnt signaling differently, pointing to both pathways being critical (Abdelhamed *et al.* 2019). Importantly, it is not simply whether cilia are present, as another JSRD mouse model, *Talpid3*, lacks cilia yet displays a JSRD-like small cerebellar vermis (Bashford and Subramanian 2019). Thus, mouse models are incredibly informative yet point to the enormous complexity underlying the MTS.

Here we investigate the role of ARL13B in relation to Hh signaling in two major features of the MTS: targeting of the SCPs to the thalamus and hypoplasia of the cerebellar vermis. We explore these processes using a series of mouse alleles through which we first define the roles of Hh signaling and ARL13B in SCP projections. Subsequently, we untangle the role of ARL13B from within and outside of the cilium and investigate a JS-causing patient allele. Taken together, our data illuminate the roles of ARL13B in MTS etiology and the complexity in modeling aspects of the MTS in mouse.

MATERIALS & METHODS

Mouse lines

All mice were cared for in accordance with NIH guidelines and Emory University's Institutional Animal Care and Use Committee (IACUC). Lines used were *Nex-Cre* (C3H/HeJ-*Neurod6*^{tm1(cre)Kan}) [MGI:2668659], *Brn4-Cre* (C3H/HeJ-*Tg(Pou3f4-cre)32Cren*) [MGI:2158470], *Smo*^{flox} (C3H/HeJ-*Smo*^{tm2AMC}) [MGI:2176256], *Arl13b*^{flox} (C3H/HeJ-*Arl13b*^{tm1Tc}) [MGI:4948239], *Arl13b*^{V358A} (C57BL/6J-*Arl13b*^{em1Tc}) [MGI:6256969], and *Arl13b*^{R79Q} (C57BL/6J-*Arl13b*^{em2Tc}) [MGI:6279301]. Note that *Arl13b*^Δ is the deletion allele resulting from germline deletion of the conditional *Arl13b*^{flox} allele. Genotyping was as previously described (Heydemann *et al.* 2001; Goebbels *et al.* 2006; Nolan-Stevaux *et al.* 2009; Su *et al.* 2012; Gigante *et al.* 2020).

To generate the R79Q mutation in *Arl13b*, a CRISPR gRNA (ATTATTATGCTGAATCCTATGG; PAM sequence is italicized) targeting exon 3 of the *Arl13b* locus along with a 180bp donor oligo (5'-CTCCCACTGTTGGCTTTTCTAAAA-TTGATCTGAGACAAGGAAAGTTCCAAGTTACCATCTTTGACTTAGGAGGTGGAAAAAGAATTCAGGGCATATGGAAGAATTATTATGCTGAATCCTATGGGGTAATATTTGTTGTGGATTCCAGTGATGAGGAGAGAATGGAAGAAACAAAGGAGA-3'; underlined bases are engineered) were designed to generate a G-to-A change creating the R79Q point mutation as well as A-to-C and T-to-A silent changes to create a *NdeI* restriction site that could be used for genotyping (Millipore Sigma). The gRNA (50 ng/ul), ssDNA oligo donor (50 ng/ul) and wild type *Cas9* mRNA (10 ng/ul) were injected into 1-cell C57BL/6J zygotes and subsequently transplanted at the 2-cell stage into C57BL/6J pseudopregnant females by the Emory Transgenic and Gene Targeting Core; all

reagents were purchased from Millipore Sigma. Genomic DNA from toes was amplified via PCR using primers (5'-TCACTTGCAACAGAGCATCC-3') and (5'-ACAGCTCTGCCCCGTGTTTAC-3') located upstream and downstream of the donor oligo breakpoints; products were sequenced with the forward (first) primer. A single founder animal heterozygous for both the R79Q mutation and NdeI restriction site was identified with no additional editing. Subsequent allele-specific genotyping of progeny was performed on ear punch or yolk sac using the following primers: Fwd-wt primer: 5'-GGAGGTGGAAAAAGAATaCg-3'; Fwd-mut primer: 5'-gctctatggctgGGAGGTGGAAAAAGAATTga-3'; Rev primer: 5'-AGTGCTAAGACACCCGAGGA-3'. PCR bands at 142bp (wild type) and/or 154bp (mutant) were produced, due to the addition of 12 non-templated bases to the 5' end of the Fwd-mut primer (lowercase). Note the 3' ends of the two forward primers differed in the base that codes for the R to Q change (final nucleotide of the primer) and includes a "wobble" base (lowercase) to provide allele-specific amplification after the first round of PCR (Gaudet *et al.* 2009). In order to breed away any potential off-target edits, the founder was backcrossed to C57BL/6J for three generations with at least two independent meiotic opportunities for recombination in each generation.

Tract tracing injections and analysis

Tract tracing experiments were performed according to a protocol approved by Emory University's Institutional Animal Care and Use Committee (IACUC). Male and female mice at postnatal day 90 or older were used for tract tracing experiments. At least 3 mice of each genotypic group were analyzed in experiments (exact N included in

Figures 1-3, S1). Mice were anesthetized with inhaled isoflurane and maintained under anesthesia throughout the procedure. Animals were secured in the stereotax, and the scalp was opened with bregma and lambda aligned to flat skull position. Dorsal thalamus injections were targeted to the bregma (AP:-0.70, ML:+1.13, DV:-3.28, Angle:0°) and ventral thalamus injections were targeted using coordinates to the bregma (AP:-0.70, ML:-3.11, DV:-4.69, Angle:25°). Ventral injection targeting includes a 25° angle to avoid pulling dye through the dorsal thalamus upon needle removal. Then, a 5 ul Hamilton microsyringe was lowered to target and target was injected with lysine fixable dextran tetramethylrhodamine neuroanatomical tracer (fluoro-Ruby, 10,000 MW, ThermoFisher Scientific D1817). Animals received 0.05 - 0.5 ul injections of 10% dextran tetramethylrhodamine in sterile phosphate buffered saline (PBS, pH = 7.25) unilaterally at a rate of 0.1 ul/minute. The 0.5 ul injection volume initially used resulted in high background (non-specific) fluorescence. Later surgeries were conducted with 0.05 ul of dye, which resulted in lower background and retention of a strong signal in the DCN.

Seven days post-procedure, mice were perfused with 50mls of PBS followed by 30mls of 4% paraformaldehyde (PFA). Brains were kept in PFA overnight and subsequently placed in 30% sucrose in 0.1M phosphate buffer (pH 7.3) for cryoprotection for at least 48 hours. Brains were embedded in Tissue-Tek OCT compound (Sakura) for coronal cryostat sectioning. Sections were 60 microns thick and processed through 70% ethanol dehydration and 0.1% sudan black autofluorescence quencher, rehydrated in PBS and treated with DAPI to stain nuclei prior to fluorescence imaging. Images were taken on a Lionheart FX automated microscope (Biotek) or at 5x magnification on a Leica DM6000B microscope (Leica) using SimplePCI imaging

software (Hamamatsu). The Leica images were subsequently stitched together to reveal the entirety of the brain section in Fiji (Schindelin *et al.* 2012) or Photoshop (Adobe). Surgical injection sites were assessed to ensure dye was present at the desired injection site. If the injection was off-target or dye at the injection site was not seen, samples were removed from analysis. Cerebellar images from injections were evaluated for DCN staining with assessor blinded to genotype. The number of injections that resulted in fluorescent DCN for each injection site (dorsal and ventral thalamus) were compared between mutant and control genotypes using a two-sided Fisher's exact test (PRISM 8.2.0).

Phenotypic analysis of embryos

Timed matings of mice were performed to generate somite-matched embryos at embryonic day 10.5 (E10.5). Embryos were dissected in cold PBS and processed for immunofluorescence staining as previously described (Constable *et al.* 2020). Primary antibodies used were: mouse anti-Shh (5E1, 1:10), mouse anti-Pax6 (PAX6, 1:100) (Developmental Studies Hybridoma Bank), rabbit anti-Olig2 (AB9610, 1:500, Millipore Sigma), and mouse anti-Arl13b (N295B/66, 1:1500, NeuroMab). Multiple sections from three embryos of each genotype were examined.

Analysis of anatomical measures

Weanling age (P20-P24) male and female mice were sacrificed and brains were harvested and fixed in 4% PFA overnight at 4°. Ten sex-matched pairs of mouse brains were collected for each genotype; control animals were either wild type or heterozygous

for the *Arl13b* point mutation or floxed allele (so *Arl13b*^{R79Q/+}, *Arl13b*^{V358A/+} or *Arl13b*^{flox/+}) or lacked *Brn4-Cre* (for *Arl13b*^{Brn4-Cre}). Brains were imaged on a tilted stage to present a surface view of the cerebellum, with a standard ruler in frame to confirm scale, using a dissecting microscope (Leica MZFLIII). Measurements were made in FIJI (Schindelin *et al.* 2012) with the investigator blind to genotype. Whole cerebellar width was measured at the widest part of the cerebellum, coinciding with lobule CI or CII. Vermis width was calculated by measuring the widest part of lobule VII (Deshpande *et al.* 2020). Hemisphere and vermis heights were measured at the longest rostro-caudal point of the hemisphere or at the midline of the vermis (see Figure 5). Body and brain weights were measured on a standard lab scale. For each sex- and age-matched pair, the ratio of mutant to control measures were calculated and graphed. A one-sample t-test was performed after transforming the ratio data to a log(2) scale to normalize data distribution (PRISM 8.2.0). Using the vermis width measures from our control brains (mean = 2.6mm, SD = 0.18), a power calculation for a one-sided t-test estimated 90% power to detect a 10% difference with 9-10 paired samples for a nominal alpha=0.05 (<https://clincalc.com/stats/samplesize.aspx> and <https://www.stat.ubc.ca/~rollin/stats/ssize/n2.html>).

Data availability

Mouse lines are available upon request. Supplemental material available at figshare. Figure S1 contains diagrams illustrating the complete results of the injection and tract tracing experiments. Figure S2 contains a schematic of the CRISPR strategy.

The authors affirm that all data necessary for confirming the conclusions of the article are present within the article and figures.

RESULTS

SMO is required for normal SCP projection to the dorsal thalamus

In order to test whether proper projection of SCPs requires Hh signaling, we compared the SCP tracts in mice in which we deleted the gene encoding the obligate Hh transducer, *Smoothed* (*Smo*), to control animals. As *Smo* null embryos die during embryogenesis, we deleted *Smo* specifically in the projection neurons by generating *Nex-Cre;Smo^{fl/fl}* mice, which we refer to as *Smo^{Nex-Cre}* (Zhang *et al.* 2001; Caspary *et al.* 2002). *Nex-Cre* initiates CRE recombinase expression at E11.5, as the precursor cells of the deep cerebellar nuclei (DCN) begin to migrate and become specified (Fink *et al.* 2006; Goebbels *et al.* 2006). In the mature cerebellum, the SCPs project rostrally from the DCN (illustrated in Figure 1A). After entering the midbrain, the SCPs cross the midline and again turn rostrally to project to two positions in the rostral thalamus: one tract takes a slight dorsal path and the other tract remains in the same plane; for simplicity here, we term these projection sites the dorsal and ventral thalamus, respectively (Bohne *et al.* 2019). To examine the SCP tracts, we used retrograde tract tracing in which we performed stereotaxic injections of a lipophilic dextran dye into either the dorsal (Figure 1B) or ventral (Figure 1C) thalamus and allowed the dye to diffuse through the axons to the associated neuron's cell body (~7 days); we then sacrificed the animal and examined the cerebellum for evidence of the lipophilic dye indicating tracing.

We found that both dorsal and ventral thalamus injections resulted in visible clusters of dye-stained cells in the contralateral DCN, and not the ipsilateral DCN, in control animals indicating the retrograde tract tracing reliably labelled the SCPs in our hands (Figures 1D-E, S1A-B, dorsal: 6/8; ventral: 8/11). In the *Smo^{Nex-Cre}* mice, the results differed depending on whether we injected in the dorsal or ventral thalamus (Figures 1F-G, S1C-D). In the ventral thalamus injections, we detected dye-stained clusters of cells in the contralateral DCN but not the ipsilateral DCN, indicating normal SCP projection to the ventral thalamus (Figure 1G, 4/4 injections). This indicates that at least some SCPs cross the midline. In the dorsal thalamus injections, we could not detect dye-stained clusters of cells in either the contralateral or ipsilateral DCN (Figure 1F, 0/6 injections) suggesting that SCPs lacking SMO do not project to the dorsal thalamus. These data implicate SMO as critical for proper projection of the SCPs to the dorsal thalamus.

ARL13B is required for normal SCP projection to the dorsal thalamus

Given that ARL13B regulates vertebrate Hh signaling in a variety of contexts, we next assessed ARL13B's role in proper SCP projection. In order to delete *ARL13B* specifically in projection neurons, we generated *Nex-Cre;Arl13b^{fl/fl}* mice, which we refer to as *Arl13b^{Nex-Cre}*. We performed dorsal and ventral thalamus injections for retrograde tract tracing to examine the SCPs (Figures 1H-I, S1C-D). In the ventral thalamus injections of *Arl13b^{Nex-Cre}* mice, we found dye-stained clusters of cells in the contralateral DCN consistent with normal SCP projections crossing the midline and projecting to the ventral thalamus (Figure 1I, 4/4 injections). In contrast, in the dorsal

thalamus injections of *Arl13b*^{Nex-Cre} mice, we generally did not detect dye-stained clusters of cells in either the contralateral or ipsilateral DCN, suggesting that the SCPs lacking ARL13B do not project to the dorsal thalamus (Figure 1H, 1/7 injections). These data link ARL13B function to normal SCP projection. Furthermore, they reveal a similar phenotype in *Smo*^{Nex-Cre} and *Arl13b*^{Nex-Cre} mice.

ARL13B does not function from within cilia to mediate SCP guidance

ARL13B and the other 35 genes implicated in Joubert syndrome associate with the cilium or centrosome leading to the assumption that protein dysfunction from these locales underlies JSRD phenotypes (Parisi 2019). To directly ask whether ARL13B mediates SCP guidance to the dorsal thalamus from within cilia, we examined a mouse expressing a cilia-excluded variant of ARL13B, ARL13B^{V358A} (Figure 2) (Gigante *et al.* 2020). We previously demonstrated that ARL13B^{V358A} retains all known ARL13B biochemical activity, is undetectable in cilia yet transduces vertebrate Hh signaling normally (Mariani *et al.* 2016; Gigante *et al.* 2020). We found that either dorsal or ventral thalamus injections resulted in visible clusters of dye-stained cells in the contralateral DCN in control (Figures 2A-B, S1G-H, dorsal: 3/3; ventral: 4/4) and *Arl13b*^{V358A/V358A} (Figure 2C-D, dorsal: 4/4; ventral: 3/3) animals. In the context of the previous result showing that *Arl13b*^{Nex-Cre} mice display abnormal SCP projections to the dorsal thalamus, these data demonstrate that ARL13B does not function from within cilia to regulate SCP projections.

SCP projection is normal in mice expressing a Joubert-causing allele, Arl13b^{R79Q}

In order to understand the relationship between ARL13B and MTS formation in Joubert syndrome, we generated a mouse expressing the JSRD-causing R79Q mutation (Figure 3A). We used CRISPR/Cas9 editing to change the conserved residue in the mouse genome (Figures 3B, S2). This amino acid change disrupts ARL13B's GEF activity for ARL3 (Gotthardt *et al.* 2015; Ivanova *et al.* 2017). We found *Arl13b*^{R79Q/R79Q} mice were viable and fertile. We bred the *Arl13b*^{R79Q} allele to the null *Arl13b*^Δ allele to make *Arl13b*^{R79Q/Δ} animals, which we found survived to adulthood. As *Arl13b*^{Δ/Δ} are embryonic lethal, this genetically demonstrates that *Arl13b*^{R79Q} is a hypomorphic allele of ARL13B (Su *et al.* 2012).

To assess the role of ARL13B^{R79Q} in SCP guidance, we performed dorsal and ventral thalamus dye injections in control and *Arl13b*^{R79Q/R79Q} mice. We identified visible clusters of dye-stained cells in the contralateral DCN in control (Figures 3C-D, S1E-F, dorsal: 4/4; ventral: 3/3) and *Arl13b*^{R79Q/R79Q} (Figure 3E-F, dorsal: 3/3; ventral: 3/4) animals. Thus, despite the constitutive expression of the JSRD-causing allele throughout development, we did not detect a SCP projection defect in the *Arl13b*^{R79Q/R79Q} mouse model. In the context of the abnormal SCP projections to the dorsal thalamus that we identified in the *Smo*^{Nex-Cre} and *Arl13b*^{Nex-Cre} mice, this result indicates that the *Arl13b*^{R79Q} allele does not disrupt SMO function or any ARL13B function that regulates SMO.

Arl13b^{R79Q/R79Q} mice display normal *Shh* signal transduction in neural tube patterning

To further investigate the role of the *Arl13b*^{R79Q} allele in Hh signaling, we examined embryonic neural patterning as it is exquisitely sensitive to alterations in Shh

activity (Chiang *et al.* 1996; Briscoe and Ericson 1999). We generated E10.5 embryos and stained neural tube sections with antibodies against Shh, Olig2 and Pax6 (Figure 4). As expected in wild type embryos, we observed Shh expression in the ventral-most cells (the floorplate), Olig2 expression in lateral cells and Pax6 expression dorsally. We also saw the established abnormal cell fates in null *Arl13b*^{ΔΔ} embryos: loss of Shh staining in the floorplate (Figure 4D), dorsal and ventral expansion of Olig2 expression (Figure 4H), and a dorsal shift in Pax6 expression (Figure 4L) (Su *et al.* 2012). We found both *Arl13b*^{R79Q/R79Q} (Figure 4B, F, J) and *Arl13b*^{R79Q/Δ} (Figure 4C, G, K) embryos displayed neural patterning indistinguishable from wild type embryos, indicating the *Arl13b*^{R79Q} allele is not dosage-sensitive and does not disrupt Shh signaling in determining neural cell fate. In addition, ARL13B^{R79Q} protein localized to cilia (Figure 4N), consistent with previous results (Li *et al.* 2010; Humbert *et al.* 2012; Li *et al.* 2016; Mariani *et al.* 2016).

Arl13b^{R79Q/R79Q} mice display normal cerebellar width

The lack of a SCP projection phenotype in the *Arl13b*^{R79Q/R79Q} mice surprised us since JS patients display the MTS. In addition to defects in the SCPs, the MTS is due to an underdeveloped cerebellar vermis, so we examined the width of the cerebellum and the cerebellar vermis (Figure 5) (Aguilar *et al.* 2012). To quantify cerebellar width, we performed analysis of surface-facing anatomical measurements validated to be sufficiently sensitive to detect small differences in vermis width (Deshpande *et al.* 2020). Briefly, we measured cerebellar width as well as cerebellar vermis width (widest part of lobule VII) of fixed whole mount dissected brains. For each sex- and age-matched pair,

we calculated the ratio of the width measurements from mutant to control, which we compared to a hypothetical value of 1 (indicating no difference between groups). We detected no differences in the overall cerebellar width or the cerebellar vermis width between control and *Arl13b*^{R79Q/R79Q} mice of either sex (Figure 5B). Furthermore, we found no difference in body or brain weight for *Arl13b*^{R79Q/R79Q} mutants compared to controls (Figure 5B). Thus, unlike patients carrying *ARL13B*^{R79Q/R79Q}, *Arl13b*^{R79Q/R79Q} mice do not display a detectable growth deficit in the cerebellar vermis.

Global cerebellar hypoplasia is observed in mice lacking Ar13b in all neurons

Cerebellar size is well established to be regulated, in part, via Shh signaling which controls proliferation of the cerebellar granule precursor cells (Kenney and Rowitch 2000; Chizhikov *et al.* 2007). In order to better understand the vermis hypoplasia phenotype seen in JBTS patients in relation to ARL13B, we wanted to investigate how ARL13B regulates cerebellar width. To do so, we crossed the *Brn4-Cre* allele into the conditional null *Arl13b*^{fl/fl} background, called *Arl13b*^{Brn4-Cre} (Figure 6). *Brn4-Cre* initiates expression at E8.5 throughout the neuroectoderm so the cerebellum develops in the absence of ARL13B (Heydemann *et al.* 2001; Hazen *et al.* 2012). We again calculated width ratios using surface-facing anatomical measurements and found the overall width of the cerebellum was 6% reduced in both females and males lacking ARL13B compared to control littermates at weaning (Figure 6B, p<0.1). More striking, in the cerebellar vermis we detected a 27% reduction in width in female and a 33% reduction in male mutants compared to controls (Figure 6B, p<0.0001). *Arl13b*^{Brn4-Cre} mice develop hydrocephaly just after weaning which often leads to death. This is

reflected in the body and brain weight ratios: while the mutant mice had slightly lower body weights, they had comparatively heavier brains (Figure 6B). From these data, we conclude that loss of *ARL13B* leads to a modest global cerebellar width deficit and a more pronounced cerebellar vermis width reduction.

Arl13b^{V358A/V358A} mice display normal cerebellar width

JS is classified as a ciliopathy due to the majority of causative genes encoding proteins that, like ARL13B, are associated with cilia. In order to better understand the role of ciliary ARL13B in cerebellar size, we examined cerebellar width in the mice expressing the cilia-excluded variant ARL13B^{V358A} (Figure 7) (Gigante *et al.* 2020). We detected no difference in the overall cerebellar width or that of the cerebellar vermis between control and *Arl13b*^{V358A/V358A} mice (Figure 7B). Thus, ARL13B does not control cerebellar width from within cilia.

DISCUSSION

Here we demonstrate that complete loss of *Arl13b* function in mouse can account for two aspects of the MTS: aberrant SCP thalamic targeting and cerebellar vermis hypoplasia. We expand the role of Hh signaling as a critical guidance cue by showing it is required for proper SCP projection to the dorsal thalamus. Our finding that the SCP phenotype is similar in *Smo*^{Nex-Cre} and *Arl13b*^{Nex-Cre} mice is consistent with a model whereby ARL13B regulates SCP projections to the dorsal thalamus via a SMO-dependent mechanism. In line with previous work showing that ARL13B does not function from within cilia to regulate Shh-guided axon guidance, we found normal SCP

thalamic targeting in mice expressing only a cilia-excluded ARL13B variant (Ferent *et al.* 2019). By mutating a conserved arginine to glutamine, we generated a mouse expressing a mutation linked to JSRD in humans and observed no change in vertebrate Hh signaling (Cantagrel *et al.* 2008). Additionally, we identified no defects in *Arl13b*^{R79Q/R79Q} SCP projections. Whereas complete ARL13B deletion (*Arl13b*^{Bm4-Cre}) in the cerebellum led to global hypoplasia, we show the cerebellum of *Arl13b*^{R79Q/R79Q} mice is not significantly different in width compared to controls.

Overall, our data indicate ARL13B function is critical for both SCP targeting and controlling cerebellar vermis width. At one level, our data implicate Hh signaling in the etiology of the MTS since we show that SCP targeting requires SMO. However, at another level, our data indicate that Hh-independent pathways are at play as we don't observe Hh-dependent neural tube patterning defects in the presence of the JSRD-causing *Arl13b*^{R79Q} allele. In other mouse models of JSRD where the hypoplasia is specific to the cerebellar vermis, Wnt signaling is affected (Lancaster *et al.* 2011). Thus, the MTS could be due to disruption of different pathways in the SCPs and the vermis. This would imply that the 35 JSRD implicated genes all affect the distinct pathways in a similar manner (Parisi 2019). Alternatively, the MTS may form due to alterations in any of a few pathways – and it is even possible that alterations in one pathway could impact other pathways – or the ability of cells to respond to those other pathways. Such a model is hinted at by previous work showing interplay between the Hh and Wnt pathways underlying the severity of hindbrain phenotypes (Hagemann and Scholpp 2012; Bashford and Subramanian 2019). While we haven't detected any changes in Wnt response in the absence of ARL13B function, we may have not examined the

relevant biological process or used a sensitive enough readout (Horner and Caspary 2011). Parallel reasoning would thus suggest that while *Arl13b*^{R79Q/R79Q} mice clearly transduce Hh reasonably well, there may be subtle changes in Shh signaling or changes that influence Wnt signaling. Our data are consistent with the complexity exhibited by other JSRD mouse models examined to date (Delous *et al.* 2007; Garcia-Gonzalo *et al.* 2011; Roberson *et al.* 2015; Bashford and Subramanian 2019).

In patients, the SCP targeting deficit appears more severe than what we observed in the mice. Available methods for live-imaging and examination of post-mortem tissues suggest a range of SCP decussation defects, with some tracts appearing thickened on the ipsilateral side relative to their DCN (Yachnis and Rorke 1999; Poretti *et al.* 2007). Surprisingly, in the mutant mice we infer midline crossing of the SCPs. In the case of the *Smo*^{Nex-Cre} conditional mice, it is formally possible that the SCPs do not rely on SMO for midline crossing but only for subsequent targeting to the dorsal thalamus. The fact that the *Arl13b*^{Nex-Cre} conditional mice display a highly similar phenotype to the *Smo*^{Nex-Cre} conditional mice makes this less likely, since ARL13B is directly implicated in JSRD and regulates SMO-dependent axon guidance in other contexts (Cantagrel *et al.* 2008; Ferent *et al.* 2019). It is also plausible that the protein turnover driven by *Nex-Cre* completed after midline crossing occurred. *Nex-Cre* expression initiates at E11.5 in the cells on the rhombic lip of the cerebellar anlage as they start to migrate and be specified before occupying the deep cerebellar nuclei (Fink *et al.* 2006; Goebbels *et al.* 2006). We expect deletion would occur in the precursors and therefore the neurons of the DCN would not express protein. Finally, it is possible that the mouse is not a valid system in which to model the SCP midline crossing defect.

This might explain why we saw no defects in the SCP targeting of the *Arl13b*^{R79Q/R79Q} mice, as this is a constitutive mutation that requires no protein turnover, yet homozygous expression of ARL13B^{R79Q} in patients results in the molar tooth sign, which has been associated with failed decussation of white matter tracts (Quisling *et al.* 1999). Indeed, other mouse mutants such as *Cep290* and *Ahi1* which recapitulate the cerebellar vermis hypoplasia, also do not display midline crossing defects in the SCPs (Lancaster *et al.* 2011). Whether this is due to anatomical distinctions between the cerebellum in mouse and human or the genetic background on which these models were examined are open questions. Recent work highlights clear molecular and temporal differences between mouse and human cerebellar development (Haldipur *et al.* 2019; Behesti *et al.* 2021).

Examining SCP projections is labor intensive and it has not been done systematically among the JSRD mouse models (Bashford and Subramanian 2019; Guo *et al.* 2019). While previous work showed that *Arl13b*^{Nex-Cre} and *Inpp5e*^{Nex-Cre} mice exhibit SCP targeting deficits, here we pinpoint the *Arl13b*^{Nex-Cre} defect as specific to the projection to the dorsal thalamus (Guo *et al.* 2019). The projection to the ventral thalamus remains intact, suggesting there is not a generalized deficit in axon outgrowth within the tract. The work on the *Arl13b*^{Nex-Cre} and *Inpp5e*^{Nex-Cre} SCP targeting deficits argue that PI3K/Akt signaling from within cilia led to the tract defects (Guo *et al.* 2019). However, we found that cilia-excluded ARL13B mediated SCP targeting normally. These conflicting results could be explained by differences in the experimental details as the data supporting ciliary ARL13B function used a human ARL13B viral expression construct to rescue conditionally-deleted mice whereas we used genetic mutations

engineered at the endogenous locus in this study. Alternatively, these data could indicate that ARL13B plays an important cellular role in the ciliary trafficking of key components needed for the PI3K/Akt pathways.

JSRD-causing mutations in ARL13B are generally restricted to the GTPase domain of the protein, although two residues outside that domain are implicated in disease (Cantagrel *et al.* 2008; Bachmann-Gagescu *et al.* 2015; Thomas *et al.* 2015; Shaheen *et al.* 2016; Rafiullah *et al.* 2017). Based on other ARL proteins, ARL13B likely assumes distinct conformations upon the binding either GDP or GTP, permitting different binding partners or altering affinities for binding partners (Pasqualato *et al.* 2002; Miertzschke *et al.* 2014). None of the tested JSRD-causing mutations (R79Q, Y86C or R200C) disrupt GTP binding or hydrolysis, however, all three mutations disrupt ARL13B function as an ARL3 GEF (Ivanova *et al.* 2017). The arginine at this position is located within the Switch 2 region of ARL13B and is conserved in humans, mice, zebrafish and *Chlamydomonas* (Figure 3B). We were surprised that we were unable to detect any defects in our mouse, as we expected to model some aspect of Joubert syndrome. Humans with homozygous R79Q mutation exhibit motor and ocular defects (among other symptoms), whereas *arl13b*-null zebrafish injected with human ARL13B-R79Q mRNA only partially rescue the ciliopathy phenotypes of curved body axis and cystic kidney (Cantagrel *et al.* 2008). These species-dependent differences emphasize the importance of analyzing mutations expressed from the endogenous promoter over the course of normal development. A recent study linked *arl13b* disruption during zebrafish development to reduced granule and Purkinje cells through a down-regulation of Wnt signaling (Zhu *et al.* 2020). Given that complete deletion of ARL13B impacts

broader biological processes in the cerebellum than the R79Q mutation and that the null mutant misregulates Hh signaling whereas R79Q does not, we conclude that a subset of ARL13B function is disrupted in JSRD.

Acknowledgements

We are grateful to L. Mariani for her initial work on this project, R.E. Van Sciver and E. Gigante for critical comments on the manuscript and J.G. Mulle for the statistical consults.

Funding

This work was supported by funding from National Institutes of Health grants R01NS090029, R01GM110663 and R35GM122549 to T.C. and T32GM008490 and F31NS101806 to S.K.S. with additional support from the Emory University Integrated Cellular Imaging Microscopy Core of the Emory Neuroscience NINDS Core Facilities grant, P30NS055077. This study was supported in part by the Mouse Transgenic and Gene Targeting Core (TMF), which is subsidized by the Emory university School of Medicine and is one of the Emory Integrated Core Facilities. Additional support was provided by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number UL1TR000454. The content is solely the responsibility of the authors and does not necessarily reflect the official views of the National Institutes of Health

Conflicts of Interest

526 The authors have no competing interests to declare.

527

528 **Author contributions statement**

529 Author Contributions: Conceptualization T.C.; Methodology S.K.S. and A.B.L.;

530 Validation S.K.S. and A.B.L.; Formal Analysis S.K.S. and A.B.L.; Investigation S.K.S.

531 and A.B.L.; Writing – Original Draft S.K.S. and T.C.; Writing – Review & Editing S.K.S.,

532 A.B.L. and T.C.; Visualization S.K.S. and A.B.L.; Supervision T.C.; Project

533 Administration T.C.; Funding Acquisition T.C.

534

535 **References**

536 Abdelhamed, Z. A., D. I. Abdelmottaleb, M. E. El-Asrag, S. Natarajan, G. Wheway et al.,

537 2019 The ciliary Frizzled-like receptor Tmem67 regulates canonical Wwnt/beta-

538 catenin signalling in the developing cerebellum via Hoxb5. *Sci Rep.* **9**: 5446.

539 doi:10.1038/s41598-019-41940-5

540 Abdelhamed, Z. A., G. Wheway, K. Szymanska, S. Natarajan, C. Toomes et al., 2013

541 Variable expressivity of ciliopathy neurological phenotypes that encompass

542 Meckel-Gruber syndrome and Joubert syndrome is caused by complex de-

543 regulated ciliogenesis, Shh and Wnt signalling defects. *Hum Mol Genet.* **22**:

544 1358-1372. doi:10.1093/hmg/dds546

545 Aguilar, A., A. Meunier, L. Strehl, J. Martinovic, M. Bonniere et al., 2012 Analysis of

546 human samples reveals impaired SHH-dependent cerebellar development in

547 Joubert syndrome/Meckel syndrome. *Proc Natl Acad Sci U S A.* **109**: 16951-

548 16956. doi:10.1073/pnas.1201408109

549 Alkanderi, S., E. Molinari, R. Shaheen, Y. Elmaghloob, L. A. Stephen et al., 2018 ARL3
550 mutations cause Joubert syndrome by disrupting ciliary protein composition. *Am*
551 *J Hum Genet.* **103**: 612-620. doi:10.1016/j.ajhg.2018.08.015

552 Arts, H. H., D. Doherty, S. E. C van Beersum, M. A. Parisi, S. J. F. Letteboer et al., 2007
553 Mutations in the gene encoding the basal body protein RPGRIP1L, a
554 nephrocystin-4 interactor, cause Joubert syndrome. *Nat Genet.* **39**: 882-888.
555 doi:10.1038/ng2069

556 Asadollahi, R., J. E. Strauss, M. Zenker, O. Beuing, S. Edvardson et al., 2018 Clinical
557 and experimental evidence suggest a link between KIF7 and C5orf42-related
558 ciliopathies through Sonic Hedgehog signaling. *Eur J Hum Genet.* **26**: 197-209.
559 doi:10.1038/s41431-017-0019-9

560 Bachmann-Gagescu, R., J. C. Dempsey, I. G. Phelps, B. J. O'Roak, D. M. Knutzen et
561 al., 2015 Joubert syndrome: A model for untangling recessive disorders with
562 extreme genetic heterogeneity. *J Med Genet.* **52**: 514-522.
563 doi:10.1136/jmedgenet-2015-103087

564 Bachmann-Gagescu, R., J. C. Dempsey, S. Bulgheroni, M. L. Chen, S. D'Arrigo et al.,
565 2020 Healthcare recommendations for Joubert syndrome. *Am J Med Genet A.*
566 **182**: 229-249. doi:10.1002/ajmg.a.61399

567 Bashford, A. L., and V. Subramanian, 2019 Mice with a conditional deletion of Talpid3
568 (KIAA0586) - a model for Joubert syndrome. *J Pathol.* **248**: 396-408.
569 doi:10.1002/path.5271

570 Bay, S.N., A. B. Long, and T. Caspary, 2018 Disruption of the ciliary GTPase Arl13b
571 suppresses Sonic hedgehog overactivation and inhibits medulloblastoma

formation. *Proc Natl Acad Sci U S A*. **115**: 1570-1575.
doi:10.1073/pnas.1706977115

Behesti, H. A. Kocabas, D. E. Buchholz, T. S. Carroll, and M. E. Hatten, 2021 Altered temporal sequence of transcriptional regulators in the generation of human cerebellar granule cells. bioRxiv. doi:10.1101/2021.01.17.427030 (Preprint posted January 17, 2021).

Bielas, S. L., J. L. Silhavy, F. Brancati, M. V. Kisseleva, L. Al-Gazali et al., 2009 Mutations in INPP5E, encoding inositol polyphosphate-5-phosphatase E, link phosphatidyl inositol signaling to the ciliopathies. *Nat Genet*. **41**: 1032-1036. doi:10.1038/ng.423

Bimonte, S., A. De Angelis, L. Quagliata, F. Giusti, R. Tammaro et al., 2011 Odf1 is required in limb bud patterning and endochondral bone development. *Dev Biol*. **349**: 179-191. doi:10.1016/j.ydbio.2010.09.020

Bohne, P., M. K. Schwarz, S. Herlitze, and M. D. Mark, 2019 A new projection from the deep cerebellar nuclei to the hippocampus via the ventrolateral and laterodorsal thalamus in mice. *Front Neural Circuits*. **13**: 51. doi:10.3389/fncir.2019.00051

Brancati, F., B. Dallapiccola, and E. M. Valente, 2010 Joubert syndrome and related disorders. *Orphanet J Rare Dis*. **5**: 20. doi:10.1186/1750-1172-5-20

Breslow, D. K., S. Hoogendoorn, A. R. Kopp, D. W. Morgens, B. K. Vu et al., 2018 A CRISPR-based screen for Hedgehog signaling provides insights into ciliary function and ciliopathies. *Nat Genet*. **50**: 460-471. doi:10.1038/s41588-018-0054-7

594 Briscoe, J., and J. Ericson J, 1999 The specification of neuronal identity by graded
595 Sonic Hedgehog signalling. *Semin Cell Dev Biol.* **10**: 353-362.
596 doi:10.1006/scdb.1999.0295

597 Cantagrel, V., J. L. Silhavy, S. L. Bielas, D. Swistun, S. E. Marsh et al., 2008 Mutations
598 in the cilia gene ARL13B lead to the classical form of Joubert syndrome. *Am J*
599 *Hum Genet.* **83**: 170-179. doi:10.1016/j.ajhg.2008.06.023

600 Casoni, F., L. Croci, F. Vincenti, P. Podini, M. Riba et al., 2020 ZFP423 regulates early
601 patterning and multiciliogenesis in the hindbrain choroid plexus. *Development.*
602 **147**: dev190173. doi:10.1242/dev.190173

603 Caspary, T., M. J. Garcia-Garcia, D. Huangfu, J. T. Eggenschwiler, M. R. Wyler et al.,
604 2002 Mouse Dispatched homolog1 is required for long-range, but not juxtacrine,
605 Hh signaling. *Curr Biol.* **12**: 1628-1632. doi:10.1016/s960-9822(02)01147-8

606 Caspary, T., C. E. Larkins, and K. V. Anderson, 2007 The graded response to Sonic
607 Hedgehog depends on cilia architecture. *Dev Cell.* **12**: 767-778.
608 doi:10.1016/j.devcel.2007.03.004

609 Charron, F., E. Stein, J. Jeong, A. P. McMahon, and M. Tessier-Lavigne, 2003 The
610 morphogen sonic hedgehog is an axonal chemoattractant that collaborates with
611 netrin-1 in midline axon guidance. *Cell.* **113**: 11-23. doi:10.1016/s0092-
612 8674(03)00199-5

613 Chavez, M., S. Ena, J. Van Sande, A. de Kerchove d'Exaerde, S. Schurmans et al.,
614 2015 Modulation of ciliary phosphoinositide content regulates trafficking and
615 Sonic hedgehog signaling output. *Dev Cell.* **34**: 338-350.
616 doi:10.1016/j.devcel.2015.06.016

617 Chiang, C., Y. Litingtung, E. Lee, K. E. Young, J. L. Corden et al., 1996 Cyclopia and
618 defective axial patterning in mice lacking Ssonic hedgehog gene function.
619 *Nature*. **383**: 407-413. Doi:10.1038/383407a0

620 Chih, B., P. Liu, Y. Chinn, C. Chalouni, L. G. Komuves et al., 2011 A ciliopathy complex
621 at the transition zone protects the cilia as a privileged membrane domain. *Nat*
622 *Cell Biol.* **14**: 61-72. doi:10.1038/ncb2410

623 Chizhikov, V. V., J. Davenport, Q. Zhang, E. K. Shih, O. A. Cabello et al., 2007 Cilia
624 proteins control cerebellar morphogenesis by promoting expansion of the granule
625 progenitor pool. *J Neurosci.* **27**: 9780-9789. doi:10.1523/JNEUROSCI.5586-
626 06.2007

627 Christopher, K. J., B. Wang, Y. Kong, and S. D. Weatherbee, 2012 Forward genetics
628 uncovers Transmembrane protein 107 as a novel factor required for ciliogenesis
629 and Sonic hedgehog signaling. *Dev Biol.* **368**: 382-392.
630 doi:10.1016/j.ydbio.2012.06.008

631 Constable, S., A. B. Long, K. A. Floyd, S. Schurmans, and T. Caspary, 2020 The ciliary
632 phosphatidylinositol phosphatase Inpp5e plays positive and negative regulatory
633 roles in Shh signaling. *Development.* **147**: dev183301. doi:10.1242/dev.183301

634 Consugar, M. B., V. J. Kubly, D. J. Lager, C. J. Hommerding, W. C. Wong et al., 2007
635 Molecular diagnostics of Meckel–Gruber syndrome highlights phenotypic
636 differences between MKS1 and MKS3. *Hum Genet.* **121**: 591-599.
637 doi:10.1007/s00439-007-0341-3

638 Dahmane, N., and A. Ruiz i Altaba, 1999 Sonic hedgehog regulates the growth and
639 patterning of the cerebellum. *Development.* **126**: 3089-3100.

640 Davey, M. G., I. R. Paton, Y. Yin, M. Schmidt, F. K. Bangs et al., 2006 The chicken
641 talpid3 gene encodes a novel protein essential for Hedgehog signaling. *Genes*
642 *Dev.* **20**: 1365-1377. doi:10.1101/gad.369106

643 Delous, M., L. Baala, R. Salomon, C. Laclef, J. Vierkotten et al., 2007 The ciliary gene
644 RPGRIP1L is mutated in cerebello-oculo-renal syndrome (Joubert syndrome type
645 B) and Meckel syndrome. *Nat Genet.* **39**: 875-881. doi:10.1038/ng2039

646 Deshpande, O., R. Z. Lara, O. R. Zhang, D. Concepcion, and B. A. Hamilton, 2020
647 ZNF423 patient variants, truncations, and in-frame deletions in mice define an
648 allele-dependent range of midline brain abnormalities. *PLoS Genet.* **16**:
649 e1009017. doi:10.1371/journal.pgen.1009017

650 Doherty, D., 2009 Joubert syndrome: Insights into brain development, cilium biology,
651 and complex disease. *Semin Pediatr Neurol.* **16**: 143-154.
652 doi:10.1016/j.spen.2009.06.002

653 Dowdle, W. E., J. F. Robinson, A. Kneist, M. S. Sirerol-Piquer, S. G. M. Frints et al.,
654 2011 Disruption of a ciliary B9 protein complex causes Meckel syndrome. *Am J*
655 *Hum Genet.* **89**: 94-110. doi:10.1016/j.ajhg.2011.06.003

656 Ferent, J., S. Constable, E. D. Gigante, P. T. Yam, L. E. Mariani et al., 2019 The ciliary
657 protein Arl13b functions outside of the primary cilium in Shh-mediated axon
658 guidance. *Cell Rep.* **29**: 3356-3366.e3353. doi:10.1016/j.celrep.2019.11.015

659 Fink, A. J., C. Englund, R. A. M. Daza, D. Pham, C. Lau et al., 2006 Development of
660 the deep cerebellar nuclei: Transcription factors and cell migration from the
661 rhombic lip. *J Neurosci.* **26**: 3066-3076. doi:10.1523/JNEUROSCI.5203-05.2006

662 Friestad, K-A. M., E. Molinari, M. Thoresen, S. A. Ramsbottom, F. Hughes et al., 2019
663 A CEP104-CSPP1 complex is required for formation of primary cilia competent in
664 Hedgehog signaling. *Cell Rep.* **28**: 1907-1922.e6
665 doi:10.1016/j.celrep.2019.07.025

666 Garcia-Gonzalo, F. R., K. C. Corbit, M. S. Sirerol-Piquer, G. Ramaswami, E. A. Otto et
667 al., 2011 A transition zone complex regulates mammalian ciliogenesis and ciliary
668 membrane composition. *Nat Genet.* **43**: 776-784. doi:10.1038/ng.891

669 Garcia-Gonzalo, F. R., S. C. Phua, E. C. Roberson, G. Garcia 3rd, M. Abedin et al.,
670 2015 Phosphoinositides regulate ciliary protein trafficking to modulate Hedgehog
671 signaling. *Dev Cell.* **34**: 400-409. doi:10.1016/j.devcel.2015.08.001

672 Gaudet, M., A-G. Fara, I. Beritognolo, and M. Sabatti, 2009 Allele-specific PCR in SNP
673 genotyping. *Methods Mol Biol.* **578**: 415-424. doi:10.1007/978-1-60327-411-1_26

674 Gigante, E. D., M. R. Taylor, A. A. Ivanova, R. A. Kahn, and T. Caspary, 2020. ARL13B
675 regulates Sonic hedgehog signaling from outside primary cilia. *eLife.* **9**: e50434.
676 doi:10.7554/eLife.50434

677 Goebbels, S., I. Bormuth, U. Bode, O. Hermanson, M. H. Schwab et al., 2006 Genetic
678 targeting of principal neurons in neocortex and hippocampus of NEX-Cre mice.
679 *Genesis.* **44**: 611-621. doi:10.1002/dvg.20256

680 Gotthardt, K., M. Lokaj, C. Koerner, N. Falk, A. Giessl et al., 2015 A G-protein
681 activation cascade from Arl13b to Arl3 and implications for ciliary targeting of
682 lipidated proteins. *eLife.* **4**: e11859. doi:10.7554/eLife.11859

683 Guo, J., J. M. Otis, S. K. Suci, C. Catalano, L. Xing et al., 2019 Primary cilia signaling
684 promotes axonal tract development and is disrupted in Joubert syndrome-related
685 disorders models. *Dev Cell*. **51**: 759-774 e755. doi:10.1016/j.devcel.2019.11.005

686 Hagemann, A. I. H., and S. Scholpp, 2012 The tale of the three brothers – Shh, Wnt,
687 and Fgf during development of the thalamus. *Front Neurosci*. **6**: 76.
688 doi:10.3389/fnins.2012.00076

689 Haldipur, P., K. A. Aldinger, S. Bernardo, M. Deng, A. E. Timms et al., 2019
690 Spatiotemporal expansion of primary progenitor zones in the developing human
691 cerebellum. *Science*. **366**: 454-460. Doi: 10.1126/science.aax7526

692 Hata, A., J. Seoane, G. Lagna, E. Montalvo, A. Hemmati-Brivanlou et al., 2000 OAZ
693 uses distinct DNA- and protein-binding zinc fingers in separate BMP-Smad and
694 Olf signaling pathways. *Cell*. **100**: 229-240. doi:10.1016/s0092-8674(00)81561-5

695 Hazen, V. M. , M. G. Andrews, L. Umans, E. B. Crenshaw 3rd, A. Zwijsen et al., 2012
696 BMP receptor-activated Smads confer diverse functions during the development
697 of the dorsal spinal cord. *Dev Biol*. **367**: 216-227.
698 doi:10.1016/j.ydbio.2012.05.014

699 Heydemann, A., L. C. Nguyen, and E. B. Crenshaw 3rd, 2001 Regulatory regions from
700 the Brn4 promoter direct LACZ expression to the developing forebrain and neural
701 tube. *Brain Res Dev Brain Res*. **128**: 83-90. doi:10.1016/s0165-3806(01)00137-7

702 Hopp, K., C. M. Heyer, C. J. Hommerding, S. A. Henke, J. L. Sundsbak et al., 2011
703 B9D1 is revealed as a novel Meckel syndrome (MKS) gene by targeted exon-
704 enriched next-generation sequencing and deletion analysis. *Hum Mol Genet*. **20**:
705 2524-2534. doi:10.1093/hmg/ddr151

706 Horner, V. L., and T. Caspary, 2011 Disrupted dorsal neural tube BMP signaling in the
707 cilia mutant *Arl13b* hnn stems from abnormal Shh signaling. *Dev Biol.* **355**: 43-
708 54. doi:10.1016/j.ydbio.2011.04.019

709 Huang, S., J. Laoukili, M. T. Epping, J. Koster, M. Hölzel et al., 2009 ZNF423 is
710 critically required for retinoic acid-induced differentiation and is a marker of
711 neuroblastoma outcome. *Cancer Cell.* **15**: 328-340.
712 doi:10.1016/j.ccr.2009.02.023

713 Huangfu, D., A. Liu, A. S. Rakeman, N. S. Murcia, L. Niswander et al., 2003 Hedgehog
714 signalling in the mouse requires intraflagellar transport proteins. *Nature.* **426**: 83-
715 87. doi:10.1038/nature02061

716 Humbert, M. C., K. Weihbrecht, C. C. Searby, Y. Li, R. M. Pope et al., 2012 ARL13B,
717 PDE6D, and CEP164 form a functional network for INPP5E ciliary targeting. *Proc*
718 *Natl Acad Sci U S A.* **109**: 19691-19696. doi:10.1073/pnas.1210916109

719 Hynes, A. M., R. H. Giles, S. Srivastava, L. Eley, J. Whitehead et al., 2014 Murine
720 Joubert syndrome reveals Hedgehog signaling defects as a potential therapeutic
721 target for nephronophthisis. *Proc Natl Acad Sci U S A.* **111**: 9893-9898.
722 doi:10.1073/pnas.1322373111

723 Ivanova, A. A., T. Caspary, N. T. Seyfried, D. M. Duong, A. B. West et al., 2017
724 Biochemical characterization of purified mammalian ARL13B protein indicates
725 that it is an atypical GTPase and ARL3 guanine nucleotide exchange factor
726 (GEF). *J Biol Chem.* **292**: 11091-11108. doi:10.1074/jbc.M117.784025

727 Kenney, A. M., and D. H. Rowitch, 2000 Sonic hedgehog promotes G(1) cyclin
728 expression and sustained cell cycle progression in mammalian neuronal

729 precursors. *Mol Cell Biol.* **20**: 9055-9067. doi:10.1128/mcb.20.23.9055-
730 9067.2000

731 Lai, C. K., N. Gupta, X. Wen, L. Rangell, B. Chih et al., 2011 Functional
732 characterization of putative cilia genes by high-content analysis. *Mol Biol Cell.*
733 **22**: 1104-1119. doi:10.1091/mbc.E10-07-0596

734 Lan, Y., and R. Jiang, 2009 Sonic hedgehog signaling regulates reciprocal epithelial-
735 mesenchymal interactions controlling palatal outgrowth. *Development.* **136**:
736 1387-1396. doi:10.1242/dev.028167

737 Lancaster, M. A., D. J. Gopal, J. Kim, S. N. Saleem, J. L. Silhavy et al., 2011 Defective
738 Wnt-dependent cerebellar midline fusion in a mouse model of Joubert syndrome.
739 *Nat Med.* **17**: 726-731. doi:10.1038/nm.2380

740 Li, Y., Q. Wei, Y. Zhang, K. Ling, and J. Hu, 2010 The small GTPases ARL-13 and
741 ARL-3 coordinate intraflagellar transport and ciliogenesis. *J Cell Biol.* **189**: 1039-
742 1051. doi:10.1083/jcb.200912001

743 Li, Y., X. Tian, M. Ma, S. Jerman, S. Kong et al., 2016 Deletion of ADP ribosylation
744 factor-like GTPase 13B leads to kidney cysts. *J Am Soc Nephrol.* **27**: 3628-3638.
745 doi:10.1681/ASN.2015091004

746 Lipinski, R. J., C. Song, K. K. Sulik, J. L. Everson, J. J. Gipp et al., 2010 Cleft lip and
747 palate results from Hedgehog signaling antagonism in the mouse: Phenotypic
748 characterization and clinical implications. *Birth Defects Res A Clin Mol Teratol.*
749 **88**: 232-240. doi:10.1002/bdra.20656

750 Maria, B. L., K. B. Hoang, R. J. Tusa, A. A. Mancuso, L. M. Hamed et al., 1997 “Joubert
751 syndrome” revisited: key ocular motor signs with magnetic resonance imaging
752 correlation. *J Child Neurol.* **12**: 423-430. doi:10.1177/088307389701200703

753 Mariani, L. E., M. F. Bijlsma, A. A. Ivanova, S. K. Suci, R. A. Kahn et al., 2016 Arl13b
754 regulates Shh signaling from both inside and outside the cilium. *Mol Biol Cell.* **27**:
755 3780-3790. doi:10.1091/mbc.E16-03-0189

756 Miertzschke, M., C. Koerner, M. Spoerner, and A. Wittinghofer, 2014 Structural insights
757 into the small G-protein Arl13b and implications for Joubert syndrome. *Biochem*
758 *J.* **457**: 301-311. doi:10.1042/BJ20131097

759 Munoz-Estrada, J., and R. J. Ferland, 2019 Ahi1 promotes Arl13b ciliary recruitment,
760 regulates Arl13b stability and is required for normal cell migration. *J Cell Sci.*
761 **132**: jcs230680. doi:10.1242/jcs.230680

762 Nolan-Steva, O., J. Lau, M. L. Truitt, G. C. Chu, M. Hebrok et al., 2009 Gli1 is
763 regulated through Smoothened-independent mechanisms in neoplastic
764 pancreatic ducts and mediates PDAC cell survival and transformation. *Genes*
765 *Dev.* **23**: 24-36. doi:10.1101/gad.1753809

766 Nuovo, S., I. Bacigalupo, M. Ginevrino, R. Battini, E. Bertini et al., 2020 Age and sex
767 prevalence estimate of Joubert syndrome in Italy. *Neurology.* **94**: e797-e801.
768 doi:10.1212/WNL.0000000000008996

769 Onimaru, H., Y. Kumagawa, and I. Homma, 2006 Respiration-related rhythmic activity
770 in the rostral medulla of newborn rats. *J Neurophysiol.* **96**: 55-61.
771 doi:10.1152/jn.01175.2005

772 Otto, E. A., K. Tory, M. Attanasio, W. Zhou, M. Chaki et al, 2009 Hypomorphic
773 mutations in meckelin (MKS3/TMEM67) cause nephronophthisis with liver
774 fibrosis (NPHP11). *J Med Genet.* **46**: 663-670. doi:10.1136/jmg.2009.066613

775 Parisi, M. A, 2019 The molecular genetics of Joubert syndrome and related ciliopathies:
776 The challenges of genetic and phenotypic heterogeneity. *Transl Sci Rare Dis.* **4**:
777 25-49. doi:10.3233/TRD-190041

778 Parisi, M. A., D. Doherty, P. F. Chance, and I. A. Glass, 2007 Joubert syndrome (and
779 related disorders) (OMIM 213300). *Eur J Hum Genet.* **15**: 511-521.
780 doi:10.1038/sj.ejhg.5201648

781 Pasqualato, S., L. Renault, and J. Cherfils, 2002 Arf, Arl, Arp and Sar proteins: A family
782 of GTP-binding proteins with a structural device for 'front-back' communication.
783 *EMBO Rep.* **3**: 1035-1041. doi:10.1093/embo-reports/kvf221

784 Poretti, A., E. Boltshauser, T. Loenneker, E. M. Valente, F. Brancati et al., 2007
785 Diffusion tensor imaging in Joubert syndrome. *AJNR Am J Neuroradiol.* **28**:
786 1929-1933. doi:10.3174/ajnr.A0703

787 Quisling, R. G., A. J. Barkovich, and B. L. Maria, 1999 Magnetic resonance imaging
788 features and classification of central nervous system malformations in Joubert
789 syndrome. *J Child Neurol.* **14**: 628-635. doi:10.1177/088307389901401002

790 Rafiullah, R., A. B. Long, A. A. Ivanova, H. Ali, S. Berkel et al., 2017 A novel
791 homozygous ARL13B variant in patients with Joubert syndrome impairs its
792 guanine nucleotide-exchange factor activity. *Eur J Hum Genet.* **25**: 1324-1334.
793 doi:10.1038/s41431-017-0031-0

794 Ramsbottom, S. A., P. E. Thelwall, K. M. Wood, G. J. Clowry, L. A. Devlin et al., 2020
795 Mouse genetics reveals Barttin as a genetic modifier of Joubert syndrome. *Proc*
796 *Natl Acad Sci U S A.* **117**: 1113-1118. doi:10.1073/pnas.1912602117

797 Reiter, J. F., and W. C. Skarnes, 2006 Tectonic, a novel regulator of the Hedgehog
798 pathway required for both activation and inhibition. *Genes Dev.* **20**: 22-27.
799 doi:10.1101/gad.1363606

800 Roberson, E. C., W. E. Dowdle, A. Ozanturk, F. R. Garcia-Gonzalo, C. Li et al., 2015
801 TMEM231, mutated in orofacioidigital and Meckel syndromes, organizes the
802 ciliary transition zone. *J Cell Biol.* **209**: 129-142. doi:10.1083/jcb.201411087

803 Sang, L., J. J. Miller, K. C. Corbit, R. H. Giles, M. J. Brauer et al., 2011 Mapping the
804 NPHP-JBTS-MKS protein network reveals ciliopathy disease genes and
805 pathways. *Cell.* **145**: 513-528. doi:10.1016/j.cell.2011.04.019

806 Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair et al., 2012 Fiji: an
807 open-source platform for biological-image analysis. *Nat Methods.* **9**: 676-682.
808 doi:10.1038/nmeth.2019

809 Schrick, J. J., P. Vogel, A. Abuin, B. Hampton, and D. S. Rice, 2006 ADP-ribosylation
810 factor-like 3 is involved in kidney and photoreceptor development. *Am J Pathol.*
811 **168**: 1288-1298. doi:10.2353/ajpath.2006.050941

812 Schwarz, N., A. Lane, K. Jovanovic, D. A. Parfitt, M. Aguila et al., 2017 Arl3 and RP2
813 regulate the trafficking of ciliary tip kinesins. *Hum Mol Genet.* **26**: 2480-2492.
814 doi:10.1093/hmg/ddx143

815 Shaheen, R., K. Szymanska, B. Basu, N. Patel, N. Ewida et al., 2016 Characterizing
816 the morbid genome of ciliopathies. *Genome Biol.* **17**: 242. doi:10.1186/s13059-
817 016-1099-5

818 Srour, M., F. F. Hamdan, J. A. Schwartzentruber, L. Patry, L. H. Ospina et al., 2012
819 Mutations in TMEM231 cause Joubert syndrome in French Canadians. *J Med*
820 *Genet.* **49**: 636-641. doi:10.1136/jmedgenet-2012-101132

821 Su, C-Y., S. N. Bay, L. E. Mariani, M. J. Hillman, and T. Caspary, 2012 Temporal
822 deletion of Arl13b reveals that a mispatterned neural tube corrects cell fate over
823 time. *Development.* **139**: 4062-4071. doi:10.1242/dev.082321

824 Thomas, S., V. Cantagrel, L. Mariani, V. Serre, J-E. Lee et al., 2015 Identification of a
825 novel ARL13B variant in a Joubert syndrome-affected patient with retinal
826 impairment and obesity. *Eur J Hum Genet.* **23**: 621-627.
827 doi:10.1038/ejhg.2014.156

828 Thomas, S., M. Legendre, S. Saunier, B. Bessières, C. Alby et al., 2012 TCTN3
829 mutations cause Mohr-Majewski syndrome. *Am J Hum Genet.* **91**: 372-378.
830 doi:10.1016/j.ajhg.2012.06.017

831 Valente, E. M., F. Brancati, and B. Dallapiccola, 2008 Genotypes and phenotypes of
832 Joubert syndrome and related disorders. *Eur J Med Genet.* **51**: 1-23.
833 doi:10.1016/j.ejmg.2007.11.003

834 Valente, E. M., C. V. Logan, S. Mougou-Zerelli, J. H. Lee, J. L. Silhavy et al., 2010
835 Mutations in TMEM216 perturb ciliogenesis and cause Joubert, Meckel and
836 related syndromes. *Nat Genet.* **42**: 619-625. doi:10.1038/ng.594

- Vierkotten, J., R. Dildrop, T. Peters, B. Wang, and U. Rüther, 2007 Ftm is a novel basal body protein of cilia involved in Shh signaling. *Development*. **134**: 2569-2577. doi:10.1242/dev.003715
- Weatherbee, S. D., L. A. Niswander, and K. V. Anderson, 2009 A mouse model for Meckel syndrome reveals Mks1 is required for ciliogenesis and Hedgehog signaling. *Hum Mol Genet*. **18**: 4565-4575. doi:10.1093/hmg/ddp422
- Wechsler-Reya, R. J., and M. P. Scott, 1999 Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron*. **22**: 103-114. doi:10.1016/s0896-6273(00)80682-0
- Wu, C., M. Yang, J. Li, C. Wang, T. Cao et al., 2014 Talpid3-binding centrosomal protein Cep120 is required for centriole duplication and proliferation of cerebellar granule neuron progenitors. *PLoS One*. **9**: e107943. doi:10.1371/journal.pone.0107943
- Yachnis, A. T., and L. B. Rorke, 1999 Neuropathology of Joubert syndrome. *J Child Neurol*. **14**: 655-659. doi:10.1177/088307389901401006
- Zhang, X. M., M. Ramalho-Santos, and A. P. McMahon, 2001 Smoothed mutants reveal redundant roles for Shh and Ihh signaling including regulation of L/R symmetry by the mouse node. *Cell*. **106**: 781-792.
- Zhu, J., H.-T. Wang, Y.-R. Chen, L.-Y. Yan, Y.-Y. Han et al., 2020 The Joubert syndrome gene arl13b is critical for early cerebellar development in zebrafish. *Neurosci Bull*. **36**: 1023-1034. doi:10.1007/s12264-020-00554-y

Figure legends

Figure 1: SCPs lacking *Arl13b* or *Smo* fail to project to the dorsal thalamus. (A-C)

Schematics of injections and fluorescent tracer diffusion shown horizontally (A) or sagittally (B-C). (A) Red dashed arrow depicts dye path in a successful injection from injection site (red X) caudal through the brain and across the midline and into the contralateral cerebellar DCN (red arrowhead). Grey background boxes indicate area of subsequent images: the injection site (INJ) and cerebellum (DCN). (D-I) Representative images of dorsal (D, F, H) or ventral (E, G, I) thalamus injection site (top panel) and cerebellum (middle panel) with the DCN in hatched white circle and magnified (bottom panel) with recoloring to black and white to aid visualization. The retrograde fluorescent tracer is pink-red and sections are stained with DAPI (blue). Numbers indicate the number of positively stained DCN clusters (DCN traced) out of the total number of injected animals. Note that no tracing was observed on ipsilateral side to injection. (D, E) Fluorescent tracer injection in *Smo^{fl/+}; Arl13b^{fl/+}; Nex-Cre* control animals resulted in contralateral DCN staining in (D) 6/8 dorsal thalamus injections and (E) 8/11 ventral thalamus injections. (F, G) Fluorescent tracer injection in *Smo^{Nex-Cre}* animals resulted in contralateral DCN staining in (F) 0/6 dorsal thalamus injections and (G) 4/4 ventral thalamus injections. (H, I) Fluorescent tracer injection in *Arl13b^{Nex-Cre}* animals resulted in contralateral DCN staining in (H) 1/7 dorsal thalamus injections and (I) 4/4 ventral thalamus injections. Two-tailed Fisher's exact test was performed; only the dorsal injection results were significantly different from the controls (F: $p < 0.01$; H: $p < 0.05$)

Figure 2: SCPs expressing cilia-excluded ARL13B^{V358A} project normally to both the dorsal and ventral thalamus. (A-D) Representative images of dorsal (A, C) or

ventral (B, D) thalamus injection site (top panel) and cerebellum (middle panel) with the DCN in hatched white circle and magnified (bottom panel) with recoloring to black and white to aid visualization. Numbers indicate the number of positively stained DCN clusters out of the total number of injected animals. Note that no tracing was observed on the injection's ipsilateral side. (A, B) Fluorescent tracer injection in *Arl13b*^{V358A/+} control animals resulted in contralateral DCN staining in (A) 3/3 dorsal thalamus injections and (B) 4/4 ventral thalamus injections. (C-D) Fluorescent tracer injection in *Arl13b*^{V358A/V358A} animals resulted in contralateral DCN staining in (C) 4/4 dorsal thalamus injections (Fisher's exact test, two-tailed, ns, p>0.9) and (D) 3/3 ventral thalamus injections (ns, p>0.9).

Figure 3: SCPs expressing JS allele *Arl13b*^{R79Q} project normally to both the dorsal and ventral thalamus. (A) Schematic showing the protein domain structure of Arl13b. The R79Q mutation occurs in a highly conserved subregion of the GTPase domain, Switch 2 (red). (B) Alignment of protein sequence surrounding amino acid 79 of mouse Arl13b (red highlight) showing that arginine is conserved across multiple species. Sequences used in protein alignment: *Homo sapiens* NP_878899.1, *Mus musculus* NP_080853.3, *Danio rerio* NP_775379.1, *Chlamydomonas reinhardtii* XP_001691430.1. (C-F) Representative images of dorsal (C, E) or ventral (D, F) thalamus injection site (top panel) and cerebellum (middle panel) with the DCN in hatched white circle and magnified (bottom panel) with recoloring to black and white to aid visualization. Numbers indicate the number of positively stained DCN clusters out of the total number of injected animals. Note that no tracing was observed on the injection's ipsilateral side.

(C, D) Fluorescent tracer injection in *Arl13b*^{R79Q/+} control animals resulted in contralateral DCN staining in (C) 4/4 dorsal thalamus injections and (D) 3/3 ventral thalamus injections. (E-F) Fluorescent tracer injection in *Arl13b*^{R79Q/R79Q} animals resulted in contralateral DCN staining in (E) 3/3 dorsal thalamus injections (Fisher's exact test, two-tailed, ns, p>0.9) and (F) 3/4 ventral thalamus injections (ns, p>0.9).

Figure 4: Mouse embryos expressing JS allele *Arl13b*^{R79Q} display normal cell patterning in the neural tube. Shh, Olig2, Pax6 and *Arl13b* staining of E10.5 *Arl13b*^{+/+} (n=3), *Arl13b*^{R79Q/R79Q} (n=3), *Arl13b*^{R79Q/Δ} (n=3), and *Arl13b*^{Δ/Δ} (n=3) mouse neural tubes. Scale bar = 100 micrometers in A-L or 10 micrometers in M-P. (A-D) Shh is visible in the notochord and floorplate of (A) *Arl13b*^{+/+}, (B) *Arl13b*^{R79Q/R79Q} and (C) *Arl13b*^{R79Q/Δ} neural tubes but absent from the floorplate of (D) *Arl13b*^{Δ/Δ} embryos. (E-H) Olig2 stains the motor neuron precursor domain in (E) *Arl13b*^{+/+}, (F) *Arl13b*^{R79Q/R79Q} and (G) *Arl13b*^{R79Q/Δ} embryos and stains an expanded domain in (H) *Arl13b*^{Δ/Δ} embryos. (I-L) Pax6 expression is visible in the dorsal neural tube in (I) *Arl13b*^{+/+}, (J) *Arl13b*^{R79Q/R79Q} and (K) *Arl13b*^{R79Q/Δ} neural tubes but shifted dorsally in (L) *Arl13b*^{Δ/Δ} neural tubes. (M-P) *Arl13b* is localized to cilia visible in the ventral lumen of (M) *Arl13b*^{+/+}, (N) *Arl13b*^{R79Q/R79Q} and (O) *Arl13b*^{R79Q/Δ} neural tubes but absent from (P) *Arl13b*^{Δ/Δ} embryos.

Figure 5: *Arl13b*^{R79Q} mutation does not affect cerebellar width. (A) Representative surface-facing images of cerebella from control and *Arl13b*^{R79Q/R79Q} mutant mice. Scalebar = 1mm, black line indicates where cerebellar width was measured, red line indicates where vermis width was measured, gray line indicates where cerebellar

hemisphere height was measured, blue line indicates where vermis height was measured. (B) Ratios of *Arl13b*^{R79Q/R79Q} mutant to control measurements of sex- and age-matched pairs of mice showed no significant difference. Cerebellar and vermis measurements were determined from surface views (symbol colors match the lines in A), body and dissected brain weights were determined using a standard lab scale. Each symbol represents a pair of sex- and age-matched animals: females are represented as circles and males are represented as triangles.

Figure 6: Pan-neuronal deletion of *Arl13b* results in a small cerebellum. (A)

Representative images of cerebella from control and *Arl13b*^{Bm4-Cre} mutant mice. Scalebar = 1mm. (B) The ratio of *Arl13b*^{Bm4-Cre} mutant to control cerebellar width was reduced in both females and males (one-sample t-test: p<0.1, single asterisk). The ratio of *Arl13b*^{Bm4-Cre} mutant to control vermis width was reduced in female and male age matched pairs (one-sample t-test: p<0.0001, four asterisks). While *Arl13b*^{Bm4-Cre} mutants had moderately lower body weights than controls (one-sample t-test: p<0.1, single asterisk), their brains were slightly heavier, likely due to hydrocephaly. Each symbol represents a pair of sex- and age-matched animals: females are represented as circles and males are represented as triangles.

Figure 7: Mice expressing cilia-excluded *Arl13b*^{V358A} have cerebella of normal

width. (A) Representative images of cerebella from control and *Arl13b*^{V358A/V358A} mutant mice. Scalebar = 1mm. (B) Ratios of *Arl13b*^{V358A/V358A} mutant to control measurements of sex- and age-matched pairs of mice showed no significant differences (one-sample t-

952 test). Each dot represents a pair of sex- and age-matched animals: females are
953 represented as circles and males are represented as triangles.
954