1 **Main Manuscript for**

2 **Sleep Loss Impacts the Interconnected Brain-Body-Mood Regulation**

3 **of Cardiovascular Function in Humans**

4

5 Adam J. Krause¹, Raphael Vallat¹, Eti Ben Simon¹, Matthew P. Walker^{1,2*}

6

7 ¹ Center for Human Sleep Science, Department of Psychology, University of California, Berkeley,
8 California 94720-1650, USA.

9 ² Helen Wills Neuroscience Institute, University of California, Berkeley, California 94720-1650,
10 USA

11 *Corresponding author Matthew P. Walker
12 **Email:** mpwalker@berkeley.edu

13

14

15 **This file includes:**

16 Main Text
17 Figures 1 to 4
18 Supplementary Tables 1 to 2

19

20

21

22

23

Abstract

Poor sleep is associated with hypertension, a major risk factor for cardiovascular disease^{1,2}. However, the mechanism(s) through which sleep loss impacts blood pressure remain largely unknown, including the inter-related brain and peripheral body systems that regulate vascular function³. In a repeated-measures experimental study of 66 healthy adult participants, we demonstrate four core findings addressing this question. First, a night of sleep loss significantly increased blood pressure—both systolic and diastolic, yet this change in vascular tone was independent of any increase in heart rate. Second, sleep loss compromised functional brain connectivity within regions that regulate vascular tone. Third, sleep-loss related changes in brain connectivity and vascular tone were significantly inter-dependent, with changes in brain nodes explaining the shift towards hypertension. Fourth, sleep-loss related changes in mood, specifically reductions in positive and amplification in negative states, each demonstrated an interaction with the impairments in brain connectivity and blood pressure. Together, these findings support an embodied framework in which sleep loss confers increased risk of cardiovascular disease through interactions between brain homeostatic control, mood-state and blood pressure.

Main Text

Introduction

Poor sleep is associated with hypertension, which is a primary risk factor for cardiovascular disease ^{1,2}. Epidemiological studies indicate that both habitual short sleep duration and poor-quality sleep are associated with a greater incidence of hypertension, even when controlling for numerous other risk factors ^{2,4}.

Prospective longitudinal studies have demonstrated that short sleep duration and worse quality sleep predict a higher future risk of hypertension ^{2,5}. Manipulations using partial or total sleep deprivation causally and consistently increase blood pressure ^{6,7}. Such evidence supports a possible link between the coinciding increase in sleep deficiency and increasing rates of cardiovascular disease in numerous first world nations ⁸.

However, the mechanistic pathway(s) through which insufficient sleep increases blood pressure and thus cardiovascular disease risk remains unclear. This is especially true regarding a possible interaction between impaired central brain and peripheral body systems that are known to control vascular blood pressure. Indeed, the brain plays a causal role in the regulation of peripheral physiology, including that of blood pressure and heart rate ³. Core to this control are a discrete set of viscerosensory brain regions that that map and also instigate changes in vascular function.

Defining this cortical and subcortical network have been animal models demonstrating that direct stimulation of the amygdala ⁹, the cingulate ¹⁰, the insula ^{11,12} and medial prefrontal cortex (MPFC) ¹⁰ all elicit changes in blood pressure. In humans, activity within, and connectivity between, the amygdala, cingulate, and insular cortices

aids in the regulation of arterial blood pressure, as well as event-related reactivity of vascular blood pressure ¹³.

Of this collection of regions, the insular cortex and MPFC are of particular importance in the regulation of vascular state. Both regions are involved in the central brain regulation of autonomic state, including the mapping of current autonomic balance and modulating autonomic outflow ^{3,11,14}. Activity in the insula and MPFC have further been linked with increased vasoconstriction in subjects with a history of coronary artery disease ¹⁵, and ischemic injury to the insula induces endothelial dysfunction and inflammation ¹⁶. Similarly, lesioning of the MPFC increases plasma levels of adrenocorticotrophic hormone and corticosterone ¹⁷, both important stress-response factors which can impact cardiovascular function ¹⁸. The insula and MPFC are unique in the regulation of cardiovascular function through direct connections to brainstem areas ^{19,20}. Finally, the brain, including the insula and MPFC, can further impact vascular function through its influence on immune function as vascular endothelial cells are sensitive to inflammatory processes ^{21,22}.

Although the peripheral autonomic nervous system is a known regulatory pathway for vascular and endothelial function control ²³, how the autonomic nervous system interacts with these central brain viscerosensory networks under conditions of sleep loss, resulting in a shift towards hypertension, remains uninvestigated.

Building on this collective evidence, and addressing this unresolved question, here, we sought to test the hypothesis that one pathway through which insufficient sleep instigates a shift from normotensive towards a hypertensive increase in blood pressure is through an altered relationship with central functioning of brain vascular control networks and the peripheral body cardiovascular system. Specifically, we tested the

prediction that elevations in blood pressure caused by a night of sleep deprivation are significantly associated with changes in the functional connectivity state of the above-described a priori vascular control networks of the human brain, indicative of a brain-body mechanistic inter-relationship.

In short (and see Methods for details), 66 healthy adult participants who were otherwise normotensive, were enrolled in a counterbalanced, repeated-measures design involving two conditions: one night of sleep (Sleep Rested) and one night of sleep loss (Sleep Deprivation). In each condition, participants had their cardiovascular state, including blood pressure and heart rate, assessed at circadian-matched morning timepoints, and received a resting-state functional magnetic resonance imaging (fMRI) scan following sleep or sleep deprivation—the latter being a recognized method for mapping the state of network connectivity of the human brain ²⁴.

Results

Sleep Loss & Cardiovascular Function

We first tested the hypothesis that sleep deprivation significantly alters key vascular, cardiac and autonomic metrics linked with cardiovascular disease²⁵: systolic and diastolic blood pressure, heart rate, and heart rate variability.

Supportive of the hypothesis and prior findings^{1,2,4,5}, sleep deprivation significantly increased systolic blood pressure, relative to the same individuals under sleep rested conditions (**Fig. 1A**; Systolic: sleep-rested mean = 107.6 mmHg ± 11.8 SD, sleep-deprived mean = 112.1 mmHg ± 12.8 SD, $p = 0.003$). This was similarly true of

Figure 1. Cardiovascular outcomes. **A**, Following sleep deprivation, there was an increase in both systolic and diastolic blood pressure, relative to the sleep rested condition. **B**, Sleep deprivation resulted in a reduction in heart rate, relative to the sleep-rested condition. **C**, Sleep deprivation resulted in an increase in heart rate variability (pNN50), relative to the sleep-rested condition. Error bars indicate SEM. * $p < 0.05$.

114 diastolic blood pressure, which also increased significantly following sleep loss (**Fig. 1A**;
 115 Diastolic: sleep-rested mean = 71.1 mmHg $\pm$ 7.3 SD, sleep-deprived mean = 73.7
 116 mmHg $\pm$ 7.9 SD, $p = 0.03$).

117 These changes in blood pressure occurred in the absence of an increase in heart
 118 rate. Specifically, there was an overall reduction of resting heart rate under conditions of
 119 sleep loss, relative to the sleep rested condition (**Fig 1B**; sleep-rested mean = 67.5 bpm
 120 $\pm$ 11.6 SD, sleep-deprived mean = 62.9 bpm $\pm$ 10.2 SD, $p < 0.001$), a finding that is
 121 consistent with previous reports regarding sleep loss^{6,26-29}, but see³⁰.

122 Corresponding to the decrease in heart rate, there was an increase in heart rate
 123 variability under conditions of sleep deprivation, relative to the sleep-rested condition
 124 (**Fig 1C**; pNN50: sleep-rested mean = 27.5% $\pm$ 20.5% SD, sleep-deprived mean =

42.0% $\pm$ 19.1%, $p < 0.0001$). Since the pNN50 reflects vagal modulation of the heart³¹, this observed increase in heart rate variability under conditions of sleep loss is consistent with increased parasympathetic input to the heart linked to the state of extended wakefulness^{26,28}.

Further advancing a dissociation between increased peripheral vascular blood pressure in the absence of accelerated heart rate, there was no significant association between the differential changes in blood pressure and heart rate between conditions ([sleep-rested] – [sleep deprivation]) for either systolic or diastolic measures (**Fig 2A,B**; Systolic: $r = 0.06$, $p = 0.61$; Diastolic: $r = 0.03$, $p = 0.84$). A similar lack of association was observed between the sleep-loss changes in heart rate variability and in blood pressure (**Fig 2C,D**; Systolic: $r = -0.09$, $p = 0.48$; Diastolic: $r = -0.05$, $p = 0.69$). In contrast, there was a significant association between the sleep-loss related reduction in heart rate and the corresponding increase in heart rate variability ($r = -0.48$, $p < 0.0001$).

Figure 2. Associations between cardiovascular outcomes. Scatterplots represent non-significant correlations between sleep-loss changes (deprived – rested) in cardiovascular outcomes, including **(A)** heart rate and systolic blood pressure, **(B)** heart rate and diastolic blood pressure, **(C)** heart rate variability (pNN50) and systolic blood pressure, and **(D)** heart rate variability (pNN50) and diastolic blood pressure.

138

139 Sleep Loss & Viscerosensory Resting Brain Connectivity

140 Next, we assessed functional connectivity within the a priori network known to be
141 associated with cardiovascular control in humans and non-human primates, including

142 the amygdala, anterior cingulate, insula, and ventromedial and medial prefrontal
 143 cortices^{3,9-13}. Within this network, twelve connections were found to be significantly
 144 altered following FDR-correction for multiple tests under conditions of sleep deprivation,
 145 relative to the sleep rested state ($q = 0.05$)³². These included connectivity pathways
 146 between the insula and anterior cingulate, the insula and MPFC, the anterior cingulate
 147 and the MPFC, the anterior cingulate and the amygdala, the ventromedial frontal and
 148 MPFC, and MPFC and amygdala (**Fig 3A,B, Table S1**).

Figure 3. Changes in brain connectivity in regions of interest. Connections shown are significant following FDR-correction ($q = 0.05$). **A**, Connections that are significantly increased in the sleep deprived condition relative to the sleep-rested condition. **B**, Connections that are significantly decreased in the sleep-deprived condition relative to the sleep rested condition.

149 Of these twelve, there was a significant reduction in connectivity between the
150 ventromedial frontal and MPFC following sleep deprivation, relative to a night of normal

sleep ($t(65) = 3.2$, $p = 0.008$, FDR-corrected, **Fig. 3B**). The remaining eleven region of interest (ROI) pairs demonstrated a significant increase in their connectivity (**Fig. 3A**, **Table S1**), with the connectivity changes all jointly sharing at least one of two nodes: the insula and MPFC. First, the insula had significantly increased connectivity with the anterior cingulate cortex ($p = 0.01$, FDR-corrected). Second, there were increases in connectivity between the insula and the MPFC ($p < 0.0008$, FDR-corrected). The MPFC demonstrated increased connectivity to both the anterior cingulate in the right hemisphere ($p < 0.0001$, FDR-corrected), and amygdala in both hemispheres ($p < 0.02$, FDR-corrected). Details of these connectivity changes and their significance are described in **Supplement**.

Sleep Loss, Cardiovascular Function & Viscerosensory Resting Brain Connectivity

We next tested whether these two sets of identified changes—alterations in brain connectivity and alterations in peripheral blood pressure were significantly inter-related. First, greater sleep-loss-related increases in insula connectivity were significantly associated with a greater rise in systolic blood pressure, especially between the insula and the MPFC ($r(60) = 0.25$; $p = 0.05$; **Fig. 4A**). Second, sleep-loss-related decreases in MPFC connectivity with the amygdala was significantly related to increases in diastolic blood pressure ($r(61) = -0.29$, $p = 0.02$; **Fig. 4B**). Therefore, sleep-loss changes in systolic and diastolic blood pressure were associated with alterations in brain connectivity, with a clustering around changes in insular and MPFC connectivity, respectively. Unlike the association between brain and blood pressure, no associations

Figure 4. Brain connectivity and cardiovascular changes. **A**, Scatterplot represents significant positive correlation between sleep-loss related increases in insula and MPFC connectivity and sleep-loss related increases in systolic blood pressure. **B**, Scatterplot represents significant negative correlation between sleep-loss increases in MPFC and amygdala connectivity and sleep-loss related increases in diastolic blood pressure.

174 between sleep-loss related changes in heart rate or heart rate variability and changes in
175 functional connectivity were observed (all $p > 0.05$).

To better understand the specificity of central brain changes and cardiovascular changes, post-hoc analyses examined associations between sleep-loss related alterations in four non-a-priori, though well-recognized, larger-scale resting-state brain networks: the Default-Mode Network (DMN), Dorsal Attention Network (DAN), Ventral Salience Network (VSN), and Somatomotor Network (SMN)³³. Neither differences in connectivity within or between these large-scale resting-state networks showed significant associations with the sleep-loss related increases in blood pressure, the decrease in heart rate, or the increase in heart rate variability (all $p > 0.05$). Therefore, local changes within the specific a priori cardiovascular control network^{3,9-13}, rather than global changes in larger-scale brain networks³³, best accounted for changes in peripheral blood pressure.

Emotion: Interactions with Brain Connectivity & Cardiovascular Function

Independent of sleep, mood and anxiety have well-recognized relationships with peripheral autonomic activity, including changes in cardiac and vascular activity³⁴. Mood states and anxiety have also been related to changes in activity and functional connectivity of some nodes of the cardiovascular control network, specially the insular and MPFC nodes that are central to emotional brain processes³⁵⁻³⁷. Furthermore, both positive and negative mood states are deleteriously impacted by sleep loss³⁸.

Building on these inter-connected links, we finally tested the hypothesis that changes in mood and anxiety caused by sleep loss would be significantly associated with, rather than independent of, the sleep-loss related changes in cardiovascular state and cardiovascular control brain networks.

As expected^{38,39}, sleep deprivation was associated with a worsening of mood: a reduction in positive mood ($t(65) = 6.8$, $p < 0.0001$) and an increase in negative mood ($t(65) = -5.2$, $p < 0.0001$), relative to the sleep rested condition. Sleep loss also resulted in an increase in state anxiety, relative to the sleep rested state ($t(65) = -4.3$, $p < 0.0001$).

Demonstrating affective specificity, sleep-loss increases in negative mood state were associated with sleep-loss increases in blood pressure, such that the greater the negative mood increase, the greater the increase in diastolic blood pressure, though this was not significant following FDR correction ($r(59) = 0.32$, $p = 0.18$; FDR-corrected). Moreover, sleep-loss changes in heart rate and heart rate variability were not associated with changes in any of the affective measures, (all $p > 0.24$, FDR-corrected), once again demonstrating specificity to changes in vascular control, rather than heart contractility rate. The association between negative mood and systolic blood pressure was not significant ($r(60) = 0.12$, $p = 0.51$, FDR-corrected).

Sleep-loss related reduction in positive mood was instead significantly associated with the sleep-loss change in connectivity between the left MPFC and ventromedial frontal cortex ($r(65) = 0.32$, $p = 0.01$; and trend levels with changes in left insula cortex and right MPFC connectivity ($r(65) = -0.24$, $p = 0.06$). Connectivity changes in the MPFC and insula regions also demonstrated significant associations with sleep-loss related changes in systolic and diastolic blood pressure, fitting with the roles of these areas in both affective and cardiovascular regulation^{3,11,14,35-37}. No associations were evident for changes in negative mood (all $p > 0.26$).

Counter to our experimental hypothesis, however, sleep-loss changes in anxiety were not significantly associated with changes in a priori brain vascular control networks (**Table S2**).

Therefore, a dissociable influence of mood was observed, with sleep-loss amplifications in negative mood state selectively associated with the amplifications in peripheral body blood pressure, and the loss of positive mood state was expressly associated with impairments in viscerosensory brain connectivity. Such findings lend further support to an embodied framework of sleep loss in which changes in brain, body, and mental state are mutually related in their association with the sleep-loss induced shift towards hypertension.

Discussion

Taken together, the current study demonstrates that (1) experimental sleep loss significantly and selectively increases peripheral blood pressure, independent of any increase in heart rate, (2) sleep loss simultaneously compromises functional brain connectivity within regions that help regulate vascular tone—the viscerosensory network, (3) the sleep-loss related changes in viscerosensory functional brain connectivity and in peripheral vascular tone were significantly inter-dependent, with the changes in brain nodes explaining the vascular shift towards hypertension, and (4) sleep-loss-related reductions in positive state and amplification in negative state each showed a significant additive interactions with the selective impairments in brain network activity and peripheral hypertension caused by sleep loss, consistent with mental-state influence.

Sleep Loss and Cardiovascular State

Chronic sleep disturbances, including sleep disorders such as insomnia and sleep apnea, are recognized risk factors for hypertension and cardiovascular disease^{1,2,4,5}. Moreover, experimental sleep loss causally increase blood pressure^{6,7}. However, the underlying mechanism(s) by which poor sleep increases blood pressure are poorly understood.

The heart is known to receive both sympathetic and parasympathetic input, yet most vessels (arteries and veins) only receive sympathetic innervation⁴⁰. Based on the dissociation observed, one parsimonious mechanism is that increases in blood pressure caused by sleep deprivation are triggered by an increase in sympathetic tone, while the decrease in heart rate reflects a separable shift in cardiac-mediated autonomic balance towards the influence of parasympathetic input⁴¹. That is, an autonomic dissociation caused by sleep loss, wherein increased sympathetic output dominates in peripheral vessels, yet the opposite is true in the heart. This possibility is also supported by the sleep-loss related increase in heart rate variability—a measure reflecting vagal (parasympathetic) modulation of the heart³¹. Indeed, the high sleep pressure resulting from sleep deprivation increases parasympathetic cardiac drive slowing heart rate^{26,28}.

Sleep Loss & Resting Brain Connectivity

Peripheral functions of the body, including the regulation of the autonomic nervous system, are causally influenced by a set of brain regions involved in viscerosensory and cardio-modulatory processes^{3,14,42}. The second component of our experimental hypothesis sought to determine whether changes in connectivity within specific brain networks that regulate cardiovascular state occurred under conditions of

sleep deprivation, and whether such changes were related to the observed changes in cardiovascular function.

At a main effects level, sleep loss resulted in a collection of connectivity changes within our a priori set of brain regions, the majority of which showed increases in their nodal connectivity. These connectivity changes all jointly shared at least one of two nodes: the insula and MPFC. The insular cortex is not only sensitive to sleep status^{43,44}, but is of central importance in the registration and control of bodily states, including cardiovascular state^{3,11-13}. The MPFC, as with the insula, is also sensitive to sleep quality and quantity^{39,45-47}, and orchestrates cardiovascular-challenge responses¹³.

Our findings corroborate and extend previous reports, indicating that sleep-loss-related changes in insula and MPFC connectivity may reflect impairment of brain networks involved in the coordination of cardiovascular physiology. Fitting this proposition, these brain regions are highly interconnected with several cortical, subcortical, and brainstem regions that dictate autonomic activity^{3,11-14}.

In contrast to the dominant increases in connectivity caused by sleep loss, one pathway showed a decrease in connectivity under conditions of sleep loss: the link between the MPFC and ventromedial frontal cortex. Both regions have been previously associated with sympathetic nervous system drive and the central autonomic network of the brain¹⁴. Lesioning of the MPFC is associated with increased hypothalamic-pituitary-adrenal axis stress responses¹⁷, which is of central importance for the integration of brain and vascular function¹⁸. Furthermore, reduced activation in the MPFC has been associated with increased stress-induced vasoconstriction¹⁵.

Sleep Loss, Cardiovascular State & Resting Brain Connectivity

Adding to the findings of sleep-loss related changes in cardiovascular state and brain-network connectivity, we further demonstrate that these changes are significantly associated. First, the increase in systolic blood pressure was associated with increased connectivity between two regions involved in the regulation of vascular tone: the insula and MPFC. Both the insula and MPFC are involved in the representation and generation of visceral states, including cardiovascular reactions⁴⁸. Thus, these two regions, along with a network of corticolimbic and brainstem systems, play instrumental roles in tuning cardiovascular and autonomic adjustments in response to contextual demands^{13,42,48}. Indeed, the insula and MPFC are multi-synaptically connected to pre-autonomic nuclei which innervate both the heart and vasculature^{49,50}. There is also evidence that increased resting activity in the insula and MPFC predicts greater evoked blood pressure reactions, interpreted as signaling increased sympathetic-mediated blood pressure reactivity⁴⁸.

A different part of the cardiovascular control network was observed to be associated with the sleep-loss increase in diastolic blood pressure. Here, the magnitude of sleep-loss increase in diastolic blood pressure was significantly associated with decreased connectivity between the MPFC and amygdala. The amygdala is instrumental in coordinating behavioral and physiological adjustments through connections to cortical, hypothalamic, and brainstem nuclei involved in stress processing and autonomic-cardiovascular control⁴⁸. Additionally, the amygdala can impact blood pressure through influence over the baroreflex, which serves to maintain blood pressure through autonomic regulation of heart rate, cardiac output, and vascular resistance, and is associated with preclinical atherosclerosis^{9,51}. Thus, the reduction in amygdala

connectivity to other regions involved in cardiovascular control (MPFC) could represent a degree of disorganization of brain homeostatic functions.

The vascular system is subject behavioral, autonomic, and endocrine regulation⁵². The complexity of these interacting pathways suggests that altered blood pressure control is due to a failure of a central control system, namely the brain⁵³. Notably, in the current data, the decrease in heart rate was not associated with the increase in blood pressure. Furthermore, blood pressure-related changes in functional connectivity in the brain were similarly not associated with heart rate. Taken together, it is likely that multiple, perhaps counteractive, processes underlie the conflicting vascular and cardiac responses to sleep loss, fitting with a loss of central brain regulation.

Sleep Loss, Mood States, Brain, and Body

The final component of the study examined inter-relationships with changes in mood state and anxiety. Independent lines of research have demonstrated that sleep-loss detrimentally impacts affective states, including mood³⁸. Moreover, mood states are known to causally alter cardiovascular function, including peripheral blood pressure³⁴. In addition, the regulation of positive and negative mood states is further controlled by overlapping brain regions to those that are known to form the basis of central brain cardiovascular control, notably the insula and MPFC³⁵⁻³⁷.

Fitting with prior findings^{38,39}, sleep loss significantly worsened mood state through a reduction in positive mood and an increase in negative mood. We extend these findings, establishing that sleep-loss related increases in negative mood are further related to physiological changes within the body, here the shift towards hypertension.

Indicating potential mood-type specificity, we further show that, this affective link to visceral function was specific to peripheral-body vascular tone and was significant for the increase in negative mood, and not the decrease in positive mood. Instead, the link to visceral function and the sleep-loss reduction in positive mood was specific to the changes in functional brain connectivity, notably the viscerosensory network regions of the insula and MPFC.

These latter finding may therefore suggest a dissociable influence of affective mental state on vascular regulation under conditions of sleep loss: negative mood state linked to peripheral body vascular function, yet the loss of positive mood state linked to central brain visceral regulation. That there is significant overlapping functionality of the MPFC and insula in both emotional and viscerosensory brain regulation³⁵⁻³⁷ further suggests a convergence of regulatory function in these brain regions underpinning an embodied framework of sleep loss, one in which sleep deprivation impacts affective and cardiovascular function through impairment of core regulatory centers in the brain.

More generally, our results indicate the presence of a coupled relationship between vascular responses to sleep loss, brain function and mental-state changes in affect. Such findings impress the previously proposed conception of sleep loss triggering broad homeostatic biological distress caused by allostatic overlap⁵⁴. Pragmatically, these findings highlight the target of sleep as a risk factor, and modifiable intervention target, determining cardiovascular disease risk, here through regulation of central brain and mental-state mechanisms that control cardiovascular function with the goal of maintaining vascular tone within a normotensive range.

Methods and Materials

Experimental Model and Subject Details

Sixty-six healthy adults aged 18-24 years (mean $\pm$ SD: 20.7 $\pm$ 1.7, 52% female) completed a repeated-measures cross-over design (described below). There was no influence of sex on cardiovascular outcomes ($p > 0.64$). Participants abstained from caffeine and alcohol for 72 hours before and during the entire course of the study and kept a normal sleep-wake rhythm (7-9 hours of sleep per night) with sleep onset before 1:00 A.M. and rise no later than 9:00 A.M. for the three nights before the study participation, as verified by sleep logs and actigraphy.

Exclusion criteria, assessed using a prescreening semi-structured interview, were as follows: a history of previously-diagnosed sleep disorders, neurological disorders, closed head injury, Axis 1 psychiatric disorders according to the DSM-IV-TR criteria (encompassing mental disorders, including depression, anxiety disorders, bipolar disorder, attention deficit disorder, and schizophrenia), history of drug abuse and current use of antidepressant or hypnotic medication, nicotine use, consumption of more than five alcoholic drinks per week, crossing of time zones in the 3 months before the study, and general contraindications to MRI. Participants also had blood pressures within normotensive ranges, confirmed with a blood pressure reading of less than 120/80 mmHg at the time of consenting⁵⁵. Participants who reported sleeping <7 hours per night or consuming three or more daily caffeine-containing drinks were also excluded from entering the study. The Pittsburgh Sleep Quality Index⁵⁶ was employed to determine the quality of recent sleep history, and participants considered poor sleepers (global score > 5) were excluded. The study was approved by the UC Berkeley Committee for the Protection of Human Subjects, with all participants providing written consent.

385

386 **Experimental Design**

387 To test the experimental hypotheses, participants entered a repeated-measures
388 study design (**Fig 1**), including two sessions—one following a normal night of sleep and
389 one after 24 hours of total sleep deprivation. The two sessions were separated by at
390 least 7 days (mean $\pm$ SD: 9.8 ± 3.8), with the order of the sleep-rested and deprived
391 conditions counterbalanced across subjects.

392 In the sleep deprived session, participants arrived at the laboratory at 10:00 P.M.
393 and were continuously monitored throughout the enforced waking period. Activities
394 during the sleep deprivation condition were limited to use of the internet, email, short
395 walks, reading, movies of low emotionality, and playing board games, thereby providing
396 a standardized regimen of waking activity without undue stress. Participants also
397 avoided exercise during the experimental sessions. In the sleep-rested session,
398 participants came to the laboratory at 8:00 P.M. and were prepared for an 8-hour night of
399 sleep (~11:00 P.M. to 7:30 A.M. $\pm$ 30 min). In both conditions, mood and anxiety were
400 assessed in the morning at the time of the cardiovascular assessment using the Positive
401 and Negative Affective Scale and State-Trait Anxiety Inventory questionnaires,
402 respectively^{57,58}. The following morning, participants' cardiovascular state was measured
403 followed by fMRI scanning.

404

405 **Cardiovascular Assessments**

In each condition, cardiovascular state was assessed to determine blood pressure, heart rate, and heart rate variability. These assessments were performed approximately 1 hour after awakening, and at circadian-matched times in the sleep rested and sleep deprivation conditions for each participant.

Blood pressure measurement was performed with Omron BP742 blood pressure monitor (Omron Healthcare Europe BV, Hoofddorp, The Netherlands), fitted to the participants' nondominant arm over the brachial artery. Participants were instructed to remain motionless with open eyes while sitting with feet flat on the floor during the recording with rested cuffed arm at heart height, consistent with standard blood pressure recording guidelines⁵⁹. Two consecutive measurements were taken and the mean of these two measurements was used for analyses.

Immediately following blood pressure assessment, participants were fitted with electrodes for bipolar electrocardiography (ECG), and cardiac activity was recorded for five minutes. Electrodes were applied in a lead II format with the reference electrode placed below the right clavicle parallel to the right shoulder and a second electrode placed on the torso at the fourth intercostal space on the left side parallel to the left hip. Similar to the blood pressure recording, participants sat motionless with open eyes while sitting with feet flat on the floor. All data were screened for measurement artifacts and beat-to-beat intervals extracted using ARTiiFACT software⁶⁰, and all flagged interbeat intervals were visually checked. If confirmed as artifactual, they were deleted and substituted by means of cubic spline interpolation of neighboring intervals. The time-domain measure pNN50 was calculated for each participant, which is the percentage of differences between adjacent interbeat intervals (NN) that are greater than 50ms and represent a recommended measure of vagally-mediated heart rate variability³¹.

430

431 **fMRI preprocessing**

432 Preprocessing was performed using fMRIPrep 1.2.5 (RRID:SCR_016216). The
433 T1-weighted image was corrected for intensity non-uniformity using
434 N4BiasFieldCorrection and used as a reference throughout the analysis. Spatial
435 normalization to the ICBM 152 Nonlinear Asymmetrical template was performed through
436 nonlinear registration with antsRegistration using brain-extracted version of both T1-
437 weighted volume and template (ANTs 2.2.0 RRID:SCR_004757). Brain tissue
438 segmentation of cerebrospinal fluid (CSF), white-matter (WM) and gray-matter (GM) was
439 performed on the brain-extracted T1-weighted image using fast (FSL 5.0.10,
440 RRID:SCR_002823).

441 For fMRI data, the mean BOLD image was co-registered to the T1-weighted
442 reference using flirt. Head-motion parameters with respect to the BOLD reference
443 (transformation matrices, and six corresponding rotation and translation parameters)
444 were estimated before any spatiotemporal filtering using mcflirt. BOLD runs were slice-
445 time corrected using 3dTshift from AFNI (RRID: SCR_005927), and the BOLD time-
446 series (including slice-timing correction when applied) were resampled to
447 MNI152NLin2009cAsym standard space. Known time-series factors requiring covariate
448 accommodation were calculated based on the preprocessed BOLD: framewise
449 displacement (FD), DVARS and three region-wise global signals. The three global
450 signals are extracted within the CSF, the WM, and the whole-brain masks. Additionally,
451 a set of physiological regressors were extracted to allow for component-based noise
452 correction (CompCor⁶¹). Finally, regression was used to estimate and remove the whole-

brain global signal time series, and functional images were smoothed using a 6 mm FWHM Gaussian kernel.

Statistical and fMRI analysis

Connectivity analyses were performed using the CONN toolbox version 19b (www.nitrc.org/projects/conn, RRID:SCR_009550). First, denoising was performed by regressing the six motion parameters, FD, and physiological noise components. The resulting BOLD time series was band-pass filtered (0.008 – 0.09 Hz), followed by region of interest (ROI) analysis, specifically ROI-to-ROI connectivity analysis.

Based on the experimental hypotheses, analyses focused on an a priori set of brain ROIs known to be involved central brain cardiovascular control^{3,9-14}. These ROIs were drawn from anatomical atlases (Harvard-Oxford Cortical and Subcortical Atlas) included in the CONN toolbox, and were extracted using an ICA approach. For each ROI, the connectivity to all other ROIs was calculated for each subject in each condition separately, generating a symmetric correlation matrix wherein each element is the correlation between two ROIs in each condition and each participant.

In addition to ROI-to-ROI connectivity analyses, control analyses for specificity examined connectivity within and between additional prototypical brain networks of the Default-Mode Network (DMN), Dorsal Attention Network (DAN), Ventral Salience Network (VSN), and Somatomotor Network (SMN)³³. Within-network connectivity was calculated for each network by extracting the mean resting-state BOLD time-series from each voxel within a region of a network and calculating the mean Pearson correlation with all other regions of the network. Between-network connectivity was similarly

calculated for each network by extracting the mean resting-state BOLD time-series from every voxel within a network, and then calculating the correlation with the mean time-series of all other networks.

The resulting correlation coefficients from these fMRI analyses across participants in each condition were then statistically compared (two-tailed paired t-test) to determine the effect of sleep loss on brain connectivity. Consistent with imaging recommendations to correct for multiple statistical comparisons among the eight ROIs and four networks, the Benjamini-Hochberg procedure for controlling FDR was applied, with $q = 0.05^{32}$. This procedure addresses the problem of simultaneously performing multiple significance tests by using an adaptive stepwise procedure to control the expected ratio of erroneous rejections to the number of total rejections. Only connectivity changes that survived the FDR criterion after correction were considered statistically significant and used in analyses. Pearson correlations were employed to test the relationships between sleep loss-related changes in brain connectivity and sleep loss-related changes in cardiovascular state. All statistical analyses were performed in Python using the Pingouin package⁶².

Acknowledgments: We thank Olivia G. Murillo for her assistance in running the study.

Author Contributions: AJK, RV, EBS, and MPW designed research. AJK, RV, and EBS performed research. AJK analyzed data. AJK and MPW wrote the paper with contributions from all authors.

498 **Declaration of Interests:** Dr. Walker serves as a consultant for and has equity interest
499 in Bryte, Oura, Shuni, and StimScience.

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

References

1. Stokes 3rd J, Kannel WB, Wolf PA, D'Agostino RB, Cupples LA. Blood pressure as a risk factor for cardiovascular disease. The Framingham Study--30 years of follow-up. *Hypertension*. 1989;13(5_supplement):I13.
2. Gangwisch JE. A review of evidence for the link between sleep duration and hypertension. *American journal of hypertension*. 2014;27(10):1235-1242.
3. Critchley HD, Mathias CJ, Dolan RJ. Neuroanatomical basis for first-and second-order representations of bodily states. *Nature neuroscience*. 2001;4(2):207-212.
4. Pepin J-L, Borel A-L, Tamisier R, Baguet J-P, Levy P, Dauvilliers Y. Hypertension and sleep: overview of a tight relationship. *Sleep medicine reviews*. 2014;18(6):509-519.
5. Jackowska M, Steptoe A. Sleep and future cardiovascular risk: prospective analysis from the English Longitudinal Study of Ageing. *Sleep medicine*. 2015;16(6):768-774.
6. Kato M, Phillips BG, Sigurdsson G, Narkiewicz K, Pesek CA, Somers VK. Effects of sleep deprivation on neural circulatory control. *Hypertension*. 2000;35(5):1173-1175.
7. Lusardi P, Mugellini A, Preti P, Zoppi A, Derosa G, Fogari R. Effects of a restricted sleep regimen on ambulatory blood pressure monitoring in normotensive subjects. *American journal of hypertension*. 1996;9(5):503-505.
8. Wolk R, Gami AS, Garcia-Touchard A, Somers VK. Sleep and cardiovascular disease. *Current problems in cardiology*. 2005;30(12):625-662.
9. Dampney R. Functional organization of central pathways regulating the cardiovascular system. *Physiological reviews*. 1994;74(2):323-364.

- 549 10. Pool JL, Ransohoff J. Autonomic effects on stimulating rostral portion of cingulate
550 gyri in man. *Journal of neurophysiology*. 1949;12(6):385-392.
- 551 11. Yasui Y, Breder CD, Safer CB, Cechetto DF. Autonomic responses and efferent
552 pathways from the insular cortex in the rat. *Journal of Comparative Neurology*.
553 1991;303(3):355-374.
- 554 12. Oppenheimer SM, Gelb A, Girvin JP, Hachinski VC. Cardiovascular effects of
555 human insular cortex stimulation. *Neurology*. 1992;42(9):1727-1727.
- 556 13. Gianaros PJ, Sheu LK, Matthews KA, Jennings JR, Manuck SB, Hariri AR.
557 Individual differences in stressor-evoked blood pressure reactivity vary with
558 activation, volume, and functional connectivity of the amygdala. *Journal of*
559 *Neuroscience*. 2008;28(4):990-999.
- 560 14. Valenza G, Sclocco R, Duggento A, et al. The central autonomic network at rest:
561 uncovering functional MRI correlates of time-varying autonomic outflow.
562 *Neuroimage*. 2019;197:383-390.
- 563 15. Shah A, Chen C, Campanella C, et al. Brain correlates of stress-induced
564 peripheral vasoconstriction in patients with cardiovascular disease.
565 *Psychophysiology*. 2019;56(2):e13291.
- 566 16. Balint B, Jaremek V, Thorburn V, Whitehead SN, Sposato LA. Left atrial
567 microvascular endothelial dysfunction, myocardial inflammation and fibrosis after
568 selective insular cortex ischemic stroke. *International journal of cardiology*.
569 2019;292:148-155.
- 570 17. Diorio D, Viau V, Meaney MJ. The role of the medial prefrontal cortex (cingulate
571 gyrus) in the regulation of hypothalamic-pituitary-adrenal responses to stress.
572 *Journal of Neuroscience*. 1993;13(9):3839-3847.

- 573 18. Burford NG, Webster NA, Cruz-Topete D. Hypothalamic-pituitary-adrenal axis
574 modulation of glucocorticoids in the cardiovascular system. *International journal*
575 *of molecular sciences*. 2017;18(10):2150.
- 576 19. Fryszak RJ, Neafsey EJ. The effect of medial frontal cortex lesions on
577 cardiovascular conditioned emotional responses in the rat. *Brain research*.
578 1994;643(1-2):181-193.
- 579 20. An X, Bandler R, Öngür D, Price J. Prefrontal cortical projections to longitudinal
580 columns in the midbrain periaqueductal gray in macaque monkeys. *Journal of*
581 *Comparative Neurology*. 1998;401(4):455-479.
- 582 21. Chen P, Chen F, Chen G, et al. Inflammation is associated with decreased
583 functional connectivity of insula in unmedicated bipolar disorder. *Brain, Behavior,*
584 *and Immunity*. 2020;89:615-622.
- 585 22. Vecchiarelli HA, Gandhi CP, Gray JM, Morena M, Hassan KI, Hill MN. Divergent
586 responses of inflammatory mediators within the amygdala and medial prefrontal
587 cortex to acute psychological stress. *Brain, behavior, and immunity*. 2016;51:70-
588 91.
- 589 23. Amiya E, Watanabe M, Komuro I. The relationship between vascular function
590 and the autonomic nervous system. *Annals of Vascular Diseases*. 2014;7(2):109-
591 119.
- 592 24. Van Den Heuvel MP, Pol HEH. Exploring the brain network: a review on resting-
593 state fMRI functional connectivity. *European neuropsychopharmacology*.
594 2010;20(8):519-534.
- 595 25. Benetos A, Rudnichi A, Thomas F, Safar M, Guize L. Influence of heart rate on
596 mortality in a French population: role of age, gender, and blood pressure.
597 *Hypertension*. 1999;33(1):44-52.

- 598 26. Holmes AL, Burgess HJ, Dawson D. Effects of sleep pressure on endogenous
599 cardiac autonomic activity and body temperature. *Journal of Applied Physiology*.
600 2002;92(6):2578-2584.
- 601 27. Franzen PL, Gianaros PJ, Marsland AL, et al. Cardiovascular reactivity to acute
602 psychological stress following sleep deprivation. *Psychosomatic medicine*.
603 2011;73(8):679.
- 604 28. Vaara J, Kyröläinen H, Koivu M, Tulppo M, Finni T. The effect of 60-h sleep
605 deprivation on cardiovascular regulation and body temperature. *European journal*
606 *of applied physiology*. 2009;105(3):439-444.
- 607 29. Zhong X, Hilton HJ, Gates GJ, et al. Increased sympathetic and decreased
608 parasympathetic cardiovascular modulation in normal humans with acute sleep
609 deprivation. *Journal of applied physiology*. 2005;98(6):2024-2032.
- 610 30. Lusardi P, Zoppi A, Preti P, Pesce RM, Piazza E, Fogari R. Effects of insufficient
611 sleep on blood pressure in hypertensive patients: a 24-h study. *American journal*
612 *of hypertension*. 1999;12(1):63-68.
- 613 31. Task Force of ESC. Heart rate variability: standards of measurement,
614 physiological interpretation, and clinical use. *Circulation*. 1996;93(5):1043-1065.
- 615 32. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and
616 powerful approach to multiple testing. *Journal of the Royal statistical society:*
617 *series B (Methodological)*. 1995;57(1):289-300.
- 618 33. Yeo BT, Krienen FM, Sepulcre J, et al. The organization of the human cerebral
619 cortex estimated by intrinsic functional connectivity. *Journal of neurophysiology*.
620 2011.

- 621 34. Räikkönen K, Matthews KA, Flory JD, Owens JF, Gump BB. Effects of optimism,
622 pessimism, and trait anxiety on ambulatory blood pressure and mood during
623 everyday life. *Journal of personality and social psychology*. 1999;76(1):104.
- 624 35. Namkung H, Kim S-H, Sawa A. The insula: an underestimated brain area in
625 clinical neuroscience, psychiatry, and neurology. *Trends in neurosciences*.
626 2017;40(4):200-207.
- 627 36. Fornito A, Malhi GS, Lagopoulos J, et al. Anatomical abnormalities of the anterior
628 cingulate and paracingulate cortex in patients with bipolar I disorder. *Psychiatry*
629 *Research: Neuroimaging*. 2008;162(2):123-132.
- 630 37. Bijsterbosch J, Smith S, Forster S, John OP, Bishop SJ. Resting state correlates
631 of subdimensions of anxious affect. *Journal of Cognitive Neuroscience*.
632 2014;26(4):914-926.
- 633 38. Harvey AG, Murray G, Chandler RA, Soehner A. Sleep disturbance as
634 transdiagnostic: consideration of neurobiological mechanisms. *Clinical*
635 *psychology review*. 2011;31(2):225-235.
- 636 39. Simon EB, Rossi A, Harvey AG, Walker MP. Overanxious and underslept. *Nature*
637 *Human Behaviour*. 2020;4(1):100-110.
- 638 40. Gordan R, Gwathmey JK, Xie L-H. Autonomic and endocrine control of
639 cardiovascular function. *World journal of cardiology*. 2015;7(4):204.
- 640 41. Paton J, Boscan P, Pickering A, Nalivaiko E. The yin and yang of cardiac
641 autonomic control: vago-sympathetic interactions revisited. *Brain Research*
642 *Reviews*. 2005;49(3):555-565.
- 643 42. Critchley HD. Neural mechanisms of autonomic, affective, and cognitive
644 integration. *Journal of comparative neurology*. 2005;493(1):154-166.

- 645 43. Goldstein AN, Greer SM, Saletin JM, Harvey AG, Nitschke JB, Walker MP. Tired
646 and apprehensive: anxiety amplifies the impact of sleep loss on aversive brain
647 anticipation. *Journal of Neuroscience*. 2013;33(26):10607-10615.
- 648 44. Sämann PG, Tully C, Spoormaker VI, et al. Increased sleep pressure reduces
649 resting state functional connectivity. *Magnetic Resonance Materials in Physics,*
650 *Biology and Medicine*. 2010;23(5-6):375-389.
- 651 45. Jiang B, He D, Guo Z, Gao Z. Effect-size seed-based d mapping of resting-state
652 fMRI for persistent insomnia disorder. *Sleep and Breathing*. 2019:1-7.
- 653 46. Greer SM, Goldstein AN, Walker MP. The impact of sleep deprivation on food
654 desire in the human brain. *Nature communications*. 2013;4:2259.
- 655 47. Goldstein-Piekarski AN, Greer SM, Saletin JM, Walker MP. Sleep deprivation
656 impairs the human central and peripheral nervous system discrimination of social
657 threat. *Journal of Neuroscience*. 2015;35(28):10135-10145.
- 658 48. Gianaros PJ, Sheu LK. A review of neuroimaging studies of stressor-evoked
659 blood pressure reactivity: emerging evidence for a brain-body pathway to
660 coronary heart disease risk. *Neuroimage*. 2009;47(3):922-936.
- 661 49. Augustine JR. Circuitry and functional aspects of the insular lobe in primates
662 including humans. *Brain research reviews*. 1996;22(3):229-244.
- 663 50. Öngür D, Price JL. The organization of networks within the orbital and medial
664 prefrontal cortex of rats, monkeys and humans. *Cerebral cortex*. 2000;10(3):206-
665 219.
- 666 51. De Ferrari GM, Sanzo A, Bertoletti A, Specchia G, Vanoli E, Schwartz PJ.
667 Baroreflex sensitivity predicts long-term cardiovascular mortality after myocardial
668 infarction even in patients with preserved left ventricular function. *Journal of the*
669 *American College of Cardiology*. 2007;50(24):2285-2290.

- 670 52. Oparil S, Zaman MA, Calhoun DA. Pathogenesis of hypertension. *Annals of*
671 *internal medicine*. 2003;139(9):761-776.
- 672 53. Jennings JR, Zanstra Y. Is the brain the essential in hypertension? *Neuroimage*.
673 2009;47(3):914-921.
- 674 54. McEwen BS. Sleep deprivation as a neurobiologic and physiologic stressor:
675 allostasis and allostatic load. *Metabolism*. 2006;55:S20-S23.
- 676 55. Whelton PK, Carey RM, Aronow WS, et al. 2017
677 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for
678 the prevention, detection, evaluation, and management of high blood pressure in
679 adults: a report of the American College of Cardiology/American Heart
680 Association Task Force on Clinical Practice Guidelines. *Journal of the American*
681 *College of Cardiology*. 2018;71(19):e127-e248.
- 682 56. Buysse DJ, Reynolds III CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh
683 Sleep Quality Index: a new instrument for psychiatric practice and research.
684 *Psychiatry research*. 1989;28(2):193-213.
- 685 57. Watson D, Clark LA, Tellegen A. Development and validation of brief measures
686 of positive and negative affect: the PANAS scales. *Journal of personality and*
687 *social psychology*. 1988;54(6):1063.
- 688 58. Spielberger CD. State-Trait anxiety inventory. *The Corsini encyclopedia of*
689 *psychology*. 2010:1-1.
- 690 59. Muntner P, Shimbo D, Carey RM, et al. Measurement of blood pressure in
691 humans: a scientific statement from the American Heart Association.
692 *Hypertension*. 2019;73(5):e35-e66.

- 693 60. Kaufmann T, Sütterlin S, Schulz SM, Vögele C. ARTiiFACT: a tool for heart rate
694 artifact processing and heart rate variability analysis. *Behavior research*
695 *methods*. 2011;43(4):1161-1170.
- 696 61. Behzadi Y, Restom K, Liao J, Liu TT. A component based noise correction
697 method (CompCor) for BOLD and perfusion based fMRI. *Neuroimage*.
698 2007;37(1):90-101.
- 699 62. Vallat R. Pingouin: statistics in Python. *Journal of Open Source Software*.
700 2018;3(31):1026.

701

702

703

704

705

706

707

708

709

710

711

712

Supplementary Materials

Supplementary Table S1. Resting Brain Connectivity

Significant condition effects (sleep rested < > sleep deprived) on brain connectivity, with all condition effects shown significant following FDR-correction. ACC, anterior cingulate cortex; MPFC, medial prefrontal cortex; vmPFC, ventromedial prefrontal cortex; HR, heart rate; HRV, heart rate variability (pNN50).

Source	Brain Region	Condition Effects (P-value)
Deprived > Rested		
Insula (R)	ACC (L/R)	0.01
	MPFC (R)	<0.001
	MPFC (L)	<0.001
Insula (L)	MPFC (R)	<0.001
	MPFC (L)	<0.001
MPFC (R)	ACC (L/R)	<0.001
	MPFC (L)	0.01
	Amygdala (R)	<0.005
	Amygdala (L)	<0.001
MPFC (L)	Amygdala (L)	0.02
ACC (L/R)	Amygdala (L)	0.01
Rested > Deprived		
MPFC (L)	vmPFC (L/R)	0.002

Supplementary Table S2.

Connectivity Associations with Affect. Columns show the Pearson correlation between sleep-loss related changes in brain connectivity and changes in affective state, including mood (positive and negative) and anxiety. * significant following FDR-correction. ACC, anterior cingulate cortex; MPFC, medial prefrontal cortex; vmPFC, ventromedial prefrontal cortex.

Source	Brain Region	Brain Correlation with Affect (R-value)		
Deprived > Rested		Positive Mood	Negative Mood	Anxiety
Insula (R)	ACC (L/R)	-0.03	-0.09	0.07
	MPFC (R)	-0.20	-0.05	0.18
	MPFC (L)	-0.15	-0.05	0.08
Insula (L)	MPFC (R)	-0.24	-0.001	0.13
	MPFC (L)	-0.16	0.07	0.13
MPFC (R)	ACC (L/R)	-0.05	0.11	0.001
	MPFC (L)	-0.05	0.10	-0.13
	Amygdala (R)	-0.003	-0.14	-0.02
	Amygdala (L)	-0.03	-0.13	-0.13
MPFC (L)	Amygdala (L)	0.05	-0.04	-0.10
ACC (L/R)	Amygdala (L)	-0.02	-0.03	0.04
Rested > Deprived				
MPFC (L)	vmPFC (L/R)	0.32*	0.06	-0.18

A

B

C

D

A

B

A

B