Title: Will Current Protected Areas Harbour Refugia for Threatened Arctic Vegetation Types until 2050?
A First Assessment

Authors: Merin R. Chacko¹, Ariane K.A. Goerens¹, Jacqueline Oehri¹, Elena Plekhanova¹ and Gabriela Schaepman-Strub¹

¹ Department of Evolutionary Biology and Environmental Studies, University of Zurich
Winterthurerstrasse 190, 8057 Zurich, Switzerland

Correspondence:

Gabriela Schaepman-Strub
University of Zurich
Winterthurerstrasse 190, 8057 Zurich, Switzerland
gabriela.schaepman@ieu.uzh.ch
+41 44 635 4806

Funding: This work was supported by the University Research Priority Programme Global Change and Biodiversity of the University of Zurich.

Author approvals: All authors have seen the approved the manuscript, and confirm that it has not been accepted or published elsewhere.

Declaration of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Keywords: global change, arctic conservation, tundra vegetation, CAVM, vegetation shifts, climate change refugia

Abstract

Arctic vegetation types provide food and shelter for fauna, support livelihoods of Northern peoples, and are tightly linked to climate, permafrost soils, lakes, rivers, and the ocean through carbon, energy, water, and nutrient fluxes. Despite its significant role, a comprehensive understanding of climate change effects on Arctic vegetation is lacking. We compare the 2003 baseline with existing 2050 predictions of circumpolar Arctic vegetation type distributions and demonstrate that abundant vegetation types with a proclivity for expansion contribute most to current protected areas. Applying IUCN criteria, we categorize five out of the eight assessed vegetation types as threatened by 2050. Our analyses show that current protected areas are insufficient for the mitigation of climate-imposed threats to these Arctic vegetation types. Therefore, we located potential climate change refugia, areas where vegetation may remain unchanged, at least until 2050, providing the highest potential for safeguarding threatened vegetation types. Our study provides an essential first step to assessing vegetation type vulnerability in the Arctic, but is based on predictions covering only 46% of Arctic landscapes. The co-development of new protective measures by policymakers and indigenous peoples at a pan-Arctic scale requires more robust and spatially complete vegetation predictions. This is essential as increasing pressures from resource exploration and rapid infrastructure development complicate the road to a sustainable development of the rapidly thawing and greening Arctic.

1. Introduction

The Arctic experiences climate warming at twice the global mean^{1,2}, leading to observed shifts in the distribution and composition of Arctic vegetation³ that are projected to intensify in the future⁴. These changes may threaten not only endemic plant species but entire vegetation types and the associated ecosystem functions. Vegetation types provide habitats for sessile and migratory animal species⁵, support livelihoods of Northern peoples⁶ and take part in various feedbacks involving climate^{7–10}, permafrost soils^{7,9,11}, lakes, rivers, and the ocean through water, energy and carbon fluxes^{12,13}.

Arctic vegetation is characterised by small vascular plants, bryophytes and lichens forming distinct plant communities, which are classified according to their functional types into fifteen vegetation types of the Circumpolar Arctic Vegetation Map (2003 CAVM)¹⁴. The southern border to the Arctic tundra is the treeline, which has been both observed and projected to advance northwards, decreasing the total

extent of the tundra while simultaneously displacing southern Arctic vegetation types^{15–18}. Due to decreasing snow cover and phenological changes, primary productivity and vascular plant biomass have been increasing, specifically that of tall shrubs⁹. This shrubification of the tundra has been widespread, resulting in a reduction in the abundance of less productive lichen- and moss-dominated vegetation types^{9,17,19–22}. Widespread greening of the Arctic due to climate warming has long been identified^{23,24}. However, spectral browning and increased heterogeneity of Arctic greening at the circumpolar scale have been observed in recent years^{17,25}. Indeed, a modelling study forecasted that 48–84% of Arctic vegetation types will have shifted by 2050 due to the effects of climate change⁴.

Novel pressures beyond those posed by warming are arising in the Arctic due to increasing anthropogenic presence. Historically, human land use and modification have been relatively low or non-existent in the Arctic biome. The human footprint map classifies most of the Arctic as under low pressure, and the human modification index demonstrates that the tundra remains thus far one of the last true terrestrial wild places on Earth^{26,27}. However, human interest in Arctic commodities such as oil and gas are increasing due to rising global energy needs, and with it also the extent of disturbances that terrestrial ecosystems experience due to exploration and infrastructure development^{6,28}. The effects of human disturbances and landscape processes such as climate and biota shifts on the distribution and composition of vegetation over time must be explicitly included in effective conservation efforts. As vegetation in the Arctic is more fragile and requires longer times to recover from perturbation in comparison to southern vegetation²⁸, it is both an ecological, political and economic imperative to have plans in place to mitigate and minimise these disturbances in order to provide a path towards the sustainable development of the Arctic.

As of 2016, 20.2% of the terrestrial Arctic area is protected to some degree²⁹. Although conservation efforts in the Arctic are well-developed, their focus is generally at the species level rather than the scale of ecosystems³⁰. In other biomes, the impact of climate change-induced biota shifts on conservation efforts has been recognised, and adaptive conservation strategies have been developed, though not widely implemented³¹. A recent systematic study identified areas with high potential for the persistence of multiple biodiversity elements under climate change in North America, and demonstrated that at the biome-scale, ~80% of areas within the top quintile of future conservation importance lacked formal protection, though this study excluded the High Arctic due to lacking data³². Additionally, needs in the North are influenced by rapid change at global scales and can no longer be addressed solely by local

actions³⁰. Indeed, local studies of Arctic plant communities do not always mirror observational trends over more extended temporal periods, which tend to be heterogeneous and complex^{17,24}. Conservation actions which operate at various scales of space, time and biological organisation may be required to effectively prevent the loss of potentially vulnerable ecosystems and their functions.

Refugia have become a focus of interest in conservation, as they have been demonstrated to enable the persistence of biodiversity over longer temporal scales and changing climates while retaining ecosystem and habitat functions^{33,34}. Vegetation refugia can be defined as areas where existing vegetation will remain within current suitable climate conditions³⁵. Refugia may serve as a means for giving plant communities the time needed to allow for local adaptation to new environmental states by providing habitats where the effects of climate change are least felt in the short-term³⁶. The identification and protection of refugia may facilitate the persistence of retreating vegetation types under projected anthropogenic climate change^{37,38}.

Presently, protected areas in the Arctic have not been established with climate change-induced vegetation shifts in mind. Consequently, there is no consensus as to the current state of vegetation vulnerability in the Arctic. Additionally, as Arctic vegetation shifts, protected areas of today may no longer protect the same vegetation to the same extent in the future. Vegetation distribution projections for the year 2050³⁹ provide the opportunity to locate vegetation refugia in the Arctic. Conservation according to vegetation refugia could, in effect, protect species at higher trophic levels in the food web through trophic cascades. However, to date, a comprehensive overview of vegetation refugia and their protected status in the Arctic has been unavailable.

The objective of this study is to establish an overview of vegetation type abundance in Arctic protected areas and assess their vulnerability. For this purpose, we utilised the 2003 CAVM as the baseline abundance of Arctic vegetation types within and outside of current protected areas. We then determined the distribution of Arctic vegetation types within protected areas using previously existing 2050 vegetation scenarios³⁹. Furthermore, we assessed the risk of collapse of the vegetation types following the IUCN Red List of Ecosystems criteria for the baseline and the future⁴⁰. Lastly, to inform conservation efforts, we located potential refugia for the vegetation types that had been identified as threatened.

2. Materials and Methods

2.1. Baseline status of vegetation type abundance in protected areas

We defined the network of protected areas in the Arctic according to their extent in the Map of Arctic Protected Areas (MAPA)²⁹. The protected areas were treated as one network regardless of the six management categories within the map, as our focus was on the collective pan-Arctic protection status. We utilised the 2003 CAVM to determine the baseline distribution of Arctic vegetation types within protected areas, which included fifteen distinct vegetation types (Appendix A) as well as the non-vegetative glacier class^{14,41}. We intersected the 2003 CAVM with MAPA and computed a zonal histogram. This resulted in the pixel count of the protected area for each vegetation type in order to obtain the total and absolute areas of the vegetation types present within protected areas.

2.2. Future status of vegetation type abundance in protected areas

The bases for all analyses concerning future vegetation type distribution were the 36 Maps of Future Arctic Vegetation Distribution (MFAVD), which predicted climate change-induced vegetation shifts for 2050⁴. Two machine learning methods for ecological niche modelling, three climate models, two emissions scenarios, and three tree dispersal scenarios had been applied to create 36 predictions⁴. The MFAVDs are the result of all possible combinations of the parameters named above. Areas covered by glaciers, barren lands and wetlands were excluded in the predictions, but tree cover was incorporated to illustrate the northward treeline shift of the taiga. We intersected all 36 MFAVDs with MAPA to obtain the complete range of possible outcomes. We selected an MFAVD that we deemed to be most realistic (hereafter 'realistic model'): random forest machine learning because it was the one with slightly higher accuracy⁴; the intermediate of the three global climate models (i.e. CSIRO); the higher of the two emissions scenarios, A2a⁴²; and the intermediate tree dispersal of 20 km, as it represents the highest rate at which trees have been observed to disperse northward in Alaska and Canada^{16,43}.

We analysed the eight vegetation types available in the MFAVDs out of the fifteen originally present in the 2003 CAVM, as barren (four classes) and wetlands (three classes) had been excluded from the predictions⁴. To assess the future distribution and protection status of Arctic vegetation, we converted MAPA from a polygon to a 4.5-km resolution raster to match the spatial resolution of the MFAVDs and attributed the protection status to every pixel. Finally, we summed the pixels and obtained the number of pixels per total as well as per protected area for each vegetation type.

2.3. Vulnerability of Arctic vegetation types following IUCN criteria

We applied the “IUCN Red List of Ecosystems Categories and Criteria” to quantify the vulnerability of Arctic vegetation types⁴⁰. The IUCN provides guidelines to assess the risk of ecosystem collapse, which are designed to be applicable for various kinds of ecosystems. Here, we defined ecosystems as the CAVM vegetation types, as they represent over 400 plant communities classified into broader functional groups⁴¹. The IUCN assigns one of three graded categories to threatened ecosystems: “vulnerable”, “endangered” and “critically endangered”. We did not differentiate here between the two graded categories for unthreatened ecosystems (“not threatened” and “least concern”) and instead simply summarised them as “not threatened”.

Five criteria are used to determine the IUCN risk category of the ecosystem: “reduction in geographic distribution”, “restricted in geographic distribution”, “environmental degradation”, “disruption of biotic processes or interactions”, and “quantitative analysis that estimates the probability of ecosystem collapse”⁴⁰. It is recommended that as many criteria as possible are assessed, as the ultimate classification of risk is determined as the highest of the five criteria⁴⁰. At the pan-Arctic scale, data were only available for “restricted distribution” and “decline of distribution”; therefore, this study could only assess these two criteria. Hence, our results represent the minimum risk status that each vegetation type can be assigned. Analyses of the remaining three criteria could potentially result in a higher risk category.

To determine the “restricted distribution” criterion classification, the number of 10×10 km grid cells occupied by the ecosystem must be calculated. An ecosystem is classified as “vulnerable”, “endangered”, or “critically endangered” if it occupies at most 50, 20 or 2 grid cells, respectively⁴⁴. After the total extent of each vegetation type within the 2003 CAVM was calculated in square kilometres, the extents were divided by 100 to determine the number of 10×10 grid cells occupied by the vegetation type across the pan-Arctic extent. The “decline of distribution” criterion is determined by the predicted relative reduction in the distribution of an ecosystem over fifty years. An ecosystem is classified as “vulnerable”, “endangered”, or “critically endangered”, if the reduction in its spatial extent is at least 30%, 50% or 80%, respectively⁴⁴. In this analysis, the relative changes in area were calculated according to the changes between the 2003 CAVM¹⁴ and the 2050 MFAVDs⁴; therefore, the classification is based on the predicted relative reduction over 47 rather than 50 years.

2.4. Identification of potential refugia for threatened vegetation types

For the threatened vegetation types, we identified areas which may serve as refugia, i.e. where a vegetation type is present in the baseline and is predicted to persist at least until 2050. We overlaid the realistic model with the 2003 CAVM and selected the areas where the pixels remained unchanged to create a map of areas with persistent vegetation types. We did not consider areas where these vegetation types had been predicted to shift to, and only considered the areas where the vegetation remained the same, as a) the predictions of future vegetation distribution contain a measure of uncertainty, b) there is evidence of lag effects in species-scale responses to climate change⁴⁵, and c) considerable variability in their ability to track new climatic niches exists⁴⁶.

Spatial data were analysed using ArcMap 10.5.1⁴⁷ and RStudio 1.2.5033^{48,49}. Maps were converted into the Lambert Azimuthal Equal Area Polar Projection to preserve area, a crucial feature for this analysis.

3. Results

3.1. Baseline status of vegetation type abundance in protected areas

The terrestrial pan-Arctic region covers over 7.1 million km², 20.2% of which is protected. Of the vegetated areas (4.7 million km²), 21% — or approximately 977,000 km² — fall within protected areas. Additionally, a significant part of the protected areas encompass glaciated areas (777,000 km²), but could not be considered within the scope of this study due to a lack of predictions.

Our analysis demonstrates that the fifteen vegetation types have substantially different absolute and relative spatial abundances in the pan-Arctic tundra (Figure 1). The most abundant vegetation type (S1, erect dwarf-shrub tundra) covers over six times more area than the least abundant vegetation type (W1, sedge/grass, moss wetland), 626,000 km², and 102,000 km², respectively. The abundances of the vegetation types within protected areas generally mirror their abundance at the pan-Arctic scale. For example, 11.4% of Arctic vegetation is composed of the G3 vegetation type (nontussock sedge, dwarf-shrub, moss tundra), which accordingly covers 11.3% of protected areas. Notable exceptions do occur, such as the wetland types. In comparison to their total extents (W1: sedge/grass, moss wetland, 2.0%; W2: sedge, moss, dwarf-shrub wetland, 2.7%; W3: sedge, moss, low-shrub wetland, 3.7%), they are overrepresented within protected areas (W1, 3.0%; W2, 4.8%; W3, 6.7%). Contrastingly, though 10.8% of Arctic vegetation is composed of the vegetation type B2 (cryptogam barren complex, bedrock), it spans only 2.5% of the protected area. In terms of absolute extents, the most abundant types at pan-

Arctic scales are overrepresented in protected areas, while the least abundant types are underrepresented.

3.2. Predicted vegetation type abundance in protected areas by 2050

Compared to the baseline abundances of vegetation types, the assessment of predicted abundances across protected areas by 2050 includes only eight vegetation types (see section 2.2). Predictions for the vegetation types within the barren (25.6% of non-glaciated terrestrial protected areas, 260,000 km²) and wetland (14.5% of non-glaciated terrestrial protected areas, 148,000 km²) classes were unavailable. Additionally, as the nearly 777,000 km² of glaciated protected areas were lacking in predictions, they were excluded from the study. The results differed considerably between the 36 MFAVDs: between 48,000 (0.1%) and 1,826,000 km² (39.9%) of the total area covered by tundra vegetation in the 2003 CAVM is predicted to be replaced by taiga⁴. Hence, considerable differences also resulted in the predictions of single vegetation types as a function of MFAVDs. We restrict the presentation of results to our selected realistic model and present results of the other MFAVDs as an envelope around this model result. Of the eight vegetation types, five are predicted to decline in their total area according to the realistic model, accompanied by a mirrored decline in their abundance within the protected area network (Figure 2). In contrast, two southern vegetation types (S2: low shrub tundra and G3: nontussock sedge, dwarf-shrub moss tundra) are predicted to gain in area, both within and outside of protected areas. Vegetation type G4 (tussock sedge, dwarf shrub, moss tundra) showed a decrease in total area, yet its abundance within the protected area network increased.

3.3. Vulnerability of Arctic vegetation types following IUCN criteria

To determine the risk of collapse for the eight vegetation types, we analysed two spatial criteria: “restriction of distribution” and “decline of distribution”; the final assigned category of risk was determined as the higher of these two. The “restriction of distribution” criterion diagnosed none of the eight vegetation types as threatened because each vegetation type has an extent of at least 5,000 km². The “decline in distribution” criterion assigned different results (Figure 3). Depending on the MFAVD, the specific risk status assigned to each type varied. Under the realistic model, one vegetation type (P1: prostrate dwarf-shrub, herb tundra) was classified as critically endangered. Four vegetation types were classified as endangered (G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra; S1: erect dwarf-shrub tundra), and three as not threatened (G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock sedge, dwarf-shrub, moss tundra;

S2: low shrub tundra). The threatened vegetation types (i.e., those classified at least as endangered) are generally the more northern types with sparse, low-growing, non-vascular vegetation.

3.4. Potential refugia for threatened vegetation types

For the five vegetation types (G1, G2, P1, P2, S1) classified as threatened under the realistic model, we determined refugia, regions where these vegetation types are predicted to persist until 2050 (Figure 4, Appendix C). The total area of refugia ranges from < 9,000 km² (P2) to > 100,000 km² (S1), accounting for 2.1% and 8.7% of currently protected areas. The refugia are scattered over Canada, Greenland, Norway, Russia, and the USA (Figures 4a-d).

4. Discussion

We evaluated the capability of the current Arctic network of protected areas for the conservation of Arctic vegetation types. Our results demonstrate that all Arctic vegetation types are found within the protected areas, and their relative abundance significantly varied, but generally followed the same pattern as seen at pan-Arctic scale. Noticeably, wetlands tend to be protected to an above-average extent; this is likely due to conservation efforts focusing on Wetlands of International Importance (Ramsar sites) related to their protection as breeding areas for migrating birds²⁹.

The impacts of climate change may demand the adaptation of protected areas^{31,50}, as they may no longer harbour the same vegetation types. Thus, we analysed the protection and threat status of the vegetation type abundances predicted for the year 2050. The two most productive vegetation types (G3 and S2) occupying the lower southern bioclimatic zones were demonstrably increasing in extent. The five vegetation types which were predicted to decline drastically currently occupy the three more northern bioclimatic zones³⁹ and are characterised by sparse and low-growing vegetation⁵¹. Our results demonstrate that these northern vegetation types also decrease in their representation within protected areas. We attribute this to two reasons. First, protected areas are generally situated in more southern regions rather than the northernmost edges of the Arctic, where these northern vegetation types can find sanctuary. This interpretation is validated by the fact that the refugia we located are clustered around the northernmost edges of the terrestrial Arctic. In our realistic model, we observe them being especially abundant in the Canadian Archipelago, which contains most of the terrestrial High Arctic. The climate will stay unsuitable for southern vegetation types in these regions, at least until 2050. Secondly, Arctic protected areas were not established with the conservation of vegetation types in mind⁶;

therefore, it stands to reason that vegetation types of conservation importance have not been recognised and hence prioritised.

4.1. Implications for flora, fauna and indigenous people if vegetation types are left unprotected

Vegetation shifts have been observed to have outcomes for species and communities across trophic levels^{5,9}. Threatened vegetation types may lead to the increased vulnerability of endemic flora and fauna, especially those that occur only in the High Arctic, which are most at risk. For example, there is evidence that vegetation exerts bottom-up control on caribou populations, as increasing shrubification decreases pasture quality⁵². This decline is exacerbated by the loss of lichen-rich vegetation types, which negatively affect caribou populations due to a loss of winter foraging options^{53,54} and energy through increased methane emission if lichens are reduced in their diet⁵⁵. Indigenous knowledge also shows that caribou populations have already adopted more northern migration patterns⁵⁶. This can, in turn, have negative consequences for indigenous communities that depend on caribou for food and as economic resources^{54,57}. Without focused conservation efforts on maintaining these vegetation types, the species depending on them will also become threatened with extinction.

4.2. Implications of land use change on threatened vegetation types if left unprotected

Climate warming and technological advances have opened up the Arctic as a new frontier in economic development. Increasing interest in commodities such as oil, gas, and mineral resources will naturally lead to increased infrastructure requirements in the Arctic, which may, in turn, intensify disturbances of Arctic ecosystems. Surface disturbances such as road networks and settlements have led to permafrost degradation through soil warming⁵⁸. Off-road vehicles like tundra tractors can destroy endemic vegetation, and leave the tracks visible for decades as vegetation struggles to recover⁵⁸. A study of the Alaska North Slope demonstrated that 34% of the area was affected by oil development by 2010⁵⁹. Further exploration has already been proposed, such as a 3-D seismic survey covering 63,000 km of trails within the Arctic National Wildlife Refugia in Alaska⁶⁰. A recent study demonstrated that this could result in mid-to-high level impacts on 122 km² of the area in question, leading to increased thermokarst formation and erosion, and negatively impacting moist vegetation types⁶¹. Additionally, vehicle tires are capable of significantly increasing the dispersal distance of southern invaders, such as *Salix lanata*, which has been found to occur in many road-side areas, replacing lichen and moss cover⁵⁹. Local warming due to the settlements also generate pockets of suitable habitats in latitudes where southern vegetation could not otherwise persist⁵⁸.

Permafrost thaw is critically threatening Arctic infrastructures⁶² and their collapse can lead to dramatic consequences for terrestrial ecosystems and indigenous livelihoods. As the MFAVDs did not account for human land use modification, this study severely understates their effects on vegetation shifts. At the pan-Arctic scale, a combination of economic, geopolitical, climatic, infrastructure and ecological factors lead to uncertainty in the future spatial distribution of development pressures, and their degree of impact⁶³. Therefore, it is essential to predefine conservation areas for the currently unprotected and threatened vegetation types before widespread development begins in the increasingly accessible Arctic. Complementing protected areas, biodiversity conservation schemes developed for other regions of the world, such as around mining sites, need to be adapted to Arctic conditions to prevent loss of vegetation where industrial development outside of protected areas is allowed.

4.3. Implications of using climate change refugia for land conservation and management

Globally protected areas are 10.6% richer in species diversity than non-protected areas⁶⁴. Nevertheless, conserving biodiversity using protected areas poses to be difficult under climate change. A study assessing the protected areas of Canada found that climate change will lead to over 40% of the protected areas experiencing a change in biome type, with the total extent of the Canadian tundra standing to decline by 38-79%⁶⁵. In order to most efficiently use limited resources and protect threatened vegetation, efforts in in situ management must be focused on areas where the vegetation is likely to persist under global change, such as climate change refugia for vegetation. There is scepticism over the use of refugia as areas where vegetation may retreat until conditions in the surrounding environment become more favourable. Vegetation shifts due to climate warming are projected to continue and intensify; therefore, the time periods required until favourable conditions return may be longer than the existence of these refugia. However, they may serve a powerful purpose by buying time for the climate adaptation of vulnerable species and ecological communities³⁶. Adjusting Arctic protection efforts in mitigating the consequences of global change with a focus on relatively transient climate change refugia may aid in a long-term transformation of these threatened vegetation types into novel community assemblages (i.e. ecological replacements) that have adapted to these environmental changes, while performing the same ecological and habitat functions³⁶. This process has been documented in the late Quaternary⁶⁶.

4.4. Limitations and urgent need for comprehensive predictions of Arctic vegetation type distribution

The aim of this study is to construct a first assessment of the protected status of Arctic vegetation types in light of climate change based on existing vegetation predictions. These predictions³⁹ were based on the 2003 CAVM, a vector map with polygons of 14 km minimum diameter⁴¹; thus for the sake of consistency, it was utilised by this study instead of the more recent raster version (2019 CAVM) that is resolved at 1 km⁵¹. Nevertheless, these disparities did not affect the results regarding the baseline risk status of the vegetation types; the general pattern of abundant vegetation types being better represented within protected areas holds, regardless of the map used (Appendix B).

Increasing spatial resolution in future predictions could reveal pockets of vegetation occurring outside of the main macroclimatic niche of a vegetation type. Such pockets could accelerate the spread of vegetation types into surrounding areas should the climate become more favourable. Potentially, southern vegetation types could then disperse northward at faster rates than modelled⁴, increasing the displacement of northern vegetation types and potentially our risk classifications. This displacement may be balanced by potential northern micropockets which would increase the total extent of refugia. Nevertheless, these disparities did not affect the results regarding the present risk status of the vegetation types. Though the “decline in distribution” criterion could not be tested as MFAVDs do not exist for the 2019 CAVM, none of the vegetation types of the 2019 CAVM showed a restricted distribution and were classified as unthreatened under the IUCN “restriction of distribution” criterion.

Due to limited data availability in the Arctic, we could only analyse two of the five IUCN criteria. Therefore, the assigned risk category only represents the minimal risk, and could potentially be higher, both in the present and in the future. Though it is recommended that as many criteria as possible are assessed, the IUCN Red List of Ecosystems was created with the purpose of flexibility in the use of data containing variations in quality and coverage⁴⁰. Therefore, this risk assessment still provides valuable information as a starting point for future assessments as data availability increases.

The MFAVDs only rendered predictions for 46% of the pan-Arctic area. Specifically, they excluded glaciers (including nunataks), barren lands, and wetlands⁴. Subsequently, we were unable to assess the future distribution of wetland and barren vegetation types, classify their risk of collapse, and locate potential refugia. Therefore, our results do not take into consideration the potential of the assessed

vegetation types to disperse into areas currently covered by these unassessed vegetation types, and vice versa.

Glaciers currently cover nearly one-third of the Arctic⁴¹, but extensive climate change-induced glacier melt is predicted^{67,68}. Vegetation succession following retreating glaciers in the Arctic follows the same patterns as elsewhere; however, these processes occur at larger temporal scales and may be less relevant for the next thirty years^{69,70}. Vegetation growth response to warming is slower in the High Arctic than in the Low Arctic due to lower growing season temperatures, seasonal length and nutrient availability⁷¹. In Svalbard, for instance, vascular plant establishment was highly limited for the first century after glacial retreat⁶⁹. Additionally, vegetation can take up to 500 years to reach equilibrium after glacial retraction in Greenland¹². Therefore, the inclusion of glaciers would have likely had limited impacts on the results for threatened vegetation types shown here for 2050. Nevertheless, they would be crucial for imminent studies which investigate the future of the terrestrial Arctic at longer timescales.

One-quarter of the terrestrial Arctic consists of barren lands⁴¹, which were excluded in this analysis. Therefore, the northern vegetation types may be less threatened than indicated by this study because they may expand into currently barren areas. The inclusion of barren lands in future studies would likely lead to predictions of northern vegetation types being slightly more abundant and increasing over centennial periods. However, due to the slow response in the development of soils and vegetation, the time scales at which this northern expansion into barren lands occurs may be longer than the 47-year span of our study, and this expansion may be limited, as barren lands lack in soil sufficient enough to support vegetation⁴.

Wetlands constitute 7% of the Arctic vegetated area⁴¹. Predictions for future wetland distribution are particularly difficult due to the complexity of the hydrological processes regulating them⁷². Permafrost thaw, increasing precipitation, ice melt, and evaporation due to warming influence the abundance and distribution of surface water, indirectly affecting wetlands^{13,73}. In the southern areas with discontinuous permafrost, ponds are generally in decline as permafrost thaw leads to increasing drainage^{74–76}. Conversely, thawing of continuous permafrost in the High Arctic has been shown to lead to an increasing abundance of thaw ponds⁷⁵. In the Canadian Arctic, this trend of thermokarst pond formation has been shown to occur at rapid paces⁷⁷. This pattern of wetland development means that generally, southern vegetation types would have more potential areas of expansion, whereas northern areas may become

further displaced by expanding wetland. We hypothesise that the inclusion of wetlands in this study could consequently have a reinforcing effect on the observed patterns of threatened vegetation types. While wetlands are crucial because of the abundance of migratory birds dependent on them, they make up a relatively small portion of the vegetated pan-Arctic and are generally better protected than other vegetation types. However, this may change significantly in the future due to climate change. Our study highlights the need for further focus on these ill-understood vegetation types in longer-term predictions.

The future vegetation shifts⁴ were predicted under various assumptions about climate change and tree dispersal rate scenarios, which, though necessary, carry over to this study. Uncertainties in future vegetation distributions are inherent to predictions that are based on climate models with many uncertain assumptions, where one model is no more or less valid than the other⁴². Our selection of the realistic future model with regard to climate and emission scenario is, therefore, subjective. There is a large spread of the vegetation type abundances across the models, making the results, especially for vegetation types G1, G2, G4 and S1, more uncertain (figure 2 and appendix D). However, the general trend of increase in southern vegetation types and decrease in northern vegetation types still holds within and outside of protected areas, as demonstrated for most results by the minimum and maximum scenario envelope.

While land surface schemes for earth system models and vegetation development models are being adapted to Arctic tundra conditions, parameterisations at the Arctic tundra vegetation type level, including all vegetation types, need to be implemented, and biotic and abiotic interactions integrated with holistic modelling approaches, at pan-Arctic scale. Furthermore, inclusion of human land use scenarios (e.g. reindeer herding densities, potential mining sites, road development) will be necessary in future modelling efforts to allow for informed policies and decisions.

5. Conclusions

This study identifies baseline and future abundances of Arctic vegetation types within protected areas as well as potential climate change refugia for threatened vegetation types using the 2003 CAVM and existing predictions of future Arctic vegetation distribution. Though uncertainties exist within the maps provided here, the general trends seen in this study are valuable in guiding future conservation efforts. Further studies on the development of Arctic vegetation types which include the whole Arctic — including wetlands, barrens and ice-covered areas — are urgently needed. They could provide complete and

precise findings to improve and adapt conservation efforts to climate change-induced biota shifts. Moreover, we recommend independent validation of these hypothetical predictions to better incorporate climate change refugia into the designation of protected areas. This could potentially be achieved through recent validation methods which compare predictions of refugia to local measures of species richness, endemic species persistence, genetic diversity, plant functional traits and/or demographic variables⁷⁸.

This study aims to shift the focus of pan-Arctic protection efforts towards higher scales of biological organisation, from species to communities and ecosystems, and in this case, vegetation types. Endemic tundra vegetation types face the threat of ecosystem collapse due to global change. The establishment of refugia for the vegetation types identified here could protect them as refugia have done over the history of life on earth^{66,67}. The protection of vegetation refugia would additionally have positive effects on Arctic fauna and climate regulation through bottom-up biotic and abiotic interactions. However, the integration of vegetation types for climate change-adapted conservation in the Arctic requires urgent collaboration between policymakers and indigenous peoples, as the area becomes increasingly under pressure from exploration and rapid infrastructure development. As experienced in the recent decade, Arctic change is rapid. Extreme events, including extreme winter precipitation and summer drought, are already affecting Arctic ecosystems through major disturbance events such as extensive flooding and fires, adding yet another dimension of abrupt change to mitigate in the future. Without a plan already in place for the protection of these critically important Arctic landscapes, we cannot enable the sustainability of economic and structural development in what is increasingly no longer one of the world's last truly wild places.

References

1. Walsh, J. E., Overland, J. E., Groisman, P. Y. & Rudolf, B. Ongoing climate change in the arctic. *Ambio* (2011) doi:10.1007/s13280-011-0211-z.
2. Collins, M. *et al.* Long-term climate change: Projections, commitments and irreversibility. in *Climate Change 2013: The Physical Science Basis. IPCC Working Group I Contribution to AR5.* (2013). doi:10.1017/CBO9781107415324.024.
3. Serreze, M. C. *et al.* Observational evidence of recent change in the northern high-latitude environment. *Clim. Change* (2000) doi:10.1023/A:1005504031923.
4. Pearson, R. G. *et al.* Shifts in Arctic vegetation and associated feedbacks under climate change. *Nat. Clim. Chang.* (2013) doi:10.1038/nclimate1858.
5. Wheeler, H. C., Høye, T. T. & Svenning, J. C. Wildlife species benefitting from a greener Arctic are most sensitive to shrub cover at leading range edges. *Glob. Chang. Biol.* (2018) doi:10.1111/gcb.13837.
6. CAFF. *Arctic Biodiversity Assessment. Status and trends in Arctic biodiversity. Conservation of Arctic Flora and Fauna. Arctic Biodiversity Assessment* (2013).
7. Juszak, I., Erb, A. M., Maximov, T. C. & Schaepman-Strub, G. Arctic shrub effects on NDVI, summer albedo and soil shading. *Remote Sens. Environ.* (2014) doi:10.1016/j.rse.2014.07.021.
8. Loranty, M. M., Goetz, S. J. & Beck, P. S. A. Tundra vegetation effects on pan-Arctic albedo. *Environ. Res. Lett.* (2011) doi:10.1088/1748-9326/6/2/029601.
9. Myers-Smith, I. H. *et al.* Shrub expansion in tundra ecosystems: Dynamics, impacts and research priorities. *Environ. Res. Lett.* (2011) doi:10.1088/1748-9326/6/4/045509.
10. Swann, A. L., Fung, I. Y., Levis, S., Bonan, G. B. & Doney, S. C. Changes in arctic vegetation amplify high-latitude warming through the greenhouse effect. *Proc. Natl. Acad. Sci. U. S. A.* (2010) doi:10.1073/pnas.0913846107.
11. Bonfils, C. J. W. *et al.* On the influence of shrub height and expansion on northern high latitude climate. *Environ. Res. Lett.* (2012) doi:10.1088/1748-9326/7/1/015503.

- 450 12. Lunt, D. J., de Noblet-Ducoudré, N. & Charbit, S. Effects of a melted greenland ice sheet on
451 climate, vegetation, and the cryosphere. *Clim. Dyn.* (2004) doi:10.1007/s00382-004-0463-4.
- 452 13. Woo, M. K. & Young, K. L. High Arctic wetlands: Their occurrence, hydrological characteristics
453 and sustainability. in *Journal of Hydrology* (2006). doi:10.1016/j.jhydrol.2005.07.025.
- 454 14. CAVM Team. Circumpolar Arctic Vegetation Map. (1:7,500,000 scale), Conservation of Arctic
455 Flora and Fauna (CAFF) Map No. 1. *US Fish and Wildlife Service.(Anchorage, AK). Available*
456 *at <http://www.geobotany.uaf.edu/cavm/>[Verified 1 Augus15 July 2015]* (2003).
- 457 15. Chapin, F. S. & Starfield, A. M. Time lags and novel ecosystems in response to transient
458 climatic change in arctic Alaska. *Clim. Change* (1997) doi:10.1023/A:1005337705025.
- 459 16. Lloyd, A. H., Rupp, T. S., Fastie, C. L. & Starfield, A. M. Patterns and dynamics of treeline
460 advance on the Seward Peninsula, Alaska. *J. Geophys. Res. D Atmos.* (2002)
461 doi:10.1029/2001jd000852.
- 462 17. Bjorkman, A. D. *et al.* Status and trends in Arctic vegetation: Evidence from experimental
463 warming and long-term monitoring. *Ambio* (2020) doi:10.1007/s13280-019-01161-6.
- 464 18. Rundqvist, S. *et al.* Tree and shrub expansion over the past 34 years at the tree-line near
465 Abisko, Sweden. *Ambio* (2011) doi:10.1007/s13280-011-0174-0.
- 466 19. Blok, D. *et al.* Shrub expansion may reduce summer permafrost thaw in Siberian tundra. *Glob.*
467 *Chang. Biol.* (2010) doi:10.1111/j.1365-2486.2009.02110.x.
- 468 20. Elmendorf, S. C. *et al.* Plot-scale evidence of tundra vegetation change and links to recent
469 summer warming. *Nat. Clim. Chang.* (2012) doi:10.1038/nclimate1465.
- 470 21. Sturm, M., Racine, C. & Tape, K. Increasing shrub abundance in the Arctic. *Nature* (2001)
471 doi:10.1038/35079180.
- 472 22. Loranty, M. M. & Goetz, S. J. Shrub expansion and climate feedbacks in Arctic tundra.
473 *Environmental Research Letters* (2012) doi:10.1088/1748-9326/7/1/011005.
- 474 23. Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G. & Nemani, R. R. Increased plant growth
475 in the northern high latitudes from 1981 to 1991. *Nature* (1997) doi:10.1038/386698a0.

- 476 24. Myers-Smith, I. H. *et al.* Complexity revealed in the greening of the Arctic. *Nature Climate*
477 *Change* (2020) doi:10.1038/s41558-019-0688-1.
- 478 25. Phoenix, G. K. & Bjerke, J. W. Arctic browning: extreme events and trends reversing arctic
479 greening. *Global Change Biology* (2016) doi:10.1111/gcb.13261.
- 480 26. Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch-Mordo, S. & Kiesecker, J. Managing
481 the middle: A shift in conservation priorities based on the global human modification gradient.
482 *Glob. Chang. Biol.* (2019) doi:10.1111/gcb.14549.
- 483 27. Venter, O. *et al.* Global terrestrial Human Footprint maps for 1993 and 2009. *Sci. Data* (2016)
484 doi:10.1038/sdata.2016.67.
- 485 28. Kumpula, T., Pajunen, A., Kaarlejärvi, E., Forbes, B. C. & Stammer, F. Land use and land
486 cover change in Arctic Russia: Ecological and social implications of industrial development.
487 *Glob. Environ. Chang.* (2011) doi:10.1016/j.gloenvcha.2010.12.010.
- 488 29. CAFF & PAME. *Arctic Protected Areas: Indicator Report, 2017: Conservation of Arctic Flora*
489 *and Fauna and Protection of the Arctic Marine Environment.* (2017).
- 490 30. Chapin, F. S., Sommerkorn, M., Robards, M. D. & Hillmer-Pegram, K. Ecosystem stewardship:
491 A resilience framework for arctic conservation. *Glob. Environ. Chang.* (2015)
492 doi:10.1016/j.gloenvcha.2015.07.003.
- 493 31. Heller, N. E. & Zavaleta, E. S. Biodiversity management in the face of climate change: A
494 review of 22 years of recommendations. *Biological Conservation* (2009)
495 doi:10.1016/j.biocon.2008.10.006.
- 496 32. Stralberg, D., Carroll, C. & Nielsen, S. E. Toward a climate-informed North American protected
497 areas network: Incorporating climate-change refugia and corridors in conservation planning.
498 *Conserv. Lett.* (2020) doi:10.1111/conl.12712.
- 499 33. Tzedakis, P. C., Lawson, I. T., Frogley, M. R., Hewitt, G. M. & Preece, R. C. Buffered tree
500 population changes in a Quaternary refugium: Evolutionary implications. *Science* (80-.).
501 (2002) doi:10.1126/science.1073083.
- 502 34. Morelli, T. L. *et al.* Managing climate change refugia for climate adaptation. *PLoS One* **11**, 1–

- 503 17 (2016).
- 504 35. Thorne, J. H. *et al.* Vegetation refugia can inform climate-adaptive land management under
505 global warming. *Front. Ecol. Environ.* **18**, 281–287 (2020).
- 506 36. Morelli, T. L. *et al.* Climate-change refugia: biodiversity in the slow lane. *Front. Ecol. Environ.*
507 **18**, 228–234 (2020).
- 508 37. Taberlet, P. & Cheddadi, R. Ecology: Quaternary refugia and persistence of biodiversity.
509 *Science* (2002) doi:10.1126/science.297.5589.2009.
- 510 38. Noss, R. F. Beyond kyoto: Forest management in a time of rapid climate change.
511 *Conservation Biology* (2001) doi:10.1046/j.1523-1739.2001.015003578.x.
- 512 39. Pearson, R. G. *et al.* Predicted Arctic vegetation distribution shifts under future climate change.
513 (2013) doi:10.18739/A2V935.
- 514 40. Bland, L. M., Keith, D. A., Miller, R. M., Murray, N. J. & Rodriguez, J. P. *Guidelines for the*
515 *application of IUCN Red List of ecosystems categories and criteria. Guidelines for the*
516 *application of IUCN Red List of ecosystems categories and criteria* (2015).
517 doi:10.2305/iucn.ch.2016.rle.1.en.
- 518 41. Walker, D. A. *et al.* The Circumpolar Arctic vegetation map. *J. Veg. Sci.* (2005)
519 doi:10.1111/j.1654-1103.2005.tb02365.x.
- 520 42. IPCC. *IPCC Special Report on Emissions Scenarios (SRES) of Working Group III.* (2000)
521 doi:citeulike-article-id:9904924.
- 522 43. Lescop-Sinclair, K. & Payette, S. Recent Advance of the Arctic Treeline Along the Eastern
523 Coast of Hudson Bay. *J. Ecol.* (1995) doi:10.2307/2261175.
- 524 44. Dudley, N. *Guidelines for applying protected area management categories. Guidelines for*
525 *applying protected area management categories* (2008). doi:10.2305/iucn.ch.2008.paps.2.en.
- 526 45. Stewart, L. *et al.* The regional species richness and genetic diversity of Arctic vegetation reflect
527 both past glaciations and current climate. *Glob. Ecol. Biogeogr.* (2016) doi:10.1111/geb.12424.
- 528 46. La Sorte, F. A. & Jetz, W. Tracking of climatic niche boundaries under recent climate change.

- 529 *J. Anim. Ecol.* (2012) doi:10.1111/j.1365-2656.2012.01958.x.
- 530 47. ESRI. ArcMap 10.5.1 [GIS Software]. (2017).
- 531 48. R Development Core Team. R: A language and environment for statistical computing. *Vienna,*
532 *Austria* (2017) doi:R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-
533 0, URL <http://www.R-project.org>.
- 534 49. Team RStudio. RStudio: Integrated Development for R. *Rstudio Team, PBC, Boston, MA URL*
535 *<http://www.rstudio.com/>* (2020).
- 536 50. Hannah, L., Midgley, G. F. & Millar, D. Climate change-integrated conservation strategies.
537 *Glob. Ecol. Biogeogr.* (2002) doi:10.1046/j.1466-822X.2002.00306.x.
- 538 51. Raynolds, M. K. *et al.* A raster version of the Circumpolar Arctic Vegetation Map (CAVM).
539 *Remote Sens. Environ.* (2019) doi:10.1016/j.rse.2019.111297.
- 540 52. Fauchald, P., Park, T., Tømmervik, H., Myneni, R. & Hausner, V. H. Arctic greening from
541 warming promotes declines in caribou populations. *Sci. Adv.* (2017)
542 doi:10.1126/sciadv.1601365.
- 543 53. Cornelissen, J. H. C. *et al.* Global change and arctic ecosystems: Is lichen decline a function of
544 increases in vascular plant biomass? *J. Ecol.* (2001) doi:10.1046/j.1365-2745.2001.00625.x.
- 545 54. Joly, K., Jandt, R. R. & Klein, D. R. Decrease of lichens in Arctic ecosystems: The role of
546 wildfire, caribou, reindeer, competition and climate in north-western Alaska. *Polar Research*
547 (2009) doi:10.1111/j.1751-8369.2009.00113.x.
- 548 55. Hansen, K. K., Sundset, M. A., Folkow, L. P., Nilsen, M. & Mathiesen, S. D. Methane
549 emissions are lower from reindeer fed lichens compared to a concentrate feed. *Polar Res.* **37**,
550 (2018).
- 551 56. Ksenofontov, S., Backhaus, N. & Schaepman-Strub, G. 'There are new species': indigenous
552 knowledge of biodiversity change in Arctic Yakutia. *Polar Geogr.* **42:1**, 34–57 (2019).
- 553 57. Nuttall, M. *et al.* Ch.12 Hunting, herding, fishing, and gathering: Indigenous peoples and
554 renewable resource use in the Arctic. *Arct. Clim. Impact Assess.* (2005).

- 555 58. Forbes, B. C., Ebersole, J. J. & Strandberg, B. Anthropogenic disturbance and patch dynamics
556 in circumpolar arctic ecosystems. *Conservation Biology* (2001) doi:10.1046/j.1523-
557 1739.2001.015004954.x.
- 558 59. Raynolds, M. K. *et al.* Cumulative geoeological effects of 62 years of infrastructure and
559 climate change in ice-rich permafrost landscapes, Prudhoe Bay Oilfield, Alaska. *Glob. Chang.*
560 *Biol.* (2014) doi:10.1111/gcb.12500.
- 561 60. SAExploration. Marsh Creek 3D plan of operationswinter seismic survey.
562 [https://eplanning.blm.gov/public_projects/nepa/111085/153349/187888/Marsh_Creek_Plan_of](https://eplanning.blm.gov/public_projects/nepa/111085/153349/187888/Marsh_Creek_Plan_of_Operations_Submitted_May2018.pdf)
563 [_Operations_Submitted_May2018.pdf](https://eplanning.blm.gov/public_projects/nepa/111085/153349/187888/Marsh_Creek_Plan_of_Operations_Submitted_May2018.pdf) (2018).
- 564 61. Raynolds, M. K. *et al.* Landscape impacts of 3D-seismic surveys in the Arctic National Wildlife
565 Refuge, Alaska. *Ecol. Appl.* **0**, 1–20 (2020).
- 566 62. Hjort, J. *et al.* Degrading permafrost puts Arctic infrastructure at risk by mid-century. *Nat.*
567 *Commun.* **9**, (2018).
- 568 63. Walker, D. A. & Peirce, J. L. *Rapid Arctic Transitions due to Infrastructure and Climate*
569 *(RATIC): A contribution to ICARP III.* (2015).
- 570 64. Gray, C. L. *et al.* Local biodiversity is higher inside than outside terrestrial protected areas
571 worldwide. *Nat. Commun.* (2016) doi:10.1038/ncomms12306.
- 572 65. Lemieux, C. J. & Scott, D. J. Climate change, biodiversity conservation and protected area
573 planning in Canada. *Can. Geogr.* (2005) doi:10.1111/j.0008-3658.2005.00103.x.
- 574 66. Jackson, S. T. & Overpeck, J. T. Responses of plant populations and communities to
575 environmental changes of the late Quaternary. *Paleobiology* (2000)
576 doi:10.1017/s0094837300026932.
- 577 67. Alley, R. B. *et al.* History of the Greenland Ice Sheet: paleoclimatic insights. *Quat. Sci. Rev.*
578 (2010) doi:10.1016/j.quascirev.2010.02.007.
- 579 68. Hinzman, L. D. *et al.* Evidence and implications of recent climate change in Northern Alaska
580 and other Arctic regions. *Clim. Change* (2005) doi:10.1007/s10584-005-5352-2.
- 581 69. Hodkinson, I. D., Coulson, S. J. & Webb, N. R. Community assembly along proglacial

- 582 chronosequences in the high Arctic: Vegetation and soil development in north-west Svalbard.
- 583 *J. Ecol.* (2003) doi:10.1046/j.1365-2745.2003.00786.x.
- 584 70. Raynolds, M. K. & Walker, D. A. Effects of deglaciation on circumpolar distribution of arctic
- 585 vegetation. *Can. J. Remote Sens.* (2009) doi:10.5589/m09-006.
- 586 71. Walker, M. D. *et al.* Plant community responses to experimental warming across the tundra
- 587 biome. *Proc. Natl. Acad. Sci. U. S. A.* (2006) doi:10.1073/pnas.0503198103.
- 588 72. Walvoord, M. A. & Kurylyk, B. L. Hydrologic Impacts of Thawing Permafrost-A Review. *Vadose*
- 589 *Zo. J.* (2016) doi:10.2136/vzj2016.01.0010.
- 590 73. White, D. M., Craig Gerlach, S., Loring, P., Tidwell, A. C. & Chambers, M. C. Food and water
- 591 security in a changing arctic climate. *Environ. Res. Lett.* (2007) doi:10.1088/1748-
- 592 9326/2/4/045018.
- 593 74. Riordan, B., Verbyla, D. & McGuire, A. D. Shrinking ponds in subarctic Alaska based on 1950-
- 594 2002 remotely sensed images. *J. Geophys. Res. Biogeosciences* (2006)
- 595 doi:10.1029/2005JG000150.
- 596 75. Smith, L. C. Disappearing Arctic Lakes. *Science* (80-.). (2005) doi:10.1126/science.1108142.
- 597 76. Yoshikawa, K. & Hinzman, L. D. Shrinking thermokarst ponds and groundwater dynamics in
- 598 discontinuous permafrost near Council, Alaska. *Permafr. Periglac. Process.* (2003)
- 599 doi:10.1002/ppp.451.
- 600 77. Farquharson, L. M. *et al.* Climate Change Drives Widespread and Rapid Thermokarst
- 601 Development in Very Cold Permafrost in the Canadian High Arctic. *Geophys. Res. Lett.* (2019)
- 602 doi:10.1029/2019GL082187.
- 603 78. Barrows, C. W. *et al.* Validating climate-change refugia: empirical bottom-up approaches to
- 604 support management actions. *Front. Ecol. Environ.* **18**, 298–306 (2020).
- 605

Figures

Figure 1. The 2003 baseline abundance and protection status of Arctic vegetation types. The Arctic tundra encompasses approximately 4.7 million km². The height of the bars represents the absolute extent of each vegetation type within and outside of protected areas. The percentages above the bars represent the relative abundance of the vegetation type in the pan-Arctic tundra. The percentages within the darker coloured bars (protected area) represent the relative abundance of each vegetation type in the protected area in comparison to the total protected area. The vegetation types can be summarised into barren tundra (B1: cryptogam herb barren; B2: cryptogam barren complex (bedrock); B3: noncarbonate mountain complex and B4: carbonate mountain complex), graminoid tundra (G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock sedge, dwarf-shrub, moss tundra), prostrate-shrub tundra (P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra), erect-shrub tundra (S1: erect dwarf-shrub tundra; S2: low shrub tundra) and wetlands (W1: sedge/grass, moss wetland; W2: sedge, moss, dwarf-shrub wetland; W3: sedge, moss, low-shrub wetland).

Figure 2. A comparison of the abundance and protection status of Arctic vegetation types at the baseline and in the future. The data presented refers to the realistic model scenario; error bars represent the envelope of the minimal and maximal abundance predicted by the different MFAVDs. The minimum and maximum values of the envelope are not always the result of the same MFAVDs. Depending on the vegetation type, different models predict the highest or lowest abundance. The vegetation types can be summarised into graminoid tundra (G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock sedge, dwarf-shrub, moss tundra), prostrate-shrub tundra (P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra), and erect-shrub tundra (S1: erect dwarf-shrub tundra; S2: low shrub tundra). Barren and wetland vegetation types were not assessed as predictions for 2050 were unavailable.

Figure 3. Risk status according to the 'decline in distribution' criterion of the IUCN Red List of Ecosystems Categories and Criteria under the realistic model. The vegetation types are ordered along the bioclimatic subzones they occur in, from low to high summer temperatures (left to right). The data presented refers to the realistic model scenario; error bars represent the envelope of the minimal and maximal abundance predicted by the different MFAVDs. The minimum and maximum values of the envelope are not always the result of the same MFAVDs. Depending on the vegetation type, different models predict the highest or lowest abundance. The vegetation types can be summarised into graminoid tundra (G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; G3: nontussock sedge, dwarf-shrub, moss tundra; G4: tussock sedge, dwarf-shrub, moss tundra), prostrate-shrub tundra (P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra), and erect-shrub tundra (S1: erect dwarf-shrub tundra; S2: low shrub tundra). Barren and wetland vegetation types were not assessed as no predictions for 2050 were available.

Figure 4. Refugia maps of threatened vegetation types in the Arctic (a), and magnified subsets centred on the Yamal peninsula, Russia (b); the Canadian Archipelago (c); and Alaska, USA, and North-eastern Siberia, Russia (d). The vulnerable vegetation types presented here are G1: rush/grass forb, cryptogam tundra; G2: graminoid, prostrate dwarf-shrub, forb tundra; P1: prostrate dwarf shrub, herb tundra; P2: prostrate/hemiprostrate dwarf-shrub tundra and S1: erect dwarf-shrub tundra.