THE CORRELATION OF SUBSTITUTION EFFECTS ACROSS POPULATIONS AND GENERATIONS IN THE PRESENCE OF NON-ADDITIVE FUNCTIONAL GENE ACTION

A. Legarra*, C.A. Garcia-Baccino*[†], Y.C.J. Wientjes[‡], Z.G. Vitezica*

* INRAE/INP, UMR 1388 GenPhySE, 31326, Castanet-Tolosan, France.

[†] Departamento de Producción Animal, Facultad de Agronomía, Universidad de Buenos Aires, C1417DSQ, Buenos Aires, Argentina.

[‡] Wageningen University & Research, Animal Breeding and Genomics, 6700 AH Wageningen, The Netherlands.

13 RUNNING TITLE: Correlation of QTL substitution effects

14

15 Keywords: QTL, substitution effects, epistasis, dominance, genetic distance

16

17

18 Corresponding author:

19

20 Andres Legarra,

21 andres.legarra@inrae.fr

22 +33 561285182

23 INRAE, UMR1388 GenPhySE

24 CS 52627

25 31326 Castanet Tolosan, France

26

ABSTRACT

Allele substitution effects at quantitative trait loci (QTL) are part of the basis of quantitative genetics theory and applications such as association analysis and genomic prediction. In the presence of non-additive functional gene action, substitution effects are not constant across populations. We develop an original approach to model the difference in substitution effects across populations as a first order Taylor series expansion from a “focal” population. This expansion involves the difference in allele frequencies and second-order statistical effects (additive by additive and dominance). The change in allele frequencies is a function of relationships (or genetic distances) across populations. As a result, it is possible to estimate the correlation of substitution effects across two populations using three elements: magnitudes of additive, dominance and additive by additive variances; relationships (Nei’s minimum distances or F_{st} indexes); and assumed heterozygosities. Similarly, the theory applies as well to distinct generations in a population, in which case the distance across generations is a function of increase of inbreeding. Simulation results confirmed our derivations. Slight biases were observed, depending on the non-additive mechanism and the reference allele. Our derivations are useful to understand and forecast the possibility of prediction across populations and the similarity of GWAS effects.

INTRODUCTION

One of the aims of quantitative genetics is to provide methods for prediction, for instance genomic prediction (prediction of livestock breeding values or of crop performance) or polygenic risk score (risk of a disease in humans). These predictions would ideally work across a range of populations (different breeds, future generations). Ideally, the prediction goes through a process of identifying causal genes, estimating their effects in some population, and transposing these effects to newly genotyped individuals (Lande and Thompson 1990; Meuwissen *et al.* 2001). These “gene effects” are substitution effects – the regression of the own phenotype (for polygenic risk scores) or expected progeny phenotypes (for estimated breeding values) on gene content at the locus. Being able to use substitution effects at causal genes across populations and generations is a goal of genomic prediction, QTL detection and also of causal mutation finding (Grisart *et al.* 2002).

There are several obstacles for these aims. Finding and validating causal genes and understanding their functional mechanism is extremely difficult (Grobet *et al.* 1997; Bonifati *et al.* 2003; Rupp *et al.* 2015). In practice, predictions are done using SNP markers using statistical genetics techniques. In livestock, use of markers results in very good predictions within populations, but mediocre (at best) predictions across populations, even with very sophisticated techniques (Hayes *et al.* 2009; Karoui *et al.* 2012; Porto-Neto *et al.* 2015; MacLeod *et al.* 2016). Indeed, livestock and human genetics empirical results show decreasing predictive ability with increasing genetic distance across distinct populations or generations (Liu *et al.* 2016; Martin *et al.* 2019). In humans, there is strong correlation between GWAS effect estimates across human populations, with typical values around 0.82

(Marigorta and Navarro 2013; Shi *et al.* 2021). The lack of perfect linkage disequilibrium (LD) across markers and genes has been claimed to be a reason for this decrease in accuracy. Adding extra information (more dense maps, biological prior information) should result in a better choice of markers close to causal genes, and therefore in a boost in predictive abilities across populations (Roos *et al.* 2009; MacLeod *et al.* 2016). However, in practice, the increase in predictive ability across populations is small at best (MacLeod *et al.* 2016; Moghaddar *et al.* 2019). This is against results from simulations (Roos *et al.* 2009; MacLeod *et al.* 2016) – but a problem in simulations is that typically gene effects are assumed to be biologically additive and therefore constant across populations.

We argue that, although imperfect LD is a likely cause for not being able to predict across populations, it is not the only one. In fact, substitution (statistical) additive gene action is not homogeneous across populations, even for exactly the same causal mutation. Examples in livestock genetics include myostatin gene (Aiello *et al.* 2018) or DGAT1 (Gautier *et al.* 2007). For instance, in the latter, the “K” allele had rather different substitution effects across breeds: -611, -142 and -351 kg of milk for the respective breeds Montbéliarde, Normande and Holstein, for a trait with a genetic standard deviation of ~600 kg. Although part of these differences may be due to genotype by environment interactions, it is also plausible that this is due to epistasis or dominance; for instance this is the case in DGAT1 (Streit *et al.* 2011). For instance, in the 5-loci epistatic network in Carlborg *et al.* (2006) some substitution effects at genes switch signs depending on genetic background. In *Drosophila*, estimated substitution effects of P-element insertions switched signs depending on the genetic background (Magwire *et al.* 2010; Mackay 2015).

There is indeed widespread evidence of biological epistasis (Mackay 2014). Whereas biological epistasis does not impede (on the contrary) large additive variation (Hill *et al.* 2008; Mackay 2014; Mäki-Tanila and Hill 2014), it **does imply** that substitution effects do vary across genetic backgrounds. Thus, in the presence of functional dominance and epistasis, there is no stability of substitution effects across different genetic backgrounds. Even if in all of these populations, additive variation accounts for most genetic variation, and additive substitution effects are sizable (Hill *et al.* 2008), substitution effects may differ across populations.

Recent simulations (Dai *et al.* 2020; Duenk *et al.* 2020) showed that the difference in substitution effects across populations may be quite large under non-additive biological gene action and increases with divergence of populations. It is relatively easy to derive algebraic expressions for substitution effects α assuming specific hypothesis of biological gene action. For instance, assuming biological additive and dominance effects only (Falconer and Mackay 1996) results in $\alpha = a + (q - p)d$, whereas assuming additive, dominance and additive by additive biological gene effects results, **assuming linkage equilibrium (LE)**, in $\alpha_1 = a_1 + (q_1 - p_1)d_1 + (p_2 - q_2)[aa]_{12}$ (Fuerst *et al.* 1997). Both of the approaches (simulation or analytical) are limited because the hypotheses of specific biological gene actions (*e.g.* 2-loci interaction but not 3-loci interaction) are too restrictive. Classically, these hypotheses are bypassed in quantitative genetics by working on statistical effects (Falconer and Mackay 1996; Lynch and Walsh 1998; Mäki-Tanila and Hill 2014).

The key parameter to describe the resemblance of statistical effects across populations is the correlation of substitution effects across populations. Under certain assumptions

(independence of allele frequencies and substitution effects, appropriate coding; (Wientjes *et al.* 2017)), this correlation can be estimated from SNP markers and data of two populations (Karoui *et al.* 2012; Wientjes *et al.* 2017), and can easily be accommodated into genomic prediction models (Karoui *et al.* 2012; Xiang *et al.* 2017). In human studies, the correlation is estimated through the meta-analysis of GWAS statistics (Marigorta and Navarro 2013; Shi *et al.* 2021). However, in addition to empirical results, some theory to describe the resemblance of substitution effects across different populations would be helpful to (1) better understand and quantify the change in *true* substitution effects (*i.e.* if true genes instead of markers were being used), (2) give upper bounds for genomic prediction across populations/generations, and (3) allow *a priori* planning of genomic predictions *i.e.* to include or not different subpopulations.

The aim of this work is to develop a theory to understand and predict, under a neutral scenario, the extent of change of substitution effects across space (breeds, lines) and across time (generations), without invoking or assuming specific modes of biological gene action. As a result, we obtain explicit estimators that are functions of additive, dominance and additive by additive variances, genetic distances across populations, and distributions of allele frequencies. We check and illustrate our theory using published results and simulations considering dominance and epistasis (additive by additive and complementary) from 5-loci interactions. Main factors affecting correlation of substitution effects across populations are their genetic distance and the extent of additive by additive variation, which is rarely large.

MATERIALS AND METHODS

Theory

General Theory. Here, we model difference of substitution effects across two populations as Taylor expansions around one of them, the “focal” population. Using Kojima’s method (Kojima 1959, 1961), we put additive substitution effects as a function of differences in allele frequencies across populations, “focal” additive substitution effects, and second order (dominance, additive by additive epistasis) statistical effects. From here, we show that the correlation of substitution effects across populations is (approximately) a function of their differentiation (or genetic distance), the additive, dominant and additive by additive genetic variances, the average heterozygosities and average squared heterozygosities. All these parameters can be estimated in real populations. In the following, we try to stick to Mäki-Tanila and Hill (2014) notation. Many details are given as Appendix in the Supplementary Material. Main notation is presented in Table 1.

Table 1. Notation.

$\alpha_i^b, \alpha_i^{b'}$	Substitution effect of locus i in population b , in population b'	$\sigma_{\alpha,b}^2$	Variance of substitution effects in population b
d_i^{*b}	Dominance deviation at locus i in population b	$\sigma_{d,b}^2, \mu_{d,b}$	Variance and mean of dominance deviations in population b
$(\alpha\alpha)_{ij}^b$	Additive by additive effect of loci i, j at population b	$\sigma_{(\alpha\alpha,b)}^2$	Variance of additive by additive effects in

			population b
$(\alpha\alpha)_i^b$	Vector of additive by additive effects of locus i with all other loci in population b		
$p_i^b, p_i^{b'}$	Allele frequency of locus i in population b , in population b'		
$\mathbf{p}^b, \mathbf{p}^{b'}$	Vectors of allele frequencies in population b , in population b'		
$\bar{H}_b$	Average heterozygosity at population b , $\bar{H}_b = E\left(2p_i^b(1 - p_i^b)\right)$		
$\overline{H_b^2}$	Average squared heterozygosity at population b , $\overline{H_b^2} = E\left(\left(2p_i^b(1 - p_i^b)\right)^2\right)$		
$\overline{HH}_b$	Average product of heterozygosities at population b , $\overline{HH}_b = E_{i>j}\left(4p_i^b(1 - p_i^b)p_j^b(1 - p_j^b)\right)$		
ϵ_i	Difference in allele frequencies at locus i , $\epsilon_i = p_i^{b'} - p_i^b$	σ_ϵ^2	Variance of the difference in allele frequencies across all loci
$D_{b,b'}$	Nei's minimum genetic distance		
$\sigma_A^2, \sigma_D^2, \sigma_{AA}^2$	Additive, Dominance and Additive by Additive variances (in population b)		

157

158

159 Note that our procedure is general – it does not invoke any particular mechanism for

160 epistasis or dominance, nor knowledge of individual QTL effects and locations. We assume

161 that the population mean is a (possibly complex) function of QTL allele frequencies $\mathbf{p}$ and

162 QTL functional or biological effects. The latter, albeit unknown, are assumed to be constant

163 across populations - we therefore **do not consider** genotype by environment interactions.

164

165 Consider the correlation $r(\alpha_i^b, \alpha_i^{b'})$ of substitution effects α^b and $\alpha^{b'}$ across respective

166 populations (breeds, heterotic groups, lines, or generations) b and b' . For simplicity, the

167 allele to which α refers is random – it can be either the wild or the mutant allele. In this

168 manner the average value of α is 0, even in presence of deleterious mutations. It is known

169 that, in presence of dominant and epistatic biological interactions, the value of α depends

170 on the allele frequencies and biological effects; for instance, $\alpha_1 = a_1 + (q_1 - p_1)d_1 +$

171 $(p_2 - q_2)[aa]_{12}$ for 2-locus epistasis (Fuerst *et al.* 1997), for $p_1 = 1 - q_1$ and $p_2 = 1 - q_2$

172 frequencies at loci 1 and 2, respectively. Inspired by this, we want to write the correlation

173 $r(\alpha_i^b, \alpha_i^{b'})$ for a locus i as a function of vectors of respective allele frequencies, $\mathbf{p}^b$ and $\mathbf{p}^{b'}$,

174 but in a general manner, without defining a particular functional or biological gene action.

175 We use a first order Taylor series expansion to approximate additive substitution effects in a

176 population b' , as a function of effects in another “focal” population b and their distance. By

177 doing this, it can be shown that the difference of substitution effects between two

178 populations is (approximately) a function of the genetic distance of the two populations, and

179 the magnitude of dominance and second order epistatic variances in the focal population b .

180

181

182

183

184 To derive additive substitution effects α as function of allelic frequencies $\mathbf{p}$, we use Kojima’s

185 definition of statistical effects as first, second... derivatives of the mean of the population (μ)

186 as a function of $\mathbf{p}$. We don’t invoke any explicit function – we just presume that there is

187 one, in other words, change in allele frequencies of the population implies change in the

total average genotypic value. Using Kojima's method, the additive substitution effect at the i -th locus is the first derivative (Kojima 1959, 1961):

$$\alpha_i = \frac{1}{2} \frac{\partial \mu}{\partial p_i}$$

Higher order statistical effects implying locus i (i.e. dominance deviations and epistatic interactions) can be represented by higher order partial derivatives of μ or equivalently as derivatives of α_i . The dominance deviation at the i -th locus (that we denote as d^* to distinguish from the biological or functional effect d (Falconer and Mackay 1996)) is:

$$d_i^* = -\frac{1}{4} \frac{\partial^2 \mu}{\partial p_i^2} = -\frac{1}{2} \frac{\partial \alpha_i}{\partial p_i}$$

The negative sign comes because the dominance deviation is usually understood as a feature of *heterozygosity*, in other words, it is of opposite sign than the increase of homozygosity in ∂p_i^2 . Last, the epistatic pairwise deviation of locus i with j is

$$(\alpha\alpha)_{ij} = \frac{1}{4} \frac{\partial^2 \mu}{\partial p_i \partial p_j} = \frac{1}{2} \frac{\partial \alpha_i}{\partial p_j}$$

This is positive because it is the effect of increasing both p_i and p_j . Note that interaction of order k implies k -th order derivative with scaling factor $1/2^k$.

Kojima's method shows, therefore, that higher order effects of one locus are derivatives of lower order effects. With these elements we can make a Taylor order expansion of α_i around frequencies in the "focal" population, p^b , so that $p^{b'} = p^b + \epsilon$, such that from values of α in b and changes in allele frequencies $\epsilon = p^{b'} - p^b$ we create a function

207 $\alpha_i^{b'} \approx f(\alpha_i^b, \epsilon)$. In the Appendix (section 1.1), we show that the Taylor linear approximation
 208 of the substitution effects of population b' : $\alpha_i^{b'}$, from effects from populations $b, \alpha_i^b, d_i^{*b},$
 209 $(\alpha\alpha)_i^b$ is:

$$\alpha_i^{b'} \approx \alpha_i^b + 2\epsilon_i(-d_i^{*b}) + 2\epsilon'(\alpha\alpha)_i^b \quad (1)$$

210
 211 where we use differences in allele frequencies ϵ , d_i^{*b} is the statistical dominance deviation
 212 at the locus i and $(\alpha\alpha)_i^b$ is a vector containing epistatic substitution effects of locus i with
 213 the rest of loci. By convention we assign $(\alpha\alpha)_{ii}^b = 0$.

214
 215 From equation 1, the covariance across two populations b and b' of the two substitution
 216 effects α_i^b and $\alpha_i^{b'}$ of the locus i is:

$$\text{Cov}(\alpha_i^b, \alpha_i^{b'}) \approx \text{Cov}(\alpha_i^b, \alpha_i^b + 2\epsilon_i(-d_i^{*b}) + 2\epsilon'(\alpha\alpha)_i^b) = \text{Var}(\alpha_i^b) \quad (2)$$

217
 218 The equality holds because terms $\text{Cov}(\alpha_i^b, 2\epsilon_i(-d_i^{*b}))$ and $\text{Cov}(\alpha_i^b, 2\epsilon'(\alpha\alpha)_i^b)$ are null,
 219 given that the different statistical effects $(\alpha_i^b, d_i^{*b}$ and $(\alpha\alpha)_i^b)$ are mutually orthogonal by
 220 construction.

221 Thus $r(\alpha_i^b, \alpha_i^{b'}) = \sqrt{\frac{\text{Var}(\alpha_i^b)}{\text{Var}(\alpha_i^b)\text{Var}(\alpha_i^{b'})}}$, and now we need the variance of $\alpha_i^{b'}$ as a function of
 222 effects in population b , this is (see the Appendix, section 1.2 for details):

223

$$\begin{aligned} Var(\alpha_i^{b'}) &\approx Var(\alpha_i^b + 2\epsilon_i(-d_i^{*b}) + 2\epsilon'(\alpha\alpha)_i^b) \\ &= Var(\alpha_i^b) + 4Var(\epsilon_i) \left(Var(d_i^{*b}) + E^2(d_i^{*b}) \right) \\ &\quad + 4tr \left(Var(\epsilon')Var((\alpha\alpha)_i^b) \right) \end{aligned} \tag{3}$$

224

225

226 where $Var(\epsilon')$ and $Var((\alpha\alpha)_i^b)$ are matrices:

$$Var(\epsilon') = \begin{pmatrix} Var(\epsilon_1) & Cov(\epsilon_1, \epsilon_2) & Cov(\epsilon_1, \epsilon_n) \\ Cov(\epsilon_2, \epsilon_1) & Var(\epsilon_2) & Cov(\epsilon_2, \epsilon_n) \\ & & \dots \\ Cov(\epsilon_n, \epsilon_1) & & Var(\epsilon_n) \end{pmatrix}$$

227 which represents how loci differ. A locus that may diverge a lot has high $Var(\epsilon)$; two loci in

228 strong linkage will show non-zero $Cov(\epsilon_1, \epsilon_2)$. Also

$$Var((\alpha\alpha)_i^b) = \begin{pmatrix} Var((\alpha\alpha)_1^b) & Cov((\alpha\alpha)_1^b, (\alpha\alpha)_2^b) & Cov((\alpha\alpha)_1^b, (\alpha\alpha)_n^b) \\ Cov((\alpha\alpha)_2^b, (\alpha\alpha)_1^b) & Var((\alpha\alpha)_2^b) & Cov((\alpha\alpha)_2^b, (\alpha\alpha)_n^b) \\ & & \dots \\ Cov((\alpha\alpha)_n^b, (\alpha\alpha)_1^b) & & Var((\alpha\alpha)_n^b) \end{pmatrix}$$

229 where $Var((\alpha\alpha)_1^b)$ contains the variance of the epistatic effect of marker i with marker 1,

230 and so on. We assume $Var((\alpha\alpha)_i^b) = I \sigma_{(\alpha\alpha,b)}^2$, i.e. epistatic terms $(\alpha\alpha)_i^b$ have a variance

231 $\sigma_{(\alpha\alpha,b)}^2$ and are a priori uncorrelated to each other. In the previous product

232 $tr \left(Var(\epsilon')Var((\alpha\alpha)_i^b) \right)$, the use of the trace implies that off-diagonal elements of

233 $Var(\epsilon')$ are ignored. Thus, assuming that all diagonal elements of $Var(\epsilon')$ have the same

234 common variance σ_ϵ^2 , this results in $4tr \left(Var(\epsilon')Var((\alpha\alpha)_i^b) \right) = 4n\sigma_\epsilon^2\sigma_{(\alpha\alpha,b)}^2$

235

236

237 The expression $Var(\epsilon_i d_i^{*b}) = Var(\epsilon_i) (Var(d_i^{*b}) + E^2(d_i^{*b}))$ is detailed in the Appendix
 238 (section 1.2), and it shows that both the variability of dominance deviations $Var(d_i^{*b}) =$
 239 $\sigma_{d,b}^2$ and its mean $E(d_i^{*b}) = \mu_{d,b}$ which, if not zero, can be understood as the basis of
 240 inbreeding depression, enter into the expression. Also, we assume ϵ (change in allele
 241 frequencies, but not the allele frequencies themselves) and the different non-additive
 242 effects d_i^{*b} and $(\alpha\alpha)_i^b$ to be independent (uncorrelated). This makes sense as loci may be
 243 selected for additive effects but not for non-additive effects.

244
 245 In expression (3) we need the variances of the different genetic effects and the variance of
 246 changes in allele frequencies. We address these two terms in turn.

247
 248 We factorize the variance of statistical additive, dominant and additive by additive effects as
 249 follows (Mäki-Tanila and Hill 2014; Vitezica *et al.* 2017):

$$Var(\alpha_i^b) = \sigma_{\alpha,b}^2 = \frac{\sigma_A^2}{nH_b}$$

$$Var(d_i^{*b}) + E^2(d_i^{*b}) = \sigma_{d,b}^2 + \mu_{d,b}^2 = \frac{\sigma_D^2}{nH_b^2}$$

250 (the latter is shown in the Appendix, section 1.4) and

$$Var((\alpha\alpha)_{i,j,j>i}^b) = \sigma_{(\alpha\alpha,b)}^2 = \frac{\sigma_{AA}^2}{n(n-1)\overline{H_b}\overline{H_b}} \approx 2 \frac{\sigma_{AA}^2}{n^2\overline{H_b}\overline{H_b}}$$

251 for n the number of QTL loci and using functions of heterozygosities (more details in the
 252 Appendix, section 1.5):

$$\overline{H_b} = E(2p_i^b(1-p_i^b))$$

$$\overline{H_b^2} = E(4p_i^b(1-p_i^b)p_i^b(1-p_i^b))$$

$$\overline{HH_b} = E_{i>j} \left(2p_i^b(1-p_i^b)2p_j^b(1-p_j^b) \right) \approx \frac{1}{2} \overline{H_b} \overline{H_b}$$

253
254 Here we have assumed independence of QTL allele frequencies and QTL effects. All
255 variances and effects, as well as heterozygosities H_b , refer to the focal population b with
256 allele frequencies p^b and effects α^b . Note that we assume HWE and LE within both b and b' .
257 Of course, we don't know allele frequencies or even numbers of true causal genes, but the
258 first can be prudently guessed and the later cancel out in the following.

259
260 Consider the scalar σ_ϵ^2 , the variance of differences of allele frequency. In fact, $\sigma_\epsilon^2 =$
261 $Var(p^{b'} - p^b) = E((p^{b'} - p^b)^2) - E(p^{b'} - p^b)^2 = E((p^{b'} - p^b)^2)$ because $E(p^{b'} -$
262 $p^b) = 0$ when averaged across loci. Therefore, $\sigma_\epsilon^2 = E((p^{b'} - p^b)^2)$ which corresponds to
263 Nei's "minimum genetic distance" (Nei 1987; Caballero and Toro 2002), and, accordingly, will
264 be called $D_{b,b'} = \sigma_\epsilon^2$. The value of $D_{b,b'}$ can be estimated from marker data as $\hat{D}_{b,b'} =$
265 $\frac{1}{n} \sum (p_i^{b'} - p_i^b)^2$, although it is sensitive to the spectra of polymorphisms (e.g. SNP chips vs
266 sequencing). In addition, $D_{b,b'}$ is also the numerator of the F_{ST} fixation index, e.g.

267 Hudson's (1992) $F_{ST} = \frac{E((p_i^{b'} - p_i^b)^2)}{E(p_i^{b'}(1-p_i^b) + p_i^b(1-p_i^{b'}))}$ with an estimator

268 $\hat{F}_{ST} = \frac{\frac{1}{n} \sum_i (p_i^{b'} - p_i^b)^2}{\frac{1}{n} \sum_i (p_i^{b'}(1-p_i^b) + p_i^b(1-p_i^{b'}))}$ (Hudson *et al.* 1992; Bhatia *et al.* 2013). In principle, $D_{b,b'}$

269 and F_{ST} can be estimated from markers, but also from evolutionary distances and effective
270 population size, or both (Weir and Hill 2002; Bonhomme *et al.* 2010). Now we can rewrite
271 the last term of equation (3) as

$$4tr\left(Var(\epsilon')Var((\alpha\alpha)_i^b)\right) = 4n(\sigma_\epsilon^2\sigma_{(\alpha\alpha,b)}^2) = 4nD_{b,b'}\frac{2\sigma_{AA}^2}{n^2\bar{H}_b\bar{H}_b}$$

272 Combining expressions above, equation (3) becomes

$$Var(\alpha_i^{b'}) = \quad (4)$$

$$\begin{aligned} \frac{\sigma_A^2}{n\bar{H}_b} + 4D_{b,b'}\frac{\sigma_D^2}{n\bar{H}_b^2} + 4nD_{b,b'}2\frac{\sigma_{AA}^2}{n^2\bar{H}_b\bar{H}_b} = \\ \frac{1}{n}\left(\frac{\sigma_A^2}{\bar{H}_b} + 4D_{b,b'}\frac{\sigma_D^2}{\bar{H}_b^2} + 8D_{b,b'}\frac{\sigma_{AA}^2}{\bar{H}_b\bar{H}_b}\right) \end{aligned}$$

273

274

275 From here the correlation of α across populations is (factorizing out $\frac{1}{n}$)

$$\begin{aligned} r(\alpha_i^b, \alpha_i^{b'}) &\approx \frac{Cov(\alpha_i^b, \alpha_i^{b'})}{\sqrt{(Var(\alpha_i^b))(Var(\alpha_i^{b'}))}} = \quad (5) \\ &= \frac{Var(\alpha_i^b)}{\sqrt{(Var(\alpha_i^b))(Var(\alpha_i^{b'}))}} = \frac{\sqrt{Var(\alpha_i^b)}}{\sqrt{Var(\alpha_i^{b'})}} = \\ &= \frac{\sqrt{\frac{\sigma_A^2}{\bar{H}_b}}}{\sqrt{\frac{\sigma_A^2}{\bar{H}_b} + 4D_{b,b'}\frac{\sigma_D^2}{\bar{H}_b^2} + 8D_{b,b'}\frac{\sigma_{AA}^2}{\bar{H}_b\bar{H}_b}}} \end{aligned}$$

276

277 Factorizing out $\bar{H}_b$ we arrive to the slightly clearer expression

$$r(\alpha_i^b, \alpha_i^{b'}) \approx \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + D_{b,b'} \left(4 \frac{\bar{H}_b}{H_b^2} \sigma_D^2 + 8 \frac{1}{H_b} \sigma_{AA}^2 \right)}} \quad (6)$$

which shows well that the correlation is a function of distance across populations ($D_{b,b'}$) and weights of additive vs. non-additive variances. Note that $D_{b,b'}$ is always positive, which implies that $0 < r(\alpha_i^b, \alpha_i^{b'}) < 1$ as expected.

The quantities involved in equation (6) are (1) Nei's "minimum genetic distance" $D_{b,b'}$, which describes the similarity of populations b and b' , (2) the variance of statistical additive, dominant and additive by additive effects at the individual level $\sigma_A^2, \sigma_D^2, \sigma_{AA}^2$ in population b (3) first and second moments of heterozygosities $\bar{H}_b, \bar{H}_b^2$. All these values can be, in principle, estimated from data or "prudently guessed". For the particular case of heterozygosities, SNP markers are a poor choice and it is likely better to use a guess based on sequence or evolutionary processes (coalescence).

From the definition of F_{ST} and assuming $\bar{H}_b \approx \bar{H}_{b'}$, $D_{b,b'} \approx \frac{F_{ST}}{1-F_{ST}} \bar{H}_b$ (detailed in the Appendix, section 1.6), leading to

$$r(\alpha_i^b, \alpha_i^{b'}) \approx \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \frac{F_{ST}}{1-F_{ST}} \left(4 \frac{(\bar{H}_b)^2}{H_b^2} \sigma_D^2 + 8 \sigma_{AA}^2 \right)}} \quad (7)$$

The advantage of the F_{ST} is that it is more robust to the spectra of allele frequencies used to estimate it (Bhatia *et al.* 2013). If we further assume that dominance variance σ_D^2 is negligible, we can write a neat expression of the correlation in terms of F_{ST} :

$$r(\alpha_i^b, \alpha_i^{b'}) \approx \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \frac{8F_{ST}}{1-F_{ST}}\sigma_{AA}^2}} \approx \sqrt{1 - \frac{8F_{ST}}{1-F_{ST}}\frac{\sigma_{AA}^2}{\sigma_A^2}} \quad (8)$$

The second approximation involving $\frac{1}{1+x} \approx 1 - x$. This expression (8) plainly tells that the squared correlation of gene substitution effects across two population is (to a few degrees of approximation) a linear function of the similarity of populations and the additive by additive variance. To our knowledge, these results showing the importance of the different factors on the difference between substitution effects had been shown before only through simulations.

Thus, the algorithm to estimate *a priori* the correlation of α across populations b and b' , $r(\alpha_i^b, \alpha_i^{b'})$, is:

1. Estimate in population b
 - a. additive, dominance, and additive by additive variances
 - b. average heterozygosity $\overline{H}_b$ and average squared heterozygosity $\overline{H}_b^2$
2. Estimate Nei's distance $D_{b,b'}$ and/or F_{ST} of the two populations
3. Apply equation (6) or (7).

Consideration of directionality of substitution effects

In Evolutionary Genetics it is reasonable to think that most mutations are deleterious, thus with a negative effect of the mutant allele. In Medical Genetics reports of estimated substitution effects are also often done in terms of “susceptible” alleles. In both cases $E(\alpha) \neq 0$ or even $\alpha > 0$ for all loci. In both cases α is “oriented” and has no zero mean. In the theory above we have considered that α is the effect of a randomly drawn allele, which leads to $E(\alpha) = 0$ and enormously simplifies the algebra. In order to consider oriented α , we propose to transform the estimate of $r(\alpha_i^b, \alpha_i^{b'})$ into $r(|\alpha_i^b|, |\alpha_i^{b'}|)$, the correlation of the absolute values. Assuming that α follows a normal distribution, $|\alpha|$ follows a so-called folded normal distribution. From here, $r(|\alpha_i^b|, |\alpha_i^{b'}|)$ is obtained from $r(\alpha_i^b, \alpha_i^{b'})$ using expressions (not detailed here) in (Kan and Robotti 2017), conveniently programmed in the R package MomTrunc (Galarza *et al.* 2020). The specific R function is in the Appendix, section 1.7, and we will call it r2rabs().

Covariance across generations within one population. The definition above of two populations is general enough that we can consider any two populations, *e.g.* two breeds (Angus and Hereford), two strains (New Zealand Holstein and US Holstein) or two generations or time frames (*e.g.* animals born in 2000 vs. animals born in 2005, or animals born in 2000 vs. their descendants). There is evidence that across-generations genetic correlation decreases with (many) generations to values as low as 0.6 (Tsuruta *et al.* 2004; Haile-Mariam and Pryce 2015). Part of this is likely due to genotype by environment interactions. Anyway, part of the across-generation genetic correlation could be due to changes in the allele frequency due to drift, and therefore it can be accounted for by our model, based on the evolution of average coancestry in the breed. We develop this next.

We will talk about “generations” but ideas apply to pedigrees with overlapping generations as well. Consider that what we previously called populations b and b' are animals born at

time t_1 and t_2 , with $t_2 > t_1$. Equation (7) becomes $r(\alpha_i^{t_1}, \alpha_i^{t_2}) \approx \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \frac{F_{ST}}{1-F_{ST}} \left(4 \frac{(\bar{H}_{t_1})^2}{H_{t_1}^2} \sigma_D^2 + 8\sigma_{AA}^2 \right)}}$.

In pedigree-based context, the F_{ST} is simply half the increase in average relationship from t_1 to t_2 (Powell *et al.* 2010) which is approximately equal to the increase in inbreeding from $t_1 + 1$ to $t_2 + 1$, which in turn is approximately $(t_2 - t_1)\Delta F$ for small values of ΔF and steady increase of inbreeding. Thus, $F_{ST} \approx \frac{(t_2 - t_1)\Delta F}{2}$ to obtain

$$r(\alpha_i^{t_1}, \alpha_i^{t_2}) \approx \frac{\sqrt{\sigma_A^2}}{\sqrt{\sigma_A^2 + \frac{\frac{(t_2 - t_1)\Delta F}{2}}{1 - \frac{(t_2 - t_1)\Delta F}{2}} \left(4 \frac{(\bar{H}_{t_1})^2}{H_{t_1}^2} \sigma_D^2 + 8\sigma_{AA}^2 \right)}}$$

Assuming like in (8) small values of dominance variance and of F_{ST} , we obtain

$$r(\alpha_i^b, \alpha_i^{b'}) \approx \sqrt{1 - 4(t_2 - t_1)\Delta F \frac{\sigma_{AA}^2}{\sigma_A^2}}$$

Thus, the correlation of substitution effects decreases when there is large drift, reflected in high values of ΔF . This is the case for instance if parents of the next generation are very highly selected without restrictions in future inbreeding, resulting in a considerable change in allele frequencies over the generations.

This agrees with classical theory (Falconer and Mackay 1996): the variance in change of allele frequencies from one generation to the next is simply ΔF . Typical values of ΔF in

livestock (rabbits, pigs, cattle or sheep) are at most 0.01 per generation (Welsh *et al.* 2010; García-Ruiz *et al.* 2016; Fernández *et al.* 2017; Rodríguez-Ramilo *et al.* 2019), although this is potentially changing with genomic selection.

Examples

Estimates of correlation across populations using literature values. From values of the literature, the previous expressions, equations (6) and (7) were applied to obtain guesses of the correlation of substitution effects across different populations and generations within population. Estimates of statistical non-additive variation are scarce, as are estimates of distances across populations in terms of covariance of allele frequencies. The differentiation F_{ST} Jersey and Holstein breeds of cattle obtained from (VanRaden *et al.* 2011) is 0.06 and among Landrace and Yorkshire is 0.16 (Xiang *et al.* 2017).

Table 2 includes literature estimates for additive, dominance and epistatic variation. We used estimates for milk yield in Simmental cattle (Fuerst and Sölkner 1994) as if it was Holstein, and for litter size in a commercial pig line (Vitezica *et al.* 2018) as if it was Landrace, because we could not find estimates in the same populations. These estimates (as most estimates of non-additive variances) are inaccurate and are presented here only as examples. In the particular case of cattle, the estimated value of σ_{AA}^2 is higher than usually expected *a priori* (Hill *et al.* 2008), so we take it as an example of an extreme, but not impossible, case.

Table 2. Variance component estimates, as ratio from phenotypic variance, from literature

Species	σ_A^2	σ_D^2	σ_{AA}^2
Cattle	0.20	0.09	0.15
Pigs	0.092	0.020	0.016

Estimates for milk yield in cattle (Fuerst and Sölkner 1994) and litter size in pigs (Vitezica *et al.* 2018).

Then, we assumed three distributions for QTL allele frequencies, following Hill *et al.* (2008): a uniform distribution (“Uniform”), and a U-shaped distributions with $f(p) \propto p^{-1}(1-p)^{-1}$ with effective population size of $Ne = 50$ (“Hill”), bounded at $\left[\frac{1}{2Ne}, 1 - \frac{1}{2Ne}\right]$. We also assumed a Beta(0.04, 0.04) distribution, which is an extreme U-shaped distribution (“Extreme”) in which roughly 80% of the QTLs have minor allele frequency lower than 0.01. This results in respective values of $\overline{H_b} = (0.333, 0.107, 0.037)$ and $\overline{H_b^2} = (0.133, 0.018, 0.013)$. Using the values described above, we estimated the correlation of QTL substitution effects across breeds, $r(\alpha_i^b, \alpha_i^{b'})$, using equation (7).

Similarly, the correlation of QTL substitution effects a certain number of generations away, $r(\alpha_i^{t_2}, \alpha_i^{t_1})$, was obtained applying Equation (7). An increase $\Delta F = 0.01$ per generation was assumed, for the same assumed variances and allele frequency distributions.

Simulations. The objective of the simulation is to verify our results considering different kinds of non-additive biological gene action, and not to infer the values of the correlation of

substitution effects across populations as this requires realistic presumptions of the forms and magnitudes of epistatic interactions, something that is largely unknown.

We used *macs* (Chen *et al.* 2009) to simulate a “cattle” scenario of two domestic cattle breeds which diverged from a common ancestral population, in the lines of (Pérez-Enciso 2014; Pérez-Enciso *et al.* 2015), where a large population had a 10-fold reduction bottleneck (domestication) 2000 generations ago and a split into two populations t generations ago, where we made t oscillate between 0 and 100 by steps of 10. Parameters in the simulation were tailored (Pérez-Enciso 2014; Pérez-Enciso *et al.* 2015) to mimic observed levels of diversity in cattle (Gibbs *et al.* 2009) and lead to F_{ST} values between 0 and 0.15. We considered 100 DNA stretches of 300 Kb each. We simulated 200 individuals per population, from which we obtained allele frequencies per population. Details are provided in the Appendix, section 1.8. This provided ~400,000 segregating loci with realistic allele frequencies (L-shaped distribution of allele frequencies).

To simulate non-additive biological gene action, we considered three scenarios. All of them involved 5000 single marker polymorphisms drawn at random from the simulated ones with no particular restriction. First scenario was Complete Dominance (within locus); second one was 1000 networks of 5 loci in Complementary Epistasis, and the third one 1000 networks of 5 loci with Multiplicative Epistasis (additive by additive). We also simulated networks of 2 and of 10 loci with similar results (not shown). These scenarios are similar to Hill *et al.* (2008) but instead of considering 2 loci we consider 5. We use these forms of epistasis, as the first two ones may be interpreted as biologically meaningful gene actions, and the third one is an

extreme case for the change in substitution effect across populations. In addition, all three scenarios are analytically tractable. Complete dominance has the genotypic value of the heterozygote equal to one of the homozygotes, e.g. the presence of a single copy of the “good” allele is enough to, say, avoid the disease. Complementary epistasis can be seen as a multi-loci dominance, e.g. disease happens when there is a recessive deleterious genotype at any of the loci. Additive by additive epistasis can be understood as a pure multiplication of gene contents and has no good biological interpretation. Networks contribute additively to the total phenotype. Note that, by definition, functional gene action is the same across all populations. From the description of the non-additive gene action and some algebra in the Appendix, we are able to derive analytically the values of α , d^* and $(\alpha\alpha)$ in each population and variances σ_A^2 , σ_D^2 and σ_{AA}^2 . Details are given in the Appendix, section 1.9. For each value of t , we made 10 replicates and averaged the results.

From the true substitution effects α^b and $\alpha^{b'}$ derived above, we obtained the true value $r(\alpha^b, \alpha^{b'})$. Note that because of the coalescent simulation, all reference alleles (i.e. with frequency p) are “mutant” ones, and due to assumed dominance and epistatic actions, α are negative by construction (i.e. the mutant allele is deleterious). To conciliate this fact with our derivations, that assume that α refers to a random allele and has null means, we did two things: (1) compute a “random allele” version of $\alpha^b, \alpha^{b'}$ in which α was changed sign for “odd” loci, and (2) estimate $r(\alpha^b, \alpha^{b'})$ using the transformation of normal distribution into folded normal (Kan and Robotti 2017), i.e. the `r2rabs()` function mentioned above. Thus, we have two estimands, the correlation for the “mutant allele” effect $r_{mutant}(\alpha^b, \alpha^{b'})$ (which corresponds e.g. to typical use in Evolutionary and Medical Genetics) and the correlation for

the “random allele” effect $r_{random}(\alpha^b, \alpha^{b'})$ (which corresponds, *e.g.* to genomic selection and some GWAS). By construction, $r_{mutant}(\alpha^b, \alpha^{b'}) < r_{random}(\alpha^b, \alpha^{b'})$. For instance, in one of the scenarios $r_{mutant}(\alpha^b, \alpha^{b'}) = 0.67$ and $r_{random}(\alpha^b, \alpha^{b'}) = 0.85$.

Now we describe the estimators. We considered either the use of “all” polymorphisms, or of a “SNP” selection in which $MAF > 0.01$ across both populations simultaneously (roughly 40,000 polymorphisms), as this corresponds to typical SNP panels in genetic improvement. Then we used the equation (6) either for “all” or for “SNP”, using their frequencies to obtain $D_{b,b'}$, $\bar{H}_b$ and $\bar{H}_b^2$. In “all”, the spectra of allele frequencies of polymorphisms and of QTL is the same, but not in “SNP”. As “SNP” tends to be biased, we also considered the equation (7) using F_{ST} estimated from SNPs and $\bar{H}_b = 0.107$, and $\bar{H}_b^2 = 0.018$ from the “Hill” U-shaped distribution mentioned above. Note that F_{ST} is more robust to the spectra of allele frequencies used to estimate it. Thus, we obtained three estimators for “random allele” $r_{random}(\alpha^b, \alpha^{b'})$: $\hat{r}_{all}$, $\hat{r}_{SNP}$ and $\hat{r}_{SNPFST}$, and also the three corresponding estimators for “mutant allele” $r_{mutant}(\alpha^b, \alpha^{b'})$ applying function `r2rabs()` to $\hat{r}_{all}$, $\hat{r}_{SNP}$ and $\hat{r}_{SNPFST}$.

Data Availability Statement. Appendix associated with this manuscript is in https://figshare.com/articles/online_resource/Appendix_final_v4_pdf/14509941. Code and files for simulations can also be found in https://figshare.com/articles/software/scripts_zip/14509956. All other results can be reproduced using equations and figures from the text.

RESULTS

Estimates of correlation across populations using literature values. The results for reasonable guesses of the distribution of allele frequencies at QTL are in Table 3. The correlation of QTL substitution effects across breeds was roughly 0.85 across both species (pigs and cattle) and all assumed distributions of gene frequencies. These numbers are higher than scarce estimates of genetic correlations in the literature: 0.4 to 0.8 for production traits (Karoui *et al.* 2012), 0.3 to 0.5 for milk yield (Legarra *et al.* 2014) and 0.6 for body condition (Porto-Neto *et al.* 2015), and we consider that this is because across-breeds genetic correlations also include genotype by environment interactions. It may be argued that these studies used SNPs instead of true QTLs; however, Wientjes *et al.* (2017) in a simulation study obtained unbiased estimates of correlation across QTLs using SNP markers for a true correlation of 0.5, and (Cuyabano *et al.* 2018) showed analytically that when SNP markers yield correct estimates of true relationships at QTL, estimates of genetic parameters are unbiased.

Table 3: Estimates of correlations of QTL effects across breeds based on values from Table 2 and Equation 7, for different distributions of QTL frequencies

Species	Uniform	Hill ¹	Extreme ²
Cattle	0.83	0.82	0.84
Pigs	0.87	0.85	0.88

¹ U-shaped distribution with effective population size of 50

² Beta(0.04, 0.04) distribution

Concerning decrease of correlations across generations, results (Table 4) show a slow decrease, and this decrease is, again, not sensitive to the assumed distribution of allelic frequencies. Values in Table 4, quite close to 1, somehow disagree with few existing estimates of across-generation genetic correlations, clearly lower than 1 (Tsuruta *et al.* 2004; Haile-Mariam and Pryce 2015). Again, part of the correlation lower than 1 is probably due to non-additive gene action, and part due to genotype by environment interactions.

Table 4: Correlation of QTL effects within breed across time, based on values from Table 1 and Equation 7, for different distributions of QTL frequencies

Species	Distance in generations	Uniform	Hill ¹	Extreme ²
Cattle	1	0.98	0.97	0.98
	2	0.97	0.94	0.97
	5	0.93	0.87	0.93
	10	0.86	0.78	0.87
Pigs	1	1.00	0.99	1.00
	2	0.99	0.98	0.99
	5	0.98	0.96	0.97
	10	0.96	0.93	0.93

¹ U-shaped distribution with effective population size of 50

² Beta(0.04, 0.04) distribution

Simulations. Simulated variance components (expressed as ratios from total genetic variance σ^2) with marker genotypes and with QTL genotypes are shown in Table 5. All scenarios yield high additive genetic variances, as expected.

Table 5. Additive, dominance and additive by additive variances in the simulated population

	additive	dominance	additive by additive
Complete Dominance	252	113	0
Complementary	130	53	12
Epistasis			
Multiplicative	219	0	122

True simulated values of correlation across substitution effects $r_{random}(\alpha^b, \alpha^{b'})$ and their estimates $\hat{r}_{all}$, $\hat{r}_{SNP}$ and $\hat{r}_{SNPFst}$ are presented in Figure 1 (for α defined for random alleles) and Figure 2 (for α defined for mutant alleles). Generally speaking, Figure 1 applies to non-fitness related traits (for “random”) and Figure 2 to fitness-related traits (for “mutant”). As predicted by our derivations, there is a clear and almost linear decrease of $r(\alpha^b, \alpha^{b'})$ with increasing values of F_{ST} .

In Figure 1 (for α defined for random alleles) it can be observed that our expressions (6-7) tend to slightly over-estimate $r_{random}(\alpha^b, \alpha^{b'})$ for Complete Dominance, and Complementary Epistasis. The estimates are quite good for Multiplicative Epistasis. Estimators using SNP markers are slightly less accurate than using the complete

polymorphism spectra, and using F_{ST} from SNP markers, coupled with a guess of heterozygosities, is similar to using SNP alone to infer both distances and heterozygosities. Figure 2 (for α defined for mutant alleles) presents values of $r_{mutant}(\alpha^b, \alpha^{b'})$ and estimates $\hat{r}_{all}$, $\hat{r}_{SNP}$ and $\hat{r}_{SNPFst}$. Estimating $r_{mutant}(\alpha^b, \alpha^{b'})$ is more difficult than estimating $r_{random}(\alpha^b, \alpha^{b'})$: depending on the scenario there is more over- or under-estimation than for $r_{random}(\alpha^b, \alpha^{b'})$. In this case the effect of using all polymorphisms or SNP is more marked. However, overall, we find that our estimators explain well the decay in $r(\alpha^b, \alpha^{b'})$. The imperfect disagreement (both in Figures 1 and 2) is probably due to several wrong assumptions; we comment some of them in the Discussion.

529
530 Figure 1. Simulated (straight black line) and estimated (points) correlation across QTL
531 substitution effects of random alleles across two populations as a function of their F_{ST}
532 differentiation coefficient. r_{HatAll} : estimates with all polymorphism. r_{HatSNP} : estimates
533 using SNP-like loci. $r_{\text{HatSNPFst}}$: estimates using SNP-like loci with a correction for
534 heterozygosity. Results of 10 replicates per point with s.e.<0.01.

535

Figure 2. Simulated (straight black line) and estimated (points) correlation across QTL substitution effects of mutant alleles across two populations as a function of their F_{ST} differentiation coefficient. r_{HatAll} : estimates with all polymorphism. r_{HatSNP} : estimates using SNP-like loci. $r_{\text{HatSNPFst}}$: estimates using SNP-like loci with a correction for heterozygosity. Results of 10 replicates per point with s.e.<0.01.

DISCUSSION

The fact that most genetic variation is statistically additive is well known. However, this does not imply equality of statistical additive effects across populations, something that is possibly reflected in the difficulty of predicting breeding values or disease risks across (distinct) populations, either in animal or human genetics. Only recently has non-additive biological gene action been identified as playing a role in determining the degree to which prediction across populations is successful (Dai *et al.* 2020; Duenk *et al.* 2020). These authors presented result of simulations, where the problem is that the results are scenario specific and do not allow a good appraisal of the different factors. Having a theory allows to derive more general expressions and understanding factors influencing across-population differences in substitution effects.

In this work, we present a general, formal framework that does not depend on specific hypothesis regarding gene action or order of the epistatic interactions. In our derivation, we approached *high-order functional* epistasis by Taylor expansions, leading to expressions that involve only *low-order statistical* additive epistasis (actually pairwise). Including one extra term in our Taylor expansion would include *three-loci statistical* epistasis and so on, but extra terms would lead to more difficult expressions (covariance of allele frequencies across loci and populations), and it is expected that the magnitude of the statistical effects is smaller and smaller with higher orders of interactions (Mäki-Tanila and Hill 2014). We find reasonable agreement with our simulation and also with values from literature. However, our simulations may not be particularly realistic, something that would require considerable thinking on how to simulate biologically meaningful epistasis mechanisms for a variety of traits. We see them as building blocks of non-additive architecture. At any rate, the three

scenarios generated sizeable non-additive variation which is a challenging case for our expressions.

Three main factors influence the correlation of substitution effects between populations $r(\alpha^b, \alpha^{b'})$: the genetic similarity of the two populations, the magnitudes of additive, dominance and additive by additive variances, and the distribution of allele frequencies at QTL. We consider that showing explicitly these three factors is an achievement, as their role is implicit, yet not explicitly shown, in previous works in simulated and real data (Martin *et al.* 2019; Dai *et al.* 2020; Duenk *et al.* 2020). Now we discuss these three factors.

The distance across populations is summarized by Nei's minimum genetic distances $D_{b,b'}$ or F_{st} indexes. Under pure drift scenarios, these depend on divergence times and effective population sizes (Weir and Hill 2002; Bonhomme *et al.* 2010; Walsh and Lynch 2018).

The factors $\frac{\overline{H_b}}{H_b^2}$ and $\frac{1}{H_b}$ are weighting factors on dominance and additive by additive variances.

If the allele frequencies are modelled using symmetric $Beta(a, a)$ distributions (see

Appendix, section 1.10), these become $\frac{\overline{H_b}}{H_b^2} = \frac{1}{2^{\frac{a+1}{a}} \left(1 + \frac{a+2}{2a+2} \left(\frac{a+3}{2a+3} - 2 \right) \right)}$ and $\frac{1}{H_b} = 2 + \frac{1}{a}$. The first is

bounded between 3 (for U-shaped distributions) and 2 (for peaked distributions), so that

dominance variation does not play a big role in the difference between substitution effects

across populations unless dominance variation is much larger than additive variation,

something that seems unlikely based on theory and estimates. However, the second weight,

due to epistasis, is not bounded and is large for small values of a (e.g. 27 for $Beta(0.04, 0.04)$),

and in this case functional epistasis plays a strong role in additive variation for U-shaped distributions (Hill *et al.* 2008). The spectra of allele frequencies of causal mutations is subject to large debate but there seems to be a consensus that low frequency mutations make non-negligible contributions to genetic variance (Eyre-Walker 2010; Gibson 2012). It is unclear if this leads to extreme values of $\frac{1}{H_b}$.

This result also reinforces the findings of Duenk *et al.* (2020) and Dai *et al.* (2020) that it is mainly biological epistasis that generates changes in additive substitution effects across populations. For instance, when one population drifts, the additive by additive statistical variation enters into additive variation (Hill *et al.* 2006; Walsh and Lynch 2018).

Thus, $r(\alpha_i^b, \alpha_i^{b'})$ will be very low only if (1) populations are distinct, (2) there is *statistical* additive by additive variation and (3) QTL allele frequency distributions are U- or L- shaped.

As for the magnitudes of non-additive variance, in most conceivable cases σ_{AA}^2 and σ_D^2 are of magnitudes at most like additive variances, but more often less than half (Fuerst and Sölkner 1994; Palucci *et al.* 2007; Hill *et al.* 2008; Mäki-Tanila and Hill 2014; Vitezica *et al.* 2018), so in practice the most limiting factor is distance across breeds. Indeed, there are very few accurate estimates of non-additive variation although more are becoming available. We argue that then, and based also in our results, it will be possible to make theory and data-based choices on the possibilities of using predictions and GWAS results across populations.

Then, for reasonable assumptions about allele frequency distribution of QTLs, we have shown that the correlation of substitution effects across populations is typically around 0.8

or higher, which is higher than scarce estimates of genetic correlations across populations available in the literature, which range from 0.3 to 0.8 (Karoui *et al.* 2012; Legarra *et al.* 2014; Porto-Neto *et al.* 2015; Xiang *et al.* 2017) the difference being due, probably, to genotype by environment interaction. Our results seem rather robust to different distributions of allele frequencies. These values are high but not 1, which raises the question of how to conceive optimal strategies for across populations predictions (*e.g.* more data within breed or finer locations of causal variants across breed). This is of practical relevance, *e.g.* for genomic predictions in livestock improvement, but also in human genetics *e.g.* for the use of European-based Polygenic Risk Scores in individuals from other ancestries (Martin *et al.* 2019).

Similar results apply to the same populations across generations, in which case the correlation of substitution effects across generations goes from 0.99 (next 1-2 generations) to 0.80 (10 generations). This illustrates that, even if genomic predictions would use QTLs or markers in very tight LD with QTLs, there would still be, in the long run, a need for a continuous system of data collecting and re-estimation of effects.

In the simulations we confirmed that our estimates are reasonably good although not perfect. They are, depending on the scenario and target (random allele or mutant allele), almost unbiased, slightly biased upwards, or biased downwards. There are several reasons for the disagreement. The most obvious one is the inherent approximation of the Taylor series expansion. Secondly, splitting variances such as $Var(\epsilon_i d_i^{*b})$ or $Var(\epsilon'(\alpha\alpha)_i^b)$ into basic components implies either a strong assumption of multivariate normality and independence or a less strong one of “expectation and variance-independence” (Bohrnstedt

and Goldberger 1969). For instance, it is assumed that the variance of d_i^{*b} is not related to the magnitude of the difference of allele frequencies ϵ_i , but this is not necessarily true. Third, it is further assumed, in the factorization of genetic variances, that QTL effects and allele frequencies are independent. We have also ignored that the change ϵ is proportional to heterozygosity and therefore small at extremes values of allele frequencies. Another factor that we did not consider is the empirical evidence of an inverse relationship between heterozygosity and absolute effect at the locus (Park *et al.* 2011). It is unclear how this would affect our findings. A (rather extreme) functional gene action that generates larger α at extreme frequencies is overdominance, which is similar to our “dominance” scenario.

As for our estimators of the correlation across “negative” alleles $r_{mutant}(\alpha^b, \alpha^{b'})$, they are less robust than the estimators for a “random” allele $r_{random}(\alpha^b, \alpha^{b'})$. The reason for this is that obtaining $r_{mutant}(\alpha^b, \alpha^{b'})$ from $r_{random}(\alpha^b, \alpha^{b'})$ involves a further approximation, the normality of α .

CONCLUSIONS

We presented a coherent, approximate theory, that does not invoke any particular mechanism of gene action, to explain and appraise the change in magnitude of (additive) QTL substitution effects across populations and generations. The theory gives good approximate estimates of this correlation, that needs to be otherwise explicitly estimated.

More importantly, the theory shows that the main sources for the change of effects are relationships across populations, magnitudes of additive and first-order non-additive variances (dominance and additive by additive), and spectra of allele frequencies. These findings provide better understanding of the properties of genomic prediction methods and of quantitative genetics in general.

ACKNOWLEDGMENTS

Part of this work was while Y. Wientjes was visiting the GenPhySE unit at INRAE, Toulouse, financed by the Netherlands Organisation of Scientific Research (NWO). Authors thank INRA SelGen metaprograms EpiSel and EpiFun. This project has received funding from the European Unions' Horizon 2020 Research & Innovation programme under grant agreement N°772787 -SMARTER. We are grateful to the Genotoul Bioinformatics Platform Toulouse Midi- Pyrenees (Bioinfo Genotoul) for providing computing and storage resources. We thank editor and reviewers for very detailed feedback, and MA Toro, A Caballero, S Boitard and M Bonhomme for advice.

REFERENCES

686 Aiello D., K. Patel, and E. Lasagna, 2018 The myostatin gcene: an overview of mechanisms of
687 action and its relevance to livestock animals. *Anim. Genet.* 49: 505–519.
688 <https://doi.org/10.1111/age.12696>

689 Bhatia G., N. Patterson, S. Sankararaman, and A. L. Price, 2013 Estimating and interpreting
690 FST: The impact of rare variants. *Genome Res.* 23: 1514–1521.
691 <https://doi.org/10.1101/gr.154831.113>

692 Bohrnstedt G. W., and A. S. Goldberger, 1969 On the Exact Covariance of Products of
693 Random Variables. *J. Am. Stat. Assoc.* 64: 1439–1442.
694 <https://doi.org/10.1080/01621459.1969.10501069>

695 Bonhomme M., C. Chevalet, B. Servin, S. Boitard, J. Abdallah, *et al.*, 2010 Detecting Selection
696 in Population Trees: The Lewontin and Krakauer Test Extended. *Genetics* 186: 241–
697 262. <https://doi.org/10.1534/genetics.110.117275>

698 Bonifati V., P. Rizzu, M. J. van Baren, O. Schaap, G. J. Breedveld, *et al.*, 2003 Mutations in the
699 DJ-1 Gene Associated with Autosomal Recessive Early-Onset Parkinsonism. *Science*
700 299: 256–259. <https://doi.org/10.1126/science.1077209>

701 Caballero A., and M. A. Toro, 2002 Analysis of genetic diversity for the management of
702 conserved subdivided populations. *Conserv. Genet.* 3: 289.

703 Carlborg Ö., L. Jacobsson, P. Åhlgren, P. Siegel, and L. Andersson, 2006 Epistasis and the
704 release of genetic variation during long-term selection. *Nat. Genet.* 38: 418–420.
705 <https://doi.org/10.1038/ng1761>

706 Chen G. K., P. Marjoram, and J. D. Wall, 2009 Fast and flexible simulation of DNA sequence
707 data. *Genome Res.* 19: 136–142. <https://doi.org/10.1101/gr.083634.108>

708 Cuyabano B. C. D., A. C. Sørensen, and P. Sørensen, 2018 Understanding the potential bias of
709 variance components estimators when using genomic models. *Genet. Sel. Evol.* 50:
710 41. <https://doi.org/10.1186/s12711-018-0411-0>

711 Dai Z., N. Long, and W. Huang, 2020 Influence of Genetic Interactions on Polygenic
712 Prediction. *G3 Genes Genomes Genet.* 10: 109–115.
713 <https://doi.org/10.1534/g3.119.400812>

714 Duenk P., P. Bijma, M. P. L. Calus, Y. C. J. Wientjes, and J. H. J. van der Werf, 2020 The Impact
715 of Non-additive Effects on the Genetic Correlation Between Populations. *G3 Genes*
716 *Genomes Genet.* 10: 783–795. <https://doi.org/10.1534/g3.119.400663>

717 Eyre-Walker A., 2010 Genetic architecture of a complex trait and its implications for fitness
718 and genome-wide association studies. *Proc. Natl. Acad. Sci.* 107: 1752–1756.
719 <https://doi.org/10.1073/pnas.0906182107>

720 Falconer D. S., and T. F. C. Mackay, 1996 *Introduction to quantitative genetics*. Longman New
721 York.

722 Fernández E., J. Sánchez, R. Martínez, A. Legarra, and M. Baselga, 2017 Role of inbreeding
723 depression, non-inbred dominance deviations and random year-season effect in
724 genetic trends for prolificacy in closed rabbit lines. *J. Anim. Breed. Genet.* 134: 441–
725 452.

726 Fuerst C., and J. Sölkner, 1994 Additive and Nonadditive Genetic Variances for Milk Yield,
727 Fertility, and Lifetime Performance Traits of Dairy Cattle. J. Dairy Sci. 77: 1114–1125.
728 [https://doi.org/10.3168/jds.S0022-0302\(94\)77047-8](https://doi.org/10.3168/jds.S0022-0302(94)77047-8)

729 Fuerst C., J. W. James, J. Sölkner, and A. Essl, 1997 Impact of dominance and epistasis on the
730 genetic make-up of simulated populations under selection: a model development. J.
731 Anim. Breed. Genet. 114: 163–175. [https://doi.org/10.1111/j.1439-](https://doi.org/10.1111/j.1439-0388.1997.tb00502.x)
732 [0388.1997.tb00502.x](https://doi.org/10.1111/j.1439-0388.1997.tb00502.x)

733 Galarza C. E., L. A. Matos, D. K. Dey, and V. H. Lachos, 2020 On moments of folded and
734 doubly truncated multivariate extended skew-normal distributions. ArXiv200913488
735 Math Stat.

736 García-Ruiz A., J. B. Cole, P. M. VanRaden, G. R. Wiggans, F. J. Ruiz-López, *et al.*, 2016
737 Changes in genetic selection differentials and generation intervals in US Holstein
738 dairy cattle as a result of genomic selection. Proc. Natl. Acad. Sci. 113: E3995–E4004.

739 Gautier M., A. Capitan, S. Fritz, A. Eggen, D. Boichard, *et al.*, 2007 Characterization of the
740 DGAT1 K232A and Variable Number of Tandem Repeat Polymorphisms in French
741 Dairy Cattle. J. Dairy Sci. 90: 2980–2988. <https://doi.org/10.3168/jds.2006-707>

742 Gibbs R. A., J. F. Taylor, C. P. Van Tassell, W. Barendse, K. A. Eversole, *et al.*, 2009 Genome-
743 wide survey of SNP variation uncovers the genetic structure of cattle breeds. Sci. N.
744 Y. NY 324: 528–532.

745 Gibson G., 2012 Rare and common variants: twenty arguments. Nat. Rev. Genet. 13: 135–
746 145. <https://doi.org/10.1038/nrg3118>

747 Grisart B., W. Coppieters, F. Farnir, L. Karim, C. Ford, *et al.*, 2002 Positional candidate cloning
748 of a QTL in dairy cattle: identification of a missense mutation in the bovine DGAT1
749 gene with major effect on milk yield and composition. *Genome Res* 12: 222–231.

750 Grobet L., L. J. Royo Martin, D. Poncelet, D. Pirottin, B. Brouwers, *et al.*, 1997 A deletion in
751 the bovine myostatin gene causes the double-musled phenotype in cattle. *Nat.*
752 *Genet.* 17: 71–74. <https://doi.org/10.1038/ng0997-71>

753 Haile-Mariam M., and J. E. Pryce, 2015 Variances and correlations of milk production,
754 fertility, longevity, and type traits over time in Australian Holstein cattle. *J. Dairy Sci.*
755 98: 7364–7379. <https://doi.org/10.3168/jds.2015-9537>

756 Hayes B. J., P. J. Bowman, A. C. Chamberlain, K. Verbyla, and M. E. Goddard, 2009 Accuracy
757 of genomic breeding values in multi-breed dairy cattle populations. *Genet Sel Evol*
758 41: 51.

759 Hill W. G., N. H. Barton, and M. Turelli, 2006 Prediction of effects of genetic drift on variance
760 components under a general model of epistasis. *Theor. Popul. Biol.* 70: 56–62.
761 <https://doi.org/10.1016/j.tpb.2005.10.001>

762 Hill W. G., M. E. Goddard, and P. M. Visscher, 2008 Data and Theory Point to Mainly Additive
763 Genetic Variance for Complex Traits. *PLOS Genet.* 4: e1000008.
764 <https://doi.org/10.1371/journal.pgen.1000008>

765 Hudson R. R., M. Slatkin, and W. P. Maddison, 1992 Estimation of levels of gene flow from
766 DNA sequence data. *Genetics* 132: 583–589.

767 Kan R., and C. Robotti, 2017 On Moments of Folded and Truncated Multivariate Normal
768 Distributions. J. Comput. Graph. Stat. 26: 930–934.
769 <https://doi.org/10.1080/10618600.2017.1322092>

770 Karoui S., M. J. Carabaño, C. Díaz, and A. Legarra, 2012 Joint genomic evaluation of French
771 dairy cattle breeds using multiple-trait models. Genet Sel Evol 44: 39.

772 Kojima K., 1959 Role of epistasis and overdominance in stability of equilibria with selection.
773 Proc. Natl. Acad. Sci. U. S. A. 45: 984–989.

774 Kojima K.-I., 1961 Effects of dominance and size of population on response to mass
775 selection*. Genet. Res. 2: 177–188. <https://doi.org/10.1017/S0016672300000689>

776 Lande R., and R. Thompson, 1990 Efficiency of marker-assisted selection in the improvement
777 of quantitative traits. Genetics 124: 743–756.

778 Legarra A., G. Baloche, F. Barillet, J. M. Astruc, C. Soulas, *et al.*, 2014 Within-and across-
779 breed genomic predictions and genomic relationships for Western Pyrenees dairy
780 sheep breeds Latxa, Manech, and Basco-Béarnaise. J. Dairy Sci. 97: 3200–3212.

781 Liu Z., H. Alkhoder, F. Reinhardt, and R. Reents, 2016 Accuracy and bias of genomic
782 prediction for second-generation candidates. Interbull Bull.

783 Lynch M., and B. Walsh, 1998 *Genetics and analysis of quantitative traits*. Sinauer associates.

784 Mackay T. F. C., 2014 Epistasis and Quantitative Traits: Using Model Organisms to Study
785 Gene-Gene Interactions. Nat. Rev. Genet. 15: 22–33.
786 <https://doi.org/10.1038/nrg3627>

787 Mackay T. F. C., 2015 Epistasis for Quantitative Traits in *Drosophila*, pp. 47–70 in *Epistasis:*
788 *Methods and Protocols*, Methods in Molecular Biology. edited by Moore J. H.,
789 Williams S. M. Springer, New York, NY.

790 MacLeod I. M., P. J. Bowman, C. J. Vander Jagt, M. Haile-Mariam, K. E. Kemper, *et al.*, 2016
791 Exploiting biological priors and sequence variants enhances QTL discovery and
792 genomic prediction of complex traits. BMC Genomics 17: 144.
793 <https://doi.org/10.1186/s12864-016-2443-6>

794 Magwire M. M., A. Yamamoto, M. A. Carbone, N. V. Roshina, A. V. Symonenko, *et al.*, 2010
795 Quantitative and Molecular Genetic Analyses of Mutations Increasing *Drosophila* Life
796 Span. PLOS Genet. 6: e1001037. <https://doi.org/10.1371/journal.pgen.1001037>

797 Mäki-Tanila A., and W. G. Hill, 2014 Influence of Gene Interaction on Complex Trait Variation
798 with Multilocus Models. Genetics 198: 355–367.
799 <https://doi.org/10.1534/genetics.114.165282>

800 Marigorta U. M., and A. Navarro, 2013 High Trans-ethnic Replicability of GWAS Results
801 Implies Common Causal Variants. PLOS Genet. 9: e1003566.
802 <https://doi.org/10.1371/journal.pgen.1003566>

803 Martin A. R., M. Kanai, Y. Kamatani, Y. Okada, B. M. Neale, *et al.*, 2019 Clinical use of current
804 polygenic risk scores may exacerbate health disparities. Nat. Genet. 51: 584–591.
805 <https://doi.org/10.1038/s41588-019-0379-x>

806 Meuwissen T. H. E., B. J. Hayes, and M. E. Goddard, 2001 Prediction of total genetic value
807 using genome-wide dense marker maps. Genetics 157: 1819–1829.

808 Moghaddar N., M. Khansefid, J. H. J. van der Werf, S. Bolormaa, N. Duijvesteijn, *et al.*, 2019
809 Genomic prediction based on selected variants from imputed whole-genome
810 sequence data in Australian sheep populations. *Genet. Sel. Evol.* 51: 72.
811 <https://doi.org/10.1186/s12711-019-0514-2>

812 Nei M., 1987 *Molecular Evolutionary Genetics*. Columbia University Press.

813 Palucci V., L. R. Schaeffer, F. Miglior, and V. Osborne, 2007 Non-additive genetic effects for
814 fertility traits in Canadian Holstein cattle (Open Access publication). *Genet. Sel. Evol.*
815 39: 181. <https://doi.org/10.1186/1297-9686-39-2-181>

816 Park J.-H., M. H. Gail, C. R. Weinberg, R. J. Carroll, C. C. Chung, *et al.*, 2011 Distribution of
817 allele frequencies and effect sizes and their interrelationships for common genetic
818 susceptibility variants. *Proc. Natl. Acad. Sci.* 108: 18026–18031.
819 <https://doi.org/10.1073/pnas.1114759108>

820 Pérez-Enciso M., 2014 Genomic relationships computed from either next-generation
821 sequence or array SNP data. *J. Anim. Breed. Genet. Z. Tierzuchtung*
822 *Zuchtungsbiologie* 131: 85–96. <https://doi.org/10.1111/jbg.12074>

823 Pérez-Enciso M., J. C. Rincón, and A. Legarra, 2015 Sequence-vs. chip-assisted genomic
824 selection: accurate biological information is advised. *Genet. Sel. Evol.* 47: 43.

825 Porto-Neto L. R., W. Barendse, J. M. Henshall, S. M. McWilliam, S. A. Lehnert, *et al.*, 2015
826 Genomic correlation: harnessing the benefit of combining two unrelated populations
827 for genomic selection. *Genet. Sel. Evol.* 47: 84. [https://doi.org/10.1186/s12711-015-](https://doi.org/10.1186/s12711-015-0162-0)
828 0162-0

829 Powell J. E., P. M. Visscher, and M. E. Goddard, 2010 Reconciling the analysis of IBD and IBS
830 in complex trait studies. *Nat Rev Genet* 11: 800–805.

831 Rodríguez-Ramilo S., J. M. Elsen, and A. Legarra, 2019 Inbreeding and effective population
832 size in French dairy sheep: Comparison between genomic and pedigree estimates. *J.*
833 *Dairy Sci.* 102: 4227–4237.

834 Roos A. P. W. de, B. J. Hayes, and M. E. Goddard, 2009 Reliability of Genomic Predictions
835 Across Multiple Populations. *Genetics* 183: 1545–1553.
836 <https://doi.org/10.1534/genetics.109.104935>

837 Rupp R., P. Senin, J. Sarry, C. Allain, C. Tasca, *et al.*, 2015 A Point Mutation in Suppressor of
838 Cytokine Signalling 2 (Socs2) Increases the Susceptibility to Inflammation of the
839 Mammary Gland while Associated with Higher Body Weight and Size and Higher Milk
840 Production in a Sheep Model. *PLOS Genet.* 11: e1005629.
841 <https://doi.org/10.1371/journal.pgen.1005629>

842 Shi H., S. Gazal, M. Kanai, E. M. Koch, A. P. Schoech, *et al.*, 2021 Population-specific causal
843 disease effect sizes in functionally important regions impacted by selection. *Nat.*
844 *Commun.* 12: 1098. <https://doi.org/10.1038/s41467-021-21286-1>

845 Streit M., N. Neugebauer, T. H. E. Meuwissen, and J. Bennewitz, 2011 Short communication:
846 evidence for a major gene by polygene interaction for milk production traits in
847 German Holstein dairy cattle. *J. Dairy Sci.* 94: 1597–1600.
848 <https://doi.org/10.3168/jds.2010-3834>

849 Tsuruta S., I. Misztal, T. J. Lawlor, and L. Klei, 2004 Modeling final scores in US Holsteins as a
850 function of year of classification using a random regression model. *Livest. Prod. Sci.*
851 91: 199–207. <https://doi.org/10.1016/j.livprodsci.2003.09.016>

852 VanRaden P., K. Olson, G. Wiggans, J. Cole, and M. Tooker, 2011 Genomic inbreeding and
853 relationships among Holsteins, Jerseys, and Brown Swiss. *J. Dairy Sci.* 94: 5673–5682.

854 Vitezica Z. G., A. Legarra, M. A. Toro, and L. Varona, 2017 Orthogonal Estimates of Variances
855 for Additive, Dominance and Epistatic Effects in Populations. *Genetics*
856 *genetics.116.199406*. <https://doi.org/10.1534/genetics.116.199406>

857 Vitezica Z. G., A. Reverter, W. Herring, and A. Legarra, 2018 Dominance and epistatic genetic
858 variances for litter size in pigs using genomic models. *Genet. Sel. Evol.* 50: 1–8.

859 Walsh B., and M. Lynch, 2018 *Evolution and Selection of Quantitative Traits*. Oxford
860 University Press.

861 Weir B. S., and W. G. Hill, 2002 Estimating F-statistics. *Annu. Rev. Genet.* 36: 721–750.
862 <https://doi.org/10.1146/annurev.genet.36.050802.093940>

863 Welsh C. S., T. S. Stewart, C. Schwab, and H. D. Blackburn, 2010 Pedigree analysis of 5 swine
864 breeds in the United States and the implications for genetic conservation. *J. Anim.*
865 *Sci.* 88: 1610–1618. <https://doi.org/10.2527/jas.2009-2537>

866 Wientjes Y. C. J., P. Bijma, J. Vandenplas, and M. P. L. Calus, 2017 Multi-population Genomic
867 Relationships for Estimating Current Genetic Variances Within and Genetic
868 Correlations Between Populations. *Genetics* 207: 503–515.
869 <https://doi.org/10.1534/genetics.117.300152>

870 Xiang T., O. F. Christensen, and A. Legarra, 2017 Technical note: Genomic evaluation for
 871 crossbred performance in a single-step approach with metafounders. J. Anim. Sci. 95:
 872 1472–1480.

873

874

875

dominance , mutant allele alpha

complementary , mutant allele alpha

multiplicative , mutant allele alpha

dominance , random allele alpha

complementary , random allele alpha

multiplicative , random allele alpha