Inferring species compositions of complex fungal communities from long- and short-read sequence data

Yiheng Hu^{1†}, Laszlo Irinyi^{2,3,4}, Minh Thuy Vi Hoang^{2,3,4}, Tavish Eenjes¹, Abigail Graetz¹, Eric Stone^{1,5}, Wieland Meyer^{2,3,4,6}, Benjamin Schwessinger^{1*}, John P. Rathjen^{1*}

¹ Research School of Biology, Australian National University, Canberra, ACT, Australia

² Molecular Mycology Research Laboratory, Centre for Infectious Diseases and Microbiology, Faculty of Medicine and Health, Sydney Medical School, Westmead Clinical School, The University of Sydney, Sydney, NSW, Australia.

³ Marie Bashir Institute for Infectious Diseases and Biosecurity, The University of Sydney, Sydney, NSW, Australia.

⁴ Westmead Institute for Medical Research, Westmead, NSW Australia.

⁵ ANU-CSIRO Centre for Genomics, Metabolomics and Bioinformatics, Canberra, ACT, Australia

⁶ Westmead Hospital (Research and Education Network), Westmead, NSW, Australia.

* Correspondence: yiheng.hu@anu.edu.au; benjamin.schwessinger@anu.edu.au; john.rathjen@anu.edu.au

† Present address: Department of Microbial Interactions, IMIT/ZMBP, University of Tübingen, Tübingen, Germany

22 **Abstract**

23 Background:

24 The kingdom fungi is crucial for life on earth and is highly diverse. Yet fungi are challenging to
 25 characterize. They can be difficult to culture and may be morphologically indistinct in culture.
 26 They can have complex genomes of over 1 Gb in size and are still underrepresented in whole
 27 genome sequence databases. Overall their description and analysis lags far behind other
 28 microbes such as bacteria. At the same time, classification of species via high throughput
 29 sequencing without prior purification is increasingly becoming the norm for pathogen
 30 detection, microbiome studies, and environmental monitoring. However, standardized
 31 procedures for characterizing unknown fungi from complex sequencing data have not yet
 32 been established.

33 Results:

34 We compared different metagenomics sequencing and analysis strategies for the
 35 identification of fungal species. Using two fungal mock communities of 44 phylogenetically
 36 diverse species, we compared species classification and community composition analysis
 37 pipelines using shotgun metagenomics and amplicon sequencing data generated from both
 38 short and long read sequencing technologies. We show that regardless of the sequencing
 39 methodology used, the highest accuracy of species identification was achieved by sequence
 40 alignment against a fungi-specific database. During the assessment of classification
 41 algorithms, we found that applying cut-offs to the query coverage of each read or contig
 42 significantly improved the classification accuracy and community composition analysis
 43 without significant data loss.

Conclusion:

Overall, our study expands the toolkit for identifying fungi by improving sequence-based fungal classification, and provides a practical guide for the design of metagenomics analyses.

Introduction

Fungi are ubiquitous yet their presence and impact are often overlooked. It has been estimated that 2.2-3.8 million species inhabit planet earth [1] but only about 4% of these are catalogued [2]. Mora *et al.* estimated that there are 7.8 million and 298,0000 animal and plants species on earth with 12.3% and 72.4% of these characterised scientifically, respectively [3], which points towards a more central role in cultural awareness. In contrast, fungi are introduced to our consciousness via a brief mention in high school textbooks, or as largely side subjects in botany and microbiology courses at university [4,5]. Fungi play diverse roles throughout evolution and are particularly active in mediating the breakdown and uptake of nutrients. They constitute a major disease load to humans, causing millions of deaths per year, and wreak devastating crop losses via a constant toll of disease and epidemics and are an existential threat to many frog species [6,7]. On the other hand, fungi are or are used to manufacture delicious foods and beverages, and have saved countless lives via antibiotic production [8,9]. Therefore, a recent call was made to expand fungal research and improve our awareness of this special kingdom [10].

To progress our understanding of fungal biology we need to be able to classify more species more precisely. Fungi have been an independent kingdom since 1969 [11] with addition of further phyla in early 2000 [12–16]. Historically, its taxonomy was based on morphological

and reproductive traits but this has been surpassed by DNA-based classification which revolutionized mycology, not only refining the conventional taxonomic tree [17,18] but also standardizing the identification of new species. In the absence of whole genome data, DNA-based classification primarily exploits the internal transcribed spacer (ITS) within the ribosomal RNA genes as a highly polymorphic marker to distinguish species. It is easily amplified and sequenced due to highly conserved flanking sequences and contains a high degree of variation between even closely related species. Although a mature pipeline comprising ITS amplification, Illumina sequencing and data analysis has been established[19], several studies reported biases from the sequencing technology used and from unevenly amplified fungal marker regions [20–22]. Recently, novel strategies exploiting long-range amplification and long-read sequencing have been developed to improve these classifications [23,24]. In addition, whole genome shotgun sequencing and rapidly expanding genome databases allow mapping of newly generated DNA sequences directly to the database. This strategy allows exploitation of genetic variation throughout the genome and abandonment of the marker gene amplification step, which increases classification accuracy and reduces the biases from the estimation of relative abundance [25].

Although advanced sequencing methods allow novel strategies for fungal identification particularly from mixed samples, new demands are placed on data analysis pipelines to improve the accuracy of fungal classification. Different algorithms have been developed to classify DNA sequences at distinct taxonomic ranks based on sequence databases with taxonomic information [26–30]. For example, alignment algorithms such as Basic Local Alignment Search Tool (BLAST) [27] detect matches of each sequence to subjects of the target database along with the taxonomic information assigned to each entry. Alternatively, sequence features represented by short unique subsequences named k-mers can be derived

from sequence data and mapped to databases to identify taxa with the highest number of cross-mapping k-mers[28]. Several studies have critically assessed algorithms for species classification on simulated datasets or bacterial community datasets [31–33], but comparisons of sequencing strategies for complex fungal communities alignment using real data and different identification pipelines are extremely rare. In addition to search algorithms, the choice of database also influences classifications dramatically, but only a few studies have researched their impact [34–36]. Therefore, more comprehensive benchmarking of both classification algorithms and databases are needed to optimise identification pipelines.

Here, we assessed different combinations of algorithms and databases during processing of both short- and long-read sequencing data for the identification of taxa from complex mock fungal communities. We identified key factors that influence the accuracy of classifications, both for mock community datasets and public datasets. Optimisation of these methods also lead to more accurate community composition analysis. Our results provide guidelines for the design of sequence-based community analysis for fungal species.

Results

Construction of mock fungal community datasets

We constructed two mock communities from the same set of 44 fungal species (Supplementary Table S1). Most of these are human-associated pathogenic yeasts while some are basidiomycete pathogens. One community comprised pooled DNA (PD) from each species and the second was composed of DNA extracted from equal quantities of fungal biomass (PB) of each species that were mixed together prior to extraction. We generated four sequence

datasets for each community using Illumina and nanopore technologies, sequencing both shotgun metagenomes and targeted amplicons respectively. The data derived from each strategy are summarized in Table 1.

Alignment algorithm against a specific fungal database resulted in the most accurate fungal classifications

We compared different analysis strategies for each shotgun dataset. For nanopore datasets, we directly used the quality-controlled reads for classification. For Illumina data, we quality filtered all reads and assembled them into contigs before classification to maximize the classification accuracy. We performed both alignment and k-mer based classifications on these data using BLAST and Kraken2 [27,29] using a ‘winner-takes-all’ strategy in which the top hit was taken as the identity of the query sequence. For each algorithm, we compared the use of two reference databases: the non-redundant NCBI nucleotide database (nt) [37] and the RefSeq fungi database (RFD) [38] which only contains curated fungal genomes. We first assessed the performance of each alignment tool on both databases for each data input. We compared the concordance in the results of each pipeline at the genus level. We define concordance as the percentage of fungal genera identified by both analyses in a pairwise comparison (Figure 1A). The concordance between analyses on each dataset varied between 69% and 86% and generally, Illumina data resulted in higher concordance than did nanopore data.

We then aimed to identify the combination of algorithm and database that yielded the most accurate species identification. We used classified proportion and precision to evaluate each

classification, where $\text{Classified Proportion} = \frac{\# \text{ total basepairs classified.}}{\# \text{ total basepairs of input reads}}$, and

$\text{Precision} = \frac{\# \text{ total basepairs classified correctly}}{\# \text{ total basepairs classified}}$.

The number of total basepairs is calculated as total read length for nanopore reads and total coverage of Illumina reads to each contig [32,33]. We plotted the precision and classified proportion for each pipeline and found three regular patterns (Figures 1B and 1C): First, for each dataset, BLAST resulted in higher precision but lower classification proportion by comparison to Kraken2. Second, Illumina contigs returned higher classification proportion and precision than nanopore reads. Third, classification against the RFD database yielded higher precision than those against the nt database. In summary, BLAST alignments against the RFD database yielded the best classification strategy.

Applying cut-offs to query coverage improves classification accuracy on shotgun metagenomics datasets

We next aimed to improve our classification scheme by filtering the BLAST search results. We reasoned that restricting alignment metrics would reduce the number of false classifications. To investigate changes in classification accuracy after restricting BLAST output parameters, we first BLASTed shotgun metagenomics reads against the RFD database without applying any filter, then applied progressive cut-offs on different parameters of the BLAST results. We evaluated changes in the results based on the metrics precision, remaining rate and completeness. Precision is described above and estimates the accuracy of the classification; remaining rate captures the percentage of the input data remaining after the application of each cut-off; and completeness is the number of taxa captured relative to the total number of taxa within the mock community. We initially applied cut-offs on query length and two

alignment metrics; E-value - the number of expected hits of similar quality that could be found by chance alone; and pident – the percentage of identical matches within the region of alignment between query and subject. As shown in Figure 2A, applying progressive cut-offs to query length did not improve the precision, whilst both completeness and remaining rate diminished dramatically from very small cut-off values. Cut-offs applied to alignment E-values removed <20% of the BLAST results, whereas precision showed minor improvement, especially on nanopore datasets (Figure 2B). For Illumina data, applying cut-offs to the E-value increased the precision by around 2% but at the cost of diminished completeness. E-value cut-offs performed better on nanopore datasets, improving precision by 3% (PD) or 4% (PB) with non-identification of only a single genus from the mock community, at 10^{-250} or almost 10^{-400} respectively. Progressive cut-offs on pident yielded the best results of all three filters. For Illumina data, precision was improved by up to 8% for PB data, and completeness remained at 100% in almost all cases (Figure 2C). For nanopore datasets, pident cut-offs improved the precision by up to ~3% before sharp decreases, with a concurrent filtering of ~60% BLAST result as shown by the remaining rate. Given the characteristically high error rate of nanopore reads, we also applied cut-offs on quality scores to these data. Cut-offs applied to Phred scores did not alter the precision, while a significant proportion of the dataset was lost through filtering (Supplementary Figure S1). Overall, our results suggest that applying each filter to BLAST results performs well on either Illumina or nanopore data but not both, and that cut-offs based on query length or quality scores did not affect the precision significantly.

Given the results above, we investigated how the alignment parameters were calculated and explored other variables to improve the classifications. The BLAST E-value is calculated as $E = mn2^{-S}$ in which S is the bits score derived from the number of gaps and mismatches in the

alignment, and m and n are the query length and database total length respectively [39]. Therefore, the E-value is influenced exponentially by the alignment quality. We next investigated query coverage, a metric based on how much of the query sequence aligned to the subject. We calculated the query coverage as the number of identical matches divided by the read or contig length, and applied progressive cut-offs on this parameter for each dataset/algorithm analysis. As shown in Figure 2D, applying cut-offs on query coverage improved the precision of all four analyses significantly, and did not cause losses of completeness at smaller cut-off values. For example, at a 10% cut-off on query coverage, the precision of all four analyses was 98-99% while the completeness remained at 100% and the removed BLAST results ranged from 10-25%. This result not only supported our hypothesis that the total length of the alignment matters as much as the alignment quality, but also suggested a novel approach to improve the accuracy of fungal classification.

Improving taxa identification from published metagenomics datasets using query coverage as a filtering parameter

After improving classifications by applying cut-offs to the query coverage on the mock community datasets, we extended this strategy to try to improve the classification of published shotgun metagenomics datasets. We re-analysed ten nanopore and six Illumina shotgun metagenomics datasets [40–43]. These included host-associated fungal samples (nanopore) and host-depleted microbiome data (Illumina). Since the environmental datasets contain unknown species, we followed the concept of classification precision. We calculated the percentage of the dataset that was classified into taxa known to be included in the sample. For example, in re-analysing human clinical samples [42], we included the pathogen

(*Pneumocystis*) and the host human (*Homo*) as the true taxa, and calculated the total proportion of query sequences classified to these taxa before and after applying cut-offs on query coverage. Table 2 shows the improvement in taxonomic classification from the published datasets after applying query coverage cut-offs. We initially applied a 20% cut-off on the query coverage for all analyses, but the data loss in most cases was too high. Therefore, we applied query cut-offs that filtered around 20% of the blast result based on our results from the mock fungal community datasets (Figure 2D).

For all Illumina datasets, we downloaded the quality-controlled sequences and re-analysed them using the assembly and BLAST pipeline described above against the NCBI nt database. For the nanopore human datasets [42], we used the BLAST results taken directly from the original articles for analysis. For the infected wheat datasets [41], we downloaded the sequences and re-analysed them against the RefSeq fungal database. The precision increased for nearly all datasets after applying query coverage cut-offs (Table 2). For the Illumina microbiome datasets, we first assessed the change of proportions in fungal taxa after applying cut-offs on query coverages using the species lists identified by Donovan *et al.* [44] as confirmed taxa. We observed only marginal increase in percentages for the confirmed fungal communities, due to their low total proportions in the original samples. We then calculated the improvement in precision for the bacterial communities. The Illumina datasets were generated from swine and mouse gut microbiome samples, so we assessed the change in proportions of their core bacterial genera (a group of bacteria commonly present in swine and mouse guts [45,46]). The percentages of confirmed core bacterial genera improved by up to 5.7% after applying cut-offs on query coverage (Table 2). In addition, in the nanopore human datasets, the total percentage of reads classified as *Homo* in the three healthy individual samples were improved by applying cut-offs to query coverage. These results

indicated that this strategy may be broadly applicable not only to fungal species, but also to the classification of other eukaryotes and bacteria. One Illumina dataset (d1) and one nanopore dataset (a5) showed decreased percentages of confirmed taxa after applying query coverage cut-offs, which might be because the core microbiome species are not representing the species identified in the Illumina sample, or due to the low coverage and high error rate of nanopore data.

Benchmarking classification pipelines for amplicon datasets identified advantages of each strategy

We next assessed different strategies for the classification of ITS amplicon datasets. We amplified the ITS region from both mock communities using two different primer pairs and three technical replicates for each sample. Taking advantage of nanopore technology, we performed long-amplicon sequencing of a roughly 3 kb ribosomal RNA gene region covering part of the 28S subunit, ITS1, 5.8S subunit, ITS2 and part of the 18S subunit [19]. For Illumina sequencing we used the well-established ITS1F-ITS2 amplicon of about 300 bp in length [47]. Similar to the analysis of the shotgun datasets, we applied both k-mer and alignment-based approaches to the classification of nanopore amplicon data. We used the pair-wise alignment algorithm minimap2 as the alignment algorithm instead of BLAST due to its speed and efficiency. We tested four different databases for classification of long amplicons; the NCBI 18S and 28S databases, and two ITS databases from NCBI and UNITE, respectively [38,48]. Overall, we found that the k-mer algorithm returned much higher classification proportion than alignment for each nanopore dataset, but the highest precision (~97%) were achieved by combining the minimap2 alignment algorithm with the NCBI ITS database (Figure 3A). For

Illumina amplicon datasets, we applied the QIIME2 pipeline which is one of the most widely used strategies for ITS classification and community composition analysis[49]. The QIIME2 pipeline groups similar Illumina amplicons into sequence features before classification to reduce the demand on computational resources [50]. Since all individual Illumina reads are grouped into sequence features and all the sequence features are classified, the classification proportion of the Illumina amplicon datasets are 100%. We plotted precision rates from the QIIME2 analysis of both the PD and PB samples with their means (Figure 3B). The mean precision from either Illumina dataset were lower than that from k-mer analysis of the respective nanopore datasets.

Although the precision from the amplicon datasets were higher than that from shotgun datasets, the ITS classification did not identify all genera within the mock community, as shown by our completeness analysis (Figure 3C). The nanopore amplicons identified 68% (PD) and 63% (PB) of the total genera in the mock community, whereas the Illumina amplicon datasets covered only 25% and 41% of the genera respectively. We suspect that the low completeness from ITS classifications was due partially to the low quality of this particular dataset (Table 1) and partially due to non-uniform amplification from the different primer pairs. However, there were fewer nanopore amplicon reads than in the Illumina amplicon datasets and the completeness from the nanopore data was higher (Figure 3C). This supports the argument that long amplicons identify a wider range of species and are more accurate in species classification than short amplicons [51,52].

Cut-offs on query coverage also improve community composition analysis

We next analysed community compositions using the most accurate classification method for each dataset. Community composition refers to the identity and relative abundances of all taxa in a community. Given the observation that use of a restricted database resulted in higher classification precision, we constructed a database containing only the genomes from species within the mock community and aligned all of the data to the mock community database using BLAST. This forces the precision to 100% as any classification will belong to a species from the mock community. We then BLASTed each dataset against this database and calculated the relative abundance of each genus. We defined this as the ‘gold standard’ for community composition analysis of the mock fungal community (Figure 4A). We then compared the community composition determined from each combination of algorithms and databases with the gold standard for each dataset, and measured their differences using three statistical distance tests: Bhattacharyya distance, relative Euclidean distance and relative entropy [53–55]. Consistently, BLASTing sequences against the RFD database produced community compositions with the highest similarity to the gold standard analysis (Figure 4B).

To assess whether query coverage cut-offs also improved the community composition analysis of shotgun metagenomics data, we plotted the changes in statistical distance after progressive application of query coverage cut-offs (Figure 4). After applying cut-offs on the query coverage, the community composition improved in all cases especially for lower cut-off values. The community compositions from PB-Illumina datasets improved and turned out to be the most similar to the gold standard at query-coverage cut-offs greater than 3 - 4%, which is consistent with the changes in precision rate shown in Figure 2D. Overall, our results illustrated that applying cut-offs on query coverage did not only improve the classification accuracy, but also the community composition analysis.

297

298 **Discussion:**

299 Here we investigated the taxonomic classification from sequencing data, one of the key steps
 300 in all metagenomic workflows, with a particular focus on fungi. After assessing various
 301 combinations of algorithms and databases following different sequencing strategies, we
 302 found that the combination of BLAST with the specific RFD database always resulted in the
 303 most precise classification for all mock fungal community datasets. These classifications were
 304 further improved when applying cut-offs on query coverage including positive flow on effects
 305 on downstream community composition analysis from shotgun metagenomics datasets.

306 Despite that sampling and DNA extraction substantially influence the outcome of species
 307 classifications [56–58], choosing an appropriate sequencing strategy is the primary step
 308 towards accurately profiling a sample. For shotgun datasets, our results suggested that both
 309 short and long shotgun datasets have comparable accuracy and both higher than the
 310 amplicon datasets. However, Illumina shotgun datasets require additional steps to assemble
 311 reads into contigs before querying them against a database, and to map all reads back to the
 312 assembly to quantify the coverage. These processes are necessary to achieve accurate
 313 classification from longer contigs [59], but result in a longer sequence-to-result turnover than
 314 the long read shotgun data. In the analysis of the amplicon data, long range amplicons
 315 performs better in the classification accuracy and completeness compare to the short ITS data,
 316 consistent with other studies [51,52]. Comparing to the results from shotgun datasets, the
 317 overall completeness from the result of amplicon datasets is much lower. We think that is
 318 because we used much less amplicon data for benchmarking classification pipelines, and the

incomplete database which do not contain all taxa present in our mock community. Overall, the long read shotgun datasets returned the most accurate fungal classification.

Next, our data supported that alignment algorithm (BLAST) outperform the k-mer based approach (kraken2) in the accuracy of classification [32,60], and also compared progressive cut-offs to major alignment parameters for shotgun metagenomics data. We found that applying read length or read quality cut-offs did not improve the precision of the classification for all shotgun datasets. This observation is different with the previous study based on simulated data, which claimed that the long reads improves the accuracy of classification [60]. Cut-offs on pident slightly improved the classification accuracy for illumina datasets, but the error-prone nature of the nanopore data (~10% error rate) is also reflected in the result, as it causes the breakdown of precision when pident cut-offs reach 90% (Figure 2C).

We found that query coverage cut-off that filter out 20% blast result worked best. Unlike the E-value weighing the gaps and mismatch as the major factor effecting alignment quality, the query coverage weighs the query length as well as the number of identical matches in the assessment of the alignment quality. In this case, we can eliminate more spurious alignments that are due to a small proportion of reads with high fidelity to the reference, which are commonly found in reads containing conserved genes and repeated sequences. Interestingly, to reach the same 20% filtering threshold, we set up higher cut-offs on the query coverage (10 -20%) in mock community datasets than the real environmental datasets, including few extremely low thresholds of query coverage in the Illumina shotgun datasets. We compared other studies that use simulated data to generate metagenomics contigs for classification, and found that they used 90% query coverage cut-offs as the parameter[62–64]. Together with the different result of read length and read quality cut-offs, this observations highlighted

the difference between the use of real environmental data and the simulated data in benchmarking studies, especially for the classification of complex microbial communities.

PD and PB samples showed slightly different results in comparing statistical distance with the gold standard. After applying cut-offs on query coverage, both Bhattacharyya distance and Euclidean distance between the best practice and the gold standard classification only showed marginal decrease in PD samples, and slowly reversed as the cut-offs increase. We think that is because about 1/3 of reads were classified as *Candida* in the pooled DNA sample, so the difference on the relative abundance of one *Candida* genus between the gold standard community composition and the best practice is much higher and much more influential to the final distance than that from other genera.

Following the importance of the alignment quantity represented by the query coverage, the next question is, how to bring the low quality but high quantity alignment into consideration? Therefore, the winner-takes-all selection strategy itself can be re-designed, as the highly conserved genome regions from different species generate highly close alignment scores between the best alignment and other top alignments. In this case, a weighing statistics and the relative probability for multiple top taxonomic assignments can be explored and introduced to replace the best-hit-takes-all strategy. This will be particularly useful in connection with the rapid expansion of the fungal genome databases.

Next to the right classification tool, chosen the appropriate database significantly influences analysis outcomes [33,34]. Based on our observation, we suggest that ‘prior knowledge’ about the dataset should guide the choice of the appropriate database as this will improve the accuracy of taxonomic classifications. For example, our results suggested that the restricted database resulted in more accurate fungal classifications for shotgun

metagenomics datasets. This strategy might be appropriate if queries are initial binned into kingdoms before a more in-depth analysis with kingdom specific databases. Also, Kaehler *et al.* [65] incorporated environment-specific taxonomic abundance information into the analysis of amplicon datasets and showed that these improve classification accuracy. Similar approaches can be applied to metagenomic datasets. In addition, machine learning strategies become increasingly popular for analysing genomic data. Here taxonomic classifiers could be trained on existing labelled sequence datasets before being applied to communities with similar composition to the training datasets or to identify target species from complex communities [60,66].

Conclusion

In this study, we perform an in-depth analysis on how different sequencing strategies, classification algorithms and databases impact fungal classifications using complex real-life mock community sequencing datasets. We find that alignment algorithm (BLAST) with targeted fungal database (RFD) achieve the best classification accuracy and community composition estimates. These can be further improved by applying cut-offs on query coverage. Taken together, the findings from our benchmarking workflows have important implications for mycology studies for multiple stages of metagenomics analysis, and provided a guide to other researchers aiming to study fungal metagenomics.

Methods

Code availability

All detailed commands and scripts used in each step were summarized in <https://github.com/Yiheng323/Benchmarking-taxonomic-classification-strategies-using-mock-fungal-communities>.

Fungal harvesting, DNA extraction and construction of mock communities

Selected fungal strains were cultured onto Sabouraud dextrose agar and incubated for 48 hours at 27°C.

For the species in the PD community, an inoculating loop full of fungal cells were scraped into a 1.5 mL microfuge tube and crushed with a pestle and liquid nitrogen. Genomic DNA was then extracted using the Zymo Research *Quick-DNA* Fungal/Bacterial Miniprep Kit (cat. no. D6005 Zymo Research, Irvine, CA, USA). First, BashingBead™ Buffer was added to the crushed fungal cells and vortexed. The mixture was then filtered through a Zymo-Spin™ III-F Column and the filtrate was combined with Genomic Lysis Buffer. The mixture was filtered through a Zymo-Spin™ IICR Column and washed with DNA Pre-Wash buffer and g-DNA Wash Buffer. The DNA was eluted in nuclease free water. DNA concentration was measured using the DeNovix dsDNA Broad Range Kit (DeNovix, Wilmington, DE, USA) and 250 ng of DNA from each strain were then pooled together.

For the PB community, two inoculating loops of fungi of each species in teg mock community were scraped into a ceramic mortar. Liquid nitrogen was then poured into the mortar and the fungal mixture was crushed into a fine powder. DNA was then extracted using the Qiagen DNeasy PowerMax Soil Kit (cat. no. 12988-10 Qiagen, Hilden, Germany). PowerBead Solution and Solution C1 were added to the crushed fungal community, vortexed and centrifuged. The supernatant was then added to Solution C2, mixed and centrifuged, which was then repeated with Solution C3. The resulting supernatant was combined with Solution C4 and centrifuged

through a column. The column was then washed twice with Solution C5. Final DNA was eluted in nuclease free water and the concentration measured using the DeNovix dsDNA Broad Range Kit.

Library preparation and sequencing

The ITS1 regions of the rRNA gene were amplified with the universal fungal primers, ITS1F (CTTGGTCATTTAGAGGAAGTAA) and ITS2 (GCTGCGTTCTTCATCGATGC)[47]. Sequencing of PCR amplicons was conducted with MiSeq® System of Illumina (Illumina, San Diego, CA, USA) by the Australian Genome Research Facility. The Illumina bcl2fastq 2.18.0.12 pipeline was used to generate the sequence data. Pair-ends reads 2 × 300bp were generated up to 0.15 GB per sample for amplicon data. The Illumina amplicon data are then directly imported into QIIME2 for analysis. For shotgun Illumina datasets, we employed the same sequencing pipeline as the amplicon data, with MiSeq® and bcl2fastq 2.18.0.12 pipeline for the 2 x 300bp paired end reads. Raw shotgun Illumina reads were trimmed adapters with Trimomatic [67]. Quality controlled, paired end reads were merged and assembled to metagenomics contigs using IDBA_UD [68], which is more suitable for datasets with uneven sequencing depths of each species. After assembly, raw reads were mapped back to the contigs using bwa-mem [69], and the bam files were generated and sorted from sam files using samtools [70]. Bedtools [71] was used for generating coverage for each contig, and we used python numpy and pandas module to calculate the average coverage for each contig.

For Nanopore sequencing of both shotgun and amplicon sequencing, we used Ligation Sequencing 1D SQK-LSK108 and Native Barcoding Expansion (PCR-free) EXP-NBD103 Kits from ONT (UK), as adapted by Hu and Schwessinger [72], which was adapted from the manufacturer's instructions with the omission of DNA fragmentation and DNA repair. DNA

433 was first cleaned up using a 1× volume of Agencourt AMPure XP beads (cat. no. A63881,
434 Beckman Coulter, Indianapolis, IN, USA) following manufacturer's instructions. We then
435 eluted the beads binded DNA in 51 µl nuclease free water and quantified using NanoDrop®
436 and Quibit™ Fluorometer (Thermo Fisher Scientific, Waltham, MA, USA). DNA was end-
437 repaired (NEBNext Ultra II End-Repair/dA-tailing Module, cat. No. E7546), 1x volume beads
438 cleaned (AMPure XP beads) and eluted in 31 µl nuclease free water. Barcoding reaction was
439 performed by adding 2 µl of each native barcode and 20 µl NEB Blunt/TA Master Mix (cat. No.
440 M0367) into 18 µl DNA, mixing gently and incubating at room temperature for 10 minutes. A
441 1× volume (40 µl) Agencourt AMPure XP clean-up was then performed and the DNA was
442 eluted in 15 µl nuclease free water. Ligation was then performed by adding 20 µl Barcode
443 Adapter Mix (EXP-NBD103 Native Barcoding Expansion Kit, ONT, UK), 20 µl NEBNext Quick
444 Ligation Reaction Buffer, and Quick T4 DNA Ligase (cat. No. E6056) to the 50 µl pooled
445 equimolar barcoded DNA, mixing gently and incubating at room temperature for 10 minutes.
446 The adapter-ligated DNA was cleaned-up by adding a 0.4× volume (40 µl) of Agencourt
447 AMPure XP beads, incubating for 5 minutes at room temperature and resuspending the pellet
448 twice in 140 µl ABB provided in the SQK-LSK108 kit. The purified-ligated DNA was
449 resuspended by adding 15 µl ELB provided in the SQK-LSK108 kit and resuspending the beads.
450 The beads were pelleted again and the supernatant sequencing library was transferred to a
451 new 0.5 ml DNA LoBind tube (Eppendorf, Germany). Nanopore sequencing was carried out
452 by MinION MK1b device using R9.4.1 Flowcells. Raw fast5 files are barcode demultiplexed by
453 deepbiner (ONT), then basecalled by Guppy (v3.6.0, ONT, UK). Quality passed reads in fastq
454 files were trimmed adapters and barcodes using qcat (ONT, UK). For the long amplicon data,
455 we filtered out reads less than 2000 base pairs. All sequencing data was submitted to NCBI
456 Short Read Archive (SRA) under the bioproject PRJNA725368 including eight accessions:

SRX10705648, SRX10705649, SRX10705650, SRX10705651, SRX10705695, SRX10705696 and SRX10705697.

Genome assembly

While generating the reference genome database, we found that there were no reference genomes for *Candida rugosa*, *Candida mesorugosa* and *Cryptococcus magnus*, so we performed nanopore sequencing on pure DNA from each species and assembled their draft genomes. These assemblies were of sufficient contiguity and quality (Supplementary Table S2), so we added the new draft genomes into the reference database.

The nanopore data of *Candida rugosa*, *Candida mesorugosa* and *Cryptococcus magnus* was generated individually using Ligation Sequencing 1D SQK-LSK108 kit alone, and from independent flowcells. Data from each flowcell was basecalled and quality filtered using the same pipeline as described above. We got roughly 40X coverage for *Candida rugosa* and *Candida mesorugosa*, and 20X coverage for *Cryptococcus magnus*. Draft genomes were assembled with Flye [73] using default parameters and an estimated genome size of 20Mb. After assembly, the contigs were polished ten times with Racon [74] using nanopore reads, followed by one polishing with Medaka (ONT). Polished assembly was assessed completeness using BUSCO [75]. The assembly statistics were reported from Flye.

Database constructions

For shotgun metagenomics analysis, we used three BLAST database and three kraken databases. Two databases (nt and RFD) are from the same NCBI source, downloaded in May 2019. BLAST and kraken2 nt databases were downloaded using the updateblastdb.pl script from BLAST+ package[76] and the kraken2 program [29], respectively. The fasta files of RefSeq

479 fungal database was downloaded from the NCBI and converted to BLAST database using
480 makblastdb command from the BLAST+ package[76], and was added to the kraken2 database
481 library using kraken2 command [29]. We also build the standard kraken2 database for
482 masking the contaminated regions within the fungal genomes using kraken2 command [29].
483 To generate the mock community database with only the species from the mock community,
484 we downloaded the genomes of all species in the mock community from the NCBI according
485 to their accessions (Supplementary Table S1), and concatenated them with the three newly
486 assembled genomes of *Candida rugosa*, *Candida mesorugosa* and *Cryptococcus magnus*.
487 Following the previous pipeline [77], we then performed a kraken2 search to identify the
488 potential contaminated regions in the concatenated fasta, and masked those regions using
489 bedtools [71]. We also masked the low complexity regions using the dustmasker from BLAST+
490 package [76]. To enable new genomes to be indexed by blastn, we updated the taxonomic
491 map file by adding the fasta headers of the three new genomes and manually assigned their
492 taxonomic ID in the file. Lastly, we used the makeblastdb program to construct the mock
493 community database.

494 For amplicon data analysis, we used two versions of fungal ITS database from the NCBI and
495 UNITE, plus the fungal 18S, 28S database from the NCBI. All of them are downloaded as fasta
496 format in February 2020 and added to the kraken2 database library using kraken2 command
497 [29].

498 **Data analysis**

499 For Shotgun metagenomics datasets, we first used blastn (version 2.10.1) and kraken2
500 (version 2.0.8) to assign the NCBI taxonomic ID for each Illumina contig or Nanopore read.
501 During the classification, we found one contamination species *Purpureocillium lilacinum*

always present in all samples with a significant abundance (10-20%). Therefore, we added this species into the true species list. The best hit from BLAST or species with the highest k-mer counts for each read and/or contig was retained for further analysis. After classification, we used python pandas module to merge information from different output files, and used ete3 module [78] to assign taxonomic information to each read or contigs. The relative abundance of each classification were calculated based on the total length of Nanopore reads of total coverage of Illumina contigs. We used python numpy and math module for all statistical analysis.

For amplicon datasets, we sequenced each sample with three technical replicates. The classification workflow was different for datasets with different sequencing technologies. We only used QIME2 workflow plus the UNITE database for the Illumina amplicon data, since it is the only widely used method for classification. The paired end reads were denoised using the DADA2[79] plugin and assigned taxonomic information using the q2-feature-classifier [80] plugin. The QIME2 classifier was trained by the database sequence before classification. The classification output .qzv files were visualized by the QIME2 view website (<https://view.qiime2.org/>) and the feature-frequency csv file was extracted from the website. We then used python numpy and math module for the mathematical analysis and used seaborn module for generating figures.

For nanopore amplicon datasets, we used kraken2 as the k-mer based algorithm and minimap2 as the alignment based algorithm. The kraken2 command is the same as the kraken2 analysis for the shotgun metagenomics datasets, only using different databases. For the minimap2 analysis, we extracted the accessions of the best hits from the output files, and searched their corresponding taxonomic ID from the NCBI taxonomic map (downloaded from

https://ftp.ncbi.nih.gov/pub/taxonomy/accession2taxid/nucl_wgs.accession2taxid.gz, in

June 2020) using python pandas module. We then merge information from different output files, and used ete3 module again to assign taxonomic information to each read.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Data Availability

All sequencing data was submitted to NCBI Short Read Archive (SRA) under the BioProject PRJNA725368 including eight accessions: SRX10705648, SRX10705649, SRX10705650, SRX10705651, SRX10705695, SRX10705696 and SRX10705697.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

WM, ES, BS and JPR conceived the study and designed experiments. YH, LI and WTVH prepared the samples and generated sequencing data. YH, TE and AG performed the bioinformatics analysis. ES provided feedback on statistical analysis. All authors contributed to data analysis and manuscript writing. All authors read and approved the final manuscript.

Acknowledgements

We thank Dr. Yu Lin for the insightful discussion and valuable suggestions to our manuscript. This work was supported by computational resources provided by the Australian Government through the National Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme.

Funding

This study was supported by a National Health and Medical Research Council of Australia (NH&MRC) grant [#GNT1121936] to W.M., B.S. is supported by an Australian Research Council Future Fellowship FT180100024, and Y.H., E.S., J.R., and B.S. are supported by The Hermon Slade Foundation grant HSF_17_04.

Figure Legends

Table 1. The characteristics for each dataset.

Table 2. Assignment of published sequence data to genera after application of cut-offs to query coverage.

Figure 1. Analysis of shotgun metagenomics data. (A) Swarmplot showing the concordance in genus identification after varying either the alignment algorithm or querying different databases on different data inputs. nt = NCBI nucleotide database; RFD = RefSeq Fungi database; data inputs are indicated below the line (PD = pooled data; PB = pooled biomass); (B) Identification of fungal genera from PD samples. The classified proportion and precision were derived from different combinations of search algorithms and databases as indicated (box); (C) Identification of fungal genera from PB samples. The classification proportion and

precision were derived from the different combinations of search algorithms and databases as indicated.

Figure 2. Dynamics in precision, completeness and remaining rate after applying progressive cut-offs on BLAST alignment metrics. (A) Cut-offs applied to query length. (B) Cut-offs applied to alignment E-values. (C) Cut-offs applied to the percentage of identical matches. (D) Cut-offs applied to query coverage.

Figure 3. Benchmarking of amplicon datasets. (A) Scatter plot represented genus level classification proportion and precision for nanopore amplicon data. (B) Genus level precision of Illumina amplicon data. Classification proportion of Illumina data were 100% due to the nature of the QIIME2 pipeline (based on the UNITE ITS database). (C) Genus level completeness of both nanopore and Illumina amplicon datasets. The nanopore results are from minimap2 algorithm and uniteITS database.

Figure 4. Improving community composition analysis by applying query coverage cut-offs. (A) Experimental flowchart for analysing community compositions. (B) Statistical similarity measures between gold standard community composition and each combination of algorithms and databases. Lower values correspond to greater similarity between the samples and the gold standard. (C) Change in Bhattacharyya distance after applying cut-offs to query coverage for each dataset as indicated. The query coverage gap between each dot point is 0.5%. (D) Change in relative Euclidean distance after applying cut-offs to query coverage for each dataset. The gap between each dot point is 0.5%. (E) Change in relative entropy after applying cut-offs on query coverage for each dataset. The gap between each dot point is 0.5%.

Supplementary Table S1. Metadata of the mock fungal community

Supplementary Table S2. Assembly statistics of the draft genomes of *Candida rugosa*,
Candida mesorugosa and *Cryptococcus magnus* in the mock fungal community.

Supplementary Figure S1. Change of alignment metrics after applying cut-offs on Phred score.

Reference

1. Hawksworth DL, Lücking R. Fungal Diversity Revisited: 2.2 to 3.8 Million Species. *Fungal Kingd.* 2017;79–95.
2. Cheek M, Lughadha EN, Kirk P, Lindon H, Carretero J, Looney B, et al. New scientific discoveries: Plants and fungi. *PLANTS PEOPLE PLANET.* 2020;2:371–88.
3. Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. How Many Species Are There on Earth and in the Ocean? *PLOS Biol. Public Library of Science*; 2011;9:e1001127.
4. Freimoser F. Start teaching mycology! [Internet]. *Nat. Res. Microbiol. Community.* 2017 [cited 2021 Jan 6]. Available from: <https://naturemicrobiologycommunity.nature.com/posts/20287-start-teaching-mycology>
5. Editorial. Stop neglecting fungi. *Nat Microbiol.* Nature Publishing Group; 2017;2:1–2.
6. Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, Stukenbrock EH, et al. Threats Posed by the Fungal Kingdom to Humans, Wildlife, and Agriculture. *mBio* [Internet]. American Society for Microbiology; 2020 [cited 2021 Jan 6];11. Available from: <https://mbio.asm.org/content/11/3/e00449-20>
7. Scheele BC, Pasmans F, Skerratt LF, Berger L, Martel A, Beukema W, et al. Amphibian fungal panzootic causes catastrophic and ongoing loss of biodiversity. *Science. American Association for the Advancement of Science*; 2019;363:1459–63.
8. Naranjo-Ortiz MA, Gabaldón T. Fungal evolution: diversity, taxonomy and phylogeny of the Fungi. *Biol Rev.* 2019;94:2101–37.
9. Valverde ME, Hernández-Pérez T, Paredes-López O. Edible Mushrooms: Improving Human Health and Promoting Quality Life [Internet]. *Int. J. Microbiol. Hindawi*; 2015 [cited 2020 Oct 19]. p. e376387. Available from: <https://www.hindawi.com/journals/ijmicro/2015/376387/>
10. Kong HH, Segre JA. Cultivating fungal research. *Science. American Association for the Advancement of Science*; 2020;368:365–6.
11. Whittaker RH. New Concepts of Kingdoms of Organisms. *Science. American Association for the Advancement of Science*; 1969;163:150–60.

- 621 12. James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, Cox CJ, et al. Reconstructing the
622 early evolution of Fungi using a six-gene phylogeny. *Nature*. Nature Publishing Group;
623 2006;443:818–22.
- 624 13. White MM, James TY, O'Donnell K, Cafaro MJ, Tanabe Y, Sugiyama J. Phylogeny of the
625 Zygomycota based on nuclear ribosomal sequence data. *Mycologia*. Taylor & Francis;
626 2006;98:872–84.
- 627 14. James TY, Letcher PM, Longcore JE, Mozley-Standridge SE, Porter D, Powell MJ, et al. A
628 molecular phylogeny of the flagellated fungi (Chytridiomycota) and description of a new
629 phylum (Blastocladiomycota). *Mycologia*. Taylor & Francis; 2006;98:860–71.
- 630 15. Fischer WM, Palmer JD. Evidence from small-subunit ribosomal RNA sequences for a
631 fungal origin of Microsporidia. *Mol Phylogenet Evol*. 2005;36:606–22.
- 632 16. Keeling PJ, Luker MA, Palmer JD. Evidence from Beta-Tubulin Phylogeny that
633 Microsporidia Evolved from Within the Fungi. *Mol Biol Evol*. 2000;17:23–31.
- 634 17. Jones MDM, Forn I, Gadelha C, Egan MJ, Bass D, Massana R, et al. Discovery of novel
635 intermediate forms redefines the fungal tree of life. *Nature*. Nature Publishing Group;
636 2011;474:200–3.
- 637 18. Adl SM, Simpson AGB, Lane CE, Lukeš J, Bass D, Bowser SS, et al. The Revised Classification
638 of Eukaryotes. *J Eukaryot Microbiol*. 2012;59:429–514.
- 639 19. Raja HA, Miller AN, Pearce CJ, Oberlies NH. Fungal Identification Using Molecular Tools: A
640 Primer for the Natural Products Research Community. *J Nat Prod*. 2017;80:756–70.
- 641 20. Schirmer M, Ijaz UZ, D'Amore R, Hall N, Sloan WT, Quince C. Insight into biases and
642 sequencing errors for amplicon sequencing with the Illumina MiSeq platform. *Nucleic Acids*
643 *Res*. 2015;43:e37.
- 644 21. Filippis FD, Laiola M, Blaiotta G, Ercolini D. Different Amplicon Targets for Sequencing-
645 Based Studies of Fungal Diversity. *Appl Environ Microbiol* [Internet]. American Society for
646 Microbiology; 2017 [cited 2021 Feb 15];83. Available from:
647 <https://aem.asm.org/content/83/17/e00905-17>
- 648 22. Frau A, Kenny JG, Lenzi L, Campbell BJ, Ijaz UZ, Duckworth CA, et al. DNA extraction and
649 amplicon production strategies deeply influence the outcome of gut mycobiome studies. *Sci*
650 *Rep*. Nature Publishing Group; 2019;9:9328.
- 651 23. Heeger F, Bourne EC, Baschien C, Yurkov A, Bunk B, Spröer C, et al. Long-read DNA
652 metabarcoding of ribosomal RNA in the analysis of fungi from aquatic environments. *Mol Ecol*
653 *Resour*. 2018;18:1500–14.
- 654 24. D'Andreano S, Cuscó A, Francino O. Rapid and real-time identification of fungi up to the
655 species level with long amplicon Nanopore sequencing from clinical samples. *bioRxiv*. Cold
656 Spring Harbor Laboratory; 2020;2020.02.06.936708.

657 25. Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. Shotgun metagenomics, from
658 sampling to analysis. *Nat Biotechnol.* 2017;35:833–44.

659 26. Marcelino VR, Clausen PTLC, Buchmann JP, Wille M, Iredell JR, Meyer W, et al. CCMetagen:
660 comprehensive and accurate identification of eukaryotes and prokaryotes in metagenomic
661 data. *Genome Biol.* 2020;21:103.

662 27. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J*
663 *Mol Biol.* 1990;215:403–10.

664 28. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using exact
665 alignments. *Genome Biol.* 2014;15:R46.

666 29. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. *Genome Biol.*
667 2019;20:257.

668 30. Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S, et al. Integrating
669 taxonomic, functional, and strain-level profiling of diverse microbial communities with
670 bioBakery 3. *bioRxiv.* Cold Spring Harbor Laboratory; 2020;2020.11.19.388223.

671 31. Zielezinski A, Girgis HZ, Bernard G, Leimeister C-A, Tang K, Dencker T, et al. Benchmarking
672 of alignment-free sequence comparison methods. *Genome Biol.* 2019;20:144.

673 32. McIntyre ABR, Ounit R, Afshinnikoo E, Prill RJ, Hénaff E, Alexander N, et al. Comprehensive
674 benchmarking and ensemble approaches for metagenomic classifiers. *Genome Biol.*
675 2017;18:182.

676 33. Ye SH, Siddle KJ, Park DJ, Sabeti PC. Benchmarking Metagenomics Tools for Taxonomic
677 Classification. *Cell.* 2019;178:779–94.

678 34. Nasko DJ, Koren S, Phillippy AM, Treangen TJ. RefSeq database growth influences the
679 accuracy of k-mer-based lowest common ancestor species identification. *Genome Biol.*
680 2018;19:165.

681 35. R. Marcelino V, Holmes EC, Sorrell TC. The use of taxon-specific reference databases
682 compromises metagenomic classification. *BMC Genomics.* 2020;21:184.

683 36. Heeger F, Wurzbacher C, Bourne EC, Mazzoni CJ, Monaghan MT. Combining the 5.8S and
684 ITS2 to improve classification of fungi. *Methods Ecol Evol.* 2019;10:1702–11.

685 37. NCBI Resource Coordinators. Database resources of the National Center for Biotechnology
686 Information. *Nucleic Acids Res.* 2018;46:D8–13.

687 38. O’Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference
688 sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional
689 annotation. *Nucleic Acids Res.* 2016;44:D733–45.

690 39. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and
691 PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*
692 1997;25:3389–402.

693 40. Xiao L, Feng Q, Liang S, Sonne SB, Xia Z, Qiu X, et al. A catalog of the mouse gut
694 metagenome. *Nat Biotechnol.* Nature Publishing Group; 2015;33:1103–8.

695 41. Hu Y, Green GS, Milgate AW, Stone EA, Rathjen JP, Schwessinger B. Pathogen Detection
696 and Microbiome Analysis of Infected Wheat Using a Portable DNA Sequencer. *Phytobiomes J.*
697 Scientific Societies; 2019;3:92–101.

698 42. Irinyi L, Hu Y, Hoang MTV, Pasic L, Halliday C, Jayawardena M, et al. Long-read sequencing
699 based clinical metagenomics for the detection and confirmation of *Pneumocystis jirovecii*
700 directly from clinical specimens: A paradigm shift in mycological diagnostics. *Med Mycol.*
701 Oxford Academic; 2020;58:650–60.

702 43. Xiao L, Estellé J, Kiillerich P, Ramayo-Caldas Y, Xia Z, Feng Q, et al. A reference gene
703 catalogue of the pig gut microbiome. *Nat Microbiol.* Nature Publishing Group; 2016;1:1–6.

704 44. Donovan PD, Gonzalez G, Higgins DG, Butler G, Ito K. Identification of fungi in shotgun
705 metagenomics datasets. *PLOS ONE.* Public Library of Science; 2018;13:e0192898.

706 45. Holman DB, Brunelle BW, Trachsel J, Allen HK. Meta-analysis To Define a Core Microbiota
707 in the Swine Gut. *mSystems.* 2017;2.

708 46. Wang J, Lang T, Shen J, Dai J, Tian L, Wang X. Core Gut Bacteria Analysis of Healthy Mice.
709 *Front Microbiol* [Internet]. Frontiers; 2019 [cited 2021 Jan 6];10. Available from:
710 <https://www.frontiersin.org/articles/10.3389/fmicb.2019.00887/full>

711 47. White TJ, Bruns T, Lee S, Taylor J. AMPLIFICATION AND DIRECT SEQUENCING OF FUNGAL
712 RIBOSOMAL RNA GENES FOR PHYLOGENETICS. *PCR Protoc* [Internet]. Elsevier; 1990 [cited
713 2021 Mar 19]. p. 315–22. Available from:
714 <https://linkinghub.elsevier.com/retrieve/pii/B9780123721808500421>

715 48. Nilsson RH, Larsson K-H, Taylor AFS, Bengtsson-Palme J, Jeppesen TS, Schigel D, et al. The
716 UNITE database for molecular identification of fungi: handling dark taxa and parallel
717 taxonomic classifications. *Nucleic Acids Res.* Oxford Academic; 2019;47:D259–64.

718 49. Bharti R, Grimm DG. Current challenges and best-practice protocols for microbiome
719 analysis. *Brief Bioinform* [Internet]. 2019 [cited 2021 Jan 7]; Available from:
720 <https://doi.org/10.1093/bib/bbz155>

721 50. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible,
722 interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol.*
723 Nature Publishing Group; 2019;37:852–7.

724 51. Johnson JS, Spakowicz DJ, Hong B-Y, Petersen LM, Demkowicz P, Chen L, et al. Evaluation
725 of 16S rRNA gene sequencing for species and strain-level microbiome analysis. *Nat Commun.*
726 Nature Publishing Group; 2019;10:5029.

727 52. Krehenwinkel H, Pomerantz A, Henderson JB, Kennedy SR, Lim JY, Swamy V, et al.
728 Nanopore sequencing of long ribosomal DNA amplicons enables portable and simple
729 biodiversity assessments with high phylogenetic resolution across broad taxonomic scale.

730 GigaScience [Internet]. 2019 [cited 2021 Jan 7];8. Available from:
731 <https://doi.org/10.1093/gigascience/giz006>

732 53. Dokmanic I, Parhizkar R, Ranieri J, Vetterli M. Euclidean Distance Matrices: Essential
733 Theory, Algorithms and Applications. 2015 [cited 2021 Jan 7]; Available from:
734 <https://arxiv.org/abs/1502.07541v2>

735 54. Aherne FJ, Thacker NA, Rockett PI. The Bhattacharyya metric as an absolute similarity
736 measure for frequency coded data. *Kybernetika*. 1998;34:363–8.

737 55. MacKay DJC, Kay DJCM. Information Theory, Inference and Learning Algorithms.
738 Cambridge University Press; 2003.

739 56. Henderson G, Cox F, Kittelmann S, Miri VH, Zethof M, Noel SJ, et al. Effect of DNA
740 Extraction Methods and Sampling Techniques on the Apparent Structure of Cow and Sheep
741 Rumen Microbial Communities. *PLOS ONE*. Public Library of Science; 2013;8:e74787.

742 57. Davis A, Kohler C, Alsallaq R, Hayden R, Maron G, Margolis E. Improved yield and accuracy
743 for DNA extraction in microbiome studies with variation in microbial biomass. *BioTechniques*.
744 Future Science; 2019;66:285–9.

745 58. Douglas CA, Ivey KL, Papanicolas LE, Best KP, Muhlhausler BS, Rogers GB. DNA extraction
746 approaches substantially influence the assessment of the human breast milk microbiome. *Sci*
747 *Rep*. Nature Publishing Group; 2020;10:123.

748 59. Tamames J, Cobo-Simón M, Puente-Sánchez F. Assessing the performance of different
749 approaches for functional and taxonomic annotation of metagenomes. *BMC Genomics*.
750 2019;20:960.

751 60. Liang Q, Bible PW, Liu Y, Zou B, Wei L. DeepMicrobes: taxonomic classification for
752 metagenomics with deep learning. *NAR Genomics Bioinforma* [Internet]. 2020 [cited 2021 Jan
753 6];2. Available from: <https://doi.org/10.1093/nargab/lqaa009>

754 61. Pearman WS, Freed NE, Silander OK. Testing the advantages and disadvantages of short-
755 and long- read eukaryotic metagenomics using simulated reads. *BMC Bioinformatics*.
756 2020;21:220.

757 62. Mallawaarachchi V, Wickramarachchi A, Lin Y. GraphBin: refined binning of metagenomic
758 contigs using assembly graphs. *Bioinformatics*. 2020;36:3307–13.

759 63. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, et al. Binning
760 metagenomic contigs by coverage and composition. *Nat Methods*. Nature Publishing Group;
761 2014;11:1144–6.

762 64. Wickramarachchi A, Mallawaarachchi V, Rajan V, Lin Y. MetaBCC-LR: metagenomics
763 binning by coverage and composition for long reads. *Bioinformatics*. 2020;36:i3–11.

764 65. Kaehler BD, Bokulich NA, McDonald D, Knight R, Caporaso JG, Huttenley GA. Species
765 abundance information improves sequence taxonomy classification accuracy. *Nat Commun.*
766 Nature Publishing Group; 2019;10:4643.

767 66. Bokulich NA, Dillon MR, Bolyen E, Kaehler BD, Huttenley GA, Caporaso JG. q2-sample-
768 classifier: machine-learning tools for microbiome classification and regression. *J Open Res*
769 *Softw* [Internet]. 2018 [cited 2021 Jan 6];3. Available from:
770 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6759219/>

771 67. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
772 data. *Bioinformatics*. 2014;30:2114–20.

773 68. Peng Y, Leung HCM, Yiu SM, Chin FYL. IDBA-UD: a de novo assembler for single-cell and
774 metagenomic sequencing data with highly uneven depth. *Bioinformatics*. Oxford Academic;
775 2012;28:1420–8.

776 69. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.
777 *Bioinformatics*. 2009;25:1754–60.

778 70. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
779 Alignment/Map format and SAMtools. *Bioinformatics*. 2009;25:2078–9.

780 71. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
781 *Bioinformatics*. Oxford Academic; 2010;26:841–2.

782 72. Hu Y, Schwessinger B. Amplicon sequencing using MinION optimized from 1D native
783 barcoding genomic DNA [Internet]. *protocols.io*. 2018 [cited 2018 Sep 27]. Available from:
784 [https://www.protocols.io/view/amplicon-sequencing-using-minion-optimized-from-1d-](https://www.protocols.io/view/amplicon-sequencing-using-minion-optimized-from-1d-mhkc34w)
785 [mhkc34w](https://www.protocols.io/view/amplicon-sequencing-using-minion-optimized-from-1d-mhkc34w)

786 73. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat
787 graphs. *Nat Biotechnol*. Nature Publishing Group; 2019;37:540–6.

788 74. Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly from
789 long uncorrected reads. *Genome Res*. 2017;27:737–46.

790 75. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing
791 genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*.
792 Oxford Academic; 2015;31:3210–2.

793 76. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
794 architecture and applications. *BMC Bioinformatics*. 2009;10:421.

795 77. Lu J, Salzberg SL. Removing contaminants from databases of draft genomes. *PLOS Comput*
796 *Biol*. 2018;14:e1006277.

797 78. Huerta-Cepas J, Serra F, Bork P. ETE 3: Reconstruction, Analysis, and Visualization of
798 Phylogenomic Data. *Mol Biol Evol*. 2016;33:1635–8.

799 79. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
800 resolution sample inference from Illumina amplicon data. Nat Methods. Nature Publishing
801 Group; 2016;13:581–3.

802 80. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing
803 taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-
804 classifier plugin. Microbiome. 2018;6:90.

805

Table 1. The characteristics for each dataset.

Sample	Sequencing Tech	Sequencing Strategy	# Basepairs	# reads	# Assembled contigs	# Mapped basepairs (Gb)
PD	Illumina	Shotgun	3.91 Gb	14525058	338823	3.69
		Amplicon	66.9/95.8/106.4 Mb ^a	39374/9614/10236 ^b	N/A	N/A
	Nanopore	Shotgun	1.96 Gb	1273484	N/A	N/A
		Amplicon	71.5/72.5/86.5 Mb	26212/ 26680/ 31826 ^b	N/A	N/A
PB	Illumina	Shotgun	3.67 Gb	13623120	345009	3.44
		Amplicon	55.7/38.1/71.9 Mb ^a	23613/13828/27093 ^b	N/A	N/A
	Nanopore	Shotgun	3.78 Gb	1043343	N/A	N/A
		Amplicon	54.5/49.4/42.0 Mb	20163/ 18273/ 15502 ^b	N/A	N/A

^a The total basepairs of each technical replicate were calculated before importing into QIIME2 pipeline.

^b Number of nanopore reads or paired-end Illumina reads for technical replicate 1/replicate 2/replicate 3 after quality control.

Table 2. Assignment of published sequence data to genera after application of cut-offs to query coverage.

Sample ID	Sample description	Sequencing tech	Cut-offs on query coverage (%)	Filtered results (%)	Percentage of confirmed genera BEFORE applying cut-offs (%)	Percentage of confirmed genera AFTER applying cut-offs (%)
a1	Human sputum samples ⁴²	Nanopore	59	20.2	85.9	86.5
a2			53.2	20.1	97.9	98.5
a3			54	20.5	96.5	97.4
a4			45.5	20.1	16.2	19.8
a5			58.5	20	71.1	66.9
a6			50.4	20.1	93.6	94.7
b1	Field infected wheat samples ⁴¹		5	20	60.4	75.1
b2			0.77	19.9	34.8	43
b3			12	19.7	67	82
b4			0.61	20	5.8	6.2
c1	Pig gut microbiome samples ⁴³	Illumina	2.4	20.1	32	35.4
c2			3.3	20.2	34.2	36.6
c3			2.6	20.2	35.2	38.3
d1	Mouse gut microbiome samples ⁴⁰		3.4	19.8	29.1	24.3
d2			14	20.1	63.7	69.4
d3			4.5	20.2	38.6	42.3

A **B** **C**

A**Illumina****Nanopore** **B****Illumina****Nanopore** **C****Illumina****Nanopore** **D****Illumina****Nanopore**

A

bioRxiv preprint doi: <https://doi.org/10.1101/2021.05.02.442318>; this version posted May 2, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

B

C

D

E