1 INTRODUCTION

2 Cell to cell communication is fundamental to multicellular life. Developmental signals critical
3 for tissue patterning and formation are traditionally considered as diffuse signals that
4 pervade the extracellular matrix. However, signals have often been found to be delivered to
5 target cells or received from source cells via specialized signalling filopodia — cytonemes
6 (Kornberg and Roy, 2014).

7 Stem cells, both embryonic and adult, regulate tissue formation. Many stem cells rely on
8 Wnts as signals for self-renewal and differentiation in the balance of cell fate determination
9 during development, tissue homeostasis and disease (Garcin and Habib, 2017). Wnt ligands
10 bind receptor Fzd, alongside co-receptors Lrp5 or Lrp6, ultimately leading to β -catenin–
11 dependent transcription of target genes. In several systems (Bertrand, 2016; Goldstein et al.,
12 2006; Huang and Niehrs, 2014; Kaur et al., 2020; Sugioka et al., 2011; Walston et al., 2004),
13 including mouse embryonic stem cells (ESCs) (Habib et al., 2013), localized Wnts induce
14 oriented asymmetric cell division (ACD).

15 As cultured single cells that retain the capacity to self-renew and differentiate to all adult
16 lineages, ESCs are an ideal experimental system for studying cell fate determination
17 (Martello and Smith, 2014). By combining ESCs and trophoblast stem cells (TSCs) to
18 generate 3D structures that resemble the developing embryo, we recently identified ESC-
19 generated cytonemes that mediate the interaction with Wnt-secreting TSCs (Junyent et al.,
20 2021, 2020). To overcome the multiplicity of signals and the difficulty in visualising secreted
21 Wnt ligands, we used Wnt3a tethered to microbeads, showing that ESC-cytonemes
22 selectively recruit self-renewal-promoting Wnt3a ligands. Cytonemes interacting with
23 localised Wnt3a signals engage not only Wnt-receptors, but also ionotropic AMPA/Kainate-
24 type glutamate receptors (iGluRs) (Junyent et al., 2021, 2020). Consequently, rapid calcium
25 (Ca^{2+}) transients are generated at the cytoneme, leading to Wnt3a reactivity: Wnt3a-bead
26 recruitment to the cell body (soma), activating Wnt/ β -catenin signalling and promoting
27 asymmetric cell division (ACD) of markers of cell fate (Habib et al., 2013).

28 iGluRs are primarily studied in excitatory glutamatergic synapses between neurons, however
29 they are increasingly being found in other cell types, such as stem cells (Genever et al.,
30 1999; Morhenn et al., 1994; Skerry and Genever, 2001) and functioning in *Drosophila*
31 development (Huang et al., 2019). AMPA- and Kainate-type iGluRs are tetrameric
32 glutamate-sensitive (amongst other ligands (Traynelis et al., 2010)) ion channels.
33 AMPA/Kainate-receptors require subunits GriA3 and GriA4, or GriK1 and GriK3 to function,
34 respectively (Cull-Candy et al., 2006; Lerma and Marques, 2013). Activation increases their
35 ion permeability, inducing rapid ion influxes critical to signal transmission.

1 However, it remained unknown how the interconnected roles of Wnt-pathway and
2 AMPA/Kainate-receptor components influence highly dynamic Wnt3a interactions, and the
3 subsequent Wnt3a-mediated ACD of ESCs.

4 Here, we resolve the dynamics of localized Wnt3a ligand (Wnt3a-bead) recruitment by single
5 ESCs into three stages: initial cytoneme-mediated interactions; Wnt3a-bead interaction with
6 the cell membrane; and the subsequent effects of the Wnt3a-bead recruitment on spindle
7 orientation and cell fate determination. We generated ESCs knocked-out (KO) for the AMPA-
8 and Kainate-type iGluRs by double knock-out (dKO) of critical protein subunits (AMPA:
9 GriA3/4dKO; Kainate: GriK1/3dKO). Multiple clonal cell lines were investigated alongside
10 pharmacological inhibition, both previously- and newly-generated mutants for Wnt co-
11 receptors (Lrp5, Lrp6 and Lrp5/6 dKO (Junyent et al., 2020)), and inducible β -catenin KO
12 ESCs (Raggioli et al., 2014)).

13 We show that while Lrp6, β -catenin and AMPA/Kainate-receptors are required for the
14 population-wide cytoneme-mediated uptake of Wnt3a-beads, neither AMPA- nor Kainate-
15 receptors are important to the formation of cytonemes themselves. Lrp5, conversely, has an
16 inhibitory role in regulating cytoneme generation in single ESCs and Wnt3a-bead
17 accumulation at cell-population level. In WT ESCs, Wnt3a ligands are recruited from initial
18 cytoneme interaction site at the cell periphery, towards the centre of the cell soma, a process
19 which requires Lrp5, Lrp6, β -catenin, AMPA- and Kainate-receptors. Upon progression to
20 mitosis, WT ESCs orient the spindle towards Wnt3a-beads but not inert (chemically
21 inactivated) iWnt3a-beads. Cells lacking Lrp5, Lrp6, β -catenin, AMPA- or Kainate-receptors
22 have significantly impaired ability to orient the spindle towards the Wnt3a signal source.

23 WT ESCs divide asymmetrically in response to a Wnt3a-bead, producing a self-renewed
24 proximal ESC cell and a distal cell that is prone to differentiation (Habib et al., 2013).
25 Specifically, we demonstrate that Lrp5 is not required in Wnt3a-mediated ACD, while Lrp6
26 and β -catenin are crucial. Importantly, we distinguish that AMPA-receptors, but not Kainate-
27 receptors, are critical in the cell fate determination of Wnt3a-mediated ACD in ESCs.

28 Taken together, we show that AMPA/Kainate-type iGluR activity is incorporated throughout
29 the interaction of ESCs with localized Wnt3a signals. We dissect the dynamic interaction of
30 Wnt3a-ligand with ESC, from initial uptake to cellular division, showing that its efficient
31 regulation requires Wnt co-receptors and downstream β -catenin in concert with
32 AMPA/Kainate-receptor activity. Our multiple clone approach to component KOs suggests
33 that the downstream cellular responses to localized Wnt3a of spindle orientation and
34 asymmetric cell division are impervious to clonal variation in the initial dynamics of the

1 Wnt3a-interaction. Through this, we specify novel differences between the roles of Lrp5 and
2 6, and between AMPA- and Kainate-type iGluRs in the response to localized Wnt3a.

3 **RESULTS**

4 **Wnt co-receptors and β -catenin regulate Wnt3a-bead accumulation by modifying the** 5 **cytonemes of ESCs**

6 We aimed to investigate how components of the Wnt/ β -catenin pathway (**Fig. 1A**) affect the
7 cellular dynamics of ESCs and their capacity to respond to localized Wnt3a ligands. We
8 used our previously generated ESC lines knocked-out (KO) for the co-receptors of the
9 Wnt/ β -catenin pathway: Lrp5 (Lrp5KO), Lrp6 (Lrp6KO) or both Lrp5 and 6 double KO
10 (Lrp5/6dKO) (described in (Junyent et al., 2020), here named KO #1). An additional clone of
11 each line was newly generated to investigate the influence of clonal variation during the
12 interaction with Wnt3a signals (named KO #2, **Fig. 1 – Figure Supplement 1**). We also
13 used an ESC line with a conditional KO of the pathway effector, β -catenin (β KO) (generated
14 as described in (Junyent et al., 2020; Raggioli et al., 2014)).

15 To validate the response of the clones of the KO cell lines to exogenous Wnt3a ligands, we
16 used cells harbouring the Wnt/ β -catenin pathway reporter, 7xTCF-eGFP, in which eGFP is
17 expressed upon pathway activation (Fuerer and Nusse, 2010) (**Fig. 1B and C**). Similar to
18 our previous findings (Junyent et al., 2020), in the presence of solubilized Wnt3a ligands
19 49.07% of wild-type (WT) cells express eGFP, while only a basal level of pathway activation
20 is observed in these cells in the absence of Wnt3a (**Fig. 1C**). In comparison, Lrp5KO ESCs
21 have a significant reduction in pathway activation in response to Wnt3a (26.97% eGFP⁺
22 cells). In Lrp6KO ESCs, pathway activation is further compromised (5.61% eGFP⁺ cells).
23 Lrp5/6dKO and β KO ESCs cannot activate the Wnt/ β -catenin pathway in response to
24 solubilized Wnt3a addition (**Fig. 1B and C**). We also measured the expression of *Axin2* (a
25 universal Wnt/ β -catenin target gene) in WT or KO ESCs stimulated with soluble Wnt3a.
26 Again, this demonstrated a reduction in Wnt/ β -catenin pathway activation in Lrp5KO and
27 Lrp6KO ESCs, and the absence of responsiveness to Wnt3a in Lrp5/6dKO or β KO cells
28 (**Fig. 1D**).

29 At the single cell level, ESCs use dynamic cytonemes to contact and recruit Wnt3a ligands
30 immobilized to a microbead (Wnt3a-beads) (Junyent et al., 2021, 2020). These cytonemes
31 are shorter in β KO and Lrp6-mutant ESCs (Lrp6KO, Lrp5/6dKO) than in WT ESCs, and are
32 presented in greater numbers in β KO cells (Junyent et al., 2020) (exemplified in **Fig. 1E**,
33 **Figure 1 – Figure Supplement 2, Movies 1-5**).

1 To investigate how differences in Wnt3a-response and single-cell behaviours affect the
2 accumulation of Wnt3a-beads at the population level over a prolonged period, we cultured
3 ESCs at low density and presented them with dispersed Wnt3a-beads. In some conditions,
4 we presented the cells with inactivated Wnt3a-beads (iWnt3a-beads), dithiothreitol (DTT)-
5 treated to denature the Wnt3a protein while covalently bound to the bead. Cells were then
6 live-imaged each minute for a total of 12 hours and the percentage of cells with cytonemes
7 as well as the proportion of cells contacting Wnt3a-beads were analysed.

8 WT ESCs begin forming cytonemes upon substrate attachment: within 4 hours, over 40% of
9 WT ESCs had cytonemes (**Fig. 2A**). This was coupled with a linear increase in the
10 proportion of WT ESCs with beads in the first 4 hours of imaging, rising to 54.1% by 4h, and
11 63.5% by 11 hours (**Fig. 3A**). ESC cytonemes are ligand-selective and can efficiently recruit
12 localized Wnt3a signals but not iWnt3a-beads (Junyent et al., 2020). Indeed, the percentage
13 of WT ESCs with iWnt3a-beads did not rise and remained less than 20% for most of the time
14 period (**Fig. 3A**).

15 Interestingly, >56% of Lrp5KO ESCs generate cytonemes in 3 hours, more than WT ESCs
16 (**Fig. 2A**). These cells retained the ability to quickly accumulate Wnt3a-beads, reaching
17 ~50% cells with beads after 3h and an overall mean of 61.5% after 6 hours (**Fig. 3B**). The
18 rate of increase in the percentage of Lrp6KO (**Fig. 2B**), Lrp5/6dKO (**Fig. 2C**) and β KO ESCs
19 (**Fig. 2D**) with cytonemes is largely similar to WT, with some clonal variability in the Lrp6KO
20 cells (**Fig. 2B**). Cytonemes of single ESCs lacking Lrp6 (Lrp6KO, Lrp5/6dKO) exhibit
21 reduced reactivity – decreased Wnt3a-bead uptake following initial contact (Junyent et al.,
22 2020). Indeed, Lrp6KO and Lrp5/6dKO showed long-term impaired Wnt3a-bead
23 accumulation at the population level, remaining at ~33% cells with beads over the course of
24 12 hours. This resembled the interaction of WT cells with iWnt3a-beads (**Fig. 3C, D and F**).
25 While cells lacking β -catenin (β KO) retain the capacity to efficiently recruit Wnt3a signals in
26 single-cell–single-bead interactions (Junyent et al., 2020), they had significantly impaired
27 long-term Wnt3a-bead accumulation at the population level, reaching a maximum mean of
28 32.7% cells with beads after 12 h (**Fig. 3E and F**).

29 Together with cytoneme formation, cell motility provides a mechanism to encounter and
30 interact with the local environment. To understand whether cytoneme formation or Wnt3a-
31 bead accumulation is coupled to cellular motility, we calculated the Mean Squared
32 Displacement per minute (MSD), the average movement per minute, of single cells. Cell
33 motility in WT ESCs is minimal, with a median MSD of $1.08 \mu\text{m}^2/\text{min}$ (**Fig. 2E**). No significant
34 differences to WT were observed between receptor or β -catenin KO ESCs, with the median
35 MSD for each cell type falling between 0.74 and $1.36 \mu\text{m}^2/\text{min}$ (**Fig. 2E**).

1 Altogether, we observed that intrinsic Wnt/ β -catenin components control the generation and
2 function of cytonemes without affecting overall cellular motility. Single cell effects, such as
3 shortening of the cytonemes (in β KO ESCs) or changes in the functionality of the cytonemes
4 (Lrp6KO, Lrp5/6dKO ESCs, or WT ESCs contacting iWnt3a beads) are reflected in
5 population-level changes in the rate of Wnt3a signal accumulation over a prolonged period.

6 Despite these differences, it is important to note that, in all conditions, some cells do take up
7 and retain Wnt3a-beads, allowing further analysis of the interaction with localized Wnt3a.

8 **Wnt3a-bead positioning at the membrane is controlled by Wnt co-receptors and β -** 9 **catenin**

10 Cytoneme-mediated Wnt3a recruitment is the first stage in the process of signal reception,
11 followed by a period of dynamic cell-signal interaction at the membrane of ESCs. Unless
12 interrupted by the loss of contact (Wnt3a-bead dropping) (**Fig. 4A**), this interaction often
13 persists until cell division. The positioning of the signal-receptor complexes on the plasma
14 membrane has been shown to affect signalling activity (Rangamani et al., 2013)(Schmick
15 and Bastiaens, 2014). The Wnt3a-bead approach provides a unique opportunity to quantify
16 the dynamics and positioning of localized Wnt3a on the cell membrane, as well as the roles
17 of the Wnt pathway co-receptors and β -catenin in its regulation. Therefore, we next analysed
18 the interaction of Wnt3a-bead and ESCs after initial cytoneme-mediated recruitment.

19 We followed cells that recruit a Wnt3a-bead with a cytoneme for 3h (180 min) after Wnt3a-
20 bead recruitment, a time point sufficient to measure the kinetics of initial cell-signal
21 interaction, and activate the downstream β -catenin related machinery of the pathway
22 (Hernandez et al., 2012)(**Movie 6**). We measured the distance between the Wnt3a-bead and
23 the cell centre, reporting it as a proportion of the cell radius (**Fig. 4B**).

24 WT ESCs initially recruit Wnt3a-beads with a cytoneme, and rapidly relocate them from the
25 tip to the membrane at the base (between 0.7 and 1.0 of the cell radius), then closer to the
26 cell centre (between 0.3 and 0.7 of the cell radius). Within 1 hour, Wnt3a-beads were, on
27 average, at ~0.5 of the cell radius (**Fig. 4B and C, Figure 4 – Figure Supplement 1, Movie**
28 **6**). iWnt3a-beads are also initially moved from the WT ESC cytoneme tip to the periphery
29 (0.7 to 1.0 of the cell radius). However, iWnt3a-beads are not positioned further towards the
30 cell centre, reaching at closest ~0.7 of the cell radius, on average (**Fig. 4B and C**).

31 ESCs lacking Lrp6 (Lrp6KO, Lrp5/6dKO) or β -catenin are likewise, on average, largely
32 unable to position the Wnt3a-bead closer to the cell body (**Fig. 4B and E-G, Movie 6**). In
33 these conditions the bead is stabilized after 90 min in mean positions ranging from 0.62 to
34 0.70 for Lrp6KO and Lrp5/6dKO, to ~0.70 for β KO ESCs. Lrp5KO ESCs position the Wnt3a-

1 beads at approximately 0.6 after 120 minutes, further from the centre of the cell than WT
2 ESCs (**Fig. 4D**), with clonal differences in the initial positioning.

3 The dynamics of Wnt3a-cell interaction are also defined by the speed of Wnt3a-bead
4 movement relative to the cell. Thus, we also calculated the MSD of the bead relative to the
5 centre of the cell (**Fig. 4H**). iWnt3a-beads held by WT ESCs have a higher rate of movement
6 than Wnt3a-beads, with a median MSD of 4.02 $\mu\text{m}^2/\text{min}$ versus 3.08 $\mu\text{m}^2/\text{min}$ for Wnt3a-
7 beads (**Fig. 4I**). In mutant ESCs contacting Wnt3a-beads, differences in Wnt3a-bead
8 movement are largely clone-dependent and independent of the genotype of the cell, with the
9 median MSD ranging between 4.25 $\mu\text{m}^2/\text{min}$ in Lrp5KO (#1) and 2.61 $\mu\text{m}^2/\text{min}$ in Lrp5/6dKO
10 (#2) (**Fig. 4I**).

11 Finally, we investigated whether positioning of the Wnt3a-bead further from the cell centre
12 lead to shorter Wnt3a-cell interactions. We observed that, within the first 3 hours of
13 interaction following cytoneme-recruitment, WT ESCs drop iWnt3a-beads at a significantly
14 higher rate than Wnt3a-beads (44.4% versus 4.08%, **Fig. 4J**). However, most mutant ESCs
15 that succeed in recruiting Wnt3a-beads retain them, with rates ranging between 100% of
16 cells retaining beads in Lrp5KO #2 and 80.4% cells retaining beads in Lrp6KO #2 (**Fig. 4J**).

17 Altogether, our results show that WT ESCs recruit and retain Wnt3a-beads at the
18 membrane, where they are positioned near the centre of the cell with limited movement.
19 However, WT ESCs do not recruit iWnt3a-beads efficiently and form unstable interactions
20 with them, positioning them at the periphery of the cells, where they are also dropped at
21 higher rates. While perturbation of the pathway components has a limited effect in recruited
22 Wnt3a-bead movement and Wnt3a-bead retention, KO ESCs exhibit a significantly impaired
23 ability to position Wnt3a-beads close to the centre of the cell.

24 **AMPA/Kainate-type receptors are required for dynamic interactions with localized** 25 **Wnt3a in ESCs**

26 A crosstalk between Lrp6 and the activity of ionotropic AMPA/Kainate-type glutamate
27 receptors (iGluRs) regulates the reactivity of ESC cytonemes to Wnt3a-beads (Junyent et
28 al., 2021, 2020). Here, we aimed to assess whether iGluR activity might also regulate the
29 dynamics of Wnt3a-stem cell interaction.

30 ESCs express the AMPA subunits GriA3 and 4, and the Kainate subunits GriK1, 2, 3 and 5
31 (Junyent et al., 2021, 2020). Previous publications have shown that GriA3/4 combinations
32 can form active AMPA-receptors that channel Ca^{2+} (Brorson et al., 2004; Cull-Candy et al.,
33 2006; Liu and Zukin, 2007), while Kainate receptor formation requires the presence of either
34 GriK1 or GriK3 (Lerma and Marques, 2013).

1 To investigate the role of AMPA and Kainate iGluRs in Wnt3a-ESC interaction, we
2 pharmacologically inhibited AMPA/Kainate iGluR activity by adding cyanquinoxaline (CNQX) to
3 the media, which reduces the generation of localized, Wnt3a-mediated Ca^{2+} transients in the
4 ESC cytonemes and impairs Wnt3a-bead recruitment (Junyent et al., 2020). To dissect the
5 specific role of each type of receptor family, we generated ESC lines with a double KO for
6 GriA3/4 (GriA3/4dKO, clone #1 and #2) and GriK1/3 (GriK1/3dKO, clone #1 and #2, **Fig. 5 –**
7 **Figure Supplement 1**).

8 We initially explored the capacity of these cells to activate the Wnt/ β -catenin pathway in
9 response to exogenous Wnt3a (200 ng/mL solubilized Wnt3a or a high density of Wnt3a-
10 beads, $6.25 \mu\text{g}/\text{cm}^2$, **Fig. 5A-C**) after 24h. In these conditions, cells were cultured as a
11 monolayer, and therefore Wnt3a-cell contact is cytoneme independent (**Fig. 5C**) (Junyent et
12 al., 2021, 2020). We analysed the relative expression of *Axin2* by qPCR (for GriA3/4dKO
13 and GriK1/3dKO) or the production of eGFP in response to 7xTCF-eGFP activation by FACS
14 (for CNQX-treated cells). In all conditions, Wnt3a addition led to activation of the Wnt/ β -
15 catenin pathway, to levels comparable to WT (**Fig. 5A and B**).

16 Next, we examined whether perturbation of AMPA/Kainate-receptors by KO or
17 pharmacological inhibition affected the capacity of ESCs to contact and recruit Wnt3a-beads.
18 At the single cell level, both GriK1/3dKO and GriA3/4dKO ESCs generate thin dynamic
19 cytonemes of similar number and length to WT ESCs (**Fig. 6A and B, Fig. 1 – Figure**
20 **Supplement 1, Movie 7 and 8**). These results corroborate previous observations in CNQX-
21 treated WT ESCs (Junyent et al., 2020), suggesting that AMPA/Kainate iGluRs do not play a
22 role in cytoneme formation.

23 At the population level, GriK1/3dKO ESCs, or WT ESCs + CNQX, generate cytonemes at a
24 rate similar to WT ESCs (approx. 44% of cells with cytonemes after 4h, **Fig. 6C, E and F**),
25 while the percentage of cells with cytonemes in GriA3/4dKO ESCs is equal or higher than
26 control cells (~67% after 4h for clone #2, **Fig. 6D and F**). However, Wnt3a-bead
27 accumulation is affected in Kainate-receptor-deficient cells: GriK1/3dKO ESCs show lower
28 Wnt3a-bead accumulation than WT ESCs, with a maximum mean of 49.5% cells with beads
29 over the 12 hours (**Fig. 6G**). This phenotype is even more pronounced in WT ESC treated
30 with CNQX, reaching only a mean of 39.9% of cells with Wnt3a-beads after 12h (**Fig. 6I and**
31 **J**).

32 GriA3/4dKO ESCs show clonal variability, with clone #1 exhibiting reduced Wnt3a-bead
33 uptake, and cells from clone #2 recruiting Wnt3a-beads at a rate similar to WT cells (**Fig. 6H**
34 **and J**). This is accompanied by a significant increase in motility, with a median cell MSD of
35 $\sim 1.4 \mu\text{m}^2/\text{min}$ for GriA3/4dKO, compared to $\sim 1.1 \mu\text{m}^2/\text{min}$ for WT ESCs (**Fig. 6K**).

1 Next, we explored the dynamics of Wnt3a-bead movement after initial recruitment by
2 AMPA/Kainate receptor-deficient cells. Using high temporal resolution live imaging (as
3 described in **Fig. 4B**), we observed that both GriK1/3dKO and GriA3/4dKO ESCs rapidly
4 relocate the Wnt3a-bead from the tip of the cytoneme to the cell soma, but stabilize the
5 position of the bead, on average, significantly further from the cell centre than WT ESCs
6 (~0.62 of cell radius between 30 and 120 min, **Fig. 7A and B**, expanded in **Fig. 7 – Figure**
7 **Supplement 1**). These cells show a slight increase in Wnt3a-bead movement, in a clone-
8 dependent, genotype-independent manner (**Fig. 7D**). In CNQX-treated ESCs, where both
9 AMPA and Kainate iGluRs are inhibited, the Wnt3a-beads are positioned far from the cell
10 centre (**Fig. 7C**), and with restricted movement (**Fig. 7D**).

11 Collectively, our data indicates that KO or inhibition of AMPA/Kainate-type receptors, much
12 like KO of Wnt co-receptor Lrp6 or pathway effector β -catenin, impairs Wnt3a-bead uptake
13 and alters the dynamics of Wnt3a-ESC interaction at the cell membrane. In all conditions,
14 importantly, once Wnt3a-cell contact has been achieved, it is often sustained until division,
15 enabling downstream analysis of the interaction (**Fig. 7E**).

16 A localized Wnt3a signal can induce spindle orientation and asymmetric cell division (ACD)
17 of single ESCs (Habib et al., 2013). We next investigated WT, Wnt/ β -catenin pathway and
18 AMPA/Kainate iGluRs KO ESCs throughout the process of Wnt3a-mediated ACD.

19 **Wnt/ β -catenin pathway components regulate the orientation of the mitotic spindle**

20 Wnt3a-bead orients the spindle of dividing ESCs (Habib et al., 2013). Spindle orientation is
21 an important factor in the process of ACD, in several biological systems (Bertrand, 2016;
22 Goldstein et al., 2006; Huang and Niehrs, 2014; Kaur et al., 2020; Sugioka et al., 2011;
23 Walston et al., 2004). Spindle orientation is established and can be determined as early as
24 the anaphase (Kiyomitsu and Cheeseman, 2013). Therefore, we measured the angle (α)
25 between the minor axis of mitosis and the point of bead contact, relative to the centre of
26 division, at anaphase (**Fig. 8A and B**). In this way, 90° represents a completely Wnt3a-bead
27 orientated division (**Fig. 8B**).

28 WT ESCs divide predominantly oriented towards the Wnt3a-bead, with a most common
29 angle of 85-90° (**Fig. 8C**). This Wnt3a-biased distribution is significantly different to a
30 randomized (Wnt3a-independent) distribution (Chi-squared test, $P < 0.001$), in which division
31 angle counts are assumed to be equally distributed between 0°-30°, 30°-60° and 60°-90°.
32 Indeed, when presented with iWnt3a-beads, a randomized distribution of angles is observed
33 (Chi-squared test, $P > 0.05$), with an increased proportion of cells dividing with iWnt3a-beads
34 between 0° and 5° (**Fig. 8D and L**), where the contact area is maximized and the free
35 energy is minimized (Koltover et al., 1999).

1 Single ESCs that lack Lrp5, Lrp6, or β -catenin exhibit spindle misorientation (**Fig. 8E-K**),
 2 with significantly different angle distributions than WT ESCs + Wnt3a beads, as determined
 3 by two-sample Kolmogorov-Smirnov (K-S) statistical analysis (**Fig. 8L**). In these cells, most
 4 cells divide at angles ranging from 0° to 60° for Lrp5KO, Lrp6KO and Lrp5/6dKO ESCs (**Fig.**
 5 **8E-J**), or with randomised angle distributions (Chi squared, $P > 0.05$) for β KO ESCs (**Fig. 8K**).

6 In conclusion, disruption of the Wnt co-receptors or the downstream component β -catenin
 7 impairs the spindle orientation towards a source of localized Wnt3a during division. While
 8 localized Wnt3a controls spindle orientation, it also controls cell fate, producing a self-
 9 renewing Wnt3a-proximal ESC and a Wnt3a-distal cell prone to differentiation (Habib et al.,
 10 2013). Next, we studied whether the individual KOs affect cell fate determination during
 11 ACD.

12 **Lrp6, but not Lrp5, is critical for asymmetric cell fate determination**

13 ACD is the partitioning or enrichment of cellular components in one of the two daughter cells
 14 deriving from mitosis, leading to differences in their cell fate. Wnt3a-beads induce ACD in
 15 single ESCs (Habib et al., 2013). Using this system, quantifying the expression of
 16 pluripotency markers Nanog, Rex1 and Sox2 (**Fig. 9A-D**) allows the comparison of the cell
 17 fate of the daughter cells: a higher expression in the Wnt3a-contacting cell indicates
 18 maintenance of a naïve ESC, whereas a lower expression in the Wnt3a-distal cell indicates
 19 a shift towards a more differentiated state.

20 To analyse ACD in WT or Wnt/ β -catenin pathway KO ESCs, we seeded single cells at low
 21 density in the presence of Wnt3a- or iWnt3a-beads for 6 hours prior to fixation. Then we
 22 analysed divided cell pairs where a bead is in clear contact with only one of the two daughter
 23 cells. This ensured that only cell pairs with an asymmetric presentation of the Wnt3a signal
 24 were analysed (**Fig. 9A**). We distinguished three categories: a “Proximal” doublet consisting
 25 of a cell contacting a Wnt3a-bead (Wnt3a-proximal) with higher pluripotency marker signal,
 26 and a Wnt3a-distal cell with reduced pluripotency marker expression. Conversely, a doublet
 27 where the farthest cell expresses more pluripotency marker was named a “Distal” doublet.
 28 An even distribution, with no significant bias to either cell, was named “Distributed”. Of these,
 29 only the “Proximal” condition represents a Wnt3a-mediated ACD.

30 When contacting a Wnt3a-bead, most WT ESC doublets have “Proximal” marker expression
 31 and have therefore undergone Wnt3a-mediated ACD (**Fig. 9B-D**). On the other hand, the
 32 majority of WT ESCs dividing with iWnt3a-beads are “Distributed”, with a similar partitioning
 33 of pluripotency markers between daughter cells. While Lrp5KO ESCs have a reduced ability
 34 to precisely align the spindle towards a Wnt3a signal (**Fig. 8E and F**), their capacity for
 35 Wnt3a-mediated ACD is retained (**Fig. 9B-D**). ESCs lacking Lrp6 or β -catenin (Lrp6KO,

1 Lrp5/6dKO, β KO) however, show impaired ACD (**Fig. 9B-D**), with cell pairs predominantly
2 having “Distributed” expression of Nanog, Rex1 and Sox2 between daughter cells (**Fig. 9B-
3 D**).

4 To validate the roles of Lrp6 and Lrp5, we transiently transfected Lrp5/6dKO ESCs with a
5 plasmid encoding GFP and the *Lrp6* sequence and sorted them into GFP⁺ (Lrp5KO, Lrp6
6 overexpression, *OE*) and GFP⁻ (Lrp5/6dKO) populations (**Fig. 10A and B**). Lrp5/6dKO cells
7 overexpressing Lrp6 (GFP⁺) could not efficiently align the spindle towards the Wnt3a-bead
8 during mitosis (**Fig. 10C**). However, Wnt3a-oriented ACD was rescued in these cells, as
9 shown by an increased rate of “Proximal” doublets, quantified for Sox2 and Nanog (**Fig.
10 10D-F**). Altogether, these results reiterate differences between Lrp5 and Lrp6 in the
11 regulation of spindle orientation and ACD.

12 Considering all evidence, we have shown that the co-receptor Lrp6 and β -catenin play co-
13 dependent roles in the positioning of localized Wnt3a signals at the membrane of ESCs, as
14 well as in regulating spindle orientation and ensuring asymmetric partitioning of cell fate
15 between the daughter cells. KO of these components in ESCs results in severely impaired
16 activation of the Wnt/ β -catenin pathway, as well as impaired recruitment of Wnt3a-beads.
17 Lrp5KO ESCs retain higher capacity to activate the Wnt/ β -catenin pathway in response to
18 solubilized Wnt3a, to recruit Wnt3a beads and to induce ACD, despite a compromised
19 spindle orientation towards the Wnt3a-bead. Importantly, regardless of clonal variability
20 observed in the initial interaction stages between the cells and the Wnt3a signals, the
21 downstream phenotypes observed are strongly retained when the cells divide.

22 **AMPA-type iGluRs participate in controlling Wnt3a-oriented ACD**

23 Our data has shown that AMPA/Kainate receptors activity is required for Wnt3a signal
24 recruitment and interaction dynamics in ESCs, suggesting a pervasive crosstalk with the
25 Wnt/ β -catenin pathway activity. Next, we explored whether such effects span beyond early
26 Wnt3a-stem cell interaction in modulating the process of division.

27 Wnt3a-beads fail to orient the spindle of AMPA/Kainate receptor-defective ESCs.
28 GriK1/3dKO, GriA3/4dKO and WT ESCs treated with CNQX show misoriented spindle
29 distributions (**Fig. 11A**). In Kainate-receptor KO cells, spindle orientation angles accumulate
30 between 0°-60°, while GriA3/4dKO ESCs or CNQX-treated cells present broader,
31 randomised spindle angle distributions (Chi-squared test, $P > 0.05$). All AMPA/Kainate-type
32 iGluR-defective cells are significantly different to the Wnt3a-biased spindle orientation
33 observed in WT ESCs (K-S statistical tests, **Fig. 11A**).

1 Post-division doublet analysis revealed that, while GriK1/3dKO ESCs largely exhibit Wnt3a-
2 oriented ACD, GriA3/4dKO or CNQX treatment reduces the proportion of “Proximal”
3 doublets dividing cells (**Fig. 11C-E**).

4 To confirm these results we treated WT ESCs with NBQX, a pharmacological inhibitor
5 specific to AMPA-receptors (Goldstein and Litwin, 1993). When compared to control ESCs
6 treated with DMSO, NBQX-treated cells also fail to orientate the spindle towards a Wnt3a-
7 bead (**Fig. 11B**) or undergo Wnt3a-oriented ACD (**Fig. 11F-H**).

8 Our data indicate that AMPA/Kainate-type iGluR activity has important roles throughout the
9 interaction between ESCs and localized sources of Wnt3a. GriK1/3dKO impairs Wnt3a-bead
10 recruitment without affecting the number or length of the cytonemes, or the rate of cytoneme
11 generation in the population. AMPA-type receptors (involving GriA3/4) play lesser roles in
12 ensuring Wnt3a-bead recruitment. Yet, both AMPA and Kainate iGluRs are important to
13 correctly position the bead at the cell membrane, and to align the mitotic spindle towards the
14 Wnt3a signal. Nevertheless, precise spindle orientation is also not required for Wnt3a-
15 oriented ACD in these cells: Kainate receptors are dispensable for ACD but lack of GriA3/4
16 leads to reduced asymmetric cell fate marker segregation.

17 Together with our analysis of Wnt/ β -catenin pathway KO ESCs, our results suggests that a
18 dynamic crosstalk between AMPA/Kainate-receptor signalling and Lrp6 is not only involved
19 in signal recognition and response, as previously shown, but is important to regulate the
20 spatial organisation and specification of dividing ESCs by Wnt3a.

1 DISCUSSION

2 Cell fate and spatial positioning occur in concert to achieve proper tissue organization. In the
3 stem cell niche, responsive stem cells receive Wnts as a cue for self-renewal (Clevers et al.,
4 2014; Garcin and Habib, 2017). In a spatial context, Wnt reception has been shown to
5 induce cell polarization and asymmetric cell division (ACD), in an evolutionarily conserved
6 manner (Goldstein et al., 2006; Huang and Niehrs, 2014; Mizumoto and Sawa, 2007;
7 Ouspenskaia et al., 2016; Sugioka et al., 2011; Walston et al., 2004).

8 Understanding initial ligand recruitment to stem cell membranes may benefit regenerative
9 medicine and the targeting of Wnt-related diseases. However, the study of dynamic,
10 localized Wnt reception by single stem cells is impeded by difficulties in visualising secreted
11 Wnts. The Wnt3a-bead (Habib et al., 2013; Lowndes et al., 2017) allows a temporal analysis
12 spanning the entire duration of the stem cell—Wnt3a-bead interaction: from cell morpho-
13 motility prior to contact and initial bead recruitment by cytonemes, to the dynamic interaction
14 at the cell membrane, ultimately leading to effects on spindle orientation and ACD.

15 AMPA/Kainate-type iGluR activity, and its crosstalk with the Wnt/ β -catenin pathway, controls
16 cytoneme-mediated recruitment of Wnt signals in mouse embryonic stem cells (ESCs)
17 (Junyent et al., 2020). Additionally, components of the Wnt pathway regulate the length and
18 number of cytonemes (Junyent et al., 2020). Conversely, we show in this study that
19 knockout of AMPA- or Kainate-receptors does not affect cytoneme formation.

20 The Wnt co-receptors Lrp5 and Lrp6 have mostly been studied in the context of β -catenin–
21 dependent signalling, where they have overlapping functions (He et al., 2004; Houston and
22 Wylie, 2002; Kelly et al., 2004; Tamai et al., 2000; Wehrli et al., 2000), albeit to different
23 levels of efficiency (Holmen et al., 2002; MacDonald et al., 2011). Here, we demonstrate
24 differing roles between them: endogenous expression of Lrp6 alone in ESCs (in Lrp5KO)
25 results in an increased rate of cytoneme formation over WT ESCs, while cells lacking Lrp6
26 (Lrp6KO and Lrp5/6dKO) have a reduced rate of cytoneme formation, suggesting a distinct
27 role for Lrp6 in cytoskeleton regulation.

28 Both Lrp5 and Lrp6 contain cytoplasmic domains that can bind to regulators of actin
29 filaments (Brown et al., 1998; Krause et al., 2003; Lian et al., 2016; Zhu et al., 2012),
30 however this region of Lrp5 has just 37% identity to Lrp6, compared to 69% overall
31 sequence identity between them (MacDonald et al., 2011). We speculate that such
32 differences might contribute to their differential roles in cytoneme formation/elongation: Lrp5
33 may compete with Lrp6 for the interaction of Wnt-pathway components or cytoskeletal
34 regulators. Hence, improved Lrp6-cytoskeleton interactions may facilitate the increased rate
35 of cytoneme formation and initial Wnt3a-bead recruitment observed in Lrp5KO ESCs.

1 WT ESCs use cytonemes to efficiently recruit Wnt3a-beads, and subsequently position them
2 near the centre of the cell. iWnt3a-beads cannot be recruited with the same efficiency.
3 Additionally, iWnt3a-beads are maintained nearer the periphery of the cell, with shorter and
4 less stable interactions. AMPA/Kainate-receptor activity is involved throughout the Wnt3a-
5 interaction. AMPA- or Kainate-receptor knockout (KO) ESCs, similarly to those lacking Lrp6
6 or β -catenin, are able to recruit Wnt3a-beads via cytonemes, but over 12 hours are
7 inefficient in their population-wide accumulation of Wnt3a-signal compared to WT or Lrp5KO
8 ESCs.

9 Beyond the initial interaction between cell and Wnt3a-ligand, a role for AMPA/Kainate-
10 receptors lies in controlling the dynamics of Wnt3a on the membrane. Unlike WT ESCs,
11 neither AMPA- nor Kainate-receptor KO cells are able to relocate the Wnt3a-bead from the
12 cytoneme-periphery and retain it near the centre of the cell. Pharmacological inhibition of
13 both AMPA- and Kainate-receptors together by CNQX resulted in a similar phenotype with
14 an enhanced effect. The KO of Wnt-pathway components Lrp5, Lrp6 or β -catenin also
15 reduces the ability to position a Wnt3a-bead on the membrane near the centre of the cell.

16 Following ligand-receptor binding, Wnt-pathway component polarization (Capelluto et al.,
17 2002) and oligomerization (Cong et al., 2004) are suggested to play roles in the
18 enhancement or modulation of Wnt signalling (Bilic et al., 2007). Here, the spatial
19 stabilization and colocalization of upstream and downstream pathway components at a
20 'signalling hub' may be critical for efficient signal transmission. In fact, we previously
21 demonstrated colocalization of Lrp6 and AMPA/Kainate-receptors during ESC interaction
22 with a Wnt3a-bead (Junyent et al., 2021).

23 Membrane receptors, including glutamate receptors (Yang et al., 2006), have been
24 demonstrated to flow from membrane protrusions towards the centre of the cell. This motion
25 can be promoted by directed movement of the actin cytoskeleton (Hanley, 2014; Yu et al.,
26 2010) and its efficiency can be governed by the activation of the pathway (Hartman et al.,
27 2009). Areas of relative membrane concavity, such as the cell centre away from the
28 periphery, have also been shown to enhance membrane receptor interactions with
29 intracellular components (Rangamani et al., 2013). This process may be required for more
30 efficient signalling (Schmick and Bastiaens, 2014). Therefore, the control of the Wnt3a-bead
31 positioning in WT ESCs may facilitate efficient protein interactions required for signal
32 transduction, enhancing downstream pathway activation.

33 After Wnt3a-bead recruitment and upon progression to mitosis, WT ESCs orientate the
34 spindle towards the source of localized Wnt3a at anaphase. Perturbation of the
35 AMPA/Kainate-receptors or Wnt/ β -catenin pathway components resulted in significant

1 impairment of the alignment of the spindle to localized Wnt3a, suggesting that both
2 pathways function in concert during this process.

3 Interestingly, AMPA-receptors are involved in Wnt3a-mediated ACD. While Kainate-receptor
4 KO had no significant effect, both AMPA-receptor KO and AMPA-receptor-specific inhibition
5 by NBQX significantly compromised Wnt3a-mediated ACD. Similarly, Wnt3a-mediated ACD
6 is impaired when ESCs lack Lrp6 or β -catenin. Lrp5KO ESCs, however, much like with
7 Kainate-receptor KO, remain able to asymmetrically segregate pluripotency markers to the
8 proximal cell, despite the lack of precise alignment of the spindle towards a Wnt3a-bead at
9 anaphase. Therefore, a robust spindle orientation towards a Wnt3a-bead is not required for
10 asymmetric fate outcome. Our data further demonstrates a differential requirement for Lrp5
11 and Lrp6 in ESCs, beyond the rate of cytoneme formation, in the regulation of cell fate.
12 Interestingly, the *Drosophila* neuroblast can tolerate certain degrees of spindle
13 misorientation due to disruption in internal polarity cues during early stages of cell division.
14 These cells can overcome initial misorientation and retain the prospective cell fate of the
15 daughter cells (Knoblich et al., 1995; Peng et al., 2000; Schober et al., 1999).

16 In summary, response to localized Wnt3a reception and the downstream Wnt3a-mediated
17 ACD is not orchestrated by the Wnt/ β -catenin pathway alone, but also by crosstalk with
18 AMPA/Kainate-type glutamate-receptor activity.

19 Glutamate-receptor activity is an evolutionarily conserved pathway that links homologous
20 proteins in prokaryotes (e.g. GluR0 (Chen et al., 1999)) and plants (GLRs (Forde and
21 Roberts, 2014; Ortiz-Ramírez et al., 2017)) with metazoans (iGluRs). In mammals, they are
22 well known for their function in excitatory synaptic signalling (Niciu et al., 2012). However,
23 there is a more ancient role for glutamate signalling that pervades throughout single- and
24 multi-cellular organisms alike: chemotaxis.

25 From non-neuronal mammalian cells (Gupta and Chattopadhyay, 2009), to plant
26 reproductive cells (Ortiz-Ramírez et al., 2017) and bacteria (Brumley et al., 2019), sensitivity
27 to glutamate or related molecules can provide a spatial signal that influences cellular
28 organisation. In *E. coli*, cytoplasmic proteins are segregated to the cellular poles by
29 interaction and colocalization with glutamate-sensitive chemoreceptors (Maddock and
30 Shapiro, 1993). This therefore presents a glutamate-receptor-mediated mechanism for the
31 spatial orientation of intracellular components in response to external signals. Further,
32 glutamate acts as a nutrient (Hottes et al., 2004) and chemoattractant (Felzenberg et al.,
33 1996) to the bacterium *Caulobacter crescentus*. This bacterium undergoes ACD to replicate
34 its 'stalked' form, and to generate a motile 'swarmer' cell. Crucially, in *C. crescentus*,
35 chemoreceptors are synthesized in the pre-divisional cell (Gomes and Shapiro, 1984),

1 before being asymmetrically polarized at division to the pole of the cell along with other
 2 protein components of the swarmer cell fate (Alley et al., 1992) . It remains to be determined
 3 whether the glutamate-sensing machinery is specifically required for the ACD of *C.*
 4 *crensentus* and other bacteria. However, these findings above suggest an evolutionarily
 5 conserved link between the sensing of external chemo-attractants such as glutamate, and
 6 the polarization and compartmentalisation of cell fate components.

7 As this ancient function for glutamate-receptor activity is present in ESCs during Wnt3a-
 8 mediated ACD, it may prove to be a mechanism that influences stem cell biology, cell fate
 9 and tissue organisation throughout mammalian adult stem cells and metazoans.

10

11

12

1 MATERIALS AND METHODS

2 Cell culture

3 Wild-type W4 (129S6/SvEvTac) mouse ESCs or ESCs knock-out for Lrp5, Lrp6 or
 4 Lrp5/6dKO (generated as described in (Junyent et al., 2020)) were maintained in ESC basal
 5 media containing Advanced DMEM/F12 (cat. num. 12634028, ThermoFisher), 10% ESC-
 6 qualified foetal bovine serum (FBS, cat. num. ES-009-B, Millipore), 1% penicillin-
 7 streptomycin (cat. num. P4333, Sigma), 2 mM GlutaMax (cat. num. 35050061,
 8 ThermoFisher), 50 μ M β -mercaptoethanol (cat. num. 21985-023, ThermoFisher) and 1000
 9 U/mL recombinant Leukaemia Inhibitory Factor (LIF; cat. num. 130-095-775, Miltenyi),
 10 supplemented with with 2i: MEK inhibitor PD0325901 (1 μ M, cat. num. 130-104-170,
 11 Miltenyi) and GSK3 inhibitor CHIR99021 (3 μ M, cat. num. 130-104-172, Miltenyi).
 12 Heterozygous β -catenin deficient ($\beta^{fl/-}$) ESCs, with an inducible flox β -catenin, and stably
 13 carrying a Cre-ER-T2 cassette under the control of a chicken β -actin promoter (generated in
 14 a W4 129S6/SvEvTac background by (Raggioli et al., 2014)) were cultured in the same
 15 conditions. Media was changed daily, and cells were grown at low density until formation of
 16 mid-sized colonies before passaging (every 3-4 days). To passage ESCs, colonies were
 17 rinsed with PBS, treated with 0.25% trypsin-EDTA (T-E, cat. num. 25200056, ThermoFisher)
 18 for 4 min at 37°C, 5%CO₂, resuspended using ESC basal media and centrifuged at 1200 x g
 19 for 4 minutes. Pelleted cells were resuspended in ESC basal media, counted and seeded in
 20 a clean tissue culture-treated 6-well plate at low density (~7,000 cells/well) in complete ESC
 21 media (containing LIF and 2i). In some experiments, 24 hours before use, 2i was withdrawn
 22 and replaced with soluble Wnt3a protein (200 ng/mL).

23 β -catenin knock-out in $\beta^{fl/-}$ cells was induced through the addition of 0.1 μ g/mL 4-hydroxy-
 24 tamoxifen (4-HT, cat. num. H6278, Sigma) to the media every 24 hours for 72 hours. Cells
 25 treated for 3 days with 4-HT were used in the experiments (referred to as β KO in the text).

26 All cell lines were maintained at 37°C, 5%CO₂ and were routinely tested for mycoplasma
 27 infection.

28 Pharmacological inhibition of iGluRs

29 When indicated in the text, figures or figure legends, cells were treated with the
 30 AMPA/Kainate iGluR inhibitor cyanquixaline (CNQX, 10 μ g/mL, cat. num. C127, Sigma) or
 31 the AMPA-iGluR specific inhibitor 2,3-dioxo-6-nitro-7-sulfamoyl-benzo[f]quinoxaline (NBQX,
 32 cat. num. 373, Tocris). Drugs were dissolved in ESC complete media (no 2i) to the desired
 33 concentration and were added to the cells at the beginning of the experiment. For Wnt/ β -

1 catenin pathway activation experiments, CNQX was added to the cells 1h before solubilized
2 Wnt3a or Wnt3a-bead addition.

3 Generation of KO ESCs by CRISPR/Cas9-mediated gene editing

4 Generation of KO ESCs was achieved by CRISPR/Cas9-mediated gene editing, following
5 (Ran et al., 2013). gRNA sequences adjacent to a protospacer adjacent motif (PAM) and
6 targeted at an early conserved exon in *Lrp5*, *Lrp6*, *GriA3*, *GriA4*, *GriK1* and *GriK3* were
7 designed in Geneious 11 (Biomatters). gRNAs were scored for on- and off-target effects
8 against the whole mouse genome, and sequences with low off-target score were synthesised
9 as oligonucleotides and cloned into a PX458 plasmid (Addgene plasmid #48138). gRNA
10 sequences used can be found in **Fig. 1 – Figure Supplement 1** and **Fig. 5 – Figure**
11 **Supplement 1**.

12 WT (to generate KO for *Lrp5*, *Lrp6*, *GriA3* and *GriK3*), *Lrp6*KO (to generate the *Lrp5/6*dKO
13 line), *GriA3*KO (to generate *GriA3/4*dKO cells) or *GriK3*KO (to generate *GriK1/3*dKO) ESCs
14 were transfected with 1 µg PX458 plasmid containing the sequence for the correct gRNA,
15 using JetPrime (cat. num. 114-07, Polyplus transfection) according to manufacturer
16 instructions. After 24h, transfected cells were sorted (as described in the “Fluorescence
17 activated Cell Sorting” section), and single GFP+ cells were collected in the wells of a 96-
18 well tissue culture-treated plate, in ESC complete media with LIF and 2i. Individual clones
19 were grown until formation of colonies, then expanded by passaging twice. Genomic DNA
20 was extracted using a DNeasy Blood and Tissue extraction kit (cat. num. 69506, Qiagen)
21 and the targeted region was amplified by *Taq* polymerase PCR (cat. num. M7501,
22 Promega). Sequence verification was achieved by TOPO-cloning (cat. num. 450159,
23 ThermoFisher) of the *Taq*-PCR fragments, transformation into Top10 chemically competent
24 bacteria (cat. num. C404006, ThermoFisher) and clonal selection. At least 12 TOPO-clones
25 per mutant clone were analysed by Sanger sequencing (Source Bioscience). Clonal ESC
26 lines with homozygous mutations that generated a frame-shift mutation inducing a premature
27 stop codon were selected for the experiments. KO ESCs were cultured as described above.

28 Preparation of Wnt3a-microbeads

29 Recombinant Wnt3a proteins were produced as described before (Habib et al., 2013;
30 Junyent et al., 2020), or purchased (cat. num. 1324-WN, R&D systems). Wnt3a proteins
31 were immobilized to 2.8 µm carboxylic acid-coated Dynabeads® (cat. num. 14305D,
32 ThermoFisher), as described before (Habib et al., 2013; Junyent et al., 2020; Lowndes et al.,
33 2017). Briefly, the carboxylic acid groups on the Dynabeads® were activated by a 30-minute
34 incubation with N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide (EDC, cat. num. 39391,
35 Sigma) and N-Hydroxysuccinimide (NHS, cat. num. 56480-25G, Sigma). 50 mg/mL EDC

1 and NHS solutions were prepared in 25 mM cold 2-(N-morpholino) ethanesulfonic acid
2 (MES, cat. num. M3671-50G, Sigma), at pH5. EDC/NHS incubation was performed in
3 constant rotation at room temperature (RT). Following activation, beads were retained using
4 a magnet and washed three times with 25 mM MES buffer (pH 5). Solubilized Wnt3a protein
5 (500 ng) was diluted 1:5 in MES buffer (pH 5) and incubated with the beads for 1 hour in
6 constant rotation, at RT. Beads were washed again three times with PBS (pH 7.4) before
7 storage in media containing 10% FBS at 4°C. Inactivation of Wnt3a beads was achieved
8 through incubation with 20 mM Dithiothreitol (DTT; Life Technologies, #P2325) for 30
9 minutes at 37°C. Following incubation with DTT, beads were washed three times in PBS
10 before storage in media containing 10% FBS at 4°C (up to 10 days). Before use in the
11 experiments, Wnt3a-bead activity was validated as described before (Habib et al., 2013;
12 Junyent et al., 2020; Lowndes et al., 2017).

13 Time-course live-cell imaging

14 To analyse the dynamics of ESCs and Wnt3a-bead interaction, 2,500 ESCs/well mixed with
15 0.3 µg Wnt3a-beads/well were seeded in a tissue culture-treated, clear-bottomed, black-
16 walled 96-well plate (cat. num. 3603, Corning) in complete media containing LIF (no 2i).
17 Cells and beads were allowed to settle at 37°C, 5% CO₂ for 30 min. Then, the plate was
18 placed in a Zeiss inverted Axio Imager epifluorescence microscope, equipped with a
19 CoolSNAP HQ2 CCD camera. Between 20-30 positions containing cells and beads were
20 selected and cells were imaged using a 10x/0.3 dry objective, for brightfield, every 1 minute
21 for 12 hours. The Zen 2 (Blue edition, Zeiss) software was used for acquisition. The resulting
22 videos were analysed in Fiji (ImageJ) as described in the next sections. Three or more
23 independent videos were generated for each cell line or condition (detailed in the figure
24 legends and the source data files).

25 To image the dynamics of the cytonemes in ESCs, 5,000 ESCs were seeded in a well of an
26 ibiTreat-modified µ-Slide 8 well slide (cat. num. 80826, ibidi) in ESC complete media with
27 LIF (no 2i). Cells were incubated at 37°C, 5% CO₂ for 4h to allow cell attachment and
28 cytoneme generation. Cells were stained with CellMask Deep Red plasma membrane stain
29 (1x concentration, cat. num. C10046, ThermoFisher) for 15 min at 37°C, 5%CO₂. Cells were
30 then rinsed once with ESC complete media and imaged immediately in a Nikon Eclipse Ti
31 Inverted Spinning Disk confocal (equipped with a Yokogawa CSU-1 disk head and an Andor
32 Neo sCMOS camera, and an incubation system at 37°C, 5% CO₂). Cells were imaged using
33 a Plan Apo lambda 40x/0.95 dry objective every 10 seconds, for 10 minutes. The resulting
34 videos were processed in Fiji (ImageJ) and presented in inverted grayscale.

35 Image analysis: Number and length of the cytonemes in ESCs

Brightfield time-course videos (at 1 min resolution) were used to measure the average number and maximum length of the cytonemes in WT, GriK1/3dKO and GriA3/4dKO ESCs. Cells from each condition that attach and generate cytonemes were selected. For each cell, the number and length of their cytonemes was measured from the cell centre, 10 times in timepoints separated by 10 minutes. The distance between the cell centre and the membrane was also measured in these timepoints. The average number of cytonemes, as well as the maximum length for any cytoneme was calculated, and the average cell centre to cell membrane distance was subtracted from the values. The normalized values were reported in the figure.

Image analysis: Percentage of cells with cytonemes or Wnt3a-beads over time

Brightfield videos of ESCs and Wnt3a-beads at 1 min resolution were also used to measure the percentage of cells presenting cytonemes or contacting Wnt3a-beads. Cytoneme generation was analysed every hour for the first 4 hours of each video (timepoints 0h to 4h), while Wnt3a-bead contact was analysed for the totality of the video (timepoints 0h to 12h). Several positions were analysed for each independent video, on Fiji (ImageJ), to count the total number of cells in the frame, the number of cells with cytonemes and/or the number of cells in contact with Wnt3a-beads. Cell counts in the different positions of each video were summed to avoid position-specific differences, and the percentage values were calculated per time-point, as presented in the figures.

Image analysis: Wnt3a-bead position and movement in the cell and cell movement

Single ESCs that recruited a Wnt3a-bead with a cytoneme and retained contact with it for ≥ 180 min were selected to measure Wnt3a-bead position and movement on the membrane. The same cells were used to measure cell movement. For each cell and each minute, the position of the cell centre, the Wnt3a-bead and the membrane were registered as X and Y coordinates using the “Manual tracking” plugin on Fiji (ImageJ).

First, the distances between the cell centre and the Wnt3a-bead ($Distance_{cell-bead}$) or the cell centre and the membrane ($Distance_{cell-membrane}$, cell radius) were calculated for each minute as:

$$Distance_{cell-bead} = \sqrt{(x_{bead} - x_{cell})^2 + (y_{bead} - y_{cell})^2}$$

$$Distance_{cell-membrane} = \sqrt{(x_{membrane} - x_{cell})^2 + (y_{membrane} - y_{cell})^2}$$

Then, the ratio between the $Distance_{cell-bead}$ and the $Distance_{cell-membrane}$ (cell radius) was calculated and reported in the figure.

1 Second, cell movement was calculated as mean squared displacement (MSD), based on the
2 position of the cell centre (P_{cell}) over the 180 min analysed, as:

$$MSD_{cell}(\tau) = \langle \Delta P_{cell}(\tau) \rangle = \langle [P_{cell}(t + \tau) - P_{cell}(t)]^2 \rangle$$

3 Where τ is the interval of time between two positions (1 min).

4 Finally, the normalized position of the Wnt3a-bead relative to the position of the cell
5 (Normalized position_{bead}, NP_{bead}) was calculated for each time point (t) as:

$$Normalized\ position_{bead}(t) = NP_{bead}(t) = (x_{bead}(t) - x_{cell}(t), y_{bead}(t) - y_{cell}(t))$$

6 In this way, the movement of the Wnt3a-bead was normalized to the movement of the cell.

7 Then, the MSD of the Wnt3a-bead was calculated with $\tau = 1\ minute$ as:

$$MSD_{bead}(\tau) = \langle \Delta NP_{bead}(\tau) \rangle = \langle [NP_{bead}(t + \tau) - NP_{bead}(t)]^2 \rangle$$

8 Image analysis: Percentage of bead retention/dropping

9 Brightfield videos of ESCs at 1 min resolution were also used to measure the percentage of
10 cells that retain or drop the Wnt3a-bead after recruitment. A sample of cells from each
11 condition were observed, and the time of Wnt3a-bead uptake with a cytoneme was
12 registered (t_{uptake}). Then, the cell was observed for a further 180 min ($t_{uptake+180}$): if a cell lost
13 contact with the Wnt3a-bead before $t_{uptake+180}$, the cell was qualified as “drops”; if contact was
14 maintained beyond $t_{uptake+180}$, the cell was qualified as “retains”. Cells that divided within the
15 observed period were excluded from the analysis. The percentage of cells that drop or retain
16 Wnt3a-beads over the total number of cells analysed was presented in the figures.

17 Image analysis: Spindle orientation

18 The videos at 1 min resolution were also used to measure spindle orientation. Cells dividing
19 in contact with a Wnt3a-bead were analysed. The “Angle” tool in Fiji (ImageJ) was used to
20 measure the angle between the minor axis and the Wnt3a-bead position at anaphase for all
21 conditions (**Fig. 8, 10 and Movie 9**). Rose-plots were generated with a custom Python script.

22 Immunofluorescence and measurement of asymmetric cell division

23 3,000 ESCs/well mixed with 0.3 μ g Wnt3a-beads/well were seeded onto human fibronectin
24 (10 μ g/mL, cat. um. 7340101, VWR/SLS) coated 8-well slides (cat. num. 80828, Thistle
25 Scientific) in ESC complete media containing LIF (no 2i). Slides were incubated for 6h at
26 37°C, 5% CO₂. After incubation, media was gently aspirated, and cells were fixed with 4%
27 paraformaldehyde for 8 min at RT. Cells were then washed 3 times in staining buffer (PBS
28 with 0.1% Bovine Serum Albumin (cat. num. A8806, Sigma), 0.05% NaN₃ (cat. num. S2002,

1 Sigma) and 0.001% Tween20 (cat. num. P2287, Sigma)) for 5 minutes each. Slides were
2 incubated in primary antibody diluted in staining buffer at 4°C overnight. Then, slides were
3 carefully washed 3 times for 5 minutes in staining buffer and incubated with AlexaFluor (AF)-
4 conjugated secondary antibodies diluted in staining buffer, for 1h at RT. Final washes were
5 performed with staining buffer containing DAPI (cat. num. D1306, ThermoFisher) at 1:1000
6 dilution. Slides were carefully mounted with ProLong Gold antifade mounting media (cat.
7 num. P10144, Life Technologies) and let to solidify for 24-48h before imaging.

8 Slides were imaged on a Zeiss inverted Axio Imager epifluorescence microscope using Zen
9 2 (Blue edition, Zeiss) software and equipped with a CoolSNAP HQ2 CCD camera. A
10 40x/1.3 oil immersion objective was used. Z-stacks were taken according to the optimal
11 interval. Images were processed, deconvolved, and quantified using Volocity software
12 (Perkin Elmer) or analysed in Fiji (ImageJ).

13 Quantification involved reconstruction of the cell based on “Sum” projection and manual
14 identification of divided cells. Cells were qualified as “Proximal”, “Distributed” or “Distal” by
15 three independent researchers according to the partitioning of Nanog, Rex1 or Sox2
16 intensities between the Wnt3a-bead-proximal and -distal cell. Additionally, samples of
17 doublets were allocated by semi-automated 3D intensity quantification in the Wnt3a-bead
18 proximal and -distal cell using Volocity (Perkin Elmer) or Fiji (ImageJ). In these cases, mean
19 fluorescent intensity in the proximal and distal cells was measured for each channel and
20 normalized to background intensity. The percentage of protein partitioning in the proximal or
21 distal cell was used to allocate doublets as follows: “Proximal”, >55% of protein in bead-
22 proximal cell; “Distributed”, 55% to 45% of protein in bead-proximal cell; “Distal”, <45% of
23 protein in bead-proximal cell.

24 Antibodies

25 The antibodies (Ab) used in this study were anti-Nanog (monoclonal rat Ab, clone eBioMLC-
26 51; 1:500, cat. num. 14-5761-80, ThermoFisher), anti-Rex1 (rabbit polyclonal Ab, 14HCLC;
27 1:1500, cat. num. 710190, ThermoFisher), anti-Rex1 (rabbit polyclonal Ab; 1:500, cat. num.
28 ab28141, Abcam), anti-Sox2 (mouse monoclonal Ab, clone 245610; 1:500, cat. num.
29 MAB2018, R&D systems) and AF488-, AF555- and AF647-tagged secondary antibodies
30 (donkey; 1:1000, cat. num. A-21208, A-31572, A-31571, ThermoFisher).

31 Fluorescence Activated Cell Sorting

32 Fluorescence activated cell sorting (FACS) was employed to detect Wnt3a response in WT,
33 KO or CNQX-treated ESCs stably expressing 7xTCF-eGFP//SV40-mCherry (Junyent et al.,
34 2020). Following incubation with control media (ESC basal media), media containing

1 solubilised Wnt3a (200 ng/mL), or addition of a high density of Wnt3a- or iWnt3a-beads
2 (6.25 $\mu\text{g}/\text{cm}^2$) for 24h, cells were harvested using 0.25% T-E, filtered and analyzed using a
3 FACSFortessa system (BD Biosciences). The gating strategy included gating for SSC-FSC
4 (cells), SSC-A-SSC-W (single cells), SSC-DAPI (alive cells), SSC-PE Texas Red (mCherry+
5 cells), and SSC-FITC (eGFP). GFP+/- gate was established using CNTRL-treated and
6 Wnt3a-treated WT ESCs. Analysis was performed using the FlowJo software (FlowJo).

7 FACS was also used to purify GFP+ and GFP- populations from an Lrp5/6dKO ESC line
8 transfected with pEGFP-Lrp6 (Habib et al., 2013), or to sort single GFP+ cells transfected
9 with PX458 CRISPR/Cas9 plasmids. Cells were transfected with 1 μg pEGFP-Lrp6 or
10 PX458 using JetPrime, 24h before sorting. Then, a cell suspension was prepared as
11 described before, and cells were sorted using a FACSAriaII system (BD technologies). The
12 gating strategy included gates for cells, single cells, alive cells and GFP (as described
13 before), but laser intensities were adjusted as required. GFP+/- gates were established
14 using control (non-transfected) ESCs and cells transfected with the plasmids. For pEGFP-
15 Lrp6 sorting, only Lrp5/6dKO ESCs transfected with the plasmid were used for sorting GFP
16 and GFP+ cells. Sorted cells were collected in complete ESC media containing LIF (no 2i),
17 counted and seeded for experiments (as described before). For PX458 sorting, single cells
18 were sorted in tissue culture-treated 96-well plates in ESC complete media (with LIF and 2i).

19 Quantitative PCR

20 Quantitative PCR (qPCR) was used to measure *Axin2* expression in response to Wnt/ β -
21 catenin pathway stimulation. 2.0×10^4 WT or KO ESCs/well were seeded in a tissue culture-
22 treated 12-well plate in ESC complete media supplemented with LIF (no 2i). Plates were
23 incubated at 37°C, 5% CO₂ for 24h. Then, media was aspirated and replaced with ESC
24 complete media (LIF, no 2i) with or without 200 ng/mL solubilized Wnt3a. Cells were
25 returned to the incubator for 24h. Cell lysis, RNA extraction and purification was performed
26 with an RNeasy mini kit (cat. num. 74106, Qiagen), following manufacturer instructions. 500
27 ng RNA were retrotranscribed with a QuantiTect reverse transcription kit (cat. num. 205313,
28 Qiagen) following manufacturer instructions. qPCR was prepared with 2x TaqMan Fast
29 Universal PCR MasterMix (cat. num. 4366072, ThermoFisher) and TaqMan probes for *Axin2*
30 (Mm00443610_m1, cat. num. 4331182, ThermoFisher) and GAPDH (Mm99999915_g1, cat.
31 num. 4351370, ThermoFisher). qPCR was performed in a Bio-Rad CFX384 system (Bio-
32 Rad). The threshold-cycle values (Ct) of *Axin2* were normalized first to Ct values of GAPDH
33 (DCt), and then to the average DCt values of the WT CNTRL condition (DDCt) and were
34 reported as $2^{-\text{DDCt}}$ in the figures.

35 Statistical analysis

1 Data representation and statistical analysis were performed using Prism (GraphPad), as
2 described in the figures. ANOVA (one- or two-way) analysis, followed by Tukey's, Dunnett's
3 or Šídák's multiple comparisons test was performed to compare data between conditions.
4 For analysis of spindle orientation, the Kolmogorov-Smirnov statistical test was performed to
5 compare between conditions of non-normally distributed data and Chi-square analysis was
6 performed to compare between observed data and the expected/randomized distribution.
7 The statistical analysis used for each assay is described in the figure legends. In all
8 experiments, biological replicates (n) for each assay comprise of experiments performed
9 independently from one another, and as such are from different cultured
10 populations/passages of cells, different sessions of imaging/measurement, and independent
11 media/reagent/cell dilution. Sample sizes (N) were selected such that the power of the
12 statistical analysis performed was sufficient, in accordance with previous studies (Habib et
13 al., 2013; Junyent et al., 2020). When indicated in the figure legends, the cells analysed
14 were pooled approximately evenly from at least three independent experiments, and
15 analysis were performed on the pooled data. Outliers were considered as individual data
16 points significantly outside the range of possible/meaningful values for each assay and were
17 excluded from subsequent analysis. For all figures, symbols indicate statistical significance
18 as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Non-significance was set as $p >$
19 0.05. All numerical source data, as well as statistical analysis results including adjusted p -
20 values can be consulted in the source data file for each figure.

21

22

23

24

1 ACKNOWLEDGEMENTS

2 This work was supported by a Sir Henry Dale Fellowship (102513/Z/13/Z) to S.J.H.

3

4 CONTRIBUTION

5 S.J., J.R. and S.J.H. designed the study, performed experiments and analysed data, wrote
6 the manuscript and prepared figures. J.L.A.S, C.L.G and T-J.T. performed experiments and
7 analysed data. M.W. and J.L-B. analysed data. S.J.H, conceived the idea, led the project
8 and provided financial support for the project.

9

10 COMPETING INTERESTS

11 No competing interests.

12

13 REFERENCES

- 14 Alley MRK, Maddock JR, Shapiro L. 1992. Polar localization of a bacterial chemoreceptor.
15 *Genes Dev* **6**:825–836. doi:10.1101/gad.6.5.825
- 16 Bertrand V. 2016. B-Catenin-Driven Binary Cell Fate Decisions in Animal Development.
17 *Wiley Interdiscip Rev Dev Biol* **5**:377–388. doi:10.1002/wdev.228
- 18 Bilic J, Huang Y-L, Davidson G, Zimmermann T, Cruciat C-M, Bienz M, Niehrs C. 2007. Wnt
19 Induces LRP6 Signalosomes and Promotes Dishevelled-Dependent LRP6
20 Phosphorylation. *Science (80-)* **316**:1619–1622. doi:10.1126/science.1137065
- 21 Brorson JR, Li D, Suzuki T. 2004. Selective Expression of Heteromeric AMPA Receptors
22 Driven by Flip-Flop Differences. *J Neurosci* **24**:3461–3470.
23 doi:10.1523/JNEUROSCI.5023-03.2004
- 24 Brown SD, Twells RCJ, Hey PJ, Cox RD, Levy ER, Soderman AR, Metzker ML, Caskey CT,
25 Todd JA, Hess JF. 1998. Isolation and characterization of LRP6, a novel member of the
26 low density lipoprotein receptor gene family. *Biochem Biophys Res Commun* **248**:879–
27 888. doi:10.1006/bbrc.1998.9061
- 28 Brumley DR, Carrara F, Hein AM, Yawata Y, Levin SA, Stocker R. 2019. Bacteria push the
29 limits of chemotactic precision to navigate dynamic chemical gradients. *Proc Natl Acad*
30 *Sci U S A* **166**:10792–10797. doi:10.1073/pnas.1816621116

- 1 Capelluto DGS, Kutateladze TG, Habas R, Finkielstein C V., He X, Overduin M. 2002. The
2 DIX domain targets dishevelled to actin stress fibres and vesicular membranes. *Nature*
3 **419**:726–729. doi:10.1038/nature01056
- 4 Chen GQ, Cul C, Mayer ML, Gouaux E. 1999. Functional characterization of a potassium-
5 selective prokaryotic glutamate receptor. *Nature* **402**:817–821. doi:10.1038/45568
- 6 Clevers H, Loh KM, Nusse R. 2014. Stem cell signaling. An integral program for tissue
7 renewal and regeneration: Wnt signaling and stem cell control. *Science* **346**:1248012.
8 doi:10.1126/science.1248012
- 9 Cong F, Schweizer L, Varmus H. 2004. Wnt signals across the plasma membrane to
10 activate the β -catenin pathway by forming oligomers containing its receptors, Frizzled
11 and LRP. *Development* **131**:5103–5115. doi:10.1242/dev.01318
- 12 Cull-Candy S, Kelly L, Farrant M. 2006. Regulation of Ca²⁺-permeable AMPA receptors:
13 synaptic plasticity and beyond. *Curr Opin Neurobiol* **16**:288–297.
14 doi:10.1016/j.conb.2006.05.012
- 15 Felzenberg ER, Yang GA, Hagenzieker JG, Poindexter JS. 1996. Physiologic, morphologic
16 and behavioral responses of perpetual cultures of *Caulobacter crescentus* to carbon,
17 nitrogen and phosphorus limitations. *J Ind Microbiol Biotechnol* **17**:235–252.
18 doi:10.1007/bf01574698
- 19 Forde BG, Roberts MR. 2014. Glutamate receptor-like channels in plants: A role as amino
20 acid sensors in plant defence? *F1000Prime Rep* **6**:1–7. doi:10.12703/P6-37
- 21 Fuerer C, Nusse R. 2010. Lentiviral Vectors to Probe and Manipulate the Wnt Signaling
22 Pathway. *PLoS One* **5**:e9370. doi:10.1371/journal.pone.0009370
- 23 Garcin CL, Habib SJ. 2017. A Comparative Perspective on Wnt / β -Catenin Signalling in
24 Cell Fate Determination In: J.-P. Tassan JZK, editor. Asymmetric Cell Division in
25 Development, Differentiation and Cancer, Results and Problems in Cell Differentiation
26 **61**, DOI 10.1007/978-3-319-53150-2_15. Springer International Publishing AG. pp.
27 323–350. doi:10.1007/978-3-319-53150-2
- 28 Genever PG, Maxfield SJ, Kennovin GD, Maltman J, Bowgen CJ, Raxworthy MJ, Skerry TM.
29 1999. Evidence for a novel glutamate-mediated signaling pathway in keratinocytes. *J*
30 *Invest Dermatol* **112**:337–342. doi:10.1046/j.1523-1747.1999.00509.x
- 31 Goldstein B, Takeshita H, Mizumoto K, Sawa H. 2006. Wnt signals can function as positional
32 cues in establishing cell polarity. *Dev Cell* **10**:391–6. doi:10.1016/j.devcel.2005.12.016

- 1 Goldstein JM, Litwin LC. 1993. NBQX is a selective non-NMDA receptor antagonist in rat
2 hippocampal slice. *Mol Chem Neuropathol* **18**:145–152. doi:10.1007/BF03160028
- 3 Gomes SL, Shapiro L. 1984. Differential expression and positioning of chemotaxis
4 methylation proteins in *Caulobacter*. *J Mol Biol* **178**:551–568. doi:10.1016/0022-
5 2836(84)90238-9
- 6 Gupta R, Chattopadhyay D. 2009. Glutamate is the chemotaxis-inducing factor in placental
7 extracts. *Amino Acids* **37**:271–277. doi:10.1007/s00726-008-0143-5
- 8 Habib SJ, Chen B-C, Tsai F-C, Anastassiadis K, Meyer T, Betzig E, Nusse R. 2013. A
9 Localized Wnt Signal Orients Asymmetric Stem Cell Division in Vitro. *Science (80-)*
10 **339**:1445–1448. doi:10.1126/science.1231077
- 11 Hanley JG. 2014. Actin-dependent mechanisms in AMPA receptor trafficking. *Front Cell*
12 *Neurosci* **8**:1–8. doi:10.3389/fncel.2014.00381
- 13 Hartman NC, Nye JA, Groves JT. 2009. Cluster size regulates protein sorting in the
14 immunological synapse. *Proc Natl Acad Sci U S A* **106**:12729–12734.
15 doi:10.1073/pnas.0902621106
- 16 He X, Semenov M, Tamai K, Zeng X. 2004. LDL receptor-related proteins 5 and 6 in Wnt/ β -
17 catenin signaling: Arrows point the way. *Development* **131**:1663–1677.
18 doi:10.1242/dev.01117
- 19 Hernandez AR, Klein AM, Kirschner MW. 2012. Kinetic Responses of β -Catenin Specify the
20 Sites of Wnt Control. *Science (80-)* **338**:1337–1340. doi:10.1126/science.1228734
- 21 Holmen SL, Salic A, Zylstra CR, Kirschner MW, Williams BO. 2002. A novel set of Wnt-
22 Frizzled fusion proteins identifies receptor components that activate β -catenin-
23 dependent signaling. *J Biol Chem* **277**:34727–34735. doi:10.1074/jbc.M204989200
- 24 Hottes AK, Meewan M, Yang D, Arana N, Romero P, McAdams HH, Stephens C. 2004.
25 Transcriptional profiling of *Caulobacter crescentus* during growth on complex and
26 minimal media. *J Bacteriol* **186**:1448–61. doi:10.1128/jb.186.5.1448-1461.2004
- 27 Houston DW, Wylie C. 2002. Cloning and expression of *Xenopus* Lrp5 and Lrp6 genes.
28 *Mech Dev* **117**:337–342. doi:10.1016/S0925-4773(02)00205-8
- 29 Huang H, Liu S, Kornberg TB. 2019. Glutamate signaling at cytoneme synapses. *Science*
30 *(80-)* **363**:948–955. doi:10.1126/science.aat5053
- 31 Huang YL, Niehrs C. 2014. Polarized Wnt signaling regulates ectodermal cell fate in
32 *Xenopus*. *Dev Cell* **29**:250–257. doi:10.1016/j.devcel.2014.03.015

- 1 Junyent S, Garcin CL, Szczerkowski JLA, Trieu T-J, Reeves J, Habib SJ. 2020. Specialized
2 cytonemes induce self-organization of stem cells. *Proc Natl Acad Sci* **117**:7236–7244.
3 doi:10.1073/pnas.1920837117
- 4 Junyent S, Reeves J, Gentleman E, Habib SJ. 2021. Pluripotency state regulates cytoneme
5 selectivity and self-organization of embryonic stem cells. *J Cell Biol* **220**.
6 doi:10.1083/jcb.202005095
- 7 Kaur S, Méléneć P, Murgan S, Bordet G, Recouvreur P, Lenne P-F, Bertrand V. 2020. Wnt
8 ligands regulate the asymmetric divisions of neuronal progenitors in *C. elegans*
9 embryos. *Development* dev.183186. doi:10.1242/dev.183186
- 10 Kelly OG, Pinson KI, Skarnes WC. 2004. The Wnt co-receptors Lrp5 and Lrp6 are essential
11 for gastrulation in mice. *Development* **131**:2803–2815. doi:10.1242/dev.01137
- 12 Kiyomitsu T, Cheeseman IM. 2013. XCortical dynein and asymmetric membrane elongation
13 coordinately position the spindle in anaphase. *Cell* **154**:391.
14 doi:10.1016/j.cell.2013.06.010
- 15 Knoblich JA, Jan LY, Jan YN. 1995. Asymmetric segregation of numb and prospero during
16 cell division. *Nature* **377**:624–627. doi:10.1038/377624a0
- 17 Koltover I, Rädler JO, Safinya CR. 1999. Membrane mediated attraction and ordered
18 aggregation of colloidal particles bound to giant phospholipid vesicles. *Phys Rev Lett*
19 **82**:1991–1994. doi:10.1103/PhysRevLett.82.1991
- 20 Kornberg TB, Roy S. 2014. Cytonemes as specialized signaling filopodia. *Dev* **141**:729–736.
21 doi:10.1242/dev.086223
- 22 Krause M, Dent EW, Bear JE, Loureiro JJ, Gertler FB. 2003. Ena/VASP Proteins: Regulators
23 of the Actin Cytoskeleton and Cell Migration. *Annu Rev Cell Dev Biol* **19**:541–564.
24 doi:10.1146/annurev.cellbio.19.050103.103356
- 25 Lerma J, Marques JM. 2013. Kainate receptors in health and disease. *Neuron* **80**:292–311.
26 doi:10.1016/j.neuron.2013.09.045
- 27 Lian G, Dettenhofer M, Lu J, Downing M, Chenn A, Wong T, Sheen V. 2016. Filamin A- and
28 formin 2-dependent endocytosis regulates proliferation via the canonical wnt pathway.
29 *Dev* **143**:4509–4520. doi:10.1242/dev.139295
- 30 Liu SJ, Zukin RS. 2007. Ca²⁺-permeable AMPA receptors in synaptic plasticity and
31 neuronal death. *Trends Neurosci* **30**:126–134. doi:10.1016/j.tins.2007.01.006
- 32 Lowndes M, Junyent S, Habib SJ. 2017. Constructing cellular niche properties by localized

- 1 presentation of Wnt proteins on synthetic surfaces. *Nat Protoc* **12**:1498–1512.
- 2 doi:10.1038/nprot.2017.061
- 3 MacDonald BT, Semenov M V., Huang H, He X. 2011. Dissecting molecular differences
- 4 between Wnt coreceptors LRP5 and LRP6. *PLoS One* **6**.
- 5 doi:10.1371/journal.pone.0023537
- 6 Maddock JR, Shapiro L. 1993. Polar location of the chemoreceptor complex in the
- 7 *Escherichia coli* cell. *Science (80-)* **259**:1717–1723. doi:10.1126/science.8456299
- 8 Martello G, Smith A. 2014. The nature of embryonic stem cells. *Annu Rev Cell Dev Biol*
- 9 **30**:647–675. doi:10.1146/annurev-cellbio-100913-013116
- 10 Mizumoto K, Sawa H. 2007. Cortical β -Catenin and APC Regulate Asymmetric Nuclear β -
- 11 Catenin Localization during Asymmetric Cell Division in *C. elegans*. *Dev Cell* **12**:287–
- 12 299. doi:10.1016/j.devcel.2007.01.004
- 13 Morhenn VB, Waleh NS, Mansbridge JN, Unson D, Zolotorev A, Cline P, Toll L. 1994.
- 14 Evidence for an NMDA receptor subunit in human keratinocytes and rat cardiocytes.
- 15 *Eur J Pharmacol Mol Pharmacol* **268**:409–414. doi:10.1016/0922-4106(94)90066-3
- 16 Niciu MJ, Kelmendi B, Sanacora G. 2012. Overview of glutamatergic neurotransmission in
- 17 the nervous system. *Pharmacol Biochem Behav* **100**:656–664.
- 18 doi:10.1016/j.pbb.2011.08.008
- 19 Ortiz-Ramírez C, Michard E, Simon AA, Damineli DSC, Hernández-Coronado M, Becker JD,
- 20 Feijó JA. 2017. GLUTAMATE RECEPTOR-LIKE channels are essential for chemotaxis
- 21 and reproduction in mosses. *Nature* **549**:91–95. doi:10.1038/nature23478
- 22 Ouspenskaia T, Matos I, Mertz AF, Fiore VF, Fuchs E. 2016. WNT-SHH Antagonism
- 23 Specifies and Expands Stem Cells prior to Niche Formation. *Cell* **164**:156–169.
- 24 doi:10.1016/j.cell.2015.11.058
- 25 Peng CY, Manning L, Albertson R, Doe CQ. 2000. The tumour-suppressor genes *lgl* and *dlg*
- 26 regulate basal protein targeting in *Drosophila* neuroblasts. *Nature* **408**:596–600.
- 27 doi:10.1038/35046094
- 28 Raggioli A, Junghans D, Rudloff S, Kemler R. 2014. Beta-Catenin Is Vital for the Integrity of
- 29 Mouse Embryonic Stem Cells. *PLoS One* **9**:e86691. doi:10.1371/journal.pone.0086691
- 30 Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. 2013. Genome engineering
- 31 using the CRISPR-Cas9 system. *Nat Protoc* **8**:2281–2308. doi:10.1038/nprot.2013.143
- 32 Rangamani P, Lipshtat A, Azeloglu EU, Calizo RC, Hu M, Ghassemi S, Hone J, Scarlata S,

- 1 Neves SR, Iyengar R. 2013. XDecoding Information in Cell Shape. *Cell* **154**:1356.
- 2 doi:10.1016/j.cell.2013.08.026
- 3 Schmick M, Bastiaens PIH. 2014. The interdependence of membrane shape and cellular
- 4 signal processing. *Cell* **156**:1132–1138. doi:10.1016/j.cell.2014.02.007
- 5 Schober M, Schaefer M, Knoblich JA. 1999. Bazooka recruits inscuteable to orient
- 6 asymmetric cell divisions in Drosophila neuroblasts. *Nature* **402**:548–551.
- 7 doi:10.1038/990135
- 8 Skerry TM, Genever PG. 2001. Glutamate signalling in non-neuronal tissues. *Trends*
- 9 *Pharmacol Sci* **22**:174–181. doi:10.1016/S0165-6147(00)01642-4
- 10 Sugioka K, Mizumoto K, Sawa H. 2011. Wnt regulates spindle asymmetry to generate
- 11 asymmetric nuclear β -catenin in C. elegans. *Cell* **146**:942–954.
- 12 doi:10.1016/j.cell.2011.07.043
- 13 Tamai K, Semenov M, Kato Y, Spokony R, Liu C, Katsuyama Y, Hess F, Saint-Jeannet JP,
- 14 He X. 2000. LDL-receptor-related proteins in Wnt signal transduction. *Nature* **407**:530–
- 15 535. doi:10.1038/35035117
- 16 Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, Hansen KB,
- 17 Yuan H, Myers SJ, Dingledine R. 2010. Glutamate Receptor Ion Channels: Structure,
- 18 Regulation, and Function. *Pharmacol Rev* **62**:405–496. doi:10.1124/pr.109.002451
- 19 Walston T, Tuskey C, Edgar L, Hawkins N, Ellis G, Bowerman B, Wood W, Hardin J. 2004.
- 20 Multiple Wnt signaling pathways converge to orient the mitotic spindle in early C.
- 21 elegans embryos. *Dev Cell* **7**:831–841. doi:10.1016/j.devcel.2004.10.008
- 22 Wehrli M, Dougan ST, Caldwell K, O’Keefe L, Schwartz S, Valzel-Ohayon D, Schejter E,
- 23 Tomlinson A, DiNardo S. 2000. Arrow encodes an LDL-receptor-related protein
- 24 essential for Wingless signalling. *Nature* **407**:527–530. doi:10.1038/35035110
- 25 Yang G, Huang A, Zhu S, Xiong W. 2006. It is time to move: Role of lateral diffusion in
- 26 AMPA receptor trafficking. *J Neurosci* **26**:9082–9083. doi:10.1523/JNEUROSCI.3018-
- 27 06.2006
- 28 Yu C han, Wu HJ, Kaizuka Y, Vale RD, Groves JT. 2010. Altered actin centripetal retrograde
- 29 flow in physically restricted immunological synapses. *PLoS One* **5**:1–9.
- 30 doi:10.1371/journal.pone.0011878
- 31 Zhu Y, Tian Y, Du J, Hu Z, Yang L, Liu J, Gu L. 2012. Dvl2-dependent activation of Daam1
- 32 and RhoA regulates Wnt5a-induced breast cancer cell migration. *PLoS One* **7**.

1 doi:10.1371/journal.pone.0037823

2

3

1 FIGURE LEGENDS

2 **Figure 1. Wnt3a responsiveness and formation of dynamic cytonemes in Wnt/ β -** 3 **catenin pathway KO ESCs**

4 **A.** Simplified schematic representation of the Wnt/ β -catenin pathway. Wnt3a binding to the
5 receptor Frizzled and the co-receptors Lrp5 and Lrp6 induces the inhibition of the β -catenin
6 destruction complex (Axin2, Gsk3, Ck1, APC) through Dvl. β -catenin is stabilized and
7 translocated to the nucleus, where it binds Tcf/Lef to initiate the transcription of Wnt3a target
8 genes.

9 **B.** (*Top-left*) Schematic representation of the 7xTCF-eGFP Wnt/ β -catenin reporter. (*Top-*
10 *right* and *Bottom*) Representative dot-plots depicting the levels of eGFP expression in WT or
11 Wnt pathway KO ESCs carrying the 7xTCF-eGFP reporter and treated with CNTRL solution
12 or 200 ng/mL soluble Wnt3a for 24h. Y axis is SSC. Values in plots are percentage of GFP+
13 cells.

14 **C.** Percentage of 7xTCF-eGFP⁺ cells measured by FACS, as described in **B**. n = 6 for WT, n
15 = 3 for KO ESCs. Bars and numbers are mean, error bars indicate SEM. Stars indicate
16 statistical significance calculated by two-way ANOVA with Dunnett's multiple comparison
17 test: **** $p < 0.0001$.

18 **D.** Relative *Axin2* expression in WT or Wnt pathway KO ESCs treated with CNTRL media or
19 200 ng/mL soluble Wnt3a for 24h, measured by qPCR. n = 3. Bars and numbers are
20 geometric mean, error bars indicate geometric SD. Stars indicate statistical significance
21 calculated by two-way ANOVA with Dunnett's multiple comparison test: **** $p < 0.0001$.

22 **E.** Representative images of the cytonemes in WT, Lrp5KO (#1), Lrp6KO (#1), Lrp5/6dKO
23 (#1) and β KO ESCs. Cells were stained with Cell Mask Deep Red and imaged for 10 sec
24 every 10 min on a spinning disk confocal, with a 40x/0.95 air objective (incubation at 37°C,
25 5%CO₂). Insets (*below*) are magnifications of black dashed boxes, contrast-enhanced for
26 clarity. Yellow arrowheads indicate thin protrusions. Time is in seconds. Scale-bars are 10
27 μ m.

28 Numeric data used in the figure can be found in the Source Data file.

29 **Figure 1 – Figure Supplement 1. Genotype of Wnt co-receptor KO ESCs**

30 **A – C.** Schematic representations of the genotype of Lrp5KO (#2) (**A**), Lrp6KO (#2) (**B**) and
31 Lrp5/6dKO (#2) (**C**). “CRISPR” sequence indicates the gRNA sequence used for the gene
32 editing, red NGG sequences indicate adjacent PAM (or CCN for reverse complement). “WT”

1 sequences indicate wildtype gene and “KO” sequences indicate modified gene sequences
2 (homozygous). Dashes indicate deleted base-pairs.

3 **Figure 1 – Figure Supplement 2. Representative images of cytonemes in Wnt co-** 4 **receptor KO ESCs**

5 Representative images of the cytonemes in Lrp5KO (#2), Lrp6KO (#2), Lrp5/6dKO (#2)
6 ESCs. Cells were stained and imaged as described in **Figure 1E**. Insets (*below*) are
7 magnifications of black dashed boxes, contrast-enhanced for clarity. Yellow arrowheads
8 indicate thin protrusions. Time is in seconds. Scale-bars are 10 μ m.

9 **Figure 2. Lrp5 controls the generation of cytonemes by ESCs**

10 **A – D.** Percentage of WT, Lrp5KO (clone #1 and #2) (**A**), Lrp6KO (#1 and #2) (**B**),
11 Lrp5/6dKO (#1 and #2) (**C**) and β KO ESCs (**D**) with cytonemes at 0 to 4h after seeding. Line
12 and dots are mean, error bars are SEM. $N \geq 33$ cells, $n \geq 3$ experiments. Stars and symbols
13 in box (right of each plot) indicate statistical significance to WT ESCs, calculated by two-way
14 ANOVA with Dunnett’s multiple comparison test (for **A – C**) or multiple unpaired t-tests (for
15 **D**): ns (non-significant, $p > 0.05$), * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. The same WT
16 ESCs data is presented in **A – D** for comparison.

17 **E.** Cell movement, presented as mean squared displacement per minute (MSD) for WT
18 ESCs co-cultured with Wnt3a- or iWnt3a-beads, or Wnt/ β -catenin pathway KO ESCs co-
19 cultured with Wnt3a-beads. $n \geq 36$ cells, pooled from ≥ 3 independent experiments, were
20 tracked for 3h after attachment. Box represents median and quartiles, error bars are 10-90
21 percentile, dots are data outside 10-90 range. ns indicates non-significant ($p > 0.05$)
22 differences calculated by one-way ANOVA with Dunnett’s multiple comparison test.

23 Numeric data used in the figure can be found in the Source Data file.

24 **Figure 3. Lrp6 and β -catenin are required for Wnt3a-bead accumulation**

25 **A – E.** Percentage of WT, Lrp5KO (#1 and #2) (**B**), Lrp6KO (#1 and #2) (**C**), Lrp5/6dKO (#1
26 and #2) (**C**) and β KO (**E**) ESCs contacting Wnt3a-beads or iWnt3a beads (indicated in
27 brackets) at 0 to 12h after seeding. Line and dots are mean, error bars are SEM. $N \geq 36$
28 cells, $n \geq 3$ experiments. The same WT ESCs + Wnt3a-beads data is presented in **A – E** for
29 comparison.

30 **F.** Statistical analysis of **A – E**, calculated for all conditions against WT ESCs + Wnt3a-
31 beads. Stars and symbols indicate statistical significance calculated by two-way ANOVA with
32 Dunnett’s multiple comparison tests: ns (non-significant, $p > 0.05$), * $p < 0.05$, ** $p < 0.01$,
33 *** $p < 0.001$, **** $p < 0.0001$.

1 Numeric data used in the figure can be found in the Source Data file.

2 **Figure 4. Wnt co-receptors and β -catenin orchestrate the interaction between ESCs**
3 **and Wnt3a-beads at the membrane**

4 **A.** Representative brightfield frames of time-course live imaging sequences displaying an
5 ESC contacting a Wnt3a-bead without recruitment (*left*), contact with recruitment (*middle*)
6 and a Wnt3a-bead recruitment with subsequent dropping (*right*). Scale-bars are 20 μ m.
7 Wnt3a-beads are highlighted by white dashed circles.

8 **B.** Representative brightfield image of an ESC contacting a Wnt3a-bead (*right*). Schematic
9 representation of the cell “periphery” (1-0.7 proportion of radius, brown), 0.7-0.3 proportion of
10 radius (orange) or “centre” of the cell (0.3-0 proportion of radius, yellow), and of the radius
11 used to calculate the bead position (white line) (*right*). Scale-bar is 20 μ m. Wnt3a-bead is
12 highlighted by white dashed circles.

13 **C – G.** Wnt3a-bead position in the cell, measured every minute for 3h (180 min) after initial
14 cytoneme-mediated recruitment, and presented as ratio to cell radius. Lines are mean of $n \geq$
15 38 cells/condition, pooled approximately evenly from ≥ 3 independent experiments. Single
16 cell tracks are presented in **Figure 4 - Figure Supplement 1**. Stars or symbols are statistical
17 significance, calculated from the average Wnt3a-bead position per cell in three
18 representative time windows (0 to 30 min, 31 to 120 min and 121 to 180 min) against the
19 position of Wnt3a-beads in WT ESCs, by two-way ANOVA with Dunnett’s (**D - F**) or Šídák’s
20 (**C and G**) multiple comparison test: ns (non-significant, $p > 0.05$), * $p < 0.05$, ** $p < 0.01$,
21 **** $p < 0.0001$. The same WT ESCs + Wnt3a-beads data is presented in **C – G** for
22 comparison.

23 **H.** Representative brightfield frames of a time-course live imaging, showing an ESC
24 contacting a Wnt3a-bead (highlighted with white dashed circle). Yellow track indicates
25 Wnt3a-bead movement on the cell over time. Time is in minutes. Scale-bar is 20 μ m.

26 **I.** Wnt3a-bead movement on the cell, presented as mean squared displacement per minute
27 (MSD). Bead movement was measured every minute for 3h (180 min) after cytoneme-
28 mediated recruitment in WT ESCs contacting Wnt3a- or iWnt3a-beads, and in Wnt pathway
29 KO ESCs contacting Wnt3a-beads. Box indicates median and quartiles, error bars indicate
30 10-90 percentile, dots are data outside 10-90 percentile range. $n \geq 38$ cells pooled
31 approximately evenly from ≥ 3 independent experiments. Stars indicate statistical
32 significance against WT ESCs + Wnt3a-beads, calculated by one-way ANOVA with
33 Dunnett’s multiple comparison test: ns (non-significant, $p > 0.05$), * $p < 0.05$, ** $p < 0.01$,
34 *** $p < 0.001$.

1 **J.** Percentage of cells that drop a Wnt3a-bead or iWnt3a-bead within 3h of initial recruitment
2 with a cytoneme (grey) or retain it for longer than 3h (blue). $n \geq 43$ cells/condition. Stars
3 indicate statistical significance calculated by multiple Fisher's exact tests: ns (non-significant,
4 $p > 0.05$), * $p < 0.05$, **** $p < 0.0001$.

5 Numeric data used in the figure can be found in the Source Data file.

6 **Figure 4 – Figure Supplement 1. Details of Wnt3a-bead position in WT or Wnt/ β -**
7 **catenin pathway KO ESCs membrane**

8 **A – I.** Detailed quantification of the Wnt3a-bead position on the membrane of WT or Wnt/ β -
9 catenin KO ESCs after initial cytoneme-mediated recruitment, including single-cell tracked
10 positions (grey lines) and mean position per condition (coloured line).

11 Numeric data used in the figure can be found in the Source Data file.

12 **Figure 5. iGluR inhibition or KO does not affect cytoneme-independent Wnt3a**
13 **responsiveness**

14 **A.** Relative expression of *Axin2* mRNA in a monolayer of WT, GriA3/4dKO or GriK1/3dKO
15 ESCs treated with control media (CNTRL), solubilized Wnt3a (200 ng/mL) or a high density
16 of Wnt3a-beads for 24h. Bars are mean of $n = 3$, error bars are SEM. Stars indicate
17 statistical significance, calculated by two-way ANOVA with Tukey's multiple comparison test:
18 ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

19 **B.** Percentage of 7xTCF-eGFP⁺ cells in control- or CNQX-treated cells upon addition of
20 control solution (CNTRL), solubilized Wnt3a (200 ng/mL), Wnt3a-beads or iWnt3a-beads at
21 coating concentrations. Bars are mean of $n = 3$, error bars are SEM. Stars indicate statistical
22 significance calculated by one-way ANOVA with Šidák's multiple comparison test:
23 **** $p < 0.0001$.

24 **C.** Representative brightfield image of a monolayer of cells treated with Wnt3a-beads. Scale-
25 bar is 50 μ m.

26 Numeric data used in the figure can be found in the Source Data file.

27 **Figure 5 – Figure Supplement 1. Genotype of GriA3/4dKO and GriK1/3dKO ESCs.**

28 **A – D.** Schematic representations of the genotype of GriA3/4dKO (clone #1 and #2) (**A and**
29 **B**) and GriK1/3dKO (clone #1 and #2) ESCs (**C and D**). “CRISPR” sequence indicates the
30 gRNA sequence used for the gene editing, red NGG sequences indicate adjacent PAM.
31 “WT” sequences indicate wildtype gene and “KO” sequences indicate modified gene

1 sequences. For GriA3/4dKO (#2), two mutant sequences were detected at equal rates.
2 Dashes indicate deleted base-pairs. Asterisk (*) indicates insertion.

3 **Figure 6. iGluRs control Wnt3a-bead recruitment without affecting the generation and**
4 **characteristics of the cytonemes.**

5 **A – B.** Average number of cytonemes/cell (**A**) and maximum cytoneme length/cell (**B**) in WT,
6 GriA3/4dKO, GriK1/3dKO and CNQX-treated ESCs. Boxes are median and quartiles, error-
7 bars are 10-90 percentiles, dots are data outside range. $n \geq 25$ cells/condition (**A** and **B**).
8 “ns” are non-significant differences calculated against WT ESCs by one-way ANOVA with
9 Dunnett’s multiple comparison tests.

10 **C – E.** Percentage of WT, GriK1/3dKO (clone #1 and #2) (**C**), GriA3/4dKO (#1 and #2) (**D**)
11 and CNQX-treated ESCs (**E**) with cytonemes at 0 to 4h after seeding. Line and dots are
12 mean, error bars are SEM. $N \geq 38$ cells, $n \geq 3$ experiments. WT ESC data from **Fig. 2A – D**
13 presented for comparison.

14 **F.** Statistical analysis of **C – E**. Stars and symbols in box indicate statistical significance of
15 each condition to WT ESCs, calculated by two-way ANOVA with Dunnett’s multiple
16 comparison test (for **C** and **D**) or multiple unpaired t-tests (for **E**): ns (non-significant,
17 $p > 0.05$), $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$.

18 **G – I.** Percentage of WT, GriK1/3dKO (#1 and #2) (**G**), GriA3/4dKO (#1 and #2) (**H**) and
19 CNQX-treated ESCs (**I**) contacting Wnt3a-beads at 0 to 12h after seeding. Line and dots are
20 mean, error bars are SEM. $N \geq 36$ cells, $n \geq 3$ experiments. WT ESCs + Wnt3a-beads data
21 from **Fig. 3A – E** presented for comparison.

22 **J.** Statistical analysis of **G – I**. Stars and symbols indicate statistical significance of each
23 condition to WT ESCs, calculated by two-way ANOVA with Dunnett’s multiple comparison
24 tests: ns (non-significant, $p > 0.05$), $^{*}p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$.

25 **K.** Cell movement, presented as mean squared displacement per minute (MSD) for WT,
26 GriK1/3dKO, GriA3/4dKO and CNQX-treated ESCs. $n \geq 44$ cells, pooled from ≥ 3
27 independent experiments, were tracked for 3h after attachment. Box represents median and
28 quartiles, error bars are 10-90 percentile, dots are data outside 10-90 range. Stars indicate
29 statistical differences calculated by one-way ANOVA with Dunnett’s multiple comparison
30 test: $^{*}p < 0.05$

31 Numeric data used in the figure can be found in the Source Data file.

32 **Figure 6 – Figure Supplement 1. Representative images of cytonemes in GriK1/3dKO**
33 **and GriA3/4dKO ESCs**

1 Representative images of the cytonemes in GriK1/3dKO (clone #1 and #2) and GriA3/4dKO
2 (clone #1 and #2) ESCs. Cells were stained and imaged as described in **Figure 1E**. Insets
3 (*below*) are magnifications of black dashed boxes, contrast-enhanced for clarity. Yellow
4 arrowheads indicate thin protrusions. Time is in seconds. Scale-bars are 10 μ m.

5 **Figure 7. GriK1/3dKO, GriA3/4dKO or CNQX-treated cells retain the beads further from**
6 **the cell centre**

7 **A – C.** Wnt3a-bead position in the cell surface, measured every minute for 3h (180 min) after
8 initial cytoneme-mediated recruitment, and presented as proportion of cell radius. Lines are
9 mean of $n \geq 45$ cells/condition, pooled approximately evenly from ≥ 3 independent
10 experiments. Single cell tracks are presented in **Figure 7 - Figure Supplement 1**. Stars or
11 symbols are statistical significance, calculated from the average Wnt3a-bead position per
12 cell in three representative time windows (0 to 30 min, 31 to 120 min and 121 to 180 min)
13 against the position of Wnt3a-beads in WT ESCs, by two-way ANOVA with Dunnett's
14 multiple comparison test: ns (non-significant, $p > 0.05$), * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,
15 **** $p < 0.0001$. WT ESCs + Wnt3a-beads data from **Fig. 4C – G** presented for comparison.

16 **D.** Wnt3a-bead movement on the cell, presented as mean squared displacement per minute
17 (MSD). Bead movement was measured every minute for 3h (180 min) after cytoneme-
18 mediated recruitment in WT GriK1/3dKO, GriA3/4dKO and CNQX-treated ESCs contacting
19 Wnt3a-beads. Box indicates median and quartiles, error bars indicate 10-90 percentile, dots
20 are data outside 10-90 percentile range. $n \geq 45$ cells pooled approximately evenly from ≥ 3
21 independent experiments. Stars indicate statistical significance against WT ESCs + Wnt3a-
22 beads, calculated by one-way ANOVA with Dunnett's multiple comparison test: ns (non-
23 significant, $p > 0.05$), * $p < 0.05$, *** $p < 0.001$.

24 **E.** Percentage of cells that drop a Wnt3a-bead within 3h of initial recruitment with a
25 cytoneme (grey) or retain it for longer than 3h (blue). $n \geq 37$ cells/condition. ns is non-
26 significant differences calculated by multiple Fisher's exact tests.

27 Numeric data used in the figure can be found in the Source Data file.

28 **Figure 7 – Figure Supplement 1. Details of Wnt3a-bead position in WT, GriK1/3dKO,**
29 **GriA3/4dKO and CNQX-treated ESCs membrane**

30 **A – E.** Detailed quantification of the Wnt3a-bead position on the membrane of WT,
31 GriK1/3dKO, GriA3/4dKO and CNQX-treated ESCs after initial cytoneme-mediated
32 recruitment, including single-cell tracked positions (grey lines) and mean position per
33 condition (coloured line).

1 Numeric data used in the figure can be found in the Source Data file.

2 **Figure 8. Components of the Wnt/ β -catenin pathway regulate the orientation of the**
3 **mitotic spindle of ESCs**

4 **A.** Representative brightfield frames of a time course live-cell imaging, showing an ESC
5 contacting a Wnt3a-bead and undergoing Wnt3a-oriented division. Wnt3a-bead is black
6 sphere. Dashed black box indicates the “Anaphase” timepoint used for spindle orientation
7 analysis. Scale-bar is 10 μ m.

8 **B.** (*Left*) Representative image of the method used for spindle angle measurement relative
9 to Wnt3a-bead position. Asterisks indicate cell poles used as reference for the
10 measurement. Orthogonal white lines depict major and minor axis, yellow line points to
11 Wnt3a-bead and shows the angle measured (α). Wnt3a-bead is highlighted with yellow
12 dashed circle. (*Right*) Representative images of different ESCs dividing with Wnt3a-beads at
13 angles between 0 and 90° (indicated in the figure). Scale-bars are 10 μ m.

14 **C – K.** Rose plots depicting the distribution of spindle angles in WT ESCs dividing with
15 Wnt3a-beads (**C**), WT ESCs dividing with iWnt3a-beads (**D**) or Lrp5KO ESCs (**E** and **F**),
16 Lrp6KO ESCs (**G** and **H**), Lrp5/6dKO ESCs (**I** and **J**) and β KO ESCs (**K**) dividing with
17 Wnt3a-beads. n (number of cells analysed) is depicted in the figure for each condition.

18 **L.** Statistical comparison of the distribution of spindle angles in all conditions to the one
19 observed in WT ESCs + Wnt3a beads. Stars indicate statistical significance calculated by
20 multiple Kolmogorov-Smirnov tests: ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

21 Numeric data used in the figure can be found in the Source Data file.

22 **Figure 9. β -catenin and Lrp6 are required for Wnt3a-mediated ACD**

23 **A.** Representative images of ESC doublets after dividing in contact with a Wnt3a-bead,
24 labelled with antibodies against Nanog (cyan), Rex1 (magenta) and Sox2 (greyscale), or
25 with 4',6-diamidino-2-phenylindole (DAPI, yellow). Images are maximum intensity projections
26 of a selected Z-stack. DIC is differential interference contrast. Panels display doublets with
27 higher marker expression in the Wnt3a-bead contacting cell (*top*, “Proximal”), with equal
28 levels of expression between both cells (*middle*, “Distributed”) or with higher marker
29 expression in the cell not contacting the Wnt3a-bead (*bottom*, “Distal”). The Wnt3a-beads
30 are highlighted by yellow dashed circles. Scale-bars are 20 μ m.

31 **B – C.** Percentage of doublets of WT or Wnt/ β -catenin KO ESCs contacting Wnt3a-beads or
32 iWnt3a-beads in the “Proximal”, “Distributed” or “Distal” categories for Nanog (**B**), Rex1 (**C**)
33 or Sox2 (**D**). n $\geq$ 3 experiments, with N $\geq$ 18 doublets/n (Nanog, Rex1) or N $\geq$ 28 doublets/n

1 (Sox2). For **D**, only clone #2 of Lrp5KO, Lrp6KO and Lrp5/6dKO ESCs were analysed. Box
2 indicates median and quartiles, error bars indicate 10-90 percentile. Dots are data outside
3 range. Stars and symbols indicate statistical significance for each condition against WT
4 ESCs + Wnt3a beads, calculated by two-way ANOVA with Dunnett's multiple comparison
5 test: ns (non-significant, $p>0.05$), * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

6 Numeric data used in the figure can be found in the Source Data file.

7 **Figure 10. Lrp6 overexpression in Lrp5/6dKO ESCs rescues ACD**

8 **A.** Schematic of the experimental pipeline. Lrp5/6dKO (#2) ESCs were transfected with
9 pEGFP-Lrp6 plasmids and sorted by FACS to Lrp6-expressing (GFP⁺) and Lrp6 non-
10 expressing (GFP⁻) populations before experimentation.

11 **B.** Representative dot-plot of GFP expression levels in Lrp5/6dKO control (non-transfected)
12 ESCs or Lrp5/6dKO ESCs transfected with pEGFP-Lrp6. GFP intensity was detected using
13 a FITC filter. Y axis is PE. GFP⁻ (blue) and GFP⁺ (magenta) cells were sorted only in the
14 transfected condition.

15 **C.** Rose plots depicting the distribution of spindle angle orientations in GFP⁺ or GFP⁻ ESCs
16 dividing in contact with a Wnt3a-bead. Stars and symbols in boxes (*right*) indicate statistical
17 significance calculated by multiple Kolmogorov-Smirnov tests against GFP⁺ ESCs (*top*) or
18 WT ESCs dividing with a Wnt3a-bead (*bottom*): ns (non-significant, $p>0.05$), *** $p<0.001$,
19 **** $p<0.0001$. n (number of cells) is depicted in the figure.

20 **D.** Representative images of GFP⁺ and GFP⁻ ESCs after dividing with a Wnt3a-bead, stained
21 with antibodies against Sox2 (cyan) and Nanog (magenta), or with DAPI (yellow). Wnt3a-
22 beads are highlighted by yellow dashed circle. Scale-bars are 20 μ m.

23 **E and F.** Percentage of doublets in the "Proximal", "Distributed" or "Distal" categories
24 (described in **Fig. 9A**) in GFP⁺ and GFP⁻ ESCs stained for Sox2 (**E**) and Nanog (**F**). Mean of
25 $n = 3$ independent experiments, $N \geq 29$ doublets analysed per n , error bars are SEM. Stars
26 indicate statistical significance calculated by two-way ANOVA with Šídák's multiple
27 comparison tests: * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

28 Numeric data used in the figure can be found in the Source Data file.

29 **Figure 11. AMPA-type iGluR receptors are important for Wnt3a-oriented ACD.**

30 **A and B.** Rose plots depicting the distribution of spindle orientations in GriK1/3dKO,
31 GriA3/4dKO or CNQX-treated ESCs (**A**) or DMSO- or NBQX-treated ESCs (**B**) dividing with
32 a Wnt3a-bead. n (number of cells analysed) is stated in the figures. Stars in boxes indicate

1 statistical significance calculated by multiple Kolmogorov-Smirnov test against WT ESCs +
2 Wnt3a-beads (in **A**) or DMSO-treated ESCs + Wnt3a-beads (in **B**): ** $p < 0.01$, **** $p < 0.0001$.

3 **C – E.** Percentage of doublets in the “Proximal”, “Distributed” or “Distal” categories
4 (described in **Fig. 9A**) for WT, GriK1/3dKO, GriA3/4dKO or CNQX-treated ESCs that have
5 divided with Wnt3a-beads. Quantification for Nanog (**C**), Rex1 (**D**) or Sox2 (**E**). $n \geq 3$
6 experiments, with $N \geq 18$ doublets/ n (Nanog, Rex1) or $N \geq 30$ doublets/ n (Sox2). Box
7 indicates median and quartiles, error bars indicate 10-90 percentile. Dots are data outside
8 range. Stars and symbols indicate statistical significance calculated by two-way ANOVA with
9 Dunnett’s multiple comparison test: ns (non-significant, $p > 0.05$), * $p < 0.05$, ** $p < 0.01$,
10 *** $p < 0.001$, **** $p < 0.0001$. WT ESC data from **Fig. 9B – D** presented for comparison.

11 **F – H.** Percentage of doublets in the “Proximal”, “Distributed” or “Distal” categories
12 (described in **Fig. 9A**) for DMSO- or NBQX-treated ESCs that have divided with a Wnt3a-
13 bead. Quantification for Nanog (**F**), Rex1 (**G**) or Sox2 (**H**). $n = 4$ experiments, with $N = 30$
14 doublets/ n . Box indicates median and quartiles, error bars indicate 10-90 percentile. Stars
15 indicate statistical significance calculated by two-way ANOVA with Šídák’s multiple
16 comparison tests: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

17 Numeric data used in the figure can be found in the Source Data file.

18 CAPTIONS FOR MOVIES

19 **Movie 1. WT ESC cytonemes.** High-quality images of the dynamic cytonemes of WT ESCs.
20 Cells were stained with Cell Mask Deep Red and imaged in a spinning disk confocal
21 microscope every 10 sec, for 10 min. Images are presented as inverted grayscale. Yellow
22 boxes are magnifications, contrast-enhanced for clarity. Time is minutes and seconds. Scale
23 bar, 10 μm .

24 **Movie 2. Lrp5KO ESC cytonemes.** High-quality images of the dynamic cytonemes of
25 Lrp5KO ESCs (clone #1 and #2). Cells were stained and imaged as described for **Movie 1**.
26 Images are presented as inverted grayscale. Yellow boxes are magnifications, contrast-
27 enhanced for clarity. Time is minutes and seconds. Scale bar, 10 μm .

28 **Movie 3. Lrp6KO ESC cytonemes.** High-quality images of the dynamic cytonemes of
29 Lrp6KO ESCs (clone #1 and #2). Cells were stained and imaged as described for **Movie 1**.
30 Images are presented as inverted grayscale. Yellow boxes are magnifications, contrast-
31 enhanced for clarity. Time is minutes and seconds. Scale bar, 10 μm .

32 **Movie 4. Lrp5/6dKO ESC cytonemes.** High-quality images of the dynamic cytonemes of
33 Lrp5/6dKO ESCs (clone #1 and #2). Cells were stained and imaged as described for **Movie**

1 1. Images are presented as inverted grayscale. Yellow boxes are magnifications, contrast-
2 enhanced for clarity. Time is minutes and seconds. Scale bar, 10 μm .

3 **Movie 5. βKO ESC cytonemes.** High-quality images of the dynamic cytonemes of βKO
4 ESCs. Cells were stained and imaged as described for **Movie 1**. Images are presented as
5 inverted grayscale. Yellow boxes are magnifications, contrast-enhanced for clarity. Time is
6 minutes and seconds. Scale bar, 10 μm

7 **Movie 6. Tracking of ESCs and Wnt3a beads.** Time-course live-cell brightfield images of a
8 WT or Lrp6KO ESC interacting with a Wnt3a-bead. Cell centre and Wnt3a-bead were
9 tracked for 180 min, and the tracks are displayed. Scale bars, 20 μm .

10 **Movie 7. GriA3/4dKO ESC cytonemes.** High-quality images of the dynamic cytonemes of
11 GriA3/4dKO ESCs (clone #1 and #2). Cells were stained and imaged as described for **Movie**
12 **1**. Images are presented as inverted grayscale. Yellow boxes are magnifications, contrast-
13 enhanced for clarity. Time is minutes and seconds. Scale bar, 10 μm .

14 **Movie 8. GriK1/3dKO ESC cytonemes.** High-quality images of the dynamic cytonemes of
15 GriK1/3dKO ESCs (clone #1 and #2). Cells were stained and imaged as described for **Movie**
16 **1**. Images are presented as inverted grayscale. Yellow boxes are magnifications, contrast-
17 enhanced for clarity. Time is minutes and seconds. Scale bar, 10 μm .

18 **Movie 9. Spindle orientation in dividing ESCs with Wnt3a-beads.** Time-course live-cell
19 brightfield images of a WT ESC dividing with Wnt3a-beads at $\sim 90^\circ$, $\sim 45^\circ$ or $\sim 0^\circ$. Spindle
20 orientation angle was always measured at anaphase. White lines in stills indicate minor and
21 major axis of the cell. Yellow line indicates the position of the Wnt3a-bead relative to the
22 minor axis. Scale bars, 20 μm .

23

24

25

26

27

Figure 1

Figure 1 - Supplement 1

A

Lrp5KO (#2)

Lrp5 (Chromosome 19, NC_000085.6)

PAM

CCGGGATGTGCGGCTAGTGG

CRISPR

2bp deletion

5' CTGTTGTTTGCCAA

CCG

CCGGGATGTGCGGCTAGTGGATGCCGGC 3' WT

5' CTGTTGTTTGCCAA

ACCG

CCGG-----TAGTGGATGCCGGC 3' KO

DNA

B

Lrp6KO (#2)

Lrp6 (Chromosome 6, NC_000072.6)

PAM

ACAGAGATTTACTGGCCAAA

CRISPR

19bp deletion

5' TATAATAAAC

ACAGAGATTTACTGGCCAAA

CGGACTGACTCTGGACT 3' WT

5' TATAATAAAC

ACAGAGATTT-----TCTGGACT 3' KO

DNA

C

Lrp5/6dKO (#2)

Lrp5 (Chromosome 19, NC_000085.6)

PAM

CCGGGATGTGCGGCTAGTGG

CRISPR

2bp deletion

5' CTGTTGTTTGCCAA

CCG

CCGGGATGTGCGGCTAGTGGATGCCGGC 3' WT

5' CTGTTGTTTGCCAA

ACCGC--GGATGTGCGGCTAGTGGATGCCGGC 3' KO

DNA

Lrp6 (Chromosome 6, NC_000072.6)

PAM

GTGCCAAAGATAGAACGTGC

CRISPR

10bp deletion

5' TGGGGAGAA

GTGCCAAAGATAGAACGTGC

TGGGATGGATGGCTCA 3' WT

5' TGGGGAGAA

GTGCCAA-----TGCTGGGATGGATGGCTCA 3' KO

DNA

Figure 1 - Supplement 2

Figure 2

A

B

C

D

E

Figure 3

Figure 4

Figure 4 - Supplement 1

Figure 5

A

B

C

Figure 5 - Supplement 1

A GriA3/4dKO (#1)

bioRxiv preprint doi: <https://doi.org/10.1101/2020.06.16.154609>; this version posted May 6, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

GriA3 (Chromosome 1, NC_000098.6) **GriA3** (Chromosome 1, NC_000098.6)

GGCGGTCTCTTTTAGTCC available CRISPR CC-BY 4.0 International license CRISPR

5' GCTCCGGGCGGTCTCTTTTAGTCC TGGGGCTTTT 3' WT | DNA 5' TTCCTCAGACCTCCCTCAAGTGCTCAGGATC 3' WT | DNA

5' GCTCCGGGCGGTCTCTTTT--TCCTGGGGCTTTT 3' KO | DNA 5'-----AGGATC 3' KO | DNA

GriA4 (Chromosome 9, NC_000075.6) 13pb deletion

GGATTTTGGGGACTCGCCAT CRISPR

5' TTTTCT**GGATTTTGGGGACTCGCCAT**GGGAGCCTTT 3' WT | DNA

5' TTTTCTGGATTTTGGG-----AGCCTTT 3' KO

B GriA3/4dKO (#2)

GriA3 (Chromosome X, NC_000086.7) 2bp deletion

GGCGGTCTTCTTTTAGTCC CRISPR

5' GCTCCGGGCGGTCTTCTTTTAGTCC TGGGCTTTT 3' WT

5' GCTCCGGGCGGTCTTCTTTT –TCCTGGGCTTTT 3' KO

DNA

GriA4 (Chromosome 9, NC_000075.6) 17pb deletion/ 1bp insertion

GGATTTTGGGGACTCGCCAT CRISPR

5' **GGATTTTGGGGACTCGCCAT**GGGAGCCTTTCCGAG 3' WT

5' GGATTTTGGGGACTCGC-----G 3' KO

5' GGATTTTGGGGACTCGC*CATGGGAGCCTTTCCGAG 3' KO

DNA

C GriK1/3dKO (#1)

bioRxiv preprint doi: <https://doi.org/10.1101/082161>; this version posted May 6, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

5' TTCCTC AGACCTCCCCTCAAGTGCTC AGGATC 3'		WT	DNA
5' -----AGGATC 3'		KO	

Grik3 (Chromosome 16, NC_000070.6) 14bp deletion

CCTCCGGAGTCTGGTTT CRISPR

5' GCGGCG**CCTCCGGAGTCTGGTTT**GGGAATACTG 3' WT | DNA

5' GCGGCGCCTCCGGAGT-----CTG 3' KO

D GriK1/3dKO (#2)

GriK1 (Chromosome 4, NC_000082.6) 8bp deletion

AGACCTCCCCTCAAGTGCTC	CRISPR
5' TTCCTC AGACCTCCCCTCAAGTGCTC AGGATC 3'	WT
5' TTCCTCAGACCTCCCCTCA-----GGATC 3'	KO

DNA

GriK3 (Chromosome 16, NC_000070.6) 14bp deletion

CCTCCGGAGTCTGGTTT CRISPR

5' GCGGCG**CCTCCGGAGTCTGGTTT**GGGAATACTG 3' WT | DNA

5' GCGGCGCCTCCGGAGT-----CTG 3' KO

Figure 6

Figure 6 - Supplement 1

Figure 7

Figure 7 - Supplement 1

Figure 8

A

B

C

D

E

G

I

K

F

H

J

L

Figure 9

A

B

C

D

Figure 10

Figure 11

A

B

GriK1/3dKO (#1)

+ Wnt3a beads (n=99)

GriA3/4dKO (#1)

+ Wnt3a beads
(n=102) av

CNQX

CC-BY 4.0

DMSO

(n = 91)

C

D

Nanog

Rex1

E

Sox2

F

G

H

Nanog

Rex1

Sox2