Suppression of Global Protein Translation in SARS-CoV-2 Infection

Haripriya Parthasarathy^{*,1}, Divya Gupta^{*,1}, Abhirami P Suresh^{§,1}, Dixit Tandel^{§,1,2}, Vishal Sah^{1,2}, and Krishnan Harinivas Harshan^{€,1}

^{*}These authors contributed equally

[§]These authors contributed equally

¹CSIR-Centre for Cellular and Molecular Biology, Hyderabad, India 500007

²Academy for Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India

[€]Correspondence: hkrishnan@ccmb.res.in

Keywords

SARS-CoV-2, COVID-19, Global translation, mTORC1, Polysome profiling, 5'-TOP

ABSTRACT

The relationship of SARS-CoV-2 with the host translation remains largely unexplored. Using polysome profiling of SARS-CoV-2 infected Caco2 cells, we here demonstrate that the virus induces a strong suppression of global translation by 48 hours of infection. Heavy polysome fractions displayed substantial depletion in the infected cells, indicating the loss of major translational activities in them. Further assessment of the major pathways regulating translation in multiple permissive cell lines revealed strong eIF4E dephosphorylation accompanied by Mnk1 depletion and ERK1/2 dephosphorylations. p38MAPK showed consistent activation and its inhibition lowered viral titers, indicating its importance in viral survival. mTORC1 pathway showed the most profound inhibition, indicating its potential contribution to the suppression of global translation associated with the infection. Pharmacological activation of mTORC1 caused a drop in viral titers while inhibition resulted in higher viral RNA levels, confirming a critical role of mTORC1 in regulating viral replication. Surprisingly, the infection did not cause a general suppression of 5'-TOP translation, as evident from the continued expression of ribosomal proteins. Our results collectively indicate that the differential suppression of mTORC1 might allow SARS-CoV-2 to hijack translational machinery in its favor and specifically target a set of host mRNAs.

INTRODUCTION

Severe acute respiratory syndrome- coronavirus 2 (SARS-CoV-2) is responsible for the current pandemic COVID-19 that has been wreaking havoc across the world, infecting millions and causing the death of over 3.22 million people over the past year (1). The newest member of the family *Coronaviridae* is a β -coronavirus with an approximately 30 kb long RNA genome with positive polarity. The enveloped viral particles are approximately 120 nm in diameter. The Spike protein on the outer surface of the virions,

characteristic of coronaviruses, binds to angiotensin converting enzyme 2 (ACE2) found on the surfaces of several cells acting as the entry receptor for the virus (2). The virus enters through endocytosis and its genetic material is released into the cytosol after the endosome-lysosome fusion results in the unpacking of the virion.

After its release into the cytosol, SARS-CoV-2 RNA undergoes translation as in other positive stranded RNA viruses (3, 4). The preliminary rounds of translation synthesize long polypeptides pp1a and pp1ab from ORFs 1a and 1ab respectively. These polypeptides are later cleaved by proteases to generate about sixteen functional polypeptides which together form the replicase complex (5). In addition to these ORFs, SARS-CoV-2 codes for at least nine distinct sub-genomic mRNAs of variable lengths with common 3'-UTRs. Translation of these mRNAs is believed to be temporally regulated (6), possibly indicating its significance in the viral life-cycle.

Viruses establish a unique relationship with the host protein translation machinery. The general understanding is that viruses are total parasites on the host translation and hijack this machinery for translating their own protein. This often provides the virus an unhindered access to the machinery to keep synthesizing its proteins. However, various viruses have distinct requirements based on their nature of relationship with the hosts. Viruses such as poliovirus completely shut down host translation and use the machinery for its own translation using a cap-independent mechanism (7). Several other viruses inhibit host translation to varying degrees while allowing a set of mRNAs to translate (4, 8). Yet, some other viruses such as hepatitis C virus (HCV) do not cause an apparent suppression of host translation, but still use a cap-independent mechanism for their translation. Members of Flaviviridae have a 5' capped genome but seem to be resistant

to the translational arrest imposed by them even though it affects host mRNAs (9).
Coronaviruses are known to inhibit host protein translation (6, 10-13). Nsp1 is reported
to interfere with host translation through its interaction with 40S ribosomes (6, 14-17).
Reports also indicate that translation efficiency of viral mRNAs are not higher than the
host mRNAs, but SARS-CoV-2 mediated preferential destruction of host mRNAs lead to
their reduced translation events (6, 18). However, the molecular mechanisms remain
much elusive.

Global translation activities in higher eukaryotes are regulated by three major pathways.
mTORC1 pathway is the most studied of these and is known to regulate translation of a
sub-set of mRNAs with a 5' terminal oligo pyrimidine (TOP) stretch (19-21). mTORC1 is
active in metabolically active cells and promotes translation by facilitating the free
availability of the cap-binding protein eIF4E (22). One of the substrates of mTORC1,
eIF4E binding protein (4EBP), inhibits translation activities by sequestering eIF4E (23).
mTORC1 mediated phosphorylation of 4EBP lowers its affinity towards eIF4E thereby
making it available for cap-binding. mTORC1 also facilitates translation by
phosphorylating ribosomal protein rpS6 (24), eIF4B and helicase eIF4A through
p70S6K (25). Several viruses are reported to target mTORC1 in order to suppress host
translation activities (13). Inhibition of mTORC1 is known to cause a major drop in
active polysomes and translation activities (20, 21).

MAPKs p38 and ERK1/2 are known to regulate the phosphorylation of eIF4E through
their substrate Mnk1/2 (26, 27). Even though Mnk mediated phosphorylation of eIF4E
does not alter its affinity for the 5' cap of the mRNAs, phosphorylated eIF4E is
commonly detected in several cancers leading several researchers to hypothesize that

this phosphorylation results in preferential translation of a set of mRNAs (28, 29). A third mechanism of regulation of global translation is the phosphorylation of eIF2 α at S52, a key event leading to reduced recycling rate of eIF2 ternary complexes that is critical for new events of translation initiations (30). Four kinases known as integrated stress response kinases coordinate this phosphorylation relaying various upstream signals. Protein kinase R (PKR), a dsRNA binding protein is one of these kinases that phosphorylates eIF2 α after the detection of dsRNA replication intermediates in the cytosol. This results in severe translational suppression in the virus infected cells as demonstrated in several cases (31, 32).

Coronavirus genome is 5' capped and polyadenylated indicating that they use the cap-dependent translation machinery. However, other coronaviruses were reported to inhibit host translation by various means (11). In this study, we investigated the relationship of SARS-CoV-2 with host translation machinery and regulatory networks. We demonstrate a severe dissociation of polysomes from 48 hours of infection that remained so during the rest of the course of infection. We did not find any evidence of eIF2 α participating in this translational decline. p38MAPK was phosphorylated throughout the course of the infection and its inhibition also resulted in lower viral titer. SARS-CoV-2 targeted Mnk1 levels thereby limiting eIF4E phosphorylation. The strongest inhibition was visible in the mTORC1 pathway where its substrates 4EBP1 and ULK1 showed loss in levels and phosphorylation. Our studies demonstrate that SARS-CoV-2 infection causes severe arrest of host translation machinery most likely through strong mTORC1 inhibition without impacting its own protein synthesis and suggests that the viral mRNAs employ unique means to continue their translation under these conditions.

RESULTS

Polysome profiles of SARS-CoV-2 infected cells demonstrate severe collapse of polysomes

We performed polysome profiling of Caco2 cells infected with SARS-CoV-2 (hCoV-19/India/TG-CCMB-O2-P1/2020) at multiple time intervals to map any changes in the global translation activity. The virus established infection by 24 hours post infection (hpi) as evident from the high levels of expression of the viral nucleocapsid (N) that continued until 96 hpi (Figure 1A). The expression of spike (S) peaked at 72 hpi and dropped thenceforth. Viral RNA replication increased until 72 hpi (Figure 1B). Interestingly, no major impact on polysome profiles was seen at 24 hpi while 48 hpi marked a remarkable collapse of polysomes with a modest swelling of the 80S peaks from 48 hpi (Figure 1 C-F). The heavy polysomes were particularly affected and this trend remained true until 96 hpi (Figure 1 C-F). Polysome profiling of cells infected with another strain of SARS-CoV-2 (hCoV-19/India/TG-CCMB-L1021/2020) induced an earlier collapse of the polysomes but confirmed the impact on the polysomes (Figure S1 A-D). Even though the polysomes underwent substantial dissociation, only a moderate swelling of the 80S was visible. This could be possibly due to the reported loss of host mRNAs by a selective degradation method initiated by Nsp1 of SARS-CoV-2 (16). At the same time, the translation of viral proteins continued unaffected (Figure 1A), confirming that the polysome dissociation is specifically targeting host mRNAs.

SARS-CoV-2 infection does not cause eIF2 α phosphorylation during the suppression of translation activities

eIF2 α phosphorylation mediated inhibition of translation initiation is frequently observed in several viral infections including SARS-CoV. In addition to the activation of PKR by dsRNA, interferon has also been demonstrated to cause eIF2 α phosphorylation mediated translational arrest (33). We analyzed this modification in SARS-CoV-2 infected Caco2 cells. A modest increase in eIF2 α phosphorylation observed at 24 and 48 hpi in the infected cells disappeared soon while a prominent collapse of polysome was apparent (Figure 2). A similar observation was made in the infected Calu-3 cells as well (Figure S2A) despite a robust viral replication (Figure S2B) validating that SARS CoV-2 mediated translational arrest is not mediated through eIF2 α phosphorylation. On the other hand, the infected Huh7 cells (Figure S2C) exhibited a curious increase in eIF2 α phosphorylation throughout the course of infection, indicating a possible cell-type specific effect on the ISR pathway (Figure S2D). Since no eIF2 α phosphorylation was evident concurrent with the collapse of polysomes in Caco2, this molecule is unlikely to have contributed to the translational suppression.

ERK1/2-Mnk1/2-eIF4E is inhibited during SARS-CoV-2 infection

eIF4E phosphorylation is often targeted under several physiological conditions and in certain viral infections (4, 8). We tested if SARS-CoV-2 targets this molecule in order to suppress host translation in Caco2 cells. Viral infection impacted the levels of several of the key molecules in this pathway beyond 48 hours of infection and hence we normalized the phosphorylation of these molecules and their abundance separately with the loading control. A moderate dephosphorylation of eIF4E at S209 residue was visible from 24 hours of infection (Figure 3).

Mnk1, the kinase that phosphorylates eIF4E, is regulated by two MAPKs, p38 and ERK1/2. Either of them has been demonstrated to activate Mnk1 through its phosphorylation. Mnk1 associated with eIF4G, the scaffold initiation factor of eIF4F complex, is activated upon phosphorylation and subsequently phosphorylates eIF4E. In agreement with the eIF4E dephosphorylation, Mnk1 also underwent dephosphorylation in SARS-CoV-2 infected cells (Figure 3), suggesting that the upstream MAPKs could be targeted by the viral infection. We subsequently analyzed the activation of the two MAPKs during SARS-CoV-2 infection in Caco2 cells. Consistent with the eIF4E and Mnk1 dephosphorylations, ERK1/2 dephosphorylation was evident in the infected cells from 24 hpi onwards, indicating that the upstream signals to ERK1/2 have been targeted during the infection (Figure 3). Major dephosphorylation of ERK1/2 and eIF4E was evident from 48 hpi in Huh7 cells as well (Figure S3). These results demonstrate that ERK1/2-Mnk-eIF4E pathway is targeted by SARS-CoV-2 infection at the abundance levels of the component molecules and additionally at their phosphorylation levels.

p38MAPK phosphorylated during SARS-CoV-2 infection is beneficial to the viral replication

Unlike ERK1/2, p38MAPK was phosphorylated in SARS-CoV-2 infected cells throughout the duration. The phosphorylation increased with time, with the most intense phosphorylation detected at 96 hpi, suggesting that this MAPK might be very important for the viral activities (Figure 4A). We tested this hypothesis by inhibiting SARS-CoV-2 infected Caco2 cells for 24 hours. The effect of inhibition of eIF4E phosphorylation was less remarkable in the infected cells as compared with the mock cultures similarly

inhibited, indicating the pressure from the viral replication. As we expected, inhibition of p38MAPK, confirmed by the dephosphorylation of eIF4E (Figure 4B), resulted in significantly lower intracellular viral RNA (Figure 4C) and infectious viral titer in the supernatant (Figure 4D) as compared against the untreated control culture. These results indicated that p38MAPK is activated in SARS-CoV-2 infected cells through specific upstream signals and this molecule plays important roles in SARS-CoV-2 biology.

SARS-CoV-2 inhibits mTORC1 and depletes its key substrates

4EBP1 is a key substrate of mTORC1 through which the complex regulates translation initiation. Active mTORC1 phosphorylates T37/46 in 4EBP1, causing a reduction in its affinity for eIF4E. This phosphorylation triggers phosphorylations at additional sites and the hyperphosphorylated 4EBP1 migrates slowly as compared with the hypo- and partly phosphorylated molecules. We analyzed the kinetics of phosphorylation of 4EBP1 during SARS-CoV-2 infection in Caco2 cells. As demonstrated in Figure 5A, 4EBP1 phosphorylation was significantly reduced in SARS-CoV-2 infected cells from 48 hpi onwards. As in the case of ERK1/2-eIF4E pathway, 4EBP1 was also depleted in the infected cultures. Despite this depletion, the dephosphorylation was more intense, indicating that mTORC1 activity was inhibited. p70S6K1 and ULK1, two other major substrates of mTORC1 were also dephosphorylated in these samples, further confirming the loss of activity of the kinase complex. Interestingly, dephosphorylation was accompanied by a significant loss in the levels of all these proteins as well, suggesting that mTORC1 pathway components are also targeted for their availability in the infected cells. These results were consistent in Huh7 cells also, validating this

mechanism across cell types (Figure S4A). Recent reports have demonstrated a global decay of host mRNA possibly driven by Nsp1 during SARS-CoV-2 infection (6, 18). We investigated the association of the loss of 4EBP1 and ULK1 upon infection with a potential degradation of their transcripts using quantitative RT-PCR and surprisingly detected significantly elevated levels of their transcripts in the infected cells indicating the involvement of post-transcriptional regulations (Figure 5 B and C). Thus, these transcripts are not part of the host mRNAs specifically degraded by viral proteins. These results demonstrate that SARS-CoV-2 targets mTORC1 pathway by suppressing its activity as well by targeting the expression of the key molecules in the pathway. Active viral translation during severe inhibition of mTORC1 indicates that mTORC1 is dispensable for the translation of SARS CoV-2 proteins.

Since mTORC1 regulates translation of a large number of transcripts including those encoding ribosomal proteins through 5' TOP elements, we asked if the inhibition of mTORC1 pathway negatively impacts ribosomal biogenesis. Analysis of ribosomal proteins rpS3, rpL13a and rpL26 revealed that their expressions are not affected by SARS-CoV-2 infection (Figure 5D). Thus, despite a strong polysome dissociation and inhibition of mTORC1, ribosomal protein synthesis goes on unabated indicating that inhibition of mTORC1 activity is not affecting the translation of 5' TOP mRNAs. This part of the data suggests that SARS-CoV-2 brings about translational suppression through a remarkable inhibition of mTORC1 and the suppression could be selectively targeting a set of mRNAs.

mTORC1 restricts SARS CoV-2 replication

Since SARS CoV-2 infection caused strong suppression of mTORC1, we investigated whether this inhibition benefits the virus. Huh7 cells infected with SARS CoV-2 for 24 hours were treated with 10 μ M MHY1485 to activate mTORC1. The drug failed to induce mTORC1 activity (4EBP1 phosphorylation) in the virus infected cells while its activation was detected in the mock-infected cells (Figure 6A), indicating that the virus infection overrides the activation of mTORC1 by the drug. Interestingly, activation of mTORC1 resulted in decreased intracellular RNA as well as infectious titer of the virus (Figure 6 B and C respectively). In agreement with this observation, a moderate drop in the nucleocapsid levels was also visible (Figure 6A). These observations suggest that lower mTORC1 activity is beneficial for SARS-CoV-2 replication.

Next, we inhibited mTORC1 by Torin1 and investigated its effect on the infection. After infecting the cells with SARS CoV-2 for 2 hours, they were treated with 750 nM Torin1 until 24 hpi before analyzing the intracellular viral RNA. mTORC1 inhibition, confirmed by the dephosphorylation of 4EBP1 (Figure 6D), caused a two-fold increase in intracellular viral RNA levels (Figure 6E), strengthening the observations made in the preceding experiment that mTORC1 inhibition favors the viral replication. Our results indicate that mTORC1 inhibition might facilitate SARS-CoV-2 replication.

DISCUSSION

Several studies have indicated that SARS-CoV-2 infection suppresses host protein translation (2-4, 6, 12). While some have speculated this observation based on the reports from similar β -coronaviruses, others have implicated this based on the host mRNA degradation mediated by SARS-CoV-2 Nsp1 (6, 18). Nsp1 was also shown to associate with 40S ribosomes and block the entry of mRNAs (15). Our study provides a

detailed map of the impact of SARS-CoV-2 on global translation and the signal pathways that regulate the process. Polysome profile kinetics provided striking evidence of the suppression of host translation from around 48 hpi.

Even as other studies have reported global degradation of host mRNAs (6, 17), we have not come across any evidence that testifies this observation from our studies. Widespread host mRNA degradation would have resulted in the accumulation of the short and free nucleotides in mRNP fractions that our studies have not observed. Similar studies done in our laboratory using a flavivirus JEV show a significant swelling in 80S peaks concurrent with polysome dissociation as infection progressed, which wasn't as apparent in SARS-CoV-2 infected cells (data not shown). The fact that the 80S peak did not undergo any shortening at the later time intervals suggested that a significant fraction of mRNAs are still associated with monosomes and could be translation-ready, as is evidenced by a sustained maintenance of lighter polysomes throughout the course of infection. Thus, a considerable proportion of the host mRNA population is likely to be intact despite being subject to specific degradation by viral factors. Justifying this claim, *4EBP1* and *ULK1* mRNAs were detected at significantly higher levels in the infected cells. This could have been a reflection of their transcriptional activation or enhanced stabilization of the transcripts, either of which indicates that they are not subject to degradation. Nsp1 mediated blocking of the host mRNAs from accessing 40S ribosomes might also have resulted in significant drop in the 80S assembly. However, a clear enlargement of 80S fraction was visible in cells expressing Nsp1 (17) indicating that the regulation is more complex. Interestingly, no such information is available for MERS in the literature. Further detailed studies are

necessary to understand the larger impact of SARS-CoV-2 infection on 80S and polysome assembly.

We have observed a systematic depletion of several host proteins during the course of viral infection, particularly at later stages. Majority of these included substrates of mTORC1 and members of MAPK pathway. Since 4EBP1 and ULK1 were not subject to mRNA degradation, it is very apparent that post-transcriptional and post-translational mechanisms targeting specific host proteins are quite pervasive in SARS-CoV-2 infected cells.

mTORC1 was strongly inhibited by SARS-CoV-2. Targeting mTORC1 seems to be more concerted and with purpose since the substrates were also depleted at protein level. Justifying this point, conditions of lower mTORC1 activities promoted viral replication and its activation lowered the titers. It appears that post-transcriptional regulations play a role in their abundance in the infected cells. The implication of lower availability of 4EBP1 on the translation of host and viral mRNAs is unclear at this stage. Lower abundance of this inhibitory molecule could be interpreted to be facilitating eIF4F assembly and capped translation. However, the lower activity of mTORC1 also resulted in lower p70S6K phosphorylation indicating that the net impact of its inhibition results in reduced polysome assembly and translation activities. Interestingly, ribosomal proteins that we tested remained abundantly available in the infected cells and this might be important for the translation of viral proteins. Thus, it appears that mTORC1 inhibition does not target all 5'TOP mRNAs but must be targeting a select set of mRNAs without compromising the requirements of the virus.

How mTORC1 inhibition is brought about by SARS-CoV-2 is unclear. A recent study (34) reported that SARS-CoV-2 rewires metabolic pathways in the infected cells that results in enhanced mTORC1 activity. However, this study was limited to 24 hpi which is quite early in the context of an ongoing infection. Our study also indicated an early, albeit modest, activation of mTORC1. However, the inhibition accompanied by the loss of substrates at later time points was very consistent and strong in more than one cell line. In the context of altering metabolic activities during infection, it appears that the metabolic networks are manipulated differently during the distinct phase of infection and this may have a significant bearing on the outcome of infection.

eIF4E phosphorylation is dependent on the activities of ERK1/2 and p38MAPK. It is curious to note that only ERK1/2, but not p38MAPK, was dephosphorylated by SARS-CoV-2 mediated signaling activities. Unpublished results from our laboratory have indicated synergistic regulation of Mnk1 by these MAPKs. Curiously, Mnk1 was also targeted at the protein level by the virus and this must have significantly impacted eIF4E phosphorylation. Since eIF4E phosphorylation is understood to affect only a select set of mRNAs translationally (35), we believe that its contribution to the global suppression of translation activities caused by SARS-CoV-2 infection could be limited and more studies are necessary to determine its impact. The consequence of p38MAPK phosphorylation and possible activation of this molecule in SARS-CoV-2 infection is very evident from the inhibition studies. The drop in viral titer was modest, but proportionate to the magnitude of inhibition. Whether this has any impact on the translation of viral proteins is to be determined.

It is intriguing why SARS-CoV-2 infection does not induce eIF2 α phosphorylation. eIF2 α is phosphorylated by one of its four kinases most of which are activated upon various stress exerted on the cell. RNA viruses often impart intense stress on ER that is relayed to PERK (36, 37). PKR, one of the dsRNA sensors is often activated by RNA viral infections. These observations indicate that SARS-CoV-2 depends on the canonical mechanism of translation initiation that requires the availability of active ternary complexes, which eIF2 α is a part of. Since eIF2 α phosphorylation results in the inhibition of new initiation events that would adversely affect the translation of viral transcripts as well, SARS-CoV-2 might have evolved strategies to bypass this modification.

Materials and Methods

Antibodies and inhibitors

All primary antibodies were purchased from Cell Signaling Technologies except the anti-SARS Spike antibody (Novus Biologicals; NB100-56578) and anti-SARS-CoV-2 Nucleocapsid (Thermo Fisher; MA5-29982). HRP-conjugated anti-rabbit and anti-mouse secondary antibodies were purchased from Jackson ImmunoResearch. Torin1 and MHY1485 were from Sigma, whereas the p38 VIII inhibitor was from Cayman Chemicals.

Cell culture

Vero (CCL-81) African green monkey kidney epithelial cells, Huh7 human hepatoma cells and Calu3 lung adenocarcinoma cells were cultured in Dulbecco's Modified

Eagle's Medium (DMEM; from Gibco) with 10% Fetal Bovine Serum (FBS; Hyclone) and 1× penicillin-streptomycin cocktail (Gibco) at 37°C and 5% CO₂. Colorectal adenocarcinoma Caco2 cells, were grown in DMEM supplemented with 20% FBS and 1× antibiotic. Cells were continuously passaged at 70-80% confluency and mycoplasma contamination was monitored periodically.

SARS-CoV-2 Infection and quantification

Two Indian isolates of SARS-CoV-2 strains were used in this study (GSSAID id: EPI_ISL_458075 and EPI_ISL_458046) (38, 39). All the viral cultures were propagated in Vero (CCL-81) cells in serum and antibiotics free conditions. Caco2, Huh7 or Calu-3 cells were infected at 1 MOI for 2 hours in serum-free conditions after which the media was replaced with complete media and further incubated until the time of harvesting. At the time of harvesting, the cells were first trypsinized and collected separately for protein and RNA study. The intracellular and extracellular RNA from cells was isolated using respective kits (MACHEREY-NAGEL GmbH & Co. KG) and the SARS-CoV-2 RNA was quantified using a commercial kit (LabGun™ COVID-19 RT-PCR Kit) following manufacturers protocol in Roche LightCycler 480. For intracellular SARS-CoV-2 RNA, the normalization was performed against GAPDH after preparing cDNA in two-step reactions (Primescript, Takara Bio). The infectious viral particle numbers in the supernatant were quantified using plaque-forming unit (PFU/mL) assay. Briefly, the supernatant was log diluted (10^{-1} - 10^{-7}) in 1× serum-free DMEM and used for infecting Vero monolayer grown in six- or twelve-well plates. 2 hpi, the cells briefly washed and were overlaid with agarose: DMEM mix (in 1:1 ratio; 2 x DMEM with 5% FBS and 1% penicillin-streptomycin mixed with equal volumes of 2% LMA), after which the plates

were incubated undisturbed for 6 days at 37°C. Later, the cells were fixed with 4% formaldehyde and stained with crystal violet. The clear zones were counted and PFU was calculated as PFU/mL.

Inhibitions and infection

Torin1 inhibition and MHY1485 activation were done in Huh7 cells. For the Torin1 inhibition experiment, 0.45×10^6 cells were seeded in a six-well format and 24 hours later the cells were infected with SARS-CoV-2, at 1 MOI for 2 hours in serum-free media. Later, the infection media was replaced with serum sufficient media containing 750 nM Torin1 or DMSO, and incubated for 22 hrs. At the end of the treatment, the cells were harvested and protein and RNA were prepared. For the activation of mTORC1, cells were treated with MHY1485 at 24 hpi at 10 μ M concentration and harvested at 48 hpi. The p38 inhibition was carried out in Caco2 cells similar to the MHY1485 experiment. The intracellular and extracellular RNA were subjected to qRT-PCR, and the protein lysates were subjected to western blotting for confirmation of inhibition or activation.

Polysome preparation

Polysomes were fractionated as explained elsewhere (40). Caco2 cells were grown in 175cm² flasks till 70% confluency and subsequently infected with SARS-CoV-2 at 1 MOI. Media was changed after 2 hours, and cells were harvested at 24, 48 72, and 96 hpi, along with mock-infected cells grown alongside for each time point.

The cells were incubated for 5-10 minutes, harvested and washed twice with a solution of ice-cold 1xPBS containing 100 μ g/mL cycloheximide, to freeze the polysomes on the

mRNAs. They were subsequently lysed in polysome lysis buffer containing 20 mM Tris-Cl pH 8.0, 140 mM KCl, 1.5 mM MgCl₂, 0.5 mM DTT, 1% Triton X-100, 1× protease inhibitor, 0.5 mg/L heparin, 100 µg/mL cycloheximide, and RNase inhibitor. Crude RNA was quantified using a spectrophotometer, and 90 µg was layered onto 11 mL of 10-50% linear sucrose gradient (20 mM Tris-Cl pH 8.0, 140 mM KCl, 1.5 mM MgCl₂, 0.5 mM DTT, 100 µg/mL cycloheximide, 1mM PMSF, 10-50% sucrose). The resulting gradients were centrifuged in an SW41 Ti rotor (Beckman Coulter) at 35,000 r.p.m. at 4°C for 3.5 hours. The polysome samples were fractionated using Teledyne ISCO fraction collector system and absorbance measured and graphically noted at 254 nm. Polysome profiles of mock and infected cells for each time point were digitized and overlaid on Inkscape.

Immunoblotting

Protein pellets were lysed in 1 × Nonidet P-40 lysis buffer (1% Nonidet P-40, 50 mM Tris HCl, 150 mM NaCl (pH 7.5), EGTA, 1 mM sodium orthovanadate, 10 mM sodium pyrophosphate, 100 mM NaF, and 1 mM PMSF) incubated on ice for 20 minutes with intermittent vortexing and centrifuged at 13000 rpm for 15 minutes at 4°C. The supernatants containing the proteins were collected and quantified using BCA reagents (G Biosciences). Lysates were mixed with 6× denaturing dye and the proteins were resolved using SDS-PAGE and transferred to PVDF membranes. The membranes were blocked in 5% BSA dissolved in 1× TBST before the addition of primary antibodies. Primary antibodies against the proteins of interest were diluted in the blocking buffer, added to the membrane and incubated overnight at 4°C. Later, the membranes were washed in 1× TBST, secondary antibodies conjugated with HRP were

added and the blots were developed on a Bio-Rad Chemidoc MP system using SuperSignal West Pico PLUS (Thermo Fisher) and SuperSignal West Femto Maximum Sensitivity (Thermo Fisher) chemiluminescent substrate kits.

Statistical analysis

For each experiment, at least three independent replicates were used to calculate mean $\pm$ SEM, and plotted graphically wherever indicated. Statistical significance was measured using two-tailed, unpaired Student *t*-test and the resultant *p* values were represented as *, **, *** indicating *p* values $\leq$ 0.05, 0.005, and 0.0005, respectively.

Institutional biosafety

Institutional biosafety clearance was obtained by K.H.H., for the experiments pertaining to SARS-CoV-2.

Acknowledgements

We thank Mohan Singh Moodu and Amit Kumar for assisting with logistics and Karthika S Nair for her assistance with some experiments

Funding

This study was supported by internal funding from Council of Scientific and Industrial Research (CSIR), Govt. of India. D.G., and D.K received research fellowship from

CSIR. V.S received research fellowship from the Department of Biotechnology (DBT), Govt. of India.

Contributions

The experiments were conceived by H.P., D.G., and K.H.H. H.P., and D.G., performed polysome profiling. D.G., D.K., and V.S prepared SARS-CoV-2, performed infections, quantified them and analyzed data. H.P., A.P.S., and D.K performed immunoblotting. H.P., and D.G., assisted K.H.H in writing the manuscript.

References

1. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL. 2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579:270-273.
2. Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, Li F. 2020. Cell entry mechanisms of SARS-CoV-2. *Proceedings of the National Academy of Sciences* 117:11727-11734.
3. V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. 2021. Coronavirus biology and replication: implications for SARS-CoV-2. *Nature Reviews Microbiology* 19:155-170.
4. Walsh D, Mohr I. 2011. Viral subversion of the host protein synthesis machinery. *Nature Reviews Microbiology* 9:860-875.
5. Finkel Y, Mizrahi O, Nachshon A, Weingarten-Gabbay S, Morgenstern D, Yahalom-Ronen Y, Tamir H, Achdout H, Stein D, Israeli O, Beth-Din A, Melamed S, Weiss S, Israely T, Paran N, Schwartz M, Stern-Ginossar N. 2021. The coding capacity of SARS-CoV-2. *Nature* 589:125-130.
6. Finkel Y, Gluck A, Winkler R, Nachshon A, Mizrahi O, Lubelsky Y, Zuckerman B, Slobodin B, Yahalom-Ronen Y, Tamir H, Ulitsky I, Israely T, Paran N, Schwartz M, Stern-Ginossar N. 2020. SARS-CoV-2 utilizes a multipronged strategy to suppress host protein synthesis doi:10.1101/2020.11.25.398578. *bioRxiv*.
7. Etchison D, Milburn SC, Edery I, Sonenberg N, Hershey JW. 1982. Inhibition of HeLa cell protein synthesis following poliovirus infection correlates with the proteolysis of a 220,000-dalton polypeptide associated with eucaryotic initiation factor 3 and a cap binding protein complex. *J Biol Chem* 257:14806-10.
8. Jan E, Mohr I, Walsh D. 2016. A Cap-to-Tail Guide to mRNA Translation Strategies in Virus-Infected Cells. *Annu Rev Virol* 3:283-307.
9. Edgil D, Polacek C, Harris E. 2006. Dengue Virus Utilizes a Novel Strategy for Translation Initiation When Cap-Dependent Translation Is Inhibited. *Journal of Virology* 80:2976-2986.
10. Kamitani W, Huang C, Narayanan K, Lokugamage KG, Makino S. 2009. A two-pronged strategy to suppress host protein synthesis by SARS coronavirus Nsp1 protein. *Nature Structural & Molecular Biology* 16:1134-1140.

11. Nakagawa K, Lokugamage KG, Makino S. 2016. Viral and Cellular mRNA Translation in Coronavirus-Infected Cells. *Adv Virus Res* 96:165-192.
12. Banerjee AK, Blanco MR, Bruce EA, Honson DD, Chen LM, Chow A, Bhat P, Ollikainen N, Quinodoz SA, Loney C, Thai J, Miller ZD, Lin AE, Schmidt MM, Stewart DG, Goldfarb D, De Lorenzo G, Rihn SJ, Voorhees RM, Botten JW, Majumdar D, Guttman M. 2020. SARS-CoV-2 Disrupts Splicing, Translation, and Protein Trafficking to Suppress Host Defenses. *Cell* 183:1325-1339.e21.
13. Stern-Ginossar N, Thompson SR, Mathews MB, Mohr I. 2019. Translational Control in Virus-Infected Cells. *Cold Spring Harb Perspect Biol* 11.
14. Lapointe CP, Grosely R, Johnson AG, Wang J, Fernández IS, Puglisi JD. 2021. Dynamic competition between SARS-CoV-2 NSP1 and mRNA on the human ribosome inhibits translation initiation. *Proc Natl Acad Sci U S A* 118.
15. Schubert K, Karousis ED, Jomaa A, Scaiola A, Echeverria B, Gurzeler LA, Leibundgut M, Thiel V, Mühlemann O, Ban N. 2020. SARS-CoV-2 Nsp1 binds the ribosomal mRNA channel to inhibit translation. *Nat Struct Mol Biol* 27:959-966.
16. Shi M, Wang L, Fontana P, Vora S, Zhang Y, Fu TM, Lieberman J, Wu H. 2020. SARS-CoV-2 Nsp1 suppresses host but not viral translation through a bipartite mechanism. *bioRxiv* doi:10.1101/2020.09.18.302901.
17. Thoms M, Buschauer R, Ameisemeier M, Koepke L, Denk T, Hirschenberger M, Kratzat H, Hayn M, Mackens-Kiani T, Cheng J, Straub JH, Stürzel CM, Fröhlich T, Berninghausen O, Becker T, Kirchhoff F, Sparrer KMJ, Beckmann R. 2020. Structural basis for translational shutdown and immune evasion by the Nsp1 protein of SARS-CoV-2. *Science* 369:1249-1255.
18. Puray-Chavez M, Tenneti K, Vuong HR, Lee N, Liu Y, Horani A, Huang T, Case JB, Yang W, Diamond MS, Brody SL, Dougherty J, Kutluay SB. 2020. The translational landscape of SARS-CoV-2 and infected cells. *bioRxiv* doi:10.1101/2020.11.03.367516.
19. Jefferies HB, Reinhard C, Kozma SC, Thomas G. 1994. Rapamycin selectively represses translation of the "polypyrimidine tract" mRNA family. *Proc Natl Acad Sci U S A* 91:4441-5.
20. Thoreen CC, Chantranupong L, Keys HR, Wang T, Gray NS, Sabatini DM. 2012. A unifying model for mTORC1-mediated regulation of mRNA translation. *Nature* 485:109-113.
21. Hsieh AC, Liu Y, Edlind MP, Ingolia NT, Janes MR, Sher A, Shi EY, Stumpf CR, Christensen C, Bonham MJ, Wang S, Ren P, Martin M, Jessen K, Feldman ME, Weissman JS, Shokat KM, Rommel C, Ruggero D. 2012. The translational landscape of mTOR signalling steers cancer initiation and metastasis. *Nature* 485:55-61.
22. Hay N, Sonenberg N. 2004. Upstream and downstream of mTOR. *Genes Dev* 18:1926-45.
23. Ma XM, Blenis J. 2009. Molecular mechanisms of mTOR-mediated translational control. *Nature Reviews Molecular Cell Biology* 10:307-318.
24. Hara K, Yonezawa K, Weng Q-P, Kozlowski MT, Belham C, Avruch J. 1998. Amino Acid Sufficiency and mTOR Regulate p70 S6 Kinase and eIF-4E BP1 through a Common Effector Mechanism*. *Journal of Biological Chemistry* 273:14484-14494.
25. Zoncu R, Efeyan A, Sabatini DM. 2011. mTOR: from growth signal integration to cancer, diabetes and ageing. *Nature Reviews Molecular Cell Biology* 12:21-35.
26. Pyronnet S, Imataka H, Gingras AC, Fukunaga R, Hunter T, Sonenberg N. 1999. Human eukaryotic translation initiation factor 4G (eIF4G) recruits mnk1 to phosphorylate eIF4E. *Embo j* 18:270-9.
27. Joshi B, Cai AL, Keiper BD, Minich WB, Mendez R, Beach CM, Stepinski J, Stolarski R, Darzynkiewicz E, Rhoads RE. 1995. Phosphorylation of eukaryotic protein synthesis initiation factor 4E at Ser-209. *J Biol Chem* 270:14597-603.
28. Scheper GC, Proud CG. 2002. Does phosphorylation of the cap-binding protein eIF4E play a role in translation initiation? *Eur J Biochem* 269:5350-9.

29. Lazaris-Karatzas A, Montine KS, Sonenberg N. 1990. Malignant transformation by a eukaryotic initiation factor subunit that binds to mRNA 5' cap. *Nature* 345:544-7.
30. Krishnamoorthy T, Pavitt GD, Zhang F, Dever TE, Hinnebusch AG. 2001. Tight binding of the phosphorylated alpha subunit of initiation factor 2 (eIF2alpha) to the regulatory subunits of guanine nucleotide exchange factor eIF2B is required for inhibition of translation initiation. *Mol Cell Biol* 21:5018-30.
31. Rojas M, Arias CF, López S. 2010. Protein kinase R is responsible for the phosphorylation of eIF2alpha in rotavirus infection. *J Virol* 84:10457-66.
32. Bechill J, Chen Z, Brewer JW, Baker SC. 2008. Coronavirus infection modulates the unfolded protein response and mediates sustained translational repression. *J Virol* 82:4492-501.
33. Pham AM, Santa Maria FG, Lahiri T, Friedman E, Marie IJ, Levy DE. 2016. PKR Transduces MDA5-Dependent Signals for Type I IFN Induction. *PLoS Pathog* 12:e1005489.
34. Mullen PJ, Garcia G, Purkayastha A, Matulionis N, Schmid EW, Momcilovic M, Sen C, Langerman J, Ramaiah A, Shackelford DB, Damoiseaux R, French SW, Plath K, Gomperts BN, Arumugaswami V, Christofk HR. 2021. SARS-CoV-2 infection rewires host cell metabolism and is potentially susceptible to mTORC1 inhibition. *Nature Communications* 12:1876.
35. Furic L, Rong L, Larsson O, Koumakpayi IH, Yoshida K, Brueschke A, Petroulakis E, Robichaud N, Pollak M, Gaboury LA, Pandolfi PP, Saad F, Sonenberg N. 2010. eIF4E phosphorylation promotes tumorigenesis and is associated with prostate cancer progression. *Proceedings of the National Academy of Sciences* 107:14134-14139.
36. Pavo N, Romano PR, Graczyk TM, Feinstone SM, Taylor DR. 2003. Protein synthesis and endoplasmic reticulum stress can be modulated by the hepatitis C virus envelope protein E2 through the eukaryotic initiation factor 2alpha kinase PERK. *J Virol* 77:3578-85.
37. Su HL, Liao CL, Lin YL. 2002. Japanese encephalitis virus infection initiates endoplasmic reticulum stress and an unfolded protein response. *J Virol* 76:4162-71.
38. Gupta D, Parthasarathy H, Sah V, Tandel D, Vedagiri D, Reddy S, Harshan KH. 2021. Inactivation of SARS-CoV-2 by β -propiolactone Causes Aggregation of Viral Particles and Loss of Antigenic Potential. *bioRxiv* doi:10.1101/2021.04.22.441045:2021.04.22.441045.
39. Gupta D, Ahmed F, Tandel D, Parthasarathy H, Vedagiri D, Sah V, Krishna Mohan B, Daga S, Khan RA, Kondiparthi C, Savari P, Jain S, Daga J, Reddy S, Khan N, Harshan KH. 2021. Development of Equine Immunoglobulin Fragment F(ab')₂ with High Neutralizing Capability against SARS-CoV-2. *bioRxiv* doi:10.1101/2021.03.09.434030:2021.03.09.434030.
40. George A, Panda S, Kudmulwar D, Chhatbar SP, Nayak SC, Krishnan HH. 2012. Hepatitis C virus NS5A binds to the mRNA cap-binding eukaryotic translation initiation 4F (eIF4F) complex and up-regulates host translation initiation machinery through eIF4E-binding protein 1 inactivation. *J Biol Chem* 287:5042-58.

Figure 1

Figure 1. Polysome profiles of SARS-CoV-2 infected cells demonstrate severe collapse of polysomes. (A) Immunoblot analysis of mock and SARS-CoV-2 infected Caco2 cells over 24, 48, 72, and 96 hours of infection with SARS-CoV-2. Cell lysates were electrophoresed by SDS-PAGE and probed for viral proteins, Nucleocapsid and Spike. (B) Relative fold change in SARS-CoV-2 E gene, quantified through qRT-PCR, across the four time points. Graph represents data from 3 sets and is plotted as mean ± SEM. *p*-values were calculated using Student's *t*-test and represented as * and **, indicating *p*-values ≤ 0.05 and 0.005, respectively. (C-F) Polysome profiles of Caco2 cells infected with SARS-CoV-2 for 24-, 48-, 72- and 96 hours. The cells were treated with 100 µg/mL CHX before harvesting and lysed in polysome lysis buffer. Equal quantities of lysates were layered onto continuous sucrose gradients ranging from 10-50%, subjected to ultracentrifugation, and fractionated along with measuring absorbance at 254 nm. The digitized profiles for infected and uninfected samples for each time point were overlaid to assess any change in global translation levels.

Figure 2

Figure 2. The suppression of translation activities in SARS-CoV-2 infection does not involve eIF2α phosphorylation. Immunoblot analysis of mock and SARS-CoV-2 infected Caco2 cells assessing kinetics of phosphorylation of eIF2α at S52 and its expression, each normalized against GAPDH. Intensities of phosphorylated and total proteins were separately normalized against GAPDH and the values are represented above the corresponding panel. The phosphorylated residues are marked against the respective panel.

Figure 3

Figure 3. ERK1/2-Mnk1-eIF4E axis is inhibited during SARS-CoV-2 infection.

Representative immunoblot showing phosphorylation kinetics of ERK1/2, Mnk1 and eIF4E in Caco2 cells, normalised against GAPDH, of the panel displayed. Intensities of phosphorylated and total proteins were separately normalized against GAPDH and the values are represented above the corresponding panel. The phosphorylated residues are marked against the respective panel.

Figure 4

Figure 4. Phosphorylated p38MAPK during SARS-CoV-2 infection is beneficial to the viral replication. (A) Immunoblots showing phosphorylation status of p38 at T180/Y182 position along the course of SARS-CoV-2 infection at 24-, 48-, 72-, and 96 hours in Caco2 cells, along with densitometry data of phosphorylation and expression. As in the previous sections, densitometric intensities of phosphorylated and the total proteins were separately normalized against the loading control. (B) SARS-CoV-2 infected cells were treated with p38 VIII inhibitor (p38i) at 10 μM concentration for 24 hours beginning at 24 hpi until harvesting. The inhibition was scored by dephosphorylation status of eIF4E and viral protein abundance under p38 inhibited environment was also studied. (C) Relative fold change in SARS-CoV-2 intracellular RdRp and E gene RNA, in DMSO and p38-inhibited cells. (D) Infectious virion measure in DMSO and p38-inhibited supernatants, quantified as PFU/mL. Graphs represent data from at least three sets and are plotted as mean ± SEM. *p*-values are represented as ** and ***, indicating *p*-values ≤ 0.005 and 0.0005, respectively.

Figure 5

Figure 5. SARS-CoV-2 inhibits mTORC1 and depletes its key substrates. (A) Immunoblots representing status of mTORC1 kinase activity in mock and infected Caco2 cells through its different substrates- 4EBP1, ULK1, and p70 S6 kinase. The black arrows indicate the appropriate bands corresponding to p70 S6K. Densitometry was performed to measure the intensities of the bands and phosphorylated and total

protein intensities were separately normalized against the loading control. (B and C) Relative RNA abundance of 4EBP1 (B) and ULK1 (C) in SARS-CoV-2 infected Huh7 cells, respectively. After the infections, total RNA was prepared from the cells and converted to cDNA after which the specific transcripts were quantified by qPCR. (D) Immunoblot analysis of ribosomal proteins S3, L13a, and L26, over the course of infection in Huh7 cells. Densitometric values of each band was normalized against the loading control and the corresponding values are given above the panels.

Figure 6

Figure 6. mTORC1 restricts SARS CoV-2 replication. (A) Mock and SARS-CoV-2 infected Huh7 cells were treated with either DMSO or 10 μ M MHY1485 at 24 hpi and harvested at 48 hpi, and successful activation was assessed by increased phosphorylation of 4EBP1 in mock cells treated with MHY1485. Viral protein translation under the conditions of infection was measured through Nucleocapsid levels. (B) Relative fold change in SARS-CoV-2 intracellular RNA in infected cells treated with DMSO or 10 μ M MHY1485. Viral genes E and RdRp were measured from the RNA samples isolated from the infected or the mock control cells and the relative fold change was represented graphically. (C) Relative measure of infectivity of the supernatant represented as PFU/mL from SARS-CoV-2 infected Huh7 cells treated with DMSO or

10 μ M MHY1485. (D) Mock and SARS-CoV-2 infected Huh7 cells were infected with SARS-CoV-2 for two hours, treated with either DMSO or 750 nM Torin1 upto 24 hpi and harvested. Similar set up in mock cells was used. Inhibition was assessed by drop in 4EBP1 phosphorylation in Torin1 treated mock cells. Viral translation was indicated by Nucleocapsid. (E) Relative abundance of intracellular viral RNA (RdRp and E) from CoV-2 infected Huh7 cells treated with either DMSO or 750 nM Torin1. Graphs are plotted as mean $\pm$ SEM and *p*-values are represented as * indicating *p*-values $\leq$ 0.05.

Supplementary figure S1

Figure S1: (A-D) Polysome profiles of Caco2 cells infected with another strain of SARS-CoV-2 (hCoV-19/India/TG-CCMB-L1021/2020) and processed as mentioned previously in Figure 1. Infected cells were collected at 24-, 48-, 72-, and 96 hpi and were analyzed. Panels A-D represent polysome profiles from these times points in the order of increase time intervals.

Supplementary figure S2

Figure S2: SARS-CoV-2 infection does not cause eIF2α phosphorylation during the suppression of translation activities: (A, C) Representative immunoblots showing phosphorylation and expression of eIF2α in Calu3 (A), and Huh7 (C) cells, respectively, over 12, 24, 48 and 72 hours. Each panel has densitometric values of fold change in phosphorylation of eIF2α and expression, individually normalized to GAPDH, of the panel displayed. (B, D) qRT-PCR data depicting relative fold change in SARS-CoV-2 E gene, along SARS-CoV-2 infection in Calu3 (B), and Huh7 (C) cells. Graphs represent data from 3 sets and are plotted as mean ± SEM. *p*-values are represented as * and **, indicating *p*-values ≤ 0.05 and 0.005, respectively.

Supplementary figure S3

Figure S3: ERK1/2-Mnk1/2-eIF4E axis is inhibited during SARS-CoV-2 infection. Representative immunoblot of phosphorylation levels of ERK1/2, Mnk1 and eIF4E in Huh7 cells, with densitometric analysis depicting fold change in phosphorylation of ERK1/2 and eIF4E, and fold change in their expression.

Supplementary figure S4

Figure S4: SARS-CoV-2 inhibits mTORC1 and depletes its key substrates. (A) Immunoblots of mTOR substrates, 4EBP1 and ULK1, from mock and infected Huh7 cells showing their phosphorylation as well as expression as indicated in the densitometric values above each blot.