Hybridization underlies localized trait evolution in cavefish

Moran, Rachel L.,¹ Jaggard, James B.^{2,3}, Roback, Emma Y.¹, Rohner, Nicolas^{4,5}, Kowalko, Johanna E.⁶, Ornelas-García, C. Patricia⁷ McGaugh, Suzanne E.^{1*}, Keene, Alex C.^{2*}

1. Department of Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul, MN
2. Department of Biological Sciences, Florida Atlantic University, Jupiter FL
3. Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA
4. Stowers Institute for Medical Research, Kansas City, MO
5. Department of Molecular & Integrative Physiology, KU Medical Center, Kansas City, KS
6. Harriet L. Wilkes Honors College, Florida Atlantic University, Jupiter FL
7. Departamento de Zoología, Instituto de Biología, Universidad Autónoma de México, CP 04510, Mexico City, Mexico

Summary

Compared to selection on new mutations and standing genetic variation, the role of gene flow in generating adaptive genetic variation has been subject to much debate. Theory predicts that gene flow constrains adaptive evolution via natural selection by homogenizing allele frequencies among populations and introducing migrant alleles that may be locally maladaptive¹. However, recent work has revealed that populations can diverge even when high levels of gene flow are present²⁻⁴ and that gene flow may play an underappreciated role in facilitating local adaptation by increasing the amount of genetic variation present for selection to act upon⁵⁻⁸. Here, we investigate how genetic variation introduced by gene flow contributes to adaptive evolution of complex traits using an emerging eco-evolutionary model system, the Mexican tetra (*Astyanax mexicanus*). The ancestral surface form of the Mexican tetra has repeatedly invaded and adapted to cave environments. The Chica cave is unique in that it contains several pool microenvironments inhabited by putative hybrids between surface and cave populations⁹, providing an opportunity to investigate the dynamics of complex trait evolution and gene flow on a local scale. Here we conduct high-resolution genomic mapping and analysis of eye morphology and pigmentation in fish from multiple pools within Chica cave. We demonstrate that hybridization between cave and surface populations contributes to highly localized variation in behavioral and morphological traits. Analysis of sleep and locomotor behaviors between individual pools within this cave revealed reduced sleep associated with an increase in ancestry derived from cave populations, suggesting pool-specific ecological differences may drive the highly-localized evolution of sleep and locomotor behaviors. Lastly, our analyses uncovered a compelling example of convergent evolution in a core circadian clock gene in multiple independent cavefish lineages and burrowing mammals, indicating a shared genetic mechanism underlying circadian disruption in subterranean vertebrates. Together, our results provide insight into the evolutionary mechanisms that promote adaptive genetic variation and the genetic basis of complex behavioral phenotypes involved in local adaptation.

Main Text

A rapidly growing body of research demonstrates that gene flow, both among populations within species and between different species, is more common than previously thought and can play a key role in the evolutionary process by impeding or promoting adaptive divergence^{5,6,10–12}. Hybrid zones resulting from interbreeding between lineages that occupy different environmental extremes offer a powerful means to detect targets of selection in the genome underlying complex, locally adapted traits. Natural variation present in recombinant hybrids can be leveraged through admixture mapping^{13,14}, which identifies associations between genetic ancestry and trait variation in admixed populations formed by interbreeding between two or more diverged lineages. This association-based approach was first developed to uncover the genetic basis of diseases in humans following the observation that the frequencies of some disease-causing variants differ substantially among populations^{15,16}. Recent advances in sequencing technology and statistical approaches have made it feasible to apply admixture mapping to identify adaptive loci underlying ecological divergence in plant and animal models of evolution^{17–24}. However, previous studies in plants and animals have focused on hybrids formed between distinct species with substantial genetic divergence and reproductive isolation, making it difficult to identify regions associated with ecologically relevant traits versus intrinsic incompatibilities²⁵. The application of admixture mapping to models of trait evolution has the potential to define fundamental interactions between genetic and environmental variation that shape evolution. Studies that apply whole genome sequencing to hybrids formed between interbreeding lineages in the earliest stages of divergence are likely to provide the most insight into the genetic basis of evolutionary change^{24,26}, but are currently lacking.

The Mexican tetra, *Astyanax mexicanus*, is a powerful model system for investigating the genetic and evolutionary basis of trait development and behavior^{27–31}. Surface populations inhabit rivers

from Texas to Mexico and have invaded caves multiple times, resulting in at least 30 populations of cave-morphs in the Sierra de El Abra region of Northeast Mexico^{9,32}. At least two independent lineages of surface fish, commonly referred to in the literature as “old” and “new” lineages, have invaded caves within the past roughly 200,000 years^{33–37}. Cavefish populations have converged on numerous morphological traits that are thought to be adaptive in the cave environment, including albinism and eye loss³⁸. In addition, cavefish have repeatedly evolved multiple behavioral changes, including sleep loss, which may increase time allocated to foraging in nutrient-poor cave environments^{29,39}. Recently, the application of molecular genetic approaches has led to the identification of genetic factors that regulate some of these trait differences, but the evolutionary mechanisms underlying these genetic differences remain poorly understood.

Cave and surface populations are interfertile under laboratory conditions, and we recently identified a surprising amount of historical and contemporary gene flow between surface and cave populations³⁶. The presence of admixture between populations raises the possibility that gene flow is a critical driver of trait evolution³⁶. Unlike most caves, the Chica cave contains fish that appear to exhibit high levels of phenotypic variation, and fish are present across four pools that differ in proximity to the cave entrance, nutrient input, and physicochemical properties (e.g., dissolved oxygen) (Fig. 1A)^{9,40–42}. Historical surveys suggested that fish exhibit a morphological gradation in troglotic traits across pools, potentially shaped by environmental variation within the cave and ongoing influx of surface and cave morphs from underground waterways that feed into the cave⁹. Thus, this cave provides a natural system to study the effects of hybridization on trait evolution across a variable environment. Quantification of trait variation and formal tests for hybridization have yet to be conducted. Here we leverage robust differences in behavior and morphology between surface and cavefish populations of Mexican tetras, combined with whole genome sequencing, to investigate the ancestry of putative hybrids in Chica cave and to examine the genetic basis of trait variability across a heterogeneous environment.

Figure 1. (A) Map of Chica cave modified with permission from ⁴³. (B) Collection locations for cave and surface populations. For the two surface populations, the collection location for Río Choy is represented by a light blue circle and the collection location for Rascón is represented by a dark blue circle. (C) Representative images of wild-caught fish. Scale bar denotes 1 cm. (D) Representative images of eye

morphology variations in Chica pools 1 and 2 and complete eye loss in wild-caught Pachón and Los Sabinos cave populations is denoted with “n.d.” for “no data” since there are no eyed fish present in these two populations. (E) Eye diameter is reduced in Chica pool 2 fish compared to pool 1 (* $p < 0.05$, Unpaired t-test, $t=1.88$, $df=17$). Eye size was corrected to body length. (F) Eye morphology in Chica fish fits within expected outcome. Chica 1: observed 60% eye 40% no eye; expected 55% eye, 45% no eye, $P>0.45$ Binomial test. Chica 2: observed 40% eye, 60% no eye; expected 24% eye, 76% no eye, $P>0.34$ Binomial test. (D) Pigment quantification showing differences in melanin pigmentation in different populations ($p < 0.001$, KW statistic=18.04, Kruskal-Wallis test with Dunn’s multiple comparison test: Chica vs Pachón, $p < 0.001$; Chica vs Los Sabinos, $p=0.88$; Pachón vs Los Sabinos, $p < 0.001$). Pigmentation are more variable among different cave populations, Brown-Forsythe test, $P=0.03$; Bartlett’s test, $P=0.04$.

We first conducted morphological and population genomic analysis to verify whether Chica fish represent hybrids between surface and cave populations. We also asked whether variation in morphological traits and allele frequencies are present between pool microenvironments within Chica cave. We collected adult fish from two adjacent pools in Chica cave that are partially hydrologically separate, referred to as Pool 1 and Pool 2 (Fig. 1A). We also collected adult fish from two non-admixed caves in the Sierra de El Abra regions, Pachón and Los Sabinos (Fig 1B,C). We scored wild-caught fish for two morphological traits, eye size and pigmentation, which previously have been qualitatively described as showing high variation within Chica cave compared to other cave populations. In agreement with laboratory stock populations, eyes were absent in wild-caught fish from Pachón and Los Sabinos caves (Figure 1D, Extended Data Figure 1). In contrast, the presence or absence of eyes was highly variable in wild-caught fish from both pools within Chica cave. Overall eye diameter was significantly larger in Chica Pool 1 fish compared to Pool 2 individuals ($p < 0.05$, Unpaired t-test, $t=1.69$, $df=17$). Additionally, the sample from Chica Pool 1 contained more fish with eyes present (60%) than those with no eyes (40%) while the sample from Chica Pool 2 contained fewer fish with eyes (40%) and increased numbers with no eyes (60%) (Figure 1F). Binomial analysis demonstrated that the observed rate of the eye phenotype fit within expected outcome range (Chica Pool 1, $p = 0.45$; Chica Pool 2, $p = 0.34$).

We observed reduced melanin pigmentation levels in all cavefish, but these reductions vary among different cave populations. Pachón cavefish are considered largely albinic, while Los Sabinos retain vestigial melanocytes that result in a reduced pigmentation pattern compared to surface morphs^{44,45}. Quantification of melanin pigmentation in wild-caught Pachón and Los Sabinos revealed comparable patterns to previous reports in lab-reared fish^{44,45} (Figure 1G). Although a number of pigmented individuals were present within the wild-caught Pachón and Los Sabinos populations, we observed low overall levels in the variability of melanin patterns within these cave populations. Interestingly, robust differences in the number of melnophores were observed between different populations of cavefish (Dunn's multiple comparison, Chica vs. Pachon, $p < 0.01$; Pachon vs Los Sabinos, $p < 0.01$ Fig 1G; Extended Data Figure 1). Interestingly, the standard deviations between populations was significantly different, indicating that Chica cavefish are more variable in melanin morphology patterns (Brown-Forsythe test, $p=0.03$; Bartlett's test, $p=0.04$). Taken together, these findings support the hypothesis that fish from Chica cave are the result of surface-cave hybridization and exhibit a high degree of phenotypic variability that differs between microenvironments within the Chica cave.

To identify the genetic basis of phenotype variability, we used whole-genome resequencing to conduct admixture analyses and genomic ancestry mapping in Chica cavefish to test for evidence of hybridization. We also used whole-genome resequencing to examine population structure between fish from Pools 1 and 2 within Chica cave, three other cave populations (Pachón, Los Sabinos, and Tinaja), and two surface populations (Río Choy and Rascón) (Fig. 2A,B). Our analyses revealed ongoing gene flow between Chica cave, Río Choy, and Tinaja cave populations (Fig. 2C), and confirmed that the Chica cave population represents a hybrid swarm resulting from over 2,000 generations of interbreeding between the nearby surface fish (from Río Choy/Tampaón) and southern El Abra cavefish (Fig. 2D; Extended Data Tables 7-8; Supplementary Information). This analysis indicated low levels of overall genome-wide

divergence between fish from Chica cave Pool 1 and Pool 2, suggesting that gene flow is high among pools within Chica cave (Fig. 2E). Supporting this notion, all Chica individuals exhibited highly similar global ancestry proportions from surface versus cave parental populations (Pool 1 Cave Ancestry: Mean $\pm$ SE = 0.755 ± 0.004 ; Pool 2 Cave Ancestry: Mean $\pm$ SE = 0.756 ± 0.003 ; see Supplementary Information). Furthermore, the length distribution of ancestry tracts derived from the surface parental population did not differ between Chica pools (Extended Data Tables 7-8). Together, this indicates that gene flow from the surface population does not differ significantly between pools.

Despite high levels of gene flow between pools, analysis of sequence divergence revealed several highly localized regions of genomic divergence between fish from adjacent pools within Chica cave, reflecting the morphological differences we observed between these pools (Fig. 2E). In genomic outlier windows where absolute genetic distance (i.e., Dxy) between Chica pools was in the top 5% of values, fish from Pool 2 were more likely to harbor alleles derived from the nearby southern El Abra cave populations (i.e., Tinaja) compared to Pool 1 (Wilcoxon rank sum test: $W = 2.6511e+13$, $p < 2.2e-16$). Specifically, 50.96% (371 out of 728) of the outlier windows with exceptionally high genetic divergence between pools contained a higher proportion of sites derived from cave (i.e., Tinaja) ancestry in Pool 2, whereas 39.56% (288 out of 728) had a higher proportion of sites derived from cave ancestry in Pool 1. The remaining 9.47% (69 out of 728) of outlier windows did not exhibit differences in ancestry between pools, which may be due to genetic differences that have accumulated via drift and/or regions that were not ancestry informative (i.e., the Hidden Markov Model was unable to accurately delimit ancestry blocks). We observed a positive correlation between the difference in local ancestry between pools and genetic distance between pools (Pearson's correlation: $r=0.0012$, $n=7,345,340$, $p=0.0011$), indicating that the greater proportion of cave ancestry maintained in Pool 2 compared to Pool 1 drives genetic

differences between the pools. Together, this demonstrates that gene flow has played a key role in driving genetic variation in this system and may facilitate evolution on a local scale.

Figure 2. (A) ADMIXTURE barplot showing ancestry proportions for K=5. (B) Biplot of scores for the first two PCs. Note

that individuals from Chica cave Pool 1 and Pool 2 overlap and individuals from Tinaja cave and Los Sabinos cave overlap. (C) Treemix tree with three migration events and rooted with the outgroup, *A. aeneus*. New lineage surface population (Río Choy) groups with *A. aeneus*, and old lineage surface (Rascón) and caves (Chica and Tinaja) all group together. Migration events are present between Chica cave and the geographically close surface population, Río Choy, and between Tinaja and Chica caves. (D) Local ancestry derived from surface (Río Choy, blue) versus cave (Tinaja, red) parental populations in hybrid fish from Chica cave. Each row represents a diploid individual with two haplotypes stacked on top of one another. (E) Absolute genetic divergence (Dxy) between fish from Chica cave Pool 1 versus Pool 2 in 50 kb windows across the genome. Locations are indicated for several top candidate genes with high divergence between Chica pools and biological functions related to sleep/circadian cycle (purple), eye size/morphology (green), metabolism (orange), and pigmentation (pink), or that are pleiotropically involved in two or more of these pathways (black) (see Extended Data Table 11). The 95th percentile is delimited by a horizontal line.

Out of all functionally annotated protein-coding genes, those with a high degree of sequence divergence between pools (i.e., genetic distance in the top 5% of all genes) associated well with the characteristic suite of phenotypic differences typically present between non-admixed cave and surface populations (Extended Data Table 9). These genes with exceptional divergence between pools were significantly enriched for ontologies related to traits that differ between non-admixed surface and cave populations, including pigmentation (*bloc1s3*, *cd63*, *psen2*, *sox10*, *tspan36*, *meox1*, *asip2b*, *arl6*, *rab38c*), eye development and light detection (*c1qa*, *itgam*, *rho*, *pde6ha*), sensory processing by the lateral line neuromast (*rsph9*, *shisa2a*), metabolism (*parp9*, *sult6b1*), and sleep and the circadian cycle (*btbd9*, *srd5a1*, *mc3r*, *chrnb5a*, *galn*, *gna11b*, *rorca*, *cry1a*, *cry1b*) (Fisher's exact tests, $p < 0.05$; Supplementary Information, Extended Data Tables 9-10). This set of 26 outlier genes with ontologies related to phenotypic differences observed between non-admixed cave and surface populations provided strong candidates for local adaptation. We used a deep convolutional neural network approach implemented in diploS/HIC⁴⁶ to formally test for signatures of selection on these genes. We found that 21 (81%) of the 26 candidate genes occur within regions of the genome that appear to have experienced selective sweeps in one or both Chica pools (Extended Data Table 11). Furthermore, 20 (77%) of the 26 candidate genes for local adaptation also show extreme divergence (genetic distance in the top

5% of all genes) in one or more comparisons between cave and surface population pairs that do not show evidence of recent admixture (Extended Data Table 9), potentially pointing to general trends in the genetic underpinnings of cave evolution. These genes are strong candidates underlying the morphological differences in eye size and pigment we observed between pools within Chica cave, and suggest that adaptive behavioral (i.e., sleep) differences may also be present between pools. Taken together, our results indicate that hybridization may interact with varying selection pressures between different pool microenvironments within Chica cave to recapitulate phenotypic differences associated with divergent selection between cave and surface environments.

We observed a number of additional factors that further suggest the genetic differences identified in candidate genes for local adaptation in Chica cave could have functional consequences. We used *in silico* prediction with SIFT⁴⁷ and VEP⁴⁸ to identify mutations with deleterious effects that occur at higher frequencies in cavefish and re-analyzed transcriptional data obtained from Tinaja and Río Choy fry at 30 days post fertilization (dpf)⁴⁹. These analyses revealed striking patterns of differential expression between lab-raised fry from surface versus cave populations and coding variants that affect protein function (see Supplementary Information, Extended Data Tables 11-13). Out of the 26 candidate genes with ontologies related to cave adapted phenotypes, 21 were expressed at 30 dpf. Of those 21 genes, 14 (67%) showed significant differential expression between cave and surface fish (Extended Data Table 13). We also identified putatively deleterious coding changes in five of our candidate genes with ontologies associated with sleep and the circadian cycle. One notable mutation is present in the gene cryptochrome circadian regulator 1a (*cry1a*), a transcriptional repressor. Cryptochromes play a highly conserved role in circadian clock regulation across plants and animals⁵⁰. Knockout of *cry1a* results in defects in locomotor activity and behavioral rhythms in zebrafish⁵¹, and mutations in the human *cry1* ortholog are associated with a circadian rhythm sleep disorder (delayed sleep phase syndrome, DSPS)⁵². We observed

that *cry1a* exhibits a nonsynonymous mutation, R263Q, that is present in Chica, Tinaja, and Pachón cave populations but not in Río Choy or Rascón surface populations. To determine whether this mutation is unique to the cavefish lineages, we examined an alignment of 284 CRY1 orthologs across 266 animal species (including invertebrates) downloaded from Ensembl (<https://useast.ensembl.org/>). We also downloaded the CRY1 ortholog for the Somalian cavefish *Phreatichthys andruzzii* that was available on NCBI (Accession: ADL62679.1). Remarkably, we found that the R263Q mutation is present in four distantly related cyprinid species from Somalia (*Phreatichthys andruzzii*)⁵³ and China (the blind barbel *Sinocyclocheilus anshuiensis*, the golden-line barbel *Sinocyclocheilus grahami*, and the horned golden-line barbel *Sinocyclocheilus rhinoceros*), as well as two burrowing rodent species (the naked mole rat, *Heterocephalus glaber*, and the common degu, *Octodon degus*) (Fig. 3). *Phreatichthys* and *Sinocyclocheilus* cyprinid cavefish have convergently evolved troglomorphic traits that are shared by *Astyanax* characin cavefish, including reduction or loss of eyes and pigment, and disrupted circadian cycles^{53–55}. The naked mole rat has also evolved many of the same characteristic traits associated with life in the dark, including reduced eye size and function, a disrupted circadian clock, and loss of sleep⁵⁶. Our *in silico* analyses indicated that the R263Q mutation is predicted to be deleterious to protein function (Extended Data Table 11). This is supported by the observation that this position is otherwise highly conserved across plants and animals and occurs within the FAD binding domain of CRY^{57,58} (Fig. 3A). Our findings provide compelling evidence that the R263Q mutation in the core circadian clock gene *cry1* has convergently evolved up to five times in cavefish and burrowing mammals (Fig. 3B,C), indicating that a common genetic mechanism may contribute to disruption of sleep behavior and circadian rhythm in subterranean vertebrates.

Figure 3. (A) Model of *Astyanax* Pachon cavefish CRY1A protein based on crystal structure of mouse CRY1 (PDB: 6kx7). The model for the *Astyanax* Pachon cavefish protein was generated with SWISS-MODEL and the comic structure was visualized with VMD (version 1.9.4). The location of R263Q (in the α 10 within the FAD binding pocket) is indicated with an arrow. This image was made with VMD/NAMD/BioCoRE/JMV/other software support. VMD/NAMD/BioCoRE/JMV/ is developed with NIH support by the Theoretical and Computational Biophysics group at the Beckman Institute, University of Illinois at Urbana-Champaign. (B) Species tree for 23 animal species, selected to include subterranean lineages and their epigeal relatives (based on the species tree available from Ensembl release 102 and ^{55,59,60}). Branches where the R263Q mutation has evolved are highlighted in green. Illustrations depict *Astyanax* Mexican cavefish (Pachon cavefish top, Tinaja cavefish bottom), degu, and naked mole rat. (C) Section of multiple sequence alignment for CRY1 orthologs spanning sites 187 – 289 in the *Astyanax* CRY1A protein. The arginine to glutamine mutation at *Astyanax* site 263 is indicated with a black box.

Together, the enrichment of genes with ontologies related to sleep and the circadian cycle in our list of candidates with exceptionally high divergence between Chica pools and our observation of large-effect, cave-derived mutations in several of these genes suggests that sleep behavior may differ between fish from different pools within Chica cave. Multiple laboratory-bred cavefish populations exhibit convergence on sleep loss and increased locomotor activity^{29,39}. While these behavioral differences are proposed to enhance foraging opportunity in nutrient-poor cave environments²⁸, sleep has not been assayed in wild caught fish and it is not known whether sleep and activity differ based on local cave environments. To directly test whether the genomic differences we observed in candidate genes for sleep differences within Chica cave associate with functional differences, we analyzed behavioral variation in wild-caught fish from Chica Pool 1 and Pool 2. We also assayed non-admixed cavefish from Los Sabinos and Pachón for comparison. We measured sleep duration and locomotor activity in wild-caught fish under standard laboratory settings. We observed that total sleep in wild-caught Pachón cavefish is significantly reduced compared to Los Sabinos cavefish, similar to what is observed in fish derived from these populations in the laboratory (Fig. 4 A,B) (Dunn's multiple comparison, $p < 0.01$). This provides evidence that the sleep loss observed in lab-reared stocks is replicated in wild-caught fish^{39,61}. The duration of sleep in Chica fish from Pool 1 was significantly greater than sleep in fish from Pool 2 (Dunn's multiple comparison, $p < 0.05$). The increased sleep duration from Chica Pool 1 fish was caused by an increase in number of sleep bouts compared to fish from both Chica Pool 2 and Pachón cave (Fig. 4B) (Dunn's multiple comparison, Chica pool 2, $p < 0.05$, Pachon $p < 0.05$). Sleep bout length was reduced in fish from Chica Pool 2 and Pachón cave populations compared to fish from Chica Pool 1 (Fig 4C), though this comparison was not statistically significant. These differences in sleep cannot be explained by hyperactivity, as the average activity during periods of wakefulness (waking activity) did not differ between any of the populations (Fig. 4E). Therefore, hybrid Chica fish exhibit pool-specific differences in sleep, with fish from Pool 2 largely phenocopying Pachón cavefish and fish from Pool 1 exhibiting a greater

sleep duration, similar to what has been previously observed in laboratory stocks of surface fish^{29,39}. These results reveal the presence of behavioral differences between adjacent pools within Chica cave, with Pool 2 being more cavefish-like than Pool 1. This is in agreement with our genomic analyses, which found more cave ancestry maintained in Pool 2 compared to Pool 1 specifically in genomic regions of high divergence that contained genes related to phenotypes putatively implicated in local adaption in this system.

Figure 4. Sleep variation between and within wild-caught *A. mexicanus* cave populations. (A) Twenty-four hour sleep profiles in Chica Pool 1, Chica Pool 2, Pachón and Los Sabinos fish (B) Total sleep duration is variable among

different populations of wild-caught fish (Kruskal-Wallis test, $P < 0.001$, KW statistic = 17.55). Chica Pool 1 fish sleep significantly longer than Chica Pool 2 fish (Dunn's multiple comparison, $p < 0.05$). Wild-caught Pachón cavefish show reduced sleep compared to Chica Pool 1 (Dunn's multiple comparison, $p < 0.05$). (C) Number of sleep bouts is variable in different cave populations (Kruskal-Wallis test, $P < 0.05$, KW statistic = 10.62. Chica Pool 2 and Pachón fish have reduced sleep bout numbers compared to Chica Pool 1 (Dunn's multiple comparison, Chica Pool 2, $p < 0.05$, Pachón, $p < 0.05$). (D) Sleep bout duration is not altered in any population of cavefish (Kruskal-Wallis test, $P > 0.34$, KW statistic = 3.46). (E) Waking activity is not altered among cave populations (Kruskal-Wallis test, $P > 0.3$, KW statistic = 3.65).

We conclude that genetic admixture across different environments produced substantially different phenotypes in very close geographic proximity. Our findings also imply that stronger selection against surface ancestry in Pool 2 of Chica cave could be maintaining differences in genes important to local cave pool via adaptation, rather than differential gene flow of each Chica pool and the local surface population. These results are remarkable given that the pools are separated by just 10 m. Additionally, we found that genes which are exceptionally divergent between the two cave pools associate well with the phenotypic differences observed. Notably, genes identified as highly divergent across pool microenvironments are also associated with phenotypes involved in nonadmixed surface and cave populations. These findings provide unprecedented insight into the genetic basis of local adaptation across varying environments in hybrid populations. Lastly, we identified a coding variant in the core circadian clock gene *cry1* that has convergently evolved in other distantly related cave-dwelling fish species and burrowing mammals. This suggests that a common genetic mechanism may contribute to disrupted circadian rhythms across multiple subterranean vertebrate lineages that have adapted to live in constant darkness.

A rapidly growing body of work is demonstrating that introgressive hybridization often drives patterns of phenotypic evolution and may play an integral role in the evolutionary processes of local adaptation and speciation¹². A number of studies have shown that behavioral variation can result from introgressive hybridization (e.g., song in hybrid Darwin's finches⁶², mate choice in

hybrid baboons⁶³, defensive behavior in hybrid honey bees²³), providing new substrate for selection to act upon. Hybridization has also been proposed to have influenced behavioral novelty in humans^{64,65}. Here we mapped the genetic basis of complex behavioral differences resulting from the interplay between hybridization and selection in a vertebrate system. This demonstrates that hybridization may play an underappreciated role in shaping behavioral variation and evolution. Therefore, the identification of hybridization-mediated evolution in the *A. mexicanus* that inhabit Chica cave establishes this system as a model to study the genetic basis of evolution in complex behavioral and morphological traits.

Methods

Study system

Evidence suggests that there have been at least two colonization events of northern Mexico by surface dwelling *A. mexicanus*, typically referred to as the “old” and “new” lineage. One lineage of surface fish colonized the caves in the El Abra region and a separate lineage of surface fish subsequently colonized the northern Guatemala region and western Micos region caves of Northeastern Mexico. While we now know that these two lineages and their invasion of the caves were not timed in line with the “old” and “new” designations³⁷, we use this shorthand here since these labels are consistent with past work^{33,34,36,66}. The surface fish within the Rascón/Gallinas river system are most similar to the old lineage cavefish and were likely isolated from colonization by the new lineage surface fish due to a 105 m vertical waterfall³⁷. Cavefish within the El Abra region that descended from old lineage of surface ancestors are now within close geographic proximity to surface fish from the new lineage.

Fish occupy multiple pools within Chica cave that naturally differ in ecology. Whether the Chica cave population came from the old or new lineage stock has been the subject of much debate in the cavefish community. Fish from Chica cave show higher genetic differentiation from the rest of the El Abra cave populations, which some have interpreted as evidence of an independent invasion event³⁴. However, this pattern could also be explained by hybridization with local surface populations⁶⁷. In accordance with this hypothesis, recent phylogenetic analyses have revealed that fish from Chica cave possess new lineage mitochondrial DNA and old lineage nuclear DNA, indicative of historical introgression^{33,35}.

Identifying the genetic underpinnings of behavioral evolution can be especially challenging in natural populations^{68,69}. A genomic signature of local adaptation is most detectable when gene flow is high among populations in different environments^{70,71}, as gene flow homogenizes the background level of divergence between populations while selection maintains differentiation at regions important to local adaptation. High levels of gene flow between the Chica cave and surface population together with strong selection for adaptation to the cave environment are predicted to shape patterns of divergence across the genome and provide insight into the genes important for maintaining cave phenotypes. Therefore, this system provides the unique opportunity to investigate the genetic basis of adaptive traits.

Sequencing and Genotyping

We used whole genome resequencing and population genomic analyses to (1) characterize population structure and genetic relationships between and within the Chica cavefish, three other cavefish populations, and two surface populations (2) identify candidate regions for local adaptation with high levels of genetic differentiation between Chica pools, and (3) test for signatures of introgression between Chica cave and other nearby cave and surface populations. All sequencing used in these analyses originated from wild-caught fish collected from two

adjacent pools within Chica cave (Pool 1, approximately 91 m from the entry, and Pool 2, approximately another 10 m into the cave; Fig. 1A).

Fin clips were collected from adult fish from Chica cave in 2015 and stored in 80% ethanol. We sequenced a total of 19 *A. mexicanus* samples from Chica cave (five from Pool 1 and 14 from Pool 2) using 125 bp paired end reads on an Illumina HiSeq 2500 at the University of Minnesota Genomics Center. Fin clips were collected from adult fish from Los Sabinos cave (n=3) in 2015 and were sequenced using 150 bp paired end reads on an Illumina NovaSeq S4. Genomic libraries for all Chica samples and two of the Los Sabinos samples (Sabinos_T3076_S26 and Sabinos_T3093_S27) were prepared using Illumina TruSeq v3 Nano DNA Sample Prep Kits. The genomic library for the third Los Sabinos sample (Sabinos1) was prepared using a Chromium Genome Library Kit and Gel Bead Kit v2 and a Chromium Genome Chip Kit v2. We obtained whole genome resequencing data for Pachón cavefish samples (n = 10) from a previously published study³⁷.

To investigate recent patterns of introgression between Chica cave and surface fish, we also obtained *A. mexicanus* sequence data from fish from one other cave population in the El Abra region that is not heavily admixed (Tinaja, n = 10), a nearby new lineage surface population (Río Choy, n = 9), and an old lineage surface population (Rascón, n = 8)³⁷. It has been hypothesized that caves within the southern El Abra region exchange migrants through subterranean connections, and Tinaja was previously shown to contain fish with mostly cave-like phenotypes. Thus, Tinaja cavefish sequence can provide a reference to identify cave alleles in the putative hybrid swarm present in Chica cave.

Río Choy contains new lineage surface fish and is a tributary of the Tampaón River, which is believed to be the source of surface fish in Chica cave. Rascón is a tributary of the Gallinas River,

and contains old lineage surface fish³⁷. Thus, including genomic data from Tinaja, Río Choy, and Rascón in our analyses provides a means to test for recent introgression between new lineage surface fish and old lineage cave fish within Chica cave. Previously published data from a closely related congener, *Astyanax aeneus* (n=1)³⁷, was also included to serve as an outgroup in tests for introgression. Pachón, Tinaja, Río Choy, Rascón, and *A. aeneus* samples were all previously sequenced as 100 bp paired end reads on an Illumina HiSeq2000 at The University of Minnesota Genomics Center³⁷. Raw sequencing data for these samples was downloaded from NCBI (SRA Accession Numbers SRP046999, SRR4044502, and SRR4044501).

We conducted genotype calling following the GATK Best Practices⁷²⁻⁷⁴ (Extended Data Table 1). Adapters were trimmed from raw reads using Cutadapt v1.2.1. We trimmed samples for quality using Trimmomatic v0.30 and specified a minimum quality score of 30 across a 6 bp sliding window and discarded reads with a length of <40 nucleotides. Reads were aligned to the surface *Astyanax mexicanus* genome (*Astyanax_mexicanus*-2.0, downloaded from NCBI) using bwa v0.7.4⁷⁵. We used Picard v2.3.0 (<http://broadinstitute.github.io/picard/>) to remove duplicates and add read group information and used samtools v1.7⁷⁶ to split de-duplicated bams into mapped and unmapped reads. Mapped bams were used to generate per-individual gvcfs with the Genome Analysis Tool Kit (GATK) v3.7.0 HaplotypeCaller tool. We used the GenotypeGVCFs tool in GATK v3.8.0 to produce vcf files for each chromosome and unplaced scaffolds that include all individuals (and include invariant sites). The SelectVariants and VariantFiltration tools in GATK v3.8.0 were used to apply hard filters. We subset vcfs for each chromosome and unplaced scaffolds into invariant, SNPs, and mixed/indel sites and applied filters separately following GATK best practices (Supplementary Table 1). We then used the MergeVcfs tool in GATK v4.1.4 to recombine all subset VCFs for each chromosome and unplaced scaffold. Indels and the 3 bp region around each indel were removed using a custom python script. We used the vcftools⁷⁷ --exclude-bed option to remove repetitive regions identified by WindowMasker and RepeatMasker⁷⁸. We

also used vcftools to only retain biallelic SNPs, to remove sites with greater than 20% missing data within each population, and to remove variants with a minor allele frequency <1%. This resulted in retaining a total of 225,462,242 sites throughout the genome, 3,337,738 of which were SNPs.

Population Structure

To quantify the number of distinct genetic clusters (i.e., populations) present among the *A. mexicanus* cave and surface populations, we used ADMIXTURE v1.3.0 and Principal Components Analysis (PCA). For these analyses, we applied a more stringent missing data filter, only retaining sites with <10% missing data. To control for linkage between SNPs that cluster locally on a given chromosome, we thinned SNPs to 1 kb apart and did not include unplaced scaffolds. This resulted in a set of 678,637 SNPs. We ran ADMIXTURE for each value of K from two through nine and estimated the best value of K using the Cross Validation (CV) procedure in ADMIXTURE. The best K was chosen as the value that had the lowest CV error. We used Plink v1.90 to conduct the PCA. For this analysis, we again thinned SNPs to 1 kb apart, but included all placed and unplaced scaffolds. This resulted in a set of 733,979 SNPs.

We calculated absolute genetic divergence (Dxy) and relative genetic divergence (Fst) between populations and nucleotide diversity (Pi) within populations in non-overlapping 50kb windows across the genome using the python script popgenWindows.py (https://github.com/simonhmartin/genomics_general/blob/master/popgenWindows.py).

Fst can be influenced by heterogeneous genetic diversity between populations, and Herman et al.³⁷ demonstrated that low Pi in caves can inflate relative divergence estimates in *A. mexicanus*. We therefore chose to use Dxy, which is not affected by levels of nucleotide diversity within populations, to identify regions of high genetic divergence between Chica pools.

We calculated Dxy on a site-by-site basis using a custom python script. This allowed us to calculate mean Dxy for each gene in the *A. mexicanus* genome annotation (v101, downloaded from ftp://ftp.ensembl.org/pub/release-101/gtf/astyanax_mexicanus/).

Genome-wide tests for introgression

The population of fish within Chica cave has been hypothesized to be a hybrid swarm between cavefish originating from other caves in the El Abra region (which enter into Chica cave via a subterranean connection) and surface fish from the nearby Río Choy/Tampaón river system⁹. To formally test this hypothesis, we conducted genome-wide tests for introgression between Chica cavefish and Tinaja cavefish and between Chica cavefish and Río Choy surface fish. We first used Treemix v1.13⁷⁹ to confirm relationships between our focal populations and to visualize migration events between populations. Treemix builds a bifurcating tree to represent population splits and also incorporates migration events, which are represented as “edges” connecting population branches. We first built the maximum likelihood tree (zero migration events) in Treemix and then ran Treemix sequentially with one through five migration events. For this analysis, we included individuals from Chica, Río Choy (new lineage surface), Rascón (old lineage surface), and Tinaja (old lineage cave) *A. mexicanus* populations and the *A. aeneus* individual (outgroup), and SNPs were thinned to 1kb apart. We supplied this set of 700,502 biallelic SNPs to Treemix, rooted with *A. aeneus*, and estimated the covariance matrix between populations using blocks of 500 SNPs. Samples Tinaja_E, Tinaja_6, and Rascon_6 were excluded from this analysis because ADMIXTURE indicated that they were likely early generation hybrids. We calculated the variance explained by each model (zero through five migration events) using the R script `treemixVarianceExplained.R`⁸⁰.

To test our hypothesis that Chica represents a hybrid population resulting from admixture between the nearby old lineage cave and new lineage surface populations, we used Dsuite v0.4⁸¹ to conduct formal tests for introgression between (1) Chica cavefish and Tinaja cavefish, and (2) between Chica cavefish and Río Choy surface fish. If no gene flow is occurring between the fish in Chica cave and the local surface population, we predict that fish from Chica (which has previously been shown to group phylogenetically with old lineage cavefish populations) should share more derived alleles with fish from Rascón (a surface population that is more geographically distant from Chica but also old lineage) than fish from Río Choy (a surface population that is geographically close to Chica cave but is new lineage). For this analysis, we supplied the set of 700,502 biallelic SNPs to Dsuite and specified *A. aeneus* as the outgroup. We again excluded three samples from Tinaja and Rascón with apparent hybrid ancestry. We used the Dsuite program Dtrios to calculate Patterson's D statistic for all possible trios of populations using the ABBA-BABA test⁸². The ABBA-BABA test quantifies whether allele frequencies follow those expected between three lineages (e.g., sister species P1 and P2, and a third closely related species, P3) under expectations for incomplete lineage sorting (ILS). Observing a greater proportion of shared derived alleles between P1 and P3 but not P2 or between P2 and P3 but not P1 than what would be expected by chance (i.e., ILS) indicates introgression. Dsuite requires a fourth population, P4, to serve as an outgroup and determine which alleles are ancestral versus derived. Ancestral alleles are labeled as "A" and derived alleles are labeled as "B". ABBA sites are those where P2 and P3 share a derived allele, and ABAB sites are those where P2 and P4 share a derived allele. The D statistic is calculated as the difference in the number of ABBA and BABA sites relative to the total number of sites examined. Dsuite uses jackknifing of the null hypothesis that no introgression has occurred (D statistic = 0) to calculate a p-value for each possible trio of populations.

Dsuite also calculates the admixture fraction, or f_4 -ratio, which represents the covariance of allele frequency differences between P1 and P2 and between P3 and P4. If no introgression has occurred since P1 and P2 split from P3 and P4, then $f_4 = 0$. The f_4 statistic is positive, this suggests a discordant tree topology indicative of introgression.

Local Ancestry Inference

Hybrid genomes exhibit a mosaic of ancestry from their parental populations. A number of recent studies have shown that hybridization interacts with recombination and selection to shape patterns of local ancestry along chromosomes^{83–87}. Non-random distributions of local ancestry in hybrid populations can indicate selection. Our goal here was to visualize patterns of introgression across the genome in Chica cavefish and determine whether more surface ancestry is present in Chica Pool 1 compared to Chica Pool 2. We used Hidden Markov Model (HMM) and fine-scale SNP mapping approaches to calculate ancestry proportions globally (i.e. genome-wide) and locally (at each base pair along each of the 25 chromosomes) in both Chica pools. To determine whether Pool 1 (nearer to the cave entrance) carries a higher proportion of surface ancestry compared to Pool 2 (deeper in the cave) at regions of the genome important to cave adaptation, we also asked whether regions of high divergence between pools exhibit higher differences in local ancestry.

We implemented a HMM-based approach in Loter⁸⁸ to infer genome-wide local ancestry in the Chica individuals. Tinaja and Río Choy served as the parental cave and surface populations, respectively, for the initial training stage of the HMM. We excluded two Tinaja samples that showed putative evidence of admixture³⁷. This analysis allowed us to estimate global ancestry proportions and mean minor and major parent tract lengths for each individual. Ancestry tract lengths were converted from base pairs to Morgans using the median genome-wide recombination rate of median recombination rate of 1.16 cM/Mb obtained from a previously

published genetic map for *A. mexicanus*⁸⁹. We then estimated the number of generations since the onset of admixture (T_{admix}) in each pool using the following equation:

$$T_{\text{admix}} = 1/(L_M * p_B)$$

where L_M is the mean ancestry tract length from the minor parent in Morgans and p_B is the proportion of the genome derived from the major parent (the probability of recombining)^{90–92}.

We next used a chromosome painting approach with ancestry-informative sites to validate the delimitation of ancestry blocks detected by the HMM and to visualize patterns of introgression across the Chica cavefish genomes. This approach provides a lower level of resolution for ancestry block delimitation but with higher power to classify regions as derived from either parental genome. We identified alleles that were differentially fixed in Río Choy and Tinaja parental populations and had no missing data using the script `get_fixed_site_gts.rb` (https://github.com/mmatschiner/tutorials/blob/master/analysis_of_introgression_with_snp_data/src/get_fixed_site_gts.rb). We thinned SNPs to be a minimum of 1 kb apart and mapped these ancestry-informative sites in the Chica samples using the script `plot_fixed_site_gts.rb` (https://github.com/mmatschiner/tutorials/blob/master/analysis_of_introgression_with_snp_data/src/plot_fixed_site_gts.rb).

Synthesizing patterns of genetic divergence and local ancestry

To quantify and visualize patterns of divergence between Pool 1 and Pool 2, we calculated summary statistics (D_{xy} , F_{st} , P_i) in non-overlapping 50kb windows across the genome using the python script `popgenWindows.py` (https://github.com/simonhmartin/genomics_general/blob/master/popgenWindows.py). We also

calculated Dxy on a site-by-site basis using a custom python script (Cave_fish_Dxy.py). We asked whether there was an association between differences in local ancestry between pools and absolute genetic divergence (Dxy) between pools within outlier windows (which included coding and non-coding regions) and in coding regions alone.

We identified outlier windows as any 50kb window with a Dxy value above the 95th percentile ($Dxy > 0.0035$). Within each outlier window, we calculated the difference in local ancestry between fish from Pool 1 and Pool 2 at each site. We used a Wilcoxon rank sum test to identify whether ancestry differed within these regions between fish from Pool 1 versus Pool 2. We used Pearson's correlation implemented in R (v4.0.2) to test for an association between difference in local ancestry and sequence divergence (Dxy) at each site between Chica Pool 1 and Pool 2 within outlier windows.

We calculated summary statistics for the coding region of each gene (i.e., max, median and mean Dxy, number of variant and invariant sites) within the *A. mexicanus* annotation (v101, downloaded from ftp://ftp.ensembl.org/pub/release-101/gtf/astyanax_mexicanus/) using a custom python script (Dxy_Summary_per_gene_ensemblGTF.py). This allowed us to rank genes by relative level of differentiation between Pool 1 and Pool 2. From this ranked list, we considered all genes with a mean Dxy above the 95th percentile ($Dxy > 0.00276$) as putative candidates for cave adaptation.

We used the GO Consortium Gene Ontology Enrichment Analysis tool (<http://geneontology.org/>) to ask whether any categories of biological processes were overrepresented in our set of outlier genes. We used the human (*Homo sapiens*) reference database for this analysis (20,851 genes). Fisher's exact tests were performed to determine whether the number of genes associated with

a given ontology were over- or under-represented in our set of outlier genes relative to the reference database.

We identified coding variants present among both Chica pools, Tinaja, and Río Choy and predicted the consequence of each variant on protein function using *in silico* computational analysis with the SIFT (sorting intolerant from tolerant) algorithm⁴⁷ and the Ensembl Variant Effect Predictor (VEP) software suite⁴⁸. SIFT uses sequence homology and data on the physical properties of a given protein to predict whether an amino acid substitution will be tolerated or deleterious. VEP performs annotation and analysis of genomic variants to predict impact on the protein sequence (i.e., modifier, low, moderate, or high).

Preliminary analyses indicated that one of our top candidate genes with high sequence divergence between Chica pools (*cry1a*) harbored a putative deleterious coding mutation (R263Q). To determine whether this variant is derived in cavefish and assess whether it occurs at evolutionarily conserved sites, we used the *Astyanax* surface fish genome annotation to obtain the CDS for *cry1a* from our population genomic data. We searched Ensembl for gene orthologs available in other animal species, including human, mouse, zebrafish, staghorn coral (*Acropora millepora*), thale cress (*Arabidopsis thaliana*), and three cyprinid cavefish species from China, the blind barbel (*Sinocyclocheilus anshuiensis*), the golden-line barbel (*Sinocyclocheilus grahami*), and the horned golden-line barbel (*Sinocyclocheilus rhinoceros*). We also downloaded the CDS for *cry1a* from another cyprinid cavefish species from Somalia, *Phreatichthys andruzzii*, from NCBI. We conducted a multiple species alignment for all 285 *cry1* orthologs using Muscle⁹³. While investigating the R263Q mutation in *cry1a*, we identified a misassembly in the *Astyanax mexicanus* surface genome (*Astyanax_mexicanus*-2.0, downloaded from NCBI) affecting exons 9-13 of the *cry1a* coding region (*cry1a* CDS: 14,394-15,659 bp). Further investigation revealed that a portion of the coding region (*cry1a* CDS: 268-597 bp) was missing from the Pachon cavefish

genome assembly (*Astyanax_mexicanus*-1.0.2, downloaded from NCBI). To confirm the mutation we identified in our population genomic data, we downloaded previously published *cry1a* mRNA sequences with complete CDS from Chica cave, Pachon cave, and Micos River (NCBI accession #s KF737846- KF737848). Aligning our population genomic data to the mRNA allowed us to verify that the correct exon coordinates were used around the mutation of interest. To visualize the location of the R263Q mutation, we created a 3D model of the *Astyanax* Pachon cavefish CRY1A protein in SWISS-MODEL⁹⁴ using mouse CRY1 crystal structure (PDB: 6kx7). We imported the model into VMD (version 1.9.4) for visualization. To visualize the phylogenetic relationship between lineages with the R263Q mutation and identify putative instances of convergent evolution, we constructed a species tree that included 23 animal species (subterranean lineages and their close relatives) based on the species tree available from Ensembl release 102 and^{55,59,60}.

To test for signatures of selection in regions of the genome containing outlier genes, we used diploS/HIC⁴⁶ to detect and classify selective sweeps. diploS/HIC uses a powerful supervised machine learning approach to identify windows in the genome that have undergone “soft” sweeps (selection on standing genetic variation) or “hard” sweeps (selection on new mutations) with high accuracy. We first simulated selective sweeps using discoal⁹⁵ and then used the simulated data to train diploS/HIC. We provided diploS/HIC with a VCF containing the 3,337,738 SNPs showing <20% missing data across all populations and a masked version of the surface fish genome. We generated feature vectors for both Chica pools using the default settings of 11 sub-windows across a 1,100,000 Mb region (i.e., each window was 100,000 kb). diploS/HIC ran predictions using the feature vectors to classify each window as neutral (no evidence of a selective sweep), linkedSoft (loci near a window that has undergone a soft sweep), linkedHard (loci near a window that has undergone a hard sweep), Soft (loci that have undergone a soft sweep), or Hard (loci that have undergone a hard sweep). Windows lacking sufficient SNP data to make a prediction were labeled as “NA”.

To further investigate whether outlier genes between Chica pools are associated with phenotypic differences between cavefish and surface fish, we examined differential expression between lab-raised fry from Río Choy surface populations using a recently published RNAseq data set⁴⁹ (SRA Project Accession #PRJNA421208). Briefly, batches of fry were sacrificed every 4 hrs between 6 am and 10 pm (i.e., 0 hrs, 4 hrs, 8 hrs, 16 hrs, 20 hrs; sample size mean $\pm$ SE across time points: $n_{\text{Choy}} = 5.33 \pm 0.33$, $n_{\text{Tinaja}} = 5.83 \pm 0.17$) at 30 days post-fertilization (dpf) for whole-body RNA extraction and sequencing⁴⁹. Read counts for each gene across each sample were calculated as described in ⁴⁹. We used DESeq2⁹⁶ to calculate Log2(cavefish/surface fish) values. We considered genes to be differentially expressed if they had a Benjamini–Hochberg adjusted p-value < 0.05. Gene with less than 100 counts across all samples were excluded from the analyses.

Fish collection and maintenance for phenotyping

We phenotyped wild-caught fish from Pools 1 and 2 within Chica cave and from two other cave populations, Pachón and Los Sabinos, which served as controls. Adult fish were collected from in 2015, during the dry season. The fish used in these analyses were the same fish that were sampled in 2015 for genomic sequencing analyses. The fish were transported and housed in the aquatic facility at Universidad Autónoma de Querétaro in 24 hour constant darkness. Fish were fed 1-2 times daily with dry flakes and kept at 23°C. These conditions were maintained throughout housing and experimental conditions for consistency. All fish were inspected for overall health, and any exhibiting signs of health or stress issues were excluded from experimental tests. All the samples were collected under the auspices of the permit SGPA/DGVIS/0266/15, delivered by SEMARNAT. After the completion of behavioral assays, fin clips were collected from all Chica individuals for use in genomic sequencing as described above.

Sleep behavior phenotyping

Fish were maintained in the lab for 8 months prior to behavioral assays. Adult fish were recorded in standard conditions in 10L tanks with custom-designed partitions that allowed for five fish (2L/fish) to be individually housed in each tank as previously described³⁹. Recording chambers were illuminated with custom-designed IR LED source (Infrared 850 nm 5050 LED Strip Light, Environmental Lights). After a 4–5 day acclimation period, behavior was recorded for 24 hr beginning ZT0-ZT2. Videos were recorded at 15 frames/sec using a USB webcam (LifeCam Studio 1080 p HD Webcam, Microsoft) fitted with a zoom lens (Zoom 7000, Navitar). An IR high-pass filter (Edmund Optics Worldwide) was placed between the camera and the lens to block visible light. Videos were recorded using Virtualdub, a video-capturing software (Version 1.10.4) and were subsequently processed using Ethovision XT 9.0 (Noldus, IT). Water temperature and chemistry were monitored throughout recordings, and maintained at standard conditions in all cases. Ethovision tracking was set up as previously described³⁹. Data was processed using Perl scripts (v5.22.0, developed on-site) and Excel macro (Microsoft)³⁹. These data were used to calculate sleep information by finding bouts of immobility of 60 s and greater, which are highly correlated with increased arousal threshold, one of the hallmarks of sleep³⁹.

Morphological Characterization

Melanophores were quantified from bright-field images captured from each side of the body. Areas were chosen based on previous literature⁹⁷ (i.e., caudal fin area, adipose fin area, dorsal area, eye cup area, anal fin area, infra-orbital area; see Extended Data Figure 1). Briefly, images were loaded into Fiji ImageJ (v. 1.7, National Institutes of Health, Bethesda, MD). Images were color inverted in the selected area and using a preset noise tolerance allowed for melanophores to be automatically quantified by using pixel light intensity. If any melanophores were not counted, they were then manually added. Each image was analyzed by two different researchers to assure no significant discrepancies in quantifying, and the population of origin was blind to the

researchers. All final quantifications were corrected to body length to account for different sized fish.

Eye presence and size were determined from images acquired on a handheld digital microscope (Dinoscope Pro AM4111T). Images were analyzed in Fiji ImageJ. Each image was inspected for the presence of an eye by two investigators, and the population of origin was blind to the researchers. Eye size was calculated in ImageJ by creating an ROI for the eye diameter and dividing this number by the length of the body to correct for overall size differences.

Acknowledgements

We thank the University of Minnesota Genomics Center for their guidance and performing the cDNA library preparations and Illumina HiSeq 2500 sequencing. The Minnesota Supercomputing Institute (MSI) at the University of Minnesota provided resources that contributed to the research results reported within this paper. Funding was supported by NIH (1R01GM127872-01 to SEM, ACK, and NR) and NSF award IOS 165674 to ACK, and a US-Israel BSF award to ACK, and NSF IOS-1933076 to JK, SEM, and NR. Fish were collected under CONAPESCA permit PPF/DGOPA - 106 / 2013 to Claudia Patricia Ornelas García and SEMARNAT permit 02241 to Ernesto Maldonado. We thank the Mexican government for providing the collecting permit to R.B. in 2008 (DGOPA.00570.288108-0291).

References Cited

1. Lenormand, T. Gene flow and the limits to natural selection. *Trends in Ecology and Evolution* (2002). doi:10.1016/S0169-5347(02)02497-7

- 729 2. Marques, D. A. *et al.* Genomics of Rapid Incipient Speciation in Sympatric Threespine
730 Stickleback. *PLoS Genet.* **12**, 1–34 (2016).
- 731 3. Hoekstra, H. E., Krenz, J. G. & Nachman, M. W. Local adaptation in the rock pocket
732 mouse (*Chaetodipus intermedius*): natural selection and phylogenetic history of
733 populations. *Heredity (Edinb)*. **94**, 217–228 (2005).
- 734 4. Fitzpatrick, S. W. *et al.* Gene Flow Constrains and Facilitates Genetically Based
735 Divergence in Quantitative Traits. *Copeia* (2017). doi:10.1643/CI-16-559
- 736 5. Suarez-Gonzalez, A., Lexer, C. & Cronk, Q. C. B. Adaptive introgression: A plant
737 perspective. *Biol. Lett.* (2018). doi:10.1098/rsbl.2017.0688
- 738 6. Hedrick, P. W. Adaptive introgression in animals: Examples and comparison to new
739 mutation and standing variation as sources of adaptive variation. *Molecular Ecology*
740 (2013). doi:10.1111/mec.12415
- 741 7. Racimo, F., Sankararaman, S., Nielsen, R. & Huerta-Sánchez, E. Evidence for archaic
742 adaptive introgression in humans. *Nature Reviews Genetics* (2015). doi:10.1038/nrg3936
- 743 8. Samuk, K. *et al.* Gene flow and selection interact to promote adaptive divergence in
744 regions of low recombination. *Mol. Ecol.* (2017). doi:10.1111/mec.14226
- 745 9. Mitchell, R. W., Russell, W. H. & Elliott, W. R. *Mexican eyeless characin fishes, genus*
746 *Astyanax: environment, distribution, and evolution*. (Texas Tech Press, 1977).
- 747 10. Mallet, J. Hybridization as an invasion of the genome. *Trends Ecol. Evol.* **20**, 229–237
748 (2005).
- 749 11. Abbott, R. *et al.* Hybridization and speciation. *J. Evol. Biol.* **26**, 229–246 (2013).
- 750 12. Taylor, S. A. & Larson, E. L. Insights from genomes into the evolutionary importance and
751 prevalence of hybridization in nature. *Nature Ecology and Evolution* (2019).
752 doi:10.1038/s41559-018-0777-y
- 753 13. Buerkle, C. A. & Lexer, C. Admixture as the basis for genetic mapping. *Trends Ecol. Evol.*
754 **23**, 686–694 (2008).

14. Crawford, J. E. & Nielsen, R. Detecting adaptive trait loci in nonmodel systems: divergence or admixture mapping? *Mol. Ecol.* **22**, 6131–6148 (2013).
15. Chakraborty, R. & Weiss, K. M. Admixture as a tool for finding linked genes and detecting that difference from allelic association between loci. *Proc. Natl. Acad. Sci. U. S. A.* (1988). doi:10.1073/pnas.85.23.9119
16. Risch, N. Mapping genes for complex diseases using association studies with recently admixed populations. *Am J Hum Genet Suppl* **51**, 13 (1992).
17. Rieseberg, L. H., Whitton, J. & Gardner, K. Hybrid zones and the genetic architecture of a barrier to gene flow between two sunflower species. *Genetics* **152**, 713–727 (1999).
18. Luttikhuisen, P. C., Drent, J., Peijnenburg, K. T. C. A., Van Der Veer, H. W. & Johannesson, K. Genetic architecture in a marine hybrid zone: Comparing outlier detection and genomic clines analysis in the bivalve *Macoma balthica*. *Mol. Ecol.* (2012). doi:10.1111/j.1365-294X.2012.05586.x
19. Malek, T. B., Boughman, J. W., Dworkin, I. & Peichel, C. L. Admixture mapping of male nuptial colour and body shape in a recently formed hybrid population of threespine stickleback. *Mol. Ecol.* (2012). doi:10.1111/j.1365-294X.2012.05660.x
20. vonHoldt, B. M., Kays, R., Pollinger, J. P. & Wayne, R. K. Admixture mapping identifies introgressed genomic regions in North American canids. *Mol. Ecol.* (2016). doi:10.1111/mec.13667
21. Brelsford, A., Toews, D. P. L. & Irwin, D. E. Admixture mapping in a hybrid zone reveals loci associated with avian feather coloration. *Proc. R. Soc. B Biol. Sci.* (2017). doi:10.1098/rspb.2017.1106
22. Bresadola, L. *et al.* Admixture mapping in interspecific *Populus* hybrids identifies classes of genomic architectures for phytochemical, morphological and growth traits. *New Phytol.* (2019). doi:10.1111/nph.15930
23. Calfee, E., Agra, M. N., Palacio, M. A., Ramírez, S. R. & Coop, G. Selection and

- hybridization shaped the rapid spread of African honey bee ancestry in the americas.
PLoS Genet. (2020). doi:10.1371/journal.pgen.1009038
24. Powell, D. L. *et al.* Natural hybridization reveals incompatible alleles that cause melanoma in swordtail fish. *Science* (80-.). (2020). doi:10.1126/science.aba5216
25. Gompert, Z., Mandeville, E. G. & Buerkle, C. A. Analysis of Population Genomic Data from Hybrid Zones. *Annu. Rev. Ecol. Evol. Syst.* (2017). doi:10.1146/annurev-ecolsys-110316-022652
26. Ravinet, M. *et al.* Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. *J. Evol. Biol.* **30**, 1450–1477 (2017).
27. Kowalko, J. E. *et al.* Loss of schooling behavior in cavefish through sight-dependent and sight-independent mechanisms. *Curr. Biol.* **23**, 1874–1883 (2013).
28. Yoshizawa, M., Gorički, Š., Soares, D. & Jeffery, W. R. Evolution of a behavioral shift mediated by superficial neuromasts helps cavefish find food in darkness. *Curr. Biol.* **20**, 1631–1636 (2010).
29. Duboué, E. R., Keene, A. C. & Borowsky, R. L. Evolutionary convergence on sleep loss in cavefish populations. *Curr. Biol.* (2011). doi:10.1016/j.cub.2011.03.020
30. Bibliowicz, J. *et al.* Differences in chemosensory response between eyed and eyeless *Astyanax mexicanus* of the Rio Subterráneo cave. *Evodevo* (2013). doi:10.1186/2041-9139-4-25
31. Aspirasa, A. C., Rohnera, N., Martineau, B., Borowsky, R. L. & Tabina, C. J. Melanocortin 4 receptor mutations contribute to the adaptation of cavefish to nutrient-poor conditions. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 9668–9673 (2015).
32. Espinasa, L. *et al.* Discovery of two new *astyanax* cavefish localities leads to further understanding of the species biogeography. *Diversity* (2020). doi:10.3390/d12100368
33. Ornelas-García, C. P., Domínguez-Domínguez, O. & Doadrio, I. Evolutionary history of the fish genus *astyanax* baird & Girard (1854) (Actinopterygii, Characidae) in

- 807 mesoamerica reveals multiple morphological homoplasies. *BMC Evol. Biol.* (2008).
- 808 doi:10.1186/1471-2148-8-340
- 809 34. Bradic, M., Beerli, P., García-de León, F. J., Esquivel-Bobadilla, S. & Borowsky, R. L.
- 810 Gene flow and population structure in the Mexican blind cavefish complex (*Astyanax*
- 811 *mexicanus*). *BMC Evol. Biol.* **12**, 9 (2012).
- 812 35. Strecker, U., Hausdorf, B. & Wilkens, H. Parallel speciation in *Astyanax* cave fish
- 813 (Teleostei) in Northern Mexico. *Mol. Phylogenet. Evol.* **62**, 62–70 (2012).
- 814 36. Coghill, L. M., Darrin Hulsey, C., Chaves-Campos, J., García de Leon, F. J. & Johnson,
- 815 S. G. Next generation phylogeography of cave and surface *Astyanax mexicanus*. *Mol.*
- 816 *Phylogenet. Evol.* **79**, 368–374 (2014).
- 817 37. Herman, A. *et al.* The role of gene flow in rapid and repeated evolution of cave-related
- 818 traits in Mexican tetra, *Astyanax mexicanus*. *Mol. Ecol.* (2018). doi:10.1111/mec.14877
- 819 38. Moran, D., Softley, R. & Warrant, E. J. Eyeless Mexican Cavefish Save Energy by
- 820 Eliminating the Circadian Rhythm in Metabolism. *PLoS One* **9**, e107877 (2014).
- 821 39. Yoshizawa, M. *et al.* Distinct genetic architecture underlies the emergence of sleep loss
- 822 and prey-seeking behavior in the Mexican cavefish. *BMC Biol.* (2015).
- 823 doi:10.1186/s12915-015-0119-3
- 824 40. Romero, A. Introgressive hybridization in the *Astyanax fasciatus* (Pisces: Characidae)
- 825 population at La Cueva Chica. *NSS Bull.* (1983).
- 826 41. Breder, C. M. Descriptive ecology of La Cueva Chica, with especial reference to the blind
- 827 fish, *Anoptichthys*. *Zoologica* **27**, 7–16 (1942).
- 828 42. Bridges, W. The blind fish of La cueva Chica. *Bull. New York Zool. Soc* **43**, 74–97 (1940).
- 829 43. Elliott, W. R. Cave Biodiversity and Ecology of the Sierra de El Abra Region. in *Biology*
- 830 *and Evolution of the Mexican Cavefish* 59–76 (Elsevier, 2015). doi:10.1016/B978-0-12-
- 831 802148-4.00003-7
- 832 44. Şadoğlu, P. A mendelian gene for albinism in natural cave fish. *Experientia* (1957).

doi:10.1007/BF02161111

45. Wilkens, H. Evolution and Genetics of Epigean and Cave Astyanax fasciatus (Characidae, Pisces). in *Evolutionary Biology* (1988). doi:10.1007/978-1-4613-1043-3_8
46. Kern, A. D. & Schrider, D. R. DiploS/HIC: An updated approach to classifying selective sweeps. *G3 Genes, Genomes, Genet.* (2018). doi:10.1534/g3.118.200262
47. Kumar, P., Henikoff, S. & Ng, P. C. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat. Protoc.* (2009). doi:10.1038/nprot.2009.86
48. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* (2016). doi:10.1186/s13059-016-0974-4
49. McGaugh, S. E., Passow, C. N., Jaggard, J. B., Stahl, B. A. & Keene, A. C. Unique transcriptional signatures of sleep loss across independently evolved cavefish populations. *J. Exp. Zool. Part B Mol. Dev. Evol.* (2020). doi:10.1002/jez.b.22949
50. Cashmore, A. R., Jarillo, J. A., Wu, Y. J. & Liu, D. Cryptochromes: blue light receptors for plants and animals. *Science (80-.).* **284**, 760–765 (1999).
51. Hirayama, J. *et al.* The clock components Period2, Cryptochrome1a, and Cryptochrome2a function in establishing light-dependent behavioral rhythms and/or total activity levels in zebrafish. *Sci. Rep.* (2019). doi:10.1038/s41598-018-37879-8
52. Patke, A. *et al.* Mutation of the Human Circadian Clock Gene CRY1 in Familial Delayed Sleep Phase Disorder. *Cell* **169**, 203-215.e13 (2017).
53. Ceinos, R. M. *et al.* Mutations in blind cavefish target the light-regulated circadian clock gene, period 2. *Sci. Rep.* (2018). doi:10.1038/s41598-018-27080-2
54. Cavallari, N. *et al.* A blind circadian clock in cavefish reveals that opsins mediate peripheral clock photoreception. *PLoS Biol.* (2011). doi:10.1371/journal.pbio.1001142
55. Yang, J. *et al.* The Sinocyclocheilus cavefish genome provides insights into cave adaptation. *BMC Biol.* **14**, 1 (2016).

56. Riccio, A. P. & Goldman, B. D. Circadian rhythms of locomotor activity in naked mole-rats (*Heterocephalus glaber*). *Physiol. Behav.* **71**, 1–13 (2000).
57. Czarna, A. *et al.* XStructures of drosophila cryptochrome and mouse cryptochrome1 provide insight into circadian function. *Cell* (2013). doi:10.1016/j.cell.2013.05.011
58. Beale, A. *et al.* Circadian rhythms in Mexican blind cavefish *Astyanax mexicanus* in the lab and in the field. *Nat. Commun.* (2013). doi:10.1038/ncomms3769
59. Colli, L., Paglianti, A., Berti, R., Gandolfi, G. & Tagliavini, J. Molecular phylogeny of the blind cavefish *Phreatichthys andruzzii* and *Garra barreimiae* within the family Cyprinidae. *Environ. Biol. Fishes* **84**, 95–107 (2009).
60. Patterson, B. D. & Upham, N. S. A newly recognized family from the Horn of Africa, the Heterocephalidae (Rodentia: Ctenohystrica). *Zool. J. Linn. Soc.* **172**, 942–963 (2014).
61. Jaggard, J. *et al.* The lateral line confers evolutionarily derived sleep loss in the Mexican cavefish. *J. Exp. Biol.* **220**, 284–293 (2017).
62. Lamichhaney, S. *et al.* Rapid hybrid speciation in Darwin’s Finches. *Science* (80-.). (2018). doi:10.1126/science.aao4593
63. Bergman, T. J., Phillips-Conroy, J. E. & Jolly, C. J. Behavioral variation and reproductive success of male baboons (*Papio anubis* x *papio hamadryas*) in a hybrid social group. *Am. J. Primatol.* (2008). doi:10.1002/ajp.20467
64. Ackermann, R. R. *et al.* Hybridization in human evolution: Insights from other organisms. *Evolutionary Anthropology* (2019). doi:10.1002/evan.21787
65. Ackermann, R. R., Mackay, A. & Arnold, M. L. The Hybrid Origin of “Modern” Humans. *Evol. Biol.* (2016). doi:10.1007/s11692-015-9348-1
66. Dowling, T. E., Martasian, D. P. & Jeffery, W. R. Evidence for multiple genetic forms with similar eyeless phenotypes in the blind cavefish, *Astyanax mexicanus*. *Mol. Biol. Evol.* **19**, 446–455 (2002).
67. Avise, J. C. & Selander, R. K. Evolutionary genetics of cave-dwelling fishes of the genus

885 Astyanax. *Evolution* (N. Y). 1–19 (1972).

886 68. Fitzpatrick, M. J. *et al.* Candidate genes for behavioural ecology. *Trends in Ecology and*
887 *Evolution* (2005). doi:10.1016/j.tree.2004.11.017

888 69. Niepoth, N. & Bendesky, A. How Natural Genetic Variation Shapes Behavior. *Annu. Rev.*
889 *Genomics Hum. Genet.* **21**, (2020).

890 70. Hoban, S. *et al.* Finding the genomic basis of local adaptation: pitfalls, practical solutions,
891 and future directions. *Am. Nat.* **188**, 379–397 (2016).

892 71. Tigano, A. & Friesen, V. L. Genomics of local adaptation with gene flow. *Mol. Ecol.* **25**,
893 2144–2164 (2016).

894 72. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
895 next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

896 73. Depristo, M. A. *et al.* A framework for variation discovery and genotyping using next-
897 generation DNA sequencing data. *Nat. Genet.* (2011). doi:10.1038/ng.806

898 74. Van der Auwera, G. A. *et al.* From FastQ Data to High-Confidence Variant Calls: The
899 Genome Analysis Toolkit Best Practices Pipeline. *Curr. Protoc. Bioinforma.* **43**, 11.10.1–
900 11.10.33 (2013).

901 75. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
902 transform. *Bioinformatics* **25**, 1754–1760 (2009).

903 76. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
904 2078–2079 (2009).

905 77. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
906 (2011).

907 78. McGaugh, S. E. S. *et al.* The cavefish genome reveals candidate genes for eye loss. *Nat.*
908 *Commun.* (2014). doi:10.1038/ncomms6307

909 79. Pickrell, J. & Pritchard, J. Inference of population splits and mixtures from genome-wide
910 allele frequency data. *Nat. Preced.* **1** (2012).

- 911 80. Card, D. *RADpipe*. (GitHub repository, 2015). doi:10.5281/zenodo.17809
- 912 81. Malinsky, M. Dsuite-fast D-statistics and related admixture evidence from VCF files.
913 *BioRxiv* **634477**, (2019).
- 914 82. Green, R. E. *et al.* A draft sequence of the Neandertal genome. *Science* (80-.). **328**,
915 710–722 (2010).
- 916 83. Hohenlohe, P. A., Bassham, S., Currey, M. & Cresko, W. A. Extensive linkage
917 disequilibrium and parallel adaptive divergence across threespine stickleback genomes.
918 *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 395–409 (2012).
- 919 84. Wang, J., Street, N. R., Scofield, D. G. & Ingvarsson, P. K. Variation in Linked Selection
920 and Recombination Drive Genomic Divergence during Allopatric Speciation of European
921 and American Aspens. *Mol. Biol. Evol.* **33**, 1754–1767 (2016).
- 922 85. Schumer, M. *et al.* Natural selection interacts with recombination to shape the evolution
923 of hybrid genomes. *Science* (80-.). **360**, 656–660 (2018).
- 924 86. Janzen, T., Nolte, A. W. & Traulsen, A. The breakdown of genomic ancestry blocks in
925 hybrid lineages given a finite number of recombination sites. *Evolution* (N. Y). (2018).
926 doi:10.1111/evo.13436
- 927 87. Leitwein, M. *et al.* The role of recombination on genome-wide patterns of local ancestry
928 exemplified by supplemented brook charr populations. *Mol. Ecol.* (2019).
929 doi:10.1111/mec.15256
- 930 88. Dias-Alves, T., Mairal, J. & Blum, M. G. B. Loter: A software package to infer local
931 ancestry for a wide range of species. *Mol. Biol. Evol.* (2018). doi:10.1093/molbev/msy126
- 932 89. O'Quin, K. E., Yoshizawa, M., Doshi, P. & Jeffery, W. R. Quantitative Genetic Analysis of
933 Retinal Degeneration in the Blind Cavefish *Astyanax mexicanus*. *PLoS One* **8**, (2013).
- 934 90. Gravel, S. Population genetics models of local ancestry. *Genetics* **191**, 607–619 (2012).
- 935 91. Jin, W., Li, R., Zhou, Y., Genetics, S. X.-E. J. of H. & 2014, undefined. Distribution of
936 ancestral chromosomal segments in admixed genomes and its implications for inferring

population history and admixture mapping. *nature.com*

92. Schumer, M., Cui, R., Powell, D. L., Rosenthal, G. G. & Andolfatto, P. Ancient hybridization and genomic stabilization in a swordtail fish. *Mol. Ecol.* **25**, 2661–2679 (2016).
93. Edgar, R. C. MUSCLE: A multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **32**, 1792–1797 (2004).
94. Waterhouse, A. *et al.* SWISS-MODEL: Homology modelling of protein structures and complexes. *Nucleic Acids Res.* (2018). doi:10.1093/nar/gky427
95. Kern, A. D. & Schrider, D. R. Discoal: Flexible coalescent simulations with selection. *Bioinformatics* (2016). doi:10.1093/bioinformatics/btw556
96. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).
97. Stahl, B. A., Sears, C. R., Ma, L., Perkins, M. & Gross, J. B. Pmela and Tyrp1b contribute to melanophore variation in Mexican Cavefish. in *Origin and Evolution of Biodiversity* (2018). doi:10.1007/978-3-319-95954-2_1

963
964
965
966
967
968
969
970

971 **Supplementary Information**

972 *Sequencing and Genotyping*

973 Sequencing resulted in a mean $\pm$ SE of 187,777,319 $\pm$ 3,047,876 reads per individual for the 19
974 Chica samples and 331,445,356 $\pm$ 248,640,606 reads per individual for the three Los Sabinos
975 samples. After quality filtering and mapping, all 60 samples had a mean $\pm$ SE genome-wide depth
976 of coverage of 10.50 $\pm$ 0.53X (Extended Data Table 2).

977

978 *Population Structure*

979 We examined population genetic structure among populations of cave and surface fish using PCA
980 and ADMIXTURE. Populations examined (i.e., Río Choy, Los Sabinos, Tinaja, Rascón, Pachón,
981 and Chica) generally showed separation from one another into distinct genetic clusters in both
982 PCA and ADMIXTURE analyses (Figure 2A,B; Extended Data Figure 2). The first three principal
983 components from the PCA explained over 45% of the total variance in the data (Extended Data
984 Figures 2,3; Extended Data Table 3), and separated the populations based on lineage and
985 ecotype (surface or cave). For the ADMIXTURE analysis, comparison of cross validation error for
986 K=2-9 indicated an optimal K of 5 (Extended Data Figure 4). Fish from Pool 1 and Pool 2 within
987 Chica cave clustered together in the ADMIXTURE analysis and samples from both pools overlap
988 completely in the PCA, indicating low overall levels of genetic divergence between the pools

(Extended Data Figure 5). Los Sabinos and Tinaja are neighboring caves in the El Abra region and individuals from these clustered together in both analyses (Figure 2A,B; Extended Data Figure 2).

Average nucleotide diversity (π) across the genome did not differ between pools within Chica cave ($\pi = 0.0021$ for Chica Pool 1 and Pool 2; Extended Data Table 4). Notably, nucleotide diversity within Chica cave was 2-3X higher compared to other caves (Extended Data Table 4) and absolute genetic divergence between pools within Chica cave ($D_{xy} = 0.0020$) is comparable to that observed among other cave populations (Extended Data Table 5). Despite the fact that we observed no sites differentially fixed between Chica Pool1 and Pool 2, indicative that gene flow is ongoing, we observed several peaks of high sequence divergence between pools (Fig. 2E).

Genome-wide tests for introgression

The results of two independent tests for introgression (implemented in Treemix and Dsuite) indicated hybridization between Chica cavefish and the nearby surface and cave lineages. Treemix first builds a bifurcating population tree and then fits for migration “edges” between branches. We chose the optimal number of migration events between populations in Treemix by examining the variance explained by models with zero through five migrations allowed. Adding one migration increased the proportion of variance explained from 0.59 to 0.92. The variance explained reached 1 and plateaued at three migration events (Extended Data Figure 6). Phylogenetic relationships were as expected based on previous analyses³⁷. Río Choy (new lineage) grouped with *A. aeneus*, and Rascón, Tinaja, and Chica (all old lineage) grouped together. We observed evidence of migration events between Río Choy surface fish with Chica cavefish and also between Tinaja cavefish and Chica cavefish (Figure 2C).

The results of the D statistic and f4 ratio tests implemented in Dsuite also confirmed introgression between Chica and Tinaja and between Chica and Río Choy. We were particularly interested in asking whether Chica has experienced recent gene flow with the local surface population, Río Choy. We observed that Río Choy and Chica share more derived sites (BBAA) than Chica and Rascón (ABBA) (Extended Data Table 6). Because Chica and Rascón populations are both derived from old lineage surface stock of *A. mexicanus*, whereas Río Choy is derived from new lineage surface stock, this pattern is indicative of introgression. Positive f4 ratios were also observed in all possible trios between Río Choy, Chica, Rascón, and Tinaja, indicative of introgression (Extended Data Table 6). Thus, our analyses support that Chica was originally colonized from an old lineage stock, but has subsequently experienced substantial introgression with new lineage surface populations.

Local Ancestry Inference

To characterize patterns of introgression in Chica cavefish genomes, we used HMM-based ancestry inference and SNP mapping at ancestry-informative sites. The HMM-based approach implemented in Loter revealed that the genomes of individuals from Chica cave were composed of approximately 75% cavefish (i.e. Tinaja) ancestry and 25% surface fish (i.e. Río Choy) ancestry (Extended Data Figure 6). The two pools within Chica cave exhibited nearly identical global ancestry proportions corresponding to surface (Pool 1 Surface Ancestry: Mean $\pm$ SE = 0.245 $\pm$ 0.004; Pool 2 Surface Ancestry: Mean $\pm$ SE = 0.244 $\pm$ 0.003) versus cave (Pool 1 Cave Ancestry: Mean $\pm$ SE = 0.755 $\pm$ 0.004; Pool 2 Cave Ancestry: Mean $\pm$ SE = 0.756 $\pm$ 0.003) parental populations. Surface ancestry tract length distribution was similar in both Chica pools (Extended Data Table 7). The timing since the onset of admixture also did not differ between pools (Mean $\pm$ SE generations since hybridization: Pool 1 = 2315 $\pm$ 19, Pool 2 = 2294 $\pm$ 17) (Extended Data Table 8).

We identified 89,810 sites with complete fixation of different alleles in the parental populations (i.e., Río Choy and Tinaja). We mapped these sites across all 25 diploid chromosomes for each of the 19 Chica fish. In general, ancestry was highly admixed across Chica genomes and showed a pattern in agreement with the findings of the HMM-based ancestry inference (Fig. 2D; Extended Data Figure 7).

Gene Expression

To investigate whether outlier genes may play a functional role in generating phenotypic variation associated with local adaptation, we examined differences in gene expression between lab-reared fry from Río Choy surface and Tinaja cave populations at multiple time points⁴⁹. Out of a total set of 27,420 genes, 18,958 were expressed in fry at 30 dpf, and 11,901 (63%) of expressed genes showed significant differential expression (Benjamini–Hochberg adjusted p-value < 0.05) between cave and surface fish during at least one time point (Extended Data Tables 12-13). For the subset of 706 genes in the top 5 % of Dxy values between Chica Pool 1 and Pool 2, 586 were expressed in fry at 30 dpf, and 389 (66%) of expressed genes showed significant differential expression between cave and surface fish during at least one time point (Extended Data Tables 12-13). This represents an enrichment of differentially expressed genes in the outlier genes compared to the entire genome-wide data set (χ^2 square test: $\chi^2_{1,19,544} = 3.168$, $p = 0.031$). Out of 26 top Dxy genes with ontologies related to cave adapted phenotypes (i.e., sleep/circadian cycle, light detection, eye size/morphology, metabolism, pigmentation), 21 were expressed at 30 dpf. Of those 21 genes, 14 (67%) showed significant differential expression between cave and surface fish during at least one time point (Extended Data Table 13).

Phenotyping

The pattern we documented of more surface-like phenotypes near the entrance of the cave (in Pool 1) and more cave-like phenotypes deeper in the cave (in Pool 2) (Fig. 1C,D; Fig. 4) contrasts

the findings of previous studies. Over eight decades ago,⁴² and ⁴¹ documented more cave-like phenotypes in the front of the cave (Pool 1) and more surface-like fish deeper in the cave (Pools 2-4). It was speculated that both cavefish and surface fish from nearby populations were entering Chica cave subterraneously near the lower pools, and that the connection between Pool 1 and Pools 2-4 was dynamic. A lack of connectivity between Pool 1 and the other pools was hypothesized to create a harsher, low nutrient environment towards the front of the cave, resulting in increased survival of fish with cave adapted phenotypes in this pool. However,⁴³ recently suggested that surface fish may have access to the entrance of the cave via runoff from nearby drainages. Indeed, river-dwelling fish species (i.e., cichlids and poeciliids) were observed near the entrance of the cave in Pool 1 when fish were collected for the present study in 2015. This suggests that surface fish may have access to Chica cave Pool 1 directly via the entrance and also via a deeper, subterraneous connection. Future studies using mark-recapture would provide a more definitive answer.

1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113

Extended Data Tables

Extended Data Table 1. GATK filters applied to variant and invariant sites.

Invariant sites	SNPs	Mixed/indels
QD < 2.0	QD < 2.0	QD < 2.0
FS > 60.0	FS > 200.0	FS > 200.0
MQ < 40.0	ReadPosRankSum < -20.0	ReadPosRankSum < -20.0

1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139

Extended Data Table 2. Read counts and coverage info.

https://docs.google.com/spreadsheets/d/1_3eFqa2BuOZWYVwU9J1_vDAEt9D3s4ySBtOFSUo_w3Ek/edit#gid=0

1140
1141
1142
1143
1144
1145
1146
1147
1148 **Extended Data Table 3.** Eigenvalues for PC 1-20 from PCA containing all six *Astyanax*
1149 populations (i.e., Chica, Río Choy, Los Sabinos, Pachón, Rascón, and Tinaja).

PC	EigenValue
1	3.49399
2	2.69367
3	2.48907
4	1.45645
5	1.17887
6	0.800319
7	0.796332
8	0.795117
9	0.789957
10	0.786389
11	0.764248
12	0.393822
13	0.390711
14	0.383791
15	0.378667
16	0.376847
17	0.337861
18	0.318567
19	0.2882
20	0.279671

1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169

Extended Data Table 4. Nucleotide diversity (π) within populations.

Population	n	π
Chica Pool 1	5	0.0021
Chica Pool 2	14	0.0021
Río Choy	9	0.0028
Pachón	10	0.0008
Rascón	10	0.0013
Los Sabinos	3	0.0007
Tinaja	8	0.0008

1170
1171
1172
1173
1174
1175
1176
1177

1178 **Extended Data Table 5.** Absolute genetic divergence (Dxy) and relative genetic divergence (Fst)
1179 between populations. See Extended Data Table 4 for population sample sizes.

Population 1	Population 2	Dxy	Fst
Chica Pool1	Chica Pool2	0.0021	0.0000
Chica Pool1	Río Choy	0.0033	0.1440
Chica Pool1	Pachón	0.0030	0.3501
Chica Pool1	Rascón	0.0030	0.2592
Chica Pool1	Los Sabinos	0.0024	0.2029
Chica Pool1	Tinaja	0.0024	0.2364
Chica Pool2	Río Choy	0.0033	0.1651
Chica Pool2	Pachón	0.0030	0.3693
Chica Pool2	Rascón	0.0030	0.2923
Chica Pool2	Los Sabinos	0.0024	0.1154
Chica Pool2	Tinaja	0.0024	0.2477
Río Choy	Pachón	0.0035	0.3654
Río Choy	Rascón	0.0033	0.2507
Río Choy	Sabinos	0.0034	0.1900
Río Choy	Tinaja	0.0034	0.3555
Pachón	Rascón	0.0026	0.4572
Pachón	Los Sabinos	0.0020	0.2832
Pachón	Tinaja	0.0020	0.4430
Rascón	Los Sabinos	0.0023	0.2002

Rascón	Tinaja	0.0023	0.4003
Los Sabinos	Tinaja	0.0008	0.0164

Extended Data Table 6. Results of D statistic and f4 ratio tests for introgression. *A. aeneus* served as the outgroup. BBAA = derived alleles shared by P1 and P2. ABBA = derived alleles shared by P2 and P3. BABA = derived alleles shared by P1 and P3. Significant p-values (<0.05) indicate evidence of introgression between Choy, Chica, Rascón, and Tinaja.

P1	P2	P3	D statistic	Z score	p-value	f4 ratio	BBAA	ABBA	BABA
Río Choy	Chica	Rascón	0.27893	21.4698	0	0.173087	8409.82	4100.59	2311.94
Río Choy	Tinaja	Chica	0.14629	7.98383	7.09E-16	0.262394	2195.94	9381.5	6986.95
Chica	Tinaja	Rascón	0.409914	33.6245	0	0.230212	8116.3	3375	1412.52
Rascón	Tinaja	Río Choy	0.2558	23.5379	0	0.11872	5652.15	3207.89	1901.03

Extended Data Table 7. Summary statistics for minor parent (i.e., Río Choy surface fish) ancestry tract lengths in base pair and estimated mean $\pm$ SE number of generations since the onset of admixture (T_{admix}) in Chica cave Pool 1 (n=5) and Pool 2 (n=14).

Population	Min	Max	Median	Mean	SE	T_{admix}
Pool 1	1,350	2,069,769	29,372	48,042	263	2315 $\pm$ 19
Pool 2	1,031	2,077,917	29,561	48,435	162	2294 $\pm$ 17

Extended Data Table 8. The mean minor parent (i.e. surface) tract lengths and mean major parent (i.e. cave) global ancestry proportions were used to infer the timing since admixture (T_{admix}).

Pool	ID	MeanSurface TractLengthBP	MeanSurfaceTract LengthMorgans	MeanCaveGlobal Ancestry	T_{admix}
1	Chica5_1	47096.4	0.000546	0.770	2377.275
1	Chica5_2	49508.8	0.000574	0.759	2294.262
1	Chica5_3	49475.3	0.000574	0.750	2324.321
1	Chica5_4	51214.4	0.000594	0.744	2261.418
1	Chica5_5	49574.2	0.000575	0.751	2315.212
2	Chica1_1	51893.7	0.000602	0.739	2246.508
2	Chica1_10	48084.2	0.000558	0.755	2374.725
2	Chica1_11	48674.3	0.000565	0.762	2323.384
2	Chica1_12	47755.7	0.000554	0.755	2389.959
2	Chica1_13	51812.5	0.000601	0.768	2166.720
2	Chica1_14	52055.8	0.000604	0.755	2193.243
2	Chica1_2	50207.8	0.000582	0.761	2257.289
2	Chica1_3	48756.4	0.000566	0.754	2343.562
2	Chica1_4	49882.2	0.000579	0.755	2289.383
2	Chica1_5	48534.3	0.000563	0.767	2315.360
2	Chica1_6	49014.7	0.000569	0.773	2276.677
2	Chica1_7	48674.4	0.000565	0.755	2345.607
2	Chica1_8	49867.9	0.000578	0.741	2333.061
2	Chica1_9	51125.9	0.000593	0.745	2262.986

1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259

Extended Data Table 9. Absolute genetic divergence (Dxy) for each gene in comparisons between Chica Pool 1 and Pool 2, Choy and Tinaja, Río Choy and Pachón, and Rascón and Pachón.
https://docs.google.com/spreadsheets/d/1yottC4COSed0BGbgjfWui0e4RLgnYa0Hiq_YO_fCc3k/edit#gid=2005687085

1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285

Extended Data Table 10. Results of GO Analysis on genes with highest genetic divergence (top 5% Dxy) between Chica Pool 1 and Chica Pool 2.
<https://drive.google.com/file/d/1RFayjPilvs8D-aXMa28abXI6Ghc5dx5t/view?usp=sharing>

1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311

Extended Data Table 11. Gene descriptions, phenotypes, and results of differential expression, SIFT, VEP, and diploS/HIC selection analyses for candidate genes with ontologies related to cave-adapted phenotypes and in the top 5% of Dxy values between Chica pools.

https://docs.google.com/spreadsheets/d/170S6uCleErV--V5uEzRDpUexXvwB5H_Y59_bD0_hTDk/edit?usp=sharing

1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337

Extended Data Table 12. Results of differential expression analysis across all time points.
[https://docs.google.com/spreadsheets/d/1Xmtlj965TAzRtgz7iQBEDCto132VceVIFK2_er7D3TQ/
edit?usp=sharing](https://docs.google.com/spreadsheets/d/1Xmtlj965TAzRtgz7iQBEDCto132VceVIFK2_er7D3TQ/edit?usp=sharing)

Extended Data Table 13. Summary statistics for differential expression analysis in 30 dpf fry from lab-raised cavefish (Tinaja cave) versus surface fish (Río Choy) stock across six timepoints. Summaries are shown for the entire dataset of all genes and a subset of genes within the top 5% of Dxy values between Chica pools.

Summary Statistics	0hr	4hr	8hr	12hr	16hr	20hr	Mean	SE
N Choy	6	4	6	5	6	5	5.33	0.33
N Tinaja	6	5	6	6	6	6	5.83	0.17
Total Genes	27420	27420	27420	27420	27420	27420	-	-
Genes Retained	18597	16252	18364	17747	17240	17501	17617	344
Genes Padj<0.05	3691	2913	7404	8358	6969	3932	5545	937
Prop. Genes Padj<0.05	0.20	0.18	0.40	0.47	0.40	0.23	0.31	0.05
Chica Top 5% Dxy Genes	706	706	706	706	706	706	-	-
Genes Retained	575	522	573	562	550	556	556	8
Genes Padj<0.05	133	103	239	277	225	142	187	28
Prop. Genes Padj<0.05	0.23	0.20	0.42	0.49	0.41	0.26	0.33	0.05

1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375

Extended Data Figures

Extended Data Figure 1. Eye and pigment morphology variations among cave populations. (A)

Diagram of areas used for pigmentation quantifications. (B) Brightfield images of fish bodies

showing pigmentation across multiple cave populations. (C) Pigment count quantifications by area. Eye cup: Kruskal-Wallis test, $P < 0.001$, KW statistic = 24.28. Chica 1 vs Chica 2, $p < 0.05$; Chica 1 vs Pachon, $p < 0.001$. Infra-Orbital: Kruskal-Wallis test, $P < 0.001$, KW statistic = 22.70. Chica 1 vs Chica 2, $p < 0.05$; Chica 1 vs Pachon, $p < 0.001$. Dorsal area: Kruskal-Wallis test, $P < 0.001$, KW statistic = 25.66. Chica 1 vs Pachon, $p < 0.001$; Chica 2 vs Pachon, $p < 0.05$; Pachon vs Los Sabinos, $p < 0.001$. Adipose area: Kruskal-Wallis test, $P < 0.001$, KW statistic = 23.65. Chica 1 vs Chica 2, $p < 0.05$; Chica 1 vs Pachon, $p < 0.001$, Pachon vs Los Sabinos, $p < 0.01$. Anal fin: Kruskal-Wallis test, $P < 0.01$, KW statistic = 12.44. Chica 2 vs Pachon, $p < 0.05$, Pachon vs Los Sabinos, $p < 0.01$. Tail area: Kruskal-Wallis test, $P < 0.001$, KW statistic = 16.87. Chica 1 vs Pachon, $p < 0.001$, Chica 1 vs Los Sabinos, $p < 0.05$.

Extended Data Figure 2. Biplots of scores for the first three PCs from the PCA on SNPs from cave (Chica, Pachón, Tinaja, Los Sabinos) and surface (Río Choy and Rascón) populations. Note that individuals from Chica Pool 1 and Pool 2 overlap and individuals from Tinaja and Los Sabinos overlap.

Extended Data Figure 3. Percentage of variance explained for PCs 1-20 from PCA containing all six *Astyanax* populations (i.e., Chica, Río Choy, Los Sabinos, Pachón, Rascón, and Tinaja).

1429

1430 **Extended Data Figure 4.** Cross validation (CV) error calculated in ADMIXTURE for values of K

1431 ranging from 2-9. A value of 5 is indicated to be the best estimate for the true number of

1432 populations clusters because it exhibits the lowest CV error.

1433

1434

1435

1436

1437

1438

1439

1446

1447 **Extended Data Figure 6.** Variance explained for 0-5 migration events in Treemix. Variance
 1448 explained plateaus at 3 migration events.

1449

1450

Extended Data Figure 7. Local ancestry tracts in Chica samples inferred using a Hidden Markov Model approach along each of the 25 chromosomes. Yellow represents cave ancestry and purple represents surface ancestry. The y axis shows haplotypes 1 - 38, with haplotypes 0 - 27 corresponding to Chica Pool 2 (n = 14 diploid individuals), and haplotypes 28 - 38 corresponding to Chica Pool 1 (n = 5 diploid individuals). The x axis shows bp position along each chromosome.

Extended Data Figure 8. Physical morphology in cave populations of *A. mexicanus*. A. Pachón cavefish weigh significantly less than Chica pool 1 and 2 and Los Sabinos Kruskal-Wallis test, $P < 0.01$, KW statistic = 15.19. Chica 1 vs Pachón, $p < 0.01$; Chica 2 vs Pachón, $p < 0.05$, Pachón vs Los Sabinos, $p < 0.05$. B. Body height from dorsal fin to stomach is smaller in Pachón cavefish compared to Chica Pool 1 and 2 as well as Los Sabinos. Kruskal-Wallis test, $P < 0.01$, KW statistic = 11.90. Chica 1 vs Pachón, $p < 0.05$; Pachón vs Los Sabinos, $p < 0.05$. C. Body length measured from mouth to tail is significantly smaller in Pachón cavefish compared to all other cave populations. Kruskal-Wallis test, $P < 0.001$, KW statistic = 17.84. Chica 1 vs Pachón, $p < 0.01$; Chica 2 vs Pachón, $p < 0.01$, Pachón vs Los Sabinos, $p < 0.05$. D. Body mass is significantly larger in Pachón cavefish compared to other populations Kruskal-Wallis test, $P < 0.01$, KW

1470 statistic = 14.41. Chica 1 vs Pachon, $p < 0.01$; Chica 2 vs Pachon, $p < 0.05$, Pachon vs Los

1471 Sabinos, $p < 0.05$.

1472