Original research article

Syntrophy between fermentative and purple phototrophic bacteria for carbohydrate-based wastewater treatment

Marta Cerruti¹, Guillaume Crosset-Perrotin¹, Mythili Ananth¹, Jules L. Rombouts^{1,2}
and David G. Weissbrodt^{1,*}

¹ Department of Biotechnology, Delft University of Technology, Delft, The Netherlands

² Nature's Principles B.V., Den Haag, The Netherlands

* Correspondence: Prof. David Weissbrodt, Weissbrodt Group for Environmental Life Science Engineering, Environmental Biotechnology Section, Department of Biotechnology, TNW Building 58, van der Maasweg 9, 2629 HZ Delft, The Netherlands, Tel: +31 15 27 81169; E-mail: d.g.weissbrodt@tudelft.nl

ABSTRACT

Fermentative chemoorganoheterotrophic bacteria (FCB) and purple photoorganoheterotrophic bacteria (PPB) are two interesting microbial guilds to process carbohydrate-rich wastewaters. Their interaction has been studied in axenic pure cultures or co-cultures. Little is known about their metabolic interactions in open cultures. We aimed to harness the competitive and syntrophic interactions between PPB and FCB in mixed cultures. We studied the effect of reactor regimes (batch or continuous, CSTR) and illumination modes (continuous irradiation with infrared light, dark, or light/dark diel cycles) on glucose conversions and the ecology of the process. In batch, FCB outcompeted (>80%) PPB, under both dark and infrared light conditions. In CSTR, three FCB populations of *Enterobacteriaceae*, *Lachnospiraceae* and *Clostridiaceae* were enriched (>70%), while *Rhodobacteraceae* relatives of PPB made 30% of the community. Fermentation products generated from glucose were linked to the dominant FCB. Continuous culturing at a dilution rate of 0.04 h⁻¹ helped maintain FCB and PPB in syntrophy: FCB first fermented glucose into volatile fatty acids and alcohols, and PPB grew on fermentation products. Direct supply of carboxylates like acetate under infrared light enriched for PPB (60%) independent of reactor regimes. Ecological engineering of FCB- and PPB-based biorefineries can help treat and valorize carbohydrate-based waste feedstocks.

Keywords: photoorganoheterotrophs; chemoorganoheterotrophs; syntrophy; mixed-culture fermentation; resource recovery

38 **GRAPHICAL ABSTRACT**

INTRODUCTION

In the natural environment, bacteria evolved to occupy diverse niches (Steindler et al., 2020). Microorganisms form metabolic associations to grow under nutrient-limiting conditions (Pearman et al., 2008). Anthropogenic environments like biological wastewater treatment plants (WWTPs) involve relatively high concentrations of organics (up to 200 g COD L⁻¹) and other nitrogenous and phosphorous nutrients, and complex compositions of carbon sources like cellulose and undigested fibers from food. In agri-food industrial wastewaters, most of organics are derived from simple carbohydrates (*e.g.*, glucose, fructose or xylose) and polymeric sugars such as starch, cellulose and lignocellulosic derivatives (Ghosh et al., 2017).

The wastewater treatment sector transitions to develop bioprocesses that combine nutrient removal with bioproduct valorization. Environmental biotechnologies are studied to treat and valorize nutrient-rich wastewaters by unravelling the biochemical mechanisms that can drive product formation (Kleerebezem & van Loosdrecht, 2007). The efficiency of mixed-culture processes is based on proper selection of microbial guilds needed to metabolize the wastes. Two guilds across chemo- and photoorganoheterotrophic groups are particularly attractive to process carbohydrate- and carboxylate-based feedstocks.

Fermentative chemoorganoheterotrophic bacteria (FCB) degrade carbohydrates through fermentation pathways, producing a spectrum of valuable compounds such as carboxylates (*e.g.*, volatile fatty acids – VFAs, lactic acid), ethanol and H₂. Their microbial niches are harnessed in non-axenic mixed-culture fermentation processes to valorize, *e.g.*, lignocellulosic sugars into targeted fermentation products like lactate or ethanol (Rombouts et al., 2020).

Purple phototrophic bacteria (PPB) are versatile microorganisms, involving a diversity of phototrophic and chemotrophic metabolisms depending on environmental and physiological conditions (**Table 1**). They grow preferentially as photoorganoheterotrophs, harvesting their energy from infrared (IR) light and using a variety of carbon sources from carboxylates to carbohydrates to

alcohols (Imam et al., 2011; Okubo et al., 2005; D. Puyol et al., 2017). PPB can also grow as chemotrophs using organic or inorganic substrates (Hunter et al., 2009). As photoheterotrophs, they can achieve high biomass yields up to 1 g COD_x g⁻¹ COD_s (Daniel Puyol & Batstone, 2017), that can be exploited to valorize anabolic products like biomass, single-cell proteins used in animal/human feed, or industrially relevant compounds like carotenoids.

Table 1. Metabolic versatility of purple phototrophic bacteria (PPB). PPB can grow by combining a diversity of substrates and redox conditions. PPB grow primarily as photoorganoheterotrophs using VFAs as preferential carbon sources. but they are able to grow also on glucose with low growth rates. The metabolisms targeted in this study for the conversion of glucose by combination of fermentation and photoorganoheterotrophy are highlighted in grey.

Metabolism	Energy source (photo-/chemo-)	Electron donors (organo-/litho-)	Oxidized e-donors	Carbon source (hetero-/auto-)	Electron acceptors	Reduced e-acceptors
<i>Under light conditions</i>						
Anoxygenic photoorganoheterotrophy	Light photons	Reduced organic	Oxidized organic, CO ₂	Organic	Endogenous compound, CO ₂ fixation	Biomass
Anoxygenic photolithoautotrophy	Light photons	Reduced inorganic H ₂ H ₂ S Fe ²⁺ NO ₂ ⁻ H ₂ O	Oxidized inorganic H ₂ O SO ₄ ²⁻ Fe ³⁺ NO ₃ ⁻ O ₂	Inorganic (CO ₂)	Endogenous compound, CO ₂ fixation	Biomass
Oxygenic photolithoautotrophy	Light photons	H ₂ O	O ₂	Inorganic (CO ₂)	Endogenous compound, CO ₂ fixation	Biomass
Photofermentation (not growth associated)	Light photons	Excess electrons energized from substrate	-	-	H ⁺	H ₂
<i>Under dark conditions</i>						
Aerobic-respiring chemoorganoheterotrophy	Chemical redox reaction by aerobic respiration	Reduced organic	CO ₂	Organic	O ₂	H ₂ O Biomass
Anaerobic-respiring chemoorganoheterotrophy	Chemical redox reaction by anaerobic respiration	Reduced organic	CO ₂	Organic	NO ₃ ⁻ SO ₄ ²⁻	N ₂ S ⁰ , HS ⁻ , H ₂ S Biomass
Fermentative chemoorganoheterotrophy	Chemical redox reaction by fermentation	Reduced organic (e.g., fermentable sugars)	Pyruvate or CO ₂	Organic	Endogenous compound	Fermentation products (e.g., acetate, H ₂), Biomass

77

78 Further studies evaluated PPB ability to do photofermentation, coupling glucose degradation to H₂
 79 production. Combinations of fermentative chemoorganoheterotrophy, photoorganoheterotrophy and
 80 photofermentation processes have been tested for carbohydrate degradation and conversion. In (i)
 81 *two-stage dark fermentation and photoorganoheterotrophy*, the process is carried in two separate
 82 reactors. In the first unit, pure cultures of FCB degrade carbohydrates into VFAs that are then fed in
 83 the second unit to produce PPB biomass in pure cultures (Ghimire et al., 2015). In (ii) *single-stage*
 84 *fermentation*, glucose is degraded by pure-cultures of PPB, with longer contact times compared to
 85 the two-stages fermentation and the requirement of a pre-sterilized feed stream and axenic
 86 environment (Abo-Hashesh & Hallenbeck, 2012). In (iii) *single-stage dark and photofermentation*,
 87 attempts have been made to co-cultivate defined species of FCB and PPB to simultaneously degrade
 88 carbohydrates and convert the produced VFAs to biomass (Rai & Singh, 2016). Despite the
 89 advantages of this technique, including lower investment and operation costs compared to the two-
 90 staged fermentation, it still requires an axenic environment.

91 In this context of associating fermentation and photoorganoheterotrophy, one can also wonder
 92 whether PPB, due to their metabolic versatility, could be selected in the mixed culture to perform
 93 both dark fermentation and photoorganoheterotrophy, or whether FCB would be more efficient on
 94 fermentation, prior to supplying fermentation products for the PPB to grow.

95 Open mixed cultures have the advantage, compared to traditional industrial biotechnology
 96 approaches, to not require sterilization of the inflow, therefore substantially reducing the costs
 97 (Kleerebezem & van Loosdrecht, 2007). Based on ecological selection principles, microorganisms
 98 can be selected to specifically target substrate degradation, bioproduct valorization or both.

99 Many studies have focused on H₂ production with FCB and PPB pure cultures or with co-cultures in
 100 axenic conditions. Little is known about the community dynamics in open mixed cultures. The
 101 interactions between populations and their metabolic interdependency have not been unraveled.

Here, we evaluated the ecological association of PPB and FCB in carbohydrate-fed mixed cultures. We addressed the selection mechanisms, competitive and syntrophic interactions of the two microbial guilds across reactor (batch, chemostat) and IR light irradiance (light, dark, dark/light) regimes using either glucose or acetate as model carbohydrate and carboxylate substrates that harbor the same degree of reduction ($4 \text{ mol e}^- \text{ C-mol}^{-1}$). With this microbial ecology insights, we tested the possibility to treat carbohydrate-rich wastewaters in a single-stage mixed-culture process assembling FCB and PPB in syntrophy.

MATERIAL AND METHODS

Mixed-culture bioreactor systems

To evaluate microbial competition between PPB and fermenters, two mixed-culture bioreactor regimes (batch and continuous culturing), two substrates with the same degree of reduction per carbon mole (acetate and glucose), and three light patterns (continuous illumination, continuous dark, and light / dark switches with a ratio of 12h / 8h) were applied. The reactor performances were measured by quantitative biotechnology endpoints (rates, yields). Microbial selection and community dynamics were tracked by 16S amplicon sequencing. The experimental design is shown in **Table 2**.

Table 2. Combinations of operational conditions tested to address selection, competition, and interaction mechanisms between FCB and PPB.

Reactor regime	Organic substrate	Illumination	Input concentration of organic matter (gCOD L ⁻¹)
Batch	Acetate	Continuous	10
Batch	Acetate	No (dark)	10
Batch	Glucose	Continuous	10
Batch	Glucose	No (dark)	10
CSTR	Acetate	Continuous	5 / 10
CSTR	Glucose	Continuous	5
CSTR	Acetate	16h light / 8h dark	10
CSTR	Glucose	16h light / 8h dark	10
CSTR	No carbon sources	16h light / 8h dark	-

Compositions of cultivation media

Under all conditions, the cultures were provided with a mineral medium composed of (per L): 0.14 g KH₂PO₄, 0.21 g K₂HPO₄, 1 g NH₄Cl, 2 g MgSO₄·7H₂O, 1 g CaCl₂·2H₂O, 1 g NaCl, and 2 mL of trace elements and 2 mL of vitamins solutions. The stock solution of vitamins was composed of 200 mg thiamine-HCl, 500 mg niacin, 300 mg p-amino-benzoic acid, 100 mg pyridoxine-HCl, 50 mg biotin and 50 mg vitamin B12 per liter. The trace elements solution was made of (per L): 1100 mg Na₂EDTA·2 H₂O, 2000 mg FeCl₃·6 H₂O, 100 mg ZnCl₂, 64 mg MnSO₄·H₂O, 100 mg H₃BO₃, 100 mg CoCl₂·6 H₂O, 24 mg Na₂MoO₄·2 H₂O, 16 mg CuSO₄·5 H₂O, 10 mg NiCl₂·6 H₂O and 5 mg NaSeO₃. The cultivation medium was buffered at pH 7.0 with 4 g L⁻¹ 4-(2-hydroxyethyl)-1piperazineethanesulfonic acid (HEPES). Sodium acetate and glucose were used as carbon sources,

in a concentration of 5 or 10 g COD L⁻¹ depending on experimental conditions (**Table 2**), mimicking concentrations of moderately loaded agri-food wastewaters. Acetate was provided as C₂H₃O₂·3H₂O (18.28 g L⁻¹) and glucose as C₆H₁₂O₆·H₂O (9.68 g L⁻¹).

Batches operational conditions

PPB biomass from an in-house enrichment culture grown in a sequencing batch reactor (SBR) (Cerruti et al., 2020) was used to inoculate 100-ml anaerobic bottles. The cultures were flushed with 99% argon gas to maintain anaerobic conditions (Linde, NL, >99% purity). Every batch bottle was duplicated. Half of the batch bottles were exposed to dark. The other half were exposed to IR light (> 700 nm) supplied by a halogen lamp (white light) with a power of 120 W whose wavelengths below 700 nm were filtered out by two filter sheets (Black Perspex 962, Plasticstocktist, UK). Every batch was incubated in a closed shaker (Certomat® BS1, Sartorius Stedim Biotech, Germany) at 30±1 °C under dark conditions and at 37±1 °C under light conditions: these differences in temperature were due to an uncontrolled heat derived from the lamps. The flasks were kept under agitation at 170 rpm. The cultures were fed with 10 g COD L⁻¹ of either glucose or acetate.

Continuous culture conditions

A 2.5-L reactor with 2 L of working volume connected to a controller system (In-Control and Power units, Applikon, Netherlands) was used to tune the syntrophy of FCB and PPB under continuous conditions. The pH was set at 7.0 and regulated with NaOH at 0.25 mol L⁻¹ and HCl 1 mol L⁻¹ and the temperature was maintained at 30 °C with a heat exchanger.

The reactor was irradiated from two opposite sides with 2 incandescent lights filtered for wavelengths $\lambda > 700$ nm by two sheets (Black Perspex 962, Plasticstocktist, UK) to promote PPB growth. The light intensity on the reactor wall was 270 W m⁻² and was measured with a pyranometer (CMP3, Kipp & Zonen, The Netherlands).

After start-up under batch mode and PPB enrichment with acetate, the dilution rate was set at 0.04 h^{-1} , equal to less than half of the $\mu_{\max}$ of the PPB representative genus *Rhodopseudomonas* (0.1 h^{-1}) (Cerruti et al., 2020).

Four combinations of energy patterns and carbon sources were tested: (i) continuous light with acetate, (ii) continuous light with glucose, (iii) light/ dark (16/8 h) cycles with acetate, and (iv) light/ dark (16/8 h) cycles with glucose. Light and dark alternation was controlled with an automatic switch time controlled (GAMMA, The Netherlands).

After 3 months of continuous cultivation, the carbon source feed was stopped, while phosphate and nitrogen sources were maintained for a period of 4 days (ca 4.5 HRTs), in order to evaluate the hypothesis that PPB can grow on fermentation products produced by FCB.

Mass spectrometry for off-gas analyses

The off-gas of all CSTR was connected to a mass spectrometer (ThermoFisher, Prima BT Benchtop MS) which was used to measure the production rates of CO_2 and H_2 .

Sampling

All samples were collected in 2 ml Eppendorf tubes, centrifuged at $10000 \times g$ for 3 minutes. The biomass pellet was separated from the supernatant and stored at -20°C until further analysis. The supernatant was filtered with $0.2 \mu\text{m}$ filters (Whatman, US), and stored at -20°C until further analysis. In the batch cultures, the samples were collected every 24 h. During the CSTR, 5 to 9 samples were collected for each condition.

Biomass determination

Biomass concentrations were measured over the time course of the batch and continuous reactors by spectrophotometry (Biochrom, Libra S11, US) absorbance at 660 nm (A_{660}). A calibration curve was

established to correlate A_{660} to g volatile suspended solids (gVSS) concentration: $c(\text{gVSS L}^{-1}) = 0.61 A_{660}$. Biomass gVSS was measured taking samples from the liquid phase, filtering them using 0.2 μm filters (Whatman, 435 USA) and drying the wet filters in a 105 °C stove for 24 h. The filters were then incinerated at 550 °C for 2 h, and weighted.

Amplicon sequencing

DNA was extracted from 0.5-7 mg biomass using DNeasy Powersoil microbial extraction kit (Qiagen, Hilden, Germany) accordingly to manufacturer's instructions and stored at -20°C. The compositions of the bacterial communities of the mixed liquors were characterized from the samples collected by V3-V4 16S rRNA gene-based amplicon sequencing as in (Cerruti, Stevens, et al., 2020). The fastq files provided from Novogene were analysed with QIIME2 pipeline (Bolyen et al., 2019).

Chromatography analysis of substrate conversions and fermentation products

Substrate depletion (acetate and glucose) and fermentation products formation (succinate, propionate, formate, lactate, acetate, butyrate) and were scanned and monitored using high-performance liquid chromatography (HPLC) (Waters, 2707, NL) equipped with an Aminex HPX-87H column (BioRad, USA) column. The eluent used was H_3PO_4 (1.5 mmol L^{-1}) at a flowrate of 0.6 mL min^{-1} and at a temperature of 60°C. The compounds were detected by refraction index (Waters 2414, US) and UV (210 nm, Waters 2489, US). Ethanol was quantified through gas-chromatography as in (Julius L Rombouts et al., 2019a).

Wavelength scans of biomass samples to track PPB

Wavelength scan of biomass samples were done using a spectrophotometer (DR3900, Hach, Germany) in order to evaluate the presence of the typical PPB photopigments (carotenoids at 400-

500 nm and bacteriochlorophyll at ca 850 nm). Absorbance profiles were measured at each wavelength between 320 nm and 999 nm.

Solver for pathway utilization prediction

To evaluate the fractions of metabolic fermentation and phototrophic pathways, a numerical model was set up which could estimate pathway fractions through minimizing the residual sum squared error (SSQ) using Microsoft Excel 16.48, using the following formula:

$$\min(SSQ) = \min\left(\sqrt{\sum_{i=1}^n \frac{(\hat{y}_i - y_i)^2}{n}}\right) \quad (\text{eq. 1})$$

Where $\hat{y}_i$ is the modelled yield value and y_i is the measured yield value, and n is the amount of metabolic compounds that are considered. The estimated yields were calculated using different fractions of metabolic pathways, which could be varied by the solver to minimize the SSQ (**Table 3**). This SSQ value was minimized using the Generalized Reduced Gradient (GRG) Nonlinear solver available in Excel. The pathways that were used varied per enrichment, depending on the microbial community structure, assuming only the activity of relevant catabolic and anabolic pathways. To simulate the batch enrichments the equations 1.1-1.8 were used. To simulate the CSTRs, reactions 1.9-1.10 were also added (Table 3). The result of the pathway modelling was used to create **Figure 7**, visualizing in a semi-quantitative way the flux distribution of the different pathways.

Table 3. Putative reactions included in the numerical model of metabolic reactions along FCB and PPB pathways. The reactions 1.1-1.8 were used to evaluate the contribution of each pathway under batch conditions. The reactions 1.9 and 1.10 were included in the simulation of the CSTRs.

Microbial guilds	Pathways	Reactions
FCB	Glucose $\rightarrow$ Biomass + CO ₂	(1.1)
	Glucose $\rightarrow$ Ethanol + CO ₂	(1.2)
	Glucose + H ₂ O $\rightarrow$ Acetate + Ethanol + H ⁺ + H ₂ + CO ₂	(1.3)
	Glucose $\rightarrow$ Butyrate + CO ₂ + H ₂ O	(1.4)
	Glucose $\rightarrow$ Succinate	(1.5)
	Glucose $\rightarrow$ Lactate + H ⁺	(1.6)
	Lactate $\rightarrow$ Acetate + Propionate + CO ₂ + H ₂ O	(1.7)
FCB + PPB	Formate $\rightarrow$ CO ₂ + H ₂	(1.8)
PPB	Acetate $\rightarrow$ Biomass + CO ₂	(1.9)
	CO ₂ + H ₂ O $\rightarrow$ Biomass + H ₂	(1.10)

RESULTS & DISCUSSION

The batch and continuous reactor experiments revealed key insights on the microbial community engineering principles governing the selection, competition and interaction between FCB and PPB in function of substrates, light patterns, and reactor regimes.

Batch conditions

In mixed-culture batch mode, PPB were selected on acetate and IR light

Under batch conditions, the acetate-fed cultures grew only when subjected to light, reaching a biomass concentration of 1 gVSS L⁻¹ ($\mu = 0.0411 \pm 0.0007$ h⁻¹), with a yield of biomass on substrate

239 of 0.60 ± 0.09 C-mmol $\times$ C-mmol $^{-1}$. Little growth was reported in the six days of cultivation for the
 240 acetate cultures in the dark, with a biomass specific growth rate (μ) of 0.006 h $^{-1}$.
 241 The inocula of the batches originated from a PPB in-house enrichment culture operated as SBR in
 242 parallel to the experiments. The inoculum seeded in the acetate-fed batches presented a relative
 243 abundance above 60% (vs. total amplicon sequencing read counts) of the purple non-sulfur genus
 244 *Rhodopseudomonas*. In the acetate-fed batches, *Rhodopseudomonas* sp. reached as high as 80% of
 245 the total community after 6 days of cultivation under IR light. Despite the low growth,
 246 chemoorganoheterotrophs of the family of the *Moraxellaceae* showed the highest relative abundance
 247 (80%) in the dark (**Figure 1A**).
 248 Acetate is one of the preferred substrates for PPB (Blankenship et al., 1995). It is assimilated into
 249 biomass by many PPB species, through the tricarboxylic acids cycle (Pechter et al., 2016). Acetate
 250 and IR light selected for *Rhodopseudomonas* under acetate/light conditions. Among the PPB,
 251 *Rhodopseudomonas* has the highest growth rate on acetate ($\mu_{\max} = 0.15$ h $^{-1}$) (Cerruti, Ouboter, et al.,
 252 2020) and therefore was predominant at the end of the batch cultivation. The growth rate in the
 253 acetate-fed cultures under dark conditions was 7 times lower than under light conditions, and the
 254 biomass concentration after 6 days was 270 times lower than under light conditions. Under dark
 255 conditions, acetate can only be assimilated in presence of suitable electron acceptors (such as S^0 or
 256 SO_4^{2-}) in order to produce ATP for growth (Madigan et al., 1982). In the batches, little external
 257 electron acceptor was present in the medium (3 mmol SO_4^{2-} L $^{-1}$). This can explain the very low
 258 biomass growth under dark conditions compared to the light one.

259

260 *In mixed-culture batch mode, FCB were selected on glucose*

261 In the glucose-fed batches a higher biomass concentration was reached in the dark (1 gVSS L $^{-1}$) than
 262 in the light (0.4 gVSS L $^{-1}$), with a specific μ of 0.070 ± 0.001 h $^{-1}$ and 0.020 ± 0.002 h $^{-1}$ respectively.

The inoculum of glucose-fed batches was dominated by the purple sulfur genus *Thiobaca*, due to microbial community variations in the parent reactor. In the glucose batch cultivation, the genus of *Enterobacter* was predominant, above 90% of the total sequencing reads (**Figure 1A**), both under light and dark conditions. The pH was not controlled in batch cultivations: in the glucose-fed batches pH dropped to 4 within 48 h, despite the presence of the HEPES buffer (**Figure 1B**). The *Enterobacteriaceae* family is known to be predominant in anaerobic environment in presence of non-limiting concentration of monomeric carbohydrates, such as glucose (de Vrije & Claassen, 2003) (de Vrije & Claassen, 2003). Rombouts et al., 2019 reported an enrichment of 75% of the genus *Enterobacter* in a glucose-fed SBR, in nutrient, temperature and pH conditions similar to the one here presented.

Figure 1. A: Microbial community composition in batches. PPB are dominant at the beginning of the cultivation (ACE INIT and GLU INIT). After 6 days of cultivation under acetate feed, PPB were dominant under light conditions (ACE LIGHT). Under acetate-dark conditions there was little grow from the genus *Moraxellaceae*. Under glucose feed, independently of the light conditions, the genus *Enterobacteriaceae* was dominant. **B:** pH evolution in the glucose-fed batches. Both under light and under dark conditions the pH dropped to 4 in the first 2 days of incubation.

Ethanol was the main fermentation product under glucose-fed batch conditions

Under acetate-fed batch conditions in the light, acetate was fully depleted after 6 days of incubation. Under glucose-fed batch conditions the biomass yield over glucose were 0.052 ± 0.002 C-mmolx C-mmol⁻¹ and 0.134 ± 0.011 C-mmolx C-mmol⁻¹ under light and under dark conditions respectively. After six days of cultivation, in the glucose-fed batches in the dark, ethanol was the main fermentation product, with a yield of 0.20 ± 0.10 C-mmol_{et} C-mmol⁻¹, followed by succinate (0.07 ± 0.03 C-mmol_{succ} C-mmol⁻¹) and acetate (0.03 ± 0.01 C-mmol_{ace} C-mmol⁻¹). In the glucose-fed batches under light conditions, after 6 days of cultivation 30% of the initial glucose was still present in the reaction broth, and carbon balances could not be properly closed (>90% recovery). Ethanol was the main fermentation product (0.10 ± 0.07 C-mmol_{et} C-mmol⁻¹), followed by lactate (0.05 C-mmol_{lac} C-mmol⁻¹) and succinate (0.04 ± 0.02 C-mmol_{succ} C-mmol⁻¹).

A solver enabled to evaluate the theoretical contribution of 8 putative fermentation pathways to the overall carbon and electron balances. Under batch conditions the phototrophic pathways were not included in the simulation, as PPB relative abundance was very low as measured by amplicon sequencing (~1%) (**Figure 1A**). The predicted contribution of the singular pathways to the overall yields of the batches under light and dark conditions were comparable. Under batch conditions, the dominant metabolic pathway was identified as homo-ethanol production form glucose (**Table 3**, Eq. 1.2). This catabolism is associated with the fermentative *Enterobacteriaceae* (Jang et al., 2017).

Wavelength scan is commonly used as an indirect measure for the presence of PPB in the cultures (Fiedor et al., 2016), as photopigments absorb photons with specific wavelengths. Absorbance peaks around 400-500 nm represent the carotenoids series, and the peaks around 850 nm the bacteriochlorophyll a (Stomp et al., 2007). In the acetate-based batches in the light the aforementioned peaks were clearly detected at 450 nm and 800-850 nm. In the acetate-fed batches in

the dark, almost no growth was reported in the 6 days incubation and peak for photopigments was detected at 800 and 850 nm. In the glucose-fed batches, both dark and light, no peaks were observed (Figure 2A/B). These results suggested the selective growth of PPB in illuminated acetate batches, and of non-phototrophic organisms in the glucose-batches.

Figure 2. Wavelength scan from 300 to 1000 nm. Absorbance units normalized for the biomass concentration (as A.U. at 660nm). **A:** acetate-fed batches. The typical PPB peaks (at ca 850 nm) were clearly visible under light conditions. **B:** glucose-fed batches. No peak was present. **C:** continuous light glucose-fed CSTR. The presence of PPB was confirmed by the presence of their typical peaks. **D:** dark / light glucose-fed CSTR. The presence of PPB was confirmed by the presence of their typical peaks.

Enterobacteriaceae were enriched due to higher growth rates

The maximum biomass specific growth rate ($\mu_{\max}$) is a key factor driving selection mechanisms in substrate-rich environments (Winkler et al., 2017), if no significant intermediate storage of substrate

is displayed. FCB-like species of the genus *Enterobacter* present a $\mu_{\max}$ of 0.45 – 0.80 h⁻¹ in mineral medium conditions with low sugar concentrations (<10 g_{COD} L⁻¹) (Füchslin et al., 2012; Rombouts et al., 2019). When fermenting carbohydrates like glucose, *Rhodopseudomonas capsulata* grows at a $\mu_{\max}$ of 0.014 h⁻¹ (Conrad & Schlegel, 1978). In batches, where the substrate is not limiting, the organisms with higher uptakes rates will be selected above organisms with high biomass yields (Rombouts et al 2019). According to the Herbert-Pirt relation, when the substrate uptake is directly coupled with growth, the organisms with the highest growth rate dominate. Using these studies, the genus *Enterobacter* is estimated to have a $\mu_{\max}$ about 32 times higher than *Rhodopseudomonas* under glucose feed, making it successful in the competition for glucose in batch.

Under batch conditions, both in the light and in the dark, the family of the *Enterobacteriaceae* was enriched (>90%). The species belonging to the family of the *Enterobacteriaceae* ferment sugars through either the Embden-Meyerhof pathway or the pentose phosphate pathway, with a net production of ethanol (Jang et al., 2017). Ethanol can be used as substrate for growth by PPB, but with a strict pH regulation (pH = 7) (Inui et al., 1995).

Due to lactate production, which is a major byproduct of ethanol fermentation (Converti & Perego, 2002), and insufficient pH buffer, the pH dropped to 4 within the first 2 days of cultivation, as the pK_a of lactic acid is 3.8 (**Figure 1B**). Fermentative bacteria can still grow at low pH (pH < 5.5) (Tsuji et al., 1982). PPB grow instead in a pH range between 6.5 and 9.0, with an optimum at 7.0 (Pfennig, 1977). The low pH (pH = 4) resulting from the glucose-fed batch experiments could have prevented the PPB to uptake and grow on the alcohols and VFAs produced by *Enterobacter*, leading to a low relative abundance (~ 1%) in the glucose-fed batches. We postulate that with a strict pH control to 7, PPB will be able to grow on fermentation products of *Enterobacteriaceae*.

The genus *Enterobacter* can grow in a range of temperature between 20 and 45°C (Gill & Suisted, 1978), with an optimum growth temperature at 40°C (Tanisho, 1998). The temperature differences

here reported for glucose-fed batches under dark and light (ca. 7°C difference) should not have affected its specific growth rates. On the other hand, light has been used as a control method for fermentative organism proliferation (D’Ercole et al., 2016; Gwynne & Gallagher, 2018). As IR light was provided to the illuminated cultures, we postulate its inhibitory action on the *Enterobacteriaceae* family, resulting in lower growth rates (3.5 times lower) compared to the dark cultures. How this inhibition by photons actually works on cellular and molecular level remains to be settled by future work.

Continuous culture

Under continuous-flow regime, four different light and substrate conditions were imposed (1- acetate under light, 2- acetate under light / dark cycles, 3- glucose under light, and 4- glucose under light and dark cycles). The pH was controlled at 7.0 to favor the enrichment of PPB.

CO₂ production rate increased with an increased biomass growth

The CO₂ partial pressure in the off-gas of the CSTR was monitored to evaluate the growth state of the biomass. For PPB, CO₂ production is a side product of acetate assimilation (Sadaie et al., 1997). In acetate-fed CSTRs, CO₂ production was constant under continuous illumination (0.0111 ± 0.0004 C-mmol_{CO₂} C-mmol⁻¹) (**Figure 3A**). When dark / light cycles were applied, the CO₂ production decreased to 0 C-mmol h⁻¹ during the 8 h dark periods and increased again to 0.3 C-mmol h⁻¹ during the 16 h light phases (**Figure 3B**). The CO₂ production during the light phases indicated that acetate was continuously assimilated, whereas the sudden decrease during the dark periods (1/3 less within the first hour of dark) indicated that the metabolic activity was switched off. An oscillation in the CO₂ production rate was already observed in *Rhodospseudomonas* pure cultures under CSTR (Cerruti et al., 2020). In both cases, when the substrate was provided in excess, PPB grew with a growth rate close to $\mu_{\max}$ (0.1 h⁻¹) during the light periods. In the dark, PPB halted their metabolic activity.

Both under continuous illumination and dark/light cycles, a change in CO₂ production was observed 24-48 h after the switch from acetate feed to glucose feed. In the CSTR under continuous illumination, a peak of CO₂ production reached 1.2 C-mmol h⁻¹ 24 h (ca 1 HRT) after the switch to the glucose feed and stabilized at 0.6 C-mmol h⁻¹ 2 days later (**Figure 3A**). Under light / dark cycles a peak of CO₂ production (up to 5.6 C-mmol h⁻¹) was observed after two full cycles (3.8 HRTs from the glucose switch). After 8 HRTs from the switch to the glucose feed, the CO₂ production rate presented oscillations between light and dark periods. The CO₂ production rate increased of 0.5 C-mmol h⁻¹ at the beginning of the light phase and it decreased of 0.4 C-mmol h⁻¹ in every dark period (**Figure 3B**). This behavior was observed for 8 HRTs, until the pH dropped to 2 due to a technical failure.

Figure 3. A: CO₂ production rates under acetate-fed and glucose-fed CSTR continuous light conditions. An increase of ca 1 mmol h⁻¹ was observed 3 days after the switch to glucose feed, and then stabilized. **B:** CO₂ production rates under acetate continuous light, acetate dark / light and glucose dark / light conditions. Under acetate continuous illumination CO₂ was produced at constant rate. Under acetate dark / light CO₂ was produced only during the light periods. Under glucose dark / light conditions, a peak in production rate was observed after 3 days from the switch to the glucose-feed. The production rates increased during dark periods and decreased during light periods.

Ethanol was the main fermentation product also in glucose-fed CSTR

Under glucose-fed CSTR conditions, the biomass yield over glucose was 0.22 ± 0.02 C-mmol_X C-mmol_S⁻¹ and 0.45 ± 0.12 C-mmol_X C-mmol_S⁻¹ under continuous illumination and light / dark cycles respectively. Ethanol was the main measured fermentation product (0.18 ± 0.06 and 0.11 ± 0.10 C-mmol_{et} C-mmol_S⁻¹ under continuous light and dark / light cycles respectively). In the CSTR under continuous illumination also acetate was measured (0.21 ± 0.15 C-mmol_{ace} C-mmol_S⁻¹) and butyrate (0.06 ± 0.05 C-mmol_{but} C-mmol_S⁻¹). In the CSTR under dark / light cycles, formate (0.11 ± 0.01 C-mmol_{form} C-mmol_S⁻¹) and butyrate (0.10 ± 0.01 C-mmol_{but} C-mmol_S⁻¹) were measured (**Figure 4**).

Figure 4. Yields of the glucose-fed cultures. Under batch conditions, the main fermentation products were ethanol and succinate. Under light conditions, glucose was not fully depleted. Under CSTR conditions, the main fermentation products were ethanol, formate and acetate.

Under CSTR conditions, photoautotrophic CO₂ fixation (photosynthesis) and photoheterotrophic acetate assimilation were added to the solver matrix, to evaluate the contribution of PPB to the total carbon and energy balances. Under all CSTR conditions the major fermentation pathway involved

the glucose fermentation through the acetyl-CoA pathway (**Figure 5**) (83% and 55% for continuous illumination and dark / light cycles respectively). The solver estimated a high contribution of the photoautotrophic pathway in the CSTR under continuous illumination, and of the photoheterotrophic pathway under dark / light cycles.

Figure 5. Pathways involved under the different reactor regimes. The thickness of the arrows represents the predicted yield toward a specific metabolic route, based on the calculations of the solver. The reactions with the blue background are performed by FCB. The reaction with the purple background is performed by both FCB and PPB. The reactions with the pink background are performed by PPB. **A:** batch conditions with continuous illumination. **B:** batch conditions in the dark. **C:** CSTR under continuous illumination. **D:** CSTR conditions under light / dark cycles. Under batch conditions the pathways used were mainly directed toward biomass and ethanol production. Under CSTR ethanol was the main fermentation product, but the production of propionate, succinate and lactate was predicted. The phototrophic pathways were not included for the batch conditions but were included in the carbon and electron balance of the CSTRs.

PPB were enriched in the acetate-fed CSTR

Under continuous illumination and with acetate as carbon and electron source PPB formed the dominant guild (on average 68 ± 21 %) (**Figure 6**). The genus *Rhodopseudomonas* was predominant at the beginning of the CSTR (65%), while its relative abundance decreased to 40% after one week of cultivation (8.5 HRT). Simultaneously, the genus *Rhodobacter* got enriched (from 13 to 23%). A further enrichment of *Rhodobacter* was observed once the light/dark cycles were applied, reaching around 65% of the total population after 15.6 HRTs, while *Rhodopseudomonas* abundance was stable below 10%.

In the acetate-fed CSTR under continuous light the biomass concentration was not constant. Once the dark / light cycles were applied, the biomass production followed the irradiation pattern. The biomass reached a concentration of 44.75 ± 1.24 C-mmol L⁻¹ after the light periods, and it decreased to 37.58 ± 0.09 C-mmol L⁻¹ after the dark periods. Under anaerobic conditions, PPB do not grow in the dark in absence of an external electron donor (Madigan et al., 1982; Yen & Marrs, 1977), but growth is restored once the light is present (Cerruti et al., 2020; Zhi et al., 2019). The biomass concentration increased without stabilizing in the light periods, as in (Cerruti et al., 2020). Acetate was not consumed during the dark phases, and its concentration increased due to continuous supply in the CSTR. In a chemostat, the growth rate of the microbial community is defined by the dilution rate and the concentration of the limiting compound, if no biofilm is formed (Kuenen, 2019). An increase in biomass concentration during the illumination times implied that the growth rate of the microorganisms (0.1 h⁻¹) was higher than the dilution rate (0.04 h⁻¹), as a consequence of high residual acetate concentration. During the dark cycles, acetate consumption stopped, acetate accumulated, to be used in light cycles by the PPB organisms. This a higher growth rate than the applied dilution rate, which fits with standard Monod kinetics for growth.

Figure 6. Relative abundance under CSTR conditions. Under acetate conditions, PPB were dominant community, with a shift from *Rhodospseudomonas* under continuous illumination to *Rhodobacter* under light / dark conditions. Under glucose feed, the FCB were dominant, especially the genera *Enterobacteriaceae* and *Lachnospiraceae*.

FCB were enriched in the glucose-fed CSTR

In the CSTR, FCB accounted for more of the 50% of the sequences under glucose feed, regardless the light condition. In CSTR, light and dark cycles had an impact on the selection process of the FCB. Under continuous illumination, the family of *Lachnospiraceae* was predominant (around 30 % of the total community, with a peak at 75%), followed by the family of *Enterobacteriaceae* (15%) (**Figure 6**). Under light/dark cycles *Enterobacteriaceae* was the most predominant (32 ± 17 %), along with *Clostridiaceae* (18 ± 15 %).

Among the PPB guild, *Rhodobacter* was the only PPB to significantly persist in the glucose-fed CSTR, with an average relative abundance of 19.5 ± 14 % (**Figure 6**) independently of the light conditions. The persistence of PPB in the glucose-fed CSTRs was further proven by the wavelength

scan. Peaks for the bacteriochlorophyll and the carotenoids were identified at 400-450, 800 and 850 nm for both the continuous illumination and the light / dark cycles (**Figure 2C/D**).

The microbial community defined the fermentation products production

The families of *Clostridiaceae*, *Enterobacteriaceae* and *Lachnospiraceae* are able to ferment glucose. They compete for glucose through different fermentation pathways (Grimmler et al., 2011; Horiuchi et al., 2002). *Lachnospiraceae* and *Clostridiaceae* are phylogenetically and morphologically heterogeneous families belonging to the phylum of *Firmicutes* (De Vos et al., 2005). Glucose fermentation through the acetyl-CoA pathway is typical of *Clostridiaceae* (Aristilde et al., 2015). In human gut microbiota, they contribute to sugar fermentation to lactate and short-chain fatty acids production (Venegas et al., 2019). Butyrate was produced only in the CSTR under continuous illumination (0.06 ± 0.05 C-mmol C-mmols⁻¹). The *Lachnospiraceae* and *Clostridiaceae* families ferment glucose into primarily butyrate, acetate and ethanol (Rombouts et al., 2019b; Temudo et al., 2008; Valk et al., 2018). Butyrate is produced through the butyryl-CoA dehydrogenase gene, responsible for the conversion of crotonyl-CoA to butyryl-CoA and the butyryl-CoA:acetyl-CoA transferase (Venegas et al., 2019). According to the NCBI database (NCBI, 2021), the gene encoding for this enzyme is present in the phylum of *Firmicutes* that comprises *Clostridiaceae* and *Lachnospiraceae* relatives, but not in the proteobacterial family of the *Enterobacteriaceae*, explaining the absence of butyrate in the glucose-fed batches.

Under dark / light CSTR conditions, the relatively high production of propionate (0.10 ± 0.01 C-mmol C-mmols⁻¹, 17 times higher than under continuous illumination), can be linked to the abundance of *Clostridiaceae*, which are propionate producers (Johns, 1952). *Clostridium* species can present a $\mu_{\max}$ of 0.25 h⁻¹ (Gomez-Flores et al., 2015) with primary fermentation products including ethanol, butyrate, acetate and propionate (Huang et al., 1986; Lamed et al., 1988). *Clostridium beijerincki* has been reported to decrease the H₂ production when exposed to light intensities above

200 W m⁻², with a shift from a preferential production of butyric acid to acetic acid (Zagrodnik & Laniecki, 2016). The continuous illumination under CSTR conditions might have resulted in an inhibition of *Clostridiaceae*, with an enrichment of other *Firmicutes* like *Lachnospiraceae*, through an unknown mechanism.

Formate and propionate production was 10 times higher under light/ dark conditions compared to continuous illumination. Acetate production was instead 12 times lower under dark / light cycles compared to the continuous illumination (**Figure 4**). This was probably linked to a higher abundance family of the *Enterobacteriaceae* under dark / light cycles compared to the continuous illumination (31% and 18% respectively) (**Figure 6**). *Enterobacteriaceae* present a $\mu_{\max}$ 0.5-1 h⁻¹ (Khanna et al., 2012; Martens et al., 1999) and are known to ferment sugars to primarily ethanol, with lactate and acetate as major byproduct during ethanol fermentation (Converti & Perego, 2002).

PPB grew on FCB fermentation products

After 3 months of CSTR operation, to test the hypothesis that PPB grow on the fermentation products of FCB, the glucose feed was stopped while nitrogen and phosphate were still provided. Along the 4 days of operation, the relative abundance of the PPB shifted from ca 10 to 50% (**Figure 7A**). The biomass concentration sharply decreased already 2 h after stopping the glucose feed, stabilizing at a concentration of 15.7 ± 2 C-mmolx L⁻¹. Butyrate, ethanol, formate, lactate, propionate and succinate were washed out of the system at the imposed dilution rate (0.04 h⁻¹). Acetate was depleted in 50 h, 4 times faster than the theoretical washout rate (**Figure 7B**). In absence of glucose as carbon source, the highly active fermentative organisms were not growing. The increase in the PPB relative abundance proved the interaction with FCB, showing that PPB utilize VFAs as preferred substrate and can selectively grow on FCB fermentation products.

Figure 7. A: Microbial composition after the stop to the glucose feed. Over time, the FCB relative abundance decreased over time, while the *Rhodobacteraceae* increased. **B:** Fermentation products concentration after the glucose feed-stop. After 55 h all the fermentation products were depleted.

Outlook

Metabolic strategies for microbial enrichment

Microorganisms use a combination of adaptive reactions to coexist with other organisms (Panikov, 2010). Based on the adaptation techniques, two major types of organisms can be identified (Andrews & Harris, 1986; Cowan et al., 2000): r- and K-strategists. The r-strategists show high growth and conversion rates, in low-populated and resources-rich environments. K-strategists thrive in highly populated environments, where the resources are limited. They exhibit lower growth rates but high substrate conversion yields.

Under the different reactor regimes, the microbial communities followed this postulate.

Enterobacteriaceae were predominant under non-limiting conditions can be classified as r-

organisms. *Clostridiaceae* and *Lachnospiraceae* showed high growth rates in nutrient-limiting environments and can be classified as K-strategists. Among the PPB guild, *Rhodopseudomonas* showed a high-rate behaviour, dominating in substrate-rich environments. *Rhodobacter* showed high efficiency typical of the K-organisms, being able to survive also to nutrient-limited environments, as the CSTRs (Table 4).

Table 4. Reported maximum growth rates of the main FCB and PPB in the systems under glucose and acetate feed and adaptation strategies.

Substrates	Maximum growth rate $\mu_{\max}$ (h ⁻¹)				
	Fermentative chemoorganoheterotrophic bacteria			Purple photoorganoheterotrophic bacteria	
	(FCB)			(PPB)	
	<i>Enterobacteriaceae</i>	<i>Lachnospiraceae</i>	<i>Clostridiaceae</i>	<i>Rhodopseudomonas</i>	<i>Rhodobacter</i>
Glucose	0.3 – 1	0.3	0.25	0.014	0.1
	(Hasona et al., 2004; Khanna et al., 2012)	(Soto-Martin et al., 2020)	(Gomez-Flores et al., 2015)	(Conrad & Schlegel, 1978)	(Imam et al., 2011)
Acetate	-	-	-	0.15	0.05
				(Cerruti et al., 2020)	(Kopf & Newman, 2012)
Selection strategy	r-strategist (on glucose)	K-strategist (on glucose)	K-strategist (on glucose)	r-strategist (on acetate)	K-strategist (on acetate)

Kinetic parameters define the microbial selection mechanisms in CSTRs (Kuenen, 2019). In the CSTRs, the dilution rate was relatively low (0.04 h⁻¹) compared to the growth rates of the fermentative organisms (0.5-1 h⁻¹), as it was set to retain PPB in the system. In a continuous culture, the substrate (in this case glucose) is the limiting compound, and the organisms with higher affinity for it will be predominant (Andrews & Harris, 1986). Within the guild of fermentative organisms, *Clostridiaceae*

and *Lachnospiraceae* are expected to have a lower affinity constant (K_s) and therefore have established over *Enterobacteriaceae*. The controlled pH and an appropriate dilution factor ($D = 0.04 \text{ h}^{-1}$) allowed the persistence of PPB. In particular, *Rhodobacter* was enriched in the glucose-fed CSTRs regardless the illumination conditions. We propose here a syntrophic association of FCB and PPB for glucose conversion toward CO_2 and biomass (**Figure**). Glucose was first converted into acetate, ethanol, formate and lactate, and lactate was further transformed into acetate and propionate. All the fermentation products can be potentially used by PPB for growth.

Syntrophic associations between FCB and PPB

The interaction between fermentative organisms and purple bacteria is an example of syntrophic interaction. Syntrophy has been shown in biofilms and sludges, where a mass transport of compounds occurs between organisms (Rodríguez et al., 2006). Here, we showed the establishment of a metabolic cooperativity that lead to a multispecies consortium, combining sugar degradation with VFAs assimilation. In nature, PPB are present in many aquatic environments where IR light is present, and they have been found also in anthropogenic environments such as WWTPs. They harness their ability to grow on a diverse range of compounds, creating a wide metabolic net of interaction with other organisms.

The syntrophy between FO and PPB is not only important at an ecological level but can also be exploited for wastewater treatment. The FCB are used to treat food and agricultural wastes, to degrade sugars and produce biofuels as bioethanol and hydrogen gas of VFAs (Li et al., 2015; Thapa et al., 2015, 2019). Further efforts have been put to selectively direct PPB bioproduct formation toward one or another metabolic route (Puyol et al., 2017), with success. PPB, thanks to high biomass yields (Alloul et al., 2018), harbor a major industrial potential for the production of antioxidants and single-

cell proteins. It has been proven that their biomass can be used as feed for shrimps without protein extraction (Qin et al., 2018).

Advantages of single-sludge vs. two-sludge processes for conversions of glucose by FCB and PPB

The single-step bioprocess implemented here has the advantage to simultaneously treat carbohydrate-rich wastewaters, factually removing the organic pollutants, and enriching organisms with high industrial potential, by combining the metabolic properties of FCB and PPB in one single sludge. An integrated single-stage process combining fermentation and purple photoorganoheterotrophy would allow to treat carbohydrate-rich wastewater in only one tank. The illumination conditions did not affect the syntrophy between FCB and PPB. A discontinuous illumination would reduce the operational costs compared to a continuously irradiated system (Qin et al., 2018). Understanding interactions between the FCB and PPB microbial guilds will help design efficient processes for carbohydrates-based wastewater treatment and valorization. If the aim is then to maximize and valorize the PPB biomass, a two-sludge process can be efficient to first ferment carbohydrates in the first tank selecting for FCB and then supplying the fermentation products like VFAs (whose spectrum can be engineered by mastering selection conditions) to the PPB tank.

Conclusions

FCB and PPB interact as syntrophic organisms. FCB degrade glucose into VFAs, lactate and ethanol. Acetate was assimilated into biomass by PPB. The operational conditions are crucial to establish an interaction between the guilds. FCB are more efficient fermenters than PPB (although their metabolic versatility can allow for fermentation). Under glucose-fed batch regime FCB were enriched over PPB. Under glucose-fed CSTR conditions PPB population got enriched up to 30% in syntrophic association

with FCB. An appropriate dilution rate (0.04 h^{-1}) and pH regulation to 7 enabled to enrich PPB in glucose-rich environments by enabling their metabolic coupling with FCB.

ACKNOWLEDGEMENTS

This study was funded by the start-up grant of the TU Delft Department of Biotechnology (Prof. David Weissbrodt). The authors warmly acknowledge Dirk Geerts, Zita van der Krogt and Ben Abbas for their technical assistance in fermentation and molecular biology labs at TU Delft.

CONFLICT OF INTEREST STATEMENT

The authors share no conflict of interest.

REFERENCES

- Abo-Hashesh, M., & Hallenbeck, P. C. (2012). Microaerobic dark fermentative hydrogen production by the photosynthetic bacterium, *Rhodobacter capsulatus* JP91. *International Journal of Low-Carbon Technologies*, 7(2), 97–103. <https://doi.org/10.1093/ijlct/cts011>
- Alloul, A., Ganigué, R., Spiller, M., Meerburg, F., Cagnetta, C., Rabaey, K., & Vlaeminck, S. E. (2018). Capture–Ferment–Upgrade: A Three-Step Approach for the Valorization of Sewage Organics as Commodities. *Environmental Science & Technology*, 52(12), 6729–6742 <https://doi.org/10.1021/acs.est.7b05712>
- Andrews, J. H., & Harris, R. F. (1986). r- and K-Selection and Microbial Ecology. *Advances in Microbial Ecology* (pp. 99–147). https://doi.org/10.1007/978-1-4757-0611-6_3
- Aristilde, L., Lewis, I. A., Park, J. O., & Rabinowitz, J. D. (2015). Hierarchy in pentose sugar metabolism in *Clostridium acetobutyricum*. *Applied and Environmental Microbiology*, 81(4), 1452–1462. <https://doi.org/10.1128/AEM.03199-14>
- Blankenship, R., Madigan, M., & Bauer, C. (1995). *Anoxygenic Photosynthetic Bacteria* (R. E. Blankenship, M. T. Madigan, & C. E. Bauer (eds.); Vol. 2). *Springer Netherlands*. <https://doi.org/10.1007/0-306-47954-0>
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, C., Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome

612 data science using QIIME 2. *Nature Biotechnology* 37(8), 852–857.
613 <https://doi.org/10.1038/s41587-019-0209-9>

614 Cerruti, M., Ouboter, H. T., Chasna, V., M van Loosdrecht, M. C., & Weissbrodt, D. G. (2020).
615 Effects of light / dark diel cycles on the photoorganoheterotrophic metabolism of
616 *Rhodopseudomonas palustris* for differential electron allocation to PHAs and H₂. *bioRxiv*
617 2020.08.19.258533. <https://doi.org/10.1101/2020.08.19.258533>

618 Cerruti, M., Stevens, B., Ebrahimi, S., Alloul, A., Vlaeminck, S. E., & Weissbrodt, D. G. (2020).
619 Enriching and aggregating purple non-sulfur bacteria in an anaerobic sequencing-batch
620 photobioreactor for nutrient capture from wastewater. *Frontiers in Bioengineering and*
621 *Biotechnology*, 8, 1432. <https://doi.org/10.1101/2020.01.08.899062>

622 Conrad, R., & Schlegel, H. G. (1978). Regulation of Glucose, Fructose and Sucrose Catabolism in
623 *Rhodopseudomonas capsulata*. *Journal of General Microbiology*, 105(2), 315–322.
624 <https://doi.org/10.1099/00221287-105-2-315>

625 Converti, A., & Perego, P. (2002). Use of carbon and energy balances in the study of the anaerobic
626 metabolism of *Enterobacter aerogenes* at variable starting glucose concentrations. *Applied*
627 *Microbiology and Biotechnology*, 59(2–3), 303–309. [https://doi.org/10.1007/s00253-002-](https://doi.org/10.1007/s00253-002-1009-5)
628 1009-5

629 Cowan, S. E., Gilbert, E., Liepmann, D., Keasling, J. D., Bioengineering Graduate Program, J., &
630 Francisco, S. (2000). Commensal Interactions in a Dual-Species Biofilm Exposed to Mixed
631 Organic Compounds. In *Applied and Environmental Microbiology* (Vol. 66, Issue 10).
632 <http://aem.asm.org/>

633 D’Ercole, S., Spoto, G., Trentini, P., Tripodi, D., & Petrini, M. (2016). In vitro inactivation of
634 *Enterococcus faecalis* with a led device. *Journal of Photochemistry and Photobiology B:*
635 *Biology*, 160, 172–177. <https://doi.org/10.1016/j.jphotobiol.2016.04.015>

636 De Vos, P., Garrity, G., Jones, D., Krieg, N. R., Ludwig, W., Rainey, F. A., Schleifer, K.-H., &
637 Whitman, W. B. (2005). The Firmicutes. In D. J. Brenner, N. R. Krieg, & J. T. Staley (Eds.),
638 *Bergey’s Manual® of Systematic Bacteriology*. Springer US. [https://doi.org/10.1007/0-387-](https://doi.org/10.1007/0-387-29298-5)
639 29298-5

640 de Vrije, G. J., & Claassen, P. A. M. (2003). Bio-methane & bio-hydrogen : status and perspectives
641 of biological methane and hydrogen production (J. H. Reith & R. H. Wijffels (eds.)). *Dutch*
642 *Biological Hydrogen Foundation*.

643 Fuchsli, H. P., Schneider, C., & Egli, T. (2012). In glucose-limited continuous culture the
644 minimum substrate concentration for growth, S_{min}, is crucial in the competition between the
645 enterobacterium *Escherichia coli* and *Chelatobacter heintzii*, an environmentally abundant
646 bacterium. *The ISME Journal*, 6(4), 777–789. <https://doi.org/10.1038/ismej.2011.143>

647 Ghimire, A., Frunzo, L., Pirozzi, F., Trably, E., Escudie, R., Lens, P. N. L., & Esposito, G. (2015).
648 A review on dark fermentative biohydrogen production from organic biomass: Process

parameters and use of by-products. *Applied Energy*, 144, 73–95.
<https://doi.org/10.1016/j.apenergy.2015.01.045>

Ghosh, S., Dairkee, U. K., Chowdhury, R., & Bhattacharya, P. (2017). Hydrogen from food processing wastes via photofermentation using Purple Non-sulfur Bacteria (PNSB) – A review. *Energy Conversion and Management*, 141, 299–314.
<https://doi.org/10.1016/j.enconman.2016.09.001>

Gill, C. O., & Suisted, J. R. (1978). The effects of temperature and growth rate on the proportion of unsaturated fatty acids in bacterial lipids. *Journal of General Microbiology*, 104(1), 31–36.
<https://doi.org/10.1099/00221287-104-1-31>

Gomez-Flores, M., Nakhla, G., & Hafez, H. (2015). Microbial kinetics of *Clostridium termitidis* on cellobiose and glucose for biohydrogen production. *Biotechnology Letters*, 37(10), 1965–1971.
<https://doi.org/10.1007/s10529-015-1891-4>

Grimmler, C., Janssen, H., Krauße, D., Fischer, R. J., Bahl, H., Dürre, P., Liebl, W., & Ehrenreich, A. (2011). Genome-wide gene expression analysis of the switch between acidogenesis and solventogenesis in continuous cultures of *Clostridium acetobutylicum*. *Journal of Molecular Microbiology and Biotechnology*, 20(1), 1–15. <https://doi.org/10.1159/000320973>

Gwynne, P. J., & Gallagher, M. P. (2018). Light as a Broad-Spectrum Antimicrobial. *Frontiers in Microbiology*, 9, 119. <https://doi.org/10.3389/fmicb.2018.00119>

Hasona, A., Kim, Y., Healy, F. G., Ingram, L. O., & Shanmugam, K. T. (2004). Pyruvate Formate Lyase and Acetate Kinase Are Essential for Anaerobic Growth of *Escherichia coli* on Xylose. *Journal of Bacteriology*, 186(22), 7593–7600. <https://doi.org/10.1128/JB.186.22.7593-7600.2004>

Horiuchi, J. I., Shimizu, T., Tada, K., Kanno, T., & Kobayashi, M. (2002). Selective production of organic acids in anaerobic acid reactor by pH control. *Bioresource Technology*, 82(3), 209–213. [https://doi.org/10.1016/S0960-8524\(01\)00195-X](https://doi.org/10.1016/S0960-8524(01)00195-X)

Huang, L. I., Forsberg, C. W., & Gibbins, L. N. (1986). Influence of External pH and Fermentation Products on *Clostridium acetobutylicum* Intracellular pH and Cellular Distribution of Fermentation Products. *Applied and Environmental Microbiology*, 51(6), 1230–4.
<https://doi.org/10.1128/aem.51.6.1230-1234.1986>

Hunter, C. N., Beatty, F. D., Thurnauer, M. C., & Thomas, J. (2009). The Purple Phototrophic Bacteria. Springer Netherlands. 1013 pp. <https://doi.org/10.1007/978-1-4020-8815-5>

Imam, S., Yilmaz, S., Sohmen, U., Gorzalski, A. S., Reed, J. L., Noguera, D. R., & Donohue, T. J. (2011). IRsp1095: A genome-scale reconstruction of the *Rhodobacter sphaeroides* metabolic network. *BMC Systems Biology*, 5(July). <https://doi.org/10.1186/1752-0509-5-116>

Inui, M., Momma, K., Matoba, R., Ikuta, M., Yamagata, H., & Yukawa, H. (1995). Characterization of alcohol-assimilating photosynthetic purple non-sulfur bacteria and cloning

685 of molecular chaperones from a purple non-sulfur bacterium. *Energy Conversion and*
686 *Management*, 36(6–9), 767–770. [https://doi.org/10.1016/0196-8904\(95\)00117-V](https://doi.org/10.1016/0196-8904(95)00117-V)

687 Jang, J. W., Jung, H. M., Im, D. K., Jung, M. Y., & Oh, M. K. (2017). Pathway engineering of
688 *Enterobacter aerogenes* to improve acetoin production by reducing by-products formation.
689 *Enzyme and Microbial Technology*, 106, 114–118.
690 <https://doi.org/10.1016/j.enzmictec.2017.07.009>

691 Johns, A. T. (1952). The Mechanism of Propionic Acid Formation by *Clostridium propionicum*.
692 *Journal of General Microbiology*, 6(1–2), 123–127. [https://doi.org/10.1099/00221287-6-1-2-](https://doi.org/10.1099/00221287-6-1-2-123)
693 123

694 Khanna, N., Kumar, K., Todi, S., & Das, D. (2012). Characteristics of cured and wild strains of
695 *Enterobacter cloacae* IIT-BT 08 for the improvement of biohydrogen production.
696 *International Journal of Hydrogen Energy*, 37(16), 11666–11676.
697 <https://doi.org/10.1016/j.ijhydene.2012.05.051>

698 Kleerebezem, R., & van Loosdrecht, M. C. (2007). Mixed culture biotechnology for bioenergy
699 production. *Current Opinion in Biotechnology*, 18(3), 207–212.
700 <https://doi.org/10.1016/j.copbio.2007.05.001>

701 Kopf, S. H., & Newman, D. K. (2012). Photomixotrophic growth of *Rhodobacter capsulatus*
702 SB1003 on ferrous iron. *Geobiology*, 10(3), 216–222. [https://doi.org/10.1111/j.1472-](https://doi.org/10.1111/j.1472-4669.2011.00313.x)
703 4669.2011.00313.x

704 Kuenen, G. J. (2019). Continuous cultures (chemostats). In *Encyclopedia of Microbiology* (pp. 743–
705 761). Elsevier. <https://doi.org/10.1016/B978-0-12-801238-3.02490-9>

706 Lamed, R. J., Lobos, J. H., & Su, T. M. (1988). Effects of Stirring and Hydrogen on Fermentation
707 Products of *Clostridium thermocellum*. *Applied and Environmental Microbiology*, 54(5),
708 1216–1221. <https://doi.org/10.1128/aem.54.5.1216-1221.1988>

709 Li, L., Li, K., Wang, Y., Chen, C., Xu, Y., Zhang, L., Han, B., Gao, C., Tao, F., Ma, C., & Xu, P.
710 (2015). Metabolic engineering of *Enterobacter cloacae* for high-yield production of
711 enantiopure (2R,3R)-2,3-butanediol from lignocellulose-derived sugars. *Metabolic*
712 *Engineering*, 28, 19–27. <https://doi.org/10.1016/j.ymben.2014.11.010>

713 Madigan, M., Cox, J. C., & Gest, H. (1982). Photopigments in *Rhodospseudomonas capsulata* cells
714 grown anaerobically in darkness. *Journal of Bacteriology*, 150(3), 1422–1429.
715 <https://doi.org/10.1128/JB.150.3.1422-1429.1982>

716 Martens, D. E., Béal, C., Malakar, P., Zwietering, M. H., & Van't Riet, K. (1999). Modelling the
717 interactions between *Lactobacillus curvatus* and *Enterobacter cloacae*: I. Individual growth
718 kinetics. *International Journal of Food Microbiology*, 51(1), 53–65.
719 [https://doi.org/10.1016/S0168-1605\(99\)00095-1](https://doi.org/10.1016/S0168-1605(99)00095-1)

720 NCBI. (n.d.). *Gene [Internet]. Bethesda (MD): National Library of Medicine (US), National Center*

721 for *Biotechnology Information*; 2004 – [cited 2021 05 13]. Available from:
722 <https://www.ncbi.nlm.nih.gov/gene/No Title>.

723 Okubo, Y., Futamata, H., & Hiraishi, A. (2005). Distribution and Capacity for Utilization of Lower
724 Fatty Acids of Phototrophic Purple Nonsulfur Bacteria in Wastewater Environments. *Microbes*
725 *and Environments*, 20(3), 135–143. <https://doi.org/10.1264/jsme2.20.135>

726 Panikov, N. S. (2010). Microbial Ecology. In L. K. Wang, V. Ivanov, & J.-H. Tay (Eds.),
727 *Environmental Biotechnology* (pp. 121–191). Humana Press. [https://doi.org/10.1007/978-1-](https://doi.org/10.1007/978-1-60327-140-0_4)
728 60327-140-0_4

729 Pearman, P. B., Guisan, A., Broennimann, O., & Randin, C. F. (2008). Niche dynamics in space
730 and time. *Trends in Ecology and Evolution*, 23(3), 149–158.
731 <https://doi.org/10.1016/j.tree.2007.11.005>

732 Pechter, K. B., Gallagher, L., Pyles, H., Manoil, C. S., & Harwood, C. S. (2016). Essential genome
733 of the metabolically versatile alphaproteobacterium *Rhodopseudomonas palustris*. *Journal of*
734 *Bacteriology*, 198(5), 867–76. <https://doi.org/10.1128/JB.00771-15>

735 Pfennig, N. (1977). Phototrophic green and purple bacteria: a comparative, systematic survey.
736 *Annual review of microbiology*, 31, 275–290.
737 <https://doi.org/10.1146/annurev.mi.31.100177.001423>

738 Puyol, D., Barry, E. M., Hülsen, T., & Batstone, D. J. (2017). A mechanistic model for anaerobic
739 phototrophs in domestic wastewater applications: Photo-anaerobic model (PAnM). *Water*
740 *Research*, 116, 241–253. <https://doi.org/10.1016/j.watres.2017.03.022>

741 Puyol, Daniel, & Batstone, D. J. (2017). Editorial: Resource Recovery from Wastewater by
742 Biological Technologies. *Frontiers in Microbiology*, 8, 998.
743 <https://doi.org/10.3389/fmicb.2017.00998>

744 Qin, C., Lei, Y., & Wu, J. (2018). Light/dark cycle enhancement and energy consumption of tubular
745 microalgal photobioreactors with discrete double inclined ribs. *Bioresources and*
746 *Bioprocessing*, 5(1). <https://doi.org/10.1186/s40643-018-0214-8>

747 Rai, P. K., & Singh, S. P. (2016). Integrated dark- and photo-fermentation: Recent advances and
748 provisions for improvement. *International Journal of Hydrogen Energy*, 41(44), 19957–
749 19971. <https://doi.org/10.1016/j.ijhydene.2016.08.084>

750 Rodríguez, J., Kleerebezem, R., Lema, J. M., & Van Loosdrecht, M. C. M. (2006). Modeling
751 product formation in anaerobic mixed culture fermentations. *Biotechnology and*
752 *Bioengineering*, 93(3), 592–606. <https://doi.org/10.1002/bit.20765>

753 Rombouts, Julius L, Mos, G., Weissbrodt, D. G., Kleerebezem, R., & Van Loosdrecht, M. C. M.
754 (2019a). Diversity and metabolism of xylose and glucose fermenting microbial communities in
755 sequencing batch or continuous culturing. *FEMS Microbiology Ecology*, 95(2), 233.
756 <https://doi.org/10.1093/femsec/fiy233>

- 757 Rombouts, Julius L, Mos, G., Weissbrodt, D. G., Kleerebezem, R., & Van Loosdrecht, M. C. M.
758 (2019b). The impact of mixtures of xylose and glucose on the microbial diversity and
759 fermentative metabolism of sequencing-batch or continuous enrichment cultures. *FEMS*
760 *Microbiology Ecology*, 95, 112. <https://doi.org/10.1093/femsec/fiz112>
- 761 Rombouts, Julius Laurens, Kranendonk, E. M. M., Regueira, A., Weissbrodt, D. G., Kleerebezem,
762 R., & Loosdrecht, M. C. M. (2020). Selecting for lactic acid producing and utilising bacteria in
763 anaerobic enrichment cultures. *Biotechnology and Bioengineering*, 117(5), 1281–1293.
764 <https://doi.org/10.1002/bit.27301>
- 765 Sadaie, M., Nagano, T., Suzuki, T., Shinoyama, H., & Fujii, T. (1997). Some Properties and
766 Physiological Roles of Phosphoenolpyruvate Carboxylase in *Rhodopseudomonas* sp. No. 7
767 Grown on Ethanol under Anaerobic-light Conditions. *Bioscience, Biotechnology and*
768 *Biochemistry*, 61(4), 625–630. <https://doi.org/10.1271/bbb.61.625>
- 769 Soto-Martin, E. C., Warnke, I., Farquharson, F. M., Christodoulou, M., Horgan, G., Derrien, M.,
770 Faurie, J.-M., Flint, H. J., Duncan, S. H., & Louis, P. (2020). Vitamin Biosynthesis by Human
771 Gut Butyrate-Producing Bacteria and Cross-Feeding in Synthetic Microbial Communities.
772 *mBio*, 11(4), e00886-20. <https://doi.org/10.1128/mBio.00886-20>
- 773 Steindler, L., Rubin-Blum, M., Lavy, A., Deines, P., Hammerschmidt, K., & Bosch, T. C. G.
774 (2020). Exploring the Niche Concept in a Simple Metaorganism. *Frontiers in Microbiology*,
775 11, 1942. <https://doi.org/10.3389/fmicb.2020.01942>
- 776 Stomp, M., Huisman, J., Stal, L. J., & Matthijs, H. C. P. (2007). Colorful niches of phototrophic
777 microorganisms shaped by vibrations of the water molecule. *The ISME Journal*, 1(4), 271–
778 282. <https://doi.org/10.1038/ismej.2007.59>
- 779 Tanisho, S. (1998). Hydrogen Production by Facultative Anaerobe *Enterobacter aerogenes*. In O. R.
780 Zaborsky, J. R. Benemann, T. Matsunaga, J. Miyake, & A. San Pietro (Eds.), *BioHydrogen*
781 (pp. 273–279). Springer US. https://doi.org/10.1007/978-0-585-35132-2_35
- 782 Temudo, M. F., Muyzer, G., Kleerebezem, R., & Van Loosdrecht, M. C. M. (2008). Diversity of
783 microbial communities in open mixed culture fermentations: Impact of the pH and carbon
784 source. *Applied Microbiology and Biotechnology*, 80(6), 1121–1130.
785 <https://doi.org/10.1007/s00253-008-1669-x>
- 786 Thapa, L. P., Lee, S. J., Park, C., & Kim, S. W. (2019). Metabolic engineering of *Enterobacter*
787 *aerogenes* to improve the production of 2,3-butanediol. *Biochemical Engineering Journal*,
788 143, 169–178. <https://doi.org/10.1016/j.bej.2018.12.019>
- 789 Thapa, L. P., Lee, S. J., Yang, X., Lee, J. H., Choi, H. S., Park, C., & Kim, S. W. (2015). Improved
790 bioethanol production from metabolic engineering of *Enterobacter aerogenes* ATCC 29007.
791 *Process Biochemistry*, 50(12), 2051–2060. <https://doi.org/10.1016/j.procbio.2015.09.007>
- 792 Tsuji, A., Kaneko, Y., Takahashi, K., Ogawa, M., & Goto, S. (1982). The Effects of Temperature
793 and pH on the Growth of Eight Enteric and Nine Glucose Non-Fermenting Species of Gram-

794 Negative Rods. *Microbiology and Immunology*, 26(1),15-24. <https://doi.org/10.1111/j.1348->
795 0421.1982.tb00149.x

796 Valk, L. C., Frank, J., de la Torre-Cortés, P., van 't Hof, M., van Maris, A. J. A., Pronk, J. T., & van
797 Loosdrecht, M. C. M. (2018). Galacturonate metabolism in anaerobic chemostat enrichment
798 cultures: Combined fermentation and acetogenesis by the dominant sp. nov. “*Candidatus*
799 Galacturonibacter soehngenii.” *Applied and Environmental Microbiology*, 84(18), e01370-18.
800 <https://doi.org/10.1128/AEM.01370-18>

801 Venegas, D. P., De La Fuente, M. K., Landskron, G., González, M. J., Quera, R., Dijkstra, G.,
802 Harmsen, H. J. M., Faber, K. N., & Hermoso, M. A. (2019). Short chain fatty acids (SCFAs)
803 mediated gut epithelial and immune regulation and its relevance for inflammatory bowel
804 diseases. *Frontiers in Immunology*, 10, 277. <https://doi.org/10.3389/fimmu.2019.00277>

805 Winkler, M.-K. H., Boets, P., Hahne, B., Goethals, P., & Volcke, E. I. P. (2017). Effect of the
806 dilution rate on microbial competition: r-strategist can win over k-strategist at low substrate
807 concentration. *PLOS One*, 12(3), e0172785. <https://doi.org/10.1371/journal.pone.0172785>

808 Yen, H. C., & Marrs, B. (1977). Growth of *Rhodopseudomonas capsulata* under anaerobic dark
809 conditions with dimethyl sulfoxide. *Archives of Biochemistry and Biophysics*, 181(2), 411–
810 418. [https://doi.org/10.1016/0003-9861\(77\)90246-6](https://doi.org/10.1016/0003-9861(77)90246-6)

811 Zagrodnik, R., & Laniecki, M. (2016). An unexpected negative influence of light intensity on
812 hydrogen production by dark fermentative bacteria *Clostridium beijerinckii*. *Bioresource*
813 *Technology*, 200, 1039–1043. <https://doi.org/10.1016/j.biortech.2015.10.049>

814 Zhi, R., Yang, A., Zhang, G., Zhu, Y., Meng, F., & Li, X. (2019). Effects of light-dark cycles on
815 photosynthetic bacteria wastewater treatment and valuable substances production. *Bioresource*
816 *Technology*, 274, 496–501. <https://doi.org/10.1016/j.biortech.2018.12.021>

817

818