Functional or vestigial? The genomics of the pineal gland in *Xenarthra*

Raul Valente^{1,2}, Filipe Alves^{3,4}, Isabel Sousa Pinto^{1,2}, Raquel Ruivo¹, L. Filipe C. Castro^{1,2}

¹CIMAR/CIIMAR - Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Avenida General Norton de Matos, S/N, 4450-208 Matosinhos, Portugal.

²FCUP - Department of Biology, Faculty of Sciences, University of Porto (U. Porto), Rua do Campo Alegre, Porto, Portugal.

³MARE - Marine and Environmental Sciences Centre, ARDITI, Madeira, Portugal

⁴OOM - Oceanic Observatory of Madeira, Funchal, Portugal

*Correspondence to: filipe.castro@ciimar.up.pt, +351 223401800, ORCID ID: 0000-0001-7697-386X

Abstract

Vestigial organs are historical echoes of past phenotypes. Determining whether a specific organ constitutes a functional or vestigial structure can be a challenging task, given that distinct levels of atrophy may arise between and within lineages. The mammalian pineal gland, an endocrine organ involved in melatonin biorhythmicity, represents a classic example, often yielding contradicting anatomical observations. In Xenarthra (sloths, anteaters and armadillos), a peculiar mammalian order, the presence of a distinct pineal organ was clearly observed in some species (i.e. Linnaeus's two-toed sloth) but undetected in other closely related species (i.e. brown-throated sloth). In the nine-banded armadillo, contradicting evidence supports either functional or vestigial scenarios. Thus, to untangle the physiological status of the pineal gland in Xenarthra, we used a genomic approach to investigate the evolution of the gene hub responsible for melatonin synthesis and signaling. We show that both synthesis and signaling compartments are eroded and were lost independently. Additionally, by expanding our analysis to 157 mammal genomes we offer a comprehensive view showing that species with very distinctive habitats and lifestyles have convergently evolved a similar phenotype: Cetacea, Pholidota, Dermoptera, Sirenia and Xenarthra. Our findings suggest that the recurrent inactivation of melatonin genes correlates with pineal atrophy, and endorse the use of genomic analyses to ascertain the physiological status of suspected vestigial structures.

Keywords: Gene Loss; Vestigiality; Pineal Gland; Xenarthra

1. Introduction

Understanding the evolution of organ reduction, atrophy or indeed complete loss is a fascinating quest, dating back to the seminal work of Charles Darwin, *On the Origin of Species* (Darwin, 1859). Yet, to identify a structure as vestigial, described as a trait with no function, operating sub-optimally, or even with a modified function from that originally served, is no easy undertaking (Werth, 2014; Allmon & Ross, 2018): often yielding contradictory anatomical descriptions (e.g., Jacob et al., 2000; Nweeia et al., 2012). The increasing availability of whole genome sequences, on the other hand, provides novel tools to untangle genomic signatures impacting organ reduction or loss (e.g. Zoonomia Consortium, 2020). A key question is thus to understand how genomic changes impact these processes. Among such signatures we find, more commonly than initially anticipated, gene loss episodes: such as in the morphological simplification of the urochordate *Oikopleura dioica*, the eye regression observed in cave-dwelling populations of the teleost *Astyanax mexicanus*, the loss of gastric glands in disparate vertebrate species or the loss of sebaceous glands in some mammalian lineages (e.g., Olson, 1999; Castro et al., 2014; Albalat & Cañestro, 2016; Cronk, 2009, Guijarro-Clarke et al. 2020; McGaugh et al., 2014; Springer et al. 2018; Lopes-Marques et. a 2019a; Themudo et al., 2020; Springer et al., 2021).

A remarkable example of inconsistent observations in the functionality versus vestigiality of an organ can be found in the anatomical observations of the pineal gland in mammals (Ralph, 1975). The pineal gland is a small endocrine organ present in the brain and playing a central role in the development of entrainment behaviors through the action of melatonin (circadian rhythmicity). From a physiological standpoint, melatonin synthesis occurs in a specialized cell type, the pinealocyte, through an enzymatic cascade involving the arylalkylamine N-acetyltransferase (*Aanat*) and N-acetylserotonin methyltransferase (*Asmt*) enzymes (Klein et al., 1997; Simonneaux & Ribelayga, 2003); subsequent signaling uses a set of high affinity receptors, *Mtnr1A* and *Mtnr1B*, involved in the response of the clock machinery to melatonin stimulation, leading to local and overt phase shifts (Figure 1) (Axelrod et al., 1964; Lewy et al., 1980; Reppart et al., 1996). Although anatomical studies clearly support a well-defined pineal gland in most mammals, in lineages such as cetaceans, mole rats and sirenians a true pineal gland seems to be absent; yet, some equivocal

observations exist, ranging from complete absence to detectable presence of this gland in some species or in individuals within a species (Ralph, 1975; Ralph et al., 1985; Kim et al., 2011; Panin et al., 2012). Conflicting evidence reporting measurable levels of circulating melatonin (i.e. bottlenose dolphin) shed further doubt (Panin, 2012). Interestingly, gene loss signatures were identified in these lineages, supporting the loss-of-function of melatonin synthesis, a hallmark of pineal function, and/or signalling (Fang et al., 2014; Huelsmann et al., 2019; Lopes-Marques et al., 2019b), further demonstrating the power of genome analysis towards the clarification of organ function. The presence of a functional pineal gland is also contentious in Xenarthrans (armadillos, anteaters and sloths), a relatively understudied taxonomic group characterized by its intriguing nature (Figure 1; Oksche, 1965; Benítez et al., 1994; Superina & Loughry, 2015; Freitas et al., 2019) and representing one of the earliest-branching clades of placental mammals (Murphy et al., 2007; O’Leary et al., 2013). Xenarthrans are considered *imperfect* homeotherms, given their poor ability to adjust body temperature (Mc Nab, 1979; 1980; 1985). This inaptitude for thermal regulation, possibly related with their low metabolic rate and low energetic content diet, makes Xenarthrans’ activity patterns highly affected by air temperature, with potential effects in their circadian cycles (Chiarello, 1998; Giné et al., 2015; Maccarini et al., 2015; Di Blanco et al., 2017). While a recent report clearly identified pineal glands in the six-banded armadillo (*Euphractus sexcinctus*), Linnaeus’s two-toed sloth (*Choloepus didactylus*), and in the southern tamandua (*Tamandua tetradactyla*), a distinct pineal was not found or was reported missing in species such as southern long-nosed armadillo (*Dasypus hybridus*), pale-throated sloth (*Bradypus tridactylus*), giant anteater (*Myrmecophaga tridactyla*) or big hairy armadillo (*Chaetophractus villosus*) (Benítez et al., 1994; Ferrari, 1998; Freitas 2019). However, in the nine-banded armadillo (*Dasypus novemcinctus*) inconsistent reports advocate for either the presence or absence of a genuine pineal gland (Harlow et al., 1981; Freitas et al., 2019). Also, variable concentrations of circulating serum melatonin during the 24 h day–night cycle have been detected in this species, raising the hypothesis of an extrapineal source for melatonin production (Figure 1; Harlow et al., 1981; 1982). With the emergence of various whole-genome sequences from Pilosa (sloths and anteaters) (e.g., Uliano-Silva et al., 2019) and Cingulata (armadillos) (e.g., Lindblad-

Toh et al., 2011), Yin (et al., 2021) have recently reported the molecular erosion of *Aanat* in *Xenarthra*; yet, no attempt was made to expand this analysis to the full melatonin-related gene hub. Thus, we are now able to interrogate whether the gene repertoire of circadian rhythmicity is modified in this lineage and clarify the physiological status of the pineal gland within this group.

B

Order	Melatonin Blood levels	Pineal Gland Anatomy	Habits
Cingulata	Circadian	Inconclusive	Cathemeral
Pilosa	No information	Inconclusive	Cathemeral

Figure 1: Melatonin synthesis and signaling. Melatonin, generally described as a phase marker of the circadian clock, is initially synthesized from tryptophan which is converted into serotonin (Pévet, 2002). The final steps of this synthetic pathway include a two-step metabolization of the intermediate serotonin into melatonin, a process catalyzed by *Aanat* and *Asmt* (Klein et al., 1997; Simonneaux & Ribelayga, 2003). In mammals, *Mtnr1a* and *Mtnr1b* receptors, are involved in the response of the clock machinery to melatonin stimulation (Reppert et al., 1996) (A). Summary of the available information

regarding Xenarthran's melatonin levels, pineal gland presence and habits (B). (Illustrations used elements from Servier Medical Art: <https://smart.servier.com/>)

2. Material and Methods

2.1 Sequence collection

To clarify the functional status of *Aanat*, *Asmt*, *Mtnr1a* and *Mtnr1b* in 8 Xenarthran species (Supplementary Data 1), the genomic loci were retrieved for gene annotation using three strategies (e.g. Alves et al., 2019; Lopes-Marques et al., 2019b): a) for species with annotated genes, the genomic sequence of the target gene (ranging from the upstream to the downstream flanking genes) was collected directly from NCBI; b) for species with annotated genomes but lacking the annotation of the target gene, the genomic region between two conserved flanking genes (downstream and upstream) was directly collected and c) for unannotated genomes, blastn searches were performed, using as query a set of three genes, including *Homo sapiens* (human) target gene coding sequence (CDS), as well as those of the flanking genes in the same species. From the blast results, the best matching genome scaffold corresponding to the consensus hit across those obtained per each query sequence was retrieved. When no consensual blast hit was obtained, all hits corresponding to the *H. sapiens* CDS query were inspected, the aligning regions submitted to a back-blast search against the nucleotide database of NCBI, with the matching genomic sequence corresponding to the gene of interest being the one selected (when existing). When several matchings were found, the best genomic scaffold (yielding the highest query coverage and identity value) was collected for annotation.

For the 156 non-Xenarthran mammals with annotated genomes (Supplementary Data 1), the first two strategies described above were adopted to obtain the genomic region corresponding to the target gene. In Dugong (*Dugong dugon*), since no annotation is currently available, the genomic sequence containing the target gene was retrieved via blastn searches.

2.2 Gene Annotation and Mutational Validation

The open reading frames of the mammalian orthologues of *Aanat*, *Asmt*, *Mtnr1a* and *Mtnr1b* were investigated using PseudoChecker (pseudochecker.ciimar.up.pt), an

online platform suitable for gene inactivation inference (Alves et al., 2020). For this purpose, the human gene orthologue was used as a comparative coding sequence input (NCBI Accession ID regarding human *Aanat*: NM_001088.3; *Asmt*: NM_001171039.1; *Mtnr1a*: NM_005958.4; *Mtnr1b*: NM_005959.5) to deduce the coding status of a given candidate gene in each target species. By making use of PseudoIndex - a user assistant metric built into the PseudoChecker pipeline that rapidly estimates the erosion condition of the tested genes – putative ORFs of the orthologous gene from each target species were assigned a discrete value from 0 to 5, with 0 suggesting a fully functional gene and 5 complete inactivation (Alves et al., 2020). When PseudoIndex was higher than 2, we proceeded to manual annotation and validation of possible disrupting mutations as previously described by Lopes-Marques et al. (2019a, 2019b, 2019c). Briefly, by using *H. sapiens* CDS for each target gene as reference, each exon was isolated and mapped to the genomic region of the candidate pseudogenes using Geneious Prime (2019.2.3) “map to reference” tool. The aligned regions were individually screened for ORF disrupting mutations (exon deletions, sequence frameshifts and premature stop codons) and identified mutations were annotated. Mutational validation was performed through retrieval of raw sequencing reads in (at least) two independent Sequence Read Archive (SRA) projects (when available).

2.3 RNA-seq analysis

Transcriptomic analysis was performed as previously described by Lopes-Marques (et al., 2019c). Succinctly, RNA-seq datasets of multiple tissues were obtained from SRA projects to inspect the functional condition of each target gene in Xenarthran species (when available) and Human (*H. sapiens*) (Supplementary Data 2). Transcriptomic reads recovered through blastn, were mapped to corresponding references genomes using the “map to reference” tool from Geneious Prime (2019.2.3) and manually removed if presenting poor alignment. Finally, reads were classified as spliced reads (spanning over two exons), exon-intron reads or exonic reads depending on the genomic region they mapped.

3. Results

3.1 Erosion of melatonin-related genes in Xenarthra

To infer the coding state of the melatonin synthesis genes in armadillos, anteaters and sloths, we compared the genomic regions containing *Aanat* and *Asmt* in *H. sapiens* to the full genomes of 8 eight Xenarthran species and *L. africana* (Supplementary Data 1). Analysis using PseudoChecker (Alves et al., 2020), showed that all analysed species presented a PseudoIndex equal to 5 (Supplementary Data 3) - thus suggesting that the ORF of *Aanat* and *Asmt* includes inactivating mutations. Subsequent manual annotation of all collected Xenarthran genomic sequences revealed *Aanat* and *Asmt* gene erosion across all analysed species (Supplementary Data 4). Regarding *Aanat*, in agreement with Yin (et al., 2021), we found numerous ORF-disrupting mutations, including a conserved 1-nucleotide deletion in exon 1 in Armadillos and a conserved 2-nucleotide deletion in exon 3 in Sloths (Figure 2; Supplementary Data 4). On the other hand, although Yin (et al., 2021) were not able to recover the genomic sequence containing the *Aanat* CDS in Anteaters, we uncovered, among other disruptive mutations, a conserved in-frame premature stop codon in exon 2. The identified mutations were next validated by searching at least one ORF disruptive mutation per species in the corresponding SRAs; reads corroborating the identified mutations were systematically found (Supplementary Data 5). The analysis of *Asmt* in Xenarthra also revealed variable disruption patterns across Xenarthra. In cingulatan, we found similar mutational events, however not conserved within members of this group. Specifically, in *D. novemcinctus* exons 1 to 4 and 7 were not found, possibly due to poor genome coverage or complete exon deletion (Figure 2). Moreover, in southern three-banded armadillo (*Tolypeutes matacus*) several insertions/deletions (indels) have been identified in exon 6, contrasting with *D. novemcinctus* where a validated in-frame premature stop codon in the same exon was detected (Figure 2; Supplementary Data 4 and 6). In Vermilingua (anteaters), we were only able to recover exons 4 and 5 in the Giant anteater (*Myrmecophaga tridactyla*) which provide a range of mutations with predicted disruptive effects (Figure 2; Supplementary Data 4). For Folivora (sloths), across several identified ORF-disrupting mutations, a trans-species conserved 4-nucleotide insertion in exon 6 was revealed and further validated by SRA searches (Supplementary Data 4 and 6). RNA-Seq analysis in Linnaeus's two-toed sloth

(*Choloepus didactylus*) *Aanat* yielded a high proportion of exon-intron reads versus spliced reads, in clear contrast with the pattern found in *H. sapiens* (Supplementary Data 7). In the case of *Asmt*, no transcriptomic reads were recovered for *C. didactylus*. Similarly, SRA transcriptome searches were unable to retrieve reads of *D. novemcinctus* for both genes.

Figure 2: Schematic representation of the identified *Aanat* and *Asmt* genes ORF abolishing mutations in Xenarthra orders (Cingulata and Pilosa). Phylogenetic trees were calculated in www.timetree.org; last accessed March 13, 2021 using species list. Silhouettes were sourced from Phylopic (<http://phylopic.org>). Sequence alignments of the identified conserved disruptive mutations in both *Aanat* and *Asmt* genes of Xenarthra.

We next examined the genes *Mtnr1a* and *Mtnr1b*, that encode G-protein coupled receptors responsible for melatonin signalling. In *D. novemcinctus*, the *Mtnr1a* coding status could not be accessed likely due to fragmentation of the respective genomic region (presence of sequencing gaps (Ns)). On the other hand, for both species comprising the two-toed sloth group (*Choloepus* sp.), we were able to identify a

validated 8-nucleotide deletion and a 20-nucleotide deletion in exon 2 together with a 4-nucleotide insertion in exon 1 (Figure 3; Supplementary Data 4 and 8). Curiously, in elephant (*Loxodonta africana*) exon 2 was not found despite the completeness of the assembly in the *Mtnr1a* region. The analysis of *Mtnr1b* CDS in *T. matacus* and Screaming hairy armadillo (*Chaetophractus vellerosus*) uncovered several inactivating mutations, including a conserved premature stop codon that truncates exon 2 (Figure 3). Pilosa species (sloths and anteaters) *Mtnr1b* gene annotation revealed the presence of several ORF disrupting mutations, of note a single transversal mutation present in all analysed species, namely a 2-nucleotide deletion in exon 2 (Figure 3; Supplementary Data 4). This mutation was investigated and validated in both *Choloepus* species (Supplementary Data 9). Searches for transcriptomic evidence for *Mtnr1a* and *Mtnr1b* in *C. didactylus* retrieved a low number of reads, mostly corresponding to immaturely mRNA (Supplementary Data 7).

3.2 Melatonin-related genes are inactivated in other non-xenarthran mammals

We next expanded our analysis to non-Xenarthran mammal genomes (157), to address the coding status of *Aanat*, *Asmt*, *Mtnr1a* and *Mtnr1b* (Supplementary Data 1). Sequence search and analysis for *Aanat* returned a total of 11 species with no annotation of a *Aanat*-like sequence: *Bison bison bison* (American bison), *Bos indicus* (Zebu), *Bos mutus* (Wild yak), *Bubalus bubalis* (Water buffalo), *Camelus ferus* (Wild bactrian camel), *Odocoileus virginianus texanus* (White-tailed deer), *Pantholops hodgsonii* (Tibetan antelope), *Sus scrofa* (Wild boar), *Myotis davidii* (David's myotis) and *Myotis lucifugus* (little brown bat). For the latter, we were not able to retrieve the genomic locus containing the target gene, given that both upstream and downstream flanking genes are also not annotated. Analysis using PseudoChecker (Alves et al., 2020), showed that 32 species non-Xenarthran mammals presented a PseudoIndex higher than 2 (Supplementary Data 3). From these species, members of Cetacea (cetaceans) and Pholidota (pangolins) presented among their members, a conserved (and validated) in-frame premature stop codon in Exon 1 (Supplementary Data 4 and 10). Moreover, we also found ORF-disrupting mutations in Exon 1 of velvety free-tailed bat (*Molossus molossus*), Kuhl's pipistrelle (*Pipistrellus kuhlii*) and Sunda flying lemur (*Galeopterus variegatus*) with the latter being validated through

SRA genomic reads (Supplementary Data 4 and 10). In *D. dugon*, several disruptive mutations were identified, such as an eight-nucleotide insertion in exon 2 and the presence of a stop codon in exon 3 (Supplementary Data 4 and 10).

Figure 3: Schematic representation of the identified *Mtnr1a* and *Mtnr1b* genes ORF abolishing mutations in Xenarthra orders (Cingulata and Pilosa). Phylogenetic trees were calculated in www.timetree.org; last accessed March 13, 2021 using species list. Silhouettes were sourced from Phylopic (<http://phylopic.org>). Sequence alignments of the identified conserved disruptive mutations in both *Mtnr1a* and *Mtnr1b* genes of Xenarthra.

Regarding *Asmt*, in 17 species the genomic fragments containing *Asmt*-like nucleotide sequences were not recovered given the lack of annotation for both target and flanking genes (Supplementary Data 3). For this gene, 74 species displayed a PseudoIndex higher than 2 (Supplementary Data 3), the majority due to fragmentation of the genomic region (presence of Ns), true absence of exons, poor alignment identity or incompleteness of the scaffold in the *Asmt* genomic region (Supplementary Data 4). Gene lesion events were found and validated mostly in Cetaceans, *G. variegatus*, *Manis* sp. (pangolins) and some Rodentia, with the latter showing poor alignment identity with the reference (Figure 4; Supplementary Data 4 and 11). Other examples of species presenting disruptive mutations but with no SRA validation (given that no genomic independent SRAs projects are available) include Brandt's bat (*Myotis brandtii*), the Prairie vole (*Microtus ochrogaster*) with 2-nucleotide deletion in exon 1 and the Nancy Ma's night monkey (*Aotus nancymae*)

with single nucleotide deletions in exon 2 (Supplementary Data 4). In the case of *D. dugon*, no ORF-disrupting mutations were found for *Asmt*, however, not all the exons were recovered due to incompleteness of the scaffold (Supplementary Data 4). We next expanded our search to understand whether melatonin signaling genes would be compromised in non-Xenarthran mammals. For *Mtnr1a*, a total of 44 species exhibited a PseudoIndex higher than 2 (Supplementary Data 3), yet manual curation revealed ORF-disrupting mutations only in pangolins (validated through SRA Projects; Supplementary Data 12) (Huelsmann et al., 2019), Hawaiian monk seal (*Neomonachus schauinslandi*) and *D. dugon* (Supplementary Data 4). In cetaceans a different scenario emerged, with several exons completely absent (Supplementary Data 4) (Huelsmann et al., 2019; Lopes-Marques et al., 2019b). In *Mtnr1b*, a total of 69 non-Xenarthran mammals displayed a PseudoIndex higher than 2 (Supplementary Data 3). However, and contrary to the pattern found for *Mtnr1a*, annotation of the collected sequences revealed *Mtnr1b* gene erosion across multiple species, mostly affecting the Carnivora and Cetaceans but also Pholidota, Sirenia and some Primates (Figure 4; Supplementary Data 4). Examples of conserved inactivating mutations were found in bears (*Ursus* sp.) with an in-frame premature stop codon in exon 1, weasels (*Mustela* sp.) sharing several indels in exon 2 and pangolins with a single nucleotide deletion and an in-frame premature stop codon in exon 1 (Supplementary Data 4 and 13). Other species with ORF-disruptive mutations include *Nannospalax galili* (northern Israeli blind subterranean mole rat), exhibiting a single nucleotide deletion in exon 1, *D. dugon* with a conserved fourteen-nucleotide deletion in exon 2 or *G. variegatus* with an indel also in exon 1 (Supplementary Data 4 and 13). Detailed characterization of each target gene in mammals is available in Supplementary Data 4 and the minutiae of SRA validation can be found in Supplementary Data 10, 11, 12 and 13.

Figure 4: Mutational landscape of melatonin synthesis and signalling genes along the mammalian tree. For each gene, we represented in green the orders where no ORF-disrupting mutations (frameshift mutations, in-frame premature stop codons, loss of canonical splicing site or exon deletions) were found across all members. On the other hand, orders where all members presented ORF-disrupting mutations are highlighted in red. In orders (and in genes) with no consensual disruption pattern, number of species presenting a coding/non-coding sequence were depicted respectively. Species where no SRA validation was possible, were not included in this figure. * indicates the presence of contradictory reports in *Mtnr1a* for *Trichechus manatus latirostris*. Phylogenetic relationships were adapted from Vazquez et al. (2018). (A) Summary characterization of mammalian lineages presenting complete molecular erosion of melatonin synthesis and signalling genes, regarding their sleep type, habitat and lifestyle. (B)

4. Discussion

Here, we set out to investigate how evolutionary genomic signatures might untangle the physiological status of controversial vestigial structures, using the pineal gland as a case study (Pévet, 2002). For this, we addressed the evolution of a melatonin-related gene hub, encompassing melatonin synthesis and signaling genes, in Xenarthra and other mammals. Our results strongly suggest a complete landscape of gene loss in Xenarthra, which further reinforce reports suggesting the lack of a pineal gland in several members of this superorder (Quay, 1965; Harlow et al., 1981; Benítez

et al., 1994, Ferrari 1998; Freitas 2019). On the other hand, in species in which a pineal gland was described (e.g., Freitas et al., 2019), the present data suggests that, despite the anatomical observations, the canonical pineal gland physiology leading to melatonin secretion is likely disrupted. Nevertheless, similarly to what was described for *Tursiops truncatus* (bottlenose dolphin) (Panin et al., 2012), previous radioimmunoassay methods have reported the presence of melatonin circulating in *D. novemcinctus* (Harlow et al., 1981), implying either the existence of independent pathways for melatonin synthesis and signaling (Slominski et al., 2003; Tan et al., 2016) or possible acquisition of melatonin from food sources (Tan et al., 2010). This strong genomic signature leading to the anatomical and/or physiological atrophy of this endocrine gland might be viewed as an adaptive solution to overcome physiological limitations. Described as cathemeral (irregular daily activity pattern) and heterothermic species (Eisenberg & Redford, 1999) with limited capacity to regulate their body temperature, Xenarthrans' movements are heavily influenced by air temperature (Gregeor, 1985; Camilo-Alves & Mourão, 2006; Giné et al., 2015; Attias et al., 2018). Thus, to reduce such energetic costs, Xenarthrans may have suffered reductive episodes, allowing behavioral strategies to overcome unfavorable environmental conditions and mitigate thermal limitations (Yin et al., 2021). Accordingly, convergent disruptive patterns with lineages also presenting labile body temperature and suggestive bizarre sleeping patterns (Pholidota; Mc Nab, 1984; Heath & Hammel, 1986; Weber et al., 1986; Imam et al., 2018; Yin et al., 2021) or living in environments with specific thermal constraints (Cetacea and *Trichechus manatus latirostris* (Florida manatee); Huelsmann et al., 2019; Lopes-Marques et al., 2019b; Yin et al., 2021), make it plausible to hypothesize that, in these species, inactivation of melatonin-related genes can be related with changes in their circadian rhythmicity. In addition, pseudogenization of these genes possibly paralleled loss of other circadian rhythm related genes, namely Cortistatin gene, that encodes a pleiotropic neuropeptide with an important role in sleep physiology (Valente et al., 2021). Given that the evolution of melatonin-related genes should be directly linked with pineal gland function, by inferring their coding status we were able to deduce if the organ constitutes an evolutionary vestige, despite the conflicting anatomical reports. More importantly, the present study provides a clear case-study on how genomic data can

be used to disentangle whether a specific organ constitutes a functional or vestigial structure (Hiller et al., 2012).

5. Conclusion

To date, no unequivocal inferences on the functional status of pineal gland across mammals were provided, with anatomical observations in several species from different clades presenting conflicting conclusions. However, by making use of genomic data, our results provide solid evidence for pineal gland vestigiality not only in Xenarthra, but also in other mammalian lineages. Thus, we argue that analysis of genomic changes might constitute a powerful approach to gain insights into the vestigiality of specific organs.

. Funding

This work is a result of the project ATLANTIDA (ref. NORTE-01-0145-FEDER-000040), supported by the Norte Portugal Regional Operational Programme (NORTE 2020), under the PORTUGAL 2020 Partnership Agreement and through the European Regional Development Fund (ERDF). One PhD fellowship for author R.V. (SFRH/BD/144786/2019) was granted by Fundação para a Ciência e Tecnologia (FCT, Portugal) under the auspices of Programa Operacional Regional Norte (PORN), supported by the European Social Fund (ESF) and Portuguese funds (MECTES).

. Author's contributions

Raul Valente: Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing - original draft **Filipe Alves:** Writing - review & editing **Isabel Sousa-Pinto:** Writing - review & editing **Raquel Ruivo:** Conceptualization, Methodology, Validation, Writing - review & editing **Luís Filipe Costa de Castro:** Conceptualization, Methodology, Validation, Supervision, Project administration, Resources, Writing - review & editing

. Acknowledgements

We acknowledge the various genome consortiums for sequencing and assembling the genomes.

419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450

. Declaration of Competing Interest

The authors declare that they have no competing interests.

. References

- Albalat, R., Cañestro, C., 2016. Evolution by gene loss. *Nat. Rev. Genet.* 17, 379–391.
<https://doi.org/10.1038/nrg.2016.39>
- Allmon, W.D., Ross, R.M., 2018. Evolutionary remnants as widely accessible evidence for evolution: the structure of the argument for application to evolution education. *Evol. Educ. Outreach* 11, 1. <https://doi.org/10.1186/s12052-017-0075-1>
- Alves, L.Q., Alves, J., Ribeiro, R., Ruivo, R., Castro, F. 2019. The dopamine receptor D5 gene shows signs of independent erosion in toothed and baleen whales. *PeerJ* 7, e7758. <https://doi.org/10.7717/peerj.7758>
- Alves, L.Q., Ruivo, R., Fonseca, M.M., Lopes-Marques, M., Ribeiro, P., Castro, L.F.C., 2020. PseudoChecker: an integrated online platform for gene inactivation inference. *Nucleic Acids Res.* 48, W321–W331. <https://doi.org/10.1093/nar/gkaa408>
- Attias, N., Oliveira-Santos, L.G.R, Fagan, W.F., Mourão, G., 2018. Effects of air temperature on habitat selection and activity patterns of two tropical imperfect homeotherms. *Animal Behav.* 140, 129–140.
- Axelrod, J., Wurtman, R.J., Winget, C.M., 1964. Melatonin synthesis in the Hen pineal and its control by light. *Nature* 201, 1134.
- Benítez, I., Aldana Marcos, H.J., Affanni, J.M., 1994. The encephalon of *Chaetophractus villosus*. A general view of its most salient features. *Comun. Biol.* 12, 57–73.

- Castro, L.F.C., Gonçalves, O., Mazan, S., Tay, B.-H., Venkatesh, B., Wilson, J.M., 2014. Recurrent gene loss correlates with the evolution of stomach phenotypes in gnathostome history. *Proc. R. Soc. B: Biol. Sci.*, 281
- Camilo-Alves, C.S.P., Mourão, G.M., 2006. Responses of a specialized insectivorous mammal (*Myrmecophaga tridactyla*) to variation in ambient temperature. *Biotropica* 38, 52–56.
- Chiarello, A.G., 1998. Activity budgets and ranging patterns of the Atlantic forest maned sloth *Bradypus torquatus* (Xenarthra: Bradypodidae). *J. Zool.* 246, 1–10.
- Cronk, Q., 2009. *The Molecular Organography of Plants*. Oxford University Press, Oxford.
- Darwin, C., 1859. *On the origin of species by means of natural selection or the preservation of favoured races in the struggle of life*. John Murray Press, London.
- Di Blanco, Y.E., Spørring, K.L., Di Bitetti, M.S., 2017. Daily activity pattern of re-introduced giant anteaters (*Myrmecophaga tridactyla*): effects of seasonality and experience. *Mammalia* 81, 11–21.
- Eisenberg, J.F., Redford, K.H., 1999. *Mammals of the Neotropics*. vol. 3, the central Neotropics: Ecuador, Peru, Bolivia, Brazil. University of Chicago Press, Chicago.
- Esfeld, K., Berardi, A.E., Moser, M., Bossolini, E., Freitas, L., Kuhlemeier, C., 2018. Pseudogenization and resurrection of a speciation gene. *Curr. Biol.* 28, 3776–3786.
- Fang, X., Seim, I., Huang, Z., Gerashchenko, M.V., Xiong, Z., Turanov, A.A., Zhu, Y., Lobanov, A.V., Fan, D., Yim, S.H., Yao, X., Ma, S., Yang, L., Lee, S., Buffenstein, R., Zhou, X., Krogh, A., Kim, E.B., Bronson, R.T., Sumner, R., Park, T.J., Zhang, G., Wang, J., Gladyshev, V.N., 2014. Adaptations to a subterranean environment and longevity revealed by the analysis of mole rat genomes. *Cell Rep.* 8, 1–11.

483

484 Ferrari, C.C., Marcos, H.J.A., Carmanchahi, P.D., Benítez, I., Affanni, J.M., 1998. The
485 Brain of the Armadillo *Dasypus hybridus*. A general view of its most salient features.
486 Biocell, 22, 123–140.

487

488 Freitas, L.M., dos Santos, O.P., Santos, A.L.Q., Rodrigues de Melo, F., Silveira, L.,
489 Jácomo, A.T.A., Pereira, K.F., Lima, F.C., 2019. Brain anatomy of two-toed sloth
490 (*Choloepus didactylus*, Linnaeus, 1758): A comparative gross anatomical study of
491 extant xenarthrans. Anat. Histol.
492 Embryol. 49, 130–143. <https://doi.org/10.1111/ahe.12501>

493

494 Giné, G.A.F., Cassano, C.R., de Almeida, S.S., Faria, D., 2015. Activity budget, pattern
495 and rhythm of maned sloths (*Bradypus torquatus*): responses to variations in ambient
496 temperature. Mammal Biol. 80, 459–467.
497 <https://doi.org/10.1016/j.mambio.2015.07.003>

498

499 Greigor, D.H.Jr., 1985. Ecology of the little hairy armadillo *Chaetophractus vellerosus*.
500 In: Montgomery, G.G. (Eds.), The evolution and ecology of armadillos, sloths and
501 vermilinguas. Smithsonian Institution Press, Washington D.C., pp. 397–405.

502

503 Guijarro-Clarke, C., Holland, P.W.H., Paps, J. 2020. Widespread patterns of gene loss
504 in the evolution of the animal kingdom. Nat. Ecol. Evol. 4, 519–523.
505 <https://doi.org/10.1038/s41559-020-1159-9>

506

507 Harlow, H.J., Phillips, J.A., Ralph, C.L., 1981. Daynight rhythm in plasma melatonin in a
508 mammal lacking a distinct pineal gland, the nine-banded armadillo. Gen. Comp.
509 Endocrinol. 45, 212–218. [https://doi.org/10.1016/0016-6480\(81\)90106-4](https://doi.org/10.1016/0016-6480(81)90106-4)

510

511 Harlow, H.J., Phillips, J.A., Ralph, C.L., 1982. Circadian rhythms and the effects of
512 exogenous melatonin in the ninebanded armadillo, *Dasypus novemcinctus*: a mammal
513 lacking a distinct pineal gland. Physiol. Behav. 29, 307–313.

514

- 515 Heath, M.E., Hammel, H.T., 1986. Body temperature and rate of O₂ consumption in
516 Chinese pangolins. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 250, 377–382.
- 517
- 518 Hiller, M., Schaar, B.T., Indjeian, V.B., Kingsley, D.M., Hagey, L.R., Bejerano G, 2012. A
519 “forward genomics” approach links genotype to phenotype using independent
520 phenotypic losses among related species. *Cell Rep.* 2(4), 817–23.
- 521
- 522 Huelsmann, M., Hecker, N., Springer, M.S., Gatesy, J., Sharma, V., Hiller, M., 2019.
523 Genes lost during the transition from land to water in cetaceans highlight genomic
524 changes involved in aquatic adaptations. *Sci. Adv.* 5 (9), eaaw6671.
525 <https://doi.org/10.1126/sciadv.aaw6671>
- 526
- 527 Imam, A., Bhagwandin, A., Ajao, M.S., Manger, P.R., 2018. The brain of the tree
528 pangolin (*Manis tricuspis*). V. The diencephalon and hypothalamus. *J. Comp. Neurol.*
529 527, 2413–2439. <https://doi.org/10.1002/cne.24619>
- 530
- 531 Jacob, S., Zelano, B., Gungor, A., Abbott, D., Naclerio, R., McClintock, M.K., 2000.
532 Location and gross morphology of the nasopalatine duct in human adults. *Arch.*
533 *Otolaryngol. Head Neck Surg.* 126, 741–748.
- 534
- 535 Kim, E.B., Fang, X., Fushan, A.A., Huang, Z., Lobanov, A.V., Han, L., Marino, S.M., Sun,
536 X., Turanov, A.A., Yang, P., Yim, S.H., Zhao, X., Kasaikina, M.V., Stoletzki, N., Peng, C.,
537 Polak, P., Xiong, Z., Kiezun, A., Zhu, Y., Chen, Y., Kryukov, G.V., Zhang, Q., Peshkin, L.,
538 Yang, L., Bronson, R.T., Buffenstein, R., Wang, B., Han, C., Li, Q., Chen, L., Zhao, W.,
539 Sunyaev, S.R., Park, T.J., Zhang, G., Wang, J., Gladyshev, V.N., 2011. Genome
540 sequencing reveals insights into physiology and longevity of the naked mole rat.
541 *Nature* 479, 223–227.
- 542
- 543 Klein, D., Coon, S., Roseboom, P., Weller, J.L., Bernard, M., Gastel, J.A., Zatz, M.,
544 Iuvone, P., Rodriguez, I., Bégay, V., Falcón, J., Cahill, G.M., Cassone, V.M., Baler, R.,
545 1997. The melatonin rhythm-generating enzyme: Molecular regulation of serotonin
546 N-acetyltransferase in the pineal gland. *Recent Prog. Horm. Res.* 52, 307–357.

547

548 Lewy, A.J., Wehr, T.A., Goodwin, F.K., Newsome, D.A., Markey, S.P., 1980. Light
549 suppresses melatonin secretion in humans. *Science* 210, 1267 – 1269.

550

551 Lindblad-Toh, K., Garber, M., Zuk, O., Lin, M.F., Parker, B.J., Washietl, S., Kheradpour,
552 P., Ernst, J., Jordan, G., Mauceli, E., et al., 2011. A high-resolution map of human
553 evolutionary constraint using 29 mammals. *Nature* 478, 476–482.

554

555 Lopes-Marques, M., Machado, A.M., Alves, L.Q., Fonseca, M.M., Barbosa, S., Sinding
556 M-H.S., Rasmussen, M.H., Iversen, M.R., Bertelsen, M.F., Campos, P.F., Da Fonseca,
557 R., Ruivo, R., Castro, L.F.C., 2019a. Complete inactivation of sebum-producing genes
558 parallels the loss of sebaceous glands in Cetacea. *Mol. Biol. Evol.* 36 (6), 1270–1280.
559 <https://doi.org/10.1093/molbev/msz068>.

560

561 Lopes-Marques, M., Ruivo, R., Alves, L.Q., Sousa, N., Machado, A.M., Castro, L.F.C.,
562 2019b. The singularity of Cetacea behavior parallels the complete inactivation of
563 melatonin gene modules. *Genes* 10(2), 121. <https://doi.org/10.3390/genes10020121>

564

565 Lopes-Marques, M., Alves, L.Q., Fonseca, M.M., Secci-Petretto, G., Machado, A.M.,
566 Ruivo, R., Castro, L.F.C. 2019c. Convergent inactivation of the skin-specific CC motif
567 chemokine ligand 27 in mammalian evolution. *Immunogenetics* 71(5), 363–372.
568 <https://doi.org/10.1007/s00251-019-01114-z>

569

570 Maccarini, T.B., Attias, N., Medri, I.M., Marinho-Filho, J., Mourão, G.M., 2015.
571 Temperature influences the activity patterns of armadillo species in a large
572 neotropical wetland. *Mamm. Res.* 60, 403–409.

573

574 McGaugh, S.E., Gross, J.B., Aken, B., Blin, M., Borowsky, R., Chalopin, D., Hinaux, H.,
575 Jeffery, W.R., Keene, A., Ma, L., Minx, P., Murphy, D., O’Quin, K.E., Rétaux, S., Rohner,
576 N., Searle, S.M.J., Stahl, B.A., Tabin, C., Volff, J.-N., Yoshizawa, M., Warren, W.C., 2014.
577 The cavefish genome reveals candidate genes for eye loss. *Nat Commun.* 5, 5307.

578

- Mc Nab, B.K., 1979. The influence of body size on the energetics and distribution of fossorial and burrowing mammals. *Ecology* 60, 1010–1021.
- Mc Nab, B.K., 1980. Energetics and the limits to a temperate distribution in armadillos. *J. Mammal.* 61, 606–627.
- Mc Nab, B.K., 1984. Physiological convergence amongst ant-eating and termite-eating mammals. *J. Zool. (Lond.)* 203, 485–510.
- Mc Nab, B.K., 1985. Energetics, population biology and distribution of xenarthrans living and extinct. In: Montgomery, G.G. (Ed.), *The Evolution and Ecology of Armadillos, Sloths and Vermilinguas*. Smithsonian Institution Press, Washington/London, pp. 219–232.
- Murphy, W.J., Pringle, T.H., Crider, T.A., Springer, M.S., Miller, W., 2007. Using genomic data to unravel the root of the placental mammal phylogeny. *Genome Res.* 17, 413–421.
- Nweeia, M.T., Eichmiller, F.C., Hauschka, P.V., Tyler, E., Mead, J.G., Potter, C.W., Angnatsiak, D.P, Richard, P.R., Orr, J.R., Black, S.R., 2012. Vestigial tooth anatomy and tusk nomenclature for *Monodon monoceros*. *Anat. Rec.* 295, 1006–1016.
- Oksche, A., 1965. Survey of the development and comparative morphology of the pineal organ. *Prog. Brain Res.* 10, 3–29.
- O’Leary, M.A., Bloch, J.I., Flynn, J.J., Gaudin, T.J., Giallombardo, A., Giannini, N.P., Goldberg, S.L., Kraatz, B.P., Luo, Z.-X., Meng, J., et al., 2013. The placental mammal ancestor and the post-K-Pg radiation of placentals. *Science* 339, 662–667.
- Olson, M.V., 1999. When less is more: gene loss as an engine of evolutionary change. *Am. J. Hum. Genet.* 64, 18–23.

- 611 Panin, M., Gabai, G., Ballarin, C., Peruffo, A., Cozzi, B, 2012. Evidence of melatonin
612 secretion in cetaceans: Plasma concentration and extrapineal HIOMT-like presence in
613 the bottlenose dolphin *Tursiops truncatus*. Gen. Comp. Endocrinol. 177, 238–245.
614
- 615 Pévet, P., 2002. Melatonin. Dialogues Clin. Neurosci. 4, 57–72.
616
- 617 Quay, W., 1965. Histological structure and cytology of the pineal organ in birds and
618 mammals. In: Kappers, J., Schade, J. (Eds.), Progress in brain research, Structure and
619 function of the epiphysis Cerebri, vol 10. Elsevier, New York, pp. 49–
620 86. [https://doi.org/10.1016/S0079-6123\(08\)63447-0](https://doi.org/10.1016/S0079-6123(08)63447-0)
621
- 622 Ralph, C.L., 1975. The pineal gland and geographical distribution of animals. Int. J.
623 Biometeorol. 19, 289–303.
624
- 625 Ralph, C.L., Young, S., Gettinger, R., O’Shea, T.J., 1985. Does the manatee have a
626 pineal body? Acta Zool. 66, 55–60.
627
- 628 Reppart, S.M., Weaver, D.R., Godson, C., 1996. Melatonin receptors step into the
629 light: Cloning and classification of subtypes. Trends Pharmacol. Sci. 17, 100–102.
630
- 631 Simonneaux, V., Ribelayga, C., 2003. Generation of the melatonin endocrine message
632 in mammals: A review of the complex regulation of melatonin synthesis by
633 norepinephrine, peptides, and other pineal transmitters. Pharmacol. Rev. 55, 325.
634
- 635 Slominski, A., Pisarchik, A., Semak, I., Sweatman, T., Wortsman, J., 2003.
636 Characterization of the serotonergic system in the C57BL/6 mouse skin. Eur. J.
637 Biochem. 270, 3335–3344.
638
- 639 Springer, M.S., Guerrero-Juarez, C.F., Huelsmann, M., Collin, M.A., Danil, K.,
640 McGowen, M.R., Oh, J.W., Ramos, R., Hiller, M., Plikus, M.V., Gatesy, J., 2021. Genomic
641 and anatomical comparisons of skin support independent adaptation to life in water
642 by cetaceans and hippos. Current Biology (in press).

643
644 Superina, M., Loughry, W.J., 2015. Why do xenarthrans matter? *J. Mammal.* 96, 617–
645 621. <https://doi.org/10.1093/jmammal/gyv099>
646
647 Tan, D.X., Hardeland, R., Manchester, L.C., Paredes, S.D., Korkmaz, A., Sainz, R.M.,
648 Mayo, J.C., Fuentes-Broto, L., Reiter, R.J., 2010. The changing biological roles of
649 melatonin during evolution: From an antioxidant to signals of darkness, sexual
650 selection and fitness. *Biol. Rev. Camb. Philos. Soc.* 85, 607–623.
651
652 Tan, D.X., Hardeland, R., Back, K., Manchester, L.C., Alatorre-Jimenez, M.A., Reiter,
653 R.J., 2016. On the significance of an alternate pathway of melatonin synthesis via 5-
654 methoxytryptamine: Comparisons across species. *J. Pineal Res.* 61, 27–40.
655
656 Themudo, G.E., Alves, L.Q., Machado, A.M., Lopes-Marques, M., da Fonseca, R.R.,
657 Fonseca, M., et al. (2020). Losing genes: the evolutionary remodeling of Cetacea skin.
658 *Front. Mar. Sci.* 7, 912.
659
660 Uliano-Silva, M., Winkler, S., Myers, E., Mazzoni, C., 2019. Slothomics: the first
661 chromosome-level genome of the slowest existing mammalian group. The G10K-
662 VGP/EBP 2019 Meeting Agenda, Manhattan, New York, August 2019. Poster
663 presentation
664
665 Valente, R., Alves, L.Q., Nabais, M., Alves, F., Sousa-Pinto, I., Ruivo, R., Castro, L.F.C.,
666 (2021). Convergent Cortistatin losses parallel modifications in circadian rhythmicity
667 and energy homeostasis in Cetacea and other mammalian lineages. *Genomics* 113(1),
668 1064-1070. <https://doi.org/10.1016/j.ygeno.2020.11.002>
669
670 Vazquez, J.M., Sulak, M., Chigurupati, S., Lynch, V.J., 2018. A zombie LIF gene in
671 elephants is upregulated by TP53 to induce apoptosis in response to DNA damage.
672 *Cell Rep.* 24 (7), 1765–1776.
673

Weber, R.E., Heath, M.E., White, F.N., 1986. Oxygen binding functions of blood and hemoglobin from the Chinese pangolin, *Manis pentadactyla*: possible implications of burrowing and low body temperature. *Respir. Physiol.* 64, 103–112.

Werth, A.J., 2014. Vestiges of the natural history of development: historical holdovers reveal the dynamic interaction between ontogeny and phylogeny. *Evol. Educ. Outreach* 7, 12. <https://doi.org/10.1186/s12052-014-0012-5>

Yin, D., Zhou, R., Yin, M., Chen, Y., Xu, S., Yang, G., 2021. Gene duplication and loss of AANAT in mammals driven by rhythmic adaptations, *Mol. Biol. Evol.* msab125. <https://doi.org/10.1093/molbev/msab125>

Zoonomia Consortium, 2020. A comparative genomics multitool for scientific discovery and conservation. *Nature* 587, 240–245. <https://doi.org/10.1038/s41586-020-2876-6>