URBANISATION, BIRD SPECIES RICHNESS AND ABUNDANCE WITHIN IBADAN METROPOLIS, NIGERIA

1. Festus O. Adegbola¹ (Corresponding Author)

Email address: adegbolafestuso@gmail.com

2. Taiye Adeniyi Adeyanju¹

Email address: at.adeyanju@mail1.ui.edu.ng

3. Soladoye B. Iwajomo²

Email address: siwajomo@unilag.edu.ng

4. Ibukunoluwa Augustine Ayodele¹

Email address: profibikunayodele@yahoo.com

1 Department of Wildlife and Ecotourism Management, University of Ibadan, Ibadan, Nigeria

2 Department of Zoology, University of Lagos, Lagos, Nigeria

ABSTRACT

Urbanisation is considered as one of the most profound threat to wildlife, with habitat loss and fragmentation being predominant. This study assessed the impacts of urbanisation on richness, abundance of bird species within Ibadan metropolis, Nigeria. A uniform grid of 500 square meters was installed on the map of Ibadan Metropolis using QGIS to produce 499 grid points distributed across the five urban local governments. 100 grids were selected randomly, identified with mapinR software application and surveyed with 5 point counts within each grid, established at 200m interval to avoid double counting. Each point count was observed for 5 minutes using a pair of 8x42mm binoculars within a 50m radius. Habitat variables like number of buildings, trees, paved roads, communication masts were also recorded.

A total number of 56 species of birds were observed at the end of the assessment, classified into 30 families. The test of statistics showed that there was no statistically significant difference in bird species richness between the Local Governments. The test of statistics showed that there was no statistically significant difference in bird species abundance between the local governments. The test of between-subjects effects revealed that there were no statistical significant effects when all the habitat variables were computed in the model on species richness. The number of paved roads and number of vehicles showed a significant effect on bird species abundance while others variables in the model did not exact statistically significant effects on bird species abundance.

The study therefore concluded that habitat actions due to urbanisation have not affected the richness and abundance of birds species found in all the local governments. The only habitat variables that have currently exacted significant effect on species abundance within the metropolis are number of vehicles and paved roads.

Word Count: 292

Keywords: Conservation, Species richness, Species abundance, Urbanisation

INTRODUCTION

In wildlife conservation and management, there is need for a thorough understanding of the relationships among individual organisms and their environment, as it is important in the development of ecological theories and in the implementation of conservation strategies (Walker et al. 2008). It is general knowledge among environmentalists that there is an increase in the proportion of the earth's surface now converted to human-dominated urban areas; it therefore becomes expedient to understand how urban wildlife communities within these now sophisticated ecosystem types are structured (McKinney 2002). The importance of understanding the ecological effects of urbanisation cannot be under-emphasised, especially with its rapid conversion of previously wild lands around the world. The process associated with urbanisation has profound effects on the distribution of wildlife species and their habitats (Wolff et al. 2018). Urban development is characterised by rapid population growth and profound land use transformation, leading to land conversion, which is a predominant process affecting ecological community structure and population dynamics of living organisms (Hostetler 1997). Research studies show that urban landscapes supports biotic communities in which only a few species increase in density compared to natural areas, resulting in a distinct difference the community diversity between these two landscapes (McKinney 2002).

Developing countries have a large number of wildlife existing outside the protected areas, on farmlands, and in urban areas. Among all wildlife, bird species are largely one of the most common wildlife surviving in urban communities (Gatesire et al. 2015). Birds are one of the most easily studied taxa occurring in cities worldwide as they serve as indicators variables in ecological assessment and monitoring (Magle et al. 2012). Birds are also important in maintaining ecosystems. For example, insectivorous bird species regulate disease vectors including mosquito and rodents. Pied Crow (*Corvus albus*), contributes to biomass recycling and reduce levels of disposable waste as scavengers. Frugivorous birds play a crucial role in the dispersal of seeds of fruit trees. Sun birds also helps to pollinate plants (Gatesire et al. 2015).

Studies show that bird species in different regions respond to urbanisation in a similar way with most research suggesting that bird communities are negatively impacted by urbanisation (Lin et al. 2011; Sol et al. 2014). There is a general shrinkage in species distribution as urbanisation increases, and the fact that similar bird species can be found in various urban landscape indicates

that urbanisation has a similar effect on local communities of birds irrespective of the region (Lin et al. 2011). As a result of the crucial role birds play in maintaining ecosystems and also support biodiversity, conservationists seek their protection against biological threats and protect the environment efficiently (Gatesire et al. 2015).

Primarily the development level, habitat diversity, age and diversity of vegetation present determine the richness of birds in urban areas (McKinney 2002). Native bird species that persists in urban landscapes partly depends on the actions of the landowners because the structural and vegetative characteristics of urban landscapes are largely human-influenced (McCaffrey and Mannan 2012). Urban areas have less assembly of bird species than adjacent natural areas, though some type and level of development support more native bird species than others. According to McKinney (2002), the moderate level of development and vegetation linkable to low density residential areas can support higher densities of some native birds species than other types of urban land use, also including undisturbed sites.

MATERIALS AND METHODS

STUDY AREA

The study was conducted in Ibadan metropolis, the capital of Oyo State. Ibadan metropolis, covering an area of 129.65 km², is located in South-Western Nigeria in the southeastern part of Oyo State at about 119 kilometers (74 miles) northeast of Lagos and 120 kilometers (75 miles) east of the Nigerian international border with the Republic of Benin (Popoola and Wahab 2018). Ibadan falls totally within the forest zone but close to the boundary between the forest and the derived savanna. It lies between latitude 3°3' N and 4°10' N and longitude 7°2'E and 7°40' E. (Popoola and Wahab 2018). The population of Metropolitan Ibadan is 1 338 659 according to census results for 2006 (Areola and Ikporukpo 2018). There are eleven (11) local governments in Ibadan Metropolitan area consisting of five urban local governments and six semi-urban local governments. The former are: Ibadan South East, Ibadan North East, Ibadan North West, Ibadan South West and Ibadan North Local Government Areas. Ibadan North Local Government has the largest land area among the urban Local Governments, while Ibadan North West is the smallest (Fig. 2). The second largest local government in Urban Ibadan is Ibadan South West. Ibadan metropolis is an important commercial centre and it comprises of people of different cultural and socio-economic backgrounds. General land use pattern of the Ibadan metropolitan area shows a clear distinction purely residential use for Urban Ibadan and agricultural use for semi-urban Ibadan (Salami et al. 2016).

88

89

90 Figure 1: Map of Ibadan Metropolis

91 Source: (Popoola and Wahab 2018)

BIRD ASSESSMENT

Using QGIS software application, a uniform grid of 500 by 500 metres was installed on the map of Ibadan Metropolis to produce 499 grids distributed across all the five urban local governments. 100 grids were surveyed with 5 point counts within each grid, established at 200 m interval to avoid double counting, each point count was observed for 5 minutes within a 50 m radius. Laser range finder was used to delineate the 50 m radius around each survey point, and to estimate distances to birds. Birds encountered outside the study grids were recorded only when it has never been observed in any of the grids before. When a bird could not be identified in the field, photos from a high resolution camera were taken for later identification by an expert in ornithology. During each visit, a pair of 8x42 mm binoculars was used for sighting birds. Helms field guide to the birds of western Africa (Borrow and Demey 2013) was used to identify the birds; birds call were recorded with a voice recorder and later played back for confirmation. The data were collected between October to December 2020, and observations were done in the morning and evening. Habitat variables recorded for this study, the number of buildings, number of trees, number of communication masts, ground cover, canopy cover, number of pedestrians and vehicles passing within 25 m radius around the point, describe urban form. No specific permissions were required to conduct this work. All bird surveys were conducted in areas which are open to the public; therefore there was no need to ask land managers for approval

Figure 3: Map showing survey grids

Source: QGIS Application

118 **DATA ANALYSIS**

119 Species richness and abundance was obtained by counting the total number of species and
 120 individual per species respectively as recorded within each grid. The data collected were entered
 121 and summarised in Microsoft Excel spread sheet computer software for analysis. Tables and
 122 figures were used to represent the results.

123 General Linear models method was fitted in R statistical package version 3.4.2 using test of
 124 correlation to the determine relationship between the vegetation variables, bird species richness,
 125 and bird abundance at 0.05 level of significance.

RESULTS

This chapter present the results of the data obtained for the purpose of this study. QGIS software application, using a uniform grid of 500 by 500 metres installed across all five selected urban local government in Ibadan was used in collecting the data. General Linear Model (GLM) of statistical analysis was used in analyzing the collected data and testing for statistical significant difference in bird species richness, bird abundance and the habitat variables that determines/ influence bird species richness and its abundance within the metropolis. ($\alpha = 0.05$).

The study observed 56 different species of birds which was grouped into 30 families. (See appendix)

DETERMINATION OF SPECIES RICHNESS WITHIN THE METROPOLIS

Descriptive statistics, Kruskal- Wallis H Test and Post Hoc Test were carried out to determine whether there are statistically significant differences between bird species richness and the five selected urban local government. The local government being the independent variable and bird species observed being the dependent variable.

The results showed that mean bird species richness at Ibadan North East, North West, South East and South West were estimated at 8.20 ± 1.643 , 10.84 ± 2.630 , 8.27 ± 2.724 , 9.36 ± 2.629 and 10.54 ± 5.222 respectively.

On the overall, Bird species richness for Ibadan metropolis was estimated at 9.73 ± 3.301 . (See table 1a)

The Kuskal- Wallis H Test revealed that the mean rank for Ibadan North East is **24.70**, Ibadan North is **44.74**, Ibadan North West is **25.32**, Ibadan South East is **34.55** and Ibadan South West is **36.38** (See table 1b). It implies that mean of Ibadan North will be rank first in bird species richness, followed by Ibadan South West, Ibadan South East, Ibadan North West and Ibadan North East respectively.

The test statistics table showed a chi-square value (X^2) of **8.253** and a p-value of **0.083**. At 5% C.I., the calculated p-value (0.083) is greater than 0.05. This implies that bird species richness between the different local governments is ns.

Table 1a: Descriptive Statistics on Species Richness

Descriptive Statistics

Dependent Variable: Species richness

Local Government	N	Mean	Std. Deviation
Ibadan North East	5	8.20	1.643
Ibadan North	19	10.84	2.630
Ibadan North West	11	8.27	2.724
Ibadan South East	22	9.36	2.629
Ibadan South West	13	10.54	5.222
Total	70	9.73	3.301

Table 1b: Kruskal- Wallis H Test on species richness

Kruskal-Wallis Test Rank

Local Government	N	Mean Rank
Ibadan North East	5	24.70
Ibadan North	19	44.74
Ibadan North West	11	25.32
Ibadan South East	22	34.55
Ibadan South West	13	36.38
Total	70	

Test Statistics

Species richness

Chi-Square	8.253
Df	4
Asymp. Sig.	.083

a. Kruskal Wallis Test

b. grouping variable: local government

165 **Table 1c: Post Hoc Test on Bird Species Richness**

166

Multiple Comparisons						
Dependent Variable: Species richness						
Tukey HSD						
		Mean		Sig.	95% Confidence Interval	
		Difference	Std.		Lower	Upper
(I) Local government		(I-J)	Error		Bound	Bound
Ibadan North East	Ibadan North	-2.64	1.628	.488	-7.21	1.93
	Ibadan North West	-.07	1.747	1.000	-4.97	4.83
	Ibadan South East	-1.16	1.605	.950	-5.67	3.34
	Ibadan South West	-2.34	1.704	.647	-7.12	2.44
Ibadan North Local Government	Ibadan North East	2.64	1.628	.488	-1.93	7.21
	Ibadan North West	2.57	1.227	.235	-.87	6.01
	Ibadan South East	1.48	1.014	.593	-1.37	4.32
	Ibadan South West	.30	1.166	.999	-2.97	3.57
Ibadan North West Local government	Ibadan North East	.07	1.747	1.000	-4.83	4.97
	Ibadan North	-2.57	1.227	.235	-6.01	.87
	Ibadan South East	-1.09	1.196	.891	-4.45	2.26
	Ibadan South West	-2.27	1.327	.437	-5.99	1.46
Ibadan South East	Ibadan North East	1.16	1.605	.950	-3.34	5.67

Local Government	Ibadan North	-1.48	1.014	.593	-4.32	1.37
	Ibadan North West	1.09	1.196	.891	-2.26	4.45
	Ibadan South West	-1.17	1.133	.837	-4.35	2.00
Ibadan South West	Ibadan North East	2.34	1.704	.647	-2.44	7.12
Local Government	Ibadan North	-.30	1.166	.999	-3.57	2.97
	Ibadan North West	2.27	1.327	.437	-1.46	5.99
	Ibadan South East	1.17	1.133	.837	-2.00	4.35

Based on observed means.

The error term is Mean Square (Error) = 10.490.

167

168

DETERMINATION OF SPECIES ABUNDANCE WITHIN THE METROPOLIS

The results in table 4.2a showed that *the mean bird abundance and standard deviation* was estimated at 64.40 ± 12.28 for Ibadan North East, 68.42 ± 31.39 for Ibadan North, 46.36 ± 9.63 for Ibadan North West, 53.68 ± 24.16 for Ibadan South East and 65.69 ± 51.26 Ibadan South West. Overall mean bird abundance and standard deviation is estimated at 59.52 ± 31.35 .

The Kruskal-Wallis Test showed that in term of ranking the abundance of bird species, Ibadan North East will be rank first, followed by Ibadan North, Ibadan North west, Ibadan South east and Ibadan South West respectively. (See table 2b)

The test statistics table showed a chi-square value (X^2) of **6.973** and a p-value of **0.137**. At 5% S.L., the calculated p-value is .137; which is greater than 0.05. This implies that bird species abundance between the different local governments is ns.

The multiple comparisons table showed that were no statistically significant differences in bird species abundance between any of the selected local government, as p-value in all the comparison are greater than 5%. (See table 2c)

Table 2a: Descriptive Statistics on species abundance

Descriptive Statistics

Dependent Variable: Species abundance

Local Government	N	Mean	Std. Deviation
Ibadan North East	5	64.4000	12.28007
Ibadan North	19	68.4211	31.39058
Ibadan North West	11	46.3636	9.63611
Ibadan South East	22	53.6818	24.16291
Ibadan South West	13	65.6923	51.26465
Total	70	59.5286	31.35571

Table 2b: Kruskal- Wallis H Test on Bird Species Abundance

Kruskal-Wallis Test Rank		
Local Government	N	Mean Rank
Ibadan North East	5	47.00
Ibadan North	19	42.79
Ibadan North West	11	26.27
Ibadan South East	22	32.18
Ibadan South West	13	33.85
Total	70	

Test Statistics	
Species abundance	
Chi-Square	6.973
Df	4
Asymp. Sig.	.137

a. Kruskal Wallis Test

b. grouping variable: local government

194

Table 2c

Post Hoc Test on Bird Species Abundance

Multiple Comparisons

Dependent Variable:		Species Abundance				
Tukey HSD						
					95% Confidence Interval	
		Mean Difference (I-J)	Std. Error	Sig.	Lower Bound	Upper Bound
(I) Local government	Ibadan North East	-4.0211	15.65613	.999	-47.9494	39.9073
	Ibadan North West	18.0364	16.80039	.819	-29.1026	65.1753
	Ibadan South East	10.7182	15.43214	.957	-32.5817	54.0181
	Ibadan South West	-1.2923	16.39156	1.000	-47.2841	44.6995
Ibadan North Local Government	Ibadan North East	4.0211	15.65613	.999	-39.9073	47.9494
	Ibadan North West	22.0574	11.80125	.344	-11.0548	55.1697
	Ibadan South East	14.7392	9.75539	.559	-12.6327	42.1111
	Ibadan South West	2.7287	11.21159	.999	-28.7290	34.1865
Ibadan North West Local government	Ibadan North East	-18.0364	16.80039	.819	-65.1753	29.1026
	Ibadan North	-22.0574	11.80125	.344	-55.1697	11.0548
	Ibadan South East	-7.3182	11.5024	.969	-39.5920	24.9556

			4			
	Ibadan South West	-19.3287	12.7608	.557	-55.1333	16.4759
			1			
Ibadan South East	Ibadan North East	-10.7182	15.4321	.957	-54.0181	32.5817
Local Government			4			
	Ibadan North	-14.7392	9.75539	.559	-42.1111	12.6327
	Ibadan North West	7.3182	11.5024	.969	-24.9556	39.5920
			4			
	Ibadan South West	-12.0105	10.8966	.805	-42.5845	18.5635
			2			
Ibadan South West	Ibadan North East	1.2923	16.3915	1.000	-44.6995	47.2841
Local Government			6			
	Ibadan North	-2.7287	11.2115	.999	-34.1865	28.7290
			9			
	Ibadan North West	19.3287	12.7608	.557	-16.4759	55.1333
			1			
	Ibadan South East	12.0105	10.8966	.805	-18.5635	42.5845
			2			

Based on observed means.

The error term is Mean Square (Error) = 970.245.

DETERMINATION OF HABITAT PARAMETERS THAT INFLUENCE BIRD SPECIES RICHNESS WITHIN IBADAN METROPOLIS

The test of between-subjects effects in table 3 showed whether the habitat variables (independent variables) are statistically significant in assessing and influencing bird species richness within Ibadan Metropolis.

The results revealed that there was **no** statistically significant relationship between the effect of the number of **buildings** on bird species richness, $F(1, 54) = 0.075, p = 0.786$; there was **no** statistically significant relationship between the effects of the number of **trees** on bird species richness, $F(1, 54) = 0.450, p = 0.505$; there was **no** statistically significant relationship between the effects of the number of **masts** on bird species richness, $F(1, 54) = 0.323, p = 0.572$; there was **no** statistically significant relationship between the effects of the number of **paved roads** on bird species richness, $F(1, 54) = 0.808, p = 0.373$; there was **no** statistically significant relationship between the effects of the number **lawns** on bird species richness, $F(1, 54) = 0.090, p = 0.766$; there was **no** statistically significant relationship between the effects of the percentage of **canopy covers** on bird species richness, $F(1, 54) = 0.360, p = 0.881$; there was **no** statistically significant relationship between the effects of the percentage of **ground covers** on bird species richness, $F(1, 54) = 0.043, p = 0.836$; there was **no** statistically significant relationship between the effects of the presence of **farm land** on bird species richness, $F(1, 54) = 0.30, p = 0.863$; there was **no** statistically significant relationship between the effects of the number of **vehicles** on bird species richness, $F(1, 54) = 7.437, p = 0.09$; there was **no** statistically significant relationship between the effects of the number of **pedestrians** on bird species richness, $F(1, 54) = 2.744, p = 0.103$; there was **no** statistically significant relationship between the effects of the number of **nests** on bird species richness, $F(1, 54) = 0.517, p = 0.475$.

Also, **R square of 32.9%** implies that all the independent variables are not too strong in predicting or influencing model.

We can therefore conclude that there were no statistically significant interaction between all the habitat parameters/variables and bird species richness within Ibadan Metropolis. That is all the habitat parameters/ variables did not influence bird species richness within Ibadan.

Table 3: Test of between- Subjects Effects on Species Richness and habitat variables

Test of Between- Subjects Effects

Source	Type III Sum of Squares	df	Mean Square	F	Sig. (p-value)
Corrected Model	247.093 ^a	15	16.473	1.762	.066
Intercept	317.361	1	317.361	33.952	.000
Number of Buildings	.698	1	.698	.075	.786
Number of Trees	4.202	1	4.202	.450	.505
Number of Masts	3.020	1	3.020	.323	.572
Number of Paved	7.556	1	7.556	.808	.373
Number of Lawns	.838	1	.838	.090	.766
Percentage of Canopy cover	3.362	1	3.362	.360	.551
Percentage of Ground cover	.405	1	.405	.043	.836
Presence of farmland	.280	1	.280	.030	.863
Number of vehicles	69.511	1	69.511	7.437	.009
Number of Pedestrians	25.651	1	25.651	2.744	.103
Number of nests	4.831	1	4.831	.517	.475
Error	504.750	54	9.347		
Total	7377.000	70			
Corrected Total	751.843	69			

R Squared = .329 (Adjusted R Squared = .142)

DETERMINATION OF HABITAT PARAMETERS THAT INFLUENCE BIRD SPECIES ABUNDANCE WITHIN IBADAN METROPOLIS

The results revealed that there was **no** statistically significant interaction between the effect of the number of **buildings** on bird species richness, $F(1, 54) = 0.20$, $p = 0.888$; there was **no** statistically significant interaction between the effects of the number of **trees** on bird species richness, $F(1, 54) = 2.181$, $p = 0.145$; there was **no** statistically significant interaction between the effects of the number of **masts** on bird species richness, $F(1, 54) = 0.359$, $p = 0.552$; there was statistically significant interaction between the effects of the number of **paved roads** on bird species richness, $F(1, 54) = 5.632$, $p = 0.021$; there was **no** statistically significant interaction between the effects of the number of **lawns** on bird species richness, $F(1, 54) = 1.408$, $p = 0.241$; there was **no** statistically significant interaction between the effects of the percentage of **canopy covers** on bird species richness, $F(1, 54) = 1.412$, $p = 0.240$; there was **no** statistically significant interaction between the effects of the percentage of **ground cover** on bird species richness, $F(1, 54) = 0.01$, $p = 0.981$; there was **no** statistically significant interaction between the effects of the presence of **farm land** on bird species richness, $F(1, 54) = 0.100$, $p = 0.754$; there was statistically significant interaction between the effects of the number of **vehicles** on bird species richness, $F(1, 54) = 5.042$, $p = 0.022$; there was **no** statistically significant interaction between the effects of the number of **pedestrians** on bird species richness, $F(1, 54) = 0.302$, $p = 0.585$; there was **no** statistically significant interaction between the effects of the number of **nests** on bird species richness, $F(1, 54) = 0.419$, $p = 0.520$. The results revealed that there were statistically significant interaction between the number of paved and the number of vehicles on the bird species abundance and no statistically significant interaction in all other habitat variables/parameters. (See table 4)

Also, R square of 26.2% implies that the independent variables are weak in predicting or influencing bird species abundance.

We can therefore conclude that only number of paved roads and number of vehicles exact a significant effect on bird species abundance while all other habitat variables do not exact significant influence on bird species abundance within Ibadan Metropolis. That is, all other habitat parameters/ variables did not influence bird species abundance, except numbers of paved roads and vehicles available within Ibadan metropolis.

Table 4: Test of between- Subjects Effects on Species Abundance and Habitat Variables

Test of Between- Subjects Effects					
Source	Type III Sum of Squares	df	Mean Square	F	Sig. (p - value)
Corrected Model	17805.715	15	1187.048	1.281	.246
Intercept	5930.968	1	5930.968	6.401	.014
Number of Buildings	18.417	1	18.417	.020	.888
Number of Trees	2027.731	1	2027.731	2.188	.145
Number of Masks	332.432	1	332.432	.359	.552
Number of Paved	5218.020	1	5218.020	5.632	.021
Number of Lawns	1304.756	1	1304.756	1.408	.241
Number of Canopy cover	1308.209	1	1308.209	1.412	.240
Number of Ground cover	.531	1	.531	.001	.981
Presence of farmland	92.273	1	92.273	.100	.754
Number of vehicles	5135.375	1	5135.375	5.542	.022
Number of Pedestrians	279.845	1	279.845	.302	.585
Number of nests	388.402	1	388.402	.419	.520
Error	50033.728	54	926.551		
Total	315895.000	70			
Corrected Total	67839.443	69			

R Squared = .262 (Adjusted R Squared = .058)

262 4.5 PICTURES OF BIRDS SPECIES AND ACTIVITY SIGHTED IN URBAN AREAS

263

264 Figure 4: Common Kestrel (*Falco tinnunculus*) resting on a billboard

265

266 Figure 5: Yellow Billed Kite (*Milvus aegyptius*) brooding in a nest hung on a billboard

267

268 Figure 6: Pied Crow (*Corvus albus*) hovering in the sky

269

270

271 Figure 7: Yellow Billed Kite (*Milvus aegyptius*) hovering in the sky

272

273 Figure 8: Laughing Dove (*Streptopelia semitorquata*) resting on a roof

274

275

276 Figure 9: Speckled Pigeon (*Columba guinea*) in its roost

DISCUSSION

The objective of this study was to examine the impacts of urbanization on bird species richness, abundance and the influence of habitat parameters on species richness and abundance in five selected urban local governments within Ibadan Metropolis. The researchers assessed the presence of birds using QGIS software application, of a uniform grid of 500 by 500 metres installed in some strategic areas in those local governments.

A total number of fifty-six (56) different species of birds were observed at the end of the assessment and those species were classified into Thirty (30) different family. Speckled Pigeon (*Columba guinea*) and Pied Crow (*Corvus albus*) had the highest number of individuals sighted, which were mostly found the roost around buildings' roof. The high numbers of *C.albus* might be as a result of the availability of refuse dumps littering the local governments visited. *C.albus* can be categorized as urban exploiters. These species have adapted to exploiting urban areas as seen in their close association with human habitation and dependence on human food subsidies (Labiran and Iwajomo 2018). All bird species were recorded within the surveyed grids. Few individuals of the endangered Grey Parrot *Psittacus erithacus* (BirdLife International 2019) were recorded during this survey. All other species recorded are currently categorized as Least Concerned under the IUCN Red List (IUCN 2018).

The Kruskal- Wallis H Test on the bird species richness revealed that the mean of Ibadan North Local Government has the highest and was rank first, subsequently followed by Ibadan South West, Ibadan South East, Ibadan North West and Ibadan North East Local Government respectively. The Post Hoc Test also reveal that there were no statistically significant differences in bird species richness in any of the Local Governments. This implies bird species richness does not vary in size of availability from one local government to another. Our study revealed that there were no difference in species richness which might probably due to the fact that all the selected local governments are urban centers and virtually the same ways of life are being practiced as they are all found in the same metropolis.

The Kruskal- Wahills H Test on bird species Abundance within the Local Government revealed that the mean of Ibadan North East has the highest mean rank, followed by Ibadan North, Ibadan South West, Ibadan South East and Ibadan North West Local Governments respectively. The test

of statistics showed that there was no statistically significant difference in bird species abundance between the local governments as a whole. The post Hoc Test showed there were no statistical significant difference between each unit of the local government. This implies bird species abundance does not varies or different in size of availability from one local government to another.

The Test of Between-subjects Effects was carried out to assess the influence of habitat variables on bird species richness and bird species abundance within the metropolis. There was no statistical significant effects/interaction between all the habitat variables and species richness. This implies that habitat variables do not influence bird species richness in Ibadan Metropolis. Our observed results were different compared to Iwajomo et al. (2018), where bird species richness was significantly related to the percentage of ground cover and densities of shrubs and buildings in the study area.

Furthermore, the number of Paved roads and number of Vehicles exacted a significant effect on bird species abundance while others variables under consideration did not exact statistically significant effects on bird species abundance. Generally, bird abundance has been reported to increase in response to increase in urbanization (Tietze and Arise 2018) and this increase has been attributed to the availability of food subsidies and the reduction of predation pressure (Luck and Smallbone 2010).

Thus, this study serves as a baseline to foster future research in how bird diversity is affected by urban ecosystem in Nigeria. Since urban landscapes represent a mosaic of habitats providing diverse opportunities for birds, planning efforts should seek to create and maintain an appropriate balance of habitats that provide the most opportunities for the most species. Also, for successful urban bird conservation, there is need to address the conservation needs of birds, habitat potential of various urban landscape forms, and the needs and motivations of urban residents (Labiran and Iwajomo 2018).

ACKNOWLEDGEMENT

I give all the glory, honor and adoration to the Almighty God who has proved His greatness and support to me beyond measures during and after this research work.

My sincere appreciation goes Prof. I. A. Ayodele for his fatherly advice and concern throughout the period of this research. Profound gratitude also goes to Dr. A. T. Adeyanju for his love, relentless effort and contributions towards the successful completion of this project.

My heartfelt gratitude also goes to my ever loving and supportive parents, Mr and Mrs Adegbola, for their spiritual, moral and financial contributions. I cannot forget my paternal aunt and her husband, Surveyor and Mrs Morawo for their immense support and love. Special regards also goes to Rev. Prof. R.S Babatunde and his family, whom have been a source of support and motivation for me, thanks for the prayers and contributions.

I cannot but give due praises and appreciation to my sisters Abike Oluwadamilola Adegbola, and Oluwamayowa Susan Adegbola, for their numerous contributions at several occasions throughout my stay in the University, may the good Lord bless you more and more. I give special thanks to my cousin and her husband- Engr Bolaji and Mrs Adeola Sadik for their numerous contributions. Oyebamiji Ayokuleyin Uthman, thank you for assisting me to make this work a successful one, I am forever indebted to you. I also acknowledge Azeez Grace Jesuferanmi for her contributions to this research work even at her inconvenience.

To my course mates, the FANTABULOUS CLASS OF 40, you all are the best, I wish us all greater heights and achievements in our future endeavors. The journey would not have been fun-filled and interesting without you all. Thank you for all the good and ugly times we shared, your respect and support really means a lot. I have learnt a lot and I am very grateful. You all will always be remembered.

I also appreciate the following people: Mrs Temidayo Adeyanju, Mr Michael (Love Doctor) and Ibrahim Abdullahi and everybody who have been instrumental in making this project a success.

Thank you all and God bless you.

REFERENCES

- Areola A, Ikporukpo C. 2018. Social Ecology and Urban Green Spaces in Ibadan, Nigeria. *Journal of Applied Sciences and Environmental Management* 22:
- Blair RB. 2001. Birds and butterflies along urban gradients in two ecoregions of the U.S. *In* Lockwood JL, McKinney ML, Eds. *Biotic Homogenization. Norwell (MA): Kluwer*; 33–56.
- Birdlife International. (2019). Birdlife Data Zone. Retrieved December 23, 2019, from Partnership for nature and people website: <http://www.birdlife.org/>
- Gatesire T, Nyiramana DA, Seburanga JL, Mirville MO. 2015. Bird Diversity and Distribution in relation to Urban Landscape Types in Northern Rwanda. Available at <http://www.hindawi.com/journals/tswj/2014/157824/> [Accessed on 20 December 2019]
- Hostetler M. 1997. Scale, birds and human decisions: A potential for integrative research in urban ecosystems. *Landscape and Urban Planning* 45: 15–19.
- Iwajomo S. 2018. Patterns in Bird Species Richness and Abundance in the University of Lagos, Akoka Campus. *Journal of Research in Forestry, Wildlife and Environmental*; 10
- Labiran J, Iwajomo S. 2018. Factors Influencing Bird Species Richness and Abundance in Surulere Local Government Area of Lagos State, Nigeria. *Proceedings of 6th NSCB Biodiversity Conference: Uniuoyo*: pp 183 - 188.
- Lin YP, Chang CR, Chu HJ, Cheng BY. 2011. Identifying the spatial mixture distribution of bird diversity across urban and suburban areas in the metropolis: A case study in Taipei Basin of Taiwan. *Landscape and Urban Planning* 102: 156–163.

378 Luck GW, Smallbone L. 2010. Species diversity and urbanization: patterns, drivers and
379 implications. *Cambridge University Press* 39: 256-289.

380 Magle SB, Hunt VM, Vernon M, Crooks KR. 2012. Urban wildlife research: Past, present and
381 future. *Biological Conservation* 155: 22–32.

382 McCaffrey RE, Mannan RW. 2012. How scale influences birds' responses to habitat features in
383 urban residential areas. *Landscape and Urban Planning* 105: 274–280.

384 McKinney ML. 2002. Urbanization, biodiversity and conservation. *Bioscience* 52: 883–890.

385 Popoola A, Wahab B. 2018. Climate-induced problems and adaptation strategies of urban
386 farmers in Ibadan. *Ethopian Journal of Environmental Studies & Management* 11: 31–42.

387 Salami RO, Von Meding JK, H Giggins. 2016. Assessing habitats of vulnerability in African
388 Cities: A case of poverty outing in Ibadan Metropolis, Nigeria.

389 Sol D, Gonza lez-Lagos C, Moreira D, Maspons J, L, O. 2014. Urbanisation tolerance and the
390 loss of avian diversity. *Ecology Letters* 17: 942–950.

391 Tietze DT (eds), Arise HT. 2018. *Bird Species: How they arise, modify and vanish*. Switzerland:
392 Springer.

393 Walker JS, Balling RC, Briggs JM, Katti M, Warren PS, Wentz EA. 2008. Birds of a feather:
394 Interpolating distribution patterns of urban birds. *Computers, Environment and Urban*
395 *Systems* 32: 19–28.

396 Wolff PJ, Degregorio BA, Rodriguez-Cruz V, Mulero-Oliveras E, Sperry JH. 2018. Bird
397 community assemblage and distribution in a tropical, urban ecosystem of Puerto Rico.

398 *Tropical Conservation Science* 11.

399

400	TABLE OF CONTENTS	
401	Contents	Pages
402	Abstract	1
403	Keywords	1
404	Introduction	2
405	Materials and Methods	4
406	Results	10
407	Discussion	30
408	Acknowledgement	32
409	References	33
410	Tables	36
411	Appendix	38
412	Figure Legends	41

413 **LIST OF TABLES**

414	Table	Title	Pages
415	1a	Descriptive Statistics on Species Richness	11
416	1b	Kruskal- Wallis H Test on Species Richness	12
417	1c	Post Hoc Test on Bird Species Richness	13
418	2a	Descriptive Statistics on Species Abundance	16
419	2b	Kruskal- Wallis H Test on Bird Species Abundance	17
420	2c	Post Hoc Test on Species Abundance	18
421	3	Test of between- Subjects Effects on Species Richness and habitat variables	21
422	4	Test of between- Subjects Effects on Species Abundance and Habitat variables	23

Appendix

SPECIES OF BIRDS OBSERVED AND THEIR GROUPING

	Family	Scientific Name	Species
425			
426	Apodidae	<i>Apus affinis</i>	Little Swift
427	Apodidae	<i>Cypsiurus parvus</i>	African Palm Swift
428	Apodidae	<i>Telacanthura ussheri</i>	Mottled Spinetail
429	Accipitridae	<i>Milvus migrans</i>	Yellow Billed Kite
430	Alcedinidae	<i>Ceyx pictus</i>	African Pygmy Kingfisher
431	Alcedinidae	<i>Alcedo cristata</i>	Malachite Kingfisher
432	Alcedinidae	<i>Halcyon senegalensis</i>	Woodland Kingfisher
433	Alcedinidae	<i>Halcyon malimbica</i>	Blue-breasted Kingfisher
434	Ardeidae	<i>Bubulcus Ibis</i>	Western Cattle Egret
435	Ardeidae	<i>Ardea purpurea</i>	Grey Heron
436	Bucerotidae	<i>Tockus fasciatus</i>	African Pied Hornbill
437	Bucerotidae	<i>Tockus nasutus</i>	African Grey Hornbill
438	Cisticolidae	<i>Cisticola erythrops</i>	Red-faced Cisticola
439	Columbidae	<i>Streptopelia senegalensis</i>	Laughing dove
440	Columbidae	<i>Streptopelia semitorquata</i>	Red-eyed Dove
441	Columbidae	<i>Tutur afer</i>	Blue-spotted Wood Dove
442	Columbidae	<i>Columba guinea</i>	Speckled Pigeon
443	Coraciidae	<i>Eurystomus glaucurus</i>	Broad-billed Roller

444	Corvidae	<i>Corvus albus</i>	Pied Crow
445	Corvidae	<i>Ptilostomus afer</i>	Piacpiac
446	Cuculidae	<i>Centropus senegalensis</i>	Senegal Coucal
447	Cuculidae	<i>Centropus monachus</i>	Blue-headed Coucal
448	Charadriidae	<i>Vanellus spinosus</i>	Spur-winged Lapwing
449	Dicruridae	<i>Dicrurus adsimilis</i>	Forked-tailed Drongo
450	Estrildidae	<i>Estrilda melpoda</i>	Orange-cheeked Waxbill
451	Estrildidae	<i>Logonosticta rufopicta</i>	Bar-breasted Fire Finch
452	Estrildidae	<i>Spermestes cucullatus</i>	Bronze Mannikin
453	Falconidae	<i>Falco tinnuculus</i>	Common Kestrel
454	Falconidae	<i>Falco ardosiaceus</i>	Grey Kestrel
455	Hirundinidae	<i>Hirundo aethiopica</i>	Ethiopian Swallow
456	Hirundinidae	<i>Hirundo senegalensis</i>	Mosque Swallow
457	Hirundinidae	<i>Hirundo rustica</i>	Barn Swallow
458	Hirundinidae	<i>Hirundo fuligula</i>	Rock Martin
459	Laniidae	<i>Corvinella corvina</i>	Yellow-billed Shrike
460	Meropidae	<i>Merops bullockoides</i>	White-fronted Bee Eater
461	Meropidae	<i>Merops persicus</i>	Blue-cheeked Bee-Eater
462	Musophagidae	<i>Crinifer piscator</i>	Western Plantain Eater
463	Motacillidae	<i>Motacilla aguimp</i>	African Pied Wagtail
464	Motacillidae	<i>Anthus leucophrys</i>	Plain-backed Pipit

465	Nectariniidae	<i>Cinnyris chloropygius</i>	Olive Bellied Sunbird
466	Nectariniidae	<i>Cinnyris coccinigastrus</i>	Splendid Sunbird
467	Nectariniidae	<i>Cinnyris Superbus</i>	Superb Sunbird
468	Nectariniidae	<i>Hedydipna collaris</i>	Collared Sunbird
469	Nectariniidae	<i>Cyanomitra olivacea</i>	Olive Sunbird
470	Phasianiidae	<i>Francolinus bicalcaratus</i>	Double-spurred Francolin
471	Phoeniculidae	<i>Malimbus rubricollis</i>	Red-headed Malimbe
472	Passridae	<i>Passer griseus</i>	Northern Grey-headed Sparrow
473	Phoeniculidae	<i>Phoeniculus purpureus</i>	Green Wood-hoopoe
474	Platysteiridae	<i>Platysteira cyanea</i>	Common Wattle-eye
475	Ploceidae	<i>Ploceus cucullatus</i>	Village Weaver
476	Pycnonotidae	<i>Pycnonotus barbatus</i>	Common Bulbul
477	Psittacidae	<i>Poicephalus senegalus</i>	Senegal Parrot
478	Sturnidae	<i>Lamprotornis purpure us</i>	Purple Glossy Starling
479	Sturnidae	<i>Onychognathus fulgidus</i>	Forest Chestnut-winged Starling
480	Sylviidae	<i>Camaroptera brachyuran</i>	Grey-backed Camaroptera
481	Sylviidae	<i>Prinia subflava</i>	Tawny-flanked Prinia
482	Turtidae	<i>Turdus pelios</i>	African Thrush

483

484	FIGURE LEGENDS		
485	Figure	Title	Pages
486	1	Map of Ibadan metropolis	5
487	2	Map of Ibadan Metropolis, Oyo State showing communities	6
488	3	Map showing survey grids	8
489	4	Common Kestrel (<i>Falco tinnunculus</i>) resting on a billboard	24
490	5	Yellow Billed Kite (<i>Milvus aegyptius</i>) brooding in a nest on a billboard	25
491	6	Pied Crow (<i>Corvus albus</i>) hovering in the sky	26
492	7	Yellow Billed Kite (<i>Milvus aegyptius</i>) hovering in the sky	27
493	8	Speckled Pigeon (<i>Columba guinea</i>) in its roost	28
494	9	Laughing Dove (<i>Streptopelia semitorquata</i>) resting on a roof	29