Galbut virus infection minimally influences *Drosophila melanogaster* fitness traits in a strain and sex-dependent manner

Shaun T. Cross^{1, #a, b, c}, Ali L. Brehm¹, Tillie J. Dunham¹, Case P. Rodgers¹, Grace I. Borlee¹, Mark D. Stenglein^{1, *}

¹ Center for Vector-Borne and Infectious Diseases, Department of Microbiology, Immunology, and Pathology, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO, USA

^{#a} Current Address: Cornell Institute for Host-Microbe Interactions and Disease, Cornell University, Ithaca, NY, USA

^{#b} Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, NY, USA

^{#c} Nancy E. and Peter C. Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, USA

* Corresponding author

Email: Mark.Stenglein@colostate.edu (MDS)

Abstract

Galbut virus (family *Partitiviridae*) infects *Drosophila melanogaster* and can be transmitted vertically from infected mothers or infected fathers with near perfect efficiency. This form of super-Mendelian inheritance should drive infection to 100% prevalence, and indeed galbut virus is ubiquitous in wild *D. melanogaster* populations. But on average only about 60% of individual flies are infected. One possible explanation for this apparent paradox is that a subset of flies are resistant to infection. Although galbut virus infected flies seem healthy, infection may be sufficiently costly to drive selection for resistant hosts, thereby decreasing overall prevalence. To test this hypothesis, we quantified a variety of fitness-related traits in galbut virus infected flies from two lines from the *Drosophila* Genetic Reference Panel (DGRP). Galbut virus infected flies had slightly decreased average lifespan and total offspring production, but these decreases were mostly not statistically significant. Galbut virus DGRP-517 flies pupated and eclosed slightly faster than their uninfected counterparts. Some galbut virus infected flies exhibited altered sensitivity to viral, bacterial, and fungal pathogens. Galbut virus infection produced minimal changes to host mRNA levels as measured by RNA sequencing, consistent with minimal phenotypic changes. The microbiome composition of flies was not measurably perturbed by galbut virus infection. Differences in phenotype attributable to galbut virus infection varied as a function of fly sex and DGRP strain and were generally dwarfed by larger differences attributable to strain and sex. Thus, galbut virus infection does produce measurable phenotypic changes, with changes being minor, offsetting, and possibly net negative.

Importance

Virology has largely focused on viruses that cause unmistakable phenotypic changes. But metagenomic surveys are revealing that persistent virus infections are extremely common, even in apparently healthy organisms. The extent to which these persistent viruses impact host fitness and evolution remains largely unclear. Here we study fitness impacts of a partitivirus named galbut virus that is ubiquitous in wild *D. melanogaster* populations. Despite efficient biparental vertical transmission, galbut virus is present in only just over half of wild flies. We quantified various fitness-related traits in galbut virus infected and uninfected fly lines and found that infection produced small but measurable changes in host phenotype that in aggregate may reduce fly fitness. Further studies that take advantage of this virus that naturally infects a premier model organism that is easy to study in the wild will shed further light on the persistent virus-host dynamics that may actually represent most viral infections.

Keywords: Galbut virus, *Drosophila melanogaster*, DGRP, arthropod, virus-host interactions, fitness, virome, partitivirus

Running Title: Fitness impacts of galbut virus infection

Introduction

Galbut virus is a remarkably successful persistent virus of *Drosophila melanogaster* [1–4]. Infected flies have been found on five continents and every wild population that has been tested includes some infected individuals. This degree of success is attributable to efficient biparental vertical transmission: infected mothers or infected fathers can transmit galbut virus to 100% of offspring, providing a means for infection to increase in frequency generation over generation [4,5].

Galbut virus belongs to a group of viruses, the partitiviruses (*Partitiviridae*), generally known for mild persistent infections [6–9]. Plant-infecting partitiviruses were originally called cryptic viruses (former genus *Cryptovirus*) because of their inapparent phenotypic effects [9,10]. Galbut virus infection is similarly cryptic: despite a century of *Drosophila* research and despite its ubiquity, galbut virus was only recently discovered by shotgun metagenomics [1]. Galbut virus infected flies do not exhibit obvious phenotypic differences from their uninfected counterparts, and in a population of wild-caught flies that we have maintained for three years, galbut virus has risen to and remained at 100% prevalence [4].

Nevertheless, there are indications that galbut virus may be in conflict with its host. Although infection is ubiquitous, the fraction of infected flies ranges from 13-100% in different populations, and on average only ~60% of flies are infected [1,3,4]. We also found that some *Drosophila* Genetic Reference Panel (DGRP) lines were relatively refractory to infection and multigenerational vertical transmission [4,11]. Galbut virus sequences exhibited high ratios of non-synonymous to synonymous variation (high dN/dS ratios), which could be consistent with selection driven by ongoing host-virus conflict [1]. Verdadero virus, a partitivirus that infects *Aedes aegypti*, exhibited similarly efficient biparental vertical transmission in colonized mosquitoes, but also is not at 100% prevalence in wild populations [12–14].

We hypothesized that galbut virus might exact a fitness cost that is small but sufficient to drive selection for resistant individuals. This could limit the overall success of galbut virus – and similar persistent viruses – in host populations. To test this hypothesis, we quantified a number of fitness-linked phenotypes in galbut virus infected flies. Host genotype and sex are variables that can substantially influence the outcome of infection [15–20], so we evaluated the phenotype of galbut virus infected males and females from two different DGRP strains.

Relatively little is known about arthropod-infecting partitiviruses, which have now been identified in association with a broad range of hosts including disease vectors [21–27]. Metagenomic surveys of apparently healthy free-living organisms have in general produced a flood of new virus sequences [12,28–30]. But beyond sequence description and phylogenetic placement, little is known about the biological impact of all of these newly recognized viruses. Follow-up virological studies that build upon this trove of sequence data are needed [4,27,31,32]. The ability to study a highly successful natural virus of a premier model organism represents a great opportunity to shed light on insect-infecting partitiviruses and persistent viral infections more generally.

Methods and Materials

***Drosophila* rearing and maintenance.** Flies were reared on the Bloomington *Drosophila* Stock Center (BDSC) standard cornmeal diet (<https://bdsc.indiana.edu/information/recipes/bloomfood.html>). Stocks were housed at 25°C and changed every 14 days. All experiments were performed with *Drosophila* Genetic Reference Panel (DGRP) stocks 399 and 517, acquired from the BDSC [11,33]. Generation of galbut virus infected lineages by microinjection was described previously [4].

Experimental groups consisted of galbut virus infected or uninfected DGRP 399 or DGRP 517 males or females (2 strains x 2 sexes x 2 galbut virus infection status = 8 groups). All flies were 3-5 day old virgins reared in a 12 hour light/dark cycle at 25°C, unless otherwise stated.

Quantification of galbut virus RNA levels. Total RNA was extracted from 5 day old, virgin flies using a bead-based protocol as previously described [4]. cDNA was synthesized by adding 5.5 µl of RNA to 200 pmol of a random 15-mer oligonucleotide and incubated for 5 min at 65°C, then set on ice for 1 min. A reverse transcription (RT) mixture containing the following was added (10 µL reaction volume): 1x SuperScript III (SSIII) FS reaction buffer (Invitrogen), 5 mM dithiothreitol (Invitrogen), 1 mM each deoxynucleotide triphosphates (dNTPs) (NEB), and 100 U SSIII reverse transcriptase enzyme (Invitrogen), then incubated at 42°C for 30 min, 50°C for 30 min, then at 70°C for 15 min. 90 µL of nuclease-free H₂O was added to dilute the cDNA to a final volume of 100 µL.

Quantification of major microbiome constituent DNA levels. Total DNA was extracted from 4-5 day old virgin flies. 10 flies per pool (total of 3 pools per group) were surface sterilized by vortexing in 70% ethanol for 2 minutes, followed by 2 rinses with autoclaved ddH₂O and vortexing for 1 minute. Flies were then stored at -80°C until DNA was extracted. DNA was extracted using the DNeasy Tissue and Blood extraction kit (Qiagen) following the manufacturer's protocol for insect tissues with three modifications. First, samples were added to 180 µL ATL buffer (provided in kit) along with a single BB bead and homogenized using a Qiagen TissueLyzer for 3 minutes at 30Hz rather than homogenizing by hand. Second, samples were incubated in proteinase K for a duration of 4 hours. Last, following incubation with proteinase K, samples were treated with 20 µL of RNase A (2 mg/mL; Sigma Aldrich) for 30 min at 37°C. After RNase treatment, samples were processed as stated in the protocol.

Following cDNA synthesis or DNA extraction, qPCR reactions were set-up using Luna qPCR Master Mix (NEB) following the manufacturer's protocol. The qPCR reaction was performed on LightCycler 480 (Roche) under the following protocol: 95°C for 3 min, 40 cycles of 95°C for 10s, then 60°C for 45s, and then followed by a melting curve analysis. Microbiome analysis primer sequences were predominately from Early et al. [34]. Primer sequences can be found in **S1 Table**.

Lifespan and fecundity assays were performed similar to as previously described [35]. Flies were reared in 5 replicate groups of 10 adults (5 female, 5 male). Flies were checked daily for survival of adults, and living adults were moved to fresh media every 3 days. Longevity of adults was compared using the R survival package [36]. After adults were moved, original vials containing laid eggs were kept for 14 days, after which offspring were counted and sexed.

Total egg production was measured by housing 10 male and 10 female flies in bottles with an apple agar plate coated with yeast paste (1:1 yeast and water) to promote egg laying. Egg plates were replaced every

24 hours, and the used plates containing eggs were frozen at -20°C until the eggs were counted. Plates were collected for a total of 3 days, and were performed in 3 biological replicates and 2 technical replicates. Images of egg plates were captured and eggs were counted manually using the ImageJ cell counter program [37]. All fecundity measurements were analyzed with R scripts that can be found at: https://github.com/scross92/galbut_fitness_analysis.

Developmental speed assays were performed as previously described [38]. Eggs were collected using standard apple agar plates without preservative, with a mixture of 1:1 yeast and water applied. Every hour for 6-8 hours, agar plates were discarded and replaced to encourage egg synchronization. Agar plates were replaced a final time and incubated for several hours. The plates were removed and eggs were collected using an autoclaved brush. Twenty eggs were collected and moved to non-nutritive agar plates containing 5% sucrose/2% agar with no antimicrobials added (no tegosept). An agar plate was placed inside a larger petri dish with a damp paper towel on the bottom and moved to a 25°C incubator with a 12 hour light/dark cycle. Every 2 days, yeast paste was added as a nutrition source for developing flies. Yeast were killed prior to use in the paste by microwaving for 45 seconds on high to prevent overgrowth. Plates were checked daily for pupae to determine speed of pupation. Once pupation began, plates were checked approximately every 5 hours (morning, midday, evening). Continual monitoring occurred from pupation to emergence of adults in the same ~5 hour increments hours for measuring total time it took for flies to reach the adult stage. This was performed in 6 replicates per group (strain and galbut virus infection status).

***Pseudomonas aeruginosa* oral challenge.** Flies were challenged orally with *Pseudomonas aeruginosa* as adapted from Lutter et al [39]. An overnight culture of *P. aeruginosa* (PAO1) was grown in a 200 mL culture Brain Heart infusion (BHI) broth incubated at 220 rpm at 37°C. The following day, the culture was centrifuged at 4200 g for 5 minutes until a loose pellet was formed. Excess supernatant was decanted and culture was resuspended to an OD_{600nm} of ~7 using a sterile 5% sucrose solution. Autoclaved filter disks were inoculated with 290 µL of the *P. aeruginosa* solution. Disks were placed on 5% sucrose agar vials. Control disks were inoculated with the 5% sucrose solution. Twelve flies that had been starved for 5 hours were placed in the bacteria-containing vials for each replicate. Flies that died by the end of the first day were censored from further analysis, since their deaths were likely due to starvation stress. Survival of flies was monitored daily for 12 days. Statistical analysis was performed using the R survival package [36]. A total of 3 technical replicates were performed.

Intrathoracic microbial pathogen challenges. The following pathogen challenges were performed through intrathoracic microinjection. All experimental injections were performed in 3 biological replicates (12 flies per replicate) per technical replicate, and a total of two technical replicates were performed for each pathogen. An exception is the *Staphylococcus aureus* challenge which was performed in 3 technical replicates. Control injections with 1x phosphate buffered saline (PBS) were performed in parallel. Flies were checked at 10-12 hours post-injection, and any flies that were dead at this point were assumed to have died from injection. Additionally, any flies that died from non-natural causes (for example, after getting stuck in the media) were also censored from analysis. Injected volumes, inoculum dose, and subsequent intervals for checking fly survival are stated below for the respective pathogen.

Pseudomonas aeruginosa: Flies were microinjected with *P. aeruginosa* (strain PAO1). A culture was started by inoculating 150mL of BHI broth and incubated at 220 rpm overnight at 37°C. The following day, the culture was centrifuged at 4200 g for 5 minutes until a loose pellet was formed. Excess supernatant was decanted and the culture was resuspended to an OD_{600nm} of 0.03 using 1x PBS. Flies were injected with 9.2 nL of this diluted *P. aeruginosa* culture, which corresponds to ~100 CFUs [40]. Flies were incubated overnight and checked at 24 hours post injection, 28 hours post injection, and every 2 hours from 28 to 42 hours post injection. After 42 hours post injection, flies were checked at one final time point of 52 hours post injection, at which any living flies were censored from downstream statistical analyses.

Staphylococcus aureus: Flies were microinjected with *S. aureus* (strain XEN36, Perkin Elmer). A culture was obtained by inoculating 150mL BHI broth and stirred at 220 rpm overnight at 37°C. The following day, the culture was centrifuged at 4200 g for 5 minutes until a loose pellet was formed. Excess supernatant was decanted and the culture was resuspended to an OD_{600nm} of 0.1 using 1x PBS. Flies were injected with 23 nL of this diluted *S. aureus* culture, which corresponds to ~100 CFUs [41]. Flies were checked daily until 8 days post injection, at which point any living flies were censored from downstream statistical analyses.

Drosophila C virus: *Drosophila C virus* (DCV) stocks were provided by the Andino lab at the University of California San Francisco. DCV stocks were amplified and titrated on *Drosophila* S2 cells. DCV infections of flies were performed as previously described [42]. Flies were microinjected with DCV at a titre of 100 50% Tissue Culture Infective Dose units (TCID₅₀) in a total volume of 50 nL. Flies were checked daily until 14 days post injection, at which point any living flies were censored from downstream analyses.

Candida albicans: *Candida albicans* challenge was performed as previously described [43]. *C. albicans* (strain SC5314) was obtained from ATCC. A yeast extract peptone dextrose (YPD) agar plate was streaked from the frozen glycerol stock and incubated at 30°C for 18 hours. 150mL of YPD broth was inoculated with a single colony from the YPD plate and incubated at 220 rpm overnight at 30°C until the culture was at an OD_{600nm} of ~1. The culture was centrifuged at 4200 g for 5 minutes until a loose pellet was formed, which was resuspended using 1x PBS. Yeast cells were counted with a cytometer and diluted to 10⁷ cells/mL. Flies were microinjected with 50 nL (~500 cells) of this dilution. Flies were incubated at 30°C and were checked daily until 6 days post injection, at which point any living flies were censored from analyses.

RNAseq library preparations. Pools of 10, 5-day old, virgin flies were collected, flash frozen in liquid nitrogen and stored at -80°C. Polyadenylated RNAs were enriched using the NEB Magnetic mRNA Isolation Kit according to the manufacturer's protocol. Sequencing libraries were created using Kapa HyperPrep RNA Library Prep Kit (Roche) according to manufacturer's protocol. Final library molecules had an average size of 348 base pairs, and were sequenced on an Illumina NextSeq using the NextSeq 500/550 High Output Kit v2.5, generating 75 base single-end reads.

Transcriptomic computational analysis. RNAseq datasets were first processed to remove low quality and adapter sequences using cutadapt tool [44] version 1.13 with the following settings: -a

AGATCGGAAGAGC -A AGATCGGAAGAGC -g GCTCTTCCGATCT -G GCTCTTCCGATCT -a
AGATGTGTATAAGAGACAG -A AGATGTGTATAAGAGACAG -g CTGTCTCTTATACACATCT -
G CTGTCTCTTATACACATCT, -q 30,30, --minimum-length 40, and -u 1. Remaining reads were
mapped to the *D. melanogaster* genome assembly BDGP6.28 from Ensembl using HISAT2 version 2.2.0
[45]. Read mapping was tabulated using featureCounts version 2.0.0 [46] to the BDGP6.28 gtf file with
the following settings: -s 2 -t exon -g gene_id. The resulting read count table was used as input for
differential gene expression analysis using DESeq2 version 1.26.0 [47] in R version 3.6.3 [48].
Differential gene expression analyses on the condition of galbut virus infection status and was performed
for each group (strain and sex). Gene set enrichment analyses (GSEA) were performed using the
clusterProfiler R package version 3.14.3 [49] using a pre-ranked gene list ordered by the log2 fold
changes and the 'gseGO' function. Redundant GO terms were collapsed using the 'simplify' function by
adjusted p values, with a cutoff value of 0.7, and the "Wang" measure.

Data Availability. Computational scripts for analysis of experiments can be found at
https://github.com/scross92/galbut_fitness_analysis. Raw sequencing data can be found on the NCBI
Sequence Read Archive (SRA) under Bioproject PRJNA683038.

Results

Confirmation of galbut virus infection status and galbut virus RNA levels in individual flies.

We first verified that our stocks of galbut virus-infected flies established previously were still persistently
infected [4]. We quantified galbut virus RNA levels using qRT-PCR in 20, 3-5 day old flies from each
line (10 male and 10 female), and normalized levels to those of ribosomal protein L32 mRNA (RpL32;
Fig 1). Galbut virus RNA levels were higher than those of highly expressed RpL32 mRNA in all cases
(**Fig. 1**). Median galbut virus RNA levels were 2.3-fold higher in DGRP 399 flies than in DGRP 517 flies
($p=1.6 \times 10^{-2}$), although some outlier DGRP 517 flies had higher levels (**Fig 1**). Galbut virus RNA levels
were 2.1-fold higher in DGRP 399 males than in females ($p=4.2 \times 10^{-5}$) and 1.5-fold higher in DGRP 517
males than in females ($p=1.3 \times 10^{-2}$). So, these populations remained persistently infected at 100%
prevalence, and galbut virus RNA levels varied as a function of DGRP strain and sex, with levels
generally higher in males and in DGRP 399 flies.

Fig 1. Relative galbut virus RNA levels in flies from two DGRP strains. Boxplots depicting galbut virus RNA levels relative to the housekeeping gene Rpl32 ($2^{-\Delta Ct}$ method) in DGRP 399 and DGRP 517 adult flies from strains used in this study (n = 10 per strain and sex).

Galbut virus infection does not have significant impacts on predominant microbiome constituents.

The microbiome composition of *D. melanogaster* can alter fitness [35]. It's also possible that viral and bacterial constituents of the microbiota can interact [50]. Commensal bacteria can also vary by DGRP background when reared under the same conditions [34]. We therefore tested whether the microbiomes of these populations varied as a function of galbut virus infection status. Our goals were to assess whether microbiome differences could underlie differential phenotypes in flies with and without galbut virus, and whether galbut virus infection was altering microbiome composition.

Previous shotgun metagenomic RNA sequencing of DGRP 399 and 517 flies reared in our lab had revealed that the major constituents of these flies' microbiomes included *Acetobacter persici*, *Lactobacillus brevis*, *L. planatarum*, *Corynebacterium* spp., and *Saccharomyces cerevisiae*. We quantified DNA levels of these microbes by qPCR using previously designed primers. DNA copy numbers were normalized to the single copy host gene *deformed* (*dfd*) as previously described [34]. The relative abundances of the different microbes were similar in DGRP 399 and 517 flies and in males and females (**Fig. 2**). Galbut virus infection did not produce any statistically significant differences in DNA levels of these taxa in any of the groups (**Fig 2**). This indicated that galbut virus infection did not measurably change microbiome composition and that any fitness effects of galbut virus infection were likely not mediated by changes in microbiome composition.

Fig 2. Galbut virus infection did not alter levels of major microbiome constituents in flies. Relative amounts of DNA from predominant microbiome constituents in galbut virus infected and uninfected flies were measured via qPCR from 3 replicate pools of 10 flies per pool per strain per sex. Mean DNA loads relative to single copy *dfd* gene are plotted and crossbars indicate standard deviations of replicates. Dotted lines depict linear regression fits and solid lines indicate the diagonal. No statistically significant differences between galbut virus infected and uninfected flies were identified using a Wilcoxon test.

Galbut virus slightly reduces *Drosophila* lifespan and fecundity

We compared the lifespan, fecundity, and developmental speed of galbut virus infected and uninfected flies [35,51]. Vials of newly eclosed adults (n=5 replicate vials per experimental group) were housed together in groups of 10 flies (5 males, 5 females). Galbut virus infected flies from both strains exhibited a slightly shortened mean lifespan (6.6 days shorter in DGRP 399 and 3.1 days shorter in DGRP 517), though this was not statistically significant in either strain (**Fig 3A**). These decreases in lifespan attributable to galbut virus infection status were smaller than differences attributable to DGRP strain: DGRP 517 flies lived on average 14.9 fewer days than DGRP 399 flies ($p=2.1 \times 10^{-4}$, **Fig 3B**).

Fig 3. Galbut virus infected flies exhibit slightly reduced lifespan. (A) Survival of DGRP 399 and 517 flies with or without persistent galbut virus infection. The mean and standard deviation of biological replicates is plotted. (B) Data as in A, but plotted to facilitate comparison of DGRP strains.

We compared fecundity of infected and uninfected flies by counting total adult offspring in vials containing 5 male and 5 female flies. Galbut virus infected DGRP 399 flies produced fewer offspring than their uninfected counterparts, but this was not significantly different (t-test; female offspring $p=0.77$, male offspring $p=0.83$; **Fig 4A**). Galbut virus infected DGRP 517 flies also produced fewer offspring, but the decrease was only significant for male offspring numbers (t-test; female offspring $p=0.16$, male offspring $p=0.027$; **Fig 4A**). The total offspring counts were not normalized to surviving mothers, so the lower number of offspring at later timepoints likely reflect the slightly shortened lifespan of galbut virus infected flies (**Fig 3**). Galbut virus infection did not significantly change offspring sex ratios (t-test;

DGRP 399: $p=0.63$, DGRP 517: 0.75 , **Fig. S1**). As with average lifespan, differences in total offspring number between the different DGRP strains were much larger than differences attributable to galbut virus infection status: DGRP 399 females produced on average 2.7x more offspring than DGRP 517 females ($p=2.8 \times 10^{-6}$, **Fig 4B**).

We recorded the cumulative number of eggs laid over three days when one or both parents were infected by galbut virus. There were no significant differences in the number of eggs laid when either or both parents were infected with galbut virus (ANOVA, DGRP 399: $p=0.85$, DGRP 517: $p=0.72$; **Fig 4C**).

Fig 4. Galbut virus infected flies produce slightly fewer offspring. (A) Galbut virus infected and uninfected flies were housed in batches of 5 males and 5 females per vial and cumulative number of female and male offspring per vial were counted. The mean and standard deviation of biological replicates are plotted. (B) Data as in panel A, but plotted for comparison of DGRP strains (data from galbut virus uninfected flies shown). (C) 10 male and 10 female flies 3-5 days post eclosion were crossed with different combinations of galbut virus infected mothers or fathers. The cumulative number of eggs laid per female over three days is depicted for individual replicates as points and summarized with boxplots.

We compared the developmental speed of galbut virus infected and uninfected flies by collecting eggs and monitoring the times from oviposition to pupation and oviposition to adulthood. DGRP 399 flies pupated in ~5 days and eclosed in ~9 days regardless of galbut virus infection status (**Fig 4A & 4C**). DGRP 517 flies infected with galbut virus pupated on average 7 hours faster than uninfected flies (Wilcoxon, $p=2.2 \times 10^{-16}$; **Fig 4A**). DGRP 517 infected females and males reached adulthood on average 10 and 12 hours faster than their uninfected counterparts (**Fig 4C**). As was the case with other phenotypes, development speed also varied as a function of DGRP background, with DGRP 399 flies pupating on average 7 hour faster than DGRP 517 flies (**Fig 4B**) and eclosing on average 13 hours faster (**Fig 4D**).

Fig 5. Galbut virus infected DGRP 517 flies develop slightly faster. (A) The time to pupation of individual DGRP 399 or DGRP 517 flies is plotted and the median time is indicated by a crossbar. (B) Data as in A, but plotted to enable comparison between DGRP strains. (C) The time between oviposition and eclosion for individual flies is indicated. (D) Data as in C, but plotted to enable comparison between DGRP strains. ns: not significant; **: $p < 0.01$, ***: $p < 0.001$

Galbut virus alters the susceptibility of flies to viral, bacterial, and fungal pathogens.

The microbiota present in a host can influence the outcome of subsequent infections [52–55]. Moth-infecting partitiviruses changed their host's ability to withstand infection by a pathogenic nucleopolyhedrovirus [27]. We hypothesized that galbut virus infection might alter the ability of flies to resist or tolerate infection by pathogenic microbes, which could alter the survival and consequently the

fitness of galbut virus infected flies. To test this hypothesis, we challenged galbut virus infected and uninfected flies with viral, bacterial, and fungal pathogens.

We first tested whether pre-existing galbut virus infection altered fly survival following infection by *Drosophila C virus* (DCV) [56]. Flies were challenged with 100 TCID₅₀ units of DCV through intrathoracic microinjection and checked daily for survival. Overall, there was little difference in the survival of galbut virus infected and uninfected flies. DGRP 517 female galbut virus-infected flies survived slightly longer than their uninfected counterparts, and although this effect was statistically significant, it was small in magnitude (**Fig 6A**, $p=0.028$). These DGRP strains are both *Wolbachia* negative, so improved survival could not be attributed to the known protective effects of *Wolbachia* against DCV [57–60].

We next challenged flies orally with *Pseudomonas aeruginosa*. Galbut virus infected DGRP 399 female flies were more susceptible to *P. aeruginosa* bacterial challenge (**Fig 6B**; $p=4.5 \times 10^{-6}$). Although ingestion is a more natural route of infection than microinjection, there is less experimental control over the ingested dose, which can decrease reproducibility [40]. We therefore also injected flies with ~100 CFUs of *P. aeruginosa*. Flies injected with *P. aeruginosa* died faster than those that ingested the pathogen, with most flies dead by 36 hours post injection (**Fig. 6C**). Galbut virus infected DGRP 399 females no longer died faster than their uninfected counterparts when microinjected with *P. aeruginosa* (**Fig 6C**; $p=0.14$). This suggests that interactions between galbut virus and *P. aeruginosa* may depend on the route of infection.

Since the *Drosophila* innate immune system responds differently to Gram negative and Gram positive bacteria [61], we continued our pathogen challenges by microinjecting flies with *Staphylococcus aureus*. When flies were microinjected with ~100 CFUs, galbut virus infected DGRP 399 male flies survived slightly longer than their galbut virus infected counterparts ($p=.021$) as did DGRP 517 females ($p=6.8 \times 10^{-4}$). As for DCV challenge, although these effects were statistically significant, they were small in magnitude.

As a final pathogen challenge, we injected flies with ~500 cells of the fungal pathogen *Candida albicans* [43]. Both male and female DGRP 399 galbut virus infected flies died faster than their uninfected counterparts following *C. albicans* challenge (**Fig 6E**; DGRP female $p=6.5 \times 10^{-6}$ and DGRP male $p=3.5 \times 10^{-5}$). No significant differences were observed for DGRP 517 flies (**Fig 6E**).

Fig 6. Galbut virus alters pathogen susceptibility of some flies. Survival of galbut virus infected and uninfected flies following (A) intrathoracic injection with 100 TCID₅₀ units of *Drosophila C virus* (DCV),

(B) ingestion of *Pseudomonas aeruginosa*, (C) injection of ~100 CFUs of *Pseudomonas aeruginosa*, (D) injection with ~100 CFUs of *Staphylococcus aureus*, (E) injection with ~500 *Candida albicans* cells. (F-H) Survival of flies following control inoculations. Flies were either microinjected with phosphate buffered saline (PBS) and stored at 25°C (F) or 30°C (G), or ingested sucrose (H). Galbut virus infected flies are depicted in orange and uninfected flies are in blue. ns: not significant; *: $p < 0.05$; **: $p < 0.01$, ***: $p < 0.001$.

Galbut virus induces strain and sex specific changes in the transcriptome.

We used RNA sequencing (RNA-seq) to explore transcriptional changes that could underlie the observed phenotypic differences between galbut virus infected and uninfected flies. We sequenced mRNA from pools of 10 whole adult females or males. We first performed hierarchical clustering to assess similarity between gene expression profiles in all datasets (**Fig. 7**). Sex was by far the most important variable influencing gene expression patterns: datasets from males and females were completely separated, with long branch lengths separating the clusters. This separation likely reflects the different chromosome repertoires of males and females in addition to sex-specific expression differences. Male datasets then clustered by DGRP strain, with DGRP 399 and 517 males forming separate subclusters. In DGRP 399 males, the group with the highest galbut virus RNA levels (**Fig. 1**), galbut virus infected and uninfected flies formed discrete clusters. In DGRP 517 males, the separation of infected and uninfected flies was not as clean. Gene expression patterns of female flies did not form subclusters based on DGRP strain nor galbut virus infection status (**Fig. 7**). Thus, as with other phenotypes, fly sex and DGRP strain influenced gene expression more than galbut virus infection.

Only a single gene exhibited significant differential expression (adjusted p-value (padj) < 0.05) in all flies when compared by galbut virus infection status alone. The gene was a ribosomal RNA pseudogene (28S ribosomal RNA pseudogene CR45851) and it was upregulated in all groups of infected flies. We therefore examined transcriptional responses in flies grouped by DGRP strain and sex. Within these subsets, the response to galbut virus infection varied by both the number of differentially expressed genes and those that passed a significance threshold (**Fig 8**). Given a lack of consistent fitness phenotypes across any one sex or strain in most cases, this may be unsurprising.

Fig 7. Galbut virus exerts minimal impact on overall transcriptional responses in flies. A sample distance matrix (Euclidean distances) quantifying the similarity between gene expression patterns in all datasets. Rep: biological replicates of 10 flies per replicate.

Fig 8. Volcano plots of differential gene expression in galbut virus infected flies. Plots depict the relative fold change of individual genes in galbut virus infected flies relative to uninfected flies (positive fold-change values indicate higher expression levels in galbut virus infected flies) on x axes and multiple testing corrected p-values on y axes. Individual genes that have a log2 fold change greater than 1 (orange), an adjusted p-value < 0.05 (gold), or both (green) are colored.

Among the top upregulated and downregulated genes in the experimental groups, few genes were shared between groups. Genes that were differentially expressed in more than one group included Kruppel homolog 1 (*Kr-h1*), which was significantly downregulated in galbut virus infected DGRP 399 and 517 females (**Fig 9**). This gene is a transcriptional regulator that has links to development [62–64]. Formin homology 2 domain containing (*Fhos*), which functions in development (remodeling of muscle cytoskeleton) and immune response (directs macrophage movement), was downregulated in both DGRP 399 infected females and males [65–67]. It is possible that these changes are related to the differences in developmental speed and pathogen susceptibility that we had observed (**Fig 4B; Fig 6B, E**).

Two genes with limited functional information were differentially expressed in both sexes of one or the other of the DGRP strains. Glycogen binding subunit 76A (*Gbs-76A*) was downregulated in galbut virus infected DGRP 399 flies of both sexes. This gene is inferred to play a role in the glycogen biosynthesis

pathway [68]. In male and female DGRP 517 galbut virus infected flies, gene CG14186 was downregulated. CG14186 is affiliated with the biological process of cilium assembly, but its molecular function is unknown.

Two genes that were among the list of top differentially regulated genes across groups, but in opposite directions, were CG17560 and Heat shock protein 70Bb (*Hsp70Bb*). CG17560 is predicted to have implications in metabolic processes [68]. In DGRP 399 infected females, this gene was upregulated, while in DGRP 517 infected females, it was downregulated (**Fig 9**). *Hsp70Bb* was downregulated in DGRP 399 infected females, but was upregulated in DGRP 517 infected males (**Fig 9**). Heat shock proteins accounted for a large fraction of the upregulated genes in DGRP 517 infected males. Heat shock proteins are upregulated under heat and chemical stress, but these proteins have additional antiviral functions [69].

Fig 9. Top differentially expressed genes in infected flies relative to uninfected flies as. (A) Top 15 most significantly upregulated genes (ranked by padj, with padj < 0.05) in each experimental group relative to uninfected flies (B) Top 15 most significantly downregulated genes in each experimental group relative to uninfected flies.

We performed a gene set enrichment analysis (GSEA) with the genes pre-ranked by log2 fold change using the clusterProfiler package in R [49] (Figs S2-4, Tables 1-2). Among the most significantly enriched gene ontology (GO) pathways (biological process ontologies only), pathways associated with development, morphogenesis, and metabolism were positively enriched in infected DGRP 517 flies (Table 1, S2-3 Fig). GO pathways associated with neuron development and differentiation and response to stimuli were also differentially regulated (Table 1, Figs S2-4). GO pathways under the parent GO terms reproduction (GO:0000003) and reproductive process (GO:0022414) were positively enriched in galbut virus infected flies (Tables 1 and 2). Between 9 and 35 GO pathways associated with reproduction were enriched in galbut virus infected DGRP flies. All differentially regulated pathways, except 2 (DGRP

399 females, GO:0046008; DGRP 517 males, GO:0051446), were positively enriched in infected flies compared to uninfected flies.

Table 1. Most significantly enriched gene ontology pathways (biological process ontologies only) within infected flies identified via gene set enrichment analysis. NES: Normalized enrichment score; FDR: False Discovery Rate q value (Benjamini and Hochberg adjustment); *: $q < 0.05$, **: $q < 0.01$, ***: $q < 0.001$.

Experimental group (infected)	GO term	Description	NES	FDR q value
DGRP 399 Females	GO:0006270	DNA replication initiation	2.02	**
	GO:0010529	negative regulation of transposition	1.99	**
	GO:0060856	establishment of blood-brain barrier	1.98	**
	GO:1902475	L-alpha-amino acid transmembrane transport	1.96	**
	GO:0042335	cuticle development	-1.65	**
	GO:0007606	sensory perception of chemical stimulus	-1.76	**
	GO:0040003	chitin-based cuticle development	-1.88	**
	GO:0050911	detection of chemical stimulus involved in sensory perception of smell	-2.00	**
	GO:0009593	detection of chemical stimulus	-2.07	**
	GO:0050907	detection of chemical stimulus involved in sensory perception	-2.13	**
DGRP 399 Males	GO:0009887	animal organ morphogenesis	1.98	***
	GO:0030182	neuron differentiation	1.92	***
	GO:0051128	regulation of cellular component organization	1.87	***
	GO:0003006	developmental process involved in reproduction	1.87	***
	GO:0010604	positive regulation of macromolecule metabolic process	1.73	***
	GO:0010605	negative regulation of macromolecule metabolic process	1.72	***
	GO:0009892	negative regulation of metabolic process	1.72	***
	GO:0051173	positive regulation of nitrogen compound metabolic process	1.71	***
	GO:0031325	positive regulation of cellular metabolic process	1.70	***
	GO:0051649	establishment of localization in cell	1.58	***
DGRP 517 Females	GO:0051128	regulation of cellular component organization	2.23	**
	GO:0003006	developmental process involved in reproduction	2.21	**
	GO:0048699	generation of neurons	2.19	**
	GO:0051239	regulation of multicellular organismal process	2.14	**
	GO:0023051	regulation of signaling	2.04	**
	GO:0010646	regulation of cell communication	2.03	**
	GO:0009893	positive regulation of metabolic process	2.02	**
	GO:0031325	positive regulation of cellular metabolic process	2.01	**
	GO:0009892	negative regulation of metabolic process	1.95	**
	GO:1901575	organic substance catabolic process	1.58	**
DGRP 517 Males	GO:0009887	animal organ morphogenesis	2.04	***
	GO:0010604	positive regulation of macromolecule metabolic process	1.85	***
	GO:0051239	regulation of multicellular organismal process	1.84	***
	GO:0031325	positive regulation of cellular metabolic process	1.83	***
	GO:0051173	positive regulation of nitrogen compound metabolic process	1.82	***
	GO:0009892	negative regulation of metabolic process	1.82	***

GO:0051128	regulation of cellular component organization	1.81	***
GO:0050793	regulation of developmental process	1.80	***
GO:0010605	negative regulation of macromolecule metabolic process	1.80	***
GO:0051649	establishment of localization in cell	1.44	***

Table 2. Enriched gene ontology pathways (biological process ontologies only) associated with reproduction identified via gene set enrichment analysis. NES: Normalized enrichment score; FDR: False Discovery Rate q value (Benjamini and Hochberg); *: $q < 0.05$, **: $q < 0.01$, ***: $q < 0.001$.

Experimental group (infected)	GO term	Description	NES	FDR q value	Within both strains?
DGRP 399 Females					
	GO:0030703	eggshell formation	2.48	**	+
	GO:0007304	chorion-containing eggshell formation	2.48	**	+
	GO:0007306	eggshell chorion assembly	2.37	**	+
	GO:0007292	female gamete generation	2.34	**	+
	GO:0030707	ovarian follicle cell development	2.31	**	+
	GO:0007548	sex differentiation	2.20	**	+
	GO:0009994	oocyte differentiation	2.12	**	+
	GO:0007143	female meiotic nuclear division	2.10	**	+
	GO:0046843	dorsal appendage formation	2.10	**	+
	GO:0048599	oocyte development	2.04	**	+
	GO:0048608	reproductive structure development	2.04	**	+
	GO:1903046	meiotic cell cycle process	2.02	**	+
	GO:0051321	meiotic cell cycle	2.00	**	+
	GO:1905879	regulation of oogenesis	1.96	**	+
	GO:0007308	oocyte construction	1.95	**	+
	GO:0007309	oocyte axis specification	1.91	**	+
	GO:0007293	germarium-derived egg chamber formation	1.88	**	—
	GO:0140013	meiotic nuclear division	1.87	**	+
	GO:2000241	regulation of reproductive process	1.84	**	+
	GO:0061988	karyosome formation	1.81	**	—
	GO:0061982	meiosis I cell cycle process	1.80	**	+
	GO:0007338	single fertilization	1.79	**	+
	GO:0007314	oocyte anterior/posterior axis specification	1.78	**	+
	GO:0007307	eggshell chorion gene amplification	1.78	*	+
	GO:0007315	pole plasm assembly	1.76	**	+
	GO:0008354	germ cell migration	1.74	*	+
	GO:0007298	border follicle cell migration	1.62	**	—
	GO:1905881	positive regulation of oogenesis	1.60	*	—
	GO:0046008	regulation of female receptivity, post-mating	-1.65	*	—
DGRP 399 Males					
	GO:0048515	spermatid differentiation	2.07	***	+
	GO:0007286	spermatid development	2.06	***	+
	GO:0007291	sperm individualization	1.88	***	—
	GO:0007281	germ cell development	1.88	***	+
	GO:0048232	male gamete generation	1.88	***	+
	GO:0003006	developmental process involved in reproduction	1.87	***	+

	GO:0007283	spermatogenesis	1.84	***	+
	GO:0008354	germ cell migration	1.79	**	+
	GO:0035092	sperm chromatin condensation	1.76	**	—
	GO:2000241	regulation of reproductive process	1.75	***	+
	GO:0007288	sperm axoneme assembly	1.74	**	—
	GO:0007140	male meiotic nuclear division	1.70	**	+
	GO:0030725	germline ring canal formation	1.64	*	—
	GO:0045297	post-mating behavior	1.63	*	—
	GO:0051321	meiotic cell cycle	1.54	**	+
	GO:0140013	meiotic nuclear division	1.41	*	+
DGRP 517					
Females	GO:0007292	female gamete generation	2.26	**	+
	GO:1903046	meiotic cell cycle process	2.23	**	+
	GO:0048477	oogenesis	2.22	**	—
	GO:0003006	developmental process involved in reproduction	2.21	**	+
	GO:0051321	meiotic cell cycle	2.21	**	+
	GO:0007281	germ cell development	2.21	**	+
	GO:0007143	female meiotic nuclear division	2.19	**	+
	GO:0140013	meiotic nuclear division	2.15	**	+
	GO:0030707	ovarian follicle cell development	2.10	**	+
	GO:0009994	oocyte differentiation	2.07	**	+
	GO:0045132	meiotic chromosome segregation	2.05	**	—
	GO:0048599	oocyte development	2.02	**	+
	GO:0007308	oocyte construction	1.99	**	+
	GO:0007309	oocyte axis specification	1.98	**	+
	GO:0008354	germ cell migration	1.98	**	+
	GO:0007277	pole cell development	1.97	**	—
	GO:0046843	dorsal appendage formation	1.94	**	+
	GO:0016321	female meiosis chromosome segregation	1.94	**	—
	GO:0030703	eggshell formation	1.91	**	+
	GO:0009566	fertilization	1.91	**	—
	GO:0007306	eggshell chorion assembly	1.91	**	+
	GO:0033206	meiotic cytokinesis	1.91	**	—
	GO:0007304	chorion-containing eggshell formation	1.91	**	+
	GO:0007314	oocyte anterior/posterior axis specification	1.91	**	+
	GO:0007338	single fertilization	1.88	**	+
	GO:0007315	pole plasm assembly	1.86	**	+
	GO:0008406	gonad development	1.85	**	—
	GO:0045137	development of primary sexual characteristics	1.85	**	—
	GO:2000241	regulation of reproductive process	1.80	**	+
	GO:0007297	ovarian follicle cell migration	1.80	**	—
	GO:0007307	eggshell chorion gene amplification	1.78	**	+
	GO:0061982	meiosis I cell cycle process	1.77	**	+
	GO:0045478	fusome organization	1.74	*	—
	GO:1905879	regulation of oogenesis	1.74	**	+
	GO:0035046	pronuclear migration	1.58	*	—
DGRP 517 Males	GO:0048608	reproductive structure development	2.02	***	+
	GO:0048515	spermatid differentiation	1.87	***	+
	GO:0007286	spermatid development	1.86	***	+
	GO:0048232	male gamete generation	1.70	***	+

GO:0007548	sex differentiation	1.70	**	+
GO:0007140	male meiotic nuclear division	1.68	**	+
GO:0007281	germ cell development	1.66	***	+
GO:0007283	spermatogenesis	1.65	***	+
GO:0051446	positive regulation of meiotic cell cycle	-1.67	*	-

Discussion

A major goal of this study was to understand why galbut virus, despite a high rate of vertical transmission (~100% from both parents), is maintained at a worldwide prevalence of only ~60%. We hypothesized that although galbut virus infection does not produce obvious phenotypic changes, infection might inflict enough of a fitness cost that resistant flies would experience a survival benefit. This would be analogous to an allele with a small negative selection coefficient. To test this hypothesis, we quantified multiple components of fitness in two genetic backgrounds.

Overall, galbut virus infection produced minimal measurable phenotypic effects. In some cases these would be predicted to decrease fitness, such as shortened average lifespan (**Fig. 3**) or decreased survival following fungal infection (**Fig 7E**). In other cases, trait differences such as faster development might increase the relative fitness of galbut virus infected flies (**Fig 5**).

Galbut virus infection minimally decreased lifespan and total offspring output, but the observed trends varied by DGRP strain and mostly did not rise to a statistically significant level (**Figs 3-4**). Our experiments lasted longer than the natural lifespan of *D. melanogaster*, which is estimated to be a week or less in the wild [70]. The total offspring output of galbut virus infected flies and uninfected flies only began to diverge after the parents were > 20 days old, and there was no impact on the number of eggs laid by young females over three days (**Fig 4**). Other examples of partitiviruses altering the reproductive output of their hosts include a partitivirus enhancing fecundity in *Cryptosporidium* [71], a reduction of spores from a partitivirus-infected fungus [72], and partitiviruses infecting *Spodoptera* moths that produced a major decrease in hatchling numbers [72].

Gene ontologies associated with reproduction were positively enriched in galbut virus infected flies, regardless of sex or strain (**Table 2**). It is possible that galbut virus infection manipulates reproductive pathways in a manner that contributes to efficient vertical transmission [4]. The upregulation of genes associated with oogenesis was observed in flies infected with *Drosophila melanogaster* sigma virus, which also depends on vertical transmission [73].

DGRP 517 flies infected by galbut virus pupated and reached adulthood faster than uninfected flies (**Fig 5**). An initial assumption would be that a faster developmental time, in combination with the short life of flies in the wild [70], would confer a fitness benefit. However, flies selected for faster development exhibited fitness trade-offs such as reduced body weight and size, decreased resistance to starvation and desiccation, and an overall lower egg output [74]. This highlights the difficulty of extrapolating total fitness from singly-measured traits [75].

For the most part, galbut virus infected and uninfected flies survived similarly following infection by microbial pathogens (**Fig 6**). Galbut virus infected DGRP 399 females exhibited decreased survival following ingestion, but not injection, of *Pseudomonas aeruginosa*. This difference may not be surprising as the gut epithelial immune response has key differences compared to responses to systemic infection [76]. DGRP 399 flies of both sexes exhibited increased sensitivity to the fungal pathogen *Candida albicans*. In *Drosophila*, the common microbiome constituent *Lactobacillus planatarum* decreased mortality of a fungal pathogen (*Diaporthe* sp.) by mitigating fungal toxicity and altered fly behaviour to reduce infection risk [77]. No significant changes in the DNA levels of *L. planatarum* or other major microbiome constituents was observed in galbut virus infected flies (**Fig 2**).

It is difficult to assess the net impact of these separately measured traits. Laboratory assays imperfectly recapitulate natural environments and these experiments provide a limited window into the influence of galbut virus in the wild. For most measured traits, differences associated with galbut virus infection were smaller than those attributable to different DGRP strain and sex (**Figs 3-4, Fig 7**). Nevertheless, selection can act on small differences in relative fitness, and it is possible that in aggregate galbut virus infection reduces fitness. Galbut virus is highly prevalent, exhibits a broad tissue distribution, and exists as a lifelong infection, so small phenotypic changes should not necessarily be interpreted as insignificant ones. Additional laboratory and field-based studies that track galbut virus-*Drosophila* dynamics will shed further light on the extent to which this virus and similar persistent viruses shape the evolution of their hosts in cryptic but possibly important ways.

Acknowledgements: We thank Marylee Kapuscinski and the Colorado State University Next Generation Sequencing Core Facility for assistance with sequencing and Dr. Susan Tsunoda for helpful discussion. We would like to acknowledge Drs. Raul Andino and Michel Tassetto for providing *Drosophila* C virus stocks and Dr. Brad Borlee for providing *Pseudomonas aeruginosa* and *Staphylococcus aureus* stocks.

Funding: This work was supported by Animal Health and Disease Grant No. 19HMFXXXXG039150001 / Project Accession No. 1024856 from the USDA National Institute of Food and Agriculture, allocated via the Colorado State University College of Veterinary Medicine and Biomedical Sciences College Research Council. Computational resources were supported by NIH/NCATS Colorado CTSA Grant Number UL1 TR002535. STC was supported in part by National Science Foundation (NSF) NRT grant 1450032. TJD was supported in part by the American Society for Microbiology Undergraduate Research Fellowship. Stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study. Any opinions, findings, conclusions or recommendations expressed in this paper are those of the author(s) and do not necessarily reflect the views of the funding organizations.

SUPPLEMENTAL MATERIALS:

Supplemental Fig 1. Galbut virus infection does not influence adult offspring sex ratio. Offspring collected from groups of galbut virus infected or uninfected parents from DGRP 399 and 517 strains every 14 days (see Fig 4). Offspring sex ratios from each time point were calculated by dividing total male offspring by total female offspring. No statistical significance was measured in either strain (t-test).

Supplemental Fig 2. Dot plot of differentially regulated gene ontology (GO) pathways in infected DGRP 399 female flies. A dot plot representation of the top differentially regulated GO pathways in galbut virus-infected DGRP 399 female flies as determined by gene set enrichment analysis (GSEA) using the R package “clusterProfiler”. Top 10 differentially regulated pathways are plotted in each GO category (biological function, BF; cellular component, CC; molecular function, MF). Differentially regulated pathways for these flies were either upregulated (activated) or downregulated (suppressed). Size of dots corresponds with number of differentially regulated genes (DEG; counts) identified in each

specified GO pathway. Percentage of DEGs in a given GO pathway (number of DEGs divided by total number of genes listed under the specified GO pathway) is plotted as gene ratio.

Supplemental Fig 3. Dot plot of differentially regulated gene ontology (GO) pathways in infected DGRP 399 male flies. A dot plot representation of the top differentially regulated GO pathways in galbut virus-infected DGRP 399 male flies as determined by gene set enrichment analysis (GSEA) using the R package “clusterProfiler”. Top 10 differentially regulated pathways are plotted in each GO category

(biological function, BP; cellular component, CC; molecular function, MF). All top differentially regulated pathways for these flies were upregulated (activated). Size of dots corresponds with number of differentially regulated genes (DEG; counts) identified in each specified GO pathway. Percentage of DEGs in a given GO pathway (number of DEGs divided by total number of genes listed the specified GO pathway) is plotted as gene ratio.

Supplemental Fig 4. Dot plot of differentially regulated gene ontology (GO) pathways in infected DGRP 517 female flies. A dot plot representation of the top differentially regulated GO pathways in galbut virus infected DGRP 517 female flies as determined by gene set enrichment analysis (GSEA) using the R package “clusterProfiler”. Top 10 differentially regulated pathways are plotted in each GO category (biological function, BF; cellular component, CC; molecular function, MF). All top differentially regulated pathways for these flies were upregulated (activated). Size of dots corresponds with number of differentially regulated genes (DEG; counts) identified in each specified GO pathway. Percentage of DEGs in a given GO pathway (number of DEGs divided by total number of genes listed under the specified GO pathway) is plotted as gene ratio.

Supplemental Fig 5. Dot plot of differentially regulated gene ontology (GO) pathways in infected DGRP 517 male flies. A dot plot representation of the top differentially regulated GO pathways in galbut virus-infected DGRP 517 male flies as determined by gene set enrichment analysis (GSEA) using the R package “clusterProfiler”. Top 10 differentially regulated pathways are plotted in each GO category (biological function, BF; cellular component, CC; molecular function, MF). All top differentially regulated pathways for these flies were upregulated (activated). Size of dots corresponds with number of differentially regulated genes (DEG; counts) identified in each specified GO pathway. Percentage of

DEGs in a given GO pathway (number of DEGs divided by total number of genes listed under the specified GO pathway) is plotted as gene ratio.

Supplemental Table 1: Primers used for quantifying levels of galbut virus and microbiome constituents.

Target	Forward (5'-3')	Reverse (5'-3')	Original Citation
Galbut virus	CCGTGAAGCAAGGA ATCAAT	TGCCGATTTTCTGCTCT TTT	Cross et al: https://doi.org/10.1128/JVI.01070-20
RpL32	TGCTAAGCTGTCGCA CAAATGG	TGCGCTTGTTTCGATCC GTAAC	Cross et al: https://doi.org/10.1128/JVI.01070-20
<i>Acetobacter</i> spp.	TAGCTAACGCGATAA GCACA	ACAGCCTACCCATACA AGCC	AM Early et al: https://doi.org/10.1371/journal.pone.0170332
<i>Lactobacillus brevis</i>	TCAGTTTTGAGGGGC TTACCTCTCT	GGCATCCACCATGCGC CCTT	AM Early et al: https://doi.org/10.1371/journal.pone.0170332
<i>Lactobacillus planatarum</i>	TGCGGCTGGATCACC TCCTTTC	ACTGGTTCGGTTCCAA TGGGCC	AM Early et al: https://doi.org/10.1371/journal.pone.0170332
<i>Corynebacterium</i> spp.	AAACGGGTACGCATC ACG	GGGTTGATATTCCCGT ACCC	This study
<i>Saccharomyces cerevisiae</i>	AGGAGTGCGGTTCTT TG	TACTTACCGAGGCAAG CTACA	H Chang et al: https://doi.org/10.1016/j.mimet.2007.08.013
<i>deformed (dfd)</i>	GTAGCGAAGAAACC CACCAA	ACGCTCCACTCACCTC ATTC	AM Early et al: https://doi.org/10.1371/journal.pone.0170332

References

1. Webster CL, Waldron FM, Robertson S, Crowson D, Ferrari G, Quintana JF, et al. The Discovery, Distribution, and Evolution of Viruses Associated with *Drosophila melanogaster*. *PLoS Biol*. 2015;13: e1002210–e1002210. doi:10.1371/journal.pbio.1002210
2. Webster CL, Longdon B, Lewis SH, Obbard DJ. Twenty-Five New Viruses Associated with the Drosophilidae (Diptera). *Evol Bioinform Online*. 2016;12: 13–25. doi:10.4137/EBO.S39454
3. Shi M, White VL, Schlub T, Eden J-S, Hoffmann AA, Holmes EC. No detectable effect of Wolbachia wMel on the prevalence and abundance of the RNA virome of *Drosophila melanogaster*. *Proc R Soc B*. 2018;285: 20181165. doi:10.1098/rspb.2018.1165
4. Cross ST, Maertens BL, Dunham TJ, Rodgers CP, Brehm AL, Miller MR, et al. Partitiviruses infecting *Drosophila melanogaster* and *Aedes aegypti* exhibit efficient biparental vertical transmission. *J Virol*. 2020;94: e01070-20. doi:10.1128/JVI.01070-20
5. Fine PEM. Vectors and Vertical Transmission: An Epidemiologic Perspective. *Annals of the New York Academy of Sciences*. 1975;266: 173–194. doi:10.1111/j.1749-6632.1975.tb35099.x
6. Nibert ML, Ghabrial SA, Maiss E, Lesker T, Vainio EJ, Jiang D, et al. Taxonomic reorganization of family Partitiviridae and other recent progress in partitivirus research. *Virus Research*. 2014;188: 128–141. doi:10.1016/j.virusres.2014.04.007
7. Vainio EJ, Chiba S, Ghabrial SA, Maiss E, Roossinck M, Sabanadzovic S, et al. ICTV Virus Taxonomy Profile: Partitiviridae. *Journal of General Virology*,. 2018;99: 17–18. doi:10.1099/jgv.0.000985
8. Ghabrial SA, Castón JR, Jiang D, Nibert ML, Suzuki N. 50-plus years of fungal viruses. *Virology*. 2015;479–480: 356–368. doi:10.1016/j.virol.2015.02.034
9. Roossinck MJ. Lifestyles of plant viruses. *Philos Trans R Soc Lond, B, Biol Sci*. 2010;365: 1899–1905. doi:10.1098/rstb.2010.0057
10. Boccardo G, Lisa V, Luisoni E, Milne RG. Cryptic plant viruses. *Adv Virus Res*. 1987;32: 171–214. doi:10.1016/s0065-3527(08)60477-7
11. Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The *Drosophila melanogaster* Genetic Reference Panel. *Nature*. 2012;482: 173–178. doi:10.1038/nature10811
12. Shi M, Lin X-D, Tian J-H, Chen L-J, Chen X, Li C-X, et al. Redefining the invertebrate RNA virosphere. *Nature*. 2016;540: 539–543. doi:10.1038/nature20167
13. Shi C, Beller L, Deboutte W, Yinda KC, Delang L, Vega-Rúa A, et al. Stable distinct core eukaryotic viromes in different mosquito species from Guadeloupe, using single mosquito viral metagenomics. *Microbiome*. 2019;7: 121. doi:10.1186/s40168-019-0734-2
14. Zakrzewski M, Rašić G, Darbro J, Krause L, Poo YS, Filipović I, et al. Mapping the virome in wild-caught *Aedes aegypti* from Cairns and Bangkok. *Scientific Reports*. 2018;8: 4690. doi:10.1038/s41598-018-22945-y
15. Cogni R, Cao C, Day JP, Bridson C, Jiggins FM. The genetic architecture of resistance to virus infection in *Drosophila*. *Mol Ecol*. 2016/08/26 ed. 2016;25: 5228–5241. doi:10.1111/mec.13769
16. Palmer WH, Varghese FS, van Rij RP. Natural Variation in Resistance to Virus Infection in Dipteran Insects. *Viruses*. 2018;10. doi:10.3390/v10030118
17. Chapman JR, Dowell MA, Chan R, Unckless RL. The Genetic Basis of Natural Variation in *Drosophila melanogaster* Immune Defense against *Enterococcus faecalis*. *Genes (Basel)*. 2020;11:

234. doi:10.3390/genes11020234
18. Klein SL, Flanagan KL. Sex differences in immune responses. *Nature Reviews Immunology*. 2016;16: 626–638. doi:10.1038/nri.2016.90
19. Kopp A, Duncan I, Carroll SB. Genetic control and evolution of sexually dimorphic characters in *Drosophila*. *Nature*. 2000;408: 553–559. doi:10.1038/35046017
20. Belmonte RL, Corbally M-K, Duneau DF, Regan JC. Sexual Dimorphisms in Innate Immunity and Responses to Infection in *Drosophila melanogaster*. *Front Immunol*. 2020;10: 3075–3075. doi:10.3389/fimmu.2019.03075
21. Medd NC, Fellous S, Waldron FM, Xuéreb A, Nakai M, Cross JV, et al. The virome of *Drosophila suzukii*, an invasive pest of soft fruit. *Virus Evol*. 2018;4: vey009–vey009. doi:10.1093/ve/vey009
22. Faizah AN, Kobayashi D, Isawa H, Amoa-Bosompem M, Murota K, Higa Y, et al. Deciphering the Virome of *Culex vishnui* Subgroup Mosquitoes, the Major Vectors of Japanese Encephalitis, in Japan. *Viruses*. 2020;12: 264. doi:10.3390/v12030264
23. Pettersson JH-O, Shi M, Eden J-S, Holmes EC, Hesson JC. Meta-Transcriptomic Comparison of the RNA Viromes of the Mosquito Vectors *Culex pipiens* and *Culex torrentium* in Northern Europe. *Viruses*. 2019;11: 1033. doi:10.3390/v11111033
24. Lara Pinto AZ de, Santos de Carvalho M, de Melo FL, Ribeiro ALM, Morais Ribeiro B, Dezengrini Shhessarenko R. Novel viruses in salivary glands of mosquitoes from sylvatic Cerrado, Midwestern Brazil. *PLoS One*. 2017;12: e0187429–e0187429. doi:10.1371/journal.pone.0187429
25. Sadeghi M, Altan E, Deng X, Barker CM, Fang Y, Coffey LL, et al. Virome of > 12 thousand *Culex* mosquitoes from throughout California. *Virology*. 2018;523: 74–88. doi:10.1016/j.virol.2018.07.029
26. Fauver JR, Grubaugh ND, Krajacich BJ, Weger-Lucarelli J, Lakin SM, Fakoli LS, et al. West African *Anopheles gambiae* mosquitoes harbor a taxonomically diverse virome including new insect-specific flaviviruses, mononegaviruses, and totiviruses. *Virology*. 2016;498: 288–299. doi:10.1016/j.virol.2016.07.031
27. Xu P, Yang L, Yang X, Li T, Graham RI, Wu K, et al. Novel partiti-like viruses are conditional mutualistic symbionts in their normal lepidopteran host, African armyworm, but parasitic in a novel host, Fall armyworm. *PLOS Pathogens*. 2020;16: e1008467. doi:10.1371/journal.ppat.1008467
28. Li C-X, Shi M, Tian J-H, Lin X-D, Kang Y-J, Chen L-J, et al. Unprecedented genomic diversity of RNA viruses in arthropods reveals the ancestry of negative-sense RNA viruses. *eLife*. 2015;4: e05378. doi:10.7554/eLife.05378
29. Shi M, Lin X-D, Chen X, Tian J-H, Chen L-J, Li K, et al. The evolutionary history of vertebrate RNA viruses. *Nature*. 2018;556: 197–202. doi:10.1038/s41586-018-0012-7
30. Edgar RC, Taylor J, Lin V, Altman T, Barbera P, Meleshko D, et al. Petabase-scale sequence alignment catalyses viral discovery. *bioRxiv*. 2021; 2020.08.07.241729. doi:10.1101/2020.08.07.241729
31. Canuti M, van der Hoek L. Virus discovery: are we scientists or genome collectors? *Trends Microbiol*. 2014;22: 229–231. doi:10.1016/j.tim.2014.02.004
32. Greninger AL. A decade of RNA virus metagenomics is (not) enough. *Virus Research*. 2018;244: 218–229. doi:10.1016/j.virusres.2017.10.014
33. Huang W, Massouras A, Inoue Y, Peiffer J, Ràmia M, Tarone AM, et al. Natural variation in genome architecture among 205 *Drosophila melanogaster* Genetic Reference Panel lines. *Genome*

- Res. 2014/04/08 ed. 2014;24: 1193–1208. doi:10.1101/gr.171546.113
34. Early AM, Shanmugarajah N, Buchon N, Clark AG. *Drosophila* Genotype Influences Commensal Bacterial Levels. PLOS ONE. 2017;12: e0170332. doi:10.1371/journal.pone.0170332
35. Gould AL, Zhang V, Lamberti L, Jones EW, Obadia B, Korasidis N, et al. Microbiome interactions shape host fitness. Proc Natl Acad Sci U S A. 2018/12/03 ed. 2018;115: E11951–E11960. doi:10.1073/pnas.1809349115
36. Therneau T. A Package for Survival Analysis in R. 2020. Available: <https://CRAN.R-project.org/package=survival>
37. Vieira B. Colony Counter. 2009. Available: <https://imagej.nih.gov/ij/plugins/colony-counter.html>
38. Olcott MH, Henkels MD, Rosen KL, L Walker F, Sneh B, Loper JE, et al. Lethality and Developmental Delay in *Drosophila melanogaster* Larvae after Ingestion of Selected *Pseudomonas fluorescens* Strains. PLOS ONE. 2010;5: e12504. doi:10.1371/journal.pone.0012504
39. Lutter EI, Faria MMP, Rabin HR, Storey DG. *Pseudomonas aeruginosa* cystic fibrosis isolates from individual patients demonstrate a range of levels of lethality in two *Drosophila melanogaster* infection models. Infect Immun. 2008/02/19 ed. 2008;76: 1877–1888. doi:10.1128/IAI.01165-07
40. Apidianakis Y, Rahme LG. *Drosophila melanogaster* as a model host for studying *Pseudomonas aeruginosa* infection. Nature Protocols. 2009;4: 1285–1294. doi:10.1038/nprot.2009.124
41. Duneau DF, Kondolf HC, Im JH, Ortiz GA, Chow C, Fox MA, et al. The Toll pathway underlies host sexual dimorphism in resistance to both Gram-negative and Gram-positive bacteria in mated *Drosophila*. BMC Biology. 2017;15: 124. doi:10.1186/s12915-017-0466-3
42. van Rij RP, Saleh M-C, Berry B, Foo C, Houk A, Antoniewski C, et al. The RNA silencing endonuclease Argonaute 2 mediates specific antiviral immunity in *Drosophila melanogaster*. Genes & Development. 2006;20: 2985–2995. doi:10.1101/gad.1482006
43. Davis MM, Alvarez FJ, Ryman K, Holm ÅA, Ljungdahl PO, Engström Y. Wild-Type *Drosophila melanogaster* as a Model Host to Analyze Nitrogen Source Dependent Virulence of *Candida albicans*. PLOS ONE. 2011;6: e27434. doi:10.1371/journal.pone.0027434
44. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal; Vol 17, No 1: Next Generation Sequencing Data Analysis. 2011. doi:10.14806/ej.17.1.200
45. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. Nature Methods. 2015;12: 357–360. doi:10.1038/nmeth.3317
46. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2013;30: 923–930. doi:10.1093/bioinformatics/btt656
47. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550–550. doi:10.1186/s13059-014-0550-8
48. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2020. Available: <https://www.R-project.org/>
49. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. OMICS: A Journal of Integrative Biology. 2012;16: 284–287. doi:10.1089/omi.2011.0118
50. Li N, Ma W-T, Pang M, Fan Q-L, Hua J-L. The Commensal Microbiota and Viral Infection: A Comprehensive Review. Front Immunol. 2019;10: 1551–1551. doi:10.3389/fimmu.2019.01551

51. Travers LM, Garcia-Gonzalez F, Simmons LW. Live fast die young life history in females: evolutionary trade-off between early life mating and lifespan in female *Drosophila melanogaster*. *Sci Rep*. 2015;5: 15469–15469. doi:10.1038/srep15469
52. Hedges LM, Brownlie JC, O'Neill SL, Johnson KN. Wolbachia and virus protection in insects. *Science*. 2008;322: 702. doi:10.1126/science.1162418
53. Trinder M, Daisley BA, Dube JS, Reid G. *Drosophila melanogaster* as a High-Throughput Model for Host-Microbiota Interactions. *Front Microbiol*. 2017;8: 751. doi:10.3389/fmicb.2017.00751
54. Sibley CD, Duan K, Fischer C, Parkins MD, Storey DG, Rabin HR, et al. Discerning the complexity of community interactions using a *Drosophila* model of polymicrobial infections. *PLoS Pathog*. 2008;4: e1000184. doi:10.1371/journal.ppat.1000184
55. Sansone CL, Cohen J, Yasunaga A, Xu J, Osborn G, Subramanian H, et al. Microbiota-Dependent Priming of Antiviral Intestinal Immunity in *Drosophila*. *Cell Host Microbe*. 2015;18: 571–581. doi:10.1016/j.chom.2015.10.010
56. Xu J, Cherry S. Viruses and antiviral immunity in *Drosophila*. *Dev Comp Immunol*. 2013/05/13 ed. 2014;42: 67–84. doi:10.1016/j.dci.2013.05.002
57. Martinez J, Bruner-Montero G, Arunkumar R, Smith SCL, Day JP, Longdon B, et al. Virus evolution in Wolbachia-infected *Drosophila*. *Proc Biol Sci*. 2019/10/30 ed. 2019;286: 20192117. doi:10.1098/rspb.2019.2117
58. Teixeira L, Ferreira A, Ashburner M. The bacterial symbiont Wolbachia induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol*. 2008;6: e2–e2. doi:10.1371/journal.pbio.1000002
59. Martinez J, Longdon B, Bauer S, Chan Y-S, Miller WJ, Bourtzis K, et al. Symbionts commonly provide broad spectrum resistance to viruses in insects: a comparative analysis of Wolbachia strains. *PLoS Pathog*. 2014;10: e1004369–e1004369. doi:10.1371/journal.ppat.1004369
60. Osborne SE, Leong YS, O'Neill SL, Johnson KN. Variation in antiviral protection mediated by different Wolbachia strains in *Drosophila simulans*. *PLoS Pathog*. 2009/11/13 ed. 2009;5: e1000656–e1000656. doi:10.1371/journal.ppat.1000656
61. Ferrandon D, Imler J-L, Hetru C, Hoffmann JA. The *Drosophila* systemic immune response: sensing and signalling during bacterial and fungal infections. *Nature Reviews Immunology*. 2007;7: 862–874. doi:10.1038/nri2194
62. Pecasse F, Beck Y, Ruiz C, Richards G. Krüppel-homolog, a Stage-Specific Modulator of the Prepupal Ecdysone Response, Is Essential for *Drosophila* Metamorphosis. *Developmental Biology*. 2000;221: 53–67. doi:10.1006/dbio.2000.9687
63. Fichelson P, Brigui A, Pichaud F. Orthodenticle and Kruppel homolog 1 regulate *Drosophila* photoreceptor maturation. *Proc Natl Acad Sci U S A*. 2012/04/30 ed. 2012;109: 7893–7898. doi:10.1073/pnas.1120276109
64. Shi L, Lin S, Grinberg Y, Beck Y, Grozinger CM, Robinson GE, et al. Roles of *Drosophila* Kruppel-homolog 1 in neuronal morphogenesis. *Developmental Neurobiology*. 2007;67: 1614–1626. doi:10.1002/dneu.20537
65. Montana ES, Littleton JT. Expression Profiling of a Hypercontraction-induced Myopathy in *Drosophila* Suggests a Compensatory Cytoskeletal Remodeling Response. *Journal of Biological Chemistry*. 2006;281: 8100–8109. doi:10.1074/jbc.M512468200
66. Kucherenko MM, Marrone AK, Rishko VM, Magliarelli H de F, Shcherbata HR. Stress and muscular dystrophy: A genetic screen for Dystroglycan and Dystrophin interactors in *Drosophila*

- identifies cellular stress response components. *Developmental Biology*. 2011;352: 228–242.
doi:10.1016/j.ydbio.2011.01.013
67. Lammel U, Bechtold M, Risse B, Berh D, Fleige A, Bunse I, et al. The *Drosophila* FHOD1-like formin Knittrig acts through Rok to promote stress fiber formation and directed macrophage migration during the cellular immune response. *Development*. 2014;141: 1366.
doi:10.1242/dev.101352
68. Gaudet P, Livstone MS, Lewis SE, Thomas PD. Phylogenetic-based propagation of functional annotations within the Gene Ontology consortium. *Brief Bioinform*. 2011/08/27 ed. 2011;12: 449–462. doi:10.1093/bib/bbr042
69. Merkling SH, Overheul GJ, van Mierlo JT, Arends D, Gilissen C, van Rij RP. The heat shock response restricts virus infection in *Drosophila*. *Sci Rep*. 2015;5: 12758–12758.
doi:10.1038/srep12758
70. Rosewell J, Shorrocks B. The implication of survival rates in natural populations of *Drosophila*: capture-recapture experiments on domestic species. *Biological Journal of the Linnean Society*. 2008;32: 373–384. doi:10.1111/j.1095-8312.1987.tb00438.x
71. Jenkins MC, Higgins J, Abrahante JE, Kniel KE, O’Brien C, Trout J, et al. Fecundity of *Cryptosporidium parvum* is correlated with intracellular levels of the viral symbiont CPV. *International Journal for Parasitology*. 2008;38: 1051–1055. doi:10.1016/j.ijpara.2007.11.005
72. Zhong J, Chen D, Lei XH, Zhu HJ, Zhu JZ, Da Gao B. Detection and characterization of a novel Gammapartitivirus in the phytopathogenic fungus *Colletotrichum acutatum* strain HNZJ001. *Virus Research*. 2014;190: 104–109. doi:10.1016/j.virusres.2014.05.028
73. Carpenter J, Hutter S, Baines JF, Roller J, Saminadin-Peter SS, Parsch J, et al. The Transcriptional Response of *Drosophila melanogaster* to Infection with the Sigma Virus (Rhabdoviridae). *PLOS ONE*. 2009;4: e6838. doi:10.1371/journal.pone.0006838
74. Yadav P, Sharma VK. Correlated changes in life history traits in response to selection for faster pre-adult development in the fruit fly *Drosophila melanogaster*. *J Exp Biol*. 2014;217: 580.
doi:10.1242/jeb.093864
75. Orr HA. Fitness and its role in evolutionary genetics. *Nat Rev Genet*. 2009;10: 531–539.
doi:10.1038/nrg2603
76. Buchon N, Silverman N, Cherry S. Immunity in *Drosophila melanogaster*--from microbial recognition to whole-organism physiology. *Nat Rev Immunol*. 2014;14: 796–810.
doi:10.1038/nri3763
77. Su W, Liu J, Bai P, Ma B, Liu W. Pathogenic fungi-induced susceptibility is mitigated by mutual *Lactobacillus plantarum* in the *Drosophila melanogaster* model. *BMC Microbiology*. 2019;19: 302.
doi:10.1186/s12866-019-1686-1