Phycobilisome breakdown effector NblD is required to maintain the cellular amino acid composition during nitrogen starvation

Vanessa Krauspe,^a Stefan Timm,^b Martin Hagemann,^b Wolfgang R. Hess,^{a,#}

^aGenetics and Experimental Bioinformatics, Institute of Biology 3, Faculty of Biology, University of Freiburg, 79104 Freiburg, Germany

^bPlant Physiology Department, Institute of Biosciences, University of Rostock, Rostock, 18059 Rostock, Germany;

Running Head: NblD affects metabolite pools

[#]Correspondence: Wolfgang R. Hess, wolfgang.hess@biologie.uni-freiburg.de

Keywords: amino acids, cyanobacteria, nitrogen starvation, photosynthesis, phycobilisomes, *Synechocystis* sp. PCC 6803, small proteins, stress response

Abstract

Small proteins are critically involved in the acclimation response of photosynthetic cyanobacteria to nitrogen starvation. NblD is the 66-amino-acid effector of nitrogen-limitation-induced phycobilisome breakdown, which is believed to replenish the cellular amino acid pools. To address the physiological functions of NblD, the concentrations of amino acids, intermediates of the arginine catabolism pathway and several organic acids were measured during the response to nitrogen starvation in the cyanobacterium *Synechocystis* sp. PCC 6803 wild type and in an *nblD* deletion strain. A characteristic signature of metabolite pool composition was identified, which shows that NblD-mediated phycobilisome degradation is required to maintain the cellular amino acid and organic acid pools during nitrogen starvation. Specific deviations from the wild type suggest wider-reaching effects that also affect such processes as redox homeostasis via glutathione and tetrapyrrole biosynthesis, both of which are linked to the strongly decreased glutamate pool, and transcriptional reprogramming via an enhanced concentration of 2-oxoglutarate, the metabolite co-regulator of the NtcA transcription factor. The essential role played by NblD in metabolic homeostasis is consistent with the widespread occurrence of NblD throughout the cyanobacterial radiation and the previously observed strong positive selection for the *nblD* gene under fluctuating nitrogen supply.

Importance

Cyanobacteria play important roles in the global carbon and nitrogen cycles. In their natural environment, these organisms are exposed to fluctuating nutrient conditions. Nitrogen starvation induces a coordinated nitrogen-saving program that includes the

45 breakdown of nitrogen-rich photosynthetic pigments, particularly phycobiliproteins.
 46 The small protein NbID was recently identified as an effector of phycobilisome
 47 breakdown in cyanobacteria. In this study, we demonstrate that the NbID-mediated
 48 degradation of phycobiliproteins is needed to sustain cellular pools of soluble amino
 49 acids and other crucial metabolites. The essential role played by NbID in metabolic
 50 homeostasis explains why genes encoding this small protein are conserved in
 51 almost all members of cyanobacterial radiation.

52

53 Introduction

54 The recent introduction of the antibiotic retapamulin as an inhibitor of the initiation of
 55 translation into ribosome-profiling studies in bacteria has revolutionized the view on
 56 the bacterial small proteome suggesting that a high number of previously unexplored
 57 small open reading frames (smORFs) exists in bacteria (1, 2). Small proteins, often
 58 defined as those containing 50 or fewer amino acids are too small to have own
 59 enzymatic activities and this applies also to the group of “larger” small proteins that
 60 range from 50 to 100 amino acids. Nevertheless, small proteins have been shown to
 61 fulfil multiple important functions, for example as regulatory subunits of multi-protein
 62 complexes, as modulators of enzymatic activities of larger proteins or as toxins (for a
 63 recent review see Hemm et al. (3)).

64 A large number of small proteins have been characterized in photosynthetic
 65 cyanobacteria due to the extensive work on characterizing the photosynthetic
 66 apparatus. Nineteen small proteins with fewer than 50 amino acids are directly
 67 involved in photosynthesis. These proteins constitute subunits of photosystem I
 68 (encoded by genes *psaM*, *psaJ* and *psaI* (4)), of photosystem II (genes *psbM*, *psbT*
 69 (*ycf8*), *psbI*, *psbL*, *psbJ*, *psbY*, *psbX*, *psb30* (*ycf12*), *psbN*, *psbF*, *psbK* (5, 6)),
 70 participate in photosynthetic electron transport (Cyt_b₆*f* complex; *petL*, *petN*, *petM*,
 71 *petG* (7–9)) or in photosynthetic complex I (*ndhP*, *ndhQ* (10, 11)). Other previously
 72 characterized small proteins in cyanobacteria have accessory functions to
 73 photosynthesis such as the small CAB like proteins HliA, HliB, HliC and HliD (12–
 74 14). The cytochrome *b*₆*f* complex subunit VIII, encoded by *petN* is with 29 amino
 75 acids the shortest annotated photosynthetic protein conserved in cyanobacteria (15).
 76 Additional smORFs were computationally predicted in the model cyanobacterium
 77 *Synechocystis* sp. PCC 6803 (hereafter designated *Synechocystis* 6803) (16). The

44 amino acids small protein AcnSP was characterized as regulator of the tricarboxylic acid (TCA) cycle enzyme aconitase (17).

Another set of small proteins were found to be involved in the acclimation response of the photosynthetic machinery to nitrogen (N) starvation, including NbIA1, NbIA2 and NbID, proteins of 62, 60 and 66 residues in *Synechocystis* 6803 (18, 19). Nondiazotrophic cyanobacteria, such as *Synechocystis* 6803, depend on combined nitrogen (N) sources in their environment. These bacteria favor ammonia (NH_3 or NH_4^+) but can also utilize nitrate (NO_3^-) as an inorganic N source (20, 21). Under N-limited conditions, cells can start an acclimation program, manifested as chlorosis, and finally enter a dormant state (22–26). During chlorosis, N-rich pigments are degraded, predominantly the giant light-harvesting pigment-protein complexes, the phycobilisomes (22, 23, 27). The excess carbon (C) availability during chlorosis induces the accumulation of C reserves, particularly glycogen (28). The tricarboxylic acid (TCA) cycle intermediate 2-oxoglutarate (2-OG), which is primarily used as a C skeleton for ammonium incorporation by the glutamine synthetase/glutamine 2-OG aminotransferase (GS/GOGAT) cycle (29, 30), has been shown to serve as a cellular signal metabolite to report the relative N/C status (31–35). If ammonia assimilation via the GS/GOGAT cycle is slowed due to N deficiency, 2-OG is no longer converted and accumulates in the cyanobacterial cell. As a result, 2-OG can bind to the signal transduction protein PII (36, 37), thereby releasing PipX from the PipX/PII complex (33, 35). Free PipX can subsequently associate with and activate the global nitrogen transcription regulator NtcA together with 2-OG. In *Synechocystis* 6803, NtcA directly controls 79 different genes after 4 h of N starvation (38). Depending on the location of the respective binding motifs, NtcA can either activate genes, especially those involved in nitrogen uptake and assimilation, or repress genes, as observed for the

GS inhibition factors *gifA* and *gifB* (39). Two of the NtcA-activated genes in *Synechocystis* 6803 are *nbIA1* and *nbIA2* (*ssl0452* and *ssl0453*) (38), which encode small proteins that are critically important in N starvation-induced phycobilisome breakdown (18, 40–42).

Phycobilisome breakdown in N-starved cyanobacteria is a tightly controlled process that is believed to fill cellular amino acid pools, compensating for the limited *de novo* nitrogen assimilation via the GS-GOGAT cycle. NbIA1/A2 play essential roles in the coordinated process of phycobilisome degradation during N deprivation because corresponding mutants show a so-called non-bleaching phenotype (*nbl*), i.e., they retain their phycobilisomes under N-limiting conditions (18, 40). NbIA proteins function as adaptors for the Clp protease by connecting the phycobiliproteins and the ClpC chaperone (41–44). Phycobilisome degradation can be generally divided into a rapid early phase, in which almost 50% of phycocyanin is degraded within 5 h, followed by a more sedate phase until only approximately 10% of the initial phycocyanin content is attained after 24 h of N deficiency (45). The degradation of the phycobilisome starts at the distal phycocyanin rods and extends to the allophycocyanin core, which is in proximity to the photosystems situated in the thylakoid membrane (44, 46). Previous studies showed that the inability to degrade phycobilisomes in the $\Delta nbIA1/A2$ mutant affected the composition of amino acid pools during N starvation in a specific way in *Synechocystis* 6803 (47). These authors demonstrated that the pool sizes of 12 amino acids were NbIA1/A2-dependent, while others were not.

NbID has been recently identified as another small protein determined to be important for phycobilisome turnover because the pigmentation of $\Delta nbID$ mutants was only marginally reduced under N starvation (19) resembling the non-bleaching

phenotype of $\Delta nbIA$ mutants. NblD specifically binds to the phycocyanin beta subunit (CpcB) of phycobilisomes at an early time point of nitrogen depletion (19). However, the precise biochemical function of NblD remains to be determined. In this study, we compared the metabolic composition of *Synechocystis* 6803 wild-type cells (WT) with a mutant deficient for NblD ($\Delta nbID$) over a time course of N starvation. A characteristic signature in the composition of these metabolite pools was observed that supports the essential role played by NblD in the maintenance of cellular amino acid homeostasis during N starvation. Other deviations from the WT suggest wider-reaching effects, also affecting transcriptional reprogramming via an unusually high concentration of 2-OG, the metabolite co-regulator of the NtcA transcription factor.

Results

Global responses of an *nbID* deletion mutant to nitrogen starvation.

In the present study, the amino acid pools and amounts of other important metabolites were compared in the *Synechocystis* 6803 wild type (WT) to an *nbID* deletion mutant ($\Delta nbID$). In the absence of a nitrogen source, cells of the $\Delta nbID$ strain are bleaching significantly less than the WT (19). Accordingly, we hypothesized that the soluble amino acid pools of $\Delta nbID$ cells may be exhausted more quickly than those of WT cells. This assumption was tested by metabolomics based on liquid chromatography mass spectrometry (LC-MS/MS) of three biologically independent samples for each WT and $\Delta nbID$, taken before (+N condition), and 3 h or 24 h after the transfer into N-free BG11 medium. In total, 28 compounds, including almost all amino acids (Asn and Cys were below the detection limit), intermediates of the

arginine catabolism pathway, and several organic acids, were quantified (see Table S1 for the LC-MS/MS raw data).

Cells obtained from the +N condition and 3 h after nitrogen removal showed no difference in pigmentation. After 24 h of N starvation, the WT samples showed pronounced bleaching, while the $\Delta nbID$ mutant bleached much less (Fig. 1A and B).

Comparing the spectra with previously published spectra suggests that the decrease in the 630 nm absorption maximum of phycobilisomes in $\Delta nbID$ under N starvation is greater than in the $\Delta nbIA1/nbIA2$ double mutant (18, 42), although clearly not as much as in the WT under the same condition. The degradation of phycobilisomes contributed to transiently higher levels of total soluble amino acids in *Synechocystis* 6803 WT during N starvation (Fig. 2A, Table S2). The level of free amino acids increased from approximately 261 ng OD₇₅₀⁻¹ ml⁻¹ ($\pm$ 32.4) under nitrogen-replete conditions (+N) by 65% on average after 3 h and approximately 27% (in relation to the starting value) after 24 h of N limitation (-N). In contrast, the $\Delta nbID$ mutant contained 335 ng OD₇₅₀⁻¹ ml⁻¹ ($\pm$ 43.3) of free amino acids in the +N condition and remained at a similar level after 3 h. After 24 h after nitrogen removal, the amino acid content decreased dramatically by 70% to 75% in relation to the initial content in $\Delta nbID$ (Fig. 2A). Compared to the amino acid pool, the total amounts of other quantified metabolites, primarily organic acids, did not differ significantly between cells of the WT and $\Delta nbID$ (Fig. 2B), at least in the +N condition and after 24 h. The higher total amount after 3 h is mostly due to increased 2-OG and citrate concentrations in the mutant (see below).

Changes in the composition of amino acid pools differ between WT and $\Delta nbID$.

Despite the overall trend of a decreasing total amino acid pool in mutant $\Delta nbID$ compared to WT, the direction of changes in the amounts of individual amino acids were not unidirectional (Fig. 3). According to their differential changes between WT and $\Delta nbID$ during the N starvation period, these changes could be categorized into four groups. Group 1 comprised amino acids showing no major differences between WT and $\Delta nbID$; group 2 included those amino acids that appeared different already at the +N condition, while group 3 exhibited differences between WT and $\Delta nbID$ early, after 3 h, or group 4 later, after 24 h of N starvation (Fig. 3).

Amino acids of group 1 showed similar trends in both WT and $\Delta nbID$ during nitrogen limitation, as was observed for Ala, Arg, Gln and His. The strongest deviation in the response to N starvation between the two strains was observed for Glu. The level of Glu did not decrease during the 24 h N starvation period in WT but decreased in mutant $\Delta nbID$ to approximately 17% of the initial value after 24 h of N starvation (Fig. 3). Glu is by far the most abundant soluble amino acid and serves as an amino group donor in the majority of transaminase reactions. Hence, the depletion of the Glu pool should have notable effects on the entire metabolism. Indeed, the lack of Glu extended into the pool of reduced glutathione (GSH), which is the tripeptide γ -L-glutamyl-L-cysteinylglycine; the initial increase observed in the WT was lower in cells of the mutant $\Delta nbID$ and led after 24 h into N limitation to a significantly reduced amount (Fig. 3). GSH plays an important role as a reducing agent in all living cells and was found to be essential in the acclimation of *Synechocystis* 6803 to environmental and redox perturbations (48); hence, its depletion probably should have an impact on redox homeostasis in the mutant $\Delta nbID$. The amino acids of group 3, such as Gly, Pro, Ser, Thr and Val, exhibited transiently increased levels

after 3 h in WT, which was not the case or was less pronounced in the mutant (Fig. 3). The branched chain amino acids Ile, Leu and Val share the same biosynthetic pathway, whereas Gly and Ser are closely linked to photorespiratory metabolism and can be interconverted by serine hydroxymethyltransferase (49). Phe and Thr exhibited increased levels in WT cells after 24 h of N starvation, while they returned to initial levels in $\Delta nbID$, accumulating at values of less than 50% of the WT. Finally, the aromatic amino acid Tyr from group 2 accumulated nearly 100-fold more in WT cells after 24 h of N starvation, whereas it remained at low levels in $\Delta nbID$ (Fig. 3). This finding is consistent with a previous analysis of *Synechocystis* 6803 WT after 24 h N starvation (47). Tyr represents approximately 5% of the amino acids stored in the phycobilisome (Fig. 4). The strong accumulation of Tyr in WT but not in $\Delta nbID$ suggests that it originated directly from phycobilisome degradation, assuming that in contrast to other amino acids, Tyr is not readily incorporated into the metabolism of N-starved cells.

Changes in other N-containing metabolites and organic acids

In addition to proteinogenic amino acids, many additional metabolites could be quantified. For example, ornithine, a key intermediate in the arginine catabolism pathway, was clearly depleted in $\Delta nbID$ upon N starvation compared to WT (Fig. 5). This points to the arginine catabolism pathway as another mechanism for stabilization of the Glu pool. In this pathway, the arginine-guanidine-removing enzyme (AgrE) together with proline oxidase (PutA) can produce Glu from the Arg molecules released from phycocyanin via ornithine (50–52). Therefore, the diminished ornithine pool adds another important element to the changed nitrogen homeostasis in cells of the $\Delta nbID$ strain. This assumption was also supported by

changes in the 2-OG pool, which was lower in WT than in the mutant under all conditions, indicating a relatively more N-starved state of $\Delta nbID$ cells. Moreover, we observed that the absence of NbID also affected the pool of 3-phosphoglyceric acid (3PGA), the stable CO₂ fixation product of RubisCO, which was decreased under all conditions. Excess 3PGA can be used to produce glycogen, which is highly accumulating in N-starved cells during chlorosis (53). Otherwise, some 3PGA can be diverted via phosphoglycerate mutase into the direction of lower glycolysis and the TCA cycle, which is primarily regulated by PirC (54). In contrast to the decreased flow into the oxidative branch, the reductive branch of the TCA cycle seemed to be less affected by the absence of NbID, since the pool of malate was similar in cells of the mutant and the WT (Fig. 5). However, succinate was clearly lowered in $\Delta nbID$ after 3 h under N limitation, probably due to altered consumption of succinate by the respiratory chain. In the mutant cells, the succinate pool stabilized at a similar quantity as was observed in WT cells under prolonged N depletion. Taken together, the metabolic changes observed in this study enabled us to conclude that the absence of NbID had a broad impact on primary metabolism.

Discussion

Comparison of the changes in $\Delta nbID$ with the previously characterized $\Delta nbIA1/A2$ mutant.

NbID is involved in the disassembly of phycobilisomes by targeting CpcB during acclimation to N starvation (19). In contrast to the WT, the $\Delta nbID$ mutant maintains the majority of its pigments under this condition resembling the non-bleaching phenotype of the previously reported $\Delta nbIA$ strains and suggesting that phycobilisomes are retained in these mutants to a large extent during the considered

time frame (see Fig. 1). In case of the $\Delta nbIA1/A2$ mutant, amino acid pools were analyzed after 24 h in the presence or absence of nitrogen. The authors categorized the changes in the amino acid pool sizes into $NbIA1/A2$ -dependent and -independent responses. Most amino acids (Gln, Glu, Gly, Ile, Leu, Met, Phe, Pro, Ser, Thr, Tyr, Val and also GSH) fell into the $NbIA1/A2$ -dependent category. In the present study, we analyzed the effects of the absence of $NbID$ on the amino acid and organic acid pools in the $\Delta nbID$ deletion mutant after 3 h and 24 h of N starvation. The observed differences between $\Delta nbID$ and WT in the amino acid pools were similar to those described for the $\Delta nbIA1/A2$ mutant (47) but were not identical, despite the fact that both types of proteins play a role in phycobilisome degradation. Five amino acids (Ile, Leu, Ser, Thr, Val) had after 24 h of N starvation higher levels in WT but lower levels in the $\Delta nbIA1/A2$ mutant compared to the control in N-replete media. While we observed the same changes for these 5 amino acids after 24 h N starvation, we noticed that their pools were peaking even higher in WT in the sample taken after 3 h. In the study by Kiyota et al. (47), the WT had been analyzed at even higher temporal resolution (after 1, 3, 6, 12 and 24 h of N starvation) and they reported that these 5 amino acids transiently overshoot at the 3 h value, hence our data are consistent. However, we found a transient accumulation of Ile, Leu, Ser, Thr, and Val in cells of the $\Delta nbID$ mutant like in the WT in the 3 h sample. Hence, our data allow to differentiate between an early $NbID$ -independent response for these 5 amino acids and a later response that was $NbID$ -dependent. This finding suggests that the early, very pronounced response was independent from amino acid release from the phycobilisomes, whereas they were mobilized at the later time point, which both the $\Delta nbID$ and $\Delta nbIA1/A2$ mutants failed to do.

The levels of four other amino acids (Pro, Gly, Glu, Tyr) and GSH behaved very similar in the $\Delta nbIA1/A2$ and $\Delta nbID$ mutants. However, for three amino acids (Gln, Met and Phe) we observed differences between the different non-bleaching mutants. Gln levels steadily declined in our study, both in WT and $\Delta nbID$ over the time of the experiment; hence, it appeared as an NbID-independent response (Fig. 3). For Met and Phe, similar or higher concentrations had been reported in the WT but lower in $\Delta nbIA1/A2$ upon N starvation, with Met even falling below the detection limit after 24 h (47). While we observed the same effect for the WT, in contrast the $\Delta nbID$ mutant exhibited transiently higher levels of Met and Phe after 3 h, and about the same levels after 24 h as at the beginning.

In the work by Kiyota et al. (47) some amino acids were identified as NbIA1/A2-independent (Ala, Asn, Asp, Lys and Trp). The changes in the pools sizes of these amino acids were also NbID-independent in our work. We can add Arg and His to this category, which were below detection limit in the work on NbIA, while we could not measure Asn. Moreover, the decline in Gln levels appeared as an NbID-independent response, different from the findings on NbIA.

Collectively, the overall responses of $\Delta nbIA1/A2$ and $\Delta nbID$ mutants were similar but some deviations appeared. For example, our investigation of the 3 h time point allowed to conclude that some amino acid changes are independent from amino acid releases via the phycobilisome degradation. Consequently, though NbIA1/NbIA2- and NbID-assisted phycobilisome degradation represents the most obvious protein degradation process due to visible pigment changes, faster changes in the proteome likely contribute to the dynamics of amino acid changes especially during the early phase of acclimation to N starvation.

Phycobilisome proteins and amino acid pools during nitrogen starvation.

Degradation of phycocyanin rods and their proteolytic recycling are assumed to contribute extensively to the soluble amino acid pool in the cell, making those amino acids available to be incorporated into other proteins to acclimate towards nutrient limitation. From the phycobilisome core, up to six rod antennas radiate, each of which is built by three hexameric discs connected with linker polypeptides CpcG and CpcC (55). One hexameric phycocyanin disc contains six subunits of each CpcA and CpcB.

Therefore, the most abundant phycobilisome proteins are CpcA and CpcB, which are rich in Ala, Gly, Leu, Val and Tyr. Assuming full phycobilisome turnover, including also all linker proteins and allophycocyanin core, it would provide mostly Ala (~14%), Leu, Gly (both ~8%), Ser, Val, Arg, and Asp (contributing 6-7% each to phycobilisome composition). Accordingly, the pools of these amino acids were expected to increase early during N-starvation-induced phycobilisome degradation in the WT but not in $\Delta nbID$. Such an increase was clearly observed for Tyr, whereas the changes in the amounts of the other amino acids differed less pronounced between WT and $\Delta nbID$ cells. The amounts of Gly, Leu, Ser, and Val were always higher in the WT than in the mutant under -N conditions. However, the amount of Ala, the most frequent amino acid in the phycobilisome showed a similar pool change in WT and mutant cells, suggesting its increase resulted from synthesis instead. Ala biosynthesis is derived from the transamination of pyruvate and Glu by alanine dehydrogenase (*sll1682*) or by serine-pyruvate aminotransferase using pyruvate and Ser. Therefore, Ala is directly linked to the metabolic pathways of Ser, Gly and Thr, as well as to primary C metabolism, via pyruvate. Indeed, in *Synechococcus elongatus* PCC 7942, alanine dehydrogenase was found to be

required for correct phycobilisome degradation during N starvation, resulting in lower *nblA* transcription if the gene for alanine dehydrogenase was deleted (56).

As mentioned above, besides Tyr the largest difference between WT and mutant $\Delta nblD$ was observed for the pool of Glu after 24 h of N starvation. Glu is can be regarded as the central amino acid from several perspectives. It is certainly the most relevant amino group donor for transaminations, consistent with its abundance, which is far higher than any other amino acid. In addition, Glu serves as the precursor of tetrapyrrole synthesis, leading to cytochromes and especially chlorophylls in cyanobacteria, green algae and plants upon its activation by binding to its cognate tRNA^{glu} (57–59). This C5 pathway is the sole biosynthetic pathway for tetrapyrrole biogenesis in *Synechocystis* 6803 (60, 61). Glu has an approximate 5% share in the composition of the phycobilisome proteins (Fig. 4). Therefore, it is likely that amino groups of abundant amino acids that become available upon phycobilisome degradation are utilized predominantly to stabilize the Glu pool via transaminase reactions. Initially, stabilization of the Glu pool may occur through glutamate synthase from Gln and 2-OG, which is also consistent with the relatively lower amounts of 2-OG in WT compared to $\Delta nblD$ cells (Fig. 5), then via arginine catabolism from Arg. A corresponding reduction in Gln and Arg pools is seen at 3 h and at 24 h into N starvation in both WT and $\Delta nblD$. At the later time point, additionally Asp, Gly, Leu, possibly Ala and Val, might be a source of amino groups as these amino acids are (i) relatively abundant, (ii) relatively frequent in the phycobilisome and (iii) their pools initially increased in the WT after shift to N starvation, followed by a particularly sharp reduction from 3 h to 24 h. In addition, the development of pool sizes for these amino acids showed marked differences between the WT and $\Delta nblD$.

Our results support the hypothesis that amino acid pools are regulated to a large extent independently of phycobilisome degradation by NblA1/A2 and NblD, because the cells maintain distinct amino acid pool homeostasis involving transaminase reactions, primarily using the amino groups from the released phycobilisome constituents. Furthermore, such amino acids as Ala, Asp and Lys might originate from the breakdown of several other proteins (47). Hence, even though NblA1/NblA2- and NblD-assisted phycobilisome degradation represents the most obvious protein degradation process due to visible pigment changes, additional changes in the proteome likely contribute to the recycling of amino acids during acclimation to N starvation.

Effects on the organic acids pools.

The striking differences between WT and $\Delta nblD$ observed for Glu at 24 h of N starvation may extend also further, into the amounts of certain organic acids, especially succinate. At 3 h -N, succinate levels were very low, whereas Glu levels were high; at 24 -N, succinate levels were relatively high, whereas Glu levels were very low. There are four possibilities for the relatively high succinate levels after 24 h of N starvation:

- (i) it may appear from an enhanced flux into the reductive branch of the TCA cycle working in anabolic direction in both strains, because the malate content also increased over time;
- (ii) an enhanced closure of the “open” cyanobacterial TCA cycle due to the shunt, which converts 2-OG into succinate via 2-oxoglutarate decarboxylase and succinic semialdehyde dehydrogenase reactions (62) would provide higher succinate amounts;

(iii) Succinate could also originate from the gamma-aminobutyric acid shunt closing the “open” TCA cycle in which glutamate is converted into gamma-aminobutyric acid and then via succinic semialdehyde to succinate (63);

(iv) The succinate consumption by respiration via succinate dehydrogenase could have become reduced.

The flux through the alternative TCA shunts has been regarded to be rather low under light conditions (62, 63) as applied here, therefore, the possibility (ii) appears rather unlikely. Possibility (iv) is unlikely as respiration does not decrease but increases upon N limitation (24). Consequently, the most likely scenario is that the succinate accumulation after 3 h N starvation in WT cells resulted from increased flow through the arginine catabolism pathway from phycobiliproteins (variant iii) via Glu, while the succinate accumulation after 24 h in the WT and especially pronounced in $\Delta nbID$ originates from the reductive branch of the TCA cycle (i). This scenario is supported by the N starvation-induced expression (53, 64) of *slr1022* encoding an enzyme that functions also as γ -aminobutyrate aminotransferase (63), a key enzyme in the gamma-aminobutyric acid shunt pathway interconverting Glu to succinate.

Connecting metabolic changes and the regulation of gene expression.

Several of the metabolites showing marked changes between WT and mutant $\Delta nbID$ possess signaling functions connecting metabolism and the regulation of gene expression. For example, Glu is used as a signaling molecule in green algae, mosses and higher plants sensed by the PII protein (65). Moreover, in several Gram-negative bacteria, Gln serves as a signaling molecule for N availability (reviewed by Forchhammer, 2013 (66)). More recently, it was observed that Gln also exhibits a

signaling function in *Synechocystis* 6803 and other cyanobacteria via interaction with the GlnA riboswitch governing the expression of *glnB* mRNA (67).

In contrast to Gram-negative bacteria, cyanobacteria use the TCA cycle intermediate 2-OG to sense the C/N status. 2-OG is the regulatory co-factor of NtcA and of PII, where the 2-OG accumulation transduces the low N signal leading to the activation of NtcA (36, 37). Conversely, under high N levels 2-OG is converted via the GS/GOGAT cycle to Glu, thereby the low 2-OG level diminishes the activity of NtcA. In addition, the decrease in 2-OG is characteristic for C limitation and activates the transcription factor NdhR to upregulate genes involved in the cyanobacterial inorganic C-concentrating mechanism (68, 69).

Hence, the increase in 2-OG in the WT observed during N starvation should represent a pronounced signal for activation of NtcA, and indeed, the transcriptional units within the NtcA regulon become strongly differentially regulated during this condition (19). Notably, there is an NtcA motif in the promotor region of the *nbID* gene itself (16), consistent with its role in N-limited conditions. However, compared to WT, we observed that 2-OG increased even more in $\Delta nbID$ and at every sampling point, including the +N condition (Fig. 5). Because of the function of 2-OG as the main cellular indicator metabolite to report C/N homeostasis (31–35), increased 2-OG levels therefore clearly indicated a reduced ammonia assimilating capacity of mutant $\Delta nbID$ already under N replete conditions. The likely reduced use of 2-OG in the GS/GOGAT cycle was supported by the observation that the levels of citrate and aconitate, precursors of 2-OG in the oxidative branch in the open cyanobacterial TCA cycle, were lowered in $\Delta nbID$ (Fig. 5). If the flux into the GS/GOGAT cycle would be similar, this decrease should extend into the 2-OG pool; however, its

increase indicates reduced 2-OG utilization in mutant cells, whereas 2-OG was extensively used in WT to replenish the Glu pool.

It could be expected that the higher 2-OG level in $\Delta nbID$ than in the WT cells would contribute to the stimulation of the NtcA regulon, which was observed in $\Delta nbID$ during the previous transcriptome analysis. Indeed, 3 h after the induction of N starvation, the mRNA levels of NtcA-regulated *nbIA1A2* genes were more strongly induced compared to the WT ($\log_2$ fold changes of +4.28 and +4.09 in $\Delta nbID$ and +3.71 and +3.47 in the WT), while *gifA* and *gifB* were more strongly reduced (for both genes, $\log_2$ fold change of -4.7 in $\Delta nbID$ and -2.4 in the WT control) (19). By comparison, 2-OG concentrations were around 3.14 mM and 1.16 mM in $\Delta nbID$ and WT, respectively (Table S3). Hence, a higher 2-OG concentration corresponded roughly to higher mRNA levels 3 h after the induction of N starvation. However, in the nitrogen-replete condition we measured an ~100 fold higher level of 2-OG in $\Delta nbID$ compared to WT but the *nbIA1A2* mRNA levels were only marginally higher. This result strongly suggests that the high accumulation of 2-OG alone is not sufficient to induce the complete N starvation response and an additional signal can be postulated. Furthermore, the high accumulation of 2-OG in $\Delta nbID$ already in the +N condition can be taken as evidence that NbID, in addition to its prominent function during N starvation, also plays some role under N-replete conditions. This function of NbID could be related to the remodeling of phycobilisome composition, size and number per thylakoid membrane or to the turnover of phycobilisomes occurring under N-replete conditions.

Materials and Methods

Cultivation conditions and sampling for metabolite analysis

Synechocystis 6803 PCC-M was used as the WT and background for the construction of mutants. *NbID* knockout mutants were generated by homologous replacement of the *nbID* coding sequence with a kanamycin resistance cassette (*nptII*) (19).

Strains were maintained in copper-free BG11 supplemented with 20 mM TES PUFFERAN®, pH 7.5, under continuous white light of 50 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ at 30°C.

Wild-type and $\Delta nbID$ cultures were grown to OD ~0.8 in triplicate without antibiotics before cells were pelleted by centrifugation (3,210 $\times g$, 8 minutes, room temperature), washed three times in BG11 without N and resuspended in BG11 lacking a N source. After three and 24 h of incubation in N-limited conditions, the OD was recorded, and 2 ml of each culture was collected by centrifugation (9,000 $\times g$, 1 minute, room temperature) and immersed in liquid nitrogen. As controls, samples were also obtained before N removal (+N). For extraction, 630 μl methanol containing carnitine (internal standard, 1 μg per extraction) was added, mixed for one minute and incubated for 5 minutes in a sonic water bath. After another 15 minutes of shaking at room temperature, 400 μl of chloroform was added, and the sample was incubated at 37°C for 5 minutes. Next, 800 μl ROTISOLV® LC-MS grade H₂O (Roth) was added, and the sample was mixed thoroughly. Precipitation was enabled by storage at -20°C for at least 2 h, and phase separation was subsequently achieved by centrifugation for 5 minutes at room temperature (16,800 $\times g$). The upper polar phase was collected and dried in a speed vac for 30 minutes followed by lyophilization overnight. A mock sample without any culture was subjected to the same treatments as the other samples.

471

472 **Metabolite analysis**

473 Absolute metabolite contents were quantified on a high-performance liquid
474 chromatograph mass spectrometer LCMS-8050 system (Shimadzu) as described
475 previously (70). Dried samples were dissolved in 200 µl LCMS grade water and
476 filtered through 0.2-mm filters (Omnifix-F, Braun, Germany), and 4 µl of the cleared
477 supernatant was separated on a pentafluorophenylpropyl column (Supelco
478 Discovery HS FS, 3 mm, 150 3 2.1 mm).

479 The compounds were identified and quantified using the multiple reaction monitoring
480 (MRM) values given in the LC-MS/MS method package and the LabSolutions
481 software package (Shimadzu). Authentic standard substances (Merck) at varying
482 concentrations were used for calibration, and peak areas were normalized to signals
483 of the internal standard (carnitine). The data were further normalized to the OD₇₅₀
484 measured for each sample and corrected by signals obtained from the mock sample.
485 The raw data and estimations as absolute values in nanogram × ml⁻¹ × OD₇₅₀⁻¹ are
486 given in Table S1, statistical evaluations in Table S4. These data were converted to
487 molar concentrations of metabolites per total cell volume assuming an average cell
488 diameter of 3 µm and 10⁷ cells per milliliter at OD₇₅₀ = 1 (Table S3).

489

490 **Funding**

491 We appreciate the support by the Deutsche Forschungsgemeinschaft (DFG, German
492 Research Foundation) to WRH through the graduate school MeInBio -
493 322977937/GRK2344 and to WRH and MH through the priority program “Small
494 Proteins in Prokaryotes, an Unexplored World” SPP 2002 (grants DFG HE2544/12-2

and DFG HA2002/22-2). The LC-MS equipment at the University of Rostock was financed through the Hochschulbauförderungsgesetz program (GZ: INST 264/125-1 FUGG to MH).

Conflict of interest statement

The authors declare that they have no conflicts of interest.

Author Contributions

WRH designed the project and secured funding. VK constructed and cultivated strains, MH and ST measured metabolite concentrations. VK, MH and WRH wrote the manuscript with input from all authors.

References

1. Meydan S, Marks J, Klepacki D, Sharma V, Baranov PV, Firth AE, Margus T, Kefi A, Vázquez-Laslop N, Mankin AS. 2019. Retapamulin-assisted ribosome profiling reveals the alternative bacterial proteome. *Mol Cell* 74:481-493.e6.
2. Weaver J, Mohammad F, Buskirk AR, Storz G. 2019. Identifying small proteins by ribosome profiling with stalled initiation complexes. *mBio* 10.
3. Hemm MR, Weaver J, Storz G. 2020. *Escherichia coli* small proteome. *EcoSal Plus* 9.
4. Fromme P, Melkozernov A, Jordan P, Krauss N. 2003. Structure and function of photosystem I: interaction with its soluble electron carriers and external antenna systems. *FEBS Lett* 555:40–44.
5. Guskov A, Kern J, Gabdulkhakov A, Broser M, Zouni A, Saenger W. 2009. Cyanobacterial photosystem II at 2.9-Å resolution and the role of quinones, lipids, channels and chloride. *Nat Struct Mol Biol* 16:334–342.
6. Kashino Y, Lauber WM, Carroll JA, Wang Q, Whitmarsh J, Satoh K, Pakrasi HB. 2002. Proteomic analysis of a highly active photosystem II preparation from the cyanobacterium *Synechocystis* sp. PCC 6803 reveals the presence of novel polypeptides. *Biochemistry* 41:8004–8012.
7. Baniulis D, Yamashita E, Whitelegge JP, Zatsman AI, Hendrich MP, Hasan SS, Ryan CM, Cramer WA. 2009. Structure-function, stability, and chemical modification of the cyanobacterial cytochrome b_6f complex from *Nostoc* sp. PCC 7120. *J Biol Chem* 284:9861–9869.
8. Allen JF. 2004. Cytochrome b_6f : structure for signalling and vectorial metabolism. *Trends Plant Sci* 9:130–137.
9. Schneider D, Volkmer T, Rögner M. 2007. PetG and PetN, but not PetL, are essential subunits of the cytochrome b_6f complex from *Synechocystis* PCC 6803. *Res Microbiol* 158:45–50.
10. Nowaczyk MM, Wulfhorst H, Ryan CM, Souda P, Zhang H, Cramer WA, Whitelegge JP. 2011. NdhP and NdhQ: two novel small subunits of the cyanobacterial NDH-1 complex. *Biochemistry* 50:1121–1124.
11. Schwarz D, Schubert H, Georg J, Hess WR, Hagemann M. 2013. The gene *sml0013* of *Synechocystis* sp. strain PCC 6803 encodes for a novel subunit of

- the NAD(P)H oxidoreductase or complex I that is ubiquitously distributed among Cyanobacteria. *Plant Physiol* 163:1191–1202.
12. Knoppová J, Sobotka R, Tichy M, Yu J, Konik P, Halada P, Nixon PJ, Komenda J. 2014. Discovery of a chlorophyll binding protein complex involved in the early steps of photosystem II assembly in *Synechocystis*. *Plant Cell* 26:1200–1212.
13. Komenda J, Sobotka R. 2016. Cyanobacterial high-light-inducible proteins--Protectors of chlorophyll-protein synthesis and assembly. *Biochim Biophys Acta* 1857:288–295.
14. Kufryk G, Hernandez-Prieto MA, Kieselbach T, Miranda H, Vermaas W, Funk C. 2008. Association of small CAB-like proteins (SCPs) of *Synechocystis* sp. PCC 6803 with photosystem II. *Photosyn Res* 95:135–145.
15. Hobbs EC, Fontaine F, Yin X, Storz G. 2011. An expanding universe of small proteins. *Curr Opin Microbiol* 14:167–173.
16. Baumgartner D, Kopf M, Klähn S, Steglich C, Hess WR. 2016. Small proteins in cyanobacteria provide a paradigm for the functional analysis of the bacterial micro-proteome. *BMC Microbiol* 16:285.
17. de Alvarenga LV, Hess WR, Hagemann M. 2020. AcnSP - a novel small protein regulator of aconitase activity in the cyanobacterium *Synechocystis* sp. PCC 6803. *Front Microbiol* 11:1445.
18. Baier K, Nicklisch S, Grundner C, Reinecke J, Lockau W. 2001. Expression of two *nblA*-homologous genes is required for phycobilisome degradation in nitrogen-starved *Synechocystis* sp. PCC6803. *FEMS Microbiol Lett* 195:35–39.
19. Krauspe V, Fahrner M, Spät P, Steglich C, Frankenberg-Dinkel N, Mačėk B, Schilling O, Hess WR. 2021. Discovery of a small protein factor involved in the coordinated degradation of phycobilisomes in cyanobacteria. *Proc Natl Acad Sci USA* 118:e2012277118.
20. Muro-Pastor MI, Reyes JC, Florencio FJ. 2005. Ammonium assimilation in cyanobacteria. *Photosynth Res* 83:135–150.
21. Herrero A, Muro-Pastor AM, Flores E. 2001. Nitrogen control in cyanobacteria. *J Bacteriol* 183:411–425.
22. Allen MM, Smith AJ. 1969. Nitrogen chlorosis in blue-green algae. *Archiv für Mikrobiologie* 69:114–120.

23. Görl M, Sauer J, Baier T, Forchhammer K. 1998. Nitrogen-starvation-induced chlorosis in *Synechococcus* PCC 7942: adaptation to long-term survival. *Microbiology* 144:2449–2458.
24. Krasikov V, Aguirre von Wobeser E, Dekker HL, Huisman J, Matthijs HC. 2012. Time-series resolution of gradual nitrogen starvation and its impact on photosynthesis in the cyanobacterium *Synechocystis* PCC 6803. *Physiol plant* 145:426–439.
25. Sauer J, Schreiber U, Schmid R, Völker U, Forchhammer K. 2001. Nitrogen starvation-induced chlorosis in *Synechococcus* PCC 7942. Low-level photosynthesis as a mechanism of long-term survival. *Plant Physiol* 126:233–243.
26. Spät P, Klotz A, Rexroth S, Mačák B, Forchhammer K. 2018. Chlorosis as a developmental program in cyanobacteria: The proteomic fundament for survival and awakening. *Mol Cell Proteomics* 17:1650–1669.
27. Collier JL, Grossman AR. 1992. Chlorosis induced by nutrient deprivation in *Synechococcus* sp. strain PCC 7942: not all bleaching is the same. *J Bacteriol* 174:4718–4726.
28. Gründel M, Scheunemann R, Lockau W, Zilliges Y. 2012. Impaired glycogen synthesis causes metabolic overflow reactions and affects stress responses in the cyanobacterium *Synechocystis* sp. PCC 6803. *Microbiology*, 158:3032–3043.
29. Meeks JC, Wolk CP, Lockau W, Schilling N, Shaffer PW, Chien WS. 1978. Pathways of assimilation of $^{13}\text{N}_2$ and $^{13}\text{NH}_4^+$ by cyanobacteria with and without heterocysts. *J Bacteriol* 134:125–130.
30. Florencio FJ, García-Domínguez M, Martín-Figueroa E, Crespo JL, Navarro F, Muro-Pastor MI, Reyes JC. 1998. Ammonium assimilation in cyanobacteria. The regulation of the GS-GOGAT pathway, p. 3607–3612. *In* Garab, G (ed.), *Photosynthesis: Mechanisms and Effects: Volume I–V: Proceedings of the XIth International Congress on Photosynthesis, Budapest, Hungary, August 17–22, 1998*. Springer Netherlands, Dordrecht.
31. Vázquez-Bermúdez MF, Herrero A, Flores E. 2002. 2-Oxoglutarate increases the binding affinity of the NtcA (nitrogen control) transcription factor for the *Synechococcus glnA* promoter. *FEBS Lett* 512:71–74.

- 603 32. Tanigawa R, Shirokane M, Maeda S -i., Omata T, Tanaka K, Takahashi H.
604 2002. Transcriptional activation of NtcA-dependent promoters of
605 *Synechococcus* sp. PCC 7942 by 2-oxoglutarate *in vitro*. Proc Natl Acad Sci
606 USA 99:4251–4255.
- 607 33. Espinosa J, Forchhammer K, Burillo S, Contreras A. 2006. Interaction network
608 in cyanobacterial nitrogen regulation: PipX, a protein that interacts in a 2-
609 oxoglutarate dependent manner with PII and NtcA. Mol Microbiol 61:457–469.
- 610 34. Espinosa J, Rodriguez-Mateos F, Salinas P, Lanza VF, Dixon R, de la Cruz F,
611 Contreras A. 2014. PipX, the coactivator of NtcA, is a global regulator in
612 cyanobacteria. Proc Natl Acad Sci USA 111:E2423–E2430.
- 613 35. Forcada-Nadal A, Llcer JL, Contreras A, Marco-Marn C, Rubio V. 2018. The
614 PII-NAGK-PipX-NtcA regulatory axis of cyanobacteria: A tale of changing
615 partners, allosteric effectors and non-covalent interactions. Front Mol Biosci
616 5:91.
- 617 36. Forchhammer K. 2004. Global carbon/nitrogen control by PII signal transduction
618 in cyanobacteria: from signals to targets. FEMS Microbiol Rev 28:319–333.
- 619 37. Forchhammer K. 2008. PII signal transducers: novel functional and structural
620 insights. Trends Microbiol 16:65–72.
- 621 38. Giner-Lamia J, Robles-Rengel R, Hernndez-Prieto MA, Muro-Pastor MI,
622 Florencio FJ, Futschik ME. 2017. Identification of the direct regulon of NtcA
623 during early acclimation to nitrogen starvation in the cyanobacterium
624 *Synechocystis* sp. PCC 6803. Nucleic Acids Res 45:11800–11820.
- 625 39. Garca-Domnguez M, Reyes JC, Florencio FJ. 2000. NtcA represses
626 transcription of *gifA* and *gifB*, genes that encode inhibitors of glutamine
627 synthetase type I from *Synechocystis* sp. PCC 6803. Mol Microbiol 35:1192–
628 1201.
- 629 40. Collier JL, Grossman AR. 1994. A small polypeptide triggers complete
630 degradation of light-harvesting phycobiliproteins in nutrient-deprived
631 cyanobacteria. EMBO J 13:1039–1047.
- 632 41. Karradt A, Sobanski J, Mattow J, Lockau W, Baier K. 2008. NblA, a key protein
633 of phycobilisome degradation, interacts with ClpC, a HSP100 chaperone partner
634 of a cyanobacterial Clp protease. J Biol Chem 283:32394–32403.

42. Baier A, Winkler W, Korte T, Lockau W, Karradt A. 2014. Degradation of phycobilisomes in *Synechocystis* sp. PCC6803: evidence for essential formation of an NblA1/NblA2 heterodimer and its codegradation by a Clp protease complex. *J Biol Chem* 289:11755–11766.
43. Nguyen AY, Bricker WP, Zhang H, Weisz DA, Gross ML, Pakrasi HB. 2017. The proteolysis adaptor, NblA, binds to the N-terminus of β -phycocyanin: Implications for the mechanism of phycobilisome degradation. *Photosyn Res* 132:95–106.
44. Sendersky E, Kozier N, Levi M, Moizik M, Garini Y, Shav-Tal Y, Schwarz R. 2015. The proteolysis adaptor, NblA, is essential for degradation of the core pigment of the cyanobacterial light-harvesting complex. *Plant J* 83:845–852.
45. Elmorjani K, Herdman M. 1987. Metabolic control of phycocyanin degradation in the cyanobacterium *Synechocystis* PCC 6803: a glucose effect. *Microbiology* 133:1685–1694.
46. Hu P-P, Hou J-Y, Xu Y-L, Niu N-N, Zhao C, Lu L, Zhou M, Scheer H, Zhao K-H. 2020. The role of lyases, NblA and NblB proteins and bilin chromophore transfer in restructuring the cyanobacterial light-harvesting complex. *Plant J* 102:529–540.
47. Kiyota H, Hirai MY, Ikeuchi M. 2014. NblA1/A2-dependent homeostasis of amino acid pools during nitrogen starvation in *Synechocystis* sp. PCC 6803. *Metabolites* 4:517–531.
48. Cameron JC, Pakrasi HB. 2010. Essential role of glutathione in acclimation to environmental and redox perturbations in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Physiol* 154:1672–1685.
49. Eisenhut M, Kahlon S, Hasse D, Ewald R, Lieman-Hurwitz J, Ogawa T, Ruth W, Bauwe H, Kaplan A, Hagemann M. 2006. The plant-like C2 glycolate cycle and the bacterial-like glycerate pathway cooperate in phosphoglycolate metabolism in cyanobacteria. *Plant Physiol* 142:333–342.
50. Quintero MJ, Muro-Pastor AM, Herrero A, Flores E. 2000. Arginine catabolism in the cyanobacterium *Synechocystis* sp. strain PCC 6803 involves the urea cycle and arginase pathway. *J Bacteriol* 182:1008–1015.
51. Zhang H, Yang C. 2019. Arginine and nitrogen mobilization in cyanobacteria. *Mol Microbiol* 111:863–867.

52. Zhang H, Liu Y, Nie X, Liu L, Hua Q, Zhao G-P, Yang C. 2018. The cyanobacterial ornithine-ammonia cycle involves an arginine dihydrolase. *Nat Chem Biol* 14:575–581.
53. Klotz A, Georg J, Bučinská L, Watanabe S, Reimann V, Januszewski W, Sobotka R, Jendrosseck D, Hess WR, Forchhammer K. 2016. Awakening of a dormant cyanobacterium from nitrogen chlorosis reveals a genetically determined program. *Curr Biol* 26:2862–2872.
54. Orthwein T, Scholl J, Spät P, Lucius S, Koch M, Macek B, Hagemann M, Forchhammer K. 2021. The novel PII-interactor PirC identifies phosphoglycerate mutase as key control point of carbon storage metabolism in cyanobacteria. *Proc Natl Acad Sci USA* 118:e2019988118.
55. Arteni AA, Ajlani G, Boekema EJ. 2009. Structural organisation of phycobilisomes from *Synechocystis* sp. strain PCC6803 and their interaction with the membrane. *Biochim Biophys Acta* 1787:272–279.
56. Lahmi R, Sendersky E, Perelman A, Hagemann M, Forchhammer K, Schwarz R. 2006. Alanine dehydrogenase activity is required for adequate progression of phycobilisome degradation during nitrogen starvation in *Synechococcus elongatus* PCC 7942. *J Bacteriol* 188:5258–5265.
57. Wang W-Y, Gough SP, Kannangara CG. 1981. Biosynthesis of Δ -aminolevulinate in greening barley leaves IV. Isolation of three soluble enzymes required for the conversion of glutamate to Δ -aminolevulinate. *Carlsberg Res Commun* 46:243.
58. Kannangara CG, Gough SP, Oliver RP, Rasmussen SK. 1984. Biosynthesis of Δ -aminolevulinate in greening barley leaves VI. Activation of glutamate by ligation to RNA. *Carlsberg Res Commun* 49:417.
59. Schön A, Krupp G, Gough S, Berry-Lowe S, Kannangara CG, Söll D. 1986. The RNA required in the first step of chlorophyll biosynthesis is a chloroplast glutamate tRNA. *Nature* 322:281–284.
60. O'Neill GP, Peterson DM, Schön A, Chen MW, Söll D. 1988. Formation of the chlorophyll precursor delta-aminolevulinic acid in cyanobacteria requires aminoacylation of a tRNA^{Glu} species. *J Bacteriol* 170:3810–3816.

61. Rieble S, Beale SI. 1988. Transformation of glutamate to delta-aminolevulinic acid by soluble extracts of *Synechocystis* sp. PCC 6803 and other oxygenic prokaryotes. *J Biol Chem* 263:8864–8871.
62. Zhang S, Bryant DA. 2011. The tricarboxylic acid cycle in cyanobacteria. *Science* 334:1551–1553.
63. Xiong W, Brune D, Vermaas WFJ. 2014. The γ -aminobutyric acid shunt contributes to closing the tricarboxylic acid cycle in *Synechocystis* sp. PCC 6803. *Mol Microbiol* 93:786–796.
64. Kopf M, Klähn S, Scholz I, Matthiessen JKF, Hess WR, Voß B. 2014. Comparative analysis of the primary transcriptome of *Synechocystis* sp. PCC 6803. *DNA Res* 21:527–539.
65. Chellamuthu V-R, Ermilova E, Lapina T, Lüddecke J, Minaeva E, Herrmann C, Hartmann MD, Forchhammer K. 2014. A widespread glutamine-sensing mechanism in the plant kingdom. *Cell* 159:1188–1199.
66. Forchhammer K. 2007. Glutamine signalling in bacteria. *Front Biosci* 12:10–2741.
67. Klähn S, Bolay P, Wright PR, Atilho RM, Brewer KI, Hagemann M, Breaker RR, Hess WR. 2018. A glutamine riboswitch is a key element for the regulation of glutamine synthetase in cyanobacteria. *Nucleic Acids Res* 46:10082–10094.
68. Jiang Y-L, Wang X-P, Sun H, Han S-J, Li W-F, Cui N, Lin G-M, Zhang J-Y, Cheng W, Cao D-D, Zhang Z-Y, Zhang C-C, Chen Y, Zhou C-Z. 2018. Coordinating carbon and nitrogen metabolic signaling through the cyanobacterial global repressor NdhR. *Proc Natl Acad Sci USA* 115:403–408.
69. Klähn S, Orf I, Schwarz D, Matthiessen JKF, Kopka J, Hess WR, Hagemann M. 2015. Integrated transcriptomic and metabolomic characterization of the low-carbon response using an *ndhR* mutant of *Synechocystis* sp. PCC 6803. *Plant Physiol* 169:1540.
70. Reinholdt O, Schwab S, Zhang Y, Reichheld J-P, Fernie AR, Hagemann M, Timm S. 2019. Redox-regulation of photorespiration through mitochondrial thioredoxin o1. *Plant Physiol* 181:442–457.

Figures

Figure 1. Response of the *Synechocystis* 6803 WT and $\Delta nbID$ mutant during the response to N starvation. **A.** Phenotype under nitrogen-replete (+N) conditions and after 24 h of nitrogen depletion (-N). **B.** Spectrometric measurements of cultures normalized to OD₇₅₀.

Figure 2. A. Total soluble amino acid content and **B.** total soluble metabolite content (non-proteinogenic amino acids and important intermediates) of wild-type (blue) and $\Delta nbID$ (red) cultures in nitrogen-replete (+N) and nitrogen-depleted BG11 (-N) after 3 h and 24 h. For each time point and strain, triplicates were measured, and significance ($p < 0.05$) was calculated by t-test; n.s. = not significant. The raw data are given in Table S2.

Figure 3. Development of amino acid content (in mM for Glu and μ M for all other) in WT (blue) and $\Delta nblD$ mutant (red) in nitrogen replete (+N) and nitrogen deplete (-N) medium after 3 h and 24 h. Amino acids are grouped together according to the observed differences between wild type and mutant as follows: Difference already observed in +N conditions; no difference between strains; early difference (after 3 h); later differences (24 h). Significance was calculated with a two-sample t-test with unequal variance (Welch's t-test; $p < 0.05 = *$; $p < 0.01 = **$ and $p < 0.001 = ***$) for

756 the development in each strain and between the strains at corresponding time points
 757 (see Table S4 for details).

758

759 **Figure 4.** Phycobilisome individual amino acid (aa) composition (in %) plotted
760 against their increase or decrease observed in relation to the initial values measured
761 in nitrogen sufficient conditions. **A, C.** WT after 3 h of N limitation; **B, D.** WT after 24
762 h of N limitation. **E, F.** Data for $\Delta nblD$ mutant after 3 and 24 h. Phycobilisome
763 composition was calculated based on the structure description by Arteni et al. (55),
764 assuming a common complex with 6 rods each with 3 hexameric phycocyanin discs
765 and a tricylindrical allophycocyanin core. A linear trend line shows very low
766 correlation between the variables, and a poor R-squared value indicates high
767 variation of the data in relation to this line.

Figure 5. Comparison of metabolite content (in mM for Glu, citrate, 3PGA, malate and 2-OG, in μM for all other) in WT (blue) and $\Delta nblD$ deletion mutant (red) in nitrogen replete (+N) and nitrogen deplete (-N) medium after 3 h and 24 h shown alongside their biosynthetic pathways. Succinate was below detection limit (b.d.) in the $\Delta nblD$ sample under nitrogen-replete conditions. Significance was calculated with the two-sample t-test with unequal variance (Welch's t-test; $p < 0.05 = *$; $p < 0.01 = **$ and $p < 0.001 = ***$) for the development in each strain and between the strains at corresponding time points (details in Table S4).

Supplementary Tables

Table S1. LC-MS/MS data for 18 amino acids (Cys and Asn are missing because they were below detection limit), several organic acids and GSH. Retention time (r) was individually monitored with 2 standards (line 4 and 5) and the average peak area for 1 ng (line 7), which were used to determine the ng per sample. The carnitine standard (columns E, F, G, H) was used to generate the Factor IS (internal standard), which is implemented in the normalization of the peak area for each sample.

Table S2. Total amount of all amino acids and the other soluble metabolites investigated in this study. Average, standard deviation and significance calculated by two-sample t-test (Welch's t-test) are visualized in figure 2.

Table S3. Conversion in molar concentrations of metabolites in μM (Glu in mM) per total cell volume assuming an average cell diameter of $3\ \mu\text{m}$ and 10^7 cells per milliliter at $\text{OD}_{750} = 1$.

Table S4. Welch's t-test (two-sided, unequal variation) for evaluating significant differences in the amino acids and organic acids between the strains and the development in the strains. Significant differences are marked in red ($p < 0.05$), boxed ($p < 0.01$) and bold italics ($p < 0.001$).