Deciphering microbiome impacts on fungal-microalgal interaction in a marine environment using metabolomics

Olivier Berry¹, Enora Briand², Alizé Bagot^{1,2}, Maud Chaigne^{1,2}, Laurence Meslet-Cladière³, Julien Wang¹, Olivier Grovel¹, Jeroen J. Jansen⁴, Nicolas Ruiz¹, Thibaut Robiou du Pont¹, Yves François Pouchus¹, Philipp Hess², Samuel Bertrand¹

¹ Université de Nantes, MMS, EA 2160, Nantes, France

² IFREMER, DYNECO, Laboratoire Phycotoxines, F-44000 Nantes, France

³ Laboratoire Universitaire de Biodiversité et Écologie Microbienne, Université Brest, F-29280 Plouzané, France

⁴ Radboud University, Institute for Molecules and Materials, Heyendaalseweg 135, 6525 AJ Nijmegen, The Netherlands

Corresponding author: Samuel Bertrand, Laboratoire Mer, Molécules, Santé – EA 2160, UFR des Sciences Pharmaceutiques et Biologiques, Université de Nantes, Nantes, France.

samuel.bertrand@univ-nantes.fr

Abstract

The comprehension of microbial interactions is one of the key challenges in microbial ecology. The present study focuses on studying the chemical interaction between the toxic dinoflagellate *Prorocentrum lima* PL4V strain and associated fungal strains (two *Penicillium* sp. strains and three *Aspergillus* sp) among which the *Aspergillus pseudoglaucus* strain MMS1589 was selected for further co-culture experiment. Such rarely studied interaction (fungal-microalgal) was explored in axenic and non-axenic conditions, in a dedicated microscale marine environment (hybrid solid/liquid conditions), to delineate specialized metabolome alteration in relation to the *P. lima* and *A. pseudoglaucus* co-culture in regard to the presence of their associated bacteria. Such alteration was monitored by high-performance liquid chromatography coupled to high-resolution mass spectrometry (LC-HRMS). In-depth analysis of the resulting data highlighted (1) the chemical modification associated to fungal-microalgal co-culture, and (2) the impact of associated bacteria in microalgal resilience to fungal interaction. Even if only a very low number of highlighted metabolites were fully characterised due to the poor chemical investigation of the studied species, a clear co-culture induction of the dinoflagellate toxins okadaic acid and dinophysistoxin 1 was observed. Such results highlight the importance to consider microalgal microbiome to study parameters regulating toxin production. Finally, a microscopic observation showed an unusual physical interaction between the fungal mycelium and the dinoflagellates.

Keywords: *Aspergillus pseudoglaucus*, metabolomics, microbial co-culture, microbiome, *Prorocentrum lima*

Introduction

In the marine environment, microalgae are in constant interaction, both at the intra- and inter-specific levels, which span from competition, mutualism, symbiosis, parasitism and predation [1]. These large diversity of biotic interactions are governed by trophic relationships (*e.g.* nutrients acquisition) and chemical communication mediated by diffusible signalling metabolites [2, 3]. Among those marine microalgae, toxic dinoflagellates represent particular microorganisms that may drastically impact marine ecosystems, *e.g.* reduction of fishes and algae diversity, as well as, human health hazards (seafood contamination) [4, 5]. Harmful algal blooms (HABs) are considered detrimental events, and HAB species are increasingly observed in some ecosystems, especially at human activities' proximity [4]. One HABs-related threat is diarrheic shellfish poisoning which is caused by toxins, like okadaic acid (OA). These latter polyketide toxins are produced by various marine dinoflagellates, including *Dinophysis* spp. and *Prorocentrum* spp. Such production is known to be influenced by nitrogen levels, light intensity, temperature and/or salinity [6,7]. The high energy investment required by the biosynthesis of such high molecular weight molecules point out the possible fitness advantage these toxins procure to their producers [8].

Prorocentrum lima has benthic lifestyle [9], which implies complex mechanisms to compete for nutrients, light and space [10]. Such competition requires the production of specialized metabolites, which are often evoked or revealed indirectly, while only a few studies highlight such molecules. This is partly due to technical difficulties appearing in their isolation and subsequent characterisation (*e.g.* quantities). Besides abiotic inducers, OA is involved in the toxic response to *Artemia salina* chemical cues. The exposition of *P. lima* to such predator yields OA over-production, and following impaired survival rates in predator population [11]. However, it remains unclear whether OA exerts a broad spectrum deterrence effect, since only a few number of studies directly linked OA exposure to negative predator outcomes [12]. As an example, such response is also observed with *Alexandrium minutum* in response to the copepod *Acartia tonsa* which produces copepodamides (polar lipids) yielding an

increase up to 20-fold of parasitic shellfish toxin production [13]. Besides predators, co-occurring dinoflagellates are also competitors [14]. For instance, *P. lima* was shown to produce a complex chemical mixture (including OA) possessing growth inhibitory against other co-occurring species [15]. Therefore, *P. lima* seems equipped with various competition chemical signals suited to its benthic epiphytic lifestyle.

Aside from predators and competitors, microalgae are known to interact with various bacteria with mutualistic or defensive behaviours. They can alter, through specific metabolite production, the composition of co-occurring bacterial communities [16], or the bacterial metabolism, including quorum sensing [17]. In parallel, bacteria interfere with microalgal physiology, *e.g.* through chemical signals from their quorum sensing systems. For example, *Pseudoalteromonas piscicida* releases 2-heptyl-4-quinolone when co-cultured with *Emiliana huxleyi*, causing its higher mortality rates [18]. Moreover, mutualistic evidence in microalgal-bacterial interactions has been shown [19–21]. One example is microalgal dependency on vitamin B12. Some microalgae, living with vitamin B12 bacteria producers, have lost their costly methionine synthase pathway for a less costly vitamin B12 dependent one. Consequently, such microalgae rely entirely on symbiosis with bacteria producing vitamin B12 to be able to grow [19]. In return, bacteria benefit from microalgal carbonated exudates [20]. While there is clear evidence of associated bacteria effect on microalgal metabolism, alteration of toxin production is still unclear [22–24] and only supported by few studies [25, 26].

Meanwhile, in fungi parasitic microscopic chytrid zoospores are known for using microalgae carbohydrates exudates to detect preys [27] and diatoms are able to alter these parasite life cycles after first fungal induced wound, by fast metabolism adaptation towards production of lipid derived polyenals and associated aldehydes [28]. An example of parasite induced toxin production is the up-regulation and production of microalgal self-growth-inhibitory alkaloids when invaded by *Lagenisma coscinodisci* [29]. Interestingly, no data is available on the existence of chemical interactions

between saprophytic marine fungi and microalgae while physical associations are already reported to induce microalgal aggregation in bioprocesses [30].

The study of such microbial interaction *in vitro*, remains a very challenging process [31]. Up to now, the best strategies rely on exploring bipartite interaction in controlled conditions [32]. Metabolomics, defined as a non-selective, comprehensive analytical approach for the identification and quantification of metabolites in a biological system [33], remains a method of choice to highlight chemical modification induced by microbial co-culture [33, 34]. Various examples already showed the induction of specific metabolites in microbial co-culture [34] through the expression of hidden biosynthesis gene clusters (BGCs) [35].

This study of chemical mediation between microalgae and saprophytic fungi in a marine environment, focuses on the toxin producer *P. lima* PL4V strain (known to produce OA and dinophysistoxin 1 – DTX-1). It aims at exploring (1) the presence of fungal strains within the microbiome of this epiphytic dinoflagellate, (2) the chemical mediation between *P. lima* and a fungal strain (namely *Aspergillus pseudoglaucus* MMS1589) co-occurring in culture and (3) whether the fungus may influence microalgal toxin production. Therefore, after fungal strain isolation from the *P. lima* PL4V strain, chemical modifications related to microbial co-cultures were highlighted through a metabolomic approach using a high-performance liquid chromatography coupled to high-resolution mass spectrometry (LC-HRMS).

Materials and Methods

Microalgae

Prorocentrum lima is a benthic dinoflagellate that may be considered a species complex as there is both temperate and tropical strains reported for this species [36]. The *P. lima* PL4V strain was collected by colleagues from the Instituto Espanol Oceanografico (IEO, Vigo) in the lagoon *Lago dos Nenos* at the

junction of the two Cíes Islands Illa de Monteagudo and Illa de Faro (Ria de Vigo, Galicia, Spain) [37], a temperate area, and maintained at the French Research Institute for the Exploitation of the Sea (IFREMER).

Culture media

All media used for fungal strain isolation were prepared using Reef Crystal mixture at 36 g/L (enriched marine salt from Aquarium Systems) to meet standard sea water salt concentrations and autoclaved at 121°C during 20 min. Chloramphenicol (50 mg/mL, Poly Labo, Paul Block & Cie, Strasbourg, France) was added in liquid or solid media used for fungal strain isolation, to avoid bacterial growth. Malt Extract (ME) liquid media was prepared with following concentrations: glucose 20 g/L, peptone (Biokar Diagnostics, Beauvais, France) 1 g/L, Malt Extract (Conda pronadisa, Madrid, Spain) 30 g/L, copper sulphate 5 mg/L, zinc sulphate 1 mg/L. The Dextrose Casein (DC) medium was prepared as follow: glucose 40 g/L, peptone 10 g/L. Solid equivalent of these media were also prepared. Dextrose Casein Agar (DCA) used DCA mixture (Becton, Dickinson and Company, Sparks, NV USA) 65 g/L and the Malt Extract Agar (MEA, Biokar Diagnostics) medium was composed of glucose 20 g/L, peptone 1 g/L, Malt Extract mixture (Conda pronadisa) 45 g/L, copper sulphate 5 mg/L, zinc sulphate 1 mg/L.

For the co-culture experiment, f/2 microalgal growth medium, without silica addition, was used as previously described [38]. The solid f/2 medium (f/2GA) was prepared using f/2 liquid medium complemented by 20 g/L of agar and 2 g/L of glucose. In some cases, large spectrum antibiotics mixture was added to f/2 liquid medium to provide the following final concentration in the culture medium: ampicillin 500 µg/mL, gentamycin 100 µg/mL, kanamycin 200 µg/mL [39].

Fungal isolation from microalgal biomass

To isolate fungal strains, *P. lima* PL4V biomass was inoculated on several liquid and solid culture media (ME, DC, MEA and DCA media). Fungal growth at 27 °C was evaluated every day. When fungal colonies

were observed, each colony was further collected and inoculated on DCA medium for storage and morphological identification based on macroscopic and microscopic features. The MMS1589 strain was further identified based on molecular biology information (ITS and β -tubulin sequencing) according to previously published protocol [40]. The ITS and β -tubulin sequences for MMS1589 strain were deposited in GenBank under accession numbers MN134000.1 and MN164633.1, respectively.

Co-culture growth in miniaturised liquid/solid environment

The miniaturised liquid/solid environment was prepared in tilted 50 mL flasks (Figure 1). Firstly, 10 mL f/2GA medium were added in the flask. Subsequently, the *A. pseudoglossus* MMS1589 strain was inoculated by spiking it into the middle of the f/2GA medium. After five days of fungal growth at 24°C under 12h/12h dark-light cycle using artificial light (DENNERLE Nano Light 11 watt, 6000 Kelvin, 900 lumen), 10 mL liquid f/2 medium (with or without antibiotic cocktail), and 15 mL *P. lima* PL4V inoculum in f/2 medium at 30 600 cells/mL were added to the flask. The co-culture was then incubated for six days under 12h/12h dark-light cycle. In parallel, the same design was used for both microorganisms alone (monoculture), where only *A. pseudoglossus* MMS1589 or *P. lima* PL4V was inoculated. In addition, blank conditions were performed without microorganism inoculation.

Five replicates of each experiment were performed, leading to thirty samples (with antibiotics: five co-cultures, five *A. pseudoglossus* and five *P. lima* monocultures; without antibiotics: five co-cultures, five *A. pseudoglossus* and five *P. lima* monocultures).

Figure 1. Experimental design of microscale co-culture experiments of marine benthic chemical interactions. Notice the clarity of the water when microalgal inoculum is treated with antibiotics.

Biomass extraction

After six days, for each growth experiment in liquid/solid environment, the liquid overlaying phase (LOP) was separated from the solid phase (resulting f/2 GA medium and fungus/microalgae) and further centrifuged to remove any cellular debris. Each LOP was extracted three times with 5 mL ethyl acetate (EtOAc), and the three resulting extracts were pooled and dried under vacuum using SpeedVac™ concentrator (Thermo Scientific). The solid phase was extracted twice with 5 mL EtOAc using an ultrasonic bath for 10 min prior to filtration. The two solid phase extracts were pooled and dried under vacuum using a SpeedVac™ concentrator. All extracts were weighed after evaporation to dryness.

LC-HRMS profiling

Each extract was diluted to a concentration of 1 mg dry weight/mL in HPLC-grade methanol (Biosolve Chimie, Dieuze, France). LC-HRMS profiling was achieved using a UFLC-ESI-IonTrap-TOFMS system (Shimadzu, Marne-la-Vallée, France) according to a previously published protocol [41]. Briefly, chromatographic separation was achieved using a Kinetex™ C₁₈ column (100 x 2.1 mm, 2.6 µm), at 40 °C with a 0.3 mL/min flow rate. The mobile phase is a gradient starting at 15 % acetonitrile (B) for 2 min, followed by an increase to 100 % over 23 min and further maintained for 5 min. Mass detection was set between m/z 100-1000 with fast switching between positive and negative ionisation mode (PI and NI).

In the sequences, samples were injected randomly using a dedicated Excel macro [42] and MeOH blanks and QC samples were injected every 15 samples. The QC sample was prepared by mixing 30 µL of all samples together.

Data Analysis

LC-HRMS data were exported as CDF files using LC Solution (Shimadzu) to allow automatic peak picking using MZmine 2 (ref.43). The mass detection was performed using a centroid algorithm with a noise level of 1.5E4. The chromatogram building used a minimum time span of 0.03 min, group intensity threshold of 1E5, a minimal of intensity of 1.5E4 and m/z tolerance of 30 ppm [44]. The peak deconvolution was performed using the baseline cut-off algorithm with median m/z centre calculation, a peak duration range between 0.05 and 10 min. The chromatograms were deisotoped with m/z tolerance of 1 mDa, retention time tolerance of 0.1 min and maximum charge of 3. The peak list was aligned using the Join aligner algorithm with m/z tolerance of 1 mDa or 30 ppm, retention time tolerance of 0.2 min, using the same weight for retention time and m/z importance. Subsequently, the gap-filling step was performed with a peak-finder algorithm, intensity tolerance of 1.0 m/z tolerance

of 1 mDa or 30 ppm, retention time tolerance of 0.2 min. The results were exported as a .csv file containing all peaks observed and referenced by their mass to charge ratio (m/z) and retention times (t_R) together with their respective peak areas in each sample, yielding the data matrix.

The peaks detected in blanks were withdrawn from the data matrix as background signal. The absence of ions corresponding to antibiotics and their possible anticipated metabolites using Biotransformer [45] (with phase I, II CYP450 and microbiota transformation) were confirmed. Then, the data matrix containing 6627 features was transformed to reflect full compound production by multiplying each peak area by the corresponding extract amount.

Finally, all data from one culture were concatenated to provide the final data matrix, thus data from replicated sample contained features (m/z at t_R with corrected peak areas) obtained from LOP extract profiles (NI and PI) and from solid phase extract profiles (NI and PI). Thus, all 30 samples were characterised by 13254 features.

The final data matrix was statistically analysed using R 4.0 (CRAN) with the PocheRmon function [46]. This function allows to analyse co-culture data univariately using the stats package [47], and multivariately using both the ropls package [48] and the Projected Orthogonalized CHEMical Encounter MONitoring (POChEMon) approach [49] encoded in R. All these approaches allowed to highlight statistically significant features. Highlighted features were confirmed manually in the raw LC-HMS data to remove false peaks detected during the automated peak detection step.

Annotation of statistically relevant features

The highlighted features were further annotated using traditional dereplication strategy [50] based on mass and spectral accuracy of the MS spectra. Accurate masses (up to 30 ppm for smaller peaks) were searched after adduct mass correction ($[M+H]^+$, $[M+Na]^+$, $[M-H]^-$, $[M+Cl]^-$ and $[M+formic\ acid-H]^-$) were considered [51] for compounds reported to be produced by either *Prorocentrum* sp. and *Aspergillus* sp. (also considering its teleomorph *Eurotium* sp. [52]) in the Dictionary of Natural

Products® (CRC Press, v.29.2) [53]. Furthermore, OA standard (Diagnostic Chemicals Limited, Charlottetown, CA USA) was injected in the LC-HRMS method and was detected at 14.62 min.

Results

Filamentous fungi were isolated from *Prorocentrum lima* fresh biomass

To evaluate the presence of fungi in *P. lima* microbiome, biomass from PL4V strain was used as inoculum for cultivation in fungal adapted medium [54], in presence of antibiotics. Cultivation was achieved either in liquid or solid conditions (Table S1). After few weeks, fungal colonies were observed and isolated (Figure S1). All strains were identified based on fungal morphology resulting in the isolation of two *Penicillium* sp. strains (MMS1593, MMS1596) and three *Aspergillus* sp. strains (MMS1589, MMS1591, MMS1594).

Among isolated strains, the *Aspergillus* sp. strains MMS1589 was selected for further co-culture study, due to its unusual morphology (cleistotheca and ascospores simultaneous observation) (Figure S2) (only one was selected due to expected difficulties in such experimental design, as such fungal-dinoflagellates co-culture were not previously reported). Congo red coloration revealed hyaline septate mycelium and hyaline conidiophores, bearing uniseriate totally covered vesicles. Conidia were round shaped. Interestingly, globose cleistotheca was also visible. Asci and ascospores were present next to cleistotheca. On the basis of the genes encoding fungal small subunit ribosomal RNA (SSU rRNA), Internal Transcribed Spacer 1 (ITS) and β -tubulin of MMS1589, phylogenetic analyses were performed with all sequences in Genbank. The MMS1589 ITS partial gene has 99 % of identity with the *A. pseudoglaucus* strain FJAT-31014 SSU rRNA (partial sequence) and the MMS1589 β -tubulin partial gene has 99 % of identity with the *A. pseudoglaucus* benA gene for β -1 and β -2 tubulin (partial sequence). Finally, this strain was identified as *A. pseudoglaucus*, formerly known as *Eurotium repens* [52].

Specific microscale marine environment was designed for co-culture

The study of chemical interaction between *P. lima* PL4V and *A. pseudoglaucus* MMS1589, implied the challenging design of a functional micro-environment for microalgal / fungal co-culture since both species have specific requirements for their growth. On one hand, the benthic dinoflagellate *P. lima* growth requires a liquid media (typically f/2 medium) without stirring, without organic carbon source and with a light source for night-and-day cycle. In that case, it grew as a biofilm in the bottom of the culture flask. On the other hand, fungal growth requires either liquid or solid media with an organic carbon source. However, *A. pseudoglaucus* growth in liquid medium without stirring yielded fungal colonies floating over the liquid media. Thus, it was decided to culture the fungus on solid medium (containing a low level of glucose, 2 g/L) prior to addition of liquid medium in the culture flask and *P. lima* inoculation.

When both organisms were cultivated separately in the composite liquid/solid environment with light (12h/12h night-and-day cycle), normal growth was observed (Figure 1): (1) increase of the fungal colony size at the bottom of the flask (on the surface of the agar layer) and (2) formation of a *P. lima* biofilm as usually observed in usual conditions. During *P. lima* culture, bacterial development was observed leading to a cloudy culture medium (Figure 1), as confirmed by microscopic observation (most likely due to the sugar content of the underlying solid medium and the microalgae derived organic matter). Thus, to study microalgal growth with a very limited bacterial development, antibiotics were added in parallel [39].

During co-culture, both microorganisms grew normally (based on morphologic observation). However, *P. lima* tended to gather within the fungal mycelium, possibly using it as anchor to stick to the bottom of the flask. Again, bacterial development was observed in absence of antibiotics in the liquid medium.

Aspergillus pseudoglaucus MMS1589 impacts specialized metabolite production by *Prorocentrum lima* PL4V and its associated bacteria during co-culture

The co-culture experiment was achieved between *P. lima* PL4V and *A. pseudoglaucus* MMS1589 in the devised marine microscale environment in presence and absence of antibiotics [39] to be able to delineate the clear effect *A. pseudoglaucus* on *P. lima*. After 6 days, culture was stopped and agar (bottom part) and liquid supernatant were collected and extracted separately. The agar samples contained fungal mycelium and/or dinoflagellates in a biofilm and the liquid supernatant contained only *P. lima* when inoculated (fungal spores traces were observed). To obtain a clear view of metabolites present in both types of extracts, they were profiled by LC-HRMS [41]. For sake of clarity, single strain *P. lima* culture corresponds to the dinoflagellates with (xenic) or without (axenic) their associated bacteria depending on the presence of antibiotics.

To explore the LC-HRMS data, all peaks were automatically detected using dedicated open-source software [43]. All peaks detected in both LOP and solid phase were concatenated (per samples), generating the final data matrix. The resulting data matrix was subjected to unsupervised statistical analysis through principal component analysis (PCA) to assess the self-organized metabolome differences between *P. lima* PL4V and *A. pseudoglaucus* MMS1589 monocultures, and their corresponding co-culture (Figure 2A). The first two components of the resulting PCA model correspond to 22.1% (PC1) and 11.3% (PC2) of explained variance. This model clearly highlighted the presence of 3 clusters related to *P. lima*, *A. pseudoglaucus* and their co-culture. The first component (PC1) is related to the differences between the *P. lima* and *A. pseudoglaucus* monocultures, and the second component clearly differentiates between monocultures and co-cultures.

Figure 2. Unsupervised data analysis (n=5) of LC-HRMS data showing differences between *Prorocentrum lima*, *Aspergillus pseudoglaucus* and their co-culture samples, in absence or presence of antibiotics. A) Principal Component Analysis (log transformed data, pareto scaling); B) Mixing model using Projected Orthogonalised Encounter Monitoring (POChEMon) data analysis [50]. In the mixing model, the size of the co-culture samples reflects the amount of their unexplained variance within the mixing model (the larger the point is, the higher the unexplained variance is); C) Competition model using Projected Orthogonalised Encounter Monitoring (POChEMon) data analysis. The green zone corresponds to the projected co-culture domain within the competition model.

The PCA scores plot (Figure 2A) also highlighted a clustering of samples according to xenic and axenic status for each culture conditions. As signals related to antibiotics were removed from the data, the antibiotic clustering effect could mainly be linked to metabolome alterations due to the presence or absence of bacteria in the culture medium. In the presence of antibiotics, the chemical profiles of the *P. lima* monoculture corresponded only to compounds produced by the dinoflagellates, while without antibiotics, the chemical profiles reflect the metabolome of the microalgae with its associated bacteria. Greater differences between *P. lima* metabolomes under xenic and axenic conditions were observed in comparison to fungal ones under both conditions (Figure 2A), suggesting that the presence of antibiotics should alter more importantly the microalgal metabolite profiles than the fungal ones. Furthermore, an even greater impact of the xenic condition was observed within co-cultures. Also, the absence of antibiotics shifted co-cultured replicates toward the *P. lima* culture samples. This reflects the impact of *A. pseudoglaucus* on specialized metabolites produced by *P. lima* and its bacteria and the impact of microalgae (and associated bacteria) on *A. pseudoglaucus* metabolome.

Such results highlight the importance of bacteria in the chemical inertia of *P. lima* when confronted to *A. pseudoglaucus*. However, this unsupervised analysis did not point out the specific metabolome alteration induced by fungal presence as it encompasses every variation (microbiome or antibiotics derived variations). Thus, a more co-culture centric data mining strategy was used [31], called Projected Orthogonalised CHEmical Encounter MONitoring (POChEMon) [49]. Such an approach uses an unsupervised mixing model, obtained by statistically mixing data from chemical profiles corresponding to strains grown alone. This yielded an *in-silico* co-culture model in absence of co-culture-related biotic stress. Then, projection of co-culture samples within this mixing model allowed, as for the PCA, to compare chemical profiles from the co-culture to monocultures. This approach further confirmed results from the PCA, showing that absence of antibiotics makes co-culture chemical profiles resemble more *P. lima* ones (Figure 2B). Similarly, the presence of antibiotics makes the chemical profiles of the coculture resemble more to the fungal profiles. Additionally, the POChEMon approach provided a global comparison between all co-culture experiments using the residual

information, not represented in this mixing model (Figure 2B). The size of the co-culture samples when represented is directly related to unrepresented variance in the mixing model [49]. Here, this provided a strategy of unsupervised comparison between the two co-culture conditions (with or without antibiotics). Clearly, the chemical induction is stronger in presence of bacteria (larger residue) (Figure 2B). This further highlighted the importance of the microbiome in the co-culture specific chemical interaction, with more induction related to bacterial presence.

The co-culture induction pattern can be explored by interpreting the POChEMon competition model (Figure 2C) [49]. This representation corresponded to the unsupervised analysis through PCA of the co-culture sample residue when co-culture data were projected in the mixing model. Thus, the POChEMon approach allowed to identify co-culture related metabolome modifications in every sample. In Figure 2C, two co-culture patterns were observed and well separated corresponding to culture in presence or absence of antibiotics. This further confirmed the role of bacterial metabolites in co-culture, and also highlighted fungal-interaction specific compounds (residue in antibiotics treated samples). Interestingly the overall variability among co-culture replicates is lower in presence of antibiotics in comparison to their absence.

Identification of co-culture up-regulated compounds highlights

Prorocentrum lima toxin induction

As co-culture induction is a complex mechanism, different complementary supervised statistical approaches were used to highlight induced features [55]. Data were analysed univariately using Student test and multivariately using both orthogonal projection to latent structure discriminant analysis (OPLS-DA) and the POChEMon approach [49]. In all cases the top 20 most relevant features were selected with their retention time, m/z value and the statistical approach used. Finally, a set of 51 features were selected as over-produced (or *de novo* induced) during co-culture (Table 1 and Figure S3). Among those features, 14 were detected in the *P. lima* extracts, 6 in the *A. pseudoglossus* extracts

and 34 were not detected in monocultures (ND). In addition, 12 and 42 features were observed in agar and liquid media, respectively. Most of the features were found in antibiotic-free samples as the mixing model emphasized (Figure 2B), but some of them were specifically found when antibiotics were added, hence confirming a fungal-algal-interaction related chemical induction.

Based on high mass and spectral accuracy, the highlighted features were putatively annotated (Table 1 along with annotation accuracy based on Metabolomic standard initiative – MSI [61]). Considering the most significant features, none of the highlighted features was successfully annotated as *Prorocentrum* spp. metabolites. Furthermore, fungal metabolites annotation was only possible by extending the search to the clade *Aspergillus* section *restrictii*. As expected, only a very low number of features were identified.

Besides the previously highlighted features (Table 1), the two known toxins OA and DTX-1 usually produced by *P. lima* PL4V were also studied. Both toxins were found induced in co-culture with the Student test results in the LOP, with or without added antibiotics: OA (main ion detected at $m/z = 827.458$ $[M+Na]^+$, $t_R = 14.6$ min) and DTX-1 (main ion detected $m/z = 841.473$ $[M+Na]^+$, $t_R = 17.1$ min) with a fold change of 29.7 (p -value < 0.05) and 5 (p -value < 0.05), compared to monoculture conditions, respectively (they were ranked lower than 20). Okadaic acid was further confirmed using standard. No significant variations were observed in the solid phase profiles.

Table 1. Co-culture associated features using univariate and multivariate data analyses using PocheRmon package. Features highlighted using Ttest method correspond to top 20 ones with fold changes higher than 2 with Student-test *p*-value below 5%. The OPLS-DA method select up-regulated features with VIP (variable importance in the projection) value higher than 2. Finally, the POChEMon method highlights features with SSrankProduct higher than 0.995. The drastic variable selection yielded low number of features highlighted by different analysis.

Detected (ionisation mode)	<i>m/z</i>	<i>t_R</i> (min)	Matrix	Antibiotics	Statistical method	Presence monoculture (MC)	in	Molecular (error, ppm)	formula	Annotations with reported biological sources and adducts
243.7550 (-)	1.0	A	Both		OPLS-DA	Both MC				
254.9880 (+)	1.1	A	Both		OPLS-DA	Both MC				
433.0163 (-)	9.0	Both	No		Ttest, POChEMon	ND				
435.0291 (+)	9.0	L	No		Ttest	ND				
433.0172 (-)	8.4	L	No		Ttest (∞), POChEMon	ND				
273.0787 (-)	11.0	Both	No		Ttest (∞)	ND				
564.3306 (+)	11.0	L	No		OPLS-DA	ND				
311.2244 (-)	12.1	A	Yes		POChEMon	MC Fungi		C ₁₈ H ₃₂ O ₄ (5 ppm)		Sporogenic Psi Factor Cα [56] or QMZ37-I [57] (<i>Aspergillus</i>) [M-H] ⁻
418.0254 (-)	13.0	L	No		POChEMon	ND		C ₁₅ H ₁₇ N ₃ O ₅ S ₂ (-12 ppm)		Glionitrin B [58] (<i>Aspergillus</i>) [M+Cl] ⁻
329.1390 (-)	14.0	L	Yes		Ttest	ND		C ₁₉ H ₂₂ O ₅ (-2 ppm)		Arahypin 2 or Arahypin 3 (<i>Aspergillus</i>) [M-H] ⁻
415.0061 (-)	13.2	L	No		Ttest (∞), POChEMon	ND				
544.3293 (+)	11.5	L	Yes		Ttest (∞)	MC Micro-algae				
442.2601 (+)	14.2	L	Yes		Ttest	MC Micro-algae				
606.3650 (-)	15.2	L	Yes		Ttest	ND				
522.5520 (+)	16.1	L	No		Ttest	ND				
634.3363 (-)	16.1	L	No		OPLS-DA	ND				
506.3265 (-)	16.1	L	No		POChEMon	ND				
509.2905 (-)	17.0	A	No		OPLS-DA	MC Micro-algae				
544.4092 (-)	17.0	L	No		OPLS-DA	ND				
581.3034 (-)	16.3	L	No		Ttest	Both MC				
486.2516 (-)	16.3	L	No		OPLS-DA	ND		C ₂₈ H ₃₇ NO ₄ (21 ppm)		Dodecahydro-11a-(1 <i>H</i> -indol-3-yl)-1,3a,4,9-tetramethyl-8,11c-methano-11 <i>cH</i> -benzo[<i>ef</i>]furo[3,2- <i>c</i>][1]benzoxepin-3,8a(8 <i>H</i>)-diol [59] (<i>Aspergillus</i>) [M+Cl] ⁻
580.3630 (-)	17.2	L	No		Ttest	ND				
563.3460 (-)	17.2	A	No		OPLS-DA	MC Micro-algae				
514.9216 (-)	12.8	L	No		Ttest, POChEMon	ND				
417.0203 (-)	13.7	L	No		Ttest (∞), POChEMon	MC Fungi		C ₁₉ H ₁₄ N ₂ O ₅ S (-27 ppm)		4-Methylether-4'-sulfate-xanthocillin X (Antibiotic BU 4704) [60] (<i>Aspergillus</i>) [M+Cl] ⁻
419.2678 (-)	16.4	A	No		OPLS-DA	MC Micro-algae				
508.3407 (+)	16.4	L	No		OPLS-DA	ND				
509.2910 (-)	17.3	L	No		OPLS-DA	MC Micro-algae				
485.0060 (-)	12.9	L	No		Ttest (∞)	ND				
516.9162 (-)	12.9	L	No		Ttest (∞), POChEMon	ND				
417.0222 (-)	12.9	L	No		Ttest, POChEMon	MC Fungi		C ₁₉ H ₁₄ N ₂ O ₅ S (-23 ppm)		4-Methylether-4'-sulfate-xanthocillin X [60] (<i>Aspergillus</i>) [M+Cl] ⁻
419.0378 (+)	12.9	L	No		Ttest, POChEMon	ND				
464.2782 (-)	14.8	L	No		OPLS-DA	ND				
507.3061 (+)	18.4	L	No		OPLS-DA	ND				
566.3484 (-)	15.8	L	No		POChEMon	MC Micro-algae				
509.2915 (-)	17.6	L	No		OPLS-DA	ND				
542.9179 (+)	15.9	Both	No		Ttest (∞), POChEMon	ND				
542.9179 (-)	15.9	L	No		POChEMon	ND				
523.3067 (-)	18.6	A	No		OPLS-DA	MC Micro-algae				
444.4044 (+)	22.2	L	No		Ttest	ND				
509.9216 (-)	16.9	L	No		POChEMon	ND				
509.7083 (-)	16.9	L	No		POChEMon	ND				
509.2918 (-)	18.7	A	No		OPLS-DA	ND				
542.4092 (+)	18.8	L	No		OPLS-DA	ND				
655.5283 (+)	27.2	L	No		OPLS-DA	MC Micro-algae				
545.3470 (-)	24.8	A	Both		Ttest	ND				
802.6100 (+)	26.7	L	Yes		POChEMon	MC Micro-algae				
584.4907 (+)	26.8	L	No		OPLS-DA	ND				
802.6112 (+)	26.9	L	Yes		POChEMon	MC Micro-algae				
566.4814 (-)	27.8	L	No		OPLS-DA	ND				
568.4942 (+)	27.8	L	No		OPLS-DA	ND				

A: detected in Agar; L: detected in Liquid medium; MC: monoculture; ND: not detected in any monoculture replicates; ∞: *de novo* induced features according to Ttest.

Unprecedented physical contact was observed between

Aspergillus pseudoglaucus and *Prorocentrum lima*

Visual, binocular and microscopic inspection of the co-culture samples revealed aggregation of *P. lima* within and around *A. pseudoglaucus* colonies (Figure 3AB). Only a limited number of free *P. lima* cells in biofilm stuck directly to the agar layer while most of biofilm is found directly in contact with *A. pseudoglaucus*. (Figure 3A). To further explore interaction morphology between *P. lima* PL4V and *A. pseudoglaucus* MMS1589, co-culture samples were explored using microscope. Surprisingly, an unusual physical interaction was observed between those organisms. Figure 3CD shows that the mycelium of *A. pseudoglaucus* seems to connect to the dinoflagellates at the flagella location. The red coloration of the connecting organelle indicated its fungal origin as Congo Red dye is known to highlight in red fungal membranes.

Discussion

Fungal role in the marine environment [62] is largely underexplored [63, 64]. However, fungal presence has been repeatedly observed, sometimes with rather large fungal biomasses [65–67]. The present study focuses on the interactions between the toxic dinoflagellate *P. lima* and an associated fungal strain. In fact, various fungal strains from the *Penicillium* and *Aspergillus* genera were isolated from the *P. lima* PL4V strain. The *A. pseudoglaucus* MMS1589 strain was selected to further explore chemical mediation between microalgae and fungi. Thus, a simple marine microcosm, compatible with fungal and benthic dinoflagellate life style, was devised to study such interaction. Specialized metabolites production during co-culture was studied by LC-HRMS metabolomics.

The differences of chemical alteration between axenic or xenic *P. lima* conditions clearly emphasize the effect of associated bacteria in the fungal/microalgal interaction (Figure 2). Interactions between *P. lima* and bacteria were shown to alter algal metabolites [68], and it is thus expected that in the present study the bacterial presence also significantly impacted the metabolite profile. Finally, this

showed that, in addition to dinoflagellate signals, the microbiome metabolism is also altered. Up to now, chemical induction was mostly reported in fungal-fungal, bacterial-bacterial and fungal-bacterial co-cultures [34].

Figure 3. Observation of Fungal-microalgal interaction. AB) binocular observation of co-culture with aggregated cleistotheca (pale and round in A, brown in B) and asexual sporogenic spores (dark not round). Microalgae are oval and surround fungal elements. CDE) microscopic observation of co-culture with Congo Red dye (x400). The fungal mycelium is indicated (arrows) at proximity of the apex of the microalgae where the physical interactions occur.

The chemical signal specifically related to *P. lima* and *A. pseudoglaucus* were tentatively identified. Some were reported to possess antimicrobial (asperchalsin E), antiviral (didemethylasteriquinone D, xanthocillin X) and anti-inflammatory activities (aspochalasin U). Furthermore, 5S,8R-Dihydroxy-9Z,12Z-octadecadienoic acid ($m/z = 311.2244$ at $t_R = 12.11$ min) was previously reported as a sporogenic factor of *Aspergillus* spp., which could reflect a fungal biotic stress adaptation behaviour. Unfortunately, among those highlighted features only a very limited number of compounds was putatively annotated. The reason may be that (1) frequently, microbial co-culture induces previously unreported compounds [34, 42] and that (2) metabolomes of toxic dinoflagellates are still largely unexplored (except toxins) [69]. Thus, a large effort in studying the metabolome of such organisms will be of great value to better understand their metabolism. Still, two induced compounds were unequivocally identified, OA and DTX-1, produced by *P. lima* PL4V. This is particularly interesting, as these toxins are largely related to *P. lima* toxicity by seafood contamination [70, 71]. In this experiment, signals from *A. pseudoglaucus* MMS1589 induces the over-production of such toxins. Similar results were reported with the domoic acid producer *Pseudo-nitzschia* where variation in bacterial community alters toxin levels [72, 73]. So far, there is very few evidence of toxin production alteration by associated microbiome [25, 74]. However, in this context, fungi should also be considered to play a significant role in such production regulation. Interestingly, OA and its derivatives (DTX-1 and 7-deoxyokadaic acid) are known to inhibit *Aspergillus niger* growth in a paper-disk assay [75]. Thus, OA and DTX-1 over-production may reflect that *P. lima* is tentatively controlling fungal growth; even though no evidence of fungal toxicity was observed in the presented study. Such inhibition was already observed during *P. lima* interaction with other microalgae [76]. Thus, *P. lima* toxin seems to be part of its global defence mechanism.

During the *P. lima* and *A. pseudoglaucus* interaction, aggregation was also observed (Figure 3AB). Similar aggregation followed by flocculation was previously related in biotechnology-focused fungal microalgal co-culture [77]. Such aggregation reflects the epiphytic nature of *P. lima* which needs to attach to objects in order to form biofilm [70]. This may also suggest nutrient exchange between both

organisms which could be ease by their very close proximity. However, this needs more in-depth study to be proven in the present interaction between *A. pseudoglaucus* MMS1589 and *P. lima* PL4V. Such idea was recently explored in the synthetic association between the microalgae *Chlamydomonas reinhardtii* and the yeast *Saccharomyces cerevisiae* [78]. In this specific association, the two species were forced, under specific conditions, to mutually support each other's growth. In a very recent study, during the tripartite interaction between *C. reinhardtii*, *Aspergillus nidulans* and *Streptomyces iranensis*, the fungal species was demonstrated to physically defend the microalgae from bacterial algicide compound azalomycin F by forming a protective lipid coat as observed by microscopy [79].

Besides aggregation and very unexpectedly, the microscopic observation of the interaction between *P. lima* and *A. pseudoglaucus* showed a very unusual physical contact where the fungal mycelium seems to be connected to the dinoflagellate at the flagella location (Figure 3). Such synthetic structured community formation is rather unusual in the literature and one may consider such contact as a very early stage of lichen formation [80, 81]. Lichens are classified as a microalgal-fungal symbiosis harbouring various other species, to form complex ecosystems [82]. *Aspergillus pseudoglaucus* was not previously mentioned as a lichen forming fungus [83]; however, some *Aspergillus* sp. were reported as endolichenic species having antimicrobial properties [84–86].

To conclude, this study raised many questions about microalgal/fungal interactions in the marine environment, especially in the context of toxic microalgae. Further studies are still needed to identify many of the metabolites induced either by *A. pseudoglaucus* MMS1589 or *P. lima* PL4V. It also would be interesting to further explore the mechanisms of such interactions and thus understand the physical contact observed and the OA induction reported here.

Acknowledgements

The authors would like to acknowledge the technical assistance of Véronique Séchet and Florent Malo for maintaining and providing *P. lima* cultures. The authors thank the ThalassOMICS Metabolomics

Facility – Plateforme Corsaire, Biogenouest, Nantes – for LC-HRMS analysis and metabolomics expertise. The authors would also like to acknowledge funding received from Ifremer and the regional research federation IUML for funding received for the Master studies of Alizé Bagot and Maud Chaigne.

References

1. Zhang S, Merino N, Okamoto A, Gedalanga P. Interkingdom microbial consortia mechanisms to guide biotechnological applications. *Microb Biotechnol* 2018; **11**: 833–847.
2. Puglisi MP, Sneed JM, Sharp KH, Ritson-Williams R, Paul VJ. Marine chemical ecology in benthic environments. *Nat Prod Rep* 2014; **31**: 1510–1553.
3. Brown ER, Cepeda MR, Mascuch SJ, Poulson-Ellestad KL, Kubanek J. Chemical ecology of the marine plankton. *Nat Prod Rep* 2019; **36**: 1093–1116.
4. Stewart I, Seawright AA, Shaw GR. Chapter 28: Cyanobacterial poisoning in livestock, wild mammals and birds – an overview. *Cyanobacterial Harmful Algal Blooms: State of the Science and Research Needs*, Hudnell H.K. (eds). Springer, New York, NY., p 25.
5. Grattan LM, Holobaugh S, Morris JG. Harmful algal blooms and public health. *Harmful Algae* 2016; **57**: 2–8.
6. Wang S, Chen J, Li Z, Wang Y, Fu B, Han X, et al. Cultivation of the benthic microalga *Prorocentrum lima* for the production of diarrhetic shellfish poisoning toxins in a vertical flat photobioreactor. *Bioresour Technol* 2015; **179**: 243–248.
7. Lee T, Fong F, Ho K-C, Lee F. The mechanism of diarrhetic shellfish poisoning toxin production in *Prorocentrum* spp.: physiological and molecular perspectives. *Toxins* 2016; **8**: 272.
8. Van Wagoner RM, Satake M, Wright JLC. Polyketide biosynthesis in dinoflagellates: what makes it different? *Nat Prod Rep* 2014; **31**: 1101.
9. Nishimura T, Uchida H, Noguchi R, Oikawa H, Suzuki T, Funaki H, et al. Abundance of the benthic dinoflagellate *Prorocentrum* and the diversity, distribution, and diarrhetic shellfish toxin

- p>production of
- Prorocentrum lima*
- complex and
- P. caipirignum*
- in Japan.
- Harmful Algae*
- 2020;
- 96**
- : 101687.
10. Hay ME. Marine chemical ecology: chemical signals and cues structure marine populations, communities, and ecosystems. *Annu Rev Mar Sci* 2009; **1**: 193–212.
 11. Gu S, Xiao S-W, Zheng J-W, Li H-Y, Liu J-S, Yang W-D. ABC transporters in *Prorocentrum lima* and their expression under different environmental conditions including okadaic acid production. *Mar Drugs* 2019; **17**: 259.
 12. Shaw BA, Andersen RJ, Harrison PJ. Feeding deterrent and toxicity effects of apo-fucoxanthinoids and phycotoxins on a marine copepod (*Tigriopus californicus*). *Mar Biol* 1997; **128**: 273–280.
 13. Selander E, Thor P, Toth G, Pavia H. Copepods induce paralytic shellfish toxin production in marine dinoflagellates. *Proc R Soc B* 2006; **273**: 1673–1680.
 14. Wang R, Wu J, Zhou S, Cao R, Chan LL. A preliminary study on the allelopathy and toxicity of the dinoflagellate *Karlodinium veneticum*. *Mar Pollut Bull* 2020; **158**: 111400.
 15. Sugg LM, VanDolah FM. No evidence for an allelopathic role of okadaic acid among ciguatera-associated dinoflagellates. *J Phycol* 1999; **35**: 93–103.
 16. Edwards BR, Bidle KD, Van Mooy BAS. Dose-dependent regulation of microbial activity on sinking particles by polyunsaturated aldehydes: Implications for the carbon cycle. *Proc Natl Acad Sci USA* 2015; **112**: 5909–5914.
 17. Johnson WM, Kido Soule MC, Kujawinski EB. Evidence for quorum sensing and differential metabolite production by a marine bacterium in response to DMSP. *ISME J* 2016; **10**: 2304–2316.
 18. Harvey EL, Deering RW, Rowley DC, El Gamal A, Schorn M, Moore BS, et al. A bacterial quorum-sensing precursor induces mortality in the marine coccolithophore, *Emiliana huxleyi*. *Front Microbiol* 2016; **7**.
 19. Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. Algae acquire vitamin B12 through a symbiotic relationship with bacteria. *Nature* 2005; **438**: 90–93.

20. Wagner-Döbler I, Ballhausen B, Berger M, Brinkhoff T, Buchholz I, Bunk B, et al. The complete genome sequence of the algal symbiont *Dinoroseobacter shibae*: a hitchhiker's guide to life in the sea. *ISME J* 2010; **4**: 61–77.
21. Segev E, Wyche TP, Kim KH, Petersen J, Ellebrandt C, Vlamakis H, et al. Dynamic metabolic exchange governs a marine algal-bacterial interaction. *Elife* 2016; **5**: e17473.
22. Green DH, Llewellyn LE, Negri AP, Blackburn SI, Bolch CJS. Phylogenetic and functional diversity of the cultivable bacterial community associated with the paralytic shellfish poisoning dinoflagellate *Gymnodinium catenatum*. *FEMS Microbiol Ecol* 2004; **47**: 345–357.
23. Green DH, Hart M, Blackburn SI, Bolch CJS. Bacterial diversity of *Gymnodinium catenatum* and its relationship to dinoflagellate toxicity. *Aquat Microb Ecol* 2010; **61**: 73–87.
24. Prol MJ, Guisande C, Barreiro A, Míguez B, de la Iglesia P, Villar A, et al. Evaluation of the production of paralytic shellfish poisoning toxins by extracellular bacteria isolated from the toxic dinoflagellate *Alexandrium minutum*. *Can J Microbiol* 2009; **55**: 943–954.
25. Uribe P, Espejo RT. Effect of associated bacteria on the growth and toxicity of *Alexandrium catenella*. *Appl Environ Microbiol* 2003; **69**: 659–662.
26. Ho A, Hsieh D, Qian P. Variations in paralytic shellfish toxin and homolog production in two strains of *Alexandrium tamarense* after antibiotic treatments. *Aquat Microb Ecol* 2006; **42**: 41–53.
27. Scholz B, Küpper F, Vyverman W, Ólafsson H, Karsten U. Chytridiomycosis of marine diatoms—The role of stress physiology and resistance in parasite-host recognition and accumulation of defense molecules. *Mar Drugs* 2017; **15**: 26.
28. Pohnert G. Wound-activated chemical defense in unicellular planktonic algae. *Angew Chem Int Ed* 2000; **39**: 4352–4354.
29. Vallet M, Baumeister TUH, Kaftan F, Grabe V, Buaya A, Thines M, et al. The oomycete *Lagenisma coscinodisci* hijacks host alkaloid synthesis during infection of a marine diatom. *Nat Commun* 2019; **10**: 4938.

30. Wrede D, Taha M, Miranda AF, Kadali K, Stevenson T, Ball AS, et al. Co-cultivation of fungal and microalgal cells as an efficient system for harvesting microalgal cells, lipid production and wastewater treatment. *PLOS ONE* 2014; **9**: e113497.
31. Bertrand S, Bohni N, Schnee S, Schumpp O, Gindro K, Wolfender J-L. Metabolite induction via microorganism co-culture: A potential way to enhance chemical diversity for drug discovery. *Biotechnol Adv* 2014; **32**: 1180–1204.
32. Khalil ZG, Cruz-Morales P, Licona-Cassani C, Marcellin E, Capon RJ. Inter-kingdom beach warfare: microbial chemical communication activates natural chemical defences. *ISME J* 2019; **13**: 147–158.
33. Wolfender J-L, Marti G, Thomas A, Bertrand S. Current approaches and challenges for the metabolite profiling of complex natural extracts. *J Chromatogr A* 2015; **1382**: 136–164.
34. Arora D, Gupta P, Jaglan S, Roullier C, Grovel O, Bertrand S. Expanding the chemical diversity through microorganisms co-culture: Current status and outlook. *Biotechnol Adv* 2020; **40**: 107521.
35. Schroeckh V, Scherlach K, Nutzmann H-W, Shelest E, Schmidt-Heck W, Schuemann J, et al. Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proc Natl Acad Sci USA* 2009; **106**: 14558–14563.
36. Lassus P, Chomérat N, Hess P, Nézan E. Toxic and harmful microalgae of the world ocean. 2016. International Society for the Study of Harmful Algae, Copenhagen N, Denmark.
37. Bravo I, Fernández ML, Ramilo I, Martínez A. Toxin composition of the toxic dinoflagellate *Prorocentrum lima* isolated from different locations along the Galician coast (NW Spain). *Toxicon* 2001; **39**: 1537–1545.
38. Guillard RRL, Ryther JH. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Can J Microbiol* 1962; **8**: 229–239.

39. Cho J-Y, Choi J-S, Kong I-S, Park S-I, Kerr RG. A procedure for axenic isolation of the marine microalga *Isochrysis galbana* from heavily contaminated mass cultures. *J Appl Phycol* 2002; **14**: 385–390.
40. Rédou V, Navarri M, Meslet-Cladière L, Barbier G, Burgaud G. Species richness and adaptation of marine fungi from deep-subseafloor sediments. *Appl Environ Microbiol* 2015; **81**: 3571–3583.
41. Roullier C, Bertrand S, Blanchet E, Peigné M, Robiou du Pont T, Guitton Y, et al. Time dependency of chemodiversity and biosynthetic pathways: an LC-MS metabolomic study of marine-sourced *Penicillium*. *Mar Drugs* 2016; **14**: 103.
42. Bertrand S, Schumpp O, Bohni N, Bujard A, Azzollini A, Monod M, et al. Detection of metabolite induction in fungal co-cultures on solid media by high-throughput differential ultra-high pressure liquid chromatography–time-of-flight mass spectrometry fingerprinting. *J Chromatogr A* 2013; **1292**: 219–228.
43. Pluskal T, Castillo S, Villar-Briones A, Orešič M. MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics* 2010; **11**: 395.
44. Myers OD, Sumner SJ, Li S, Barnes S, Du X. One step forward for reducing false positive and false negative compound identifications from mass spectrometry metabolomics data: new algorithms for constructing extracted ion chromatograms and detecting chromatographic peaks. *Anal Chem* 2017; **89**: 8696–8703.
45. Djoumbou-Feunang Y, Fiamoncini J, Gil-de-la-Fuente A, Greiner R, Manach C, Wishart DS. BioTransformer: a comprehensive computational tool for small molecule metabolism prediction and metabolite identification. *J Cheminformatics* 2019; **11**: 2.
46. Wang J, Bertrand S. Projected Orthogonalized CHEmical Encounter MONitoring) for mixed metabolome analysis (R function).
47. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing.

48. Thévenot EA, Roux A, Xu Y, Ezan E, Junot C. Analysis of the human adult urinary metabolome variations with age, body mass index, and gender by implementing a comprehensive workflow for univariate and OPLS statistical analyses. *J Proteome Res* 2015; **14**: 3322–3335.
49. Jansen JJ, Blanchet L, Buydens LMC, Bertrand S, Wolfender J-L. Projected Orthogonalized CHemical Encounter MONitoring (POCHEMON) for microbial interactions in co-culture. *Metabolomics* 2015; **11**: 908–919.
50. Wolfender J-L, Nuzillard J-M, van der Hooft JJJ, Renault J-H, Bertrand S. Accelerating metabolite identification in natural product research: toward an ideal combination of Liquid Chromatography–High-Resolution tandem Mass Spectrometry and NMR profiling, *in Silico* databases, and chemometrics. *Anal Chem* 2019; **91**: 704–742.
51. Nielsen KF, Månsson M, Rank C, Frisvad JC, Larsen TO. Dereplication of microbial natural products by LC-DAD-TOFMS. *J Nat Prod* 2011; **74**: 2338–2348.
52. Chen AJ, Hubka V, Frisvad JC, Visagie CM, Houbraken J, Meijer M, et al. Polyphasic taxonomy of *Aspergillus* section *Aspergillus* (formerly *Eurotium*), and its occurrence in indoor environments and food. *Stud Mycol* 2017; **88**: 37–135.
53. Chen Y, de Bruyn Kops C, Kirchmair J. Data resources for the computer-guided discovery of bioactive natural products. *J Chem Inf Model* 2017; **57**: 2099–2111.
54. Sallenave-Namont C, Pouchus YF. Toxigenic saprophytic fungi in marine shellfish farming areas. *Mycopathologia* 2000; **149**: 21–25.
55. Saccenti E, Hoefsloot HCJ, Smilde AK, Westerhuis JA, Hendriks MMWB. Reflections on univariate and multivariate analysis of metabolomics data. *Metabolomics* 2014; **10**: 361–374.
56. Mazur P, Nakanishi K, El-Zayat AAE, Champe SP. Structure and synthesis of sporogenic psi factors from *Aspergillus nidulans*. *J Chem Soc, Chem Commun* 1991; 1486.
57. Qiao M-F, Ji N-Y, Miao F-P, Yin X-L. Steroids and an oxylipin from an algicolous isolate of *Aspergillus flavus*. *Magn Reson Chem* 2011; **49**: 366–369.

58. Park HB, Kim Y-J, Park J-S, Yang HO, Lee KR, Kwon HC. Glionitrin B, a cancer invasion inhibitory diketopiperazine produced by microbial coculture. *J Nat Prod* 2011; **74**: 2309–2312.
59. Forseth RR, Amaike S, Schwenk D, Affeldt KJ, Hoffmeister D, Schroeder FC, et al. Homologous NRPS-like gene clusters mediate redundant small-molecule biosynthesis in *Aspergillus flavus*. *Angew Chem Int Ed* 2013; **52**: 1590–1594.
60. Takatsuki A, Tamura G, Arima K. New antiviral antibiotics; xanthocillin X mono- and dimethylether, and methoxy-xanthocillin X dimethylether. II Biological aspect of antiviral activity. *J Antibiot* 1968; 5.
61. Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, et al. Proposed minimum reporting standards for chemical analysis: Chemical Analysis Working Group (CAWG) Metabolomics Standards Initiative (MSI). *Metabolomics* 2007; **3**: 211–221.
62. Gleason FH, Gadd GM, Pitt JI, Larkum AWD. The roles of endolithic fungi in bioerosion and disease in marine ecosystems. I. General concepts. *Mycology* 2017; **8**: 205–215.
63. Worden AZ, Follows MJ, Giovannoni SJ, Wilken S, Zimmerman AE, Keeling PJ. Rethinking the marine carbon cycle: factoring in the multifarious lifestyles of microbes. *Science* 2015; **347**: 1257594–1257594.
64. Amend A, Burgaud G, Cunliffe M, Edgcomb VP, Ettinger CL, Gutiérrez MH, et al. Fungi in the marine environment: open questions and unsolved problems. *mBio* 2019; **10**: e01189-18, /mbio/10/2/mBio.01189-18.atom.
65. Hallegraeff G, Coman F, Davies C, Hayashi A, McLeod D, Slotwinski A, et al. Australian dust storm associated with extensive *Aspergillus sydowii* fungal “bloom” in coastal waters. *Appl Environ Microbiol* 2014; **80**: 3315–3320.
66. Li X, Xia Z, Tang J, Wu J, Tong J, Li M, et al. Identification and biological evaluation of secondary metabolites from marine derived fungi-*Aspergillus* sp. SCSIO3, cultivated in the presence of epigenetic modifying agents. *Molecules* 2017; **22**: 1302.

67. Wang Y, Sen B, He Y, Xie N, Wang G. Spatiotemporal distribution and assemblages of planktonic fungi in the coastal waters of the Bohai sea. *Front Microbiol* 2018; **9**: 584.
68. Tarazona-Janampa UI, Cembella AD, Pelayo-Zárate MC, Pajares S, Márquez-Valdelamar LM, Okolodkov YB, et al. Associated bacteria and their effects on growth and toxigenicity of the dinoflagellate *Prorocentrum lima* species complex from epibenthic substrates along mexican coasts. *Front Mar Sci* 2020; **7**: 569.
69. Zendong Z, Bertrand S, Herrenknecht C, Abadie E, Jauzein C, Lemée R, et al. Passive sampling and high resolution mass spectrometry for chemical profiling of French coastal areas with a focus on marine biotoxins. *Environ Sci Technol* 2016; **50**: 8522–8529.
70. Lawrence JE, Grant J, Quilliam MA, Bauder AG, Cembella AD. Colonization and growth of the toxic dinoflagellate *Prorocentrum lima* and associated fouling macroalgae on mussels in suspended culture. *Mar Ecol Prog Ser* 2021; **201**: 147–154.
71. Bauder AG, Cembella AD, Bricelj VM, Quilliam MA. Uptake and fate of diarrhetic shellfish poisoning toxins from the dinoflagellate *Prorocentrum lima* in the bay scallop *Argopecten irradians*. *Mar Ecol Prog Ser* 2021; **213**: 39–52.
72. Kobayashi K, Takata Y, Kodama M. Direct contact between *Pseudo-nitzschia* multiseres and bacteria is necessary for the diatom to produce a high level of domoic acid. *Fish Sci* 2009; **75**: 771–776.
73. Sison-Mangus MP, Jiang S, Tran KN, Kudela RM. Host-specific adaptation governs the interaction of the marine diatom, *Pseudo-nitzschia* and their microbiota. *ISME J* 2014; **8**: 63–76.
74. Gallacher S, Flynn KJ, Franco JM, Brueggemann EE, Hines HB. Evidence for production of paralytic shellfish toxins by bacteria associated with *Alexandrium* spp. (Dinophyta) in culture. *Appl Environ Microbiol* 1997; **63**: 239–245.
75. Nagai H, Satake M, Yasumoto T. Antimicrobial activities of polyether compounds of dinoflagellate origins. *J Appl Phycol* 1990; **2**: 305–308.

76. Windust AJ, Wright JLC, McLachlan JL. The effects of the diarrhetic shellfish poisoning toxins, okadaic acid and dinophysistoxin-1, on the growth of microalgae. *Mar Biol* 1996; **126**: 19–25.
77. Leng L, Li W, Chen J, Leng S, Chen J, Wei L, et al. Co-culture of fungi-microalgae consortium for wastewater treatment: A review. *Bioresour Technol* 2021; **330**: 125008.
78. Hom EFY, Murray AW. Niche engineering demonstrates a latent capacity for fungal-algal mutualism. *Science* 2014; **345**: 94–98.
79. Krespach MKC, García-Altares M, Flak M, Hanno Schoeler, Scherlach K, Netzker T, et al. Lichen-like association of *Chlamydomonas reinhardtii* and *Aspergillus nidulans* protects algal cells from bacteria. *ISME J* 2020; **14**: 2794–2805.
80. Ahmadjian V, Jacobs JB, Russell LA. Scanning electron microscope study of early lichen synthesis. *Science* 1978; **200**: 1062–1064.
81. Trembley ML, Ringli C, Honegger R. Morphological and molecular analysis of early stages in the resynthesis of the lichen *Baeomyces rufus*. *Mycol Res* 2002; **106**: 768–776.
82. Hawksworth DL, Grube M. Lichens redefined as complex ecosystems. *New Phytol* 2020; **227**: 1281–1283.
83. Honegger R. Lichen-forming fungi and their photobionts. In: Deising HB (ed). *The Mycota*. 2009. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 307–333.
84. Padhi S, Masi M, Panda SK, Luyten W, Cimmino A, Tayung K, et al. Antimicrobial secondary metabolites of an endolichenic *Aspergillus niger* isolated from lichen thallus of *Parmotrema ravum*. *Nat Prod Res* 2020; **34**: 2573–2580.
85. Dou Y, Wang X, Jiang D, Wang H, Jiao Y, Lou H, et al. Metabolites from *Aspergillus versicolor*, an endolichenic fungus from the lichen *Lobaria retigera*. *Drug Discov Ther* 2014; **8**: 84–88.
86. Prateeksha, Bajpai R, Yusuf MA, Upreti DK, Gupta VK, Singh BN. Endolichenic fungus, *Aspergillus quandricinctus* of *Usnea longissima* inhibits quorum sensing and biofilm formation of *Pseudomonas aeruginosa* PAO1. *Microb Pathog* 2020; **140**: 103933.