A potent myeloid response is rapidly activated in the lungs of premature Rhesus macaques exposed to intra-uterine inflammation.

Courtney M. Jackson^{1,2*}, Martin Demmert^{1,3*}, Shibabrata Mukherjee¹, Travis Isaacs¹, Jerilyn Gray⁴, Paranthaman Sentharamaikkannan⁴, Pietro Presicce⁵, Kashish Chetal⁶, Nathan Salomonis⁶, Lisa A. Miller⁷, Alan H. Jobe⁴, Suhas G. Kallapur⁵, William J. Zacharias^{4,8}, Ian P. Lewkowich¹, Hitesh Deshmukh⁴, Claire A. Chougnet^{1#}

1. Division of Immunobiology, Cincinnati Children's Hospital Research Foundation, Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA
2. Immunology Graduate Program, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA
3. Department of Pediatrics, Institute for Systemic Inflammation Research, University of Lübeck, Lübeck, Germany
4. Division of Neonatology/Pulmonary Biology, The Perinatal Institute, Cincinnati Children's Hospital Research Foundation, Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA
5. Divisions of Neonatology and Developmental Biology, David Geffen School of Medicine at the University of California Los Angeles, Los Angeles, CA USA
6. Division of Biomedical Informatics, Cincinnati Children's Hospital Research Foundation, Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA
7. California National Primate Research Center, University of California Davis, Davis, CA, Department of Anatomy, Physiology, and Cell Biology, School of Veterinary Medicine, University of California Davis, Davis, CA USA
8. Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, University of Cincinnati, Ohio, USA

*Contributed equally.

#Corresponding author: Dr. Claire Chougnet, Division of Immunobiology, Cincinnati Children's Hospital Medical Center, MLC #7038, 3333 Burnet Avenue, Cincinnati, OH 45229-3039. E-mail address: Claire.Chougnet@cchmc.org, phone number: 513-636-8847, fax number: 513-636-5355

Abstract

Intrauterine inflammation/infection (IUI), which is present in up to 40% of premature births, leads to elevated levels of pro-inflammatory mediators and microbial products within the amniotic fluid, which come in close contact to fetal mucosae. Yet, knowledge on the fetal mucosal responses to IUI exposure remains limited. To address these questions, we used a non-human primate model of IUI, in which pregnant Rhesus macaques received intra-amniotic (IA) LPS, compared with IA saline.

We found that IA LPS exposure induced a robust and rapid inflammation of the fetal lung, but not the intestine. This inflammatory response was characterized by high levels of pro-inflammatory cytokines in the lung and the alveolar wash, and a potent myeloid cell response, dominated by neutrophils and monocytes/macrophages. scRNAseq analyses of fetal lungs showed that the infiltrating (neutrophils and inflammatory monocytes) and the resident (alveolar and interstitial macrophages) myeloid cells exhibited transcriptional profiles consistent with exposure to TLR ligands, as well as to cytokines, notably IL-1 and TNF α . However, blocking IL-1 signaling or TNF α , alone or simultaneously by administering inhibitors intra-amniotically and subcutaneously to the dam only partially blunted fetal lung inflammation.

Together, our novel data indicate that the fetal innate immune system can mount a rapid multi-factorial mucosal innate response to IUI, responding both to direct signaling by bacterial products and to indirect cytokine-mediated pathways of activation. These data thus provide more mechanistic insights into the association between IUI exposure and the post-natal lung morbidities of the premature infant.

Introduction

Intrauterine inflammation/infection (IUI) is present in 25-40% of premature births¹. Inflammation of the placental membranes at the fetal-maternal interface, which is called chorioamnionitis (chorio)², leads to elevated levels of pro-inflammatory mediators and microbial products within the amniotic fluid²⁻⁵. Due to *in utero* aspiration/swallowing of the amniotic fluid^{6,7}, the mucosae of the developing fetus come in close contact with these mediators, which triggers an inflammatory response. Although such responses likely contribute to the increased incidence of post-natal pulmonary and intestinal morbidities associated with IUI exposure⁸⁻¹⁰, knowledge about the cells and pathways involved in the fetal mucosal response remains very limited, due to the inability to collect tissue samples in preterm infants.

Fetal lung and intestinal inflammation have been described in IUI animal models (rodent, rabbits, pig, and sheep). Main features include high levels of pro-inflammatory cytokines and infiltration of immune cells, as well as damage to the lung and intestine¹¹⁻¹⁸. However, the ontogeny of the immune system or the placenta structure are quite different in these species from those in humans. In contrast, non-human primates (NHP) are quite similar to humans in most biological aspects. Furthermore, many reagents developed against human antigens are cross-reactive allowing for in-depth analysis of the NHP fetal immune system¹⁹⁻²¹. Importantly, we have previously shown that intra-amniotic (IA) injection of LPS in pregnant Rhesus macaques leads to a type of inflammation that closely phenocopies severe human chorio^{22,23}. Previous studies from ours and other groups have reported elevated levels of inflammatory cytokines in the lung of IUI-exposed rhesus macaque fetuses^{24,23}, but no in-depth analysis of mucosal immune

responses was performed in these studies. Our goal herein was thus to apply state-of-the-art technologies, such as single-cell RNAseq (scRNAseq), to obtain a comprehensive view of the mucosal immune responses occurring in response to IUI.

Furthermore, this model allows us to evaluate the role inflammatory cytokines may play in these responses. Among the cytokines elevated in the amniotic fluid and cord blood in chorio, IL-1 β and TNF α appear particularly important^{4,25}. Indeed, both cytokines are capable of inducing chorio, and neutrophil infiltration into the fetal lung of Rhesus macaques when administered IA to pregnant dams^{26,27}. In addition, IL-1 β is involved in many aspects of LPS-induced fetal inflammation^{22,23,27-30}. We thus aimed at better delineating their respective role, using the cross-reactive IL-1 receptor antagonist (IL-1RA) and a blocking anti-TNF Ab (Adalimumab) given either individually, or in combination.

Herein, we show that IA LPS leads to rapid and robust inflammation of the fetal lung, but not of the fetal intestine. Notably, our analyses identified two main components, i.e., the recruitment of innate myeloid cells into the fetal lung, and the activation of resident lung myeloid cells. Although signaling by IL-1 and TNF α were identified as major transcriptional signatures in responsive myeloid cells, their individual or combined blockade only partially blunted fetal lung inflammation. Together, our novel data thus indicate that the fetal innate immune system is capable of mounting a rapid multi-factorial mucosal response to IUI, responding both to direct signaling by bacterial products such as LPS, and to indirect cytokine-mediated pathways of activation.

Results

IA LPS induces robust inflammation in the fetal lung with a minimal response in the small intestine.

To model IUI, pregnant Rhesus macaques were given intra-amniotic (IA) injection of LPS or saline at approximately 80% gestation. Sixteen hours post IA injections, the fetuses were surgically removed, and tissues collected for analysis. All the animals included in the study are described in Table S1.

We first examined the histological changes in the fetal lung and small intestine. Compared to controls, lungs from 16hr IA LPS-exposed fetuses displayed signs of inflammation with cell infiltration into the lung interstitium and decreased alveolar space (Fig. 1A). Secondary alveoli septa formation also appeared diminished (Fig. 1A). In contrast, in the jejunum of exposed fetuses, the intestinal villi, lamina propria, or the underlying mucosal area exhibited no abnormalities (Fig. S1A). Because intestinal inflammation did not develop until 3 days after IUI in other models (pig¹⁸ and sheep^{12,17,31}), we analyzed jejunum sections taken 48 and 120hrs post IA LPS, and again, found only minimal changes compared to controls (Fig. S1A).

Along with histological evaluations, mRNA expression of inflammatory cytokines was measured in lung and jejunum at 16hrs post IA LPS. TNF α , IL-6, and IL-1 β mRNA levels were greatly increased in the lung (>100-fold) of IA LPS-exposed fetuses along with increased detection of these cytokines in the alveolar wash (AW) (Fig. 1B, C). Other pro-inflammatory cytokines such as IL-8 and CCL2 were also increased (see Tables S2 and S3). In contrast, IL-10, a known anti-inflammatory molecule was only minimally increased

in the AW of LPS-exposed infants (Table S2). In agreement with the absence of histological alterations in the jejunum, mRNA expression of inflammatory cytokines was not significantly increased in this compartment (Fig. S1B). Thus, IA LPS induces a rapid inflammatory response within the fetal lung but has minimal impact on the fetal small intestine. Our in-depth analyses of fetal mucosal responses thus focused on the lung.

IA LPS triggers a large infiltration of myeloid cells into the fetal lung.

We next evaluated by scRNAseq the global transcriptional response of the lung in response to IA LPS (animals and sequencing metrics are described in Table S4). These analyses revealed distinct immune and non-immune cell populations in both control and IA LPS-exposed animals (Fig. 2A, Table S5). As our focus was on immune responses, subsequent analyses were performed on a subset of the whole dataset, containing only the immune cell clusters. (Fig. 2B). Using established lineage marker genes (see Fig. 2C), we identified the robust accrual of neutrophilic and monocyte/macrophage populations in the lung of IA LPS-exposed fetuses, compared to control animals (Fig. 2B). As expected, they highly expressed the myeloid-associated CD88 gene (*C5AR1*, Fig. 2D). These findings were confirmed by flow cytometric analyses of lung cell suspensions (see gating strategy in Fig. S2), which showed a trend toward a higher proportion of CD45⁺ hematopoietic cells, and, within this population, a significant increase in the proportion of CD88⁺ cells (Fig. 2E).

Due to the rapid response within the lung and the type of challenge used, another myeloid population of interest were dendritic cells (DCs), as DC populations have been described

in the human fetal lung³². By flow cytometry, we readily identified myeloid dendritic cells (mDCs) and plasmacytoid DCs (pDCs) in the lung, but their frequencies were unchanged by IA LPS (Fig. S3). However, these DCs appeared activated in IA LPS-exposed fetuses, as they both exhibited significantly increased HLA-DR levels (Fig. S3A-B).

In contrast to the brisk myeloid response, the frequency of B cells or different T cell populations, including regulatory T cells, was not changed by IA LPS as seen by flow cytometry and scRNAseq analyses (Fig. S4A-D). Additionally, analyses of differentially expressed genes (DEGs) also uncovered relatively few changes (n=342) in these populations between IA LPS and control animals (Fig S4B, S4D, Table S6).

Taken together, our data show that IA LPS rapidly affected fetal innate immune cell populations in the lung, mainly altering myeloid cells (neutrophils, monocytes/macrophages, DCs), while having little effect on the major lymphoid cell subsets.

IA LPS leads to the recruitment of inflammatory monocytes into the fetal lung and activates the resident macrophages.

scRNAseq analyses of the lungs of control fetuses revealed the presence of 2 macrophage populations (Fig. 3A), which we identified as alveolar and interstitial macrophages (Fig. 3B and S5). In the lung of IA LPS-exposed fetuses, another population of monocytes/macrophages with characteristics of inflammatory monocytes was also present (Fig. 3A-B and Fig. S5).

Next, we identified DEGs in alveolar and interstitial macrophages of IA LPS exposed lungs compared to controls. The majority of these DEGs were more highly expressed in LPS than in controls (n=482/620 in alveolar macrophages and n=302/324 in interstitial macrophages; Table S7, S8). Gene ontology (GO) enrichment analysis of these upregulated genes revealed several biological processes, including activation of the TLR signaling pathway, and response to cytokines, notably to IL-1 and TNF (Fig. 3C, Tables S9 and S10 for full list of genes). Furthermore, genes associated with apoptotic signaling and cell migration were particularly enriched (Fig. 3C). The fewer down-regulated genes in the alveolar and interstitial macrophages of IA LPS lungs mainly included ribosomal protein genes (Tables S6 and S7).

There was no inflammatory monocyte cluster in the control lungs, precluding DEG analysis. However, single sample gene set enrichment analysis (ssGSEA) of this population in the IA LPS-exposed lungs revealed a similar profile as that found in the alveolar and interstitial macrophages from the same lungs. Indeed, there was an enrichment of genes related to activation of the TLR, IL-1 and TNF signaling, as well as cell migration and apoptosis (Fig. 3D, Table S11 for full list of genes expressed and enrichment scores). All together, these data thus show that exposure to IUI strongly affects fetal lung resident macrophages as well as rapidly inducing the afflux of activated, inflammatory monocytes into the fetal lung.

Recruited neutrophils into the fetal lung have a profile of activation suggesting both direct LPS and indirect cytokine-mediated pathways of activation.

Our data also showed a strong neutrophilic response in the fetal lung after IA LPS. scRNAseq analyses identified *TNFAIP6* (TNF Alpha Induced Protein 6) expression as one of the most specific markers for neutrophils (Fig. 4A and Fig. S6A). *S100A8* was also found to be highly expressed in most neutrophils, although it was also present in the monocyte/macrophage cluster (Fig. 4A). The cluster was further uniquely characterized by the expression of neutrophil specific markers *AZU1*, *ITGAM*, *NCF2*. By IHC, we confirmed the presence of multiple *TNFAIP6*⁺*S100A8*⁺ cells in the IA LPS fetal lungs, which were not present in control lungs (Fig. 4B). These cells had neutrophilic morphology, and they formed aggregates within alveolar spaces (Fig. 4B-C). We confirmed these data by staining with CD68 and HLA-DR, as we had found that CD68 is highly expressed in both fetal neutrophils and monocytes/macrophages, with only the latter expressing HLA-DR. A significantly increased number of CD68⁺HLA-DR⁻ cells, with a neutrophilic morphology, were found in the IA LPS-exposed lungs, also forming aggregates (Fig. S6B). Furthermore, we found a significant number of neutrophils in the AW of IA LPS exposed fetuses, whereas they were largely absent in the AW of control fetuses [median 0.0 v. 106.3x10⁶ cells/kg in control (n=18) and IA-LPS (n=13), respectively, p<0.0001, Mann-Whitney test].

We next analyzed the transcriptomic program of the neutrophils recruited in the fetal lungs by ssGSEA. These analyses revealed the enrichment of similar processes as those found in monocytes/macrophages, notably migration, response to TLR, IL-1 and TNF, and apoptosis (Fig.4D and Table S12 for full list of genes expressed and enrichment scores).

Interestingly, and in line with other studies^{33,34}, substantial transcriptional heterogeneity could be seen within these recruited neutrophils. However, no distinct sub-clusters were

revealed by UMAP projection (Fig. 5A), suggesting different levels of maturation and/or activation. We therefore applied pseudo-time analyses as implemented in Monocle3 to characterize this heterogeneity. Genes that most determined the trajectory were calculated with spatial autocorrelation analysis (Fig. S7). This analysis revealed that cytokine and cytokine-response genes were particularly enriched close to the beginning of the trajectory, whereas expression of other effector genes (e.g., *S100A8/9*, *CAMP*, *MMP8*) was higher towards more advanced pseudo-time (Fig. 5, S7 and S8). Thus, the recruited neutrophils into the fetal lung exhibit transcriptional programs revealing different stages of activation, compatible with both direct activation by LPS and indirect cytokine-mediated pathways.

IL-1 and TNF α signaling are partially driving fetal lung inflammation.

Knowing the role of IL-1 and TNF α in chorio pathogenesis^{22,23,26-30}, and because scRNAseq analyses revealed gene signatures consistent with IL-1 and TNF α signaling in both the monocyte/macrophage and neutrophil lung populations of IA LPS animals, we blocked IL-1 signaling with the IL-1RA and TNF α with the blocking anti-TNF Ab Adalimumab, administered both subcutaneous and IA to the pregnant dams. We also tested their combined effect (see Fig. S9 for experimental scheme). Both reagents had previously been shown to be efficacious in macaques^{22,23,35}.

Cytokine levels in the AW were only partially diminished with any of the treatments (Table S2). This modest effect was confirmed by the fact that cytokine mRNA expression in the lung was also minimally decreased in all the treatment groups (Table S3). LPS-induced

histological changes (cell infiltration and lung interstitium thickening) remained present in animals given IL-1RA or anti-TNF, although lung interstitium thickening was partially decreased in animals given both inhibitors (Fig. 6A).

Infiltration of myeloid cells into the lung was not blunted by any of the treatments, as shown by both scRNAseq and flow cytometry analyses (Fig. 6B-C). Neutrophil accumulation in the AW was also not blunted ($p>0.9$, Kruskal-Wallis tests). Similar monocyte/macrophage populations were present in the lung of treated animals as in IA LPS animals (Fig. 7A). However, the blockades partially suppressed some of the biological processes activated by IA LPS (Fig. 7B and S10A-C). Notably, there was a downward trend of the expression of genes associated with the TLR, IL-1, and TNF α signaling pathways in the alveolar and interstitial macrophages, which appeared most marked in the animals that had received the combined treatment (Fig. 7B). Blockades also decreased the expression of genes associated with cell migration and apoptotic signaling in these cells (Fig. S10A-B). However, the blockades did not blunt these processes in the inflammatory monocytes (Fig. 7B and S10C).

Neutrophil migration into the lung was also not diminished by any of the blockades (Fig. 8A). As in the inflammatory monocytes, the blocking treatments did not markedly alter the neutrophilic transcriptome (Fig. 8B and S10D).

Blockades lead to a decreased frequency of the mDC population, decreasing it below that present in the control animals (Fig. S11). mDC activation, as evidenced by high HLA-DR MFI, was diminished by the treatments (Fig. S11). In that case, the strongest effect was seen in animals that received IL-1RA, alone or in combination (Fig. S11).

Discussion

Premature infants exposed to IUI are at increased risk of developing pulmonary and intestinal morbidities after birth^{9,36,37}. Herein, we applied several complementary techniques to obtain a comprehensive view of the fetal mucosal immune response to IUI in the highly relevant Rhesus macaque model.

In this model, shortly after IA LPS, robust inflammation developed in the fetal lungs, characterized by the accumulation of immune cells into the lung and increased levels of many inflammatory mediators both in the lung tissue and the AW, similar to what had previously been described in other IUI animal models^{10,11,28,29}. Myeloid cells were particularly involved, as the major cellular change occurring in the lungs of IA LPS-exposed fetuses was the massive accrual of neutrophils and inflammatory monocytes, which were absent in the control lungs. In addition, resident myeloid cells, both the alveolar and interstitial macrophages and the resident myeloid DCs, exhibited many signs of activation. Notably, scRNAseq analyses identified that response to pro-inflammatory cytokines, such as IL-1 and TNF, or to TLR, were activated in these subsets. Many of the activated biological processes were shared by the monocyte/macrophage and neutrophil populations, highlighting a general inflammatory transcriptional program. Overall, these data indicate that the fetal innate immune system can rapidly mount a strong response, a concept that remains controversial. Indeed, there have been many papers describing functional impairment in fetal/neonatal neutrophils and monocytes/macrophages as compared to their juvenile and adult counterparts (rev. in³⁸⁻⁴⁰). Described defects notably include the decreased recruitment of neutrophils and inflammatory monocytes into the lung in response to *E. coli* instillation during the same

time period, compared to juvenile mice⁴¹. In contrast, our data in a NHP model argue that fetal neutrophils and inflammatory monocytes can quickly migrate into the pulmonary environment, as these two populations became very abundant in the lung 16hrs after IA LPS. Neutrophils appeared to localize in the lung in the form of aggregates, a pattern reminiscent of that described in murine models of acute lung injury^{42,43}. Similarly, other groups had described rapid recruitment of neutrophils in fetal sheep or rhesus macaque lungs in response to IA LPS or IL-1 β ^{27,28}. It is also to be noted that the concept of defective functionality of innate immune cells in human neonates mainly arises from studying *in vitro* stimulated cord blood cells. It is therefore possible that these assays do not fully reveal their potential. Another caveat to remember when interpreting data about preterm babies' immune cell functions, is that many of these infants had just been given steroids, which affect various cell type functionality. We did not administer steroids in our studies, which may have allowed us to observe such robust immune responses.

Until recently, circulating and tissue neutrophils were considered as a homogeneous population with a defined array of functions. This concept has been challenged by many studies that revealed a high degree of variety in this population (rev. in⁴⁴⁻⁴⁹). Two main mechanisms appear to drive this diversity, an intrinsic heterogeneity already initiated in bone marrow precursors on one hand, and on the other hand a highly dynamic transcriptional and epigenetic program triggered by exposure to the environment, notably in inflammatory conditions. Our pseudo-time analyses concur with the concept of functional heterogeneity, as they show several gene expression profiles within the neutrophils recruited to the lungs of IA LPS exposed animals. For example, genes related to granule content (*CAMP*, *LCN2*) or antimicrobial defense (*S100A8*, *S100A9*) were in

general not simultaneously expressed along inflammatory mediators (*IL1A*, *IL1B*, *CCL2*, *CCL20*). Interestingly, *Bcl2A1*, a major pro-survival factor for neutrophils⁵⁰, including for LPS-activated neutrophils in the Rhesus chorio-decidua²², was also heterogeneously expressed among the lung neutrophils in LPS-exposed animals, suggesting that differential survival may affect neutrophil heterogeneity. Future studies will be needed to further characterize at the protein level neutrophil heterogeneity and its impact on fetal responses to IUI.

We were also able to evaluate the role played by IL-1 or TNF α in fetal mucosal responses to IA LPS. Our data show that blocking either or both of these pathways had only a partial effect on lung inflammation, a finding consistent with the fact that fetal responses were likely driven by both direct activation by LPS and indirect cytokine-mediated pathways. The partial amelioration of disease by IL-1RA is similar to what has been previously described in other IUI models^{22,23,28,30}, while this was, to our knowledge, the first attempt at evaluating the effect of anti-TNF alone or in combination with IL-1RA. In general, the anti-TNF Ab appeared to have broader effectiveness than IL-1RA, which may be due to the fact that this treatment decreased levels of IL-1 β expression in the lung tissue and AW, whereas IL-1RA did not affect TNF α levels (Tables S2-S3). Interestingly, the combination of both blockades did not radically ameliorate the effectiveness, which suggests that the two pathways are inter-connected, and that other mediators control the recruitment of myeloid cells into the lung. Notably, levels of CCL2, which regulates many aspects of monocyte biology, including recruitment, activation, and survival (rev. in⁵¹), remained elevated in the lung tissue and AW of treated animals. Lung expression of CXCL12, which plays a role in myeloid cell recruitment during lung injury^{52,53}, might also

be involved. Our data also show that high levels of IL-8 persisted in the lung and AW despite the blockades (Tables S2 and S3). Furthermore, many neutrophilic chemoattractant molecules, including chemotactic lipids, formyl peptides, and complement anaphylatoxins are also released during inflammation (rev. in ⁵⁴). We did not quantify these mediators in fetal lungs or AW, but because they are released through both TLR and cytokine-mediated mechanisms, it is likely that they were still released despite the blockades. Interestingly, compared to the fetal lung, myeloid cell recruitment in the placenta was significantly reduced in the presence of either IL-1RA or anti-TNF ^{22,35}, suggesting that the mechanisms controlling cellular migration might be tissue specific. Alternatively, and not exclusively, the fact that cells recruited in the placenta have a maternal origin, in contrast to fetal cells migrating to the lungs, may also have driven some of these differences.

Overall, our work in a model that phenocopies the main features of human IUI gives novel insights into the inflammation rapidly developing in the fetal lung following such exposure. One important conclusion from our new data is that the fetal immune system is highly responsive to its environment. In this model of acute sterile inflammation, we saw that fetal myeloid cells could be rapidly mobilized, migrating to the inflamed environment and displaying profiles of activation very similar to those described in adult cells. This confirms the new paradigm in the field of neonatal immunology, e.g., that fetuses have the capacity to mount a very robust innate immune response. Our data also emphasize the fact that the fetal lung is particularly responsive, as it is in close contact with the amniotic fluid, whereas the fetal gut appears less affected. This study thus provides more insights into why the most solid associations found between IUI exposure and post-natal morbidities

in humans involved mainly the lung (bronchopulmonary dysplasia, wheeziness, and asthma). Our data also show that the multi-parametric mucosal inflammation that develops in IUI-exposed fetuses is not easily controlled by blocking only one, or two inflammatory mediators, emphasizing the need for additional mechanistic studies to better understand the dynamic and complex responses that IUI triggers in exposed fetuses.

Materials and Methods.

Animals.

The Institutional Animal Care and Use Committee at the University of California, Davis approved all animal procedures. Normally cycling, adult female Rhesus macaques (*Macaca mulatta*) were time mated. At approximately 130 days of gestation (about 80% of term gestation), the pregnant dams received either 1 ml saline solution or 1mg of LPS (Sigma-Aldrich, St. Louis MO) in 1 ml saline solution by ultrasound guided IA injection. IA administration of LPS or saline was performed in mothers of similar weights and ages with fetus with similar fetal genders and gestational ages (Table S1). Fetuses were surgically delivered 16, 48 or 120 hrs later by cesarean section. Delivered fetuses were euthanized with pentobarbital and tissues collected. There were no spontaneous deaths or preterm labor in the animals. Some pregnant dams were given, in addition to IA LPS, either the IL-1RA (Kineret® Sobi, Stockholm, Sweden), or the anti-TNF Ab Adalimumab (HUMIRA® AbbVie Chicago, IL), or both. These inhibitors were administered subcutaneously (100mg or 40mg for IL-1RA and anti-TNF respectively) 3hrs before IA LPS as well as IA (50mg or 40 mg for IL-1RA and anti-TNF respectively) 1hr before IA LPS (see scheme on Fig. S9).

Histological evaluation of fetal lung and jejunum.

Inflated lung (30cm water pressure) and jejunum tissues were fixed in 10% formalin immediately following removal. After a series of alcohol and xylene washes, tissues were blocked in paraffin. Paraffin sections were stained with hematoxylin & eosin (H&E). 20x images were taken with a Nikon 90i Upright Widefield Microscope (Nikon Instruments

Inc., Melville, NY). 100µm scale bar added using Nikon Elements Software (Nikon Instruments Inc., Melville, NY).

Immunofluorescence staining of neutrophils and confocal analysis.

Upper left lung lobe sections (5µm thick) were blocked with 4 % goat serum in PBS, and then incubated 1hr at room temperature with the primary mouse-anti-human Ab diluted in 4% goat serum in PBS. Cross-reactivity of all Ab with Rhesus macaque cells was verified on adult tissues prior to their utilization to stain fetal cells. Ab were titrated for optimal detection of positive populations and MFI. Sections stained with unlabeled Ab were stained with a corresponding secondary goat-anti-mouse Ab conjugated with either Alexa Fluor 594, Alexa Fluor 488 or FITC (Life Technologies, dilution 1:400) for 1hr. As negative control, tissues were also stained with the secondary Ab only. Sections were washed and incubated with Vecta shield autofluorescence reduction kit and mounted with mounting media containing DAPI (Vector Laboratories). Images were acquired using an inverted microscope (Ti-E; Nikon, Japan) outfitted with a confocal scan-head (A1R; Nikon; Japan) at the CCHMC Confocal Imaging Core. Analyses involving deconvolution and background reduction were done with Nikon Elements Software.

Since neutrophils in the sections were mostly in clumps/aggregates, we evaluated the number of neutrophil aggregates in 10 fields per animal. The neutrophils were identified as S100A8-FITC (CF-145) and TNFAIP6-AF594 (polyclonal) (both ThermoFisher Waltham, MA) double positive cells. The average area covered by neutrophils clumps were also determined and normalized to the total DAPI area of the image per field. CD68-FITC (KP1) (Alligent Dako, Santa Clara, CA) and HLADR-AF594 (L243) (BioLegend, San Diego, CA) were also used to quantify neutrophils.

Alveolar wash analyses.

Alveolar wash was collected from the left lung. Supernatant was collected and frozen until cytokine analyses, using the NHP multiplex kit according to the manufacturer's protocol (Millipore Austin, TX). Cell pellets were cytospun and slides were stained with Diff quick. 5 fields/slide (100 cells/field) were counted twice.

Quantitative RT-PCR analyses of mucosal tissues.

From frozen lung and jejunum tissue, total RNA was extracted after homogenizing in TRIzol (Invitrogen, Carlsbad, CA). A Nanodrop spectrophotometer (Thermo Fisher Scientific) was used to measure RNA concentration and quality. 1µg of RNA was used to generate cDNA using the Verso cDNA synthesis kit (Thermo Fisher Scientific), following the manufacturer's protocol. Quantitative RT-PCR was done with a StepOnePlus real-time PCR system and rhesus specific TaqMan gene expression primers (Life Technologies, Carlsbad, CA). Eukaryotic 18S rRNA was the endogenous control used for normalization of the target RNAs. The mRNA fold change was calculated relative to the average value of the control group.

Fetal lung cell isolation.

The fetal lung was removed, and left lobe was processed into single cell suspensions as described previously⁵⁵. The large airways were removed and approximately ~500mg of airway free tissue was collected in a gentleMACS C-tube (Miltenyi Biotec, Auburn, CA) and 5mL digest buffer was added and placed on a gentleMACS octo dissociator for programmed runs. Following dissociation, the cell suspension was passed through a 100µm filter and washed with PBS. After washing, cells underwent red cell lysis

(eBioscience, San Diego, CA) along with another pass through a 40um filter after neutralization.

Single cell sequencing of fetal lung cells.

scRNA-seq analyses were done in a subset of animals (Table S4). ~50,000 cells per lung were submitted for single cell sequencing at the Cincinnati Children's Hospital Medical Center DNA Sequencing and Genotyping Core. Approximately ~16,000 cells were loaded into one channel of the Chromium system using the 3 prime v3 single cell reagent kit (10X Genomics, Pleasanton, CA). Following capture and lysis, cDNA was synthesized and amplified as per the manufacturer's protocol (10X Genomics). The amplified cDNA was used to construct Illumina sequencing libraries that were each sequenced using an Illumina HiSeq 4000 machine. Raw sequencing data was processed aligned to the Rhesus macaque reference Mmul_8.0.1 (Ensembl version 91) with Cell Ranger 3.0.2 (10X Genomics) generating expression count matrix files.

After doublet removal with the software DoubletFinder (version 2.0), an integrated analysis was performed using R (version 3.6.3) and the Seurat package (version 3.1.0, <https://satijalab.org/seurat/>), to identify common cell types across different experimental conditions. Specifically, cells with 25% or more mitochondrial transcripts were removed, as well as cells expressing fewer than 150 or more than 5000 features. After log-normalization, 5000 highly variable features were identified per sample using the vst method. Integration was performed using the Seurat FindIntegrationAnchors and IntegrateData functions with default parameter values. The integrated data was scaled, including regressing out mitochondrial percentage and cell cycle variables, followed by

application of PCA reduction. A UMAP projection was estimated based on the PCA reduction and the correlation metric. Clustering of cells was done using the Seurat functions FindNeighbors (based on first 30 PCs) and FindClusters (resolution=4). This high-resolution was initially used to define highly discrete immune populations, with the final populations subsequently aggregated into broader, higher confidence cell-types based common marker genes expressed and corresponding UMAP embeddings. To annotate these cell-populations, we performed a gene-set enrichment analysis on the aggregated cell populations using established human and mouse cell-population marker genes annotated by LungMAP, from the AltAnalyze and ToppCell databases. Immune cell population labels were further evidenced using label transfer with the software cellHarmony against a recently described reference human lung datasets^{56,57}. Clusters were orthogonally validated using the software ICGS2 in AltAnalyze, which does not require a resolution setting, to confirm their presence. Identified immune cell clusters were subset into a new Seurat object and re-clustered. For further analyses, Y-chromosomal genes were removed from the dataset because both IA LPS-exposed fetuses were male whereas animals of both sexes were present in the other experimental groups. Additionally, we checked for the expression of genes *XIST* and *TSIX*, which did not appear in our dataset and seem to not be drivers of differences observed. All identified cell populations were required to have unique reported marker genes to ensure they were not artifacts of “over-clustering”. Differential gene expression analysis was performed as implemented in Seurat. Plots were generated using the ggplot2 package (version 3.3.2). Differentially expressed genes (DEGs) between control and LPS in the monocyte/macrophage cluster with a fold change (FC) of ≥ 1.2 and adjusted p-value of $\leq$

0.05 were used for functional enrichment analysis of biological processes and pathways using the ToppFun web portal⁵⁸. Terms with higher expression in LPS are represented as positive log p-values. Scaled expression of genes for upregulated biological processes terms in the monocyte/macrophage cluster were used to plot parallel coordinate plots using the ggplot2 package.

Differential gene expression analysis was not feasible for the neutrophil or inflammatory monocyte clusters since those cells were not present in control samples (<20 cells). We thus performed single sample gene set enrichment analysis (ssGSEA) in R using the ssgsea2.0 package (github.com/broadinstitute/ssGSEA2.0) in order to identify enriched gene ontology (GO) gene sets (MSigDB v7.0). Expression values of overlapping genes of the most significant and relevant gene sets were plotted as parallel coordinate plots using ggplot2.

In order to assess possibly different states of gene expression within neutrophils after LPS exposure, single-cell trajectory analysis was performed with Monocle3 (version 0.2.3.0). Spatial autocorrelation analysis as implemented in Monocle3 was used to determine genes that most strongly vary along the pseudo-time trajectory. The Monocle algorithm is capable of ordering cells based on their transcriptomic profile in an unsupervised manner and thus arrange cells along a directional path. The order of the cells along this path represents different transcriptomic states within a biological process even if cells are obtained at only one time point during an experiment. The scRNA-seq data have been deposited in the Gene Expression Omnibus (GSE169390, reviewer token:qxatqgkyhxyrhkl).

Flow Cytometry.

A cocktail of conjugated Ab was used to phenotype lung cell suspensions for multi-parameter flow-cytometry. The following Ab were used: HLA-DR (L243), CD11c (3.9) (BioLegend, San Diego, CA); CD3 (GHI/61), CD3 (SP34-2), CD19 (SJ25C1), CD20 (2H7), CD123 (7G3), CD45 (DO58-1283) (BD Biosciences, San Jose, CA); CD8 α (RPA-T8), FoxP3 (PCH101), LIVE/DEAD Fixable Aqua dead cell stain, (eBioscience, San Diego, CA), CD88 (P12/1) (Bio-rad, Hercules, CA). Lung cell suspensions were treated with human IgG to block Fc-receptors and stained for surface markers. Cells were then washed and treated with fixation/permeabilization buffer (ThermoFisher). Following permeabilization, cells were stained for intracellular markers, washed, and resuspended in PBS. Cells were collected on a FACS Fortessa (BD Bioscience San Jose, CA). BD FACSDiva Software v8.0.1 (BD Bioscience San Jose, CA) was used for analysis.

Statistical analyses.

GraphPad Prism 8 (GraphPad Software, La Jolla, California, USA) was used to graph and analyze data for statistical significance. Data were first checked for normality and values expressed as either mean $\pm$ SEM or median and interquartile range. Statistical differences between 2 groups were analyzed using Mann-Whitney U-tests or Student's t-test. For comparison of more than two groups, Kruskal-Wallis or One-way ANOVA were used. Results were considered significantly different for p values ≤ 0.05 . However, due to the limited number of samples in some groups, we also report trends (p values between 0.05 and 0.1).

Author contributions.

C.A.C, A.H.J., H.D., S.G.K., L.A.M., I.P.J. and W.J.Z. conceived the study. C.M.J., S.M., P.P., T.I., P.S., M.C., M.D. and J.G. all performed experiments and data analysis. N.S. and K.C. did the initial pre-processing of scRNAseq 10X files. All authors critically reviewed the manuscript and approved the final version.

Acknowledgments.

The authors would like to thank the staff at the California National Primate Research Center for their outstanding technical support, and especially Paul-Michael Sosa, Jennifer Kendrick, and Sarah Davis and for invaluable help in animal management and care. We also thank Cincinnati Children's Hospital Medical Center's Research Flow Cytometry, Confocal Imaging, and DNA Sequencing and Genotyping Cores.

Grant funding: This work was funded by NIH (U01ES029234, K08HD084686, R01HL142708, K08HL140178-01A1) and CCHMC (Academic and Research Committee Grant).

References

- 1 Goldenberg, R. L., Culhane, J. F., Iams, J. D. & Romero, R. Epidemiology and causes of preterm birth. *The Lancet* **371**, 75-84, doi:10.1016/S0140-6736(08)60074-4 (2008).
- 2 Kim, C. J. *et al.* Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. *Am J Obstet Gynecol* **213**, S29-S52, doi:10.1016/j.ajog.2015.08.040 (2015).
- 3 Saji, F. *et al.* Cytokine production in chorioamnionitis. *Journal of Reproductive Immunology* **47**, 185-196, doi:[https://doi.org/10.1016/S0165-0378\(00\)00064-4](https://doi.org/10.1016/S0165-0378(00)00064-4) (2000).
- 4 Revello, R., Alcaide, M. J., Dudzik, D., Abehsera, D. & Bartha, J. L. Differential amniotic fluid cytokine profile in women with chorioamnionitis with and without funisitis. *The Journal of Maternal-Fetal & Neonatal Medicine* **29**, 2161-2165, doi:10.3109/14767058.2015.1077512 (2016).
- 5 Gomez, R. *et al.* The fetal inflammatory response syndrome. *American Journal of Obstetrics & Gynecology* **179**, 194-202, doi:10.1016/S0002-9378(98)70272-8 (1998).
- 6 Gitlin, D., Kumate, J., Morales, C., Noriega, L. & Arévalo, N. The turnover of amniotic fluid protein in the human conceptus. *American Journal of Obstetrics & Gynecology* **113**, 632-645, doi:10.1016/0002-9378(72)90632-1 (1972).
- 7 ROSS, M. G. & NIJLAND, M. J. M. Fetal Swallowing: Relation to Amniotic Fluid Regulation. *Clinical Obstetrics and Gynecology* **40**, 352-365 (1997).
- 8 Jobe, A. H. Effects of chorioamnionitis on the fetal lung. *Clin Perinatol* **39**, 441-457, doi:10.1016/j.clp.2012.06.010 (2012).
- 9 Been, J. V., Lieveense, S., Zimmermann, L. J. I., Kramer, B. W. & Wolfs, T. G. A. M. Chorioamnionitis as a Risk Factor for Necrotizing Enterocolitis: A Systematic Review and Meta-Analysis. *The Journal of Pediatrics* **162**, 236-242.e232, doi:10.1016/j.jpeds.2012.07.012 (2013).
- 10 Kallapur, S. G., Presicce, P., Rueda, C. M., Jobe, A. H. & Chougnet, C. A. Fetal immune response to chorioamnionitis. *Semin Reprod Med* **32**, 56-67, doi:10.1055/s-0033-1361823 (2014).
- 11 Kallapur, S. G., Willet, K. E., Jobe, A. H., Ikegami, M. & Bachurski, C. J. Intra-amniotic endotoxin: chorioamnionitis precedes lung maturation in preterm lambs. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **280**, L527-L536, doi:10.1152/ajplung.2001.280.3.L527 (2001).
- 12 Wolfs, T. G. A. M. *et al.* Chorioamnionitis-induced fetal gut injury is mediated by direct gut exposure of inflammatory mediators or by lung inflammation. *Am J Physiol Gastrointest Liver Physiol* **306**, G382-G393, doi:10.1152/ajpgi.00260.2013 (2014).
- 13 Gras-Le Guen, C. *et al.* Antenatal infection in the rabbit impairs post-natal growth and lung alveolarisation. *European Respiratory Journal* **32**, 1520-1528, doi:10.1183/09031936.00023708 (2008).
- 14 Pan, J. *et al.* Effects and molecular mechanisms of intrauterine infection/inflammation on lung development. *Respir Res* **19**, 93-93, doi:10.1186/s12931-018-0787-y (2018).
- 15 Wolfs, T. G. A. M. *et al.* Endotoxin induced chorioamnionitis prevents intestinal development during gestation in fetal sheep. *PLoS One* **4**, e5837-e5837, doi:10.1371/journal.pone.0005837 (2009).
- 16 Wolfs, T. G. A. M. *et al.* Antenatal ureaplasma infection impairs development of the fetal ovine gut in an IL-1-dependent manner. *Mucosal Immunology* **6**, 547-556, doi:10.1038/mi.2012.97 (2013).

- 17 Wolfs, T. G. A. M. *et al.* IL-1 α mediated chorioamnionitis induces depletion of FoxP3+ cells and ileal inflammation in the ovine fetal gut. *PLoS One* **6**, e18355-e18355, doi:10.1371/journal.pone.0018355 (2011).
- 18 Nguyen, D. N. *et al.* Prenatal Intra-Amniotic Endotoxin Induces Fetal Gut and Lung Immune Responses and Postnatal Systemic Inflammation in Preterm Pigs. *The American Journal of Pathology* **188**, 2629-2643, doi:10.1016/j.ajpath.2018.07.020 (2018).
- 19 Makori, N. *et al.* Functional and morphological development of lymphoid tissues and immune regulatory and effector function in rhesus monkeys: cytokine-secreting cells, immunoglobulin-secreting cells, and CD5(+) B-1 cells appear early in fetal development. *Clin Diagn Lab Immunol* **10**, 140-153, doi:10.1128/cdli.10.1.140-153.2003 (2003).
- 20 Messaoudi, I., Estep, R., Robinson, B. & Wong, S. W. Nonhuman primate models of human immunology. *Antioxid Redox Signal* **14**, 261-273, doi:10.1089/ars.2010.3241 (2011).
- 21 Miller, L. A., Royer, C. M., Pinkerton, K. E. & Schelegle, E. S. Nonhuman Primate Models of Respiratory Disease: Past, Present, and Future. *ILAR J* **58**, 269-280, doi:10.1093/ilar/ilx030 (2017).
- 22 Presicce, P. *et al.* IL-1 signaling mediates intrauterine inflammation and chorio-decidua neutrophil recruitment and activation. *JCI Insight* **3**, e98306, doi:10.1172/jci.insight.98306 (2018).
- 23 Rueda, C. M. *et al.* Lipopolysaccharide-Induced Chorioamnionitis Promotes IL-1-Dependent Inflammatory FOXP3+ CD4+ T Cells in the Fetal Rhesus Macaque. *J Immunol* **196**, 3706-3715, doi:10.4049/jimmunol.1502613 (2016).
- 24 Novy, M. J. *et al.* Ureaplasma parvum or Mycoplasma hominis as Sole Pathogens Cause Chorioamnionitis, Preterm Delivery, and Fetal Pneumonia in Rhesus Macaques. *Reproductive Sciences* **16**, 56-70, doi:10.1177/1933719108325508 (2009).
- 25 Døllner, H., Vatten, L., Halgunset, J., Rahimipoor, S. & Austgulen, R. Histologic chorioamnionitis and umbilical serum levels of pro-inflammatory cytokines and cytokine inhibitors. *BJOG: An International Journal of Obstetrics & Gynaecology* **109**, 534-539, doi:10.1111/j.1471-0528.2002.01028.x (2002).
- 26 Sadowsky, D. W., Adams, K. M., Gravett, M. G., Witkin, S. S. & Novy, M. J. Preterm labor is induced by intraamniotic infusions of interleukin-1 β and tumor necrosis factor- α but not by interleukin-6 or interleukin-8 in a nonhuman primate model. *American Journal of Obstetrics & Gynecology* **195**, 1578-1589, doi:10.1016/j.ajog.2006.06.072 (2006).
- 27 Kallapur, S. G. *et al.* Intra-amniotic IL-1 β induces fetal inflammation in rhesus monkeys and alters the regulatory T cell/IL-17 balance. *J Immunol* **191**, 1102-1109, doi:10.4049/jimmunol.1300270 (2013).
- 28 Kallapur, S. G. *et al.* IL-1 mediates pulmonary and systemic inflammatory responses to chorioamnionitis induced by lipopolysaccharide. *Am J Respir Crit Care Med* **179**, 955-961, doi:10.1164/rccm.200811-1728OC (2009).
- 29 Willet, K. E. *et al.* Intra-amniotic injection of IL-1 induces inflammation and maturation in fetal sheep lung. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **282**, L411-L420, doi:10.1152/ajplung.00097.2001 (2002).
- 30 Berry, C. A. *et al.* Interleukin-1 in lipopolysaccharide induced chorioamnionitis in the fetal sheep. *Reprod Sci* **18**, 1092-1102, doi:10.1177/1933719111404609 (2011).
- 31 Nikiforou, M. *et al.* Intra-amniotic Candida albicans infection induces mucosal injury and inflammation in the ovine fetal intestine. *Sci Rep* **6**, 29806-29806, doi:10.1038/srep29806 (2016).
- 32 McGovern, N. *et al.* Human fetal dendritic cells promote prenatal T-cell immune suppression through arginase-2. *Nature* **546**, 662-666, doi:10.1038/nature22795 (2017).

635 33 Xie, X. *et al.* Single-cell transcriptome profiling reveals neutrophil heterogeneity in homeostasis
636 and infection. *Nature Immunology* **21**, 1119-1133, doi:10.1038/s41590-020-0736-z (2020).

637 34 Evrard, M. *et al.* Developmental Analysis of Bone Marrow Neutrophils Reveals Populations
638 Specialized in Expansion, Trafficking, and Effector Functions. *Immunity* **48**, 364-379.e368,
639 doi:10.1016/j.immuni.2018.02.002 (2018).

640 35 Presicce, P. *et al.* TNF-Signaling Modulates Neutrophil-Mediated Immunity at the Feto-Maternal
641 Interface During LPS-Induced Intrauterine Inflammation. *Frontiers in Immunology* **11**,
642 doi:10.3389/fimmu.2020.00558 (2020).

643 36 Metcalfe, A., Lisonkova, S., Sabr, Y., Stritzke, A. & Joseph, K. S. Neonatal respiratory morbidity
644 following exposure to chorioamnionitis. *BMC Pediatr* **17**, 128-128, doi:10.1186/s12887-017-
645 0878-9 (2017).

646 37 Lau, J. *et al.* Chorioamnionitis with a fetal inflammatory response is associated with higher
647 neonatal mortality, morbidity, and resource use than chorioamnionitis displaying a maternal
648 inflammatory response only. *American Journal of Obstetrics & Gynecology* **193**, 708-713,
649 doi:10.1016/j.ajog.2005.01.017 (2005).

650 38 Kumar, S. K. M. & Bhat, B. V. Distinct mechanisms of the newborn innate immunity. *Immunology*
651 *Letters* **173**, 42-54, doi:<https://doi.org/10.1016/j.imlet.2016.03.009> (2016).

652 39 Lawrence, S. M., Corriden, R. & Nizet, V. Age-Appropriate Functions and Dysfunctions of the
653 Neonatal Neutrophil. *Frontiers in Pediatrics* **5**, doi:10.3389/fped.2017.00023 (2017).

654 40 Yu, J. C. *et al.* Innate Immunity of Neonates and Infants. *Frontiers in immunology* **9**, 1759-1759,
655 doi:10.3389/fimmu.2018.01759 (2018).

656 41 McGrath-Morrow, S. A. *et al.* The innate immune response to lower respiratory tract E. Coli
657 infection and the role of the CCL2-CCR2 axis in neonatal mice. *Cytokine* **97**, 108-116,
658 doi:10.1016/j.cyto.2017.06.002 (2017).

659 42 Park, I. *et al.* Neutrophils disturb pulmonary microcirculation in sepsis-induced acute lung injury.
660 *Eur Respir J* **53**, 1800786, doi:10.1183/13993003.00786-2018 (2019).

661 43 Zarbock, A., Singbartl, K. & Ley, K. Complete reversal of acid-induced acute lung injury by
662 blocking of platelet-neutrophil aggregation. *J Clin Invest* **116**, 3211-3219, doi:10.1172/JCI29499
663 (2006).

664 44 Filep, J. G. & Ariel, A. Neutrophil heterogeneity and fate in inflamed tissues: implications for the
665 resolution of inflammation. *American Journal of Physiology-Cell Physiology* **319**, C510-C532,
666 doi:10.1152/ajpcell.00181.2020 (2020).

667 45 Silvestre-Roig, C., Hidalgo, A. & Soehnlein, O. Neutrophil heterogeneity: implications for
668 homeostasis and pathogenesis. *Blood* **127**, 2173-2181, doi:10.1182/blood-2016-01-688887
669 (2016).

670 46 Ng, L. G., Ostuni, R. & Hidalgo, A. Heterogeneity of neutrophils. *Nature Reviews Immunology* **19**,
671 255-265, doi:10.1038/s41577-019-0141-8 (2019).

672 47 Alder, M. N. *et al.* Olfactomedin 4 marks a subset of neutrophils in mice. *Innate Immun* **25**, 22-
673 33, doi:10.1177/1753425918817611 (2019).

674 48 Hu, N. *et al.* Coexpression of CD177 and membrane proteinase 3 on neutrophils in
675 antineutrophil cytoplasmic autoantibody-associated systemic vasculitis: Anti-proteinase 3-
676 mediated neutrophil activation is independent of the role of CD177-expressing neutrophils.
677 *Arthritis & Rheumatism* **60**, 1548-1557, doi:10.1002/art.24442 (2009).

678 49 Pillay, J. *et al.* A subset of neutrophils in human systemic inflammation inhibits T cell responses
679 through Mac-1. *J Clin Invest* **122**, 327-336, doi:10.1172/JCI57990 (2012).

680 50 Vier, J., Groth, M., Sochalska, M. & Kirschnek, S. The anti-apoptotic Bcl-2 family protein A1/Bfl-1
681 regulates neutrophil survival and homeostasis and is controlled via PI3K and JAK/STAT signaling.
682 *Cell Death Dis* **7**, e2103-e2103, doi:10.1038/cddis.2016.23 (2016).

683 51 Gschwandtner, M., Derler, R. & Midwood, K. S. More Than Just Attractive: How CCL2 Influences
684 Myeloid Cell Behavior Beyond Chemotaxis. *Frontiers in Immunology* **10**,
685 doi:10.3389/fimmu.2019.02759 (2019).
686 52 Petty, J. M. *et al.* Pulmonary Stromal-Derived Factor-1 Expression and Effect on Neutrophil
687 Recruitment during Acute Lung Injury. *The Journal of Immunology* **178**, 8148-8157,
688 doi:10.4049/jimmunol.178.12.8148 (2007).
689 53 Konrad, F. M., Meichssner, N., Bury, A., Ngamsri, K.-C. & Reutershan, J. Inhibition of SDF-1
690 receptors CXCR4 and CXCR7 attenuates acute pulmonary inflammation via the adenosine A(2B)-
691 receptor on blood cells. *Cell Death Dis* **8**, e2832-e2832, doi:10.1038/cddis.2016.482 (2017).
692 54 Metzemaekers, M., Gouwy, M. & Proost, P. Neutrophil chemoattractant receptors in health and
693 disease: double-edged swords. *Cell Mol Immunol* **17**, 433-450, doi:10.1038/s41423-020-0412-0
694 (2020).
695 55 Zacharias, W. & Morrissey, E. Isolation and culture of human alveolar epithelial progenitor cells.
696 *Protoc. Exch* (2018).
697 56 Travaglini, K. J. *et al.* A molecular cell atlas of the human lung from single-cell RNA sequencing.
698 *Nature* **587**, 619-625, doi:10.1038/s41586-020-2922-4 (2020).
699 57 DePasquale, E. A. K. *et al.* cellHarmony: cell-level matching and holistic comparison of single-cell
700 transcriptomes. *Nucleic Acids Res* **47**, e138, doi:10.1093/nar/gkz789 (2019).
701 58 Chen, J., Bardes, E. E., Aronow, B. J. & Jegga, A. G. ToppGene Suite for gene list enrichment
702 analysis and candidate gene prioritization. *Nucleic Acids Res* **37**, W305-W311,
703 doi:10.1093/nar/gkp427 (2009).

704

Fig. 1. Fetal lung response 16 hours post IA LPS exposure. (A) 20x hematoxylin & eosin (H&E) stain of fetal lung from control and LPS exposed fetuses at 16 hours following IA LPS exposure; stars denote secondary septa, arrows represent infiltrating cells in alveolar space and scale bar is 100 μ m. IL-1 β , TNF α , and IL-6, mRNA expression in fetal lung tissue **(B)** and concentration in the alveolar wash **(C)** in control and 16 hours post IA LPS. Data presented as median and interquartile range, Mann-Whitney U test; ****p $\leq$ 0.0001.

Fig. 2

Fig. 2. Single cell RNAseq analysis of the fetal lung 16 hours post IA LPS exposure. The left lobe of fetal lung was processed into single cell suspensions and submitted for single cell sequencing. **(A)** UMAP from combined control and LPS-exposed animals (2 each). **(B)** Control (left) and IA LPS (right) exposed animals UMAP of re-clustered immune cell populations. **(C)** Representative dot plot of canonical cell type markers used in the identification of hematopoietic populations. **(D)** Feature plot of CD88(C5AR1) expression across hematopoietic cell populations and **(E)** percentage of total CD45⁺ within live cells and CD88⁺ within CD45⁺ cells in lung. Data presented as mean with SEM, student's unpaired t-test (CD45⁺%) or median and interquartile range, Mann-Whitney U test (CD88⁺); #0.05≥p≤0.10.

Fig. 4. Fetal lung neutrophil response to IA LPS. (A) Violin (left) and feature (right) plot of TNFAIP6 and S100A8 expression in fetal lung neutrophils and monocyte/macrophage population following IA LPS exposure. (B) Immunofluorescence of IA LPS exposed fetal lung stained for S100A8 and TNFAIP6; scale bar is 100µm. (C) Neutrophil aggregate count (left) and area (right) in the fetal lungs of control and IA LPS fetuses. Data presented as median and interquartile range, Mann-Whitney U test; **p<0.01. (D) ssGSEA analysis of enriched GO terms in neutrophils. Numbers in parentheses represent the total number of genes for each process.

Fig. 5

A bioRxiv preprint doi: <https://doi.org/10.1101/2021.05.28.444219>; this version posted May 29, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

Fig. 5. Pseudo-time analysis of fetal lung neutrophil response to IA LPS. (A) Pseudo-time trajectory analysis of IA LPS exposed fetal lung neutrophils using Monocle. **(B)** Plots of genes from GO biological pathways response to IL-1 and TLR signaling pathway across pseudo-time.

Fig. 6

A

B

C

Fig. 6. Hematopoietic cell changes in the fetal lung following blocking IL-1 and TNF signaling. (A) 20x H&E sections of fetal control, lps, lps+IL-1RA, lps+anti-TNF, and lps+IL-1RA+anti-TNF exposed animals; stars denote secondary septa, arrows represent cells in alveolar space and scale bar is 100µm. **(B)** UMAPs of the hematopoietic cells from control, lps, lps+IL-1RA, lps+anti-TNF, and lps+ IL-1RA+anti-TNF exposed animals. **(C)** CD88⁺ percentage within the CD45⁺ cells. Data presented as median and interquartile range, Kruskal-Wallis test; **p≤0.01.

Fig. 7

A

B

Alveolar Macrophages

Interstitial Macrophages

Inflammatory Monocytes

Fig. 7. Fetal lung monocyte/macrophage population across treatment conditions. (A) UMAPs of fetal lung myeloid populations and **(B)** parallel coordinate plots of scaled expression of representative genes in select biological processes across treatment conditions in alveolar, interstitial macrophages and inflammatory monocytes.

Fig. 8

Fig. 8. Fetal lung neutrophils in response to IA LPS and blockades. (A) UMAPs of the neutrophil population across treatment conditions. **(B)** Parallel coordinate plots of scaled expression of representative genes in select biological processes across treatment conditions.