1 MATING STRATEGIES OF INVASIVE VERSUS INDIGENOUS CRAYFISH: MULTIPLE
2 PATERNITY AS DRIVER FOR INVASION SUCCESS?

3

4 Running head: Mating strategies invasive vs native crayfish

5 Caterina FRANCESCONI¹, Mălina PÎRVU², Anne SCHRIMPF¹, Ralf SCHULZ¹,
 6 Lucian PÂRVULESCU²⁺, Kathrin THEISSINGER^{1,3+*}

7

8 ¹ Institute for Environmental Sciences, University of Koblenz-Landau, Fortstrasse
 9 7, 76829 Landau, Germany

10 ² Department of Biology-Chemistry, Faculty of Chemistry, Biology, Geography,
 11 West University of Timisoara, 16A Pestalozzi St., 300115 Timisoara, Romania

12 ³LOEWE Centre for Translational Biodiversity Genomics, Senckenberg
 13 Biodiversity and Climate Research Centre, Frankfurt, Germany

14

15 ⁺Authors having equal contribution

16 *Corresponding author: Kathrin Theissinger, kathrin.theissinger@senckenberg.de

17

ABSTRACT

The invasive spiny-cheek crayfish (*Faxonius limosus*) has been able to colonize many European waterbodies since its first introduction into Europe, threatening the indigenous crayfish fauna. *Faxonius limosus*' remarkable reproductive plasticity has been suggested as an important factor contributing to this species' alarming invasiveness. This is the first study comparing the reproductive strategies of an invasive (*F. limosus*) and a sympatric indigenous crayfish (*Pontastacus leptodactylus*). We investigated if and how parthenogenesis and multiple paternity contribute to the invasion process in the River Danube. Using microsatellites, we genotyped the offspring and their mothers of 11 clutches of *F. limosus* and 18 clutches of *P. leptodactylus*. While no parthenogenesis has been found in *F. limosus*' populations, multiple paternity has been detected for the first time in both species, with comparable incidence. The results of the study indicate that multiple paternity does not play a dominant role in *F. limosus*' successful colonization of the Danube. However, the presented results have to be regarded as pilot study, with a limited number of samples and loci investigated. Given the relevance of mating system knowledge for management measures, future studies with larger sample number could provide precious contributions to the conservation actions.

Keywords: *Faxonius limosus*, mating system analyses, microsatellites, invasive species, *Pontastacus leptodactylus*, Danube

INTRODUCTION

Since the mid-19th century, at least 11 non-indigenous crayfish species (NICS) have been purposefully or accidentally introduced into Europe (Jussila *et al.* 2015) and became invasive (Holdich *et al.* 2009); (Jussila *et al.* 2015). Today NICS are widespread across Western and Central Europe and they keep spreading (Kouba *et al.* 2014), causing changes in the distribution patterns of the indigenous crayfish's populations (Jussila *et al.* 2015). Crayfish represent keystone species in freshwater ecosystems, influencing the food web by being an important resource for other animals and by feeding on local vegetation and invertebrates (Reynolds *et al.* 2013). Several NICS can yield high-density populations and, even when replacing indigenous crayfish, they can have a significant effect on the indigenous biota such as benthic fish, molluscs, and macrophytes (Gherardi, 2007). Therefore, the consequences of their introduction, ranging from the danger posed to indigenous crayfish species to the stress imposed on the ecosystem structure, cannot be overlooked (Hobbs *et al.* 1989; Momot, 1995; Reynolds *et al.* 2013).

54 The North American spiny-cheek crayfish *Faxonius limosus* (Rafinesque,
55 1817) was the first non-indigenous crayfish species introduced into Europe in 1890
56 (Holdich *et al.* 2009; Filipová *et al.* 2011). Since its first introduction into Europe
57 this invasive species has been able to quickly increase its population and
58 successfully colonize many European waterbodies and can now be found in at least
59 25 European countries (Kouba *et al.* 2014; Trichkova *et al.* 2015; Govedič, 2017;
60 Kaldre *et al.* 2020). Many studies have addressed the alarming invasiveness of *F.*
61 *limosus* (e.g. Kozák *et al.* 2006; Buřič *et al.* 2013; Kouba *et al.* 2014; Pârvulescu *et al.*
62 2015). The spread of this invasive species represents a threat to all indigenous
63 crayfish (Holdich and Pöckl, 2007). This includes direct stress by competing for the
64 same resources (Holdich and Pöckl, 2007; Lele and Pârvulescu, 2017; Pacioglu *et al.*
65 2020) and by a superior aggressive behaviour (Lele and Pârvulescu, 2017; Weis,
66 2010; Pârvulescu *et al.* 2021), and indirect stress by acting as vector for the crayfish
67 plague pathogen *Aphanomyces astaci* (Holdich and Pöckl, 2007).

68 *Faxonius limosus* exhibits r-selected life strategies typical for invasive
69 crayfish species, such as short life cycle, high fecundity, early maturation, and
70 capability of taking maximum advantage from abundant resources (Lindqvist and
71 Huner, 1999; Kozák *et al.* 2007). This species possesses an exceptional
72 reproductive plasticity (Buřič *et al.* 2013). The division of the mating period in
73 spring and autumn seasons maximizes the probability of successful mating (Buřič
74 *et al.* 2013). The ability of both females and males to alternate between sexually
75 active and inactive forms allows them to direct the resources utilization toward
76 structures useful for specific life stages (Buřič *et al.* 2010b; Buřič *et al.* 2010a).
77 *Faxonius limosus* is also capable of increasing its fecundity, leading to a quick
78 growth of the population by maximizing the exploitation of the resources made
79 available by the decrease of indigenous crayfish populations (Pârvulescu *et al.*
80 2015). *Faxonius limosus* has been found capable of facultative parthenogenesis
81 under laboratory conditions (Buřič *et al.* 2011; Buřič *et al.* 2013). The occurrence
82 of parthenogenesis in the wild would allow reproduction under suboptimal
83 conditions, facilitating the colonization of new habitats from small founding
84 populations (Buřič *et al.* 2011). Lastly, long-term sperm storage (Buřič *et al.* 2013)
85 also allows the reception of spermatophores from different males, while enabling
86 this animal to circumvent adverse environmental conditions (Walker *et al.* 2002).
87 The potentially resulting multiple paternity can enhance the success of a population
88 introduced in a new environment by increasing its genetic diversity.

89 Multiple paternity is widely spread both among vertebrates and
90 invertebrates (Jennions and Petrie, 2000; Avise *et al.* 2011), with variable incidence
91 across and within species (Taylor *et al.* 2014). Multiple mating is associated with
92 considerable costs for females, such as loss of time and energy, and increased risk

of predation, diseases and injuries (Slatyer *et al.* 2012). Nonetheless, the ubiquitous nature of this reproductive strategy indicates that multiple paternity provides significant benefits (Hosken and Stockley, 2003). One of the major genetic benefits is the increased heterozygosity (Holman and Kokko, 2013; Taylor *et al.* 2014). High heterozygosity is linked to disease resistance, greater development stability, competitiveness, hatchability and survivorship, and the ability to respond to novel biotic and abiotic stimuli (Holman and Kokko, 2013; Taylor *et al.* 2014). Therefore, multiple paternity is an efficient method to increase offspring fitness (Palmer and Oldroyd, 2000; Taylor *et al.* 2014) and effective population size (Zeng *et al.* 2017). Multiple paternity has also been linked to the invasion success of non-indigenous species in reptiles (Eales *et al.* 2010), fishes (Zeng *et al.* 2017), mammals (Miller *et al.* 2010), gastropods (Le Cam *et al.* 2009; Rafajlović *et al.* 2013), insects (Ding *et al.* 2017) and malacostracans (Yue *et al.* 2010).

Many studies addressing the invasiveness of crustaceans compare indigenous and invasive species' behaviours (Weis, 2010). This approach allows for a deeper understanding of the contingency of a successful invasion (Stohlgren and Schnase, 2006; Weis, 2010). However, there is a general lack of comparative studies regarding the reproductive strategies of indigenous vs. invasive species (Weis, 2010). In this pilot study we analysed and compared, for the first time, the reproductive strategies of an invasive (*F. limosus*) and an indigenous (*Pontastacus leptodactylus* Eschscholtz 1823) crayfish species occurring in sympatry in the river Danube. *Faxonius limosus* has been detected in the Romanian Danube for the first time in 2009 after downstream dispersal from Serbia (Pârvulescu *et al.* 2009) and has partially replaced the indigenous *P. leptodactylus* in the upper Romanian Danube (Pârvulescu *et al.* 2012; Pârvulescu *et al.* 2015). We aimed to understand the role of different reproductive strategies in the success of the invasion by genotyping female crayfish and their offspring using microsatellite markers. Firstly, we investigated the presence of parthenogenesis in wild populations of *F. limosus*. Secondly, we analysed the incidence of multiple paternity in both crayfish species. We hypothesized that parthenogenesis and multiple paternity are prominent in the invasive species, thus acting as additional drivers for its invasion success.

124

125 MATERIALS AND METHODS

126 Sampling and study design

Female crayfish with eggs were collected in the upper Danube during summer 2016 (Fig. 1). The river was divided into three sectors, based on the timeline of the invasion and the coexistence state of the two species: an old-invaded sector (OID) where the invasion of *F. limosus* dates back at least ten years; an active

130

131 invasion front sector (IF) where both species coexist and *F. limosus* has been
 132 present for no more than three years at the time of samples collection; a non-invaded
 133 sector (NID) where only *P. leptodactylus* is present (Fig. 1). The crayfish were
 134 caught in the littoral area using bait-trap left overnight. In total, eleven female *F.*
 135 *limosus* and their clutches were collected for the analysis (Tab. 1). Four of them
 136 were sampled from the IF sector and seven from the OID sector. Six female *P.*
 137 *leptodactylus* and their clutches were sampled from each sector of the Danube river.
 138 Three families consisted of mother and embryos extracted from eggs, while in the
 139 other clutches (N = 15) the mothers were collected with juveniles (Tab. 1). Tissue
 140 samples were stored in 96% ethanol. The sampling was conducted according to
 141 animal welfare regulations, permits were obtained from the Iron Gates Nature Park
 142 Administration and the Regional Environmental Protection Agency.

143

144 **DNA isolation and microsatellites genotyping**

145 Maternal DNA from female pereopod tissues and progeny DNA from whole
 146 eggs or half juveniles was extracted using a modified high salt DNA extraction
 147 protocol (Aljanabi and Martinez, 1997). DNA pellets were eluted in 50 µL 1xTE
 148 Buffer.

149 For *P. leptodactylus* the genotyping was carried out using eight
 150 microsatellites loci described in Gross *et al.* (2017). The Type-it® Microsatellites
 151 PCR kit (QIAGEN) was used to amplify all loci across two multiplex PCRs:
 152 multiplex A consisted of Aast4_5, Aast4_12, Aast4_32 and multiplex B consisted
 153 of Aast4_8, Aast4_16, Aast4_26, Aast4_30, Aast4_34. The PCR reaction (6 µL)
 154 contained: 0.5 µL of Q-Solution, 2.5 µL of Type-it Multiplex PCR Master Mix, 2
 155 µL of Primer mix (0.25 µL of each primer), 1 µL of RNase-free water and 1 µL of
 156 template DNA. The PCR protocol was the following: initial denaturation step (5
 157 min, 95°C); 30 cycles of denaturation (30 sec, 94°C), annealing (90 sec, 57°C) and
 158 extension (60 sec, 72°C); final extension step (30 min, 60°C).

159 To genotype *F. limosus* six microsatellite primers were chosen following
 160 Buřič *et al.* (2011). The primers were grouped in three PCRs. Multiplex A included
 161 PclG_08, 2.12 and 3.1, multiplex B included PclG_02, PclG_26 and a single PCR
 162 was run for PclG_37. The PCR reaction was conducted under the same conditions
 163 as for *P. leptodactylus* with the exception of the annealing temperature for primers
 164 PclG_08 and PclG_37, set at 53°C. The fragment lengths were detected on a
 165 Beckman Coulter CEQ™ 8000 system. The raw data was analysed using the
 166 software GeneMarker v2.6.4 (Soft-Genetics, State College, PA, USA).

167

168 Data analyses

169 For *F. limosus*, genotypes of the eleven females and four additional adults
 170 sampled in the same survey were used to calculate population allele frequencies and
 171 to verify lack of linkage disequilibrium and agreement with the Hardy-Weinberg
 172 equilibrium. For *P. leptodactylus* allele frequencies were calculated using only 15
 173 of the 18 females analysed, as it was not possible to genotype the mothers from
 174 three clutches due to missing data. For both species allele frequencies, presence of
 175 linkage disequilibrium and agreement with the Hardy-Weinberg equilibrium of the
 176 adult crayfish were tested with GENEPOP 4.2 (Raymond and Rousset, 1995). The
 177 presence of null alleles per locus was tested with Micro-Checker (van Oosterhout
 178 *et al.* 2004). A genotyping error rate of 0.05 for *F. limosus* and 0.04 for *P.*
 179 *leptodactylus* was calculated by blindly repeating 10% of the samples per species.
 180 The occurrence of parthenogenesis in *F. limosus* was verified by comparing the
 181 mothers' genotypes to the genotypes of the respective offspring. Parthenogenesis
 182 was assumed if all offspring had the identical multilocus genotype. Parentage
 183 analysis for both species was conducted using the software GERUD2.0 (Jones,
 184 2005), which reconstructs for each progeny arrays the minimum number of fathers
 185 and their genotypes using polymorphic, codominant markers. The software ranks
 186 the parental genotype combinations by likelihood. The mother's genotype is not
 187 needed, but the offspring array has to share the same mother. GERUD2.0 does not
 188 accept missing data, therefore specimens with too much missing data were not
 189 considered for the analysis. Chi-square test was used to infer the significance of the
 190 different incidence of multiple paternity between the two crayfish species.

191 The software PrDM (Neff and Pitcher, 2002) was used to evaluate the
 192 probability of detecting multiple paternity (PrDM) based on the number of alleles
 193 per locus and their frequency, the number of samples and sires' contribution. Eight
 194 different scenarios were used to represent situations of equal males' contribution
 195 (50:50, 33.3:33.3:33.3 and 25:25:25:25), moderate skewed contribution (66.7:33.3
 196 and 57:28.5:14.5) and highly skewed contribution (70:10:10:10, 90:10 and
 197 85:5:5:5). Those scenarios were based on recommendations by Neff and Pitcher
 198 (2002) and on available multiple paternity data on crayfish (Walker *et al.* 2002; Yue
 199 *et al.* 2010). The average number of offspring per clutch was used for those
 200 simulations. The resulting values represent the power of the applied genetic markers.
 201 Higher values indicate higher probabilities to detect multiple paternity in the
 202 described scenario.

203

204 RESULTS

205 *Pontastacus leptodactylus*

No linkage disequilibrium was detected, and the population was in Hardy-Weinberg equilibrium. No evidence of null alleles was found. Locus Aast4_32 was monomorphic in our population. Excluding locus Aast4_32, the expected heterozygosity of the seven remaining loci varied from 0.14 to 0.61, with an average value of 0.49 (Tab. 2). The observed heterozygosity ranged from 0.15 to 0.67, with an average value of 0.47. Excluding locus Aast4_32, the mean number of alleles per locus was 2.57. Locus Aast4_26 was not genotyped successfully in some families (F29, F11, F05 and E22, S1 Appendix) due to missing data and was therefore excluded from the parentage analyses for the respective clutches. For the family SRB204 the parentage analysis was conducted using only four loci due to missing data from loci Aast4_12, Aast4_32, Aast4_26, Aast4_34 (S1 Appendix). For clutches E22, F05, F28 and F29 the paternity analysis was conducted with unknown mother genotype due to missing data.

Altogether, 284 offspring were successfully genotyped. The number of genotyped progeny per clutch ranged from six to 22 (with an average of 15.8). The PrDM simulations produced probabilities ranging from 0.438 to 0.955 (Tab. 3). The PrDM value of 0.438 was obtained with the 90:10 scenario, while all other probabilities were higher than 0.622. Multiple paternity was detected in two clutches (11%), F22 and SRB111, respectively sampled in the non-invaded sector and the old-invaded sector (Tab. 1). The minimum number of fathers for both clutches was two. The mean number of sires per brood was 1.1.

227

228 *Faxonius limosus*

No linkage disequilibrium among microsatellite loci was detected. Two loci (2.12 and PclG_02) were monomorphic in our population. No null alleles were detected. All loci, with the exception of locus PclG_37, were in Hardy-Weinberg equilibrium. Excluding loci 2.12 and PclG_02, the expected heterozygosity varied from 0.46 to 0.81 for the four remaining loci with an average value of 0.62, while the observed heterozygosity ranged from 0.64 to 0.82 with an average value of 0.73 (Tab. 2). Excluding the two monomorphic loci, the mean number of alleles per locus was 3.75. Locus PclG_24 was excluded from all analysis due to too much missing data.

A total number of 86 offspring was successfully genotyped for all six loci. The number of analysed progeny per clutch ranged between four and 11 (with an average of 7.8). No parthenogenesis was detected in the clutches, as the offspring genotypes exhibited paternal alleles and were no clones from their mothers (S1 Appendix). For clutches FL0049, FL0072, FL0098, FL0111 and FL0137 the paternity analysis was conducted with unknown mother genotype due to missing

244 data in some loci (S1 Appendix). The PrDM ranged from 0.362 and 0.945 (Tab. 3).
 245 The lowest probabilities were obtained in scenarios with high skewed paternal
 246 contribution (90:10 and 85:5:5:5). All the other probabilities are greater than 0.718.
 247 In total, two clutches (18%), one sampled in the OID sector and on in the IF sectors,
 248 were sired by multiple males (Tab. 1). Chi-square test indicated no significant
 249 difference in the incidence of multiple paternity in *F. limosus* and *P. leptodactylus*.

250

251 DISCUSSION

252 To our knowledge, this is the first study investigating the reproductive strategies of
 253 an invasive (*F. limosus*) and an indigenous (*P. leptodactylus*) crayfish species
 254 occurring in sympatry. No parthenogenesis has been found in *F. limosus*'
 255 populations. Nonetheless, the hypothesis of *F. limosus* resorting to this mating
 256 strategy in the wild still needs to be further investigated. The demonstrated
 257 capability of this crayfish of reproducing asexually under laboratory conditions
 258 (Buřič *et al.* 2011; Buřič *et al.* 2013) and the parthenogenesis documented in one
 259 wild population of another crayfish species (Martin *et al.* 2007) provide a sound
 260 theoretical background for this hypothesis.

261 In this study, multiple paternity has been documented in both *F. limosus* and
 262 *P. leptodactylus* for the first time. This mating strategy has been detected in 18% of
 263 the clutches of the invasive *F. limosus* and in 11% of the clutches of the indigenous
 264 *P. leptodactylus*, but the difference in the incidence of multiple paternity between
 265 the two species was not significant. However, our results are not conclusive, as the
 266 study has important limitations. The chosen markers show low genetic diversity for
 267 our populations, with three monomorphic loci across the two species and a low
 268 mean number of alleles per locus. Therefore, the resolution was too low to
 269 adequately address the research questions. Due to lack of project time and funding,
 270 it was not possible to use different markers after the initial analysis. The study was
 271 further limited by the small sample size for *F. limosus*, both in terms of number of
 272 clutches and number of offspring per clutch. The study was originally designed to
 273 include at least 20 offspring per clutch, but the sample size had to be reduced due
 274 to the impossibility of producing data for many loci of an elevated number of
 275 specimens. All these factors contributed to a low probability of detecting multiple
 276 paternity with our genetic markers and dataset.

277 Multiple paternity has been already associated with successful biological
 278 invasions and colonization of new habitats in a variety of animals, both in
 279 vertebrates (Eales *et al.* 2010; Miller *et al.* 2010; Ding *et al.* 2017) and invertebrates
 280 (Le Cam *et al.* 2009; Rafajlović *et al.* 2013; Zeng *et al.* 2017), including crayfish
 281 (Yue *et al.* 2010). It is suggested that multiple paternity leads to higher hatchability

and greater early stage survivorship, probably due to avoidance of genetic incompatibility (Zeh, 1997; Tregenza and Wedell, 1998; Newcomer *et al.* 1999). This polyandrous mating strategy, where one female mates with several males, is associated with increased heterozygosity in the offspring (Holman and Kokko, 2013; Taylor *et al.* 2014). As a result, multiple paternity may act as a buffer against the loss of genetic diversity caused by the bottleneck event that may have occurred after the invasive species' introduction. Multiple paternity has been linked to increased fertility as it could ensure an adequate sperm supply, guaranteeing the fertilization of all the eggs produced by the female (Birkhead and Pizzari, 2002; Hosken and Stockley, 2003). Such benefit could be particularly relevant, as a higher production of eggs by females of *F. limosus* has been observed in the newly invaded habitat (Pârvulescu *et al.* 2015). Females of *F. limosus* can store viable sperm for several months (Buřič *et al.* 2013). The combination of this feature with multiple mating could increase its initial effective population size in the invaded environment (Eales *et al.* 2010). All of those benefits linked to multiple paternity, when associated with *F. limosus* fast life cycle, could be decisive in the establishment of a fast growing, quickly adapting population.

Finally, the importance of acquiring deeper knowledge of *F. limosus* mating behaviours lies in the repercussion it may have both on the management of the invasive *F. limosus* and the conservation of indigenous crayfish species (e.g. Sutherland, 1998; Caro, 1999; Rogowski *et al.* 2013; Wildermuth *et al.* 2013). Management of invasive crayfish species has often produced less than encouraging results (Hyatt, 2004; Freeman *et al.* 2010; Gherardi *et al.* 2011). Part of this inefficiency (besides other reasons, like the difficulty of catching small crayfish) may be traced back to a lack of relevant knowledge on mating behaviours and life strategies of the targeted species (Rogowski *et al.* 2013). In a successful management plan for control or removal of invasive species, the rate of removal must exceed the growth of the population (Bomford and O'Brien, 1995). Therefore, the use of models to estimate population trends is essential (Kajin *et al.* 2012). The predictive value of those demographic models can be enhanced by incorporating information on the mating system (Wildermuth *et al.* 2013). This could be especially useful for species with considerable reproduction ability and plasticity such as crayfish. For these reasons, we believe that further studies should be conducted on *F. limosus*' mating strategies and on their comparison with the mating strategies of indigenous crayfish. While this study did not lead to significant results, it highlighted critical issues met while inferring *F. limosus*' and *P. leptodactylus*' mating systems. Overcoming the limitations of this study by choosing markers with higher polymorphism and bigger sample sizes would lead to more reliable results that could be useful for management plans.

321

322

SUPPLEMENTARY FILE

323 The dataset supporting the conclusions of this article is presented in Supplementary
324 material.

325

326

ACKNOWLEDGEMENTS

327 This work was supported by a grant of the Romanian National Authority for
328 Scientific Research and Innovation, UEFISCDI, project number PN-II-RU-TE-
329 2014-4-0785. We would like to thank Britta Wahl-Ermel for helping with the
330 samples analysis and Mišel Jelić for helping with the selection of the markers.

331

332

LITERATURE CITED

- 333 ALJANABI SM, MARTINEZ I (1997) Universal and rapid salt-extraction of high
334 quality genomic DNA for PCR-based techniques. *Nucleic Acids Res.* 25,
335 4692–4693.
- 336 AVISE JC, TATARENKOV A, LIU J-X (2011) Multiple mating and clutch size in
337 invertebrate brooders versus pregnant vertebrates. *Proc. Natl. Acad. Sci.*
338 *U.S.A.* 108, 11512–11517.
- 339 BIRKHEAD TR, PIZZARI T (2002) Postcopulatory sexual selection. *Nat. Rev. Genet.*
340 3, 262–273.
- 341 BOMFORD M, O'BRIEN P (1995) Eradication of Australia's vertebrate pests: a
342 feasibility study. GRIGG G, HALE P, LUNNEY D, eds. In: Conservation
343 through sustainable use of wildlife, Centre for Conservation Biology, the
344 University of Queensland, Brisbane, Qld., Australia.
- 345 BUŘIČ M, HULÁK M, KOUBA A, PETRUSEK A, KOZÁK P (2011) A successful
346 crayfish invader is capable of facultative parthenogenesis: a novel
347 reproductive mode in decapod crustaceans. *PLoS ONE* 6, e20281.
- 348 BUŘIČ M, KOUBA A, KOZÁK P (2010a) Intra-sex dimorphism in crayfish females.
349 *Zoology* 113, 301–307.
- 350 BUŘIČ M, KOUBA A, KOZÁK P (2010b) Molting and growth in relation to form
351 alternations in the male spiny- cheek crayfish *Orconectes limosus*. *Zool.*

- 352 *Stud.* 49, 28–38.
- 353 BUŘIČ M, KOUBA A, KOZÁK P (2013) Reproductive plasticity in freshwater
354 invader: from long-term sperm storage to parthenogenesis. *PLoS ONE* 8,
355 e77597.
- 356 CARO T (1999) The behaviour–conservation interface. *Trends. Ecol. Evol.* 14,
357 366–369.
- 358 DING G, XU H, OLDROYD BP, GLOAG RS (2017) Extreme polyandry aids the
359 establishment of invasive populations of a social insect. *Heredity* 119, 381–
360 387.
- 361 EALES J, THORPE RS, MALHOTRA A (2010) Colonization history and genetic
362 diversity: adaptive potential in early stage invasions. *Mol. Ecol.* 19, 2858–
363 2869.
- 364 FILIPOVÁ L, LIEB DA, GRANDJEAN F, PETRUSEK A (2011) Haplotype variation in
365 the spiny-cheek crayfish *Orconectes limosus*: colonization of Europe and
366 genetic diversity of native stocks. *J. North Am. Benthol. Soc.* 30, 871–881.
- 367 FREEMAN MA, TURNBULL JF, YEOMANS WE, BEAN CW (2010) Prospects for
368 management strategies of invasive crayfish populations with an emphasis on
369 biological control. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* 20, 211–223.
- 370 GHERARDI F (2007) Understanding the impact of invasive crayfish. Gherardi F,
371 ed. In: Biological invaders in inland waters: Profiles, distribution, and
372 threats, Springer Netherlands, Dordrecht, 507–542.
- 373 GHERARDI F, AQUILONI L, DIÉGUEZ-URIBEONDO J, TRICARICO E (2011) Managing
374 invasive crayfish: is there a hope? *Aquat. Sci.* 73, 185–200.
- 375 GOVEDIČ M (2017) First record of the spiny-cheek crayfish (*Orconectes limosus*)
376 in Slovenia – 300 km upstream from its known distribution in the Drava
377 River. *Knowl. Manag. Aquat. Ecosyst.* 7, 7.
- 378 GROSS R, KÕIV K, PUKK L, KALDRE K (2017) Development and characterization
379 of novel tetranucleotide microsatellite markers in the noble crayfish (*Astacus*
380 *astacus*) suitable for highly multiplexing and for detecting hybrids between
381 the noble crayfish and narrow-clawed crayfish (*A. leptodactylus*).
382 *Aquaculture* 472, 50–56.
- 383 HOBBS HH, JASS JP, HUNER JV (1989) A review of global crayfish introductions
384 with particular emphasis on two North American species (Decapoda,

385 Cambaridae). *Crustaceana* 56, 299–316.

386 HOLDICH DM, PÖCKL M (2007) Invasive crustaceans in European inland waters.
387 Gherardi F, ed. In: Biological invaders in inland waters: Profiles,
388 distribution, and threats, Springer Netherlands, Dordrecht, 29–75.

389 HOLDICH DM, REYNOLDS JD, SOUTY-GROSSET C, SIBLEY PJ (2009) A review of
390 the ever increasing threat to European crayfish from non-indigenous crayfish
391 species. *Knowl. Manag. Aquat. Ecosyst.*, 11.

392 HOLMAN L, KOKKO H (2013) The consequences of polyandry for population
393 viability, extinction risk and conservation. *Philos. Trans. R. Soc. Lond. B.*
394 *Biol. Sci.* 368, 20120053.

395 HOSKEN DJ, STOCKLEY P (2003) Benefits of polyandry: a life history perspective.
396 Macintyre Rand Clegg M, eds. In: Evolutionary Biology, Springer US,
397 Boston, MA, 173–194.

398 HYATT MW (2004) Investigation of crayfish control technology. *Final Report for*
399 *Arizona Game and Fish Department, Cooperative Agreement*, 1448–20181.

400 JENNIONS MD, PETRIE M (2000) Why do females mate multiply? A review of the
401 genetic benefits. *Biol. Reviews* 75, 21–64.

402 JONES AG (2005) GERUD 2.0: a computer program for the reconstruction of
403 parental genotypes from half-sib progeny arrays with known or unknown
404 parents. *Mol. Ecol. Notes* 5, 708–711.

405 JUSSILA J, VREZEC A, MAKONEN J, KORTET R, KOKKO H (2015) Invasive
406 crayfish and their invasive diseases in Europe with the focus on the virulence
407 evolution of the crayfish plague. CANNING-CLODE J, ed. In: Biological
408 Invasions in Changing Ecosystems, De Gruyter Open, Warsaw, Poland.

409 KAJIN M, ALMEIDA PJAL, VIEIRA MV, CERQUEIRA R (2012) The state of the art of
410 population projection models: from the Leslie matrix to evolutionary
411 demography. *Oecol. Austral.* 16, 13–22.

412 KALDRE K, PAAVER T, HURT M, GROSS R (2020) Continuing expansion of non-
413 indigenous crayfish species in Northern Europe: first established spiny-cheek
414 crayfish *Faxonius limosus* (Refinesque, 1817) population in Estonia.
415 *BioInvasions Rec.* 9, 127–132.

416 KOUBA A, PETRUSEK A, KOZÁK P (2014) Continental-wide distribution of crayfish
417 species in Europe: update and maps. *Knowl. Manag. Aquat. Ecosyst.*, 5.

- 418 KOZÁK P, BUŘIČ M, POLICAR T (2006) The fecundity, time of egg development
419 and juvenile production in spiny-cheel crayfish (*Orconectes limosus*) under
420 controlled conditions. *Bull. Franç. Pêche Piscic.*, 1171–1182.
- 421 KOZÁK P, BUŘIČ M, POLICAR T, HAMÁČKOVÁ J, LEPIČOVÁ A (2007) The effect of
422 inter- and intra-specific competition on survival and growth rate of native
423 juvenile noble crayfish *Astacus astacus* and alien spiny-cheek crayfish
424 *Orconectes limosus*. *Hydrobiologia* 590, 85–94.
- 425 LE CAM S, PECHENIK JA, CAGNON M, VIARD F (2009) Fast versus slow larval
426 growth in an invasive marine mollusc: does paternity matter? *J. Hered.* 100,
427 455–464.
- 428 LELE SF, PÂRVULESCU L (2017) Experimental evidence of the successful invader
429 *Orconectes limosus* outcompeting the native *Astacus leptodactylus* in
430 acquiring shelter and food. *Biologia* 72, 41.
- 431 LINDQVIST OV, HUNER JV (1999) Life history characteristics of crayfish: what
432 makes some of them good colonizers? HOLDICH D AND GHERARDI F, eds. In:
433 Crayfish in Europe as alien species, A. A. Balkema, Rotterdam, Brookfield,
434 23–30.
- 435 MARTIN P, KOHLMANN K, SCHOLTZ G (2007) The parthenogenetic Marmorkrebs
436 (marbled crayfish) produces genetically uniform offspring.
437 *Naturwissenschaften* 94, 843–846.
- 438 MILLER S, RUSSELL CJ, MACINNES EH, ABDELKRIM J, FEWSTER R (2010) Multiple
439 paternity in wild populations of invasive *Rattus* species. *New Zeal. J. Ecol.*
440 34.
- 441 MOMOT WT (1995) Redefining the role of crayfish in aquatic ecosystems. *Rev.*
442 *Fish. Sci.* 3, 33–63.
- 443 NEFF BD, PITCHER TE (2002) Assessing the statistical power of genetic analyses
444 to detect multiple mating in fishes. *J. Fish Biol.* 61, 739–750.
- 445 NEWCOMER SD, ZEH JA, ZEH DW (1999) Genetic benefits enhance the
446 reproductive success of polyandrous females. *Proc. Natl. Acad. Sci. U.S.A.*
447 96, 10236–10241.
- 448 PACIOGLU O, THEISSINGER K, ALEXA A, SAMOILĂ C, SÎRBU OI, SCHRIMPF A,
449 ZUBROD JP, SCHULZ R, PÎRVU M, LELE SF, JONES JI, PÂRVULESCU L (2020)
450 Multifaceted implications of the competition between native and invasive
451 crayfish: a glimmer of hope for the native’s long-term survival. *Biol. Inv.* 22:

452 827–842.

453 PALMER KA, OLDROYD BP (2000) Evolution of multiple mating in the genus *Apis*.
454 *Apidologie* 31, 235–248.

455 PÂRVULESCU L, PALOŞ C, MOLNAR P (2009) First record of the spiny-cheek
456 crayfish *Orconectes limosus* (Rafinesque, 1817) (Crustacea: Decapoda:
457 Cambaridae) in Romania. *North-West. J. Zool.* 5, 424–428.

458 PÂRVULESCU L, PÎRVU M, MOROŞAN LG, ZAHARIA C (2015) Plasticity in
459 fecundity highlights the females' importance in the spiny-cheek crayfish
460 invasion mechanism. *Zoology* 118, 424–432.

461 PÂRVULESCU L, SCHRIMPF A, KOZUBÍKOVÁ E, CABANILLAS RESINO S, VRÅLSTAD
462 T, PETRUSEK A, SCHULZ R (2012) Invasive crayfish and crayfish plague on
463 the move: first detection of the plague agent *Aphanomyces astaci* in the
464 Romanian Danube. *Dis. Aquat. Org.* 98, 85–94.

465 PÂRVULESCU L, STOIA DI, MIOK K, ION MC, PUHA AE, STERIE M, VEREŞ M,
466 MARCU I, MUNTEAN MD, ABUREL OM (2021) Force and boldness:
467 Cumulative assts of a successful crayfish invader. *Front. Ecol. Evol.*

468 RAFAJLOVIĆ M, ERIKSSON A, RIMARK A, HINTZ-SALTIN S, CHARRIER G, PANOVA
469 M, ANDRÉ C, JOHANNESSON K, MEHLIG B (2013) The effect of multiple
470 paternity on genetic diversity of small populations during and after
471 colonisation. *PLoS ONE* 8, e75587.

472 RAYMOND M, ROUSSET F (1995) An exact test for population differentiation.
473 *Evolution* 49, 1280.

474 REYNOLDS JD, SOUTY-GROSSET C, RICHARDSON A (2013) Ecological roles of
475 crayfish in freshwater and terrestrial habitats. *Freshw. Crayfish*, 197–218.

476 ŘÍMALOVÁ K, DOUDA K, ŠTAMBERGOVÁ M (2014) Species-specific pattern of
477 crayfish distribution within a river network relates to habitat degradation:
478 implications for conservation. *Biodivers. Conserv.* 23, 3301–3317.

479 ROGOWSKI DL, SITKO S, BONAR SA (2013) Optimising control of invasive
480 crayfish using life-history information. *Freshwater Biol.* 58, 1279–1291.

481 SCHULZ HK, SMETANA P, SCHULZ R (2002) Crayfish occurrence in relation to
482 land-use properties: implementation of a geographic information system
483 (GIS). *Bull. Fr. Pêche Piscic.*, 861–872.

484 SLATYER RA, MAUTZ BS, BACKWELL PR, JENNIONS MD (2012) Estimating

- 485 genetic benefits of polyandry from experimental studies: a meta-analysis.
486 *Biol. Rev. Camb. Philos. Soc.* 87, 1–33.
- 487 STOHLGREN TJ, SCHNASE JL (2006) Risk analysis for biological hazards: what we
488 need to know about invasive species. *Risk Anal.* 26, 163–173.
- 489 SUTHERLAND WJ (1998) The importance of behavioural studies in conservation
490 biology. *Anim. Behav.* 56, 801–809.
- 491 TAYLOR ML, PRICE TA, WEDELL N (2014) Polyandry in nature: a global analysis.
492 *Trends Ecol. Evol.* 29, 376–383.
- 493 TREGENZA T, WEDELL N (1998) Benefits of multiple mates in the cricket *Gryllus*
494 *bimaculatus*. *Evolution: Int. J. Org. Evol.* 52, 1726–1730.
- 495 TRICHKOVA T, TODOROV M, HUBENOV Z, JURAIDA P (2015) First record of
496 *Orconectes limosus* (Rafinesque, 1817) in Bulgaria. Esenias Tools.
497 http://www.esenias.org/index.php?option=com_content&task=view&id=366.
- 498 VAN OOSTERHOUT C, HUTCHINSON WF, WILLIS DP, SHIPLEY P (2004) Micro-
499 checker: software for identifying and correcting genotyping errors in
500 microsatellite data. *Mol. Ecol. Notes* 4, 535–538.
- 501 WALKER D, PORTER BA, AVISE JC (2002) Genetic parentage assessment in the
502 crayfish *Orconectes placidus*, a high-fecundity invertebrate with extended
503 maternal brood care. *Mol. Ecol.* 11, 2115–2122.
- 504 WEIS JS (2010) The role of behavior in the success of invasive crustaceans. *Mar.*
505 *Freshw. Behav. Physiol.* 43, 83–98.
- 506 WILDERMUTH RP, ANADÓN JD, GERBER LR (2013) Monitoring behavior:
507 assessing population status with rapid behavioral assessment. *Conserv. Lett.*
508 6, 86–97.
- 509 YUE GH, LE LI J, WANG CM, XIA JH, WANG GL, FENG JB (2010) High prevalence
510 of multiple paternity in the invasive crayfish species, *Procambarus clarkii*.
511 *Int. J. Biol. Sci.* 107–115.
- 512 ZEH JA (1997) Polyandry and enhanced reproductive success in the harlequin-
513 beetle-riding pseudoscorpion. *Behav. Ecol. Sociobiol.* 40, 111–118.
- 514 ZENG Y, DÍEZ-DEL-MOLINO D, VIDAL O, VERA M, GARCÍA-MARÍN J-L (2017)
515 Multiple paternity and reproduction opportunities for invasive mosquitofish.
516 *Hydrobiologia* 795, 139–151.

517 Tables

518 Tab. 1. Sampling location, sample size, offspring stage and inferred genetic paternity in
519 18 clutches of *Pontastacus leptodactylus* and 11 clutches of *Faxonius limosus*.

Species	Sector	Mother's ID	Known maternal genotype	Offspring stage	No. of offspring	No. of alleles	No. of fathers	MP
PL	NID	E06	Yes	Fertilized eggs	17	8	1	No
PL	NID	E22	No	Fertilized eggs	20	7	1	No
PL	NID	F05	No	Juveniles	18	7	1	No
PL	NID	F11	Yes	Juveniles	17	7	1	No
PL	NID	F28	No	Juveniles	18	8	1	No
PL	NID	F29	No	Juveniles	19	7	2	Yes
PL	OID	SRB061	Yes	Juveniles	18	8	1	No
PL	OID	SRB111	Yes	Juveniles	15	8	2	Yes
PL	OID	SRB141	Yes	Juveniles	19	8	1	No
PL	OID	SRB171	Yes	Juveniles	19	8	1	No
PL	OID	SRB181	Yes	Juveniles	18	8	1	No
PL	OID	SRB204	Yes	Juveniles	7	4	1	No
PL	IF	ESL1	Yes	Juveniles	10	8	1	No
PL	IF	ORS011	Yes	Juveniles	17	8	1	No
PL	IF	ORS021	Yes	Juveniles	14	8	1	No
PL	IF	ORS031	Yes	Juveniles	8	8	1	No
PL	IF	ORS041	Yes	Juveniles	6	8	1	No
PL	IF	ORS254	Yes	Fertilized eggs	22	8	1	No
FL	IF	FL0036	Yes	Juveniles	11	6	3	Yes
FL	IF	FL0049	No	Juveniles	8	6	1	No
FL	IF	FL0072	No	Juveniles	7	6	1	No
FL	IF	FL0268	Yes	Juveniles	9	6	1	No
FL	OID	FL0001	Yes	Juveniles	11	6	3	Yes
FL	OID	FL0085	Yes	Juveniles	9	6	1	No
FL	OID	FL0098	No	Juveniles	4	6	1	No
FL	OID	FL0111	No	Juveniles	8	6	1	No
FL	OID	FL0124	Yes	Juveniles	4	6	1	No
FL	OID	FL0137	No	Juveniles	6	6	1	No
FL	OID	FL0155	Yes	Juveniles	9	6	1	No

520 MP, multiple paternity; PL, *Pontastacus leptodactylus*; FL, *Faxonius limosus*; NID, non-
521 invaded sector; OID, old-invaded sector; IF, invasion front.

522 Tab. 2. Characterization of microsatellite loci for *Faxonius limosus* and *Pontastacus*
523 *leptodactylus*. Estimates are based on adult female genotypes.

Species	Locus	N	N _a	H _o	H _e
FL	3.1	11	3	0.8182	0.6405
FL	2.12	15	1	0.0000	0.0000
FL	PclG_08	14	2	0.7143	0.4592
FL	PclG_02	14	1	0.0000	0.0000
FL	PclG_26	12	3	0.7500	0.5799
FL	PclG_37	11	7	0.6364	0.8140
PL	Aast4_12	15	3	0.5333	0.5800
PL	Aast4_32	15	1	0.0000	0.0000
PL	Aast4_5	15	3	0.5333	0.5800
PL	Aast4_16	15	4	0.6000	0.6067
PL	Aast4_34	13	3	0.1538	0.1450
PL	Aast4_8	15	3	0.6667	0.4867
PL	Aast4_26	14	3	0.4286	0.5383
PL	Aast4_3	15	2	0.4000	0.4800

524 PL, *Pontastacus leptodactylus*; FL, *Faxonius limosus*; n, sample size; N_a, number of
525 alleles; H_o, observed heterozygosity; H_e, expected heterozygosity.

526

527 Tab. 3. Probability of detecting multiple paternity (PrDM) in *F. limosus* and *P.*
 528 *leptodactylus* using 6 and 8 microsatellite loci respectively. Three mating scenarios are
 529 tested: even contribution from the males, moderately skewed contribution, and highly
 530 skewed contribution toward one male. Two to four males were taken into consideration.

Number of males	Mating scenario (paternal skew)	Number of offspring	
		<i>P. leptodactylus</i>	<i>F. limosus</i>
		16	8
2	50:50	0.717	0.756
	66.7:33.3	0.698	0.718
	90:10	0.438	0.362
3	33.3:33.3:33.3	0.903	0.904
	57:28.5:14.5	0.856	0.841
4	25:25:25:25	0.955	0.945
	70:10:10:10	0.838	0.769
	85:5:5:5	0.622	0.516

531

Figure captions

Fig. 1. Sampling sites along the river Danube. NID, non-invaded sector; OID, old-invaded sector; IF, invasion front.

Figure 1