Longevity interventions temporally scale healthspan in *Caenorhabditis elegans*

Cyril Statzer^{1§}, Peter Reichert^{2§}, Jürg Dual^{2*}, and Collin Y. Ewald^{1*}

1 Laboratory of Extracellular Matrix Regeneration, Institute of Translational Medicine,
Department of Health Sciences and Technology, ETH Zürich, Schwerzenbach CH-8603,
Switzerland

2 Eidgenössische Technische Hochschule Zürich, Department of Mechanical and Process
Engineering, Institute for Mechanical Systems, Zürich CH-8092, Switzerland

§ Authors contributed equally

*Corresponding authors: collin-ewald@ethz.ch (CYE) and dual@imes.mavt.ethz.ch (JD)

Keywords: acoustophoresis, microfluidics, healthy aging, maximum muscle strength,
healthspan, sickspan, aging biomarker, lifespan machine

Abstract

Longer lived individuals, such as centenarians or longevity mutants of model organisms, have later onsets and lower incidence rates of late-life morbidities or disabilities than the average population. However, whether increased lifespan is caused by a compression of the portion of life spent in a state of morbidity, i.e., “sickspan,” is highly debated. It is unclear which health matrices are representative for measuring healthspan (time spent in good health); however, muscular performance is generally a good indicator for the health status in humans and a predictor for their mortality, regardless of the underlying cause. Here, we developed a novel microfluidic device that employs acoustophoretic force fields to quantify the maximum muscle strength and dynamic power in aging *C. elegans* populations. We found that longevity mutants have a delayed onset and lower declining rates in maximum muscle strength compared to wild type. Reconciling previous conflicting reports, we confirmed that certain longevity mutants exhibited a mild increase in relative sickspan measured by voluntary movement matrices, which is not the case when using our acoustophoretic force measurements, swim endurance, or other approaches. Using six different biomarkers for healthspan, we observed a time-dependent onset of morbidity, starting with a loss of stress resilience, a decline in dynamic power, and a decline in structural integrity, culminating finally in inactivity (lethargy) and a loss of mobility. We observed that a subset of aging biomarkers correlate with each other and maybe functionally interconnected. Surprisingly, we did not observe a compression of sickspan in longevity mutants but instead observed a temporal scaling of healthspan with diminishing returns for extreme lifespan extensions. Given the conservation of these longevity interventions, this raises the question of whether the healthspan of mammalian longevity interventions is temporally scaled as well.

Introduction

The continuously growing elderly population is projected to result in 1.5 billion people above the age of 65 globally by 2050¹. This poses a significant challenge since old age is the major risk factor for developing cancer, dementia, cardiovascular, and metabolic diseases², especially since people suffer for approximately 20% of their lifespan from one or multiple of these chronic illnesses, which are themselves accompanied by other late-life disabilities². Current estimates indicate that delaying the onset of these chronic diseases by two years would save \$7.1 trillion over the next fifty years in the US alone³. Therefore, major research efforts are dedicated to understanding how to increase the time spent in good health (*i.e.*, healthspan) and to postpone and compress the time spent suffering from age-related pathologies and chronic diseases (*i.e.*, sickspan)^{2,4–6}.

People that are more than one hundred years old, so-called centenarians, display a delayed onset and a lower incidence rate of late-life morbidities compared to people in the age bracket of 80 to 89 years^{7–12}. Genome-wide association studies have shown associations between the exceptional longevity of centenarians and aging-related genes identified in model organisms^{2,13,14}. Mutations in genes that promote longevity in model organisms, such as *C. elegans*, have been instrumental in identifying mechanisms that promote healthy aging^{2,13–15}.

A recent study has questioned this approach of using *C. elegans* longevity mutants to gain insights for promoting healthy aging or mechanisms that prolong healthspan¹⁶. Using four matrices (resilience to heat and oxidative stress, voluntary movement, and swimming performance) to assess the “health” status of aging *C. elegans*, they found that four commonly used longevity mutants outperformed wild type at any given timepoint at older ages, consistent with previous reports. However, compared to their maximum lifespan, longevity mutants displayed an increased sickspan-to-healthspan ratio compared to wild type¹⁶. Other studies have not observed an increase of sickspan in long-lived *C. elegans* mutants, except in the case of lower mobility or movement scores for the insulin/IGF-1 receptor longevity *daf-2(e1370)*

mutants^{17–20}. A large part of the “prolonged sickspan” based on the motility of these *daf-2(e1370)* mutants was attributed to behavioral preferences of food over exploration linked to *odr-10* gene expression¹⁷ or improper dauer-like quiescence behavior^{20–24}. Although all these studies showed that sickspan is not increased in longevity mutants, the question remained about how healthspan changes when the lifespan is extended. We hypothesized that using other health matrices independent of voluntary or behavioral influences, such as physical properties of muscular strength, which is one of the best predictors for all-cause mortality in humans²⁵, we might be able to quantify the health trajectory of *C. elegans* longevity mutants.

Here we confirm that voluntary movement during aging declines and this fragility is not extended in longevity mutants, except mildly in *daf-2* mutants, using high-resolution lifespan and movement measurements on plates. Instead of assessing swimming speed, we measured swimming endurance and found *daf-2* longevity mutants outperforming wild type during aging. We developed a novel microfluidic device and applied acoustophoretic force fields to quantify the maximum force and power of *C. elegans*. Using a high frequency and high power acoustic force field, it becomes possible to set up a contactless, constant in time, and uniform force field acting along the whole *C. elegans* body. Therefore, this force field challenges swimming *C. elegans* in a similar way body-weight exercises do for humans in a gravity field. Furthermore, the acoustic field showed to be suited for stimulating a swimming response of resting *C. elegans*. All longevity mutants showed delayed onset of the decline in maximum force and dynamic power during aging. We observed heterogeneity between individuals across all genotypes in the onset of age-related phenotypes, several correlated phenotypes, and a time-dependent occurrence of multiple disabilities. However, we did not find a compression of sickspan, but rather a temporal scaling of healthspan relative to their maximal lifespan across genotypes.

Results

Voluntary movement healthspan is proportionally increased by longevity interventions

To obtain highly quantitative data on lifespan and healthspan, we used a lifespan machine ²⁶. Here, we defined the “voluntary movement healthspan” as the time spent fast crawling and the “voluntary movement sickspan” as the time spent slow crawling or displaying minimal posture changes (see Materials and Methods for detailed definition). We chose *eat-2(ad1116)* as a genetic model for dietary restriction-mediated longevity, *glp-1(e2141)* as a genetic model for germ-stem-cell-less-mediated longevity, *daf-2(e1368)* and *daf-2(e1370)* as genetic models for reduced insulin/IGF-1 signaling mediated longevity. We cultured all animals at the same temperature (15°C) and in the same environment with the same food source, except *glp-1* that underwent a brief temperature upshift during development as in preparation for the lifespan assay. To avoid dauer-specific traits that occur in reduced insulin/IGF-1 signaling mutants ²², and to avoid pathogenicity from a bacterial food source ²⁰, lifespans were run at 15°C on heat-killed bacteria. Thus, the experimental setup was designed to offer optimal conditions and was kept identical while *C. elegans* genotypes were varied.

As expected, we measured a significant increase in lifespan for these long-lived mutants compared to wild type (Fig. 1a; Supplementary Table 1, Supplementary Video 1). Interestingly, our data showed that the longer-lived the mutant was, the more prolonged was the voluntary movement healthspan (Fig. 1b). Therefore, interventions that increase lifespan also increase the time spent moving fast and actively.

Relative increase for both health- and sickspan in long-lived mutants

To better understand the rescaling of the time spent in frailty in these long-lived mutants, we analyzed the fraction of slow-moving animals per day. We observed gaussian activity distributions, which were shifted along the time axis for these longevity mutants (Fig. 1c). This delayed onset of the sickspan (Fig. 1c) is consistent with the prolonged healthspan of these

long-lived mutants (Fig. 1b). However, except for dietary restricted *eat-2* mutants, the width and the area of the Gaussian distributions were bigger for long-lived mutants than wild type (Fig. 1c), suggesting an overall increase of sickspan. Thus, based on voluntary movement tracking, long-lived mutants display increased absolute health- and absolute sickspan compared to wild type. Next, we asked whether the fraction spent in health- and sickspan during the lifespan is altered. Wild-type animals spent 78% of their lifespan fast-moving and 22% slow-moving (Fig. 1d). For long-lived mutants, we recorded about 70-79% of their lifespan are spent fast-moving (Fig. 1d, Extended Fig. 1), suggesting no compression of sickspan but rather a proportional scaling of both health- and sickspan relative to their lifespan.

Heterogeneity in the length of sickspan but a fixed onset of sickspan

Since we did not observe a compression of voluntary movement sickspan of the entire population, we wondered whether individual animals that outlived their siblings would display a compressed sickspan. When we compared the sickspan traces of individual *C. elegans* for each genotype, we were surprised to measure such a vast heterogeneity (Fig. 1d), given that all these individual animals of a population are genetically identical, consume the same food, and are housed in the same environment. Only the *glp-1*-mediated longevity showed an overall compression of individual sickspan traces (Fig. 1d). For comparison among these different genotypes, we decided to use “relative age” by dividing lifespan curves into quartiles and computing the health-to-sickspan ratio for each quartile (Fig. 1d insets; Extended Data Fig. 2). Consistent with previous reports on wild type ²⁷, we found that in the first quantile of the lifespan curve, individual animals spent about 90% of their lifetime fast-moving and 10% slow-moving, indicating that these animals die young with a compressed sickspan compared to the last quartile wherein animals spent about 75% of their lifetime fast-moving and 25% slow-moving (Fig. 1d insets; Extended Data Fig. 1). Remarkably, it looks like the onset of an individual’s sickspan is a fixed event starting approximately when the first 10% of the isogenic

population starts to die (Fig. 1d). This observation suggests that up to a certain time point, the animal's physiological integrity is maintained. After this time point, there appears to be a stochastic decay resulting in a heterogenous sickspan distribution. Viewing the data using this alternative interpretation of a fixed onset of sickspan would explain why animals in the first quartile of the lifespan curve die young and spend less time in poor health, while animals in the last quartile of the lifespan curve die old and spend more time in poor health. Thus, the time spent fast- vs. slow-moving seems to have a fixed onset in time.

Voluntary movement healthspan temporally scales with lifespan except in *daf-2* mutants

The model of a fixed onset-timepoint for frailty would suggest that longevity interventions would simply delay the onset. To address this, we contrasted the number of days spent fast-moving (healthspan) for each individual as a function of their time lived (lifespan in days; Fig. 1e). We found that the time lived correlated and predicted the time spent fast-moving with an R squared of 0.7 for wild type and R squared ranging from 0.5 to 0.8 for the longevity mutants (Fig. 1e). Furthermore, the *glp-1* with an R squared of 0.8 and *daf-2(e1368)* with an R squared of 0.5 indicate lower or higher heterogeneity, respectively, compared to wild type (Fig. 1e). This is also apparent in the increased or decreased spread of data points below the regression line in Fig. 1e and by increased or decreased lengths of the individual sickspan traces in Fig. 1d, respectively. One interesting aspect to note is that individuals in quartile 2 and 3, which expire in the middle of the lifespan curves, displayed shortened healthspan relative to their lifespan, whereas individuals in the last quartile showed an extended healthspan relative to their lifespan (Fig. 1e). This might be because sicker individuals simply died earlier, leading to an enrichment of healthier-aging individuals in the last quartile (Extended Data Fig. 1, 2). Based on the high R squared values for all genotypes, we applied a linear model to investigate the relationship between health- and lifespan (Fig. 1e). Steeper linear regression lines compared to wild type would indicate an increase in health- to lifespan ratio. The slopes of the linear model

were steeper for *eat-2* and *glp-1*, but less steep for the two *daf-2* mutants compared to wild type (Fig. 1e), suggesting that *glp-1* and *eat-2* spent a larger fraction and *daf-2* mutants spent a smaller fraction of their lifespan actively moving. Since slopes of linear models can be sensitive to extreme values, we compared the population means of health- and lifespan across all genotypes (Fig. 1f). When we extrapolated the mean healthspan to mean lifespan ratio of wild type, we found that *eat-2* and *glp-1* were close to this extrapolated line, whereas the *daf-2* mutants lacked approximately three days (i.e., 7%) of mean healthspan in respect to their mean relative lifespan (Fig. 1f). Thus, we uncovered that the prolonged voluntary movement healthspan temporally scales with the prolonged lifespan for each of these longevity mutants except less stringently for *daf-2* mutants.

Exhaustion of Swimming endurance and stress resilience during aging

Thus far, our observations and interpretations on healthspan are based on the decline of voluntary movement on culturing plates in the abundance of food. Certain genotypes like *daf-2(e1370)* are less motivated to forage and display a more rapid decline in voluntary foraging behavior compared to wild type leading to the interpretation of being less healthy¹⁷. In our setting, this lower foraging behavior is less pronounced in *daf-2(e1370)* since they were cultured at 15°C, an environment that leads to less dauer program activation²². We used a second approach to assess health based on voluntary movement and conducted a swimming assay in the absence of food. Placing *C. elegans* in a water-based physiological buffer (M9 buffer) stimulates active trashing. When animals were young, they swim for more than 30 hours without a decline in activity levels (Fig. 2). In the case of older animals, this endurance capacity declines but less rapidly in longevity mutants (Fig. 2). Similarly, exposing aging animals to the oxidative-stress-inducing agent arsenite, we report a progressive decline in stress resilience of wild-type animals (Fig. 2). This decline in stress resilience is shifted to chronological older animals in long-lived strains compared to wild type (Fig. 2). For both endurance swimming and

stress resilience, aging *daf-2* mutants surpassed chronological age-matched wild-type animals (Fig. 2). Taken together, these results suggest that the observed deficit in healthspan of *daf-2* mutants using a food-dependent movement assay might be largely due to their preferences in lower foraging or exploratory behavior¹⁷ rather than a loss of capability or frailty¹⁶. Moreover, the progressive decline in endurance swimming and stress resilience during aging occurs comparably in both wild type and longevity mutants but is delayed by just 1 or 2 days of chronological age for longevity mutants (Fig. 2). This difference between two voluntary movement assays led us to develop an inducible, motivation-independent exercise platform for *C. elegans*. Our goal is to address the following shortcomings of current methods: the movement should be inducible with a strong stimulus and not dependent on secondary cues like food or intrinsic motivation to thrash, it should be measurable in a short time window to assess health in this instant, and it should directly measure a physiologically relevant parameter like maximum muscle force or functional tissue integrity. This is especially important when comparing different genotypes, which often respond differently to their environment.

Acoustophoretic characterization of *C. elegans* force and muscle power

In humans, one of the best predictors for all-cause mortality is the decline in muscle maximum force and power^{25,28,29}. However, a tool or device to quantify the maximum force and power of *C. elegans* muscles did not exist. The application potential would be immense since *C. elegans* muscle structures are strongly conserved, as in mammals, and forced maximum strength measurements to the point of collapse would be unethical in mammalian models. We developed a microfluidic device harnessing the power of acoustic standing waves (Fig. 3a, 3b, Supplementary Video 2, 3, Supplementary Methods). We have recently applied ultrasonic waves to compress, move and quantitatively characterize larval *C. elegans*³⁰. We reasoned that we could employ ultrasonic standing waves to trap and stretch out *C. elegans* in the minima of the acoustic force fields (Fig. 3c, 3d). *C. elegans* dislike being trapped and try to escape by

applying mechanical forces (body bending) against the acoustic force field (Fig. 3c, 3d). The further away from the acoustic force field minimum, the harder it gets to move against the force field (Fig. 3d). If the animal is stronger than the applied acoustic force field, then it can turn around in the microfluidic chamber (Fig. 3c), typical escaping behavior of *C. elegans* known as omega reversals³¹. Thus, the degree of deflection of the *C. elegans* body away from the acoustic force field minimum provides an estimate of the maximal muscle strength the animal can master to try to escape the acoustic trap.

Muscular strength declines in aging *C. elegans*

To quantify muscular forces, we developed a model by dividing the *C. elegans* body plan into 13 rigid links connected by joints along the animal's midline (Fig. 3e). Upon applying acoustic force fields, we measured the deflection of these 13 nodes for 30 seconds and the number of times the animal escaped the force fields (Fig. 3f). A typical exercise round is structured in up to ten cycles consisting of 30 seconds of ultrasonic force and a 5-second break (Fig. 3g). We measured the muscular forces of aging wild-type *C. elegans* (Fig. 3g). After 3-5 cycles, we observed muscle fatigue, which set in earlier the older the animals were (Fig. 3g). We observed first an increase and then a decrease in the heterogeneity of individual *C. elegans* muscular strengths (Fig. 3g). Similarly, we first saw an increase and then a progressive decline in muscle power during aging (Fig. 3g). By contrast, we found that longevity mutants performed better in terms of muscle strength and function at day 20 of adulthood (Fig. 3h). This indicates the preservation of muscle power in aging longevity mutants.

Longevity mutants showed prolonged healthspan assessed by the strength performance in longitudinal comparison to wild type

The overall force and power of a *C. elegans* depend on its muscle strength as well as its total body size. In agreement with previous reports^{32,33}, we observed adult *C. elegans* kept growing

in body size beyond the reproductive period and then shrunk during aging (Fig. 4a, 4b). A prolonged growing phase correlates with longevity³². We found that longevity mutants prolonged their growing phase and shrunk less than wild-type animals during aging (Fig. 4a, 4b). Structural integrity declines during *C. elegans* aging, such as internal organ atrophy³⁴, loss of internal pressure (Supplementary Method)³⁵, and disorganization of the exoskeleton cuticle occurs³⁶. We noticed that in the acoustic force field, *C. elegans* undergoes compression, and this compressibility stays fairly constant during aging (Fig. 4d, 4e). We conclude that although morphological changes occur during aging, the mechanical properties in regards to compression are less affected by age. This points towards muscular strength playing a key role. On average, young *C. elegans* can overcome the acoustic force field leading to an omega turn eight times each cycle (Fig. 4e, Supplementary Video 4). The ability to overcome the force field and turn in the microfluidic chip progressively declines during aging but is preserved in longevity mutants (Fig. 4e, Extended Data Fig. 3). Turning in the chip can be viewed as a measure of high-intensity muscular capacity since the animal can completely overcome the force field. It also showcases that the animal is not placid but trying to escape. Next, we assessed the overall energy per individual *C. elegans* as an assessment of overall body volume deflected against the force field. We found an increase of energy per individual until mid-age and then a decline (Fig. 4f) reminiscent of the longitudinal body size curve (Fig. 4a). In our measurements to determine the overall force and power of *C. elegans*, the body size is a confounding factor. In all our longitudinal measurements, we had included a positive control in the form of a muscle-defective mutant (CB190) that carries a mutation in the muscle myosin class II heavy chain (*unc-54*). These muscle-constriction-defective mutants were unable to perform omega turns in the chip but showed similar compressibility and longitudinal growth curves, illustrating that the rise and fall of the overall energy are confounded by the organismal growth curve. Therefore, we decided to use the dynamic power as defined as energy expenditure relative to the previous time point and normalized it by the volume of each animal (*i.e.*, volume; Fig. 4g). Using a

human analogy, total refers to how long a weight can be lifted, dynamic power only considers the process of lifting the weight without holding the weight, and dynamic power takes the weight of the person lifting the weight into account. In this way, we found that the overall force and power of longevity mutants were preserved for almost three-quarters (55-83%) compared to about one-third (30%) of their lifespan in wild type (Fig. 4g). Thus, muscular strength is maintained longer in longevity mutants.

Temporal scaling of age-related pathologies in longevity mutants

Next, we asked whether other age-related pathologies or morphological changes show any delayed onset on longevity mutants compared to wild type. We quantified 592 animals, investigated timepoints between day 0 and day 33 (12 animals on average per strain and time point) at the first two cycles of actuation (1183-time sequences), which comprised over 800'000 frames in total. We then manually quantified additional morphological changes such as intestine length and diameter, pixel intensity, wrinkles in the cuticle, and pharynx diameter in a subsampled representative subset (approx. 50'000 frames; Extended Data Fig. 4). Although not all, many age-related phenotypes were delayed in their onset and displayed a slowed decline in longevity mutants compared to wild type (Fig. 5, Extended Data Fig. 5). Rescaling phenotypic trajectories of wild type by the lifespan extension observed in the long-lived strains revealed that many closely match the trajectories observed in long-lived strains for both *daf-2* mutants (Fig. 5C). Notably, for animals' length, diameter, volume, and intestine length, phenotypic scaling was observed when comparing wild type to *daf-2(e1368)* and *daf-2(e1370)*. In the case of *eat-2*, the observed lifespan extension was too limited to draw conclusions and *glp-1* never ceased growing. The severely paralyzed myosin mutant *unc-54(e190)* displayed the opposite phenotypic trajectories than all other genotypes (Fig. 5C). When approximating the phenotypic trajectories as segmented linear fit reflecting the separated phases of growth and decline, we

observed that the starting values as young adults are often similar (Fig. 5D). However, the slopes and point of decline are shifted compared to wild type.

Investigating each phenotype in isolation is hindered by the inherent noise in the measurement as well as by the incomplete picture each phenotype provides. Furthermore, many phenotypes like length, diameter, and volume were strongly correlated. For this reason, we subjected all phenotypes to Principle Component Analysis (PCA) to study the overall age trajectory (Extended Data Fig. 6A). We traced these phenotypes of all genotypes across the PCA plot as they age (Extended Data Fig. 6B). All physiological parameters increased from young to middle-aged and then revert again as the animal reached old age. However, muscle strength density decreased steadily. Using the paralyzed mutant, we were able to establish the bottom left quadrant as a reduced health area. This was only possible when using both physiological and performance measurements and was entered only by the paralyzed strain as well as old wild-type animals. The multi-phenotype traces are also shown for each genotype individually (Extended Data Fig. 6C, 6D). Taken together, this suggested that many of these phenotypes change similarly during aging, that many are temporally scaled in longevity interventions, and that maximum muscle strength offers an orthogonal perspective on studying aging compared to physiological features. This highlights the importance of performing high-intensity muscle strength measurements when studying physiological aging and quantifying healthspan.

Longevity mutants show prolonged absolute but not relative healthspan

Our data revealed that longevity mutants stay healthier compared to wild type and experience a slower decline in physiological integrity. Indeed, dividing the lifespan of each genotype into three chronological fixed age categories: young (less than 7 days), middle (older than 8 but younger than 19 days), and old age (>20 days of adulthood), showed a progressing decline of volume-corrected work performed (Fig. 6a, 6b). Longevity mutants performed better in the middle age group than wild type, but only *daf-2(e1368)* outperformed wild type in the old age

group (Fig. 6a, 6b). Using hierarchical clustering of temporally scaled phenotypes as a complementary analysis, we found that longevity mutants often cluster with chronologically younger wild-type samples (Extended Data Fig. 7).

Integration of voluntary movement and forced maximum muscle strength quantification to yield a comprehensive understanding of *C. elegans* healthspan

Having two independent assessments of healthspan that act on very different stringency levels, we can further divide *C. elegans* healthspan into 3 divisions: prime health (passes both matrices), fragile health (passes 1 metric), and sickspan (failing both matrices) (Fig. 6c). Consistent with previous observations in other species, the maximum power drops prior to the cessation in general mobility^{25,28,29}. Muscle performance is much more improved in longevity mutants in relation to wild type compared to voluntary movement (Fig. 6c). Integrating both matrices revealed that wild type spent around 16% of their lifespan in prime health, whereas longevity mutants spent double the time in prime health (28-42%; Fig. 6c). From all four longevity genotypes, *daf-2(e1368)* appears to be the healthiest strain (Fig. 6c). Thus, combining multiple matrices of physiological and behavioral integrity is a powerful assessment of healthspan.

Discussion

Understanding the relationship between healthspan and lifespan is an important question in aging research since geroscience aims to increase the time spent in good health and to postpone and compress the time suffering from age-related pathologies and chronic diseases^{2,4,5,37}. Model organisms like *C. elegans* are used to identify longevity-promoting interventions that can then be of translational value for humans^{13,15}. There is a fierce debate whether *C. elegans* longevity interventions show compression of sickspan and are of translational value for improving healthspan or healthy aging in humans^{16–20}. In this study, we set out to develop a robust method to quantify maximum muscle strength as a highly interpretable healthspan metric with translational value. Lifespan measurements are well-established and allowed the study of hundreds of lifespan-extending compounds and genetic alterations, leading to ground-breaking discoveries. However, this development is not reflected in the area of healthspan extension. Numerous methods exist which often measure proxy phenotypes for healthspan that are also motivation dependent like pharyngeal pumping, thrashing, and others. With our approach, we were able to directly quantify muscle health in *C. elegans*. This is especially relevant since *C. elegans* is an ideal model system for large-scale genetic screening and both the microfluidic device as well as the image detection can be multiplexed. This approach could translate to the much-needed identification of (muscle) health-promoting interventions.

The microfluidic device operates using acoustophoresis to generate an acoustic force field to quantify the physical fitness and muscle strength of aging *C. elegans*. Using six different ways to assess healthspan in forms of voluntary movements, swimming endurance, stress resilience, muscular force, muscular fatigue, structural integrity/compressibility, and quantifying several age-related morphological changes, including cuticle/skin wrinkles, body, and internal organ sizes, we find that most of these phenotypic changes are postponed in longevity mutants. We observed a hierarchical and time-dependent succession in the occurrence of these phenotypes,

starting with a loss of stress resilience, then a loss of swimming endurance, decline in maximal force as indicated by overcoming acoustic field (omega turns), decline in dynamic power, seizing of body and organ growth (intestine), and then decline in voluntary movement and becoming inactive and lethargic. Using principle component analysis, we show that many of these phenotypes are strongly cross-correlated. The delay of all these age-related phenotypic changes is evident when using chronological age as a reference point for comparison of longevity mutants with wild type but disappears when using relative age as a reference. This points to the idea of temporal scaling of the healthspan. Consistent with this idea, we find that sickspan is neither compressed nor prolonged in longevity mutants compared to wild type. Thus, our quantifications suggest that *C. elegans* healthspan undergoes temporal scaling in longevity.

Aging is defined as a set of phenotypes or senescent pathologies occurring with a higher proportion in older individuals³⁸. Which senescent pathologies limit the lifespan depends on the context and are different for different species, genotypes, and environments^{38,39}. Whether our chosen set of phenotypes assessed are directly limiting or affecting lifespan is unclear. However, it is evident that not one single mechanism underlies all our measured age-related phenotypes. On the other extreme, we do not observe a “one mechanism causing one age-related pathology” mechanism. Our data shows that some phenotypes correlate and also follow a hierarchal time-dependent order of occurrence, indicating that these senescent pathologies are interconnected. This favors a mixed model of several causal mechanisms affecting multiple connected and independent senescent pathologies/phenotypes, including lifespan limiting phenotypes^{38,39}. Even if we might not measure lifespan limiting phenotypes directly, the interconnectedness of phenotypes should reveal the same picture of temporal scaling of age-related pathologies.

The lifespan of *C. elegans* can be increased by up to ten-fold⁴⁰ and decreased by 40-fold, but surprisingly the lifespan curves often follow the same rescaled distribution⁴¹.

Temporal scaling was also noted when comparing expression profiles of longevity mutants compared to wild-type *C. elegans*⁴². Aging is thought to be caused by the accumulation of damage^{14,43}. Organisms either remove, detoxify, compartmentalize, or dilute molecular damage⁴³. In dividing stem cells, yeast, or bacteria, damaged macromolecules or organelles are kept in the “mother” cell leading to accumulation of damage^{14,44,45}. Since *C. elegans* somatic cells are post-mitotic, the main driver of aging dictating the lifespan of *C. elegans* might be the accumulation of damage in non-dividing cells. In a perfect life-conditioned environment such as the laboratory setting, free from ever-changing environmental fluctuations or pathogens, the only lifespan limiting factor probably boils down to the imperfectness of biological systems⁴³, leading to accumulation of damage, stochastic decay of homeostatic mechanisms¹⁴, and consequently to age-related pathologies and phenotypes. In light of this, it might be less surprising that the onset of pathologies is strongly linked to lifespan and thereby, healthspan is rather temporally scaled than prolonged in longevity.

Whether temporal scaling also occurs in mammalian longevity needs to be determined in the future. However, certain longitudinal *C. elegans* phenotypes are comparable to human age-related phenotypes. Analogous to the *C. elegans* volume increase to peak mid-age and then decrease is that human BMI and waist circumference also follows this early-to-mid-life increase reaching a peak around 65-70 years and then declining⁴⁶. Furthermore, grip strength progressively declines after the age of 30-40⁴⁶, similarly to *C. elegans* muscular strength. This raises the question of whether non-compression of sickspan observed in *C. elegans* means or interpolates to non-compression of sickspan in humans? Since aging is universal, it is tempting to speculate that the underlying mechanisms of aging or age-dependent phenotypes are also universal. This might be a potentially erroneous or unproven extension of the observation that almost all living things age³⁸. Although phenotypic changes, such as the loss of *C. elegans* muscle force, is analogous to loss of grip strength or muscle strength loss in humans, the

underlying biological mechanisms resulting in physical weakness might be different. Our study makes no conclusion or interpolation about the compression of the sickspan in humans.

There is an accumulating body of evidence that long-lived humans are healthy during old age. For instance, 56-83% and 15-23% of centenarians, people over the age of hundred years, delay the onset of chronic age-dependent diseases and physical disabilities or were even free of such co-morbidities and frailties, respectively ^{7,10}. Centenarians have lower incidence rates of chronic illnesses compared to their 90- or 80-year old matched-controls ^{8,9,11,12}. This also extends to family members related to centenarians compared to families without centenarians ^{47,48}. Thus, centenarians have a later onset and a lower rate of incidence compared to people in their eighties, similar to our observation when comparing longevity mutants to wild-type *C. elegans*. However, since centenarians get to enjoy at least 20 more years, how would this comparison look if we were to compare relative age to chronological age? Is sickspan compressed or temporally scaled in centenarians compared to the average population? As our life expectancy doubled in the last hundred years and we are on the course of potentially reaching the limit of our lifespan ³⁷, the accompanied delayed onset of disabilities already started to decelerate in longer-lived woman compared to men ⁴⁹. On the other hand, there are several interventions that increase healthspan without increasing lifespan per se identified in mice ^{50,51} and Rhesus monkeys ⁵². Thus, studying longevity is an important first step of identifying molecular mechanisms promoting healthy aging, but our study and others ⁵⁰⁻⁵² point toward that it is crucially important for geroscience to start investigating interventions that improve healthspan directly in future studies ⁶. Initial steps in defining healthspan ^{4,5} and also tools and experimental setups, including this study, are being developed to reliably quantify healthspan ⁵³⁻⁵⁷.

In summary, we have demonstrated that *C. elegans* sickspan is neither compressed nor extended in longevity mutants providing an alternative answer to an ongoing debate in the aging field. With our measurements, we showed that previous claims that insulin/IGF-1 receptor mutants

have increased sickspan compared to wild type are correct if the voluntary movement is measured, but not the case if the muscular function or other healthspan measurements are considered that do not rely on the behavioral state of the animal. By adjusting the reference system from chronological age to relative age, we provide evidence that the healthspan of longevity mutants undergoes temporal scaling. Future studies using our acoustophoresis approach to study the role of healthspan will reveal novel strategies to improve healthy aging.

Author contributions

All authors participated in analyzing and interpreting the data and designing the experiments. CS performed lifespan assays. CYE and CS performed swimming and oxidative stress assays. PR designed and performed microfluidic assays. CS and PR analyzed the datasets jointly. CS and CYE wrote the manuscript in consultation with the other authors.

Author Information

The authors have no competing interests to declare. Correspondence should be addressed to C. Y. E. and J.D.

Acknowledgment

We thank the Ewald lab for constructive comments on the manuscript. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). Funding from the Swiss National Science Foundation PP00P3_163898 to CYE and CS.

21

(0.78) is displayed as a dashed black line. The distance of each population average is marked by a vertical line, and the difference in expected healthspan is indicated.

Extended Data Fig. 1: Paradoxon - young animals die apparently in good health.
The sickspan distribution for every quartile of the lifespan distribution is displayed as boxplots with each strain in a separate

panel (A). Animals experience progressively higher sickspan ratios with each quantile, with the two middle quantiles being the most similar. The distributions of the first and fourth quartile are shown as violin plots with their median line highlighted by a horizontal segment showcasing the increase of relative sickspan and heterogeneity with age across genotypes (B). To compare the individual genotypes for each relative age cohort, the corresponding quartiles are contrasted as boxplots for each strain and each quartile in a separate panel and indicate temporal scaling of healthspan (C). The sickspan ratio is compared across genotypes and quartiles (Mann-Whitney test) and the P-values are depicted as symbols ($ns > 0.05$, $* < 0.05$, $** < 0.01$, $*** < 0.001$, $**** < 0.0001$). When all quartiles are displayed, the median sickspan percentage of the first and last quartile of the wild type population are indicated as dashed horizontal lines. The number of observations associated with every subpopulation is displayed at the bottom of each observation. Each strain is shown in its distinct color, and increasing age is reflected by increasingly dark coloring.

Extended Data Fig. 2: Relative healthspan within early and late dying *C. elegans* groups.

To address the question whether outliers drive the overall health-to-lifespan ratio, each genotype was divided into seven groups with equally sized age bins between the first and last recorded death in each genotype's population to visualize the distribution of death events. At the top of each bar, the number of animals observed to die in this cohort and its share in the overall population is displayed (A). The healthspan distribution in each age group is shown as a violin plot (B). The average percentage of each population spent in their healthspan is displayed above each group, with the population size indicated in brackets. The health-to-lifespan diagonal is shown as a dashed line, and a loess trendline was fitted to the binned datasets with a 95% confidence interval shaded in grey. In case the sample size falls below a threshold, the violin plot was omitted. Each analysis is displayed separately for every *C. elegans* genotype, which are each shown in a separate line and in a distinct color. The spread of the healthspan distribution increases at older ages, with some animals experiencing nearly no sickspan and others spending most of their life in the sickspan portion.

Figure 2: Progressive movement decline with age in unchallenged swimming assay and when exposed to arsenite. In all panels, the time after subjecting the *C. elegans* populations to the arsenite stress is displayed on the x-axis and the movement activity of animals on the y-axis. The animal age at the start of the experiment is shown at the right of every panel. The mean activity in 14 mM arsenite is shown as a continuous line and in M9 as dashed lines. M9 was selected to avoid issues related to osmolarity. Vertical lines reflect the standard deviation at every measurement point. Two *daf-2* alleles and N2 populations were maintained at 15°C and subjected to movement quantification at three aging time points (a). The *daf-2(e1370)* allele was also assayed in a *spe-9* background and therefore maintained at 15°C until L3 and then shifted to 25°C until the start of the assay (b). The *glp-1(e2141)* strain was compared against a *spe-9* population and maintained at 25°C after population lysis until the time of measurement.

Figure 3: Experimental setup to measure maximum *C. elegans* muscle strength:

Top view of the schematic representation of the silicon/glass acoustofluidic chip, fluid in- and outlets, and the piezoelectric transducer positioned at the bottom and *C. elegans* trapped in the standing wave (A). A photograph of the back of the chip together with the metal clamps, the attached fluid connectors, temperature probe, and a piezoelectric transducer is shown (B, top). The front view of the chip provides an overview of the device shape, the microfluidic channel, and the measurement region (MR) (B, bottom). A day 4 wild type *C. elegans*, as seen by the camera in the measurement region, is displayed together with the image processing output highlighting the outline and the segmented midline of the animal as well as the channel borders and centerline (C). The effect of the acoustic field (frequency: 3.543 MHz, voltage amplitude: 76 Vpp, $\lambda/2$ mode) is shown in (C, 2), aligning the animal at the midline. The animal exercises maximum muscle power attempting to bend its head away from the midline (C, 3) to achieve a turn (C, 4), likely as an attempted escape response. The working model to quantify the *C. elegans* muscular force consists of 13 rigid links along the animal's midline, which are connected by joints. Blue arrows reflect the muscle activity acting on each joint to generate a force against the acoustic field (E). The acoustic force acting along the animal's body is modeled as the individual acoustic forces acting on the grey spheres aligned along the body and represented by black arrows, with the length of each arrow being proportional to its force or moment magnitude. The acoustic radiation potential is illustrated using a *C. elegans* cross-section highlighting the animal's four body wall muscles (D). If stretched, the animal rests at the minimum (see C 2) and, upon muscle contraction, moves upwards in its potential energy well (see C3). The four *C. elegans* frames (C 1-4) are put in the context of one 30 seconds actuation cycle showing the time-resolved total energy and mechanical work quantification for this animal (F). Red areas reflect regions of zero energy due to the acoustic field being turned off, and the blue areas indicate *C. elegans* turn movements during which muscle force estimation is paused. A *C. elegans* maximum force assessment routine consists of multiple 30 seconds exercise rounds interspersed by 5-second breaks for a total of up to ten actuation cycles. For wild type, this exercise regimen is displayed with the exercise rounds on the x-axis, the total power (time-averaged) on the y axis, as derived from the respective total energy curve in (F), and the age of the population given at the top of each facet (G). Wild type *C. elegans* is contrasted to long-lived *C. elegans* genotypes across up to ten exercise rounds focusing on the aged cohort above day 20 of adulthood (H).

N2 vs. daf-2(e1368)

Figure 4: Time course of fitness- and structural-related phenotypes for wild type and long-lived *C. elegans* under acoustic stimulation.

Acoustic compression was used to quantify phenotypes directly (# turns, energy, diameter- and volume compression, and dynamic power by volume) as well as indirectly by exploiting the non-destructive linear alignment of the animals (length, volume). Randomly selected images for N2 (A) and daf-2(e1368) (B) at different ages are displayed to illustrate the positioning of the *C. elegans* in the microfluidic measurement channel and how all matrices are shown below were obtained. Measurements are displayed as mean $\pm$ standard error for each assessed age point and subjected to local polynomial regression fitting displayed as a full line with the confidence interval set to 95% and bounded by dashed lines. For the acoustic compression only, the fitted line is shown. The length (C) and volume (D) of the animal was quantified automatically

using the entire length and area of the animal in the channel, respectively. The number of turns was quantified manually and corresponded to the number of times an animal successfully changed its orientation in the channel by 180°. The total energy of the individual was calculated using the magnitude of the lateral deflection of the animal from the channel middle (F). The compression experienced by the animal diameter (G) and volume (H) when the field is activated was computed automatically and displayed as a full line when the field is on and a dashed line when the field is off. The dynamic power/volume (I) reflects the work the animal performs against the field to change its lateral position in the channel and is normalized by the overall volume of the animal to enable comparisons across genotypes making this the most informative health parameter. The mean value for day 0 and day 33 N2 animals are indicated in black dashed lines, and the half-activity value between these two extremes is shown as a full line which is also used to deduce the health- to sickspan transition for each genotype. Using this N2 half-activity value, the individual strains reach their sickspan at approximately 10 days for N2, 28 days for *daf-2(e1368)*, 20 days for *daf-2(e1370)*, 18 days for *eat-2*, and 27 days for *glp-1* while CB190 spends its entire lifespan in its sickspan fraction. Long-lived genotypes, in general, do not experience the same linear energy density decline as wild type since their total energy decrease is slowed down, as is their growth leading to a non-linear energy density trajectory.

Extended Data Fig. 3. *C. elegans* movement in the acoustic exercise chamber.

The exercise regimen was benchmarked for wild type (N2) *C. elegans* at different ages to quantify after how many cycles they experience muscle fatigue (A, B). Every trace corresponds to one individual animal that was up to ten times subjected to 30 seconds maximum force exercise followed by a 5 seconds break (A). Thicker lines represent the population means at every time point. To quantify maximum muscle strength, we selected the first two actuation cycles (B) at which no fatigue can yet be identified. The total energy is computed over the entire body and muscle apparatus of the animal. Interestingly, the spline position most activated in this exercise is the head and tail sections (C). Focusing on the head we quantified the relative head movement for different strains at old age and also observed the same trend of decreasing movement with higher actuation cycles (D). The extreme case of the paralyzed CB190 vs. N2 in young adulthood displays that while the mutant is still able to move the head away from the midline of the field when the acoustic field is off, it is unable to move when the field is on while N2 successfully fights against the applied forces (E).

593

Figure 5. Temporal scaling of *C. elegans* aging phenotypes.

The measured values of selected phenotypes are shown using different scales on the y-axis, animal age on the x-axis, and faceted by strain (A). To address the temporal scaling of phenotypes hypothesis, the loess fit of the measured values for wild type is shown as a solid grey line, and it is temporally scaled values using the respective mean lifespan increase experienced by the respective strain are shown as a dashed line also in grey. The measured phenotype trajectory for each strain is shown in color, each in a separate panel. The maximum fitted value is marked using. White point and its increase relative to the maximum fitted values measured for wild type are shown for both the age and phenotype variables. The same phenotypes as in subfigure (C) are modeled using a piecewise linear relationship in (B). The breakpoint of the segmented fit is estimated by the model at the age value where the linear relationship between the measured phenotype and population age changes. The individual animals measured at each time point are displayed as mean $\pm$ standard error.

A diameter and length (acoustic on/off)

B Assumptions & simplifications

- Rod shaped intestine
- No volume variation during measurement
- No density and compressibility changes
- «Sphere» assumption for acoustic force calculation
- Energy calculation includes no turns folds since these periods are inaccessible to image processing
- Batchwise measurement along the lifespan and the batches are assumed to be identical.

C Intestine length, diameter, pharynx size and wrinkles

- Wrinkles occur rarely and only at very high age (> 25 days)
- Intestine and pharynx are often difficult to detect
- Quantification ability is dependent on the rotation of the animal

D Phenotype interdependence

E Intensity heterogeneity across sections

F Increasing bending angles with age

G Shrinking and wrinkle occurrence with age

H Acoustic field triggers escape response

Extended Data Fig. 4. Phenotypes quantified within the microfluidic setup:

Automated quantification of *C. elegans* diameter (D = diameter) evaluated at the middle segments colored in red and the length (L = length) of the animal along its spine colored in light blue (A). The change in diameter and length quantification as the field is switched (US = ultrasound) is indicated. The assumptions and necessary simplifications that were made to analyze the acquired frames are listed (B). Excerpts of the manual quantification of *C. elegans* intestine length and diameter, pharynx width and height, and the number of cuticle wrinkles are displayed (C). Annotation was performed in a self-developed image annotator suite based on the python programming language. Principal component analysis (PCA) of the

measured phenotypes are displayed for aging wild type (N2) *C. elegans* individuals, with each point representing one individual. The overlaying components refer to the highly correlated features length, volume, and diameter. Tissue heterogeneity and age pigments were studied by analyzing the intensity distribution across different cross-sections along the animal's fitted spine. Representative images of tissue and cuticle weakening displaying an increased bending angle of older animals compared to younger individuals as they attempt to turn in the channel (F). Similarly, older animals are smaller and sometimes display cuticle wrinkles (G). The strong stimulation of the acoustic force in the animals is shown for a representative *daf-2(e1368)* animal (H). The spline midline and head position is displayed for seconds preceding (left), and after (right), the acoustic field is activated. The animal directly responds with a strong escape response.

Extended Data Fig. 5: Rationale behind the partitioning of physical properties into scaled and unscaled phenotypes. Phenotypes displaying an age-dependent change and at least a partial temporal shift for any long-lived strain were manually classified as temporally scaled phenotypes (A, B), and phenotypes that do not satisfy these conditions are categorized as unscaled phenotypes (C, D). The quantified phenotypes are fitted using a loess model in all panels. The observed values for each individual are displayed using mean $\pm$ standard error for each timepoint (A, C). The maximum value predicted by the loess fit is shown, and the increase relative to the wild type maximum is displayed at the top of the graph.

Extended Data Fig. 6: Principal component analysis of temporally scaled phenotypes to identify similarity between phenotypes and their contributions.

The subset of temporally scaled phenotypes for each individual was subjected to principal component analysis, and the first two principle components are shown together with the variance they explain. The contribution of each phenotype to the two first principle components is displayed as vectors with the orientation reflecting the directionality and the color and length capturing the contribution of each phenotype (A). The overall progression of young to middle-aged to old individuals through the phenotype landscape is schematically illustrated. The bottom left quadrant is highlighted in red to indicate its association with animals experience a poor health status. To compare the effect of age on the clustering of the different *C. elegans* genotypes all animals were grouped into 3 age categories, and the individual genotypes were shown in different

646 *colors (B). Confidence ellipses are drawn at the level of 95% for all samples (transparent) and the sample means (non-*
647 *transparent). To compare the effect of age separately for each genotype, the three age categories are shown for each strain,*
648 *young in green, middle-aged in yellow and old animals in red (C). Complete separation of age and genotype is provided in*
649 *panel (D), highlighting the measurements for every individual animal encompassed by the 95% sample confidence ellipse.*

650

651

652

Extended Data Fig. 7: Hierarchical clustering of temporally scaled phenotypes.

The similarity between sampled genotypes and ages are displayed using hierarchical clustering. The comparison between all samples (A), between each long-lived genotype and wild type (B, D, F, H), and within each long-lived is shown (C, E, G, I). The tree is cut in five clusters when comparing all samples and in all other cases in four clusters and colored from left to right.

Age- and longevity-dependent phenotype scaling

Figure 6 Integration of voluntary movement and forced maximum muscle strength quantification to yield a comprehensive understanding of *C. elegans* healthspan.

The volume-corrected relative work performed by each genotype and grouped by age category (young ≤ 7 days, 8 < middle < 20 days, old ≥ 20 days) is shown as boxplots faceted by genotype (A) and by age category (B). P-values of selected comparisons (Mann-Whitney test) are displayed as symbols (ns > 0.05, * < 0.05, ** < 0.01, *** < 0.001, **** < 0.0001). The Population medians for young and old N2 are displayed as horizontal dashed lines across all panels. C) The measured lifespan of each *C. elegans* genotype is displayed with animal age on the x-axis, the fraction of the population that is alive on the y-axis, and the mean and standard error between plates shown as point and line range. The voluntary movement was quantified using the active vs sedentary behavior of the unstimulated animals on plates and is shown for each population as a bar at the top of each panel. Muscle health and the corresponding onset of sickspan due to reduced muscle function is depicted as a bar at the bottom of each facet. The position on the lifespan curve corresponding to the health-to-sickspan transition of either the unstimulated or stimulated healthspan quantification is marked by a white circle. The two health assessments divide the lifespan curve into three segments with decreasing health status reflected by increasing transparency.

Materials and Methods

Strains

Caenorhabditis elegans strains were maintained on NGM plates and OP50 *Escherichia coli* bacteria. The wild-type strain was N2 Bristol. Mutant strains used are described at www.wormbase.org: LGII: *eat-2(ad1116)*; LGIII: *daf-2(e1368, e1370)*, *glp-1(e2141)*.

C. elegans culturing conditions

C. elegans populations were age-synchronized by isolating eggs from gravid *C. elegans* adults, incubating them for 18 hours in M9 until hatching in the presence of 5 µg mL⁻¹ cholesterol (Sigma-Aldrich). Hatched L1-larvae were grown on standard culturing NGM OP50 plates and then shifted to 50 mM FUDR plates seeded with heat-killed OP50 plates when reaching the L4 state. Animals were maintained on FUDR plates until the measurement of different ages was taken. The *glp-1* genotype was placed at 25°C after bleach treatment and shifted back to 15°C at the L4 stage and otherwise treated equally to the other strains. Except for *glp-1* during its development, all animals were always maintained and aged at 15°C.

Automated survival assays using the lifespan machine

To compare the lifespans among wild type and long-lived mutants, we raised all animals for several generations in parallel. Automated survival analysis was conducted using the lifespan machine setup described by ²⁶. Briefly, approximately 1000 L4 animals were resuspended in M9 and transferred to NGM plates containing 50 µM 5-Fluoro-2'-deoxyuridine (FUDR) seeded with heat-killed OP50 bacteria and incubated at 15°C until day 4 of adulthood. Animals were then resuspended in M9 and transferred to fresh FUDR plates containing tight-fitting lids (BD Falcon Petri Dishes, 50x9mm), and the plates were dried with their lids open for 30 minutes after transfer. The plates were incubated for five additional days to rule out contamination and then loaded in the lifespan machine. Air-cooled Epson V800 scanners were utilized for all

experiments operating at a scanning frequency of one scan of 30 minutes. Temperature probes (Thermoworks, Utah, US) were used to monitor the temperature on the scanner flatbed and kept constant at 15°C.

Voluntary movement healthspan measured by lifespan machine

The time point at which the animal stops moving completely and irretrievably is classified as the point of death and defines the lifespan of each individual. The health- to sickspan transition is estimated by the time point when major movement ceases and exclusively head movements, posture change, and minor body movements can be observed. The animal is also required to be sedentary and remains in the rough vicinity of the area; it will ultimately die.

Microfluidics device measuring muscle strength

The detailed description of the microfluidics device development and characterization is found in the Supplementary Methods file.

Oxidative stress assay arsenite with automated wormtracker

Voluntary movement in liquid, as well as resistance to oxidative stress, was quantified using the wMicroTracker (NemaMetrix) platform. Briefly, animals were resuspended in M9 at indicated time points and washed three times with M9 by centrifugation. 30 – 40 *C. elegans* were pipetted into each well of a round-bottomed 96- well plate in either M9 or 14 mM sodium arsenite solution and assayed for 50 hours. Each genotype was always measured both in M9 as well as arsenite in parallel to enable comparisons. Activity measurements were analyzed using the R programming language, and average activity, as well as the standard deviation of wells, are shown.

Figure generation and statistics

The analysis was performed using the statistical software R. data processing and visualization was performed using the tidyverse package collection, most prominently dplyr and ggplot2. Furthermore, packages were used for lifespan analysis (survival, survminer), computing and visualizing PCA (stats and ggfortify, factoextra), fitting loess models (stats), and segmented fits (segmented), labeling (ggrepel) comparing distributions (ggpubr) and arranging figures (cowplot).

References

1. Nations, D. of E. and S. A. P. D. U. World Population Ageing 2019: Highlights. 1–46 (2019).
2. Partridge, L., Deelen, J. & Slagboom, P. E. Facing up to the global challenges of ageing. *Nature* **561**, 45–56 (2018).
3. Goldman, D. P. *et al.* Substantial health and economic returns from delayed aging may warrant a new focus for medical research. *Health affairs (Project Hope)* **32**, 1698–1705 (2013).
4. Kaeberlein, M. Translational geroscience: A new paradigm for 21st century medicine. *Translational Medicine of Aging* **1**, 1–4 (2017).
5. Kennedy, B. K. *et al.* Geroscience: linking aging to chronic disease. *Cell* **159**, 709–713 (2014).
6. Olshansky, S. J. From Lifespan to Healthspan. *JAMA : the journal of the American Medical Association* **320**, 1323–1324 (2018).
7. Ailshire, J. A., Beltrán-Sánchez, H. & Crimmins, E. M. Becoming centenarians: disease and functioning trajectories of older US Adults as they survive to 100. *The journals of gerontology Series A, Biological sciences and medical sciences* **70**, 193–201 (2015).
8. Andersen, S. L., Sebastiani, P., Dworkis, D. A., Feldman, L. & Perls, T. T. Health span approximates life span among many supercentenarians: compression of morbidity at the approximate limit of life span. *The journals of gerontology Series A, Biological sciences and medical sciences* **67**, 395–405 (2012).
9. Evans, C. J. *et al.* Place and cause of death in centenarians: a population-based observational study in England, 2001 to 2010. *PLoS medicine* **11**, e1001653 (2014).
10. Evert, J., Lawler, E., Bogan, H. & Perls, T. Morbidity profiles of centenarians: survivors, delayers, and escapers. *The Journals of Gerontology: Series A* **58**, 232–237 (2003).
11. Ismail, K. *et al.* Compression of Morbidity Is Observed Across Cohorts with Exceptional Longevity. *Journal of the American Geriatrics Society* **64**, 1583–1591 (2016).
12. Kheirbek, R. E. *et al.* Characteristics and Incidence of Chronic Illness in Community-Dwelling Predominantly Male U.S. Veteran Centenarians. *Journal of the American Geriatrics Society* **65**, 2100–2106 (2017).
13. Kenyon, C. J. The genetics of ageing. *Nature* **464**, 504–512 (2010).
14. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of aging. *Cell* **153**, 1194–1217 (2013).

- 788 15. Magalhães, J. P. de, Stevens, M. & Thornton, D. The Business of Anti-Aging Science.
789 *Trends in Biotechnology* **35**, 1062–1073 (2017).
- 790 16. Bansal, A., Zhu, L. J., Yen, K. & Tissenbaum, H. A. Uncoupling lifespan and healthspan
791 in *Caenorhabditis elegans* longevity mutants. *Proceedings of the National Academy of*
792 *Sciences* **112**, E277–86 (2015).
- 793 17. Hahm, J.-H. *et al.* *C. elegans* maximum velocity correlates with healthspan and is
794 maintained in worms with an insulin receptor mutation. *Nature communications* **6**, 8919
795 (2015).
- 796 18. Huang, C., Xiong, C. & Kornfeld, K. Measurements of age-related changes of
797 physiological processes that predict lifespan of *Caenorhabditis elegans*. *Proceedings of the*
798 *National Academy of Sciences of the United States of America* **101**, 8084–8089 (2004).
- 799 19. Stamper, B. L. N. *et al.* Movement decline across lifespan of *Caenorhabditis elegans*
800 mutants in the insulin/insulin-like signaling pathway. *Aging cell* **17**, e12704 (2018).
- 801 20. Podshivalova, K. & Kerr, R. A. How a Mutation that Slows Aging Can Also
802 Disproportionately Extend End-of-Life Decrepitude. *Cell reports* **19**, 441–450 (2017).
- 803 21. Ewald, C. Y., Landis, J. N., Abate, J. P., Murphy, C. T. & Blackwell, T. K. Dauer-
804 independent insulin/IGF-1-signalling implicates collagen remodelling in longevity. *Nature*
805 **519**, 97–101 (2015).
- 806 22. Ewald, C. Y., Castillo-Quan, J. I. & Blackwell, T. K. Untangling Longevity, Dauer, and
807 Healthspan in *Caenorhabditis elegans* Insulin/IGF-1-Signalling. *Gerontology* **64**, 96–104
808 (2018).
- 809 23. Gems, D. *et al.* Two pleiotropic classes of *daf-2* mutation affect larval arrest, adult
810 behavior, reproduction and longevity in *Caenorhabditis elegans*. *Genetics* **150**, 129–155
811 (1998).
- 812 24. Hess, M., Gomariz, A., Goksel, O. & Ewald, C. Y. In-vivo quantitative image analysis of
813 age-related morphological changes of *C. elegans* neurons reveals a correlation between
814 neurite bending and novel neurite outgrowths. *Eneuro* **6**, ENEURO.0014-19.2019 (2019).
- 815 25. Leong, D. P. *et al.* Prognostic value of grip strength: findings from the Prospective Urban
816 Rural Epidemiology (PURE) study. *Lancet* **386**, 266–273 (2015).
- 817 26. Stroustrup, N. *et al.* The *Caenorhabditis elegans* Lifespan Machine. *Nature Methods* **10**,
818 665–670 (2013).
- 819 27. Zhang, W. B. *et al.* Extended Twilight among Isogenic *C. elegans* Causes a
820 Disproportionate Scaling between Lifespan and Health. *Cell Systems* **3**, 333–345.e4 (2016).
- 821 28. Petrella, J. K., Kim, J., Tuggle, S. C., Hall, S. R. & Bamman, M. M. Age differences in
822 knee extension power, contractile velocity, and fatigability. *J Appl Physiol* **98**, 211–220
823 (2005).

824 29. Kostka, T. Quadriceps maximal power and optimal shortening velocity in 335 men aged
825 23–88 years. *Eur J Appl Physiol* **95**, 140–145 (2005).

826 30. Baasch, T. *et al.* Acoustic Compressibility of *Caenorhabditis elegans*. *Biophysical journal*
827 **115**, 1817–1825 (2018).

828 31. Donnelly, J. L. *et al.* Monoaminergic Orchestration of Motor Programs in a Complex *C.*
829 *elegans* Behavior. *Plos Biol* **11**, e1001529 (2013).

830 32. Hulme, S. E. *et al.* Lifespan-on-a-chip: microfluidic chambers for performing lifelong
831 observation of *C. elegans*. *Lab on a chip* **10**, 589–597 (2010).

832 33. Shi, C. & Murphy, C. T. Mating induces shrinking and death in *Caenorhabditis* mothers.
833 *Science (New York, NY)* **343**, 536–540 (2014).

834 34. Ezcurra, M. *et al.* *C. elegans* Eats Its Own Intestine to Make Yolk Leading to Multiple
835 Senescent Pathologies. *Current biology : CB* **28**, 2544–2556.e5 (2018).

836 35. Gilpin, W., Uppaluri, S. & Brangwynne, C. P. Worms under Pressure: Bulk Mechanical
837 Properties of *C. elegans* Are Independent of the Cuticle. *Biophysical journal* **108**, 1887–1898
838 (2015).

839 36. Essmann, C. L. *et al.* Mechanical properties measured by atomic force microscopy define
840 health biomarkers in ageing *C. elegans*. *Nature communications* **11**, 1043 (2020).

841 37. Olshansky, S. J. & Carnes, B. A. Inconvenient Truths About Human Longevity. *The*
842 *journals of gerontology Series A, Biological sciences and medical sciences* **74**, S7–S12
843 (2019).

844 38. Freund, A. Untangling Aging Using Dynamic, Organism-Level Phenotypic Networks.
845 *Cell Systems* **8**, 172–181 (2019).

846 39. Gems, D. The aging-disease false dichotomy: understanding senescence as pathology.
847 *Frontiers in genetics* **6**, 212 (2015).

848 40. Ayyadevara, S., Alla, R., Thaden, J. J. & Reis, R. J. S. Remarkable longevity and stress
849 resistance of nematode PI3K-null mutants. *Aging cell* **7**, 13–22 (2008).

850 41. Stroustrup, N. *et al.* The temporal scaling of *Caenorhabditis elegans* ageing. *Nature* **530**,
851 103–107 (2016).

852 42. Tarkhov, A. E. *et al.* A universal transcriptomic signature of age reveals the temporal
853 scaling of *Caenorhabditis elegans* aging trajectories. *Scientific reports* **9**, 7368 (2019).

854 43. Gladyshev, V. N. The origin of aging: imperfectness-driven non-random damage defines
855 the aging process and control of lifespan. *Trends in genetics : TIG* **29**, 506–512 (2013).

856 44. Lippuner, A. D., Julou, T. & Barral, Y. Budding yeast as a model organism to study the
857 effects of age. *FEMS microbiology reviews* **38**, 300–325 (2014).

45. Yang, Y. *et al.* Temporal scaling of aging as an adaptive strategy of *Escherichia coli*. *Science advances* **5**, eaaw2069 (2019).
46. Kuo, P. L. *et al.* A roadmap to build a phenotypic metric of ageing: insights from the Baltimore Longitudinal Study of Aging. *Journal of Internal Medicine* **287**, 373–394 (2020).
47. Ash, A. S. *et al.* Are Members of Long-Lived Families Healthier Than Their Equally Long-Lived Peers? Evidence From the Long Life Family Study. *The journals of gerontology Series A, Biological sciences and medical sciences* **70**, 971–976 (2015).
48. Sebastiani, P. *et al.* Families Enriched for Exceptional Longevity also have Increased Health-Span: Findings from the Long Life Family Study. *Frontiers in public health* **1**, 38 (2013).
49. Freedman, V. A., Wolf, D. A. & Spillman, B. C. Disability-Free Life Expectancy Over 30 Years: A Growing Female Disadvantage in the US Population. *American journal of public health* **106**, 1079–1085 (2016).
50. Fischer, K. E. *et al.* A cross-sectional study of male and female C57BL/6Nia mice suggests lifespan and healthspan are not necessarily correlated. *Aging* **8**, 2370–2391 (2016).
51. Garcia-Valles, R. *et al.* Life-long spontaneous exercise does not prolong lifespan but improves health span in mice. *Longevity & healthspan* **2**, 14 (2013).
52. Mattison, J. A. *et al.* Impact of caloric restriction on health and survival in rhesus monkeys from the NIA study. *Nature* **489**, 318–321 (2012).
53. Bellantuono, I. *et al.* A toolbox for the longitudinal assessment of healthspan in aging mice. *Nature Protocols* **15**, 540–574 (2020).
54. Collins, J. J., Huang, C., Hughes, S. & Kornfeld, K. The measurement and analysis of age-related changes in *Caenorhabditis elegans*. *WormBook: the online review of C elegans biology* 1–21 (2008) doi:10.1895/wormbook.1.137.1.
55. Teuscher, A. C., Statzer, C., Pantasis, S., Bordoli, M. R. & Ewald, C. Y. Assessing Collagen Deposition During Aging in Mammalian Tissue and in *Caenorhabditis elegans*. *Methods Mol Biology Clifton N J* **1944**, 169–188 (2019).
56. Haefke, A. & Ewald, C. Tail Tendon Break Time for the Assessment of Aging and Longitudinal Healthspan in Mice. *Bio-protocol* **10**, (2020).
57. Teuscher, A. C. & Ewald, C. Y. Overcoming Autofluorescence to Assess GFP Expression During Normal Physiology and Aging in *Caenorhabditis elegans*. *BIO-PROTOCOL* **8**, (2018).