Bacterial predation on T4 phages

Jean-Jacques Godon^a, Ariane Bize^b, Hoang Ngo^b, Laurent Cauquil^c, Mathieu Almeida^d,
Marie Agnès Petit^e, Olivier Zemb^c.

Affiliations

^aINRAE, Univ Montpellier, LBE, Narbonne, France, 11100

^bINRAE, Hydrosyst & Bioproc Res Unit, Antony, France, 92160

^cINRAE, Univ Toulouse, ENVT, GenPhySE, Toulouse, France, 31320

^dINRAE, MetaGenoPolis, Jouy-en-Josas, France, 78350

^eUniversité Paris- Saclay, INRAE, AgroParisTech, Micalis Institute, Jouy-en-Josas,
France, 78350

* Paste corresponding author name(s) here.

Email: olivier.zemb@inrae.fr

Author Contributions: J.J.G., O.Z., A.B. and N.H. designed and performed the
experiments; J.J.G., O.Z., L.C. and M.A. performed computational experiments; J.J.G.,
O.Z., L.C., M.A. and M.A.P. contributed lineaging data and expertise; J.J.G. and O.Z.

16 prepared the manuscript with assistance from all authors; J.J.G. and O.Z. supervised and

17 directed the research.

18 Competing Interest Statement: no competing interest.

19 **Classification:** Microbiology

20 **Keywords:** bacteriophage, Aeromonas, Stable-isotope probing

21 Abstract

22 **Background:** Bacterial consumption of viruses has never yet been
 23 reported, even though bacteria feed on almost anything. Viruses are
 24 omnipresent predators for all organisms, but have no acknowledged active
 25 biocontrol. The viral biomass undoubtedly reintegrates the trophic cycles,
 26 however the mechanisms of this phase still remain unknown.

27 **Methods:** Here, we used stable isotope probing with ^{13}C labelled T4 phages
 28 to monitor the increase of density of the bacterial DNA concomitant with
 29 the decrease of plaque forming units. We used ^{12}C T4 phages as control.

30 **Results:** T4 phage disappearance in wastewater sludge was found to occur
 31 mainly through predation by *Aeromonadacea*. Phage consumption also
 32 favours significant *in situ* bacterial growth. Furthermore, an isolated
 33 strain of *Aeromonas* was observed to grow on T4 phages as sole source of
 34 carbon, nitrogen and phosphorus.

35 **Conclusions:** bacterial species are capable of consuming bacteriophages *in*
 36 *situ*, which is likely a widespread and underestimated type of biocontrol.
 37 This assay is anticipated as a starting point for harnessing the bacterial

**potential in limiting the diffusion of harmful viruses within environments
such as gut or water.**

Introduction

For any type of bacteria, the presence of viruses may present a significant opportunity for feeding. Indeed, viruses represent 0.2 gigatons of carbon on Earth(Bar-On *et al.*, 2018). For example, the major capsid protein of the T4-like bacteriophage family is one of the most prevalent proteins in the biosphere(Comeau and Krisch, 2008). Therefore phages represent a major potential carbon source into which bacteria may tap. Furthermore, viruses are also a potential source of phosphorus(Jover *et al.*, 2014).

No bacterium preying on viruses have been described even though bacterial extracellular proteases are able to degrade certain bacteriophages in anaerobic wastewater treatment plants, in pure cultures(Mondal *et al.*, 2015) and in soil(Nasser *et al.*, 2002). In seawater, the only reported biotic pressure arises from marine ciliates that have been co-incubated with viruses and bacteria(Gonzalez and Suttle, 1993). This observation is also supported by the recent discovery of viral DNA in free-living eukaryotic cells(Brown *et al.*, 2020).

Here, we show that specific bacteria can indeed degrade T4 bacteriophages *in situ*, and we confirm this observation in pure culture.

Results

When searching for bacteriophage consumption activity, sludge from wastewater treatment plants is a relevant microbiota to investigate, as it boasts a high degradation capacity. In this work, the stable isotope probing method was applied by adding 2.2×10^{10} ^{13}C -labelled T4 phages to 200 μl of sludge corresponding to 10^8 bacteria cells. The increase in density of the bacterial DNA of the ^{12}C control bottle was then measured after the enumerated T4 phages decreased by 99%.

T4 phages were assimilated by bacteria. About 41% of the ^{13}C atoms initially present in T4 phages were accounted for in the bacterial biomass. However, only nine out of the 4046 microbial species - or more accurately Amplicon Sequence Variant (ASVs) - were labelled by the ^{13}C initially contained in the T4 phages (Fig. 1), thus suggesting that the incorporation of T4 phage is not a widespread ability. This incorporation generated growth, since the total biomass increased 2 fold after 24h concomitantly with the disappearance of the T4 phages.

The two main degraders of T4 phages were ASV1 (*Aeromonas sp.*) and ASV2 (*Tolumonas sp.*), which accounted for 5% and 29% of ^{13}C atoms found in the bacterial biomass respectively. Both belong to the *Aeromonadaceae* family and exhibit strong growth rates. Indeed, both rose from undetectable levels to 51% of the biomass, while the density of their DNA increased because they incorporated ^{13}C atoms from the isotopically

labelled T4 phages. For example, the 2×10^6 *Aeromonas* cells present after 24h contained 85% of ^{13}C atoms in their DNA whose density shifted from 1.72 g/mL to 1.75 g/mL in the bottle with ^{13}C -labeled T4 phages. The 16S rRNA sequences assigned to *Aeromonas* represented 19% and 8% of the total reads in the ^{12}C and the ^{13}C bottles respectively, thus revealing a consistent growth from initially undetectable levels.

In addition to the *Aeromonadaceae* family, two species (ASV12 and 21) belonging to the Ignavibacteriales PHOS-HE36 family, although labelled with medium strength (49 and 71%) and negligible growth (0 and 5.87×10^6 synthesized cells respectively), still gathered 5% of the ^{13}C atoms. The last 5 species with significant DNA density shifts (ASV7, 9, 20, 67 and 79) accounted for the remaining 1% of ^{13}C atoms but their weak labelling level may have resulted from indirect labelling.

To confirm the quality of *Aeromonas sp.* as a predator of T4 phages, an *Aeromonas*-selective medium was used for retrieving an *Aeromonas* colony from the initial sludge and called it *Aeromonas_isolate_007*. The analysis of the whole genome confirmed that this isolate belongs to an intermediate clade between *Aeromonas media* and *Aeromonas rivipollensis* species. *Aeromonas_isolate_007* was incubated with T4 phages as only substrate. Starting with 50 resting bacterial cells, the population reached 1.6×10^8 cells after 24 hours at 20°C while consuming 10^{11} T4 phages (Fig. 1C). No growth was observed when T4 phages were absent.

Aeromonas sp. can also capture T4 phages when their concentrations were comparable with environmental conditions: 7×10^4 T4 phages/mL decreased to 2×10^3 T4 phages/mL when incubated with *Aeromonas*_isolate_007 cells (Fig 1C). No decrease in the T7 phage has been observed in similar experiments where the T4 phage was replaced by the T7 phage.

Discussion

Aeromonas cells are present in virtually any environment (Janda and Abbott, 2010), including wastewater treatment plants where their abundance is around 0.1 % (Ye *et al.*, 2012).

Interestingly, *Aeromonas* cells have an S-layer (Noonan and Trust, 1997) associated with lipopolysaccharides (Sleytr *et al.*, 2014) and an outer membrane protein C, which are known to bind the T4 phages to the surface of *E. coli* cells (Islam *et al.*, 2019). Once captured at the surface, the phage is likely degraded by several extracellular enzymes, including DNase and protease (Janda, 1985). For example, metallo- and serine- proteases found in *Aeromonas* are involved in the degradation of large molecules such as albumin, earning the nickname of ‘Jack-of-all-trades’ due to this enzymatic versatility (Seshadri *et al.*, 2006). Finally, *Aeromonas* possesses transporters to uptake the resulting amino acids and peptides (Seshadri *et al.*, 2006).

Bacterial predation on bacteriophages is rich in consequences because bacteriophages are ultimate predators at the top of all food chains since they are not hunted. Indeed bacteriophage decay is mainly considered abiotic via adhesion to particulate material, chemical inactivation or degradation by solar radiation or passive grazing by flagellates (González and Suttle, 1993). In the oceans, this predation likely allows for the upper levels of the trophic chain carbon to access to the 7% of dissolved nitrogen, the 5% of phosphorus and the 1% of dissolved organic carbon contained in the viral particles (Jover *et al.*, 2014).

Furthermore, the diversity in bacteriophages could be partly related to the presence of phage-specific bacterial bacteriophage-hunters. Indeed, the bacterial predators of T4 phages do not appear to consume T7 bacteriophages. Therefore, brutal increase of a specific phage in the environment could be specifically controlled by a phage-eating bacterium, forming a killing-the-killer loop.

In conclusion, bacteria that are capable of eliminating specific viruses changes our vision of the food webs and represent a noteworthy avenue to explore to control harmful viruses such COVID-19, bacteriophages that disrupt dairy fermentations or rotaviruses causing diarrhoea.

Declarations

131 Ethics approval and consent to participate

132 Not Applicable

133 Consent for publication

134 Not Applicable

135 Availability of data and materials

136 High-throughput sequencing data have been deposited on NCBI
 137 (<https://www.ncbi.nlm.nih.gov/bioproject>) under accession number PRJNA650397
 138 and the genome of *Aeromonas_isolate_007* is accessible with the BioSample
 139 accession number SAMN17689348.

140 Competing Interests statement

141 The authors declare that they have no competing interests.

142 Funding

143 The work was funded by the French National Research Institute for Agriculture, Food
 144 and Environment (INRAE).

145 Contributions

J.J.G., O.Z., A.B. and N.H. designed and performed the experiments; J.J.G., O.Z., L.C. and M.A. performed computational experiments; J.J.G., O.Z., L.C., M.A. and M.A.P. contributed lineaging data and expertise; J.J.G. and O.Z. prepared the manuscript with assistance from all authors; J.J.G. and O.Z. supervised and directed the research.

Acknowledgments

We thank Caroline Achard for insights on bacteriophage capture, Martin Beaumont and Amira Bousleh for the qPCR of the internal standard. Electronic microscopy work has benefited from the facilities and expertise of MIMA2 MET – GABI, INRA, Agroparistech, 78352 Jouy-en-Josas, France. Sequencing was performed in collaboration with the GeT core facility, Toulouse, France (<http://get.genotoul.fr>) supported by France Génomique National infrastructure, funded as part of “Investissement d’avenir” program managed by the Agence Nationale pour la Recherche (contract ANR-10-INBS-09). Figure 1 was created by Biorender.

Authors' information

JJG directed the PhD of OZ in bacterial ecology in France. In a following postdoc, OZ learned stable-isotope probing in sludge in Australia and was granted a small project to study the adsorption of T2 phages in the light of electrostatics. The combination of SIP and phage expertise turned out useful to answer JJG’s question about the predation of phages together with AB, which has expertise in producing purified isotopically-labeled

phages and MAP, which could confirm the results in pure culture with JJG since an isolate was successfully isolated from the sample. MA's expertise in genome annotation was put to use to speculate about the mechanisms involved in the capture, digestion and assimilation of T4 phages by bacteria.

References

- Bar-On YM, Phillips R, Milo R (2018). The biomass distribution on Earth. *Proceedings of the National Academy of Sciences* **115**: 6506-6511.
- Brown JM, Labonté JM, Brown J, Record NR, Poulton NJ, Sieracki ME *et al* (2020). Single Cell Genomics Reveals Viruses Consumed by Marine Protists. *Front Microbiol* **11**: 524828.
- Comeau AM, Krisch HM (2008). The capsid of the T4 phage superfamily: the evolution, diversity, and structure of some of the most prevalent proteins in the biosphere. *Mol Biol Evol* **25**: 1321-32.
- Gonzalez JM, Suttle CA (1993). Grazing by Marine Nanoflagellates on Viruses and Virus-Sized Particles - Ingestion and Digestion. *Marine Ecology-Progress Series* **94**: 1-10.
- González JM, Suttle CA (1993). Grazing by marine nanoflagellates on viruses and virus-sized particles: ingestion and digestion. *Marine Ecology Progress Series* **94**: 1-10.
- Islam MZ, Fokine A, Mahalingam M, Zhang Z, Garcia-Doval C, van Raaij MJ *et al* (2019). Molecular anatomy of the receptor binding module of a bacteriophage long tail fiber. *PLOS Pathogens* **15**: e1008193.
- Janda JM (1985). Biochemical and exoenzymatic properties of *Aeromonas* species. *Diagnostic Microbiology and Infectious Disease* **3**: 223-232.

- 188 Janda JM, Abbott SL (2010). The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clin*
189 *Microbiol Rev* **23**: 35-73.
- 190 Jover LF, Effler TC, Buchan A, Wilhelm SW, Weitz JS (2014). The elemental composition of virus
191 particles: implications for marine biogeochemical cycles. *Nat Rev Microbiol* **12**: 519-28.
- 192 Mondal T, Rouch DA, Thurbon N, Smith SR, Deighton MA (2015). Factors affecting decay of
193 *Salmonella* Birkenhead and coliphage MS2 during mesophilic anaerobic digestion and air drying
194 of sewage sludge. *J Water Health* **13**: 459-72.
- 195 Nasser AM, Glozman R, Nitzan Y (2002). Contribution of microbial activity to virus reduction in
196 saturated soil. *Water Res* **36**: 2589-95.
- 197 Noonan B, Trust TJ (1997). The synthesis, secretion and role in virulence of the paracrystalline
198 surface protein layers of *Aeromonas salmonicida* and *A. hydrophila*. *FEMS Microbiol Lett* **154**: 1-
199 7.
- 200 Seshadri R, Joseph SW, Chopra AK, Sha J, Shaw J, Graf J *et al* (2006). Genome Sequence of
201 *Aeromonas hydrophila* ATCC 7966^T: Jack of All Trades. *Journal of Bacteriology* **188**:
202 8272-8282.
- 203 Sleytr UB, Schuster B, Egelseer E-M, Pum D (2014). S-layers: principles and applications. *FEMS*
204 *Microbiology Reviews* **38**: 823-864.
- 205 Ye L, Zhang T, Wang T, Fang Z (2012). Microbial structures, functions, and metabolic pathways in
206 wastewater treatment bioreactors revealed using high-throughput sequencing. *Environ Sci*
207 *Technol* **46**: 13244-52.

Figure 1. Bacterial growth on T4 phages. A; Identification of ^{13}C -labeled bacteria: the ^{13}C -labeled T4 (red) were incubated with a microbial community of wastewater treatment plant in the same conditions as the ^{12}C control (blue). **B; Bacteria present in each sample:** the barplots shows the growth of each ASVs based on 16S rDNA copies, detailing the nine bacteria assimilating T4 phages. **C; The density plots show the shift in density for ASV1 (*Aeromonas* sp.),** which was abundant in both bottles after 24h. **D; ^{13}C labelling level after 24h:** the ^{13}C -labeling level (left Y-axis), computed from the individual density shifts (right Y-axis), is reported against the initial abundance of the ASVs (X-axis). The volume of the spheres represents the abundance of each ASV in the ^{13}C bottle. **E; *Aeromonas* sp. growth on T4 phages:** *Aeromonas*_isolate_007 grew on T4 phages as sole carbon and nitrogen source. When a few *Aeromonas* cells were incubated with 10^{11} T4 phages, the colony forming units (brown) increased while the plaque-forming units (blue) decreased.

228