Genome wide association mapping of *Pyrenophora teres* f. *maculata* and *Pyrenophora teres* f. *teres* resistance loci utilizing natural Turkish wild and landrace barley populations

Shaun J. Clare^{*,1}, Arzu Çelik Oğuz^{†,1}, Karl Effertz^{*}, Roshan Sharma Poudel[‡], Deven See^{§,**}, Aziz Karakaya[†] and Robert S. Brueggeman^{*}

^{*} Department of Crop and Soil Sciences, Washington State University, Pullman, WA, United States, 99163

[†] Department of Plant Protection, Faculty of Agriculture, Ankara University, Dışkapı, Ankara, Turkey, 06110

[‡] Sygenta Seed Inc., Durham, NC, United States, 27709

[§] Wheat Health, Genetics and Quality Research Unit, Agricultural Research Service, U.S. Department of Agriculture, Pullman, WA, United States, 99163

^{**} Department of Plant Pathology, Washington State University, Pullman, WA, United States, 99163

¹These authors have contributed equally to this work and share first authorship

Correspondence: R. S. Brueggeman, Department of Crop and Soil Science, Johnson Hall Room 215, Washington State University, Pullman, WA, 99164-6420, +1 509 335 3644, bob.brueggeman@wsu.edu

Running Title: Turkish barley association mapping

Key Words: Barley landrace, wild barley, *Hordeum spontaneum*, *Pyrenophora teres* f. *teres*, *Pyrenophora teres* f. *maculata*, resistance, susceptibility, GWAS

ABSTRACT

Unimproved landraces and wild relatives of crops are sources of genetic diversity that were lost post domestication in modern breeding programs. To tap into this rich resource, genome wide association studies in large plant genomes have enabled the rapid genetic characterization of desired traits from natural landrace and wild populations. Wild barley (*Hordeum spontaneum*), the progenitor of domesticated barley (*H. vulgare*), is dispersed across Asia and North Africa, and has co-evolved with the ascomycetous fungal pathogens *Pyrenophora teres* f. *teres* and *P. teres* f. *maculata*, the casual agents of the diseases net form of net blotch and spot form of net blotch, respectively. Thus, these wild and local adapted barley landraces from the region of origin of both the host and pathogen represent a diverse gene pool to identify new sources of resistance, due to millions of years of co-evolution. The barley - *P. teres* pathosystem is governed by complex genetic interactions with dominant, recessive, and incomplete resistances and susceptibilities, with many isolate-specific interactions. Here we provide the first genome wide association study of wild and landrace barley from the Fertile Crescent for resistance to both forms of *P. teres*. A total of 14 loci, four against *P. teres* f. *maculata* and ten against *Pyrenophora teres* f. *teres*, were identified in both wild and landrace populations, showing that both are genetic reservoirs for novel sources of resistance. We also highlight the importance of using multiple algorithms to both identify and validate additional loci.

INTRODUCTION

Net blotch, caused by the ascomycetous fungal pathogen *Pyrenophora teres* (anamorph: *Drechslera teres*) is an economically important disease of barley worldwide. *P. teres* f. *teres* (*Ptt*) incites the net form of net blotch (NFNB) and *P. teres* f. *maculata* (*Ptm*) incites the spot form of net blotch (SFNB) (Smedegård-Petersen, 1971). Both forms are responsible for large crop losses that typically range between 10-40% when susceptible cultivars are grown, however, under conducive environmental conditions losses can reach 100% (Piening and Kaufmann 1969; Mathre 1982; Moya *et al.* 2018). At least one form of the disease has been reported from all barley growing regions and, in many regions, both forms are present with annual fluctuation in predominance. This presents challenges to breeders, as both *Ptt* and *Ptm* interact with host resistance/susceptibility genes differentially, thus are considered distinct and treated as different diseases when breeding for resistance. However, as further characterization of resistant/susceptibility loci continues, overlaps in host-pathogen genetic interactions in both pathosystems are becoming more prevalent.

Both *Ptt* and *Ptm* occur as genetically distinct populations and can be separated in the field based on lesion morphology. Although these two forms can be hybridized under laboratory conditions (Campbell and Crous 2003), hybridization under field conditions is extremely rare (Campbell *et al.* 2002; McLean *et al.* 2014; Akhavan *et al.* 2015; Çelik Oğuz *et al.* 2018; Poudel *et al.* 2019). However, both forms of *P. teres* undergo form specific sexual as well as asexual reproduction (Karakaya *et al.* 2004; Serenius *et al.* 2005; Akhavan *et al.* 2015; Çelik Oğuz *et al.* 2018, 2019b; Poudel *et al.* 2019). The complex nature of this reproduction system poses serious evolutionary risks for resistance breeding as populations contain diverse effector repertoires

that, in different combinations, can rapidly overcome deployed resistances (McDonald and Linde 2002). Yet, the use of resistant barley cultivars is the most environmentally friendly and economically feasible method of NFNB and SFNB control (Robinson and Jalli 1996; Afanasenko *et al.* 2009).

Wild barleys and barley landraces are important sources of resistance against diverse biotic and abiotic stresses (Allard and Bradshaw 1964; Ceccarelli 1996; Yitbarek *et al.* 1998; Ellis *et al.* 2000; Jakob *et al.* 2014; Karakaya *et al.* 2016, 2020). Wild barley (*Hordeum spontaneum*) is known as the progenitor of modern-day barley (*H. vulgare*) and grows naturally in the Fertile Crescent, regions of south and southeastern Turkey, North Africa, and Southwest Asia (Harlan and Zohary 1966; Nevo 1992; Zohary and Hopf 2000). Barley landraces gave rise to modern barley varieties (Thomas *et al.* 1998) as they were subjected to natural and artificial selection for the last 10,000 years (Ceccarelli and Grando 2000). Within the barley center of origin, Turkey is located at the ancestral hub of barley diversification regions including the Mediterranean, Horn of Africa, and the Tibetan Plateau (Muñoz-Amatriaín *et al.* 2014; Poets *et al.* 2015), with diverse barley landraces still used by Turkish farmers (Helbaek 1969; Kün 1996; Pourkheirandish and Komatsuda 2007; Ergün *et al.* 2017). Turkey is also at the center of origin of the *P. teres* pathogens, *Ptt* and *Ptm*. Barley lines can show differential responses to either or both forms of net blotch due to distinct yet complex genetic interactions with each form. Many barley genotypes may be resistant to the majority of isolates of one form yet susceptible to the alternate form of most isolates (Bockelman *et al.* 1983; Grewal *et al.* 2012). Thus, when breeding for resistance, the two forms of net blotch are treated as separate diseases (Liu *et al.* 2011; Usta *et al.* 2014; Yazıcı *et al.* 2015).

85 In barley, genetic resistance to *P. teres* was first reported in 1928 by Geschele. Since both forms
86 of net blotch were not described at that time, it was assumed that the net form was used. In
87 the first studies to genetically characterize barley – *P. teres* interactions, Tifang was the
88 resistant parent in the Tifang × Atlas cross (Schaller 1955). Mode and Schaller (1958) identified
89 the resistance genes *Pt₁*, *Pt₂* and *Pt₃* segregating in this cross. Bockelman et al. (1977) revised
90 the naming of *Pt₁*, *Pt₂* and *Pt₃* resistance genes and described the *Rpt1* (*Pt₁* and *Pt₂*), *Rpt2*
91 (novel) and *Rpt3* (*Pt₃*) loci on chromosomes 3H, 1H and 2H, respectively. Further studies
92 identified *Rpt4* on chromosome 7H (Williams et al. 1999, 2003), *Rpt5* on chromosome 6H, *Rpt6*
93 on chromosome 5H (Manninen et al. 2006), *Rpt7* on chromosome 4HL and *Rpt8* on
94 chromosome 4HS (Franckowiak and Platz 2013), however, there are over 340 QTLs previously
95 identified (Clare et al. 2020). Manninen et al. (2006) reclassified the locus *Pt_a* as *Rpt5* on
96 chromosome 6H, and subsequently, three genes/alleles have been characterized at the locus
97 (*Rpt5.f*, *Spt1.k*, *Spt1.r*) as dominant resistance or susceptibility genes (Franckowiak and Platz
98 2013; Richards et al. 2016). In multiple barley – *Ptt* genetic interaction studies, it has been
99 shown that the *Rpt5* locus is the most important resistance/susceptibility locus in this system.
100 This complex locus putatively contains multiple resistance as well as susceptibility genes that
101 have been characterized in diverse barley – *P. teres* interactions from around the world (Clare
102 et al. 2020). Because the *Rpt5* locus also shows dominant susceptibility in certain barley lines,
103 additional alleles were designated *Susceptibility to P. teres 1* (*Spt1*) by Richards et al., (2016).
104 Further, high resolution genetic mapping and positional cloning efforts have identified *Rpt5* and
105 *Spt1* candidate genes and functional validation is underway (Brueggeman et al. 2020).

Resistance to *Ptm* originally appeared to be less complicated when compared to *Ptt* due to the presence of three major loci. These three loci were identified as *Rpt4* on chromosome 7H (Williams *et al.* 1999, 2003), *Rpt6* on chromosome 5HS (Manninen *et al.* 2006) and *Rpt8* on chromosome 4HS (Friesen *et al.* 2006; Franckowiak and Platz 2013). To date, over 140 QTLs have been reported to be implicated in the *Ptm* reaction, which have been collapsed into 36 unique loci, five of which are specific to *Ptm* and the rest showing some degree of overlap with known *Ptt* loci (Clare *et al.* 2020). These five unique loci that are specific to the *Ptm* interaction are *SFNB-3H-78.53* on chromosome 2H (Burlakoti *et al.* 2017), *QRptm-4H-120-125* on 4H (Tamang *et al.* 2019), *QRptts-5H-106.00* on 5H (Adhikari *et al.* 2019), *QRptm7-3* on 7H (Wang *et al.* 2015) and *QRptm7-6/QRptm-7H-119-137* on 7H (Wang *et al.* 2015; Tamang *et al.* 2019). Considering that all currently designated resistance/susceptibility loci except for *Rpt2* (only implicated in the *Ptt* interaction) have now been implicated in both *Ptm* and *Ptt* interactions (Clare *et al.* 2020), it is with some caution that it can be concluded that host-pathogen genetic interactions with the two forms and barley should be considered distinct. Thus, for both forms, with the exception of *Rpt6*, numerous researchers have described synonyms of all loci (Clare *et al.* 2020).

Multiple genome wide association mapping studies (GWAS) have investigated NFNB resistance in barley (Wonneberger *et al.* 2017b; Richards *et al.* 2017; Amezrou *et al.* 2018; Adhikari *et al.* 2019, 2020; Rozanova *et al.* 2019; Daba *et al.* 2019; Novakazi *et al.* 2019). A large proportion of the resistance markers associated with NFNB resistance have been localized to the centromeric region of barley chromosome 6H (Richards *et al.* 2017). In GWAS for SFNB resistance, 29 (Wang *et al.* 2015), 27 (Tamang *et al.* 2015), 11 (Burlakoti *et al.* 2017) and one (Daba *et al.* 2019)

unique genomic loci were identified. Four important QTLs (*QRptm7-4*, *QRptm7-6*, *QRptm7-7* and *QRptm7-8*) were mapped into a region covering the *Rpt4* locus on chromosome 7HS (Wang *et al.* 2015). Burlakoti *et al.* (2017) identified a new and important QTL on chromosome 2HS that was predominately found in 6-rowed barley lines as compared to 2-rowed. Daba *et al.* (2019) also defined a new QTL on chromosome 6H associated with *Ptm* susceptibility in a large number of genotypes, which is a common mechanism in inverse gene-for-gene interactions with this pathogen. Vatter *et al.* (2017) performed nested association mapping for NFNB and described further interactions at the important *Rpt5* / *Spt1* locus. In barley, the newest approach to identify MTAs with *Ptt* resistance is exome QTL-seq. This approach identified a large number of MTAs on chromosomes 3H and 6H when analyzing resistant and susceptible bulks (Hisano *et al.* 2017). The resistance status of wild barley genotypes and barley landraces to *P. teres* has been reported by several research groups worldwide (Legge *et al.*, 1996; Lakew *et al.*, 1995; Endresen *et al.*, 2011; Silvar *et al.*, 2010; Neupane *et al.*, 2015; Fetch *et al.*, 2003; Jana and Bailey, 1995; Sato and Takeda, 1997; Çelik Oğuz *et al.*, 2017, 2019a, 2019c). However, molecular mapping studies of *P. teres* resistance in wild and landrace barleys have been limited (Yun *et al.* 2005; Vatter *et al.* 2017; Adhikari *et al.* 2019; Gyawali *et al.* 2019, 2020). In this study, four novel loci representing resistance to NFNB were mapped in Turkish wild barley and landraces. This study highlights the importance of surveying wild and unimproved barley lines for sources of resistance that may have been lost during domestication and modern breeding. These studies, focused on diversity in the barley primary germplasm pool, will provide new sources of resistance and associated markers to aid in deploying robust resistances against NFNB and SFNB.

MATERIALS AND METHODS

BIOLOGICAL MATERIALS

A total of 295 barley accessions comprised of 193 landraces (*H. vulgare*) and 102 wild barley (*H. spontaneum*) genotypes which were collected from different growing regions of Turkey and maintained at the Field Crops Central Research Institute and Department of Plant Protection, Faculty of Agriculture, Ankara University located in Ankara, Turkey were utilized in the analyses (Çelik Oğuz *et al.* 2017, 2019a). Three virulent *Ptm* isolates (GPS263, 13-179 and 13-167) and three *Ptt* isolates (UHK77, GPS18 and 13-130) collected from different provinces of Turkey (Çelik Oğuz and Karakaya 2017) were used for the phenotypic assessment of the 295 barley accessions.

PATHOGEN ASSAY PHENOTYPING

Phenotyping of the wild barley (*H. spontaneum*) genotypes and barley landraces (*H. vulgare*) were accomplished according to methods outlined in Çelik Oğuz *et al.* (2017) and Çelik Oğuz *et al.* (2019a).

PCR-GBS LIBRARY PREPERATION AND GENOTYPING BY SEQUENCING

Two independent custom PCR-GBS SNP marker panels containing 365 (Panel 1) (Sharma Poudel *et al.* 2018; Tamang *et al.* 2019) and 1,272 (Panel 2) (Ruff *et al.* 2020) barley SNP markers were used to genotype 295 barley accessions. Marker primers were divided into six and three total primer pools for Panel 1 and 2 for PCR amplification, respectively. PCR amplification and barcoding reactions were performed as described by Sharma Poudel *et al.* (2018) and Ruff *et al.*

(2020). Briefly, nine primary PCR amplification reactions were performed per sample. Following amplification, equal volumes of primary PCR products were pooled into 96-well plates with each well containing all amplified markers for a given DNA sample. Next, barcoding PCR reactions were performed with a universal barcoding reverse primer and unique forward barcoding primers for each sample. Following barcoding reactions, samples were pooled and purified. A final PCR reaction was performed using sequencing primers to ensure the barcoding reaction was successful. Samples taken before and after the final amplification were run on agarose gel to verify the appropriate product size and amplification. Quantification of the barcoded libraries were performed using the Qubit dsDNA HS assay kit (Life Technologies, Carlsbad, CA). Enrichments were carried out using the Ion OneTouch™ 2 System (Panel 1) and the Ion PI™ Hi-Q Sequencing 200 kit on the Ion Chef (Panel 2). Finally, samples were sequenced on the Ion Torrent Personal Genome Machine™ (Panel 1) and the Ion Proton™ (Panel 2) Systems using two Ion 318™ chips and 3 Ion PI™ chips, respectively, following the manufacturer's standard protocols. The resulting marker panels were collapsed to eliminate duplicated markers from each panel. The Morex reference map (Beier *et al.* 2017; Mascher *et al.* 2017; Monat *et al.* 2019) and iSelect consensus map (Muñoz-Amatriaín *et al.* 2014) were downloaded from The Triticeae Toolbox (T3) barley database (<https://triticeaetoolbox.org/barley/>). The Morex reference map was used to determine the absolute marker position, whereas markers not included in the Morex reference map were estimated based on the genetic position of the iSelect consensus map relative to the flanking markers.

IMPUTATION, FILTERING AND LINKAGE DISEQUILIBRIUM

Due to the heterozygosity present in the natural population, heterozygous calls (5.23%) were included in the analysis. Accessions and markers with more than 30% missing data were removed from analysis, resulting in 282 barley accessions and 530 markers. Missing data was imputed using LinkImpute, which uses a linkage disequilibrium k -nearest neighbor imputation (LD-kNNi) method (Money *et al.* 2015) in Trait Analysis by aSSociation, Evolution, and Linkage (TASSEL) 5.2.60 (Bradbury *et al.* 2007). Markers with a minor allele frequency of <0.05 were included in the analysis but were treated with caution based on best practice from the Genomic Association and Prediction Integrated Tool (GAPIT) 3.0 user manual. Linkage disequilibrium was calculated in TASSEL using a window size of 50 markers and an R^2 threshold of 0.8 resulting in 522 markers.

POPULATION STRUCTURE, KINSHIP MATRICES AND MODEL ALGORITHMS

Population structure was accounted for using STRUCTURE analysis and principle component analysis. A total of 522 markers were used for analysis of population structure. The software STRUCTURE v2.3.4 (Pritchard *et al.* 2000) was used to estimate population structure of the barley panel to create a population structure matrix (Q) to be used as a covariate. To determine the optimal number of subpopulations, an admixture ancestry model was used with a burnin of 10,000, followed by 25,000 Monte Carlo Markov Chain (MCMC) replications for $k=1$ to $k=10$ with ten iterations. STRUCTURE HARVESTER (Earl and vonHoldt 2012) was used to identify the optimal number of subpopulations using the Δk method (Evanno *et al.* 2005). The optimal k value was subsequently used to run a new STRUCTURE analysis using a burnin of 100,000 followed by 100,000 MCMC replications. An individual was deemed to be part of a population if the membership probability was >0.8 (Richards *et al.* 2017). Individuals that did not achieve a

value of 0.8 were deemed to have admixture ancestry. The final Q matrix was used as a fixed covariate in association models. Principle component analysis was conducted in R 3.6.3 using GAPIT 3.0 (Wang and Zhang 2018) with default settings. Principle components explaining at least 25% ($PC1$) and 50% ($PC5$) were used for further analysis. A naïve model using only genotypic (Supplemental File 1 and 2) and phenotypic data (Supplemental File 3) and an additional three fixed effect models accounting for population structure (Q [Supplemental Figure 4], $PC1$, and $PC5$) were all performed using the GLM method.

For initial discovery of the most appropriate method to account for random effect in the model, a kinship matrix (K) was constructed using the EMMA (Kang *et al.* 2008), Loiselle (Loiselle *et al.* 1995) and VanRaden (VanRaden 2008) algorithms in GAPIT 3.0 (Wang and Zhang 2018) with the MLM model (Yu *et al.* 2006). Based on these results, the EMMA derived K matrix (Supplemental File 5) was identified as the most powerful and used for subsequent analysis of mixed models for all isolates that included CMLM (Zhang *et al.* 2010), ECMLM (Li *et al.* 2014) and MLMM (Segura *et al.* 2012). Lastly, SUPER (Wang *et al.* 2014), FarmCPU (Liu *et al.* 2016) and BLINK (Huang *et al.* 2019) algorithms that reconstruct the kinship matrix were used for a total of nine random effect models. Due to the similarity of results, the MLM and MLMM methods were not used in further analysis. Mixed models included combinations to account for population structure (Q , $PC1$, and $PC5$), kinship (EMMA K) and algorithm methods (CMLM, ECMLM, SUPER, FarmCPU and BLINK) for a total of 15 mixed models per isolate. The mean-squared deviation (MSD) was calculated for each model (Supplemental File 6, Mamidi *et al.* 2011), however visual inspection of QQ plots were done to ensure the model was a good fit. This method was employed as models with the lowest MSD model often had highly correlated observed and

expected $-\log_{10}(p)$ values yielding zero significant markers. A Bonferroni correction was calculated at an α level of 0.01 and 0.05 for a $-\log_{10}(p)$ threshold of 4.72 and 4.02, respectively. Final Manhattan and QQ plots were generated with R 3.6.3 package *CMplot* 3.5.1 (<https://github.com/YinLiLin/R-CMplot>).

QTL IDENTIFICATION

Absolute marker positions were extracted for significant marker trait associations from the Morex reference genome (Beier *et al.* 2017; Mascher *et al.* 2017; Monat *et al.* 2019) and compared to collapsed *P. teres* loci (Clare *et al.* 2020). Significant markers were declared distinct from previously identified loci, i.e. novel, if the nearest neighboring marker that was closer to previously reported loci was not significant or if the gap to currently delimited locus exceeded 10 Mbp in physical distance when no closer marker was present.

RESULTS

PHENOTYPIC ANALYSIS

Wild barley was found to be statistically (Wilcoxon rank sum test) more resistant to *Ptm* isolate 13-179 and *Ptt* isolates GPS18 and UHK77, whereas landrace barley was shown to be statistically more resistant to *Ptm* isolate GPS263 and *Ptt* isolate 13-130 (Figure 1). There was no significant difference between landrace and wild barley to *Ptm* isolate 13-167. Despite UHK77 being statistically different between landrace and wild barley, no significant MTAs were found.

MARKER PANEL ANALYSIS

Using a 30% missing data threshold, a total of 282 of the 295 barley accessions phenotyped were adequately genotyped. Using a similarity of individual matrix, all remaining individuals were unique. Similarly, 530 of the 598 collapsed markers were deemed to have sufficient coverage across barley accessions using a 30% missing data threshold and were used in the final analysis of the six *P. teres* isolates. To eliminate markers in linkage disequilibrium, an LD R^2 threshold of 0.8 and sliding window of 50 markers was used, resulting in 522 markers for use in the final analysis.

POPULATION STRUCTURE AND LINKAGE DISEQUILIBRIUM

STRUCTURE analysis identified an optimal k value = 2, with 89 and 118 individuals in subpopulations one and two. Subpopulation one consisted of wild accessions, whereas subpopulation two consisted of landraces. A total of 75 barley accessions comprised of both landrace and wild barley had population membership probabilities of less than 0.8 and were deemed to have an admixture ancestry. The first five principle components accounted for 29.36, 9.4, 5.3, 4.0, and 3.7%, respectively, in the principle component analysis. Principle components were selected that accounted for at least 25% ($PC1$) and 50% ($PC5$) when eigenvalues were plotted on a cumulative scale.

ASSOCIATION MAPPING ANALYSIS

A total of 24 models were tested on each of the six *P. teres* isolates consisting of three *Ptm* and three *Ptt* isolates. The *Ptm* isolate 13-167 and *Ptt* isolate UHK77 contained no significant markers across all models tested. The *Ptm* isolates GPS263 and 13-179 contained one and three

276 significant markers, respectively. The *Ptt* isolates GPS18 and 13-130 both contained five
277 significant markers.

278 ***Ptm* Isolate GPS263:** Only one significant MTA was identified with *Ptm* isolate GPS263 on
279 chromosomes 5H based on the second version of the cv. Morex reference genome (Monat *et*
280 *al.* 2019). The SNP marker 12_20350 located on chromosome 5H at physical position
281 446449782 was identified at the $-\log_{10}(p)$ threshold = 4.72 in the K_{BLINK} . Marker 12_20350 is
282 embedded within the collapsed *NBP_QRptt5-1* locus (Wonneberger *et al.* 2017a; Clare *et al.*
283 2020) and was not previously shown to be associated with *Ptm* interactions.

284 ***Ptm* Isolate 13-179:** The three significant MTAs identified with *Ptm* isolate 13-179 were located
285 on chromosomes 3H, 4H and 5H based on the second version of the Morex reference genome
286 (Monat *et al.* 2019). The marker 11_20866 located on chromosome 3H (physical position
287 153156749), embedded within the collapsed *QRptms3-2* locus (Wang *et al.* 2015; Wonneberger
288 *et al.* 2017a; Koladia *et al.* 2017; Burlakoti *et al.* 2017; Vatter *et al.* 2017; Rozanova *et al.* 2019;
289 Daba *et al.* 2019; Novakazi *et al.* 2019; Clare *et al.* 2020), was identified at the $-\log_{10}(p)$
290 threshold = 4.02 in the K_{FarmCPU} and $\text{PC1}+K_{\text{FarmCPU}}$ models. The marker 11_10510 is located on
291 chromosome 4H at position 603258307 and was identified at the $-\log_{10}(p)$ threshold = 4.72 in
292 the K_{BLINK} and K_{SUPER} models and at the $-\log_{10}(p)$ threshold = 4.02 in the PC5_{GLM} , $\text{Q}+K_{\text{SUPER}}$, and
293 $\text{PC1}+K_{\text{SUPER}}$ models. In addition, the 11_10510 marker almost met the significance threshold
294 using the K_{FarmCPU} model. The 11_10510 marker is embedded within the *Rpt8* locus (Friesen *et*
295 *al.* 2006; Tamang *et al.* 2015; Richards *et al.* 2017; Vatter *et al.* 2017; Daba *et al.* 2019; Clare *et*
296 *al.* 2020). Lastly, the marker SCRI_RS_160332 is located on chromosome 5H at position

297 474799503, ~3.0 Mbp distal to the *Qrptts-5HL.1* locus (Richards *et al.* 2017). SCRI_RS_160332
 298 was identified with the K_{FarmCPU} model at the $-\log_{10}(p)$ threshold = 4.02.

299 **Ptt Isolate GPS18:** The five significant MTAs identified using *Ptt* isolate GPS18 were distributed
 300 across chromosomes 1H, 6H and 7H. The first marker 11_10176 located on chromosome 1H
 301 (position 397791042) was identified in the K_{FarmCPU} model at the $-\log_{10}(p)$ threshold = 4.02. The
 302 11_10176 marker is located 13.6 Mbp distal to the collapsed *NBP_QRptt1-1* (Wonneberger *et al.*
 303 *al.* 2017a) and 18.2 Mbp proximal to the QTL identified at 57.3-62.8 cM by Rozanova *et al.*
 304 (2019). The marker 11_20754 located on chromosome 1H (position 483805599) was identified
 305 in the K_{BLINK} and K_{FarmCPU} models at the $-\log_{10}(p)$ threshold = 4.72 and 4.02, respectively. The
 306 11_20754 marker is embedded within the *QPt.1H-1* (Vatter *et al.* 2017; Clare *et al.* 2020) locus
 307 along with *QRptts-1H-92-93* (Amezrou *et al.* 2018) and a QTL from Tamang *et al.* (2015). The
 308 third marker 12_31282 located on chromosome 7H (position 617741299), is embedded within
 309 the collapsed *QTL_{UHS-7H}* locus (König *et al.* 2013; Tamang *et al.* 2015, 2019; Wonneberger *et al.*
 310 2017a; Richards *et al.* 2017; Martin *et al.* 2018; Novakazi *et al.* 2019; Clare *et al.* 2020). The
 311 12_31282 MTA was identified using the K_{FarmCPU} model at the $-\log_{10}(p)$ threshold = 4.72. The
 312 fourth marker, 11_20972 on chromosome 6H (position 539551443) is embedded within the
 313 collapsed *AL_QRptt6-2* locus (Afanasenko *et al.* 2015; Wonneberger *et al.* 2017b; Vatter *et al.*
 314 2017; Amezrou *et al.* 2018; Clare *et al.* 2020). Marker 11_20972 was identified using the K_{BLINK}
 315 model at the $-\log_{10}(p)$ threshold = 4.02 and also nearly met the significant threshold using the
 316 K_{FarmCPU} model. The last significant marker, 12_30545 located on chromosome 7H (54934072) is
 317 embedded within the collapsed *QNFBAPR.AI/S-7Ha* locus (König *et al.* 2013; Tamang *et al.*
 318 2015; Wonneberger *et al.* 2017b; Vatter *et al.* 2017; Amezrou *et al.* 2018; Daba *et al.* 2019;

Novakazi *et al.* 2019; Clare *et al.* 2020). Marker 12_20545 was identified at the $-\log_{10}(p)$ threshold = 4.02 using the K_{FarmCPU} model. Two additional markers, 12_30250 and 12_111942 located on chromosome 3H and 4H, respectively, were almost significant at the $-\log_{10}(p)$ threshold = 4.02 in the K_{FarmCPU} model. The 12_30250 marker is embedded within the collapsed *QRpts3La* locus (Raman *et al.* 2003; Lehmensiek *et al.* 2007; Liu *et al.* 2015; Tamang *et al.* 2015, 2019; Richards *et al.* 2017; Burlakoti *et al.* 2017; Vatter *et al.* 2017; Daba *et al.* 2019). The 12_11104 marker is located 1.8 Mbp distal from the *Rpt8* locus (Tamang *et al.* 2015; Vatter *et al.* 2017; Clare *et al.* 2020) and 4.4 Mbp proximal to the *QRptm-4H-120-125* locus (Tamang *et al.* 2015, 2019).

Ptt isolate 13-130: The five significant MTAs identified using *Ptt* isolate 13-130 were distributed across chromosomes 2H, 3H, 6H and 7H. The first marker, 12_11452 located on chromosome 2H (position 34275254) was identified with the K_{BLINK} and K_{FarmCPU} models at the $-\log_{10}(p)$ threshold = 4.02. The marker 12_11452 has a minor allele frequency less than 5%, however the marker is embedded within the collapsed *SFNB-2H-8-10* locus (Tamang *et al.* 2015, 2019; Wonneberger *et al.* 2017a; Amezrou *et al.* 2018; Adhikari *et al.* 2019; Clare *et al.* 2020). The marker 11_20968, located on chromosome 3H (position 19966889), was identified in the K_{FarmCPU} model at the $-\log_{10}(p)$ threshold = 4.72. The marker 11_20968 is located 7.6 Mbp distal to the boundary of the collapsed *QPt.3H-1* locus (Vatter *et al.* 2017; Rozanova *et al.* 2019; Daba *et al.* 2019) and 24 Mbp proximal to the boundary of the collapsed *Rpt-3H-4* locus (Tamang *et al.* 2015; Richards *et al.* 2017; Daba *et al.* 2019; Novakazi *et al.* 2019; Clare *et al.* 2020). The marker 12_10662, also located on chromosome 3H (position 553445025), was identified using the K_{FarmCPU} and the $Q+K_{\text{FarmCPU}}$ models at the $-\log_{10}(p)$ threshold = 4.72 and 4.02, respectively.

The marker is located 4.6 Mbp distal to boundary of the collapsed *QRpts3La* locus (Liu *et al.* 2015; Tamang *et al.* 2015, 2019; Wonneberger *et al.* 2017a; Richards *et al.* 2017; Burlakoti *et al.* 2017; Vatter *et al.* 2017; Daba *et al.* 2019) and 5.0 Mbp from the boundary of collapsed *Rpt1* locus (Tamang *et al.* 2015; Burlakoti *et al.* 2017; Martin *et al.* 2018; Adhikari *et al.* 2019; Novakazi *et al.* 2019; Clare *et al.* 2020). The last two markers, 11_20714 and 11_11243, were both identified using the K_{FarmCPU} model at the $-\log_{10}(p)$ threshold = 4.02 and are located on chromosomes 6H (position 489619101) and 7H (position 601974526), respectively. The marker 11_20714 is located 4.1 Mbp proximal to the *QPt.6H-3* locus (Vatter *et al.* 2017) and 11_11243 is embedded within the collapsed *QRptm7-6* locus (Wang *et al.* 2015; Tamang *et al.* 2015, 2019; Wonneberger *et al.* 2017a; Clare *et al.* 2020).

ENRICHMENT ANALYSIS

Enrichment of either resistance or susceptibility alleles at each MTA were calculated for landraces and wild barley (Figure 2, Supplemental Figure 1). For *Ptm* isolate GPS263 and *Ptt* isolate 13-130, the landraces that were more resistant than the wild barley showed enrichment for the majority of the resistance alleles or a depletion of susceptibility alleles. For *Ptm* isolate 13-179 and *Ptt* isolate GPS18, the opposite was observed as the wild barley showed more resistance and enrichment for the majority of the resistance alleles or depletion of susceptibility alleles.

DISCUSSION

Both NFNB and SFNB are worldwide threats to barley production and recent evidence shows that both *Ptm* and *Ptt* have evolved to infect and threaten wheat production as well (Tóth *et al.* 2008; Mikhailova *et al.* 2010; Perelló *et al.* 2019). Wild barley and landraces from the origin of

cereal domestication represent a rich reservoir of net blotch resistance that could be integrated into elite varieties to help mitigate the threat. Analysis of the phenotypic responses of Turkish wild barley and landraces to regional *Ptm* and *Ptt* isolates showed evidence of landraces under selective pressures by the pathogen during domestication compared with wild barley as seen by the more compact distribution of the phenotypic scores (Table 1, 2, Figure 1). This demonstrates that wild barley harbors additional diversity for net blotch resistance that is not present in the landraces and could be exploited for barley variety development. This is corroborated by enrichment analysis that shows enrichment of the resistant haplotype for marker 11_20866 and 11_20754 in the wild barley lines (Figure 2, Supplemental Figure 1). However, the opposite is true for other loci such as the 11_20968 haplotype near the *QRpt-3H.1* locus, that is located approximately 20 Mbp distal of the domestication gene non-brittle rachis 1, *btr1* on the chromosome 3HS (Komatsuda *et al.* 2004; Wang *et al.* 2019). None of the wild barley accessions analyzed contained the resistance haplotype but it is enriched within landrace accessions (Figure 2, Supplemental Figure 1). The complete lack of the “resistance” marker 11_20968 haplotype in the wild barley could be explained by removal of the “susceptible” haplotype through a selective sweep within close proximity to the *btr1* region during domestication. These results show the importance of surveying both landraces and wild barley accessions since important resistance and/or susceptibility loci that interact in the barley – *P. teres* pathosystem may have been lost or gained through domestication. We have found loci that are unique to wild barley or landraces indicating the importance of analyses of the entire primary barley germplasm pool to identify new sources of resistance for future breeding efforts.

To date only a handful of studies have utilized wild or landrace barley to map resistance loci using biparental populations (Metcalf *et al.* 1970; Bockelman *et al.* 1977; Manninen *et al.* 2000, 2006; Williams *et al.* 2003; Koladia *et al.* 2017) or association mapping (Tamang *et al.* 2015; Wonneberger *et al.* 2017a; Richards *et al.* 2017; Vatter *et al.* 2017; Amezrou *et al.* 2018; Adhikari *et al.* 2019; Daba *et al.* 2019; Novakazi *et al.* 2019). Only one study has incorporated both *Ptm* and *Ptt* (Daba *et al.* 2019) and zero have investigated the wild and landrace barley specifically present within the center of origin of the Fertile Crescent. Despite using a reduced marker set (Figure 3) which will reduce the amount of marker trait associations (MTAs) identified (Cui *et al.* 2020), a total of 14 unique MTAs have been identified using modern mapping algorithms (Figure 4). However, the two isolates for which no significant MTAs were identified may be due to the low marker density utilized in these analyses. Four of the MTAs were potentially novel and two that mapped to previously identified *Ptt* resistance loci that had not been reported to be involved in *Ptm* interactions. Additionally, while the remaining MTAs may not be novel, they may represent important alleles that could be incorporated into breeding programs. Thus, the GWAS identified an abundance of net blotch resistance found in this germplasm collected from the barley center of origin.

In this study barley chromosomes 3H and 7H contained the most MTAs with three, followed by two MTAs on chromosomes 1H, 5H, and 6H and one MTA on chromosomes 2H and 4H (Figure 4, Table 3). Of the 14 MTAs identified, four were identified against the *Ptm* isolates GPS263 and 13-179. The remaining ten MTAs were identified against *Ptt* isolates. When selecting the appropriate GWAS algorithm, the BLINK algorithm identified five MTAs, whereas the FarmCPU algorithm identified eleven MTAs, of which only two MTAs overlapped in both BLINK and

FarmCPU algorithms. The SUPER algorithm identified one MTA using *Ptm* isolate 13-179 but this was also identified by the BLINK algorithm. Investigating model selection, the kinship (*K*) model identified 14 MTA, whereas the mixed kinship and population structure (*Q+K*) model only identified one MTA, showing overlap with the *K* model and therefore no unique MTA. Although we would encourage higher marker saturation in future studies to confirm locus novelty, we would suggest continued best practice of testing multiple models (naïve, *K*, *Q*, *Q+K*), along with the addition of including all modern association mapping algorithms based on the different set of MTAs identified using BLINK and FarmCPU.

The 14 MTAs identified in this study were compared to the collapsed loci of Clare et al. (2020) and a similar strategy of determining novel loci was dependent on the significance of the nearest neighbor marker and previous incorporation into a locus. Using this strategy, we have identified four potentially novel loci (*QRpt-1H.1*, *QRpt-3H.1*, *QRpt-3H.3*, *QRpt-6H.1*) in the barley – *Ptt* interaction and two loci previously described only against *Ptt* that had not been previously identified in *Ptm* interactions (*QRpt-5H.1* and *QRpt-5H.2* corresponding to *NBP_QRptt5-1* and *Qrptts-5HL.1*, respectively). The novel loci were detected on chromosomes 1H, 3H and 6H. The *QRpt-1H.1* and *QRpt-6H.1* MTAs were associated in the interaction with *Ptt* isolate GPS18. *QRpt-1H.1* is proximally flanked by the previously reported QTLs *NBP_QRptt1-1* (Wonneberger et al. 2017a) and a QTL identified at 57.3-62.8 cM (Rozanova et al. 2019) at distances of 13.6 Mbp and 18.2 Mbp, respectively, to the closest boundary of the delimited region of the loci (Clare et al. 2020). Markers located 18.2 Mbp proximal and 7.3 Mbp distal to *QRpt-1H.1* were not significant and therefore the MTA was deemed novel. Similarly, the closest locus to *QRpt-6H.1* is *Rpt5/Spt1* (Steffenson et al. 1996; Manninen et al. 2000, 2006; Raman et

429 *al.* 2003; Read *et al.* 2003; Emebiri *et al.* 2005; Friesen *et al.* 2006; Abu Qamar *et al.* 2008;
430 Grewal *et al.* 2008, 2012; St. Pierre *et al.* 2010; Cakir *et al.* 2011; Gupta *et al.* 2011; O'Boyle *et*
431 *al.* 2014; Liu *et al.* 2015; Hisano *et al.* 2017; Islamovic *et al.* 2017; Koladia *et al.* 2017;
432 Wonneberger *et al.* 2017b; Richards *et al.* 2017; Vatter *et al.* 2017; Martin *et al.* 2018; Amezrou
433 *et al.* 2018; Adhikari *et al.* 2019, 2020; Rozanova *et al.* 2019; Daba *et al.* 2019; Novakazi *et al.*
434 2019) located 10.7 Mbp distal, however, a marker 28 Mbp distal to *QRpt-1H.1* and embedded
435 within *Rpt5/Spt1* was not significant, giving us reason to believe the locus is novel. The novel
436 loci *QRpt-3H.1*, *QRpt-3H.3* and *QRpt-6H.2* were all identified with *Ptt* isolate 13-130. The locus
437 *QRpt-3H.1* is flanked by QTLs located at 12.1-17.4 cM (Rozanova *et al.* 2019) proximal and 53.42
438 cM (Tamang *et al.* 2019) distal on the Morex POPSEQ map (Mascher *et al.* 2013, 2017),
439 equating to distances of 7.6 Mbp and 24 Mbp. The nearest neighbor markers are 5.2 Mbp
440 proximal and 3.2 Mbp distal and are not significant indicating that this is potentially a novel
441 locus. Similarly the *QRpt-3H.3* locus is flanked by *QRpts3La* (Raman *et al.* 2003; Lehmsiek *et*
442 *al.* 2007; Liu *et al.* 2015; Richards *et al.* 2017; Vatter *et al.* 2017; Daba *et al.* 2019) and *Rpt1*
443 (Bockelman *et al.* 1977; Graner *et al.* 1996; Richter *et al.* 1998; Raman *et al.* 2003; Cakir *et al.*
444 2003; Manninen *et al.* 2006; Lehmsiek *et al.* 2007; Martin *et al.* 2018; Adhikari *et al.* 2019;
445 Novakazi *et al.* 2019) approximately 4.6 Mbp proximal and 5 Mbp distal. However, the nearest
446 neighbor markers are not significant at 6.2 Mbp proximal and 1.3 Mbp distal. The last novel
447 locus, *QRpt-6H.2* is 4.1 Mbp proximal to *QPt.6H-3* (Daba *et al.* 2019), however the nearest
448 marker is 8.1 Mbp proximal and not significant.

449 Two loci that were previously implicated in *Ptt* resistance were also identified as novel MTAs for
450 *Ptm* resistance in this study. The first locus, *QRpt-5H.1*, is located 3.0 Mbp distal to *Qrptts-5HL.1*

(Richards *et al.* 2017) with marker SCRI_RS_160332. Additionally, since markers covering the *Qrptts-5HL.1* locus were not included in either panel, and due to the close proximity of *Qrpt-5H.1* *Qrptts-5HL.1*, we believe that they are the same locus. The second locus, *Qrpt-5H.2*, is embedded within the *NBP_QRptt5-1* locus (Wonneberger *et al.* 2017a). The remaining loci are all embedded within previously identified loci (Table 3).

Barley is predominately grown as a feed crop (IBGSC 2012), however in the US where corn and soybeans are subsidized and used as feed crops it has been outcompeted and acreage has significantly dropped. This has pushed feed barley into less than optimal agricultural land due to its adaptability and hardiness (Brueggeman *et al.*, 2020). Quality malting barley demands premium prices because of its use in the multi-billion dollar added value brewing and distilling industry and is now the major class considered in breeding efforts in the US and Europe. However, in some regions of the world where traditional farming practices are still utilized barley is considered an important food crop (Helbaek 1969; Pourkheirandish and Komatsuda 2007; Geçit 2016; Ergün *et al.* 2017). Recent studies predicted that due to the effects of climate change including higher temperatures, altered precipitation patterns, and higher disease pressure (Dawson *et al.* 2015) the world could experience world malt barley shortages to feed the brewing and distilling industry (Xie *et al.* 2018). These predictions are beginning to be seen with the stagnating yields experienced in southern Europe (Dawson *et al.* 2015). Thus, barley breeding must maximize its potential in terms of quality and yield on the land it is currently afforded to sustain the demands for malting. One pillar of support for improving barley would be the introgression of pre-domestication resistance loci that are absent in current breeding programs to prevent substantial losses to net blotch, an important disease effecting barley

production across the globe. Here we report on the identification of novel loci from Turkish wild barley and landraces that could be introgressed into elite barley varieties.

Tables

Table 1. Phenotypic responses of wild barley and landraces to three *Ptm* isolates with absolute and percentage of accessions in each resistance class. The data is represented as the mean phenotypic score of each barley class to the respective isolate.

Table 2. Phenotypic responses of wild barley and landraces to three *Ptt* isolates with absolute and percentage of accessions in each resistance class. The data is represented as the mean phenotypic score of each barley class to the respective isolate.

Table 3. Identified significant markers from genome wide association analysis order by isolate identified, followed by chromosome and base pair position. Designation as well as predicted corresponding loci, models used to identify the marker and resistant/susceptibility alleles are also included.

Figures

Figure 1. Violin plots of phenotypic distribution of landrace (red) and wild (blue) barley to each *Pyrenophora teres* f. *maculata* (top row) and *P. teres* f. *teres* (bottom row) isolate. Width of the violin indicates the number of accessions with that phenotypic score and the black dot represents the barley class mean. Wilcoxon test significance is indicated by asterisks above each plot.

Figure 2. Enrichment dumbbell plot of enrichment of the resistance allele in landrace (red) and wild (blue) barley accession. Percentage is included below each data point and difference between the two barley classes as well as which isolate the loci was identified with.

Figure 3. Marker density plot of all markers utilized within this study showing the distribution of markers across the barley genome with a window size of 10 Mb.

Figure 4. Manhattan and QQ plots for two models that show significant markers for more than one of the *Pyrenophora teres* f. *maculata* isolates GPS263 and 13-179, and *P. teres* f. *teres* isolates GPS18 and 13-130.

Data Availability Statement

Isolates are available upon request. The authors ensure that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables. Supplemental data has been submitted to Figshare. File SF1 contains genotyping data. File SF2 contains marker positions. File SF3 contains phenotyping data. Files SF4 and SF5 contain population structure and EMMA kinship matrix, respectively. File SF6 contains the mean-squared deviations of each association mapping model tested.

Author Contributions

ACO, AK and RB conceived the study. ACO and AK carried out phenotyping and DNA extractions. KE and DS carried out sequencing for genotyping data. SC and RSP performed genotyping and analysis. SC wrote the manuscript with contributions from ACO, KE, RSP DS, AK and RB. All authors approved the final manuscript.

Acknowledgments

We thank the Western Regional Small Grains Genotyping Laboratory, specifically Travis Ruff and Marcus Hooker for the second Ion Torrent sequencing run, and to the staff and research personnel of the Central Research Institute for Field Crops (Ankara, Turkey) for their help.

Funding

The research presented in this manuscript was supported by the USDA National Institute of Food and Agriculture Hatch project 1014919, Crop Improvement and Sustainable Production Systems (WSU reference 00011).

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Supplementary Material

Supplemental Figure 1. Same as Figure 2 but inverted for susceptibility alleles. Enrichment dumbbell plot of enrichment of the susceptibility allele in landrace (red) and wild (blue) barley accessions. Percentage is included below each data point and difference between the two barley classes as well as which isolate the loci was identified with.

References

Abu Qamar, M., Z. H. Liu, J. D. Faris, S. Chao, M. C. Edwards *et al.*, 2008 A region of barley chromosome 6H harbors multiple major genes associated with net type net blotch resistance. *Theor. Appl. Genet.* 117: 1261.

532 Adhikari, A., B. J. Steffenson, M. J. Smith, and R. Dill-Macky, 2019 Genome-wide association
533 mapping of seedling net form net blotch resistance in an Ethiopian and Eritrean barley
534 collection. Crop Sci.

535 Adhikari, A., B. J. J. Steffenson, K. P. P. Smith, M. Smith, and R. Dill-Macky, 2020 Identification of
536 quantitative trait loci for net form net blotch resistance in contemporary barley breeding
537 germplasm from the USA using genome-wide association mapping. Theor. Appl. Genet.
538 133: 1019–1037.

539 Afanasenko, O. S., M. Jalli, H. O. Pinnschmidt, O. Filatova, and G. J. Platz, 2009 Development of
540 an international standard set of barley differential genotypes for *Pyrenophora teres* f.
541 *teres*. Plant Pathol. 58: 665–676.

542 Afanasenko, O. S., A. V. Koziakov, P. E. Hedlay, N. M. Lashina, A. V. Anisimova *et al.*, 2015
543 Mapping of the loci controlling the resistance to *Pyrenophora teres* f. *teres* and
544 *Cochliobolus sativus* in two double haploid barley populations. Russ. J. Genet. Appl. Res. 5:
545 242–253.

546 Akhavan, A., T. K. Turkington, B. Kebede, A. Tekauz, H. R. Kutcher *et al.*, 2015 Prevalence of
547 mating type idiomorphs in *Pyrenophora teres* f. *teres* and *P. teres* f. *maculata* populations
548 from the Canadian prairies. Can. J. Plant Pathol. 37: 52–60.

549 Allard, R. W., and A. D. Bradshaw, 1964 Implications of genotype-environmental interactions in
550 applied plant breeding. Crop Sci. 4:.

551 Amezrou, R., R. P. S. Verma, S. Chao, R. S. Brueggeman, L. Belqadi *et al.*, 2018 Genome-wide

552 association studies of net form of net blotch resistance at seedling and adult plant stages
553 in spring barley collection. Mol. Breed. 38: 58.

554 Beier, S., A. Himmelbach, C. Colmsee, X.-Q. Zhang, R. A. Barrero *et al.*, 2017 Construction of a
555 map-based reference genome sequence for barley, *Hordeum vulgare* L. Sci. Data 4:
556 170044.

557 Bockelman, H. E., E. L. Sharp, and M. E. Bjarko, 1983 Isolates of *Pyrenophora teres* from
558 Montana and the Mediterranean region that produce spot-type lesions on barley. Plant
559 Dis. 67: 696–697.

560 Bockelman, H. E., E. L. Sharp, and R. F. Eslick, 1977 Trisomic analysis of genes for resistance to
561 scald and net blotch in several barley cultivars. Can. J. Bot. 55: 2142–2148.

562 Bradbury, P. J., Z. Zhang, D. E. Kroon, T. M. Casstevens, Y. Ramdoss *et al.*, 2007 TASSEL:
563 software for association mapping of complex traits in diverse samples. Bioinformatics 23:
564 2633–2635.

565 Brueggeman, R. S., S. Solanki, G. Ameen, K. Effertz, R. S. Poudel *et al.*, 2020 Fungal diseases
566 affecting barley, pp. 1–58 in *Achieving sustainable cultivation of barley*, edited by G. Fox
567 and C. Li. Burleigh Dodds Science Publishing Limited, Cambridge, UK.

568 Burlakoti, R. R., S. Gyawali, S. Chao, K. P. Smith, R. D. Horsley *et al.*, 2017 Genome-wide
569 association study of spot form of net blotch resistance in the Upper Midwest barley
570 breeding programs. Phytopathology 107: 100–108.

571 Cakir, M., S. Gupta, C. Li, M. Hayden, D. E. Mather *et al.*, 2011 Genetic mapping and QTL

572 analysis of disease resistance traits in the barley population Baudin × AC Metcalfe. Crop
573 Pasture Sci. 62: 152–161.

574 Cakir, M., S. Gupta, G. J. Platz, G. A. Ablett, R. Loughman *et al.*, 2003 Mapping and validation of
575 the genes for resistance to *Pyrenophora teres* f. *teres* in barley (*Hordeum vulgare* L.). Aust.
576 J. Agric. Res. 54: 1369–1377.

577 Campbell, G. F., and P. W. Crous, 2003 Genetic stability of net × spot hybrid progeny of the
578 barley pathogen *Pyrenophora teres*. Australas. Plant Pathol. 32: 283–287.

579 Campbell, G. F., J. A. Lucas, and P. W. Crous, 2002 Evidence of recombination between net- and
580 spot-type populations of *Pyrenophora teres* as determined by RAPD analysis. Mycol. Res.
581 106: 602–608.

582 Ceccarelli, S., 1996 Adaptation to low/high input cultivation. Euphytica 92: 203–214.

583 Ceccarelli, S., and S. Grando, 2000 Barley landraces from the Fertile Crescent. A lesson for plant
584 breeders, pp. 51–76 in *Genes in the field, On –farm conservation of crop diversity*, edited
585 by S. B. Brush. IPGRI/IDRC/Lewis Publishers, Boca Raton, London, New York, Washington,
586 D.C.

587 Çelik Oğuz, A., and A. Karakaya, 2017 Pathotypes of *Pyrenophora teres* on barley in Turkey.
588 Phytopathol. Mediterr. 56: 224.

589 Çelik Oğuz, A., A. Karakaya, R. M. Duran, and K. Özbek., 2019a Identification of *Hordeum*
590 *spontaneum* genotypes resistant to net blotch disease. Tarım Bilim. Derg. 25: 115–122.

591 Çelik Oğuz, A., A. Karakaya, N. Ergün, and İ. Sayim, 2017 Turkish barley landraces resistant to

592 net and spot forms of *Pyrenophora teres*. *Phytopathol. Mediterr.* 56: 217–223.

593 Çelik Oğuz, A., F. Ölmez, and A. Karakaya., 2019b Genetic diversity of net blotch pathogens of
594 barley in Turkey. *Int. J. Agric. Biol.* 21: 1089–1096.

595 Çelik Oğuz, A., F. Ölmez, and A. Karakaya, 2018 Mating type idiomorphs of *Pyrenophora teres* in
596 Turkey. *Zemdirbyste-Agriculture* 105: 271–278.

597 Çelik Oğuz, A., A. Rahimi, and A. Karakaya, 2019c Seedling response of Iranian barley landraces
598 to *Pyrenophora teres* f. *teres* and *Pyrenophora teres* f. *maculata*. *Tarım Bilim. Derg.* 25: 86–
599 92.

600 Clare, S. J., N. A. Wyatt, R. S. Brueggeman, and T. L. Friesen, 2020 Research advances in the
601 *Pyrenophora teres*–barley interaction. *Mol. Plant Pathol.* 21: 272–288.

602 Cui, Z., H. Dong, A. Zhang, Y. Ruan, S. Jiang *et al.*, 2020 Denser markers and advanced statistical
603 method identified more genetic loci associated with husk traits in maize. *Sci. Rep.* 10:
604 8165.

605 Daba, S. D., R. Horsley, R. Brueggeman, S. Chao, and M. Mohammadi, 2019 Genome-wide
606 association studies and candidate gene identification for leaf scald and net blotch in barley
607 (*Hordeum vulgare* L.). *Plant Dis.*

608 Dawson, I. K., J. Russell, W. Powell, B. Steffenson, W. T. B. Thomas *et al.*, 2015 Barley: a
609 translational model for adaptation to climate change. *New Phytol.* 206: 913–931.

610 Earl, D. A., and B. M. vonHoldt, 2012 STRUCTURE HARVESTER: a website and program for
611 visualizing STRUCTURE output and implementing the Evanno method. *Conserv. Genet.*

612 Resour. 4: 359–361.

613 Ellis, R. P., B. P. Forster, D. Robinson, L. L. Handley, D. C. Gordon *et al.*, 2000 Wild barley: a
614 source of genes for crop improvement in the 21st century? J. Exp. Bot. 51: 9–17.

615 Emebiri, L. C., C. G. Platz, and D. B. Moody, 2005 Disease resistance genes in a doubled haploid
616 population of two-rowed barley segregating for malting quality attributes. Aust. J. Agric.
617 Res. 56: 49–56.

618 Endresen, D. T. F., K. Street, M. Mackay, A. Bari, and E. De Pauw, 2011 Predictive association
619 between biotic stress traits and eco-geographic data for wheat and barley landraces. Crop
620 Sci. 51: 2036–2055.

621 Ergün, N., S. Aydoğan, İ. Sayim, A. Karakaya, and A. Çelik Oğuz, 2017 Arpa (*Hordeum vulgare* L.)
622 köy çeşitlerinde tane verimi ve bazı tarımsal özelliklerin incelenmesi. Tarla Bitk. Merk.
623 Araştırma Enstitüsü Derg. 26: 180–189.

624 Evanno, G., S. Regnaut, and J. Goudet, 2005 Detecting the number of clusters of individuals
625 using the software structure: a simulation study. Mol. Ecol. 14: 2611–2620.

626 Fetch, T. G., B. J. Steffenson, and E. Nevo, 2003 Diversity and sources of multiple disease
627 resistance in *Hordeum spontaneum*. Plant Dis. 87: 1439–1448.

628 Franckowiak, J. D., and G. J. Platz, 2013 International database for barley genes and barley
629 genetic stocks. Barley Genet. Newsl. 43: 48–223.

630 Friesen, T. L., J. D. Faris, Z. Lai, and B. J. Steffenson, 2006 Identification and chromosomal
631 location of major genes for resistance to *Pyrenophora teres* in a doubled-haploid barley

632 population. Genome 49: 855–859.

633 Geçit, H. H., 2016 *Serin İklim Tahılları (Buğday, Arpa, Yulaf, Triticale)*. Ankara Üniversitesi Ziraat

634 Fakültesi Yayınları, Yayın No:1640, Ankara: Ankara.

635 Geschele, E. E., 1928 The response of barley to parasitic fungi *Helminthosporium teres* Sacc.

636 Bull. Appl. Bot. Genet. Plant Breed. 19: 19, 371– 384 (in Rev. Appl. Mycol. 8, 165).

637 Graner, A., B. Foroughi-Wehr, and A. Tekauz, 1996 RFLP mapping of a gene in barley conferring

638 resistance to net blotch (*Pyrenophora teres*). Euphytica 91: 229–234.

639 Grewal, T. S., B. G. Rossnagel, C. J. Pozniak, and G. J. Scoles, 2008 Mapping quantitative trait loci

640 associated with barley net blotch resistance. Theor. Appl. Genet. 116: 529–539.

641 Grewal, T. S., B. G. Rossnagel, and G. J. Scoles, 2012 Mapping quantitative trait loci associated

642 with spot blotch and net blotch resistance in a doubled-haploid barley population. Mol.

643 Breed. 30: 267–279.

644 Gupta, S., C. Li, R. Loughman, M. Cakir, S. Westcott *et al.*, 2011 Identifying genetic complexity of

645 6H locus in barley conferring resistance to *Pyrenophora teres* f. *teres*. Plant Breed. 130:

646 423–429.

647 Gyawali, S., R. Amezrou, R. P. S. Verma, R. Brueggeman, S. Rehman *et al.*, 2020 Seedling and

648 adult stage resistance to net form of net blotch (NFNB) in spring barley and stability of

649 adult stage resistance to NFNB in Morocco. J. Phytopathol. 168: 254–266.

650 Gyawali, S., R. Amezrou, R. P. S. Verma, R. Brueggeman, S. Rehman *et al.*, 2019 Seedling and

651 adult stage resistance to spot form of net blotch (SFNB) in spring barley and stability of

652 adult stage resistance to SFNB in Morocco. *Eur. J. Plant Pathol.* 153: 475–487.

653 Harlan, J. R., and D. Zohary, 1966 Distribution of Wild Wheats and Barley. *Science* (80-.). 153:

654 1074–1080.

655 Helbaek, H., 1969 Plant-collecting, dry-farming and irrigation agriculture in prehistoric Deh

656 Luran, pp. 383–426 in *Prehistory and human ecology of the Deh Luran plain. An early*

657 *village sequence from Khuzistan, Iran*, edited by F. Hole, K. V. Flannery, and J.A. Neely.

658 University of Michigan.

659 Hisano, H., K. Sakamoto, H. Takagi, R. Terauchi, and K. Sato, 2017 Exome QTL-seq maps

660 monogenic locus and QTLs in barley. *BMC Genomics* 18: 125.

661 Huang, M., X. Liu, Y. Zhou, R. M. Summers, and Z. Zhang, 2019 BLINK: a package for the next

662 level of genome-wide association studies with both individuals and markers in the millions.

663 *Gigascience* 8:.

664 IBGSC, 2012 A physical, genetic and functional sequence assembly of the barley genome.

665 *Nature* 491: 711.

666 Islamovic, E., P. Bregitzer, and T. L. Friesen, 2017 Barley 4H QTL confers NFNB resistance to a

667 global set of *P. teres* f. *teres* isolates. *Mol. Breed.* 37: 29.

668 Jakob, S. S., D. Rödder, J. O. Engler, S. Shaaf, H. Ozkan *et al.*, 2014 Evolutionary history of wild

669 barley (*Hordeum vulgare* subsp. *spontaneum*) analyzed using multilocus sequence data

670 and paleodistribution modeling. *Genome Biol. Evol.* 6: 685–702.

671 Jana, S., and K. L. Bailey, 1995 Responses of wild and cultivated barley from West Asia to net

672 blotch and spot blotch. Crop Sci. 35:.

673 Kang, H. M., N. A. Zaitlen, C. M. Wade, A. Kirby, D. Heckerman *et al.*, 2008 Efficient control of
674 population structure in model organism association mapping. Genetics 178: 1709 LP –
675 1723.

676 Karakaya, A., A. Çelik Oğuz, and I. Saraç Sivrikaya, 2020 Diseases occurring on Hordeum
677 bulbosum field populations at Bingöl province of Turkey. Work. Fac. Agric. Food Sci. Univ.
678 Sarajev. LXV: 75–81.

679 Karakaya, A., Y. Z. Katırcioğlu, and H. Aktaş, 2004 Studies on the biology of *Drechslera teres*
680 under Ankara conditions. Tarım Bilim. Derg. 10: 133–135.

681 Karakaya, A., Z. Mert, A. Çelik Oğuz, M. N. Ertaş, and A. Karagöz, 2016 Determination of the
682 diseases occurring on naturally growing wild barley (*Hordeum spontaneum*) field
683 populations. Work. Fac. Agric. Univ. Sarajev. 61: 291–295.

684 Koladia, V. M., J. D. Faris, J. K. Richards, R. S. Brueggeman, S. Chao *et al.*, 2017 Genetic analysis
685 of net form net blotch resistance in barley lines Clho 5791 and Tifang against a global
686 collection of *P. teres* f. *teres* isolates. Theor. Appl. Genet. 130: 163–173.

687 Komatsuda, T., P. Maxim, N. Senthil, and Y. Mano, 2004 High-density AFLP map of nonbrittle
688 rachis 1 (*btr1*) and 2 (*btr2*) genes in barley (*Hordeum vulgare* L.). Theor. Appl. Genet. 109:
689 986–995.

690 König, J., D. Perovic, D. Kopahnke, and F. Ordon, 2013 Development of an efficient method for
691 assessing resistance to the net type of net blotch (*Pyrenophora teres* f. *teres*) in winter

692 barley and mapping of quantitative trait loci for resistance. Mol. Breed. 32: 641–650.

693 Kün, E., 1996 *Tahıllar-1 (Serin İklim Tahılları)*. Ankara Üniversitesi Ziraat Fakültesi Yayınları,

694 Yayın No:1451, Ankara.

695 Lakew, B., Y. Semeane, and F. Alemayehu, 1995 Evaluation of Ethiopian barley landraces for

696 disease and agronomic characters. Rachis 14: 21–25.

697 Legge, W. G., D. R. Metcalfe, A. W. Chiko, J. W. Martens, and A. Tekauz, 1996 Reaction of

698 Turkish barley accessions to Canadian barley pathogens. Can. J. Plant Sci. 76: 927–931.

699 Lehmensiek, A., G. J. Platz, E. Mace, D. Poulsen, and M. W. Sutherland, 2007 Mapping of adult

700 plant resistance to net form of net blotch in three Australian barley populations. Aust. J.

701 Agric. Res. 58: 1191–1197.

702 Li, M., X. Liu, P. Bradbury, J. Yu, Y.-M. Zhang *et al.*, 2014 Enrichment of statistical power for

703 genome-wide association studies. BMC Biol. 12: 73.

704 Liu, Z., S. R. Ellwood, R. P. Oliver, and T. L. Friesen, 2011 *Pyrenophora teres*: profile of an

705 increasingly damaging barley pathogen. Mol. Plant Pathol. 12: 1–19.

706 Liu, Z., D. J. Holmes, J. D. Faris, S. Chao, R. S. Brueggeman *et al.*, 2015 Necrotrophic effector-

707 triggered susceptibility (NETS) underlies the barley–*Pyrenophora teres* f. *teres* interaction

708 specific to chromosome 6H. Mol. Plant Pathol. 16: 188–200.

709 Liu, X., M. Huang, B. Fan, E. S. Buckler, and Z. Zhang, 2016 Iterative usage of fixed and random

710 effect models for powerful and efficient genome-wide association studies. PLOS Genet. 12:

711 e1005767.

712 Loiselle, B. A., V. L. Sork, J. Nason, and C. Graham, 1995 Spatial genetic structure of a tropical
713 understory shrub, *Psychotria officinalis* (Rubiaceae). *Am. J. Bot.* 82: 1420–1425.

714 Mamidi, S., S. Chikara, R. J. Goos, D. L. Hyten, D. Annam *et al.*, 2011 Genome-wide association
715 analysis identifies candidate genes associated with iron deficiency chlorosis in soybean.
716 *Plant Genome* 4: 154–164.

717 Manninen, O. M., M. Jalli, R. Kalendar, A. Schulman, O. Afanasenko *et al.*, 2006 Mapping of
718 major spot-type and net-type net-blotch resistance genes in the Ethiopian barley line CI
719 9819. *Genome* 49: 1564–1571.

720 Manninen, O., R. Kalendar, J. Robinson, and A. H. Schulman, 2000 Application of BARE-1
721 retrotransposon markers to the mapping of a major resistance gene for net blotch in
722 barley. *Mol. Gen. Genet.* MGG 264: 325–334.

723 Martin, A., G. J. Platz, D. de Klerk, R. A. Fowler, F. Smit *et al.*, 2018 Identification and mapping of
724 net form of net blotch resistance in South African barley. *Mol. Breed.* 38: 53.

725 Mascher, M., H. Gundlach, A. Himmelbach, S. Beier, S. O. Twardziok *et al.*, 2017 A chromosome
726 conformation capture ordered sequence of the barley genome. *Nature* 544: 427–433.

727 Mascher, M., G. J. Muehlbauer, D. S. Rokhsar, J. Chapman, J. Schmutz *et al.*, 2013 Anchoring
728 and ordering NGS contig assemblies by population sequencing (POPSEQ). *Plant J.* 76: 718–
729 727.

730 Mathre, D. E. (ed.), 1982 *Compendium of Barley Diseases*. APS Press, St. Paul, Minnesota.

731 McDonald, B. A., and C. Linde, 2002 Pathogen population genetics, evolutionary potential, and

732 durable resistance. Annu. Rev. Phytopathol. 40: 349–379.

733 McLean, M. S., A. Martin, S. Gupta, M. W. Sutherland, G. J. Hollaway *et al.*, 2014 Validation of a
734 new spot form of net blotch differential set and evidence for hybridisation between the
735 spot and net forms of net blotch in Australia. Australas. Plant Pathol. 43: 223–233.

736 Metcalfe, D. R., K. W. Buchannon, W. C. McDonald, and E. Reinbergs, 1970 Relationships
737 between the ‘Jet’ and “Milton” genes for resistance to loose smut genes for resistance to
738 other barley diseases. Can. J. Plant Sci. 50: 423–427.

739 Mikhailova, L. A., I. G. Ternyuk, and N. V. Mironenko, 2010 *Pyrenophora teres*, an agent causing
740 wheat leaf spot. Microbiology 79: 561–565.

741 Mode, C. J., and C. W. Schaller, 1958 Two additional factors for host resistance to net blotch in
742 barley. Agron. J. 50: 15–18.

743 Monat, C., S. Padmarasu, T. Lux, T. Wicker, H. Gundlach *et al.*, 2019 TRITEX: chromosome-scale
744 sequence assembly of Triticeae genomes with open-source tools. Genome Biol. 20: 284.

745 Money, D., K. Gardner, Z. Migicovsky, H. Schwaninger, G.-Y. Zhong *et al.*, 2015 LinkImpute: Fast
746 and accurate genotype imputation for nonmodel organisms. G3 Genes|Genomes|Genetics
747 5: 2383–2390.

748 Moya, P. A., J. R. Girotti, A. V. Toledo, and M. N. Sisterna, 2018 Antifungal activity of
749 *Trichoderma* VOCs against *Pyrenophora teres*, the causal agent of barley net blotch. J.
750 Plant Prot. Res. 58: 45–53.

751 Muñoz-Amatriáin, M., A. Cuesta-Marcos, J. B. Endelman, J. Comadran, J. M. Bonman *et al.*, 2014

752 The USDA barley core collection: Genetic diversity, population structure, and potential for
753 genome-wide association studies (L. Yan, Ed.). PLoS One 9: e94688.

754 Neupane, A., P. Tamang, R. S. Brueggeman, and T. L. Friesen, 2015 Evaluation of a barley core
755 collection for spot form net blotch reaction reveals distinct genotype-specific pathogen
756 virulence and host susceptibility. Phytopathology 105: 509–517.

757 Nevo, E., 1992 Origin, evolution, population genetics and resources for breeding of wild barley,
758 *Hordeum spontaneum*, in the Fertile Crescent, pp. 19–43 in *Barley: genetics, biochemistry,*
759 *molecular biology and biotechnology*, edited by P. R. Shewry. C.A.B. International UK.

760 Novakazi, F., O. Afanasenko, A. Anisimova, G. J. J. Platz, R. Snowdon *et al.*, 2019 Genetic analysis
761 of a worldwide barley collection for resistance to net form of net blotch disease
762 (*Pyrenophora teres* f. *teres*). Theor. Appl. Genet. 132: 2633–2650.

763 O’Boyle, P. D., W. S. Brooks, M. D. Barnett, G. L. Berger, B. J. Steffenson *et al.*, 2014 Mapping
764 net blotch resistance in ‘Nomini’ and Clho 2291 barley. Crop Sci. 54: 2596–2602.

765 Perelló, A. E., L. Couretot, A. Curti, J. P. Uranga, and V. F. Consolo, 2019 First report of spot
766 lesion of wheat caused by *Pyrenophora teres* f. sp *maculata* observed in Argentina. Crop
767 Prot. 122: 19–22.

768 Piening, L., and M. L. Kaufmann, 1969 Comparison of the effectors of net blotch and leaf
769 removal on yield in barley. Can. J. Plant Sci. 49: 731–735.

770 St. Pierre, S., C. Gustus, B. Steffenson, R. Dill-Macky, and K. P. Smith, 2010 Mapping net form
771 net blotch and Septoria speckled leaf blotch resistance loci in barley. Phytopathology 100:

772 80–84.

773 Poets, A. M., Z. Fang, M. T. Clegg, and P. L. Morrell, 2015 Barley landraces are characterized by
774 geographically heterogeneous genomic origins. *Genome Biol.* 16: 173.

775 Poudel, B., M. S. McLean, G. J. Platz, J. A. McIlroy, M. W. Sutherland *et al.*, 2019 Investigating
776 hybridisation between the forms of *Pyrenophora teres* based on Australian barley field
777 experiments and cultural collections. *Eur. J. Plant Pathol.* 153: 465–473.

778 Pourkheirandish, M., and T. Komatsuda, 2007 The importance of barley genetics and
779 domestication in a global perspective. *Ann. Bot.* 100: 999–1008.

780 Pritchard, J. K., M. Stephens, and P. Donnelly, 2000 Inference of population structure using
781 multilocus genotype data. *Genetics* 155: 945–959.

782 Raman, H., G. J. Platz, K. J. Chalmers, R. Raman, B. J. Read *et al.*, 2003 Mapping of genomic
783 regions associated with net form of net blotch resistance in barley. *Aust. J. Agric. Res.* 54:
784 1359–1367.

785 Read, B. J., H. Raman, G. McMichael, K. J. Chalmers, G. A. Ablett *et al.*, 2003 Mapping and QTL
786 analysis of the barley population Sloop × Halcyon. *Aust. J. Agric. Res.* 54: 1145–1153.

787 Richards, J., S. Chao, T. Friesen, and R. Brueggeman, 2016 Fine mapping of the barley
788 chromosome 6H net form net blotch susceptibility locus. *G3 Genes | Genomes | Genetics* 6:
789 1809–1818.

790 Richards, J. K., T. L. Friesen, and R. S. Brueggeman, 2017 Association mapping utilizing diverse
791 barley lines reveals net form net blotch seedling resistance/susceptibility loci. *Theor. Appl.*

792 Genet. 130: 915–927.

793 Richter, K., J. Schondelmaier, and C. Jung, 1998 Mapping of quantitative trait loci affecting
794 *Drechslera teres* resistance in barley with molecular markers. Theor. Appl. Genet. 97:
795 1225–1234.

796 Robinson, J., and M. Jalli, 1996 Diversity among Finnish net blotch isolates and resistance in
797 barley. Euphytica 92: 81–87.

798 Rozanova, I. V, N. M. Lashina, Z. S. Mustafin, S. A. Gorobets, V. M. Efimov *et al.*, 2019 SNPs
799 associated with barley resistance to isolates of *Pyrenophora teres* f. *teres*. BMC Genomics
800 20: 292.

801 Ruff, T. M., E. J. Marston, J. D. Eagle, S. R. Sthapit, M. A. Hooker *et al.*, 2020 Genotyping by
802 multiplexed sequencing (GMS): A customizable platform for genomic selection. PLoS One
803 15: e0229207.

804 Sato, K., and K. Takeda, 1997 Net blotch resistance in wild species of *Hordeum*. Euphytica 95:
805 179–185.

806 Schaller, C. W., 1955 Inheritance of resistance to net blotch of barley. Phytopathology 45: 174–
807 176.

808 Segura, V., B. J. Vilhjálmsson, A. Platt, A. Korte, Ü. Seren *et al.*, 2012 An efficient multi-locus
809 mixed-model approach for genome-wide association studies in structured populations.
810 Nat. Genet. 44: 825–830.

811 Serenius, M., N. Mironenko, and M. O., 2005 Genetic variation, occurrence of mating types and

812 different forms of *Pyrenophora teres* causing net blotch of barley in Finland. Mycol. Res.
813 107: 809–817.

814 Sharma Poudel, R., A. F. Al-Hashel, T. Gross, P. Gross, and R. Brueggeman, 2018 Pyramiding
815 *rpg4*- and *Rpg1*-mediated stem rust resistance in barley requires the *Rrr1* gene for both to
816 function. Front. Plant Sci. 9: 1789.

817 Silvar, C., A. M. Casas, D. Kopahnke, A. Habekuß, G. Schweizer *et al.*, 2010 Screening the Spanish
818 barley core collection for disease resistance. Plant Breed. 129: 45–52.

819 Smedegård-Petersen, V., 1971 *Pyrenophora teres* f. *maculata* f. nov. and *Pyrenophora teres* f.
820 *teres* on barley in Denmark. Kgl. Vet. Landbohojsk. Arsskr. 124–144.

821 Steffenson, B. J., P. M. Hayes, and A. Kleinohs, 1996 Genetics of seedling and adult plant
822 resistance to net blotch (*Pyrenophora teres* f. *teres*) and spot blotch (*Cochliobolus sativus*)
823 in barley. Theor. Appl. Genet. 92: 552–558.

824 Tamang, P., A. Neupane, S. Mamidi, T. Friesen, and R. Brueggeman, 2015 Association mapping
825 of seedling resistance to spot form net blotch in a worldwide collection of barley.
826 Phytopathology 105: 500–508.

827 Tamang, P., J. K. Richards, A. Alhashal, R. Sharma Poudel, R. D. Horsley *et al.*, 2019 Mapping of
828 barley susceptibility/resistance QTL against spot form net blotch caused by *Pyrenophora*
829 *teres* f. *maculata* using RIL populations. Theor. Appl. Genet.

830 Thomas, W. T. B., E. Baird, J. D. Fuller, P. Lawrence, G. R. Young *et al.*, 1998 Identification of a
831 QTL decreasing yield in barley linked to Mlo powdery mildew resistance. Mol. Breed. 4:

832 381–393.

833 Tóth, B., M. Csősz, D. Kopahnke, and J. Varga, 2008 First report on *Pyrenophora teres* causing
834 lesions of wheat leaves in Hungary. Plant Pathol. 57: 385.

835 Usta, P., A. Karakaya, A. Çelik Oğuz, Z. Mert, K. Akan *et al.*, 2014 Determination of the seedling
836 reactions of twenty barley cultivars to six isolates of *Drechslera teres* f. *maculata*. Anadolu
837 Tarım Bilim. Derg. (Anadolu J. Agric. Sci.) 29: 20–25.

838 VanRaden, P. M., 2008 Efficient methods to compute genomic predictions. J. Dairy Sci. 91:
839 4414–4423.

840 Vatter, T., A. Maurer, D. Kopahnke, D. Perovic, F. Ordon *et al.*, 2017 A nested association
841 mapping population identifies multiple small effect QTL conferring resistance against net
842 blotch (*Pyrenophora teres* f. *teres*) in wild barley. PLoS One 12: e0186803.

843 Wang, X., E. S. Mace, G. J. Platz, C. H. Hunt, L. T. Hickey *et al.*, 2015 Spot form of net blotch
844 resistance in barley is under complex genetic control. Theor. Appl. Genet. 128: 489–499.

845 Wang, Q., F. Tian, Y. Pan, E. S. Buckler, and Z. Zhang, 2014 A SUPER powerful method for
846 genome wide association study. PLoS One 9: e107684.

847 Wang, Y. L., H. Ye, L. Liu, J. H. Wu, W. M. Ru *et al.*, 2019 Molecular insights on the domestication
848 of barley (*Hordeum vulgare* L.). CRC. Crit. Rev. Plant Sci. 38: 280–294.

849 Wang, J., and Z. Zhang, 2018 GAPIT Version 3: An interactive analytical tool for genomic
850 association and prediction. preprint.

851 Williams, K. J., A. Lichon, P. Gianquitto, J. M. Kretschmer, A. Karakousis *et al.*, 1999

852 Identification and mapping of a gene conferring resistance to the spot form of net blotch
853 (*Pyrenophora teres* f. *maculata*) in barley. Theor. Appl. Genet. 99: 323–327.

854 Williams, K. J., G. J. Platz, A. R. Barr, J. Cheong, K. Willsmore *et al.*, 2003 A comparison of the
855 genetics of seedling and adult plant resistance to the spot form of net blotch (*Pyrenophora*
856 *teres* f. *maculata*). Aust. J. Agric. Res. 54: 1387–1394.

857 Wonneberger, R., A. Ficke, and M. Lillemo, 2017a Identification of quantitative trait loci
858 associated with resistance to net form net blotch in a collection of Nordic barley
859 germplasm. Theor. Appl. Genet. 130: 2025–2043.

860 Wonneberger, R., A. Ficke, and M. Lillemo, 2017b Mapping of quantitative trait loci associated
861 with resistance to net form net blotch (*Pyrenophora teres* f. *teres*) in a doubled haploid
862 Norwegian barley population. PLoS One 12: e0175773.

863 Xie, W., W. Xiong, J. Pan, T. Ali, Q. Cui *et al.*, 2018 Decreases in global beer supply due to
864 extreme drought and heat. Nat. Plants 4: 964–973.

865 Yazıcı, B., A. Karakaya, A. Çelik Oğuz, and Z. Mert, 2015 Determination of the seedling reactions
866 of some barley cultivars to *Drechslera teres* f. *teres*. Bitki Koruma Bülteni 55: 239–245.

867 Yitbarek, S., L. Berhane, A. Fikadu, J. A. G. Van Leur, S. Grando *et al.*, 1998 Variation in Ethiopian
868 barley landrace populations for resistance to barley leaf scald and netblotch. Plant Breed.
869 117: 419–423.

870 Yu, J., G. Pressoir, W. H. Briggs, I. Vroh Bi, M. Yamasaki *et al.*, 2006 A unified mixed-model
871 method for association mapping that accounts for multiple levels of relatedness. Nat.

872 Genet. 38: 203–208.

873 Yun, S. J., L. Gyenis, P. M. Hayes, I. Matus, K. P. Smith *et al.*, 2005 Quantitative trait loci for

874 multiple disease resistance in wild barley. Crop Sci. 45: 2563–2572.

875 Zhang, Z., E. Ersoz, C.-Q. Lai, R. J. Todhunter, H. K. Tiwari *et al.*, 2010 Mixed linear model

876 approach adapted for genome-wide association studies. Nat. Genet. 42: 355–360.

877 Zohary, D., and M. Hopf, 2000 *Domestication of plants in the old world: the origin and spread of*

878 *cultivated plants in West Asia, Europe, and the Nile Valley*. Oxford University Press, New

879 York.

880

Table 1. Phenotypic responses of landrace and wild barley to three *Ptm* isolates with absolute and percentage of accessions in each resistance class. Mean phenotypic score of each barley class to the

Class	GPS263		13-179		13-167	
	Landrace	Wild	Landrace	Wild	Landrace	Wild
Resistant	6 (3%)	10 (10%)	0 (0%)	3 (3%)	6 (3%)	12 (12%)
Moderately Resistant	42 (24%)	38 (39%)	65 (37%)	34 (35%)	66 (37%)	50 (51%)
Intermediate	42 (24%)	28 (29%)	105 (59%)	57 (58%)	100 (56%)	35 (36%)
Moderately Susceptible	86 (49%)	21 (21%)	7 (4%)	4 (4%)	5 (3%)	1 (1%)
Susceptible	1 (1%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Mean Score	5.4	4.4	4.3	4.3	4.2	3.6

respective isolate.

Table 2. Phenotypic responses of landrace and wild barley to three *Ptt* isolates with absolute and percentage of accessions in each resistance class. Mean phenotypic score of each barley class to the respective isolate.

Class	GPS18		UHK77		13-130	
	Landrace	Wild	Landrace	Wild	Landrace	Wild
Resistant	1 (1%)	1 (1%)	1 (1%)	7 (7%)	4 (2%)	8 (8%)
Moderately Resistant	26 (15%)	32 (33%)	103 (58%)	32 (33%)	122 (69%)	44 (45%)
Intermediate	18 (10%)	14 (14%)	51 (29%)	18 (18%)	35 (20%)	21 (21%)
Moderately Susceptible	131 (74%)	51 (52%)	22 (12%)	38 (39%)	16 (9%)	25 (26%)
Susceptible	1 (1%)	0 (0%)	0 (0%)	3 (3%)	0 (0%)	0 (0%)
Mean Score	5.7	5.1	4.5	4.9	4.0	4.4

Table 3. Identified significant markers from genome wide association analysis order by isolate identified, followed by chromosome and base pair position. Designation as well as predicted corresponding loci, models used to identify the marker and resistant/susceptibility alleles are also included.

Loci	Marker	Chr ¹	Position ¹	Allele ²	Isolate	Corresponding Loci	Models Identified	LOD score ³
<i>QRpt-5H.2</i>	12_20350	5H	446449843	G/A	GPS263	<i>NBP_QRptt5-1</i>	K _{BLINK}	7.76
<i>QRpt-3H.2</i>	11_20866	3H	153156749	G/A	13_179	<i>QRptms3-2</i>	K _{FarmCPU} , PC1+K _{FarmCPU}	4.58
<i>QRpt-4H.1</i>	11_10510	4H	603258307	A/G		<i>Rpt8</i>	K _{BLINK+SUPER} , PC5 _{GLM} , Q+K _{SUPER} , PC1+K _{SUPER}	5.57
<i>QRpt-5H.1</i>	SCRI_RS_160332	5H	474799503	A/G		<i>Qrptts-5HL.1</i>	K _{FarmCPU}	4.70
<i>QRpt-1H.1</i>	11_10176	1H	397791042	G/C	GPS18	Novel	K _{FarmCPU}	4.31
<i>QRpt-1H.2</i>	11_20754	1H	483805599	C/G		<i>QPt.1H-1</i>	K _{BLINK+FarmCPU}	4.66-5.92
<i>QRpt-6H.2</i>	11_20972	6H	539551443	T/A		<i>AL_QRptt6-2</i>	K _{BLINK}	4.46
<i>QRpt-7H.1</i>	12_30545	7H	54934072	A/G		<i>QNFNBAPR.AL/S-7Ha</i>	K _{FarmCPU}	4.05
<i>QRpt-7H.3</i>	12_31282	7H	617741299	C/T		<i>QTL_{UHS}-7H</i>	K _{FarmCPU}	5.49
<i>QRpt-2H.1</i>	12_11452	2H	34275254	G/A	13_130	<i>SFNB-2H-8-10</i>	K _{BLINK+FarmCPU}	4.05-4.23
<i>QRpt-3H.1</i>	11_20968	3H	19966889	A/G		Novel	K _{FarmCPU}	5.15
<i>QRpt-3H.3</i>	12_10662	3H	553445025	A/T		Novel	K _{FarmCPU} , Q+K _{FarmCPU}	4.22-5.06
<i>QRpt-6H.1</i>	11_20714	6H	489619101	G/A		Novel	K _{FarmCPU}	4.71
<i>QRpt-7H.2</i>	11_11243	7H	601974526	A/G		<i>QRptm7-6</i>	K _{FarmCPU}	4.57

¹, Location based on the second version of the Morex assembly (Monat et al., 2019).

², Resistant/susceptible allele

³, LOD scores/ranges for BLINK and FarmCPU algorithms only

The number of SNPs within 10Mb window size

0Mb 84Mb 168Mb 252Mb 336Mb 420Mb 504Mb 588Mb 672Mb 756Mb