***Rpi-amr3* confers resistance to multiple *Phytophthora* species by recognizing a conserved RXLR effector**

Xiao Lin¹, Andrea Olave-Achury¹, Robert Heal¹, Kamil Witek¹, Hari S. Karki^{1#}, Tianqiao Song^{1#}, Chih-hang Wu^{1#}, Hiroaki Adachi^{1#}, Sophien Kamoun¹, Vivianne G. A. A. Vleeshouwers² and Jonathan D. G. Jones^{1*}

¹The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, NR4 7UH, UK

²Wageningen UR Plant Breeding, Wageningen University and Research, Droevendaalsesteeg 1, 6708 PB, Wageningen, The Netherlands

[#]Current addresses:

HSK: U.S. Department of Agriculture–Agricultural Research Service, Madison, WI 53706, U.S.A

TS: Institute of Plant Protection, Jiangsu Academy of Agricultural Sciences, Nanjing, 210014, P. R. China

CHW: Institute of Plant and Microbial Biology, Academia Sinica, Taiwan

HA: Nara Institute of Science and Technology, Ikoma 630-0192, Japan

*Corresponding author: Jonathan D. G. Jones (jonathan.jones@tsl.ac.uk)

Abstract

Diverse pathogens from the genus *Phytophthora* cause disease and reduce yields in many crop plants. Although many *Resistance to Phytophthora infestans* (*Rpi*) genes effective against potato late blight have been cloned, few have been cloned against other *Phytophthora* species. Most *Rpi* genes encode nucleotide-binding domain, leucine-rich repeat- containing (NLR) proteins, that recognize RXLR effectors. However, whether NLR proteins can recognize RXLR effectors from multiple different *Phytophthora* pathogens has rarely been investigated. Here, we report the effector AVRamr3 from *P. infestans* that is recognized by Rpi-amr3 from *S. americanum*. We show here that AVRamr3 is broadly conserved in many different *Phytophthora* species, and that recognition of AVRamr3 homologs enables resistance against multiple *Phytophthora* pathogens, including *P. parasitica* and *P. palmivora*. Our findings suggest a novel path to identifying *R* genes against important plant pathogens.

Introduction

Species in the oomycete genus *Phytophthora* cause many devastating plant diseases. For example, *P. infestans*, *P. parasitica*, *P. cactorum*, *P. ramorum*, *P. sojae*, *P. palmivora* and *P. megakarya* cause disease on potato and tomato, tobacco, strawberry, oak, soybean and cacao, respectively (Kamoun et al., 2015).

Plant immunity involves detection of pathogen-derived molecules by either cell-surface pattern recognition immune receptors (PRRs) or intracellular nucleotide-binding domain, leucine-rich repeat containing (NLR) immune receptors, that activate either pattern-triggered immunity (PTI) or effector-triggered immunity (ETI), respectively (Jones and Dangl, 2006). So far, many *Resistance to P. infestans* (*Rpi*) genes were cloned from wild *Solanum* species which confer resistance against potato late blight (Vleeshouwers et al., 2011). Many *R* genes against *P. sojae* (*Rps*) have also been mapped in different soybean accessions, and some were cloned (Sahoo et al., 2017). In tobacco, the black shank resistance genes *Phl*, *Php* and *Ph* were genetically mapped but not yet cloned; these confer race-specific resistance to *P. parasitica* (aka *P. nicotianae*) isolates (Gallup and Shew, 2010; Bao et al., 2019). For *P. palmivora*, some resistant cacao (*Theobroma cacao*) accessions were identified, but no dominant *R* genes have been defined or cloned (Thevenin et al., 2012). In summary, very few *R* genes against other *Phytophthora* pathogens have been cloned.

Solanum americanum and *S. nigrum* are highly resistant to *P. infestans* (Witek et al., 2016; Witek et al., 2021). Two *Rpi* genes of coiled-coil (CC) type, *Rpi-amr3* and *Rpi-amr1*, were cloned from different *S. americanum* accessions, and both confer broad-spectrum late blight resistance in cultivated potatoes (Witek et al., 2016; Witek et al., 2021).

In oomycetes, *Rpi* proteins typically recognize RXLR (Arg-X-Leu-Arg, X represents any amino acid)-EER (Glu-Glu-Arg) effectors that are secreted into plant cells (Rehmany et al., 2005; Wang et al., 2019). Many *Avirulence* (*Avr*) genes encoding recognized effectors from *Phytophthora* species have been identified and they are often fast-evolving and lineage-specific molecules (Jiang et al., 2008). Recently, AVRamr1 (PITG_07569), the recognized effector of *Rpi-amr1* was identified by a long read and cDNA pathogen enrichment sequencing (PenSeq) approach (Lin et al., 2020a). Surprisingly, AVRamr1 homologs were identified from *P.*

parasitica and *P. cactorum* genomes and both are recognized by all Rpi-amr1 variants (Witek et al., 2021). Similarly, AVR3a-like effectors were found in different *Phytophthora* species, including *P. capsici* and *P. sojae*, and the recognition of AVR3a homologs correlates with *P. capsici* or *P. sojae* resistance in *Nicotiana* species and soybean (Shan et al., 2004; Vega-Arreguín et al., 2014), also AVRblb2 homologs from *P. andina* and *P. mirabilis* trigger HR with Rpi-blb (Oliva et al., 2015). Remarkably, a single N336Y mutation in R3a expands its recognition specificity to *P. capsici* AVR3a homolog (Segretin et al., 2014). These reports raise intriguing questions. Could RXLR effectors be widely conserved molecules among different *Phytophthora* species? Could these effectors be recognized by the same plant receptor? Of particular interest, could this effector recognition capacity enable disease resistance?

Here we identified AVRamr3, a novel AVR protein from *P. infestans*, by screening an RXLR effector library. AVRamr3 is a broadly conserved effector found in many different *Phytophthora* species. Strikingly, the recognition of AVRamr3 not only enables resistance to *P. infestans*, but also to other economically important *Phytophthora* pathogens including *P. parasitica* and *P. palmivora*. We also show functional *Rpi-amr3* genes are widely distributed among *S. americanum* and *S. nigrum* accessions, together with the previously defined *Rpi-amr1* genes, they might underpin the “non-host” resistance of these species against *P. infestans*.

Results:

Avramr3 encodes a conserved RXLR-WY effector protein

To identify the effector recognized by Rpi-amr3, we screened an RXLR effector library (Rietman, 2011; Lin *et al.*, 2020) by *Agrobacterium tumefaciens*-mediated co-expression with Rpi-amr3 in *Nicotiana benthamiana*. By screening ~150 RXLR effectors, we found PITG_21190 specifically induces hypersensitive response (HR) with *Rpi-amr3* (Fig. 1a), and concluded PITG_21190 is *Avramr3*. *Avramr3* encodes a 339-aa protein with a signal peptide followed by RXLR, EER motifs and four predicted WY motifs (Win *et al.*, 2012) (Fig. 1b). In *P. infestans* T30-4, the expression of *Avramr3* is low during infection, however *Avramr3* upregulated in 3928A and US23 in 2-3 days after infection (Cooke *et al.*, 2012; Lin *et al.*, 2020a).

Many RXLR effectors are fast-evolving, multiple-member family proteins with extensive sequence polymorphism, such as the *Avr2* and *Avrblb2* families (Gilroy *et al.*, 2011; Oliva *et al.*, 2015). To study the sequence polymorphism of *Avramr3*, seventeen additional *Avramr3* homologs from eleven isolates were identified from published databases (KR_1, 3928A, EC1, 6_A1 and US23) (Lee *et al.*, 2020; Lin *et al.*, 2020a) or cloned by PCR (EC1, Katshaar, Pi14538, Pi88069, Pi99183 and Pi99177) (Fig. S1). The sequence alignment shows *Avramr3* is a highly conserved RXLR effector among *P. infestans* isolates (Fig. S1).

To find the domain responsible for recognition by Rpi-amr3, ten truncated *Avramr3* fragments were cloned (T1 to T10, Fig. 1b, Fig S2) in an expression vector, and transiently co-expressed with *Rpi-amr3* in *N. benthamiana*. We found four AVR_{amr3} truncations (T3, T4, T8 and T10) can be recognized by Rpi-amr3. T10 (111-240 aa) which carries the 2nd and 3rd WY motifs is the minimal region to be recognized by Rpi-amr3 but not the adjacent T9 protein (130-258 aa) (Fig. 1b and 1c). This suggests these 130 amino-acids of AVR_{amr3} T10 are sufficient for recognition by Rpi-amr3 and initiation of HR.

Figure 1. AVRamr3 is the recognized effector for Rpi-amr3.

(a). Co-expression of Rpi-amr3::GFP and AVRamr3::HIS-FLAG trigger cell death on *N. benthamiana*, but expression of Rpi-amr3::GFP or AVRamr3::HIS-FLAG individually do not induce cell death. The photo was taken 3 days after infiltration, *Agrobacterium* strain GV3101(pMP90) carrying Rpi-amr3::GFP or AVRamr3::HIS-FLAG constructs were used in this experiment. OD₆₀₀=0.5. Three biological replicates were performed with same results. (b). Cartoon of AVRamr3 (PITG_21190), a protein with 339 amino acids with a signal peptide (lemon), RXLR-EER motif (green), and an effector domain (red) with four predicted WY motifs (Details are shown in Fig. S4). T1-T10 indicate the AVRamr3 truncations used in HR assays. Those that induce HR after co-expression with Rpi-amr3 are marked by orange bars, otherwise by blue. (c). Co-expression of Rpi-amr3::GFP and AVRamr3 truncations, all truncations are tagged with C-terminal HIS-FLAG tag. T3, T4, T8 and T10 trigger cell death when co-expressed with Rpi-amr3, but not T1, T2, T5, T6, T7 and T9. Full-length AVRamr3::HIS-FLAG was used as control. OD₆₀₀=0.5. Three biological replicates were performed with same results.

Rpi-amr3 is dependent on the helper NLRs NRC2, NRC3 and NRC4

In Solanaceae, the functionality of many CC-NLR proteins requires helper NLR proteins of the NRC class (Wu et al., 2017). To test if *Rpi-amr3* is NRC-dependent, we co-expressed *Rpi-amr3* and *Avramr3* in NRC knockout *N. benthamiana* lines (nrc2/3_1.3.1, nrc4_185.9.1.3, nrc2/3/4_210.4.3)(Adachi et al., 2019; Wu et al., 2020; Witek et al., 2021). As with wild type *N. benthamiana*, we found HR on the nrc2/3_1.3.1 and nrc4_185.9.1.3 knockout lines, but not the nrc2/3/4_210.4.3 knockout line. Similarly, only nrc2/3/4_210.4.3 knockout lines show susceptibility to *P. infestans* after *Rpi-amr3* transient expression (Fig. S3). Therefore, these data suggest both *Rpi-amr3*-mediated effector recognition and resistance require NRC2, NRC3 or NRC4.

Rpi-amr3 associates with AVRamr3 in planta

To date, most Rpi proteins recognize their cognate effectors in an indirect manner, except RB and IPI-O effectors (Chen et al., 2012; Kourelis and van der Hoorn, 2018). To test the interaction between *Rpi-amr3* and *AVRamr3*, *Rpi-amr3::HA* and *AVRamr3::HIS-FLAG* epitope-tagged constructs were generated and transiently co-expressed in nrc2/3/4 knockout *N. benthamiana* leaves to avoid cell death. Protein was then extracted and bi-directional co-immunoprecipitation (Co-IP) was performed. These co-IPs indicate that *Rpi-amr3* associates with *AVRamr3* bidirectionally (Fig. 2a). We also tested their interaction using a split-luciferase assay. *Rpi-amr3::Cluc* and *AVRamr3::Nluc* constructs were transiently expressed in the nrc2/3/4 knockout *N. benthamiana*. Luciferase signal was only detected when *Rpi-amr3::Cluc* and *AVRamr3::Nluc* were co-expressed. It suggests *Rpi-amr3* physically associates with *AVRamr3 in-planta* (Fig. 2b), but not in negative controls. Our data therefore are consistent with direct interaction of *Rpi-amr3* and *AVRamr3* proteins, though do not exclude the possible involvement of additional proteins.

Figure 2. Rpi-amr3 directly interacts with AVRamr3.

(a). Rpi-amr3::HA and AVRamr3::HIS-FLAG constructs were used for bidirectional co-immunoprecipitation experiment, with Rpi-amr1-HA and AVRamr1::HIS-FLAG used as control. After HA pull down of Rpi-amr3::HA or Rpi-amr1::HA, only AVRamr3::HIS-FLAG is associated with Rpi-amr3::HA. After Flag pull down of AVRamr3::HIS-FLAG or AVRamr1-HIS-FLAG, only Rpi-amr3::HA is associated with AVRamr3::HIS-FLAG. *Agrobacterium* strain GV3101(pMP90) carrying different constructs were used for transiently expression in nrc2/3/4 knockout *Nicotiana benthamiana* line (210.4.3) to abolish the cell death phenotype. OD₆₀₀=0.5. Three biological replicates were performed with same results. (b). Rpi-amr3::Cluc and AVRamr3::Nluc constructs were used to test their interaction *in planta*, Rpi-amr1::Cluc and AVRamr1::Nluc were used as controls. The luciferase signal can only be detected upon Rpi-amr3::Cluc and AVRamr3::Nluc co-expression.

Avramr3 orthologs occur in multiple *Phytophthora* species

To study the evolution of *Avramr3* in *Phytophthora* species, we searched for *Avramr3* homologs from published *Phytophthora* and *Hyaloperonospora arabidopsidis* (Hpa) genomes. Surprisingly, we found *Avramr3* homologs in many *Phytophthora* genomes, including *P. parasitica*, *P. cactorum*, *P. palmivora*, *P. pluvialis*, *P. megakarya*, *P. lichii*, *P. ramorum*, *P. lateralis*, *P. sojae*, *P. capsici*, *P. cinnamomi*, and in *H. arabidopsidis*. Most of the *Avramr3* homologs are located at a syntenic locus (Fig. 3a). Notably, the *P. infestans* *Avramr3*-containing contig was not fully assembled; sequences are missing on the 5' side of *Avramr3* (Fig. 3a). The protein alignment of the thirteen AVRamr3 homologs is shown in Fig S4.

Figure 3. AVRamr3 is a conserved effector among different *Phytophthora* species.

(a). The synteny map of *AvrRamr3* loci from twelve different *Phytophthora* genomes. The *AvrRamr3* loci were extracted from different genomes, annotated by the gene prediction tool in EumicrobeDB, then analyzed and visualized by Clinker. *AvrRamr3* homologs are shown by purple triangles and indicated by a black arrow, the flanking genes with homology are represented by the corresponding colours. The *Phytophthora* clades are adapted from *Phytophthora* database (Rahman et al., 2014). (b). Expression of AVRamr3 homologs with HIS-FLAG tag alone does not trigger cell death on *Nicotiana benthamiana*. *Agrobacterium* strain GV3101(pMP90) carrying different constructs were used in this experiment. OD₆₀₀=0.5. Three biological replicates were performed with same results.

(c). Co-expression of AVRamr3 homologs with Rpi-amr3::GFP in *N. benthamiana*. The AVRamr3 homologs from *Phytophthora infestans* (Pi), *P. parasitica* (Pp), *P. cactorum* (Pc), *P. palmivora* (Ppal), *P. megakarya* (Pmeg), *P. litchi* (Plit), *P. sojae* (Ps), *P. lateralis* (Plat) and *P. pluvialis* (Pplu) induce cell death after co-expression with Rpi-amr3::GFP, but not AVRamr3 homologs from *P. ramorum* (Pr), *P. capsici* (Pcap) and *Hyaloperonospora arabidopsidis* (Hpa). The AVRamr3 homolog from *P. cinnamomi* (Pcin) shows an intermediate cell death. *Agrobacterium* strain GV3101(pMP90) carrying different constructs were used in this experiment. OD₆₀₀=0.5. Three biological replicates were performed with same results.

To test if those AVRamr3 homologs from different *Phytophthora* species are also recognized by Rpi-amr3, we synthesized and cloned them into an expression vector with the 35S promoter, and performed transient expression assays in *N. benthamiana*. Expressing the effectors alone

does not trigger HR in *N. benthamiana* (Fig. 3b), but AVR_{amr3} homologs from *P. parasitica*, *P. cactorum*, *P. palmivora*, *P. megakarya*, *P. lichii*, *P. sojae*, *P. lateralis* and *P. pluvialis* can induce HR when co-expressed with Rpi-*amr3*. The AVR_{amr3} homolog from *P. cinnamomi* triggers an intermediate HR, and the AVR_{amr3} homologs from *P. ramorum*, *P. capsici*, and *H. arabidopsidis* (Fig 3c) do not trigger Rpi-*amr3*-dependent HR.

To test if particularly conserved amino-acids of AVR_{amr3} are responsible for the Rpi-*amr3* recognition, we mutated eight conserved amino-acid on the AVR_{amr3} T10 region (Figure S4). However, all tested mutants are still recognized by Rpi-*amr3* (Figure S5). This result indicates the recognition specificity might not be determined by any single amino acid on AVR_{amr3}, but rather by its overall structure.

To test if other recognized AVR_{amr3} homologs also directly interact with Rpi-*amr3*, we performed co-immunoprecipitation and split-luciferase assays in *nrc2/3/4_210.4.3* knockout lines. We found all the recognized AVR_{amr3} homologs associate with Rpi-*amr3* by co-immunoprecipitation, though with varied affinity. Two unrecognized AVR_{amr3} homologs from *P. capsici* and *H. arabidopsidis* do not associate with Rpi-*amr3*. However, two unrecognized or weakly recognized AVR_{amr3} homologs from *P. ramorum* and *P. cinnamomi* also associate with Rpi-*amr3*, and the unrecognized AVR_{amr3}-T9 truncation shows a weak association (Figure S6). In contrast, the output of split-luciferase assay is fully consistent with the HR assay (Figure S7). Our data indicating that an *in-planta* receptor-ligand interaction is necessary but might not be sufficient for the activation of Rpi-*amr3* and triggering of HR.

Rpi-amr3* confers resistance to multiple *P. parasitica* and *P. palmivora* strains in *N. benthamiana

Rpi-amr3 was previously reported to confer resistance against potato late blight caused by *P. infestans* (Witek et al., 2016). Its broad effector recognition capacity suggested *Rpi-amr3* might confer resistance against additional *Phytophthora* pathogens.

To test this hypothesis, we generated *Rpi-amr3* stable transformed *N. benthamiana* lines. Two homozygous T2 lines #13.3 and #16.5 were verified to confer *P. infestans* resistance and evaluated for *P. parasitica* and *P. palmivora* resistance. Both these pathogens have a wide host range, including the model plant *N. benthamiana*.

Six *P. parasitica* isolates (R0, R1, 310, 666, 329 and 721) were tested on *N. benthamiana* carrying *Rpi-amr3*, and on wild type *N. benthamiana* plants as negative control. A suspension of zoospores was used for root inoculation (Material and Methods). We found both *N. benthamiana* – *Rpi-amr3* lines resist three *P. parasitica* isolates R1, 666 and 721, but are susceptible to R0, 310 and 329 (Fig 4). In summary, *Rpi-amr3* confers resistance against three out of six tested *P. parasitica* isolates in *N. benthamiana*.

Figure 4. Root inoculation of six *Phytophthora parasitica* isolates on *Rpi-amr3* transgenic *Nicotiana benthamiana* lines.

Representative photos for the *P. parasitica* root inoculation tests are shown. Two homozygous *N. benthamiana* - *Rpi-amr3* lines #13.3 and #16.5 were used in this experiment. Wild type *N. benthamiana* plants were used as control. Six *P. parasitica* isolates were used for root inoculation, *Rpi-amr3* confers resistance against R1, 666 and 721, but not R0, 310 and 329. 3-4 weeks *N. benthamiana* were used for the root inoculation, 3 plants/line were used for each experiment and as least three biological replicates were performed with similar results. The numbers indicate susceptible plants/total tested plants.

The *PpAvramr3* homologs from the six *P. parasitica* isolates were PCR amplified, sub-cloned and sequenced. *PpAvramr3* homologs were identified from R0, R1 and 310, 666 and 721, but

not from 329 (Fig S8). These data suggest the presence of recognized AVR_{amr3} homologs from the *Phytophthora* pathogens is necessary but not sufficient to induce Rpi-amr3 mediated resistance.

Furthermore, we tested another broad host range *Phytophthora* pathogen, *P. palmivora*, which causes major losses on many tropical tree crops like papaya, mango, cacao, coconut and palm tree. We tested seven *P. palmivora* isolates on the two Rpi-amr3 transgenic *N. benthamiana* lines by root inoculation, and wild type *N. benthamiana* was used as a control. We found Rpi-amr3 confers resistance to three out of seven tested *P. palmivora* isolates, including 7551, 7547, 7545, but not to 3914, 7548. For two other isolates 0113 and 3738, inconsistent results were obtained from the two Rpi-amr3 transgenic lines (Fig 5). To verify the presence of AVR_{amr3} homologs in these tested *P. palmivora* isolates, we PCR amplified the AVR_{amr3} homologs from genomic DNA of the seven *P. palmivora* isolates. All the tested *P. palmivora* carry PpalAVR_{amr3} variants (Figure S9). Taken together, Rpi-amr3 confers resistance to at least 3/7 tested *P. palmivora* isolates in the root inoculation assay.

Figure 5. Root inoculation of 7 *Phytophthora palmivora* isolates on Rpi-amr3 transgenic *Nicotiana benthamiana* lines.

Two homozygous *N. benthamiana* - *Rpi-amr3* lines #13.3 and #16.5 were used in this experiment, wild type *N. benthamiana* were used as control. Seven *P. parasitica* isolates were used for root inoculation, *Rpi-amr3* confer resistance against isolates 7547, 7551 and 7545, but not 3914, 7548. For isolates 0113 and 3738, we obtained some variable results for the two transgenic lines. 3-4 old weeks *N. benthamiana* were used for the root inoculation, 3 plants/line were used for each experiment and three or more biological replicates were performed with similar results.

Rpi-amr3* is widely distributed in *S. americanum* and *S. nigrum

Though susceptible accessions can be identified in detached leaf assays, most *S. americanum* and *S. nigrum* accessions show complete resistance in the field to *P. infestans*. Previously, many functional *Rpi-amr1* alleles were cloned from different *S. americanum* and *S. nigrum* accessions (Witek et al., 2021).

The identification of AVR_{amr3} allows us to investigate the distribution of *Rpi-amr3* from all *S. americanum* and *S. nigrum* accessions. In total, 54 *S. americanum* accessions and 26 *S. nigrum* accessions were tested by agro-infiltration with AVR_{amr3} for detecting functional *Rpi-amr3*. We found 43/54 tested *S. americanum* accessions show HR after AVR_{amr3} agro-infiltration (Fig 6a). Similarly, 21/26 tested *S. nigrum* accessions recognize AVR_{amr3} (Fig 6b).

To further investigate the sequence polymorphism of *Rpi-amr3* from different accessions, the *Rpi-amr3* homologs from 14 accessions were extracted from PacBio RenSeq dataset (Witek et al., 2021), including eleven accessions (SP1123, SP2272, SP2273, SP2307, SP2360, SP3399, SP3400, SP1101, SP3406 and SP3409) which respond to AVR_{amr3} and three accessions (SP1032, SP2271 and SP2275) that do not respond to AVR_{amr3}.

To test the functionality of *Rpi-amr3* from *S. americanum* and *S. nigrum*, *Rpi-amr3* homologs were PCR amplified from gDNA of three *S. americanum* accessions SP2272, SP2273 and SP3406, and from gDNA of two *S. nigrum* accessions SP1088 and SP1084. *Rpi-amr3* alleles (*Rpi-nig3* hereafter) were amplified from each of these two *S. nigrum* accessions and cloned into an expression vector with 35S promoter. We found all the seven *Rpi-amr3*/*Rpi-nig3* genes can recognize AVR_{amr3} in transient assays (Fig 6d), but not the negative control AVR_{amr1}. Compared to *Rpi-amr3* from SP1102, the amino-acid identity ranges from 82.2% to 95.7% (Fig 6c). Premature stop codons were found in *Rpi-amr3* homologs from SP2271 and SP2275 (Figure S10), which result in loss of *Rpi-amr3* function.

Taken together, these data suggest *Rpi-amr3* gene is widely distributed in diploid *S. americanum* and hexaploid *S. nigrum*, and contributes to their resistance to *P. infestans* and perhaps other *Phytophthora* pathogens.

Figure 6. Screen for AVRamr3 recognition on *S. americanum* and *S. nigrum* accessions.

(a). 54 *S. americanum* accessions were screened with *Agrobacterium* strain GV3101(pMP90) carrying 35S::AVRamr3. The accessions with cell death upon agro-infiltration are marked by red, otherwise blue. (b). 26 *S. nigrum* accessions were screened with *Agrobacterium* strain GV3101(pMP90) carrying 35S::AVRamr3. The accessions with cell death upon agro-infiltration are marked by red, otherwise blue. (c). Maximum likelihood (ML) tree of Rpi-amr3 and Rpi-nig3 proteins was made by iqtree with GTT+G4 model. The *Rpi-amr3* homologs from *S. americanum* were extracted from PacBio RenSeq assemblies (Witek et al., 2021). The four *Rpi-nig3* genes were PCR amplified from *S. nigrum* accession SP1088 and SP1084 (red). The non-functional *Rpi-amr3* homologs were marked by blue. Rpi-amr3b from SP1102 is a paralogue of Rpi-amr3, which was used as an outgroup of the phylogenetic analysis. The scale bar indicates the number of amino acid substitutions per site. The protein identities of each homolog compared to Rpi-amr3 (Rpi-amr3-1102) are shown by %. (d). Selected Rpi-amr3 homologs were cloned from three *S. americanum* accessions SP2272, SP2273, SP3406. Four Rpi-nig3 homologs were cloned from *S. nigrum* accessions SP1088 and SP1084, and co-expressed with AVRamr3 or AVRamr1 (negative control). All of them can recognize AVRamr3 in the transient assay but not AVRamr1.

Discussion

In this study, by screening an RXLR effector library of *Phytophthora infestans*, we identified and characterized a novel effector AVRamr3 (PITG_21190) that is recognized by the NLR protein Rpi-amr3 of *S. americanum*. AVRamr3 is very conserved among all tested *P. infestans* isolates, and AVRamr3 homologs were identified in twelve additional *Phytophthora* and *Hyaloperonospora arabidopsidis* genomes. These homologs are located in a syntenic region (Fig. 3a). Surprisingly, we found 9/13 tested AVRamr3 homologs can be recognized by Rpi-amr3 leading to HR in *N. benthamiana*. This finding suggests AVRamr3 is an essential effector among *Phytophthora* species, though its virulence function has yet to be determined.

According to the “zigzagzig” model of plant immunity (Jones and Dangl, 2006), the surface immune receptors like receptor-like proteins (RLPs) and receptor-like kinases (RLKs) perceive relatively conserved microbe-associated molecular patterns (MAMPs) and induce pattern-triggered immunity (PTI). Intracellular nucleotide-binding and leucine-rich repeat immune receptors (NLRs) recognize fast-evolving and lineage-specific effectors and activate effector-triggered immunity (ETI). Therefore, PTI was believed to confer broader-spectrum resistance compared to ETI. Indeed, many RLPs/RLKs recognize conserved ligands and /or confer broad-spectrum resistance, such as FLS2, EFR, RLP23, RXEG1 and ELR (Zipfel et al., 2006; Albert et al., 2015; Du et al., 2015; Wang et al., 2018). Remarkably, EFR from *Arabidopsis thaliana* enhances resistance against a range of bacterial pathogens in different crop plants, like tomato, orange and apple (Lacombe et al., 2010; Mitre et al., 2021; Piazza et al., 2021). However, apoplastic effectors can also be fast-evolving proteins, like the SCR74 family in *P. infestans* (Liu et al., 2005; Lin et al., 2020b), or *Cladosporium fulvum* AVR2, AVR4 and AVR9 (Joosten et al., 1994; Van den Ackerveken et al., 1994; Luderer et al., 2002; Westerink et al., 2004). On the other hand, MAMP-like cytoplasmic effectors/effector epitopes have been reported. For example, Sw-5b from tomato confers broad-spectrum tospovirus resistance by recognizing a conserved, 21-amino acid epitope NSm²¹ which derives from the viral movement protein NSm (Zhu et al., 2017). A recent functional pan-genome study revealed the ETI landscape of *A. thaliana* and *Pseudomonas syringae*; some *P. syringae* effectors are widely conserved. Similarly, the ETI mediated by two conserved NLRs CAR1 and ZAR1 confers resistance to 94.7% *P. syringae* strains (Laflamme et al., 2020). These observations, as well as our finding on AVRamr3 and Rpi-amr3, all support the view that pathogen molecules recognized by NLRs

can also be relatively invariant and conserved, and might contribute to broad-spectrum pathogen resistance.

To test this hypothesis, we established a *N. benthamiana* root inoculation system by using stable *Rpi-amr3* transgenic plants, and tested *P. parasitica* and *P. palmivora* which have a broad host range, and cause dramatic yield losses of many crops from different families (Meng et al., 2014; Ali et al., 2017). Importantly, we found *Rpi-amr3* does confer resistance against some *P. parasitica* and *P. palmivora* isolates. This is the first report of cloned *R* genes against *P. parasitica* and *P. palmivora* (Kourelis et al., 2021). Additionally, it is noteworthy that, in nature, many *Phytophthora* pathogens can co-inoculate the host and interspecific hybridization might occur; for example, *P. andina* was proposed to have emerged through hybridization of *P. infestans* and an unknown *Phytophthora* species (Goss et al., 2011). Natural hybrids of *P. parasitica* and *P. cactorum* were also found on infected loquat trees (Hurtado-Gonzales et al., 2017). An *R* protein that provides protection against both foliar and root *Phytophthora* pathogens of different species would be extremely valuable. However, some *Rpi-amr3* breaking *P. parasitica* and *P. palmivora* strains were also identified in our study, although most of them carry the recognized AVR_{amr3} homologs. This might be caused by silencing of the recognized effector gene like *Avrvnt1* to avoid the recognition by Rpi-vnt1, or presence of other suppressors or regulators like Avr_{cap1b} or splicing regulatory (SRE) effectors (Pais et al., 2018; Huang et al., 2020; Derevnina et al., 2021).

In this study, we also reported that Rpi-amr3 directly interacts with AVR_{amr3} and other recognized AVR_{amr3} homologs from different *Phytophthora* species. Surprisingly, the direct interaction has not led to accelerated evolution of AVR_{amr3} to evade detection, as we also observed for Rpi-amr1 and AVR_{amr1} (Lin et al., 2020a; Witek et al., 2021). This could predispose *Rpi-amr3* to function in different plant species.

Thus, *Rpi-amr3* could be deployed in Solanaceae crops like potato, tomato and tobacco against multiple *Phytophthora* diseases. However, interfamily transfer of *NLR* genes remains a challenge if *NLR* genes show “restricted taxonomic functionality” (Tai et al., 1999). The paired *NLR* genes *RPS4/RRS1* from Brassicaceae (*Arabidopsis*) can nevertheless function in other plant families like Solanaceae (tomato) and Cucurbitaceae (cucumber) against different bacterial and fungal diseases (Narusaka et al., 2013). Here, we found any of the NRC2, NRC3 or NRC4 proteins are required for Rpi-amr3 to execute its function. Thus, in the plant families

which lacking NRC homologs, such as tropical tree crops susceptible to *P. palmivora* and *P. megakarya*, co-delivery of *Rpi-amr3* and *NRC* genes might be required to defeat these *Phytophthora* diseases.

“Non-host” resistance is durable. *S. americanum* and *S. nigrum* are thought to be non-host plants of *P. infestans*, although susceptible accessions of both species have been found using DLAs. This opens the opportunity to dissect their “non-host” resistance. By using AVR_{amr3} as a probe, we found *Rpi-amr3* is widely distributed in *S. americanum* and *S. nigrum* species (Witek et al., 2021) (Fig. 6). We noticed that PITG_21190 (AVR_{amr3}) triggers HR in many *S. nigrum* accessions in a large-scale effector screening study (Dong, 2016), consistent with our findings (Fig. 6). Furthermore, we cloned four *Rpi-nig3* genes from two *S. nigrum* accessions. All the four *Rpi-nig3* homologs recognize AVR_{amr3} in our co-expression assays (Fig. 6), although their resistance to late blight needs to be evaluated individually. The wide distribution of *Rpi-amr3* and *Rpi-amr1* suggests that these two *R* genes, perhaps with other *R* genes and the NRC network in *S. americanum* and *S. nigrum*, underpin their “non-host” resistance against potato late blight. The identification of AVR_{amr3} and AVR_{amr1} can also help to explore other novel resistance genes from *S. americanum* and *S. nigrum*.

In summary, this study reveals that *Rpi-amr3* is a conserved and broad-spectrum *R* gene from *S. americanum* and its relatives. The recognition of the conserved AVR_{amr3} effectors leads to resistance against several different *Phytophthora* pathogens. This finding shows great potential for resistance breeding in many crop plants against different *Phytophthora* diseases.

References

- Adachi H, Contreras MP, Harant A, Wu C-H, Derevnina L, Sakai T, Duggan C, Moratto E, Bozkur TO, Maqbool A, et al** (2019) An N-terminal motif in NLR immune receptors is functionally conserved across distantly related plant species. *eLife* **8**: 121
- Albert I, Böhm H, Albert M, Feiler CE, Imkampe J, Wallmeroth N, Brancato C, Raaymakers TM, Oome S, Zhang H, et al** (2015) An RLP23–SOBIR1–BAK1 complex mediates NLP-triggered immunity. *NPLANTS* **1**: 15140
- Ali SS, Shao J, Lary DJ, Strem MD, Meinhardt LW, Bailey BA** (2017) *Phytophthora megakarya* and *P. palmivora*, causal agents of black pod rot, induce similar plant defense responses late during infection of susceptible Cacao pods. *Frontiers in Plant Science* **8**: 36
- Bao Y, Ding N, Qin Q, Wu X, Martinez N, Miller R, Zaitlin D, Li D, Yang S** (2019) Genetic mapping of the *Ph* gene conferring disease resistance to black shank in tobacco. *Mol Breeding* 1–10
- Cooke DEL, Cano LM, Raffaele S, Bain RA, Cooke LR, Etherington GJ, Deahl KL, Farrer RA, Gilroy EM, Goss EM, et al** (2012) Genome analyses of an aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathog* **8**: e1002940
- Derevnina L, Contreras MP, Adachi H, Upson J, Cruces AV, Xie R, Sklenar J, Menke FLH, Mugford ST, Maclean D, et al** (2021) Plant pathogens convergently evolved to counteract redundant nodes of an NLR immune receptor network. *Biorxivorg*: 2021.02.03.429184
- Dong R** (2016) Identification of the function of four genes encoding the RxLR secreted protein in *Phytophthora infestans*. PhD thesis. Shandong Agricultural University
- Du J, Verzaux E, Chaparro-Garcia A, Bijsterbosch G, Keizer LCP, Zhou J, Liebrand TWH, Xie C, Govers F, Robatzek S, et al** (2015) Elicitin recognition confers enhanced resistance to *Phytophthora infestans* in potato. *NPLANTS* 10.1038–nplants.2015.34
- Gallup CA, Shew HD** (2010) Occurrence of race 3 of *Phytophthora nicotianae* in North Carolina, the causal agent of black shank of tobacco. *Plant Disease* **94**: 557–562
- Gilroy EM, Breen S, Whisson SC, Squires J, Hein I, Kaczmarek M, Turnbull D, Boevink PC, Lokossou A, Cano LM, et al** (2011) Presence/absence, differential expression and sequence polymorphisms between *PiAVR2* and *PiAVR2*-like in *Phytophthora infestans* determine virulence on *R2* plants. *New Phytol* **191**: 763–776
- Goss EM, Cardenas ME, Myers K, Forbes GA, Fry WE, Restrepo S, Grünwald NJ** (2011) The plant pathogen *Phytophthora andina* emerged via hybridization of an unknown *Phytophthora* species and the Irish potato famine pathogen, *P. infestans*. *PLoS ONE* **6**: e24543
- Huang J, Lu X, Wu H, Xie Y, Peng Q, Gu L, Wu J, Wang Y, Reddy ASN, Dong S** (2020) *Phytophthora* effectors modulate genome-wide alternative splicing of host mRNAs to reprogram plant immunity. *Molecular Plant* **13**: 1470–1484

- 445 **Hurtado-Gonzales OP, Aragon-Caballero LM, Flores-Torres JG, Veld WMIT, Lamour**
 446 **KH** (2017) Molecular comparison of natural hybrids of *Phytophthora nicotianae* and *P.*
 447 *cactorum* infecting loquat trees in Peru and Taiwan. *Mycologia* **101**: 496–502
- 448 **Jiang RHY, Tripathy S, Govers F, Tyler BM** (2008) RXLR effector reservoir in two
 449 *Phytophthora* species is dominated by a single rapidly evolving superfamily with more
 450 than 700 members. *Proceedings of the National Academy of Sciences* **105**: 4874–4879
- 451 **Jones JDG, Dangl JL** (2006) The plant immune system. *Nature* **444**: 323–329
- 452 **Joosten MHAI, Cozijnsen TJ, de Wit PJGM** (1994) Host resistance to a fungal tomato
 453 pathogen lost by a single base-pair change in an avirulence gene. *Nature* **367**: 384
- 454 **Kamoun S, Furzer O, Jones JDG, Judelson HS, Ali GS, Dalio RJD, Roy SG, Schena L,**
 455 **Zambounis A, Panabières F, et al** (2015) The Top 10 oomycete pathogens in molecular
 456 plant pathology. *Molecular Plant Pathology* **16**: 413–434
- 457 **Kourelis J, Sakai T, Adachi H, Kamoun S** (2021) RefPlantNLR: a comprehensive collection
 458 of experimentally validated plant NLRs. *Biorxivorg*: 2020.07.08.193961
- 459 **Lacombe S, Rougon-Cardoso A, Sherwood E, Peeters N, Dahlbeck D, van Esse HP,**
 460 **Smoker M, Rallapalli G, Thomma BPHJ, Staskawicz B, et al** (2010) Interfamily
 461 transfer of a plant pattern-recognition receptor confers broad-spectrum bacterial resistance.
 462 *Nature Biotechnology* **28**: 365
- 463 **Laflamme B, Dillon MM, Martel A, Almeida RND, Desveaux D, Guttman DS** (2020) The
 464 pan-genome effector-triggered immunity landscape of a host-pathogen interaction. *Science*
 465 **367**: 763–768
- 466 **Lin X, Song T, Fairhead S, Witek K, Jouet A, Jupe F, Witek AI, Karki HS, Vleeshouwers**
 467 **VGAA, Hein I, et al** (2020a) Identification of Avramr1 from *Phytophthora infestans* using
 468 long read and cDNA pathogen-enrichment sequencing (PenSeq). *Molecular Plant*
 469 *Pathology* **18**: 547–11
- 470 **Lin X, Wang S, de Rond L, Bertolin N, Wouters RHM, Wouters D, Domazakis E, Bitew**
 471 **MK, Win J, Dong S, et al** (2020b) Divergent evolution of PcF/SCR74 effectors in
 472 oomycetes is associated with distinct recognition patterns in Solanaceous plants. *MBio* **11**:
 473 380
- 474 **Liu Z, Bos JIB, Armstrong M, Whisson SC, da Cunha L, Torto-Alalibo T, Win J, Avrova**
 475 **AO, Wright F, Birch PRJ, et al** (2005) Patterns of diversifying selection in the
 476 phytotoxin-like *scr74* gene family of *Phytophthora infestans*. *Mol Biol Evol* **22**: 659–672
- 477 **Luderer R, Takken FLW, de Wit PJGM, Joosten MHAI** (2002) *Cladosporium fulvum*
 478 overcomes Cf-2-mediated resistance by producing truncated AVR2 elicitor proteins.
 479 *Molecular Microbiology* **45**: 875–884
- 480 **Meng Y, Zhang Q, Ding W, Shan W** (2014) *Phytophthora parasitica*: a model oomycete
 481 plant pathogen. *Mycology* **5**: 43–51
- 482 **Mitre LK, Teixeira-Silva NS, Rybak K, Magalhães DM, de Souza-Neto RR, Robatzek S,**
 483 **Zipfel C, de Souza AA** (2021) The *Arabidopsis* immune receptor EFR increases resistance

to the bacterial pathogens *Xanthomonas* and *Xylella* in transgenic sweet orange. Plant Biotechnology J. doi: 10.1111/pbi.13629

Narusaka M, Kubo Y, Hatakeyama K, Imamura J, Ezura H, Nanasato Y, Tabei Y, Takano Y, Shirasu K, Narusaka Y (2013) Interfamily transfer of dual *NB-LRR* genes confers resistance to multiple pathogens. PLoS ONE 8: e55954

Oliva RF, Cano LM, Raffaele S, Win J, Bozkur TO, Belhaj K, Oh S-K, THINES M, Kamoun S (2015) A recent expansion of the RXLR effector gene *Avrblb2* is maintained in global populations of *Phytophthora infestans* indicating different contributions to virulence. MPMI 28: 901–912

Pais M, Yoshida K, Giannakopoulou A, Pel MA, Cano LM, Oliva RF, Witek K, Lindqvist-Kreuzer H, Vleeshouwers VGAA, Kamoun S (2018) Gene expression polymorphism underpins evasion of host immunity in an asexual lineage of the Irish potato famine pathogen. BMC Evol Biol 18: 1–11

Piazza S, Campa M, Pompili V, Costa LD, Salvagnin U, Nekrasov V, Zipfel C, Malnoy M (2021) The *Arabidopsis* pattern recognition receptor EFR enhances fire blight resistance in apple. Biorxivorg: 2021.01.22.427734

Rahman MZ, Uematsu S, Takeuchi T, Shirai K, Ishiguro Y, Suga H, Kageyama K (2014) Two new species, *Phytophthora nagaii* sp. nov. and *P. fragariaefolia* sp. nov., causing serious diseases on rose and strawberry plants, respectively, in Japan. J Gen Plant Pathol 80: 348–365

Rehmany AP, Gordon A, Rose LE, Allen RL, Armstrong MR, Whisson SC, Kamoun S, Tyler BM, Birch PRJ, Beynon JL (2005) Differential recognition of highly divergent downy mildew avirulence gene alleles by *RPPI* resistance genes from two *Arabidopsis* lines. The Plant Cell 17: 1839–1850

Sahoo DK, Abeysekara NS, Cianzio SR, Robertson AE, Bhattacharyya MK (2017) A novel *Phytophthora sojae* resistance *Rps12* gene mapped to a genomic region that contains several *Rps* genes. PLoS ONE 12: e0169950

Segretin ME, Pais M, Franceschetti M, Chaparro-Garcia A, Bos JIB, Banfield MJ, Kamoun S (2014) Single amino acid mutations in the potato immune receptor R3a expand response to *Phytophthora* effectors. MPMI 27: 624–637

Shan W, Cao M, Leung D, Tyler BM (2004) The *Avr1b* locus of *Phytophthora sojae* encodes an elicitor and a regulator required for avirulence on soybean plants carrying resistance gene *Rps1b*. MPMI 17: 394–403

Tai TH, Dahlbeck D, Clark ET, Gajiwala P, Pasion R, Whalen MC, Stall RE, Staskawicz BJ (1999) Expression of the *Bs2* pepper gene confers resistance to bacterial spot disease in tomato. Proceedings of the National Academy of Sciences 96: 14153–14158

Thevenin J-M, Rossi V, Ducamp M, Doare F, Condina V, Lachenaud P (2012) Numerous clones resistant to *Phytophthora palmivora* in the “Guiana” genetic group of *Theobroma cacao* L. PLoS ONE 7: e40915

- 523 **Van den Ackerveken GFJM, Dunn RM, Cozijnsen AJ, Vossen JPMJ, Van den Broek**
524 **HWJ, de Wit PJGM** (1994) Nitrogen limitation induces expression of the avirulence gene
525 *avr9* in the tomato pathogen *Cladosporium fulvum*. *Mol Gen Genet* **243**: 277–285
- 526 **Vega-Arreguín JC, Jalloh A, Bos JL, Moffett P** (2014) Recognition of an Avr3a homologue
527 plays a major role in mediating nonhost resistance to *Phytophthora capsici* in *Nicotiana*
528 species. *MPMI* **27**: 770–780
- 529 **Vleeshouwers VGAA, Raffaele S, Vossen JH, Champouret N, Oliva R, Segretin ME,**
530 **Rietman H, Cano LM, Lokossou A, Kessel G, et al** (2011) Understanding and exploiting
531 late blight resistance in the age of effectors. *Annu Rev Phytopathol* **49**: 507–531
- 532 **Wang S, McLellan H, Bukharova T, He Q, Murphy F, Shi J, Sun S, van Weymers P, Ren**
533 **Y, Thilliez G, et al** (2019) *Phytophthora infestans* RXLR effectors act in concert at diverse
534 subcellular locations to enhance host colonization. *Journal of Experimental Botany* **70**:
535 343–356
- 536 **Wang Y, Xu Y, Sun Y, Wang H, Qi J, Wan B, Ye W, Lin Y, Shao Y, Dong S, et al** (2018)
537 Leucine-rich repeat receptor-like gene screen reveals that *Nicotiana* RXEG1 regulates
538 glycoside hydrolase 12 MAMP detection. *Nat Commun* **9**: 594–12
- 539 **Westerink N, Brandwagt BF, de Wit PJGM, Joosten MHAI** (2004) *Cladosporium fulvum*
540 circumvents the second functional resistance gene homologue at the *Cf-4* locus (*Hcr9-4E*)
541 by secretion of a stable *avr4E* isoform. *Molecular Microbiology* **54**: 533–545
- 542 **Win J, Krasileva KV, Kamoun S, Shirasu K, Staskawicz BJ, Banfield MJ** (2012) Sequence
543 divergent RXLR effectors share a structural fold conserved across plant pathogenic
544 oomycete Species. *PLoS Pathog* **8**: e1002400
- 545 **Witek K, Jupe F, Witek AI, Baker D, Clark MD, Jones JDG** (2016) Accelerated cloning of
546 a potato late blight–resistance gene using RenSeq and SMRT sequencing. *Nature*
547 *Biotechnology* **34**: 656–660
- 548 **Witek K, Lin X, Karki HS, Jupe F, Witek AI, Steuernagel B, Stam R, van Oosterhout C,**
549 **Fairhead S, Heal R, et al** (2021) A complex resistance locus in *Solanum americanum*
550 recognizes a conserved *Phytophthora* effector. *NPLANTS* **7**: 198–208
- 551 **Wu C-H, Abd-El-Haliem A, Bozkur TO, Belhaj K, Terauchi R, Vossen JH, Kamoun S**
552 (2017) NLR network mediates immunity to diverse plant pathogens. *Proceedings of the*
553 *National Academy of Sciences* **114**: 8113–8118
- 554 **Wu C-H, Adachi H, la Concepcion De JC, Castells-Graells R, Nekrasov V, Kamoun S**
555 (2020) *NRC4* Gene cluster Is not essential for bacterial flagellin-triggered immunity.
556 *PLANT PHYSIOLOGY* **182**: 455–459
- 557 **Zhu M, Jiang L, Bai B, Zhao W, Chen X, Li J, Liu Y, Chen Z, Wang B, Wang C, et al**
558 (2017) The intracellular immune receptor Sw-5b confers broad-spectrum resistance to
559 tospoviruses through recognition of a conserved 21-amino acid viral effector epitope. *The*
560 *Plant Cell* **29**: 2214–2232

561 **Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JDG, Boller T, Felix G (2006)**
 562 Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts *Agrobacterium*-
 563 mediated transformation. *Cell* **125**: 749–760

564

Acknowledgements

This research was financed from BBSRC grant BB/P021646/1 and the Gatsby Charitable Foundation. We thank TSL transformation team (Matthew Smoker and Jodie Taylor), SynBio team (Mark Youles) and horticultural team (Sara Perkins, Justine Smith, Lesley Phillips and Catherine Taylor) for their support. We thank Experimental Garden and Genebank of Radboud University, Nijmegen, The Netherlands, IPK Gatersleben, Germany and Sandra Knapp (Natural History Museum, London, UK) for access to *S. americanum* and *S. nigrum* genetic diversity. We thank He Meng and Lirui Cheng from CAAS for kindly sharing the *Phytophthora parasitica* isolates R0 and R1, and Franck Panabières from INRA for kindly sharing the *Phytophthora parasitica* isolates 310, 329, 666 and 721. We thank Joe Win from TSL for maintaining the *P. palmivora* strains. We thank Paul Birch and colleagues at James Hutton Institute for making available clones of some of the effectors that were tested for AVR_{amr3} function.

Author contributions:

X.L. and J.D.G.J. designed the study. X.L., A.C.O.A., R.H., K.W., H.S.K., T.S., C.-H.W. and H.A. performed the experiments. X.L., A.C.O.A., R.H. and K.W. analysed the data. X.L. and J.D.G.J. wrote the manuscript with input from all authors. S.K. and V.G.A.A.V. contributed resources. All authors approved the manuscript.

Conflict of interest:

K. W. and J.D.G.J. are named inventors on a patent application (PCT/US2016/031119) pertaining to *Rpi-amr3* that was filed by the 2Blades Foundation on behalf of the Sainsbury Laboratory. The other authors declare no competing interests.

Supplementary files:

Figure S1: Protein alignment reveals strong conservation of *P. infestans* AVRamr3 alleles and paralogs

Figure S2: Western blot for AVRamr3 truncations with C-terminus HIS-FLAG tag.

Figure S3: Rpi-amr3 is NRC2, NRC3 or NRC4 dependent. (a) HR phenotype after expressing Rpi-amr3, AVRamr3 and Rpi-amr3+AVRamr3 in wild type, nrc2/3_1.3.1, nrc4_185.9.1.3 and nrc2/3/4_210.4.3 knockout *N. benthamiana* lines. (b) Detached leaf assay (DLA) after *Rpi-amr3* transient expression in wild type, nrc2/3_1.3.1, nrc4_185.9.1.3 and nrc2/3/4_210.4.3 knockout *N. benthamiana*. 500 zoospores of *P. infestans* 88069 were used one day after *Rpi-amr3* transient expression by agro-infiltration. The photos were taken eight days after inoculation.

Figure S4: Protein alignment of AVRamr3 homologs from different *Phytophthora* genomes, including *Phytophthora infestans* (Pi), *Phytophthora parasitica* (Pp), *Phytophthora cactorum* (Pc), *Phytophthora palmivora* (Ppal), *Phytophthora megakarya* (Pmeg), *Phytophthora litchi* (Plit), *Phytophthora sojae* (Ps), *Phytophthora lateralis* (Plat), *Phytophthora pluvialis* (Pplu), *Phytophthora ramorum* (Pr), *P. cinnamomi* (Pcin), *P. capsica* (Pcap) and *Hyaloperonospora arabidopsidis* (Hpa). The circles after the name are their recognition specificity by Rpi-amr3 in the HR assay, red: HR; black: no HR; yellow: weak HR. The RXLR and EER motifs are marked by black boxes. The predicted WY motifs are marked by blue boxes. The conserved amino acids which were selected for mutagenesis (see Figure S5) are marked by yellow arrows.

Figure S5: Mutagenesis of AVRamr3 from *Phytophthora infestans*. **(a)** Eight *Avramr3* mutants were generated by PCR and cloned into over-expression vector with 35S promoter. All of them induce HR when co-expressed with Rpi-amr3. An AVRamr3 truncation T9 was used as negative control, full-length AVRamr3 was used as positive control. **(b)** The position of each mutation is marked by asterisk, and correspond to those amino acids marked by yellow arrows in Fig S4. All the mutants are in the T10 region and are conserved among different AVRamr3 homologs from other *Phytophthora* species (see Figure S4).

Figure S6: Co-IP of Rpi-amr3 and AVRamr3 homologs. Rpi-amr3 is tagged with C-terminal GFP, and all AVRamr3 homologs are fused with a C-terminal HIS-FLAG tag. Bi-directional Co-IPs were performed by GFP-IP or Flag-IP individually then incubated with Flag-HRP or GFP-HRP antibody. ATR2 and AVRamr3 truncation T9 were used as negative controls.

Figure S7: Split luciferase for Rpi-amr3 and AVRamr3 homologs from different *Phytophthora* species. Rpi-amr3 was fused with Flag-Cluc and AVRamr3 homologs were fused with Flag-Nluc. The experiment was performed in nrc2/3/4_210.4.3 knockout *Nicotiana benthamiana* to abolish HR. **(a).** Expressing the AVRamr3::Flag-Nluc homologs alone does not show luciferase signal; Co-expression of Rpi-amr3::Flag-Cluc with AVRamr3::Flag-Nluc homologs can induce luciferase signal, specifically, *P. infestans* (Pi), *P. parasitica* (Pp), *P. cactorum* (Pc), *P. palmivora* (Ppal), *P. megakarya* (Pmeg), *P. litchi* (Plit), *P. sojae* (Ps), *P. lateralis* (Plat) and *P. pluvialis* (Pplu) interact with Rpi-amr3::Flag-Cluc; AVRamr3 homologs from *P. ramorum* (Pr), *P. cinnamomi* (Pcin), *P. capsici* (Pcap) and *Hyaloperonospora arabidopsidis* (Hpa) do not interact with Rpi-amr3 in this assay. **(b).** Western blot by FLAG antibody was performed to confirm the expression of all proteins.

Figure S8: DNA alignment of *Avramr3* homologs from different *Phytophthora parasitica* isolates. The polymorphic DNA and amino acids are highlighted.

Figure S9: DNA alignment of *AvrAmr3* homologs from different *Phytophthora palmivora* isolates. The polymorphic DNA and amino acids are highlighted.

665

666

667

668

669