Growth allometry and dental topography in Upper Triassic conodonts supports trophic differentiation and molar-like element function

Valentin Kelz¹, Pauline Guenser¹, Michele Mazza², Manuel Rigo³, Emilia Jarochovska¹

¹ Fachgruppe Paläoumwelt, GeoZentrum Nordbayern, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

² Largo Mazza, 29121, Piacenza, Italy

³ Department of Geosciences, University of Padova, Padova, Italy

Corresponding Author:

Pauline Guenser¹

Fachgruppe Paläoumwelt, GeoZentrum Nordbayern, Friedrich-Alexander-Universität Erlangen-Nürnberg, Loewenichstr. 28, 91054 Erlangen, Germany

Email address: pauline.guenser@gmail.com

Abstract

Conodont dental elements are distinguished by their high disparity and rapid morphological evolution. P₁ elements located in the pharynx are the most rapidly evolving, but their function in the animal has been only investigated in a handful of taxa and proposed to be analogous to mammal molars. This hypothesis predicts that their surface area should show positive allometry with respect to element length, as has been previously identified in 2D projections in two Carboniferous taxa. Here we apply the same method to test this hypothesis in 3D models of platform-bearing P₁ elements of two common Late Triassic taxa, *Metapolygnathus communisti* and *Epigondolella rigoi*. We further hypothesise that these commonly co-occurring taxa differed in their growth allometry, reflecting their different trophic niches. Platform length grew isometrically with respect to element length, whereas log-transformed platform area showed positive allometry with respect to element length, with slopes equal 3.86 in *M. communisti* and 4.16 in *E. rigoi*, supporting a function of the platform analogous to molars and trophic differentiation. We cross-tested the latter interpretation by dental topographic analysis using Dirichlet Normal Energy (DNE). Specimens of the adult growth stage of *E. rigoi* showed higher DNE values than specimens of the same growth stage in *M. communisti*, consistent with stronger positive allometry of platform surface and with a higher demand for energy in this species. DNE values of platform surface increased linearly in function of element length and log-transformed platform area, indicating no ontogenetic changes. Based on DNE values available for primates, those of the adult growth stages were similar to those reported for insectivores or folivores in the case of *E. rigoi* and for folivores or omnivores in the case of *M. communisti*.

Previous studies applying morphological and ultrastructural proxies for the dietary position of conodonts addressed mostly stratigraphically older conodont taxa, but our results indicate that Late Triassic species occupied the predator/scavenger niche in spite of the highly developed diversity of gnathostomes in this niche. We also show that within this broad niche, co-occurring taxa differed in their diets, which supports trophic diversification as an important driver of the remarkable disparity of their elements.

Introduction

Conodonts and their feeding apparatus

Conodonts are extinct, eel-shaped marine animals that lived from the middle Cambrian to Early Jurassic (Du et al., 2020). They are early vertebrates (Donoghue, Purnell & Aldridge, 1998) distinguished by an extensive fossil record (Foote & Sepkoski, 1999; Donoghue, 2001a). The majority of conodont fossil record consists of the phosphatic elements forming the feeding apparatus of the animal (Scott, 1934; Schmidt, 1934; Purnell, Donoghue & Aldridge, 2000). Soft tissues are rarely found and have, so far, not revealed a great diversity of body forms. Because of this, conodont taxonomy and functional morphology are based on their elements (Mazza, Cau & Rigo, 2012).

Evidence of dental function

Several lines of argumentation have been proposed to argue that elements of the conodont feeding apparatus had a tooth-like function. S and M elements, positioned in the anterior part, i.e. the mouth, are interpreted to perform a grasping function, whereas P elements, placed in the posterior part of the apparatus in the pharynx of the animal, have been proposed to have a function similar to molars of mammals (Purnell & von Bitter, 1992; Purnell, 1994; Donoghue & Purnell, 1999a). A model of conodonts being suspension feeders has also been suggested (Nicoll, 1987). In this model, elements would be covered in tissue and S and M elements would filter particles and create current. P₁ elements would only lightly mash food. This model, however, is not supported by more recent research.

Following arguments have been proposed for the dental function of conodont elements: (1) the presence of microwear patterns on conodont elements, produced *in vivo* (Purnell, 1995; Martínez-Pérez et al., 2014c); (2) occlusion models (Donoghue & Purnell, 1999a; Jones et al., 2012a; Martínez-Pérez et al., 2014c,b); (3) Finite Element Analysis (FEA) (Jones et al., 2012b; Martínez-Pérez et al., 2014c, 2016); (4) ultrastructural adaptation of conodont tissues to dental function, analogous to enamel (Donoghue, 2001b; Jones et al., 2012a); (5) in some conodont taxa, extreme sharpness has been proposed to be an adaptation to cutting function in the absence of jaws acting as levers (Jones et al., 2012b); and the last argument, examined here, (6) growth allometry (Purnell, 1993, 1994).

Trophic level of conodonts

Patterns interpreted to be produced by shearing on the surface and repaired during the animal's growth suggest macrophagy (Purnell, 1995; Donoghue & Purnell, 1999b; Shirley et al., 2018). An active, predatory lifestyle is supported by the discovery of a conodont specimen with preserved extrinsic eye musculature (Gabbott, Aldridge & Theron, 1995), which was interpreted as indicative of conodonts having pattern vision (Purnell, 1994). Calcium isotope analyses indicated that Triassic conodonts were first-level zooplanktivore and piscivore consumers (Balter et al., 2019), suggesting that in this Period, conodonts did not lead a purely predatory lifestyle, but rather were first level consumers. The study suggested that Triassic conodonts may have been scavengers of small fish.

Changes in the function of conodont elements during ontogeny

Conodont elements were retained throughout the life of an individual (Donoghue & Purnell, 1999b), recording periodic growth through apposition of crown tissue lamellae (Zhang, Aldridge & Donoghue, 1997; Dzik, 2008). Shirley et al. (2018) suggested that tooth-like function in conodont elements may have only developed after a larval stage, during which they exhibited a different feeding habit, as mechanical wear of conodont elements is only present after a certain stage of ontogenetic development. After this stage, elements appear to have had prolonged intervals of dental use and short intervals of growth, during which conodonts did not feed and their elements were covered in soft tissue, depositing new layers (Bengtson, 1976; Donoghue & Purnell, 1999b; Shirley et al., 2018).

Open questions

Conodonts changed their apparatus structure and disparity across their stratigraphic range (Dzik, 1991, 2015), possibly reflecting their evolving niches as marine ecosystems increased in complexity (Klug et al., 2010; Ginot & Goudemand, 2019). Their morphology also diversified (Martínez-Pérez et al., 2014a, 2015; Ginot & Goudemand, 2020). This would suggest evolving trophic niches under the assumption that conodont element morphology is an adaptation to their diet (Jones et al., 2012a; Guenser et al., 2019; Ginot & Goudemand, 2019; Petryshen et al., 2020). There is, however, evidence from Ca isotope analysis, suggesting the existence of competition between some taxa, which indicates that trophic niches overlapped between them (Balter et al., 2019). Since no direct evidence of what individual taxa ate could be obtained so far, trophic diversity of conodonts must be inferred from proxies, e.g. by evaluating morphological and functional diversity of food-processing elements. Here we use growth allometry, first applied to conodonts by Purnell (1993, 1994), as a proxy for the dental function.

Allometry

Allometry describes proportional relationships in bodies, usually the size of an organ to the total size of the organism. Its use has become popular during the paleontological revolution (Gould, 1966; Alberch et al., 1979) and has since become a popular tool in different fields of biology and

palaeontology. Allometric relationships can be explored at different scales. Static or size allometry describes variation among individuals of one population and age group. Ontogenetic or growth allometry describes variation in one taxon, as individuals of the taxon grow. Evolutionary allometry describes the variation between taxa, resulting from evolutionary development (Klingenberg, 1996).

Proportional growth, meaning, for example, growth of an organ and growth of the size of the animal at the same rate is called isometry. Positive allometry then describes the organ growing at a higher rate than the rest of the animal. An example of this are the long bones of limbs in humans. Negative allometry, conversely, describes the organ growing at a lower rate than the rest of the body. An example for this is the development of head height (Gould, 1966; Alberch et al., 1979; Klingenberg, 1996).

For conodonts, this means that if conodont elements had a tooth-like function, and specifically, if P₁ elements were used like molars, positive ontogenetic allometric growth of the elements platform to the animal's size can be expected and has been described in *Idiognathodus* sp. and *Gnathodus bilineatus* (Purnell, 1993, 1994). This is because tooth function is linked to surface area and because food requirements of an animal increase at a higher rate than an isometrically growing tooth (Gould, 1966).

Dirichlet Normal Energy

Under the assumption that P₁ elements had a molar-like function, elements can be further analysed with methods designed for teeth to infer the diet of the organism. One such method is The Dirichlet Normal Energy (DNE), a dental topographic analysis that is a measure for the curvature and morphological irregularity of a surface. Essentially, DNE is a measure of how much a given surface differs from a plane (Bunn et al., 2011). Surface topography is an important feature of teeth because they help in breaking down food and can be used to infer diet (Bunn et al., 2011).

In contrast to allometry, DNE is a relatively new tool and has, as of now, mostly been used to analyse the teeth of mammals, and primates in particular (Bunn et al., 2011; Winchester et al., 2014).

Aim of the study

In this work, we address the question, originally proposed by Purnell (1993), that conodont P₁ elements performed molar-like function and that this is reflected in the growth allometry of their surface *versus* length. The null hypothesis is that this relationship is isometric, what would be expected in filter feeders. In contrast to Purnell's (1993) study, which used projections of the surface on a plane, we use 3D meshes to calculate a more accurate platform area than is possible with a two dimensional approximation.

We further test the hypothesis that conodonts occupying the same environments had the same trophic position. To reject this hypothesis, slopes of the growth allometry of P₁ element surface *versus* length should differ.

Under the assumption that that P_1 platforms performed a molar-like function, we further hypothesise that (1) the diets of studied species did not change during ontogenetic growth of one species and that (2) their diets did not differ between species that lived in similar environments. This is tested using Dirichlet Normal Energy (DNE) of the platforms of the elements, as platforms are parts of elements that vary most prominently between elements and thus are inferred to reflect differences in their trophic positions (Jones et al., 2012b; Martínez-Pérez et al., 2016).

Under the null hypothesis (1), DNE values are expected to increase linearly with conodont size. Deviations from a linear relationship would suggest that diets changed across growth stages, allowing us to reject the hypothesis.

Under the null hypothesis (2), DNE values of the two species are expected to be the same. Indistinguishable (overlapping) distributions of DNE values of one growth stage between the two species would falsify the hypothesis, indicating no difference in diet.

We apply these tests to two Upper Triassic conodont species, *Metapolygnathus communisti* Hayashi, 1968 (Hayashi, 1968) and *Epigondolella rigoi* (Noyan & Kozur, 2007) which often co-occur in the same assemblages.

Materials & Methods

Geological location

We used 3D surfaces of P_1 elements of the ozarkodinid conodont species *Metapolygnathus communisti* and *Epigondolella rigoi* from Pizzo Mondello in western Sicily, Italy, where they were collected from a section of 430 m thick marine limestone that is dated to the upper Carnian to upper Norian (Mazza, Rigo & Gullo, 2012). These specimens have an average CAI (colour alteration index) of 1-1.5 which suggests minimal post depositional heating (Epstein, Epstein & Harris, 1977; Nicora et al., 2007; Mazza, Rigo & Gullo, 2012).

Repository

The studied elements are housed in the collection of the Dipartimento di Scienze della Terra “A. Desio” of the Università degli Studi di Milano. Conodont collection of Pizzo Mondello section is housed in Milan and in Padova (Department of Geosciences, University of Padova).

Micro CT-scanning

The 3D surfaces were produced through micro CT-scanning. The elements were scanned with a resolution of 1 μm with a microtomograph nanotomS (General Electric) of the AniRA-ImmOs platform, SFR Biosciences (UMS 3444), Ecole Normale Supérieure de Lyon, France. Amira© software was used for the 3D reconstruction (Guenser et al., 2019).

Identification of growth stages

The most complete specimens have been selected for 3D scanning (Guenser et al., 2019; Figure 1-Figure 2). P_1 elements were separated into six growth stages (GS) based on the maturity of morphological characters of the platform (Mazza & Martínez-Pérez, 2015). The six growth

197 stages are GS1 - early juvenile, GS2 – juvenile, GS3 – late juvenile, GS4 – early adult, GS5 –
198 adult and GS6 – late adult (Mazza & Martínez-Pérez, 2015).

199
200 Figure 1: Growth stages of *Metapolygnathus communisti*

Figure 2: Growth stages of *Epigondolella rigoi*

Growth stage 1 (early juvenile) is characterised by only a simple blade with three to five denticles and a cusp and a rounded or elongated basal cavity being present. By growth stage 2 (juvenile), elements have developed four to seven blade denticles, a very reduced lateral platform margin and a deep pit. Their size does not vary much from GS1. Growth stage 3 (late juvenile) shows six to nine blade denticles that begin to merge in the middle, the formation of a reduced platform with nodes or denticles on the platform margins not present or at least not fully developed. The pit gets shallower while the posterior kneel starts to develop. Growth stage 4 (early adult) is characterised by nine to ten almost fused blade denticles, a developed platform with ornamentation (nodes or denticles on the platform margin) still not fully developed, a narrow and shallow pit and the starting bifurcation of the kneel, if a bifurcated kneel is present in that species. By growth stage 5 (adult), all important characters of a species are present. Elements show ten to thirteen blade denticles that are now completely fused and platform ornamentation is fully developed. Growth stage 6 (late adult) shows an increase in growth compared to GS5 (10-20%), and an overdevelopment of characteristics, such as thickening of platform margins, further fusing of blade and platform denticles and the closing of the pit (Mazza & Martínez-Pérez, 2015).

Twenty-seven P₁ elements of *Metapolygnathus communisti* were used. At Pizzo Mondello, this species occurs from the upper Carnian to the lower Norian (Mazza, Nicora & Rigo, 2018). The specimens range from late juvenile to late adult (GS3-GS6), though mature elements are more abundant (Table 1).

Twenty-three P₁ elements of *Epigondolella rigoi* were used. The stratigraphic range of this species at Pizzo Mondello is from the lower Norian to the middle Norian (Mazza et al., 2010), a significantly longer interval than *Metapolygnathus communisti*. Elements range from GS2 to GS5, again, earlier ontogenetic stages are sparse (Table 1).

Table 1: Numbers of conodont P₁ element specimens by growth stages used for the study.

	<i>Metapolygnathus communisti</i>	<i>Epigondolella rigoi</i>
GS2	0	1
GS3	3	1
GS4	2	6
GS5	7	15
GS6	15	0

Methods

Growth allometry

The length of the element was used as a proxy for the size of the conodont animal, as has been done in previous studies (Purnell, 1993, 1994; Zhang et al., 2018; Ginot & Goudemand, 2019). The length of the element, as well as the length and the area of the platform, were measured using the 3D software MeshLab (Cignoni et al., 2008). The length of the element was measured from the anteriormost point of the blade in a straight line to the middle of the edge of the element's platform.

As the platform is not equally long on the two sides of the blade, its length was measured in different ways. First, as most elements are curved, the convex side of the platform was measured. In *Metapolygnathus communisti* this tended to be, though not always, the longer side of the platform. In *Epigondolella rigoi* this was almost exclusively also the longer side. As a second way of measuring, the longer side of the platform was measured, regardless of curvature (Figure 3).

Figure 3: Measurements of P_1 elements. In most cases one the convex side is also the long side.

In *M. communisti*, in both instances, the platform was measured from the most anterior part of the platform to its posterior end in a line parallel to the imagined symmetrical axis of the platform (Figure 3).

In *E. rigoi*, the platform was measured from the geniculation point to the platform's posterior end, as the anterior through margin in this species, though reduced (Mazza, Cau & Rigo, 2012) reaches quite far up the blade, especially in more mature growth stages (Figure 4).

Figure 4: Morphological and structural terms used in the description of P₁ elements from Mazza, Cau & Rigo (2012)

The measured area of the platform includes the platform itself and additionally the cusp and any carinal nodes that follow the cusp on its posterior side (Figure 3).

In *M. communisti*, specimens of earlier growth stages tend to exhibit only one posterior carinal node, which is already present in GS1. From GS3 on, a second posterior node may appear (Mazza & Martínez-Pérez, 2015). Adult specimens show 3-4 posterior carinal nodes (Mazza, Rigo & Gullo, 2012). In our measurements, however, we found that we consistently identified only 2-3 posterior carinal nodes in adult specimens. In *E. rigoi* the cusp is always the second to last denticle on the carina, followed on its posterior side by a single larger node (Mazza, Rigo & Gullo, 2012). These parts of P₁ elements were added to the measurements of the platform area, even though they are taxonomically not part of the platform, because they likely played a similar part in the processing of food as the platform itself. In *E. rigoi*, the anterior through margin was not included in the measurements of the platform area. The anterior trough margin is not present in *M. communisti* (Mazza, Rigo & Gullo, 2012). Mean, median and variance of lengths and areas of the P₁ elements were calculated for each growth stage with R Software (R Core Team, 2021). Reduced major axis regression (RMA) was calculated using the R package “smatr” (Warton et al., 2012) to examine the relationship between the length of the platform and the length of the element, as well as the logarithm of the platform area and the length of the elements. The

platform area was log-transformed to account for its increase as a square of the length of the element. RMA was chosen as a method because both variables are mutually dependent.

DNE

Dirichlet Normal Energy (DNE) measures a surface's curvature and complexity. The DNE of an object is independent of its size and orientation. The DNE equation is commonly written as follows (Bunn et al., 2011):

$$DNE = \sum e(p) \times area(p)$$

Here, $e(p)$ is the Dirichlet Energy Density at a point p . The sum of the areas of all points p (however small) on a surface is equal to the total area of the surface. Flat planes have DNE values of 0. A higher DNE value therefore expresses the elements complexity and the “sharpness” of a surface (Bunn et al., 2011).

The R package “molaR” (Pampush et al., 2016), used here, requires preparation of the meshes prior to DNE calculation. For this, the platform area was isolated in MeshLab, in the same way that it was measured.

To ensure consistency of DNE calculation, all meshes were simplified to 10 000 faces (Spradley et al., 2017) using Quadric Edge Collapse Decimation in MeshLab. They were then rescaled so that each platform area equaled 0.1 mm² to ensure comparable resolution of detail in each element's platform. If not resized, a small P₁ element would have more faces per mm² and, therefore, a higher resolution than a large element with the same number of faces. Meshes were then smoothed in Avizo using the Avizo smoothing function with lambda = 0.6 and 25 iterations.

Twenty to thirty smoothing iterations, a conservative amount when compared to other approaches (Bunn et al., 2011; Winchester et al., 2014; Spradley et al., 2017), are recommended because they eliminate scanning noise while capturing fine-scale features and avoiding the creation of artificial dimples that can be caused by oversmoothing (Spradley et al., 2017).

Different numbers of iterations (5, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, and 100) were tested on a singular specimen of *M. communisti* to determine the impact of the number of smoothing iterations (Figure 5).

Figure 5: The impact of different numbers of smoothing iterations on DNE values measured for the platform of the P₁ element NA37_A_02_GS5 (*M. communisti*). The red dot indicates 25 smoothing iterations.

The meshes were then resaved in MeshLab to avoid errors when calculating the DNE in R Software. They were imported into R Software as binary *ply* files, where they were cleaned and where the total surface DNE was calculated using the R package “molaR” (Pampush et al., 2016) with an included boundary exclusion criterion (BoundaryDiscard=”vertex”), as advised by Spradley et al. (2017). The total surface DNE is the mean of all DNE values for individual faces of a surface (Pampush et al., 2016). Mean, median and variance of the DNE were calculated for each growth stage and species with R Software (R Core Team, 2021).

Reduced major axis regression (RMA) was again used to examine the relationship between DNE and the length of the platform, as well as the log-transformed platform area in both species. Growth stage 5 was chosen for further analysis, because sufficient numbers of specimens of this growth stage were available (Table 1). Growth stage 5, representing adults, also allows for interpretations of the diet.

Results

Growth allometry

Metapolygnathus communisti

Regression analysis of P₁ elements of *M. communisti* shows that both sets of variables (the length of the platform and the length of the element, as well as the log-transformed platform area and the length of the elements) are correlated (Figure 6A-C). The regression slopes of the length of the element over that of the platform, as well as the regression slope of the length of the element over the log-transformed area of the platform, differed significantly from one, i.e. from isometric growth ($p < 0.000001$; Figure 6B, C). Although the long side of the element platform is also the concave one in most cases, there are differences between the two ways the platform area was measured. This is reflected in the slope coefficients deviating slightly from each other (Figure 6A).

Figure 6. Allometric growth of *M. communisti* (n=27) and *E. rigoi* (n=23); each dot represent an element and the colours are related to the growth stage. A. Platform length over element length for *M. communisti*; comparison of long and convex platform length. B. Platform length over element length for *M. communisti*; convex platform length only. C. Platform area over element length for *M. communisti*. D. Platform length over element length for *E. rigoi*; comparison of long and convex platform length. E. Platform length over element length for *E. rigoi*; convex platform length only. F. Platform area over element length for *E. rigoi*.

Slope coefficients for the platform length over element length were 0.95 and 1.06 (length of platform measured on the convex and on the long side, respectively). Both values are similar to the isometric value of one, which indicates proportional growth. The slope coefficient for the log-transformed platform area was 3.86, which is greater than the isometric value of two, indicating positive allometry. The platform area grew at a faster rate than the element length (Figure 6C).

Epigondolella rigoi

In this species, both sets of variables (the length of the platform and the length of the element, as well as the log transformed platform area and the length of the elements) are also correlated (Figure 6D-F). The regression slopes of the length of the element over that of the platform, as well as the regression slope of the length of the element over the log-transformed area of the platform, differed significantly from zero ($p < 0.000001$). Differences between the two ways the platform length was measured were barely visible in this species, because the long and the convex side of the platform describe the same side in almost all cases (Figure 6D).

Slope coefficients for the platform length were 0.75 and 0.73 (length of platform measured on the convex and on the long side, respectively). These values are smaller than the isometric value of one, which indicates negative allometry. Here, the platform length grew at a lower rate than the element length. Slope coefficients for the platform area was 4.16, which is greater than the isometric value of two, indicating positive allometry (Figure 6F).

Comparison of growth allometry in different species

The hypothesis of molar-like function of P₁ elements could not be rejected based on positive growth allometry of the platform area in both species. In *M. communisti*, *Idiognathodus* sp. and *Gnathodus bilineatus*, slope coefficients of the platform length close or slightly greater to the isometric value one (*Idiognathodus* sp.: 1.089; Purnell, 1993, 1994; *Gnathodus bilineatus*: 1.167; Purnell, 1994) indicate near isometric growth or slight positive allometric growth in the species. Platform length and element length grew at similar rates (Figure 7A).

Figure 7. Comparison of the allometric growth *Gnathodus bilineatus* (Purnell, 1994), *Idiognathodus* sp. (Purnell, 1993, 1994), *Epigondolella rigoi* and *Metapolygnathus communisti*. A. Platform length over element length. Platform length in *M. communisti* and *E. rigoi* was measured on the convex side. B. Platform area over element length.

In *E. rigoi*, the negative allometry of the platform length did not follow the same growth pattern as the other species. Its platform elongated at a slightly slower rate than the length of the element (Figure 7A).

Platform areas in all four species showed positive allometry over element length, but at different rates. *Idiognathodus* sp. and *Gnathodus bilineatus* had the lowest slope coefficients of 2.149 and 2.164, respectively (Purnell, 1993). The slope coefficients in *M. communisti* and *E. rigoi* were

considerably higher (Figure 7B). This may be explained by the fact that the platform area of *Idiognathodus* sp. was calculated as a two-dimensional surface, based on the platform's outline, and the platform area of *Gnathodus bilineatus* was approximated with the calculation of an ellipse with the platform's dimensions (Purnell, 1994). In contrast, platform areas of *M. communisti* and *E. rigoi* were in this study calculated based on three dimensional meshes.

DNE

DNE through ontogeny

Distributions of DNE values over element platforms are shown in Figure 8 and Figure 9. In both species, ontogenetically younger growth stages were represented by fewer specimens (Table 1), which made observations of these stages less conclusive. Nevertheless, in both *M. communisti* and *E. rigoi*, DNE values increased consistently with the element length (Figure 10A), as well as with the platform area (Figure 10B). No shifts in the growth trajectory of DNE were observed.

Figure 8: Log-transformed DNE values across growth stages GS3 to GS6 of *Metapolygnathus communisti*.

Figure 9: Log-transformed DNE values across growth stages GS2 to GS5 of *Epigondolella rigoi*.

The hypothesis that the dietary niches of both species, assessed using DNE as a proxy, remained the same throughout ontogeny could not, therefore, be rejected.

Metapolygnathus communisti

Regression analysis of P₁ elements of *M. communisti* showed that element length, as well as platform area, are correlated with DNE ($R^2 = 0.52$ and 0.61 , respectively). Regression slopes of the DNE values over element length and the log-transformed platform area differed significantly from one ($p < 0.0001$). The slope coefficients are 290.6725 and 75.41139 for element length and platform area, respectively (Figure 10A-B).

Epigondolella rigoi

Regression analysis of P₁ elements of *E. rigoi* showed that element length, as well as platform area, are correlated with DNE ($R^2 = 0.32$ and 0.43 respectively). Regression slopes of DNE over element length and the log-transformed platform differed significantly from one ($p < 0.001$). Slope coefficients were 395.6475 and 92.75222 for element length and platform area, respectively (Figure 10A-B).

Comparison of DNE values in adult elements

For the comparison of DNE values between P₁ element platforms of *M. communisti* and *E. rigoi*, growth stage 5 is of particular interest, because both species have enough specimens of this stage to allow further analysis (Table 1). Additionally, growth stage 5 contains specimens of adult

conodonts which makes their DNE values suitable for comparisons outside of conodonts. It needs to be remarked that the classification of growth stage 5 as “adult” does not consider biology, but the morphological maturity of the elements (Mazza & Martínez-Pérez, 2015). As with the other growth stages, specimens of growth stage 5 of *E. rigoi* showed higher DNE values than specimens of growth stage 5 of *M. communisti*, though there was some overlap (Figure 10C). The difference in DNE values between the two species appears big enough to reject the hypothesis that both species shared the same diet.

Figure 10: Regression of total surface DNE values over platform length (A), log-transformed platform area (B) in P₁ elements of *Epigondolella rigoi* and *Metapolygnathus communisti* and distribution of DNE values across growth stages in both species (C).

Discussion

Growth Allometry

We were able to reject the hypothesis that platform surface area increased isometrically with respect to element length in P₁ elements of *Metapolygnathus communisti* and *Epigondolella rigoi* in favour of the alternative hypothesis that the growth allometry was positive, as expected in dental organs functioning as molars. This finding supports previous finding of positive growth allometry in these organs in much older, Carboniferous, ozarkodinid taxa *Gnathodus bilineatus* and *Idiogonathodus* sp. (Purnell, 1993, 1994). It is consistent with previous research on microwear (Purnell, 1995; Martínez-Pérez et al., 2014b), enamel-like ultrastructure of conodont lamellar crown tissue (Purnell, Donoghue & Aldridge, 2000), and finite element analysis (Jones et al., 2012a).

The calculation of the platform area in three dimensions, as opposed to two dimensional approximation based on platform outline or platform dimensions (Purnell, 1994), yields more exact measurement of the area. Furthermore, measurements based on pictures suffer from distortions resulting from projecting on a plane, where differences in levelling the photographed specimens might affect the results. The effect of 3D measurements of the platform is a higher slope coefficient, meaning a more strongly positive allometry of the platform (Figure 7B).

The decision what to include in the platform for allometric analysis is worth discussing, because it is somewhat subjective. In this work, we decided to include the cusp and all carinal nodes posterior to it, because, in many cases, the cusp marked a notable transition between sharper denticles of the blade and flatter nodes on the platform. This resulted in different numbers of posterior nodes in *M. communisti*, because, in this species, the number of posterior nodes differs between growth stages (Mazza, Rigo & Gullo, 2012; Mazza & Martínez-Pérez, 2015). This variation has a greater effect in 3D measurements than it would in 2D. In *E. rigoi* the number of posterior nodes stays the same (Mazza, Rigo & Gullo, 2012).

Dirichlet Normal Energy

Ontogenetic development of DNE

In *M. communisti*, as well as in *E. rigoi*, DNE values increased linearly with body size, where element length was used as a proxy for body size (Zhang et al., 2018; Ginot & Goudemand, 2019). This does not contradict the possible presence of a larval stage, during which conodonts had a different method of feeding (Shirley et al., 2018), because all specimens that are considered here were more mature.

Comparison of DNE values in adult elements

Differences in DNE values between *M. communisti* and *E. rigoi* are great enough to merit a careful suggestion of different diets between the two species. More DNE analyses on conodonts are needed to understand the scope of DNE values in conodonts and to confidently suggest that a

certain discrepancy between DNE values of different species reflects different dietary niches. Here, the only reference values were those available for exclusively terrestrial mammals, and mostly for primates.

DNE values of growth stage 5 specimens of *M. communisti* and *E. rigoi* are similar to those reported for insectivores or folivores in the case of *E. rigoi* and for folivores or omnivores in the case of *M. communisti*, had these DNE values been observed in primates (Bunn et al., 2011; Winchester et al., 2014). Within the range of DNE values observed in primates, these values are relatively high. Though these dietary classifications are not applicable to conodonts, they may offer a general reference point for the methods of breaking down different food types. Insectivores rely on sharp cusps to apply maximal force on a small surface, in order to pierce hard insect chitin, folivores also use steeply sloped cusps to shear tough cellulose-rich leaves. (Lucas, 1979; Strait, 1997).

It is possible that conodont element platforms adapted to break down food types with similar properties. It has been suggested that conodonts may have punctured arthropod larvae (Dzik, 2021), which would be consistent with the DNE values observed in *M. communisti*.

Problems in DNE analysis of conodont elements

There are several challenges researchers face when working with DNE in conodonts. As DNE is a comparatively new tool, reference values and understanding of variability (e.g. intraspecific, ontogenetic, resulting from preservation) is limited. So far, DNE research has focused on gnathostomes, whose tooth function is, in large part, reliant on jaws acting as levers. This necessitates that the comparisons that are drawn here between conodont elements and primate molars must be viewed as extremely hypothetical. Marine environment, evolutionary distance and the lack of jaws in conodonts make it likely that tooth function is not completely analogous between the two.

Typical applications of DNE do not take into account ontogenetic development of conodonts, because, contrary to mammalian molars, in conodonts the number of denticles in P₁ element platforms increases during ontogenetic growth. This plays into the recurring problem of what to include in the conodont platform, which also has an impact on DNE. This was especially problematic in *M. communisti*, because here, in addition to denticles on the platform margins increasing in number over ontogenetic growth, the number of carinal nodes posterior to the cusp varies between growth stages.

A consistent protocol of mesh preparation prior to DNE calculation is also needed. Especially the number of smoothing iterations varies in current literature and this can have impact on DNE results (Spradley et al., 2017). This is important, because comparisons are most conclusive when drawn between data with comparable preparation. We second the recommendations by Spradley et al. (2017): a conservative number of smoothing iterations (20-30) using non-Laplace-based smoothing operators, such as that implemented in Avizo, and mesh simplification to a fixed number of faces.

Conclusions

We tested the hypothesis that conodont P₁ elements performed molar-like function by analysing the growth allometry of platform surface area *versus* length, following the protocol by Purnell (1993, 1994), but in 3D meshes rather than on 2D projections of platforms. We further used slopes of this allometric relationships to test the hypothesis that conodonts occupying the same environments shared the same trophic positions. We applied the test to 3D surfaces of P₁ elements of the ozarkodinid conodont species *Metapolygnathus communisti* and *Epigondolella rigoi* from the Upper Triassic section Pizzo Mondello in western Sicily, Italy. Platform length grew isometrically with respect to element length, whereas log-transformed platform area showed positive allometry with respect to element length, with slopes equal 3.86 in *M. communisti* and 4.16 in *E. rigoi*. These values are substantially higher than those reported for Carboniferous conodonts *Idiognathodus* sp. and *Gnathodus bilineatus* (Purnell, 1993), but this may reflect a more precise measurement of surface area in 3D models compared to 2D projections used in previous studies. Thus, it is not possible to compare these values directly, but we conclude that stronger positive allometry of platform area in *E. rigoi* indicates higher metabolic needs and might be a proxy for a higher trophic level compared to *M. communisti*. We also applied dental topographic analysis using Dirichlet Normal Energy (DNE) to the platforms in order to test the hypotheses that the diets of *M. communisti* and *E. rigoi* did not change during ontogenetic growth and that they occupied the same dietary niches. Surface DNE values increased linearly in function of element length and log-transformed platform area, indicating no ontogenetic changes. Specimens of the adult growth stage of *E. rigoi* showed higher DNE values than specimens of the same growth stage in *M. communisti*, consistent with stronger positive allometry of platform surface and with a higher demand for energy in this species. Based on DNE values available for primates, those of the adult growth stages were similar to those reported for insectivores or folivores in the case of *E. rigoi* and for folivores or omnivores in the case of *M. communisti*. Previous studies applying morphological and ultrastructural proxies for the dietary position of conodonts addressed mostly stratigraphically older conodont taxa, but our results indicate that Late Triassic species occupied the predator/scavenger niche in spite of the highly developed diversity of gnathostomes in this niche. We also show that within this broad niche, co-occurring taxa differed in their diets, which supports trophic diversification as an important driver of the remarkable disparity of their elements.

Acknowledgements

We thank Bryan Shirley for support in using Aviso and Wyatt Petryshen for advice on cleaning and saving meshes. We also thank Nicolas Goudemand for supporting the project. PG was supported by Visiting Scholarship awarded by Friedrich-Alexander-Universität Erlangen-Nürnberg. EJ was supported by Deutsche Forschungsgemeinschaft (project no JA 2718/3-1). MR was supported by DOR2054230/20 by University of Padova.

References

546 Alberch P, Gould SJ, Oster GF, Wake DB. 1979. Size and shape in ontogeny and phylogeny.
547 *Paleobiology* 5:296–317.

548 Balter V, Martin JE, Tacail T, Suan G, Renaud S, Girard C. 2019. Calcium stable isotopes place
549 Devonian conodonts as first level consumers. *Geochemical Perspectives Letters* 10:36–39.

550 Bengtson S. 1976. The structure of some Middle Cambrian conodonts, and the early evolution of
551 conodont structure and function. *Lethaia* 9:185–206. DOI: 10.1111/j.1502-3931.1976.tb00966.x.

552 Bunn JM, Boyer DM, Lipman Y, St. Clair EM, Jernvall J, Daubechies I. 2011. Comparing Dirichlet
553 normal surface energy of tooth crowns, a new technique of molar shape quantification for dietary
554 inference, with previous methods in isolation and in combination. *American Journal of Physical*
555 *Anthropology* 145:247–261. DOI: 10.1002/ajpa.21489.

556 Cignoni P, Callieri M, Corsini M, Dellepiane M, Ganovelli F, Ranzuglia G. 2008. Meshlab: an open-
557 source mesh processing tool. In: *Eurographics Italian Chapter Conference*. 129–136.

558 Donoghue PCJ. 2001a. Conodonts meet cladistics: recovering relationships and assessing the
559 completeness of the conodont fossil record. *Palaeontology* 44:65–93. DOI: 10.1111/1475-
560 4983.00170.

561 Donoghue PCJ. 2001b. Microstructural variation in conodont enamel is a functional adaptation.
562 *Proceedings of the Royal Society of London. Series B: Biological Sciences* 268:1691–1698. DOI:
563 10.1098/rspb.2001.1728.

564 Donoghue PCJ, Purnell MA. 1999a. Mammal-like occlusion in conodonts. *Paleobiology* 25:58–74.

565 Donoghue PCJ, Purnell MA. 1999b. Growth, function, and the conodont fossil record. *Geology* 27:251–
566 254. DOI: 10.1130/0091-7613(1999)027<0251:GFATCF>2.3.CO;2.

567 Donoghue PCJ, Purnell MA, Aldridge RJ. 1998. Conodont anatomy, chordate phylogeny and vertebrate
568 classification. *Lethaia* 31:211–219. DOI: 10.1111/j.1502-3931.1998.tb00509.x.

569 Du Y, Chiari M, Karádi V, Nicora A, Onoue T, Pálffy J, Roghi G, Tomimatsu Y, Rigo M. 2020. The
570 asynchronous disappearance of conodonts: New constraints from Triassic-Jurassic boundary

571 sections in the Tethys and Panthalassa. *Earth-Science Reviews* 203:103176. DOI:
572 10.1016/j.earscirev.2020.103176.

573 Dzik J. 1991. Evolution of oral apparatuses in the conodont chordates. *Acta Palaeontologica Polonica*
574 36:265–323.

575 Dzik J. 2008. Evolution of morphogenesis in 360-million-year-old conodont chordates calibrated in days.
576 *Evolution & Development* 10:769–777. DOI: 10.1111/j.1525-142X.2008.00291.x.

577 Dzik J. 2015. Evolutionary roots of the conodonts with increased number of elements in the apparatus.
578 *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* 106:29–53.

579 Dzik J. 2021. Protaspis larva of an aglaspigid-like arthropod from the Ordovician of Siberia and its
580 habitat. *Arthropod Structure & Development* 61:101026. DOI: 10.1016/j.asd.2020.101026.

581 Epstein AG, Epstein JB, Harris LD. 1977. Conodont color alteration: an index to organic metamorphism.
582 *United States Geological Survey Professional Paper* 995:1–27.

583 Foote M, Sepkoski JJ. 1999. Absolute measures of the completeness of the fossil record. *Nature* 398:415–
584 417. DOI: 10.1038/18872.

585 Gabbott SE, Aldridge RJ, Theron JN. 1995. A giant conodont with preserved muscle tissue from the
586 Upper Ordovician of South Africa. *Nature* 374:800–803. DOI: 10.1038/374800a0.

587 Ginot S, Goudemand N. 2019. Conodont size, trophic level, and the evolution of platform elements.
588 *Paleobiology* 45:458–468. DOI: 10.1017/pab.2019.19.

589 Ginot S, Goudemand N. 2020. Global climate changes account for the main trends of conodont diversity
590 but not for their final demise. *Global and Planetary Change* 195:103325. DOI:
591 10.1016/j.gloplacha.2020.103325.

592 Gould SJ. 1966. Allometry and size in ontogeny and phylogeny. *Biological Reviews* 41:587–638. DOI:
593 10.1111/j.1469-185X.1966.tb01624.x.

594 Guenser P, Souquet L, Dolédec S, Mazza M, Rigo M, Goudemand N. 2019. Deciphering the roles of
595 environment and development in the evolution of a Late Triassic assemblage of conodont
596 elements. *Paleobiology* 45:440–457. DOI: 10.1017/pab.2019.14.

597 Hayashi S. 1968. The Permian Conodonts in Chert of the Adoyama Formation, Ashio Mountains, Central
598 Japan. *Earth Science* 2:63–77.

599 Jones D, Evans AR, Rayfield EJ, Siu KKW, Donoghue PCJ. 2012a. Testing microstructural adaptation in
600 the earliest dental tools. *Biology Letters* 8:952–955. DOI: 10.1098/rsbl.2012.0487.

601 Jones D, Evans AR, Siu KKW, Rayfield EJ, Donoghue PCJ. 2012b. The sharpest tools in the box?
602 Quantitative analysis of conodont element functional morphology. *Proceedings of the Royal*
603 *Society B: Biological Sciences* 279:2849–2854. DOI: 10.1098/rspb.2012.0147.

604 Klingenberg CP. 1996. Multivariate Allometry. In: Marcus LF, Corti M, Loy A, Naylor GJP, Slice DE
605 eds. *Advances in Morphometrics*. Boston, MA: Springer US, 23–49. DOI: 10.1007/978-1-4757-
606 9083-2_3.

607 Klug C, Kröger B, Kiessling W, Mullins GL, Servais T, Fryda J, Korn D, Turner S. 2010. The Devonian
608 nekton revolution. *Lethaia* 43:465–477. DOI: 10.1111/j.1502-3931.2009.00206.x.

609 Lucas PW. 1979. The Dental-Dietary Adaptations of Mammals. *Neues Jahrbuch für Geologie und*
610 *Paläontologie Monatshefte* 8:486–512.

611 Martínez-Pérez C, Cascales-Miñana B, Plasencia P, Botella H. 2015. Exploring the major depletions of
612 conodont diversity during the Triassic. *Historical Biology* 27:503–507. DOI:
613 10.1080/08912963.2014.890192.

614 Martínez-Pérez C, Plasencia P, Cascales-Miñana B, Mazza M, Botella H. 2014a. New insights into the
615 diversity dynamics of Triassic conodonts. *Historical Biology* 26:591–602. DOI:
616 10.1080/08912963.2013.808632.

617 Martínez-Pérez C, Plasencia P, Jones D, Kolar-Jurkovšek T, Sha J, Botella H, Donoghue PCJ. 2014b.
618 There is no general model for occlusal kinematics in conodonts. *Lethaia* 47:547–555. DOI:
619 10.1111/let.12080.

620 Martínez-Pérez C, Rayfield EJ, Botella H, Donoghue PCJ. 2016. Translating taxonomy into the evolution
621 of conodont feeding ecology. *Geology* 44:247–250. DOI: 10.1130/G37547.1.

- Martínez-Pérez C, Rayfield EJ, Purnell MA, Donoghue PCJ. 2014c. Finite element, occlusal, microwear and microstructural analyses indicate that conodont microstructure is adapted to dental function. *Palaeontology* 57:1059–1066. DOI: 10.1111/pala.12102.
- Mazza M, Cau A, Rigo M. 2012. Application of numerical cladistic analyses to the Carnian–Norian conodonts: a new approach for phylogenetic interpretations. *Journal of Systematic Palaeontology* 10:401–422.
- Mazza M, Furin S, Spötl C, Rigo M. 2010. Generic turnovers of Carnian/Norian conodonts: Climatic control or competition? *Palaeogeography, Palaeoclimatology, Palaeoecology* 290:120–137.
- Mazza M, Martínez-Pérez C. 2015. Unravelling conodont (Conodonta) ontogenetic processes in the Late Triassic through growth series reconstructions and X-ray microtomography. *Bollettino della Società Paleontologica Italiana* 54:161–186.
- Mazza M, Nicora A, Rigo M. 2018. *Metapolygnathus parvus* Kozur, 1972 (Conodonta): a potential primary marker for the Norian GSSP (Upper Triassic). *Bollettino della Società Paleontologica Italiana*. DOI: 10.4435/BSPI.2018.06.
- Mazza M, Rigo M, Gullo M. 2012. Taxonomy and biostratigraphic record of the Upper Triassic conodonts of the Pizzo Mondello section (western Sicily, Italy), GSSP candidate for the base of the Norian. *Rivista Italiana di Paleontologia e Stratigrafia* 118:85–130.
- Nicoll RS. 1987. Form and function of the Pa element in the conodont animal. In: Aldridge RJ ed. *Palaeobiology of conodonts*. Chichester: Ellis Horwood, 77–90.
- Nicora A, Balini M, Bellanca A, Bertinelli A, Bowring SA, Di Stefano P, Dumitrica P, Guaiumi C, Gullo M, Hungerbuehler A, others. 2007. The Carnian/Norian boundary interval at Pizzo Mondello (Sicani Mountains, Sicily) and its bearing for the definition of the GSSP of the Norian Stage. *Albertiana* 36:102–129.
- Noyan ÖF, Kozur HW. 2007. Revision of the late Carnian–early Norian conodonts from the Stefanion section (Argolis, Greece) and their palaeobiogeographic implications. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 245:159–178.

648 Pampush JD, Winchester JM, Morse PE, Vining AQ, Boyer DM, Kay RF. 2016. Introducing molaR: a
649 New R Package for Quantitative Topographic Analysis of Teeth (and Other Topographic
650 Surfaces). *Journal of Mammalian Evolution* 23:397–412. DOI: 10.1007/s10914-016-9326-0.

651 Petryshen W, Henderson CM, De Baets K, Jarochovska E. 2020. Evidence of parallel evolution in the
652 dental elements of Sweetognathus conodonts. *Proceedings of the Royal Society B: Biological
653 Sciences* 287:20201922. DOI: 10.1098/rspb.2020.1922.

654 Purnell MA. 1993. Feeding mechanisms in conodonts and the function of the earliest vertebrate hard
655 tissues. *Geology* 21:375–377. DOI: 10.1130/0091-7613(1993)021<0375:FMICAT>2.3.CO;2.

656 Purnell MA. 1994. Skeletal ontogeny and feeding mechanisms in conodonts. *Lethaia* 27:129–138. DOI:
657 10.1111/j.1502-3931.1994.tb01567.x.

658 Purnell MA. 1995. Microwear on conodont elements and macrophagy in the first vertebrates. *Nature*
659 374:798–800. DOI: 10.1038/374798a0.

660 Purnell MA, von Bitter PH. 1992. Blade-shaped conodont elements functioned as cutting teeth. *Nature*
661 359:629–631. DOI: 10.1038/359629a0.

662 Purnell MA, Donoghue PCJ, Aldridge RJ. 2000. Orientation and anatomical notation in conodonts.
663 *Journal of Paleontology* 74:113–122. DOI: 10.1017/S0022336000031292.

664 R Core Team. 2021. R: A Language and Environment for Statistical Computing. *R Foundation for
665 Statistical Computing, Vienna, Austria*.

666 Schmidt H. 1934. Conodonten-Funde in ursprünglichem Zusammenhang. *Palaeontologische Zeitschrift*
667 16:76–85. DOI: 10.1007/BF03041668.

668 Scott HW. 1934. The zoological relationships of the conodonts. *Journal of Paleontology* 8:448–455.

669 Shirley B, Grohgan M, Bestmann M, Jarochovska E. 2018. Wear, tear and systematic repair: testing
670 models of growth dynamics in conodonts with high-resolution imaging. *Proceedings of the Royal
671 Society B: Biological Sciences* 285:20181614. DOI: 10.1098/rspb.2018.1614.

- 672 Spradley JP, Pampush JD, Morse PE, Kay RF. 2017. Smooth operator: The effects of different 3D mesh
673 retriangulation protocols on the computation of Dirichlet normal energy. *American Journal of*
674 *Physical Anthropology* 163:94–109. DOI: 10.1002/ajpa.23188.
- 675 Strait SG. 1997. Tooth use and the physical properties of food. *Evolutionary Anthropology: Issues, News,*
676 *and Reviews* 5:199–211. DOI: 10.1002/(SICI)1520-6505(1997)5:6<199::AID-
677 EVAN2>3.0.CO;2-8.
- 678 Warton DI, Duursma RA, Falster DS, Taskinen S. 2012. smatr 3– an R package for estimation and
679 inference about allometric lines. *Methods in Ecology and Evolution* 3:257–259. DOI:
680 10.1111/j.2041-210X.2011.00153.x.
- 681 Winchester JM, Boyer DM, St. Clair EM, Gosselin-Ildari AD, Cooke SB, Ledogar JA. 2014. Dental
682 topography of platyrrhines and prosimians: Convergence and contrasts. *American Journal of*
683 *Physical Anthropology* 153:29–44. DOI: 10.1002/ajpa.22398.
- 684 Zhang S, Aldridge RJ, Donoghue PCJ. 1997. An Early Triassic conodont with periodic growth? *J.*
685 *Micropalaeontol.* 16:65–72. DOI: 10.1144/jm.16.1.65.
- 686 Zhang ZT, Sun YD, Wignall PB, Fu JL, Li HX, Wang MY, Lai XL. 2018. Conodont size reduction and
687 diversity losses during the Carnian Humid Episode in SW China. *Journal of the Geological*
688 *Society* 175:1027–1031.