**Dive Performance and Aquatic Thermoregulation of the World's Smallest Mammalian
Diver, the American Water Shrew (*Sorex palustris*)**

Roman W. Gusztak,^{1,2} Robert A. MacArthur,¹ and Kevin L. Campbell^{1,†}

¹Department of Biological Sciences, University of Manitoba, Winnipeg, Manitoba R3T 2N2
Canada

²Current address: Department of Anesthesiology, University of Saskatchewan, Saskatoon,
Saskatchewan S7N 0W8 Canada

[†]Corresponding author; e-mail: Kevin.Campbell@umanitoba.ca

Running Head: Diving physiology of American water shrews

Keywords: aerobic dive limit, body oxygen store, diving behavior, energetics, hemoglobin, lung
volume, muscle buffering capacity, myoglobin, Soricidae

20 **ABSTRACT**

21 Allometry predicts that the 12–17 g American water shrew (*Sorex palustris*)—the world’s
 22 smallest mammalian diver—will have the highest diving metabolic rate coupled with the lowest
 23 total body oxygen storage capacity, skeletal muscle buffering capacity, and glycolytic potential
 24 of any endothermic diver. Consistent with expectations, and potentially owing to their low
 25 thermal inertia, water shrews had a significantly higher diving metabolic rate in 10°C (8.77 mL
 26 O₂ g⁻¹ hr⁻¹) compared to 30°C water (6.57 mL O₂ g⁻¹ hr⁻¹). Unlike larger-bodied divers, muscle
 27 myoglobin contributed minimally (7.7–12.4%) to total onboard O₂ stores of juvenile and adult
 28 water shrews, respectively, but was offset by high blood O₂ carrying capacities (26.4–26.9 vol.
 29 %). Diving was predominantly aerobic, as only 2.3–3.9% of dives in 30 and 10°C water,
 30 respectively, exceeded the calculated aerobic dive limits at these temperatures (10.8–14.4 sec).
 31 The mean voluntary dive time of water shrews during 20-min trials in 3–30°C water was 5.1±0.1
 32 sec ($N=25$, $n=1584$), with a mean maximum dive time of 10.3±0.4 sec. However, the average
 33 dive duration (6.9±0.2 sec, $n=257$) of radio-telemetered shrews exclusively foraging in a
 34 simulated riparian environment (3°C water) for 12- to 28-hr suggest that mean (but not
 35 maximum) dive times of water shrews in the wild may be longer. Mean dive duration, duration
 36 of the longest dive, and total time in water all decreased significantly as water temperature
 37 declined, suggesting that shrews employed behavioral thermoregulation to defend against
 38 immersion hypothermia. Additionally, free-diving shrews in the 24-hr trials consistently elevated
 39 core body temperature by ~1.0–1.5°C immediately prior to initiating aquatic foraging bouts, and
 40 ended these bouts when body temperature was still at or above normal resting levels (~37.8°C).
 41 We suggest this observed pre-dive hyperthermia aids to heighten the impressive somatosensory
 42 physiology, and hence foraging efficiency, of this diminutive predator while submerged.

Introduction

The three primary factors dictating the maximum underwater endurance of air-breathing mammalian divers—anaerobic capacity, total onboard oxygen stores, and the rate at which these reserves are consumed—are all strongly influenced by body size (Emmett and Hochachka, 1981; Schreer and Kovaks, 1997; Halsey et al. 2006). The mass-specific rate of O_2 consumption ($\dot{V}O_2$), for instance, varies inversely with body mass according to a scaling coefficient of -0.25 to -0.33 (Kleiber 1975; White and Seymour 2003). Conversely, the catalytic activity of glycolytic enzymes (i.e., glycogen phosphorylase, pyruvate kinase, and lactate dehydrogenase), and hence the maximum anaerobic potential of muscle, scales positively with size among mammals with an exponent of 1.09 to 1.15 (Emmett and Hochachka, 1981). Finally, body oxygen stores representing the combined oxygen reserves of lungs, blood, and muscle scale approximately isometrically with body mass (Calder 1996). Taken together, the resultant elevation in mass-specific $\dot{V}O_2$ of small amphibious species actively swimming underwater (diving metabolic rate, DMR), coupled with lower absolute oxygen stores, should lead to a disproportionate reduction in their aerobic dive performance relative to larger divers. Maximal submergence times should be further curtailed by a reduced ability to exploit anaerobic pathways for ATP production and buffer aerobic (CO_2) and anaerobic byproducts (lactic acid). As with dive endurance, the body surface area-to-volume ratio varies inversely with body size, resulting in lower mass-specific thermoregulatory costs for larger divers compared to their smaller counterparts (MacArthur 1989). Small-bodied divers also have a limited capacity for enhancing pelage or tissue insulation, while large divers are often endowed with a thick blubber layer or fur that dramatically increases whole-body insulation (Favilla and Costa 2020). All of these factors predispose small, semi-

aquatic species to higher mass-specific rates of heat loss in the aquatic medium than is the case for their larger counterparts.

Given the significant physiological challenges facing small-bodied amphibious divers, it is surprising that relatively few studies have examined the diving capacity and aquatic thermal biology of mammals weighing less than 5 kg (e.g., Dawson and Fanning 1981; MacArthur 1984a; Evans et al. 1994; Hindle et al. 2006; Harrington et al. 2012; Jordaan et al. 2021). Moreover, to our knowledge, comparable data for the smallest mammalian divers (<100 g) are available only for star-nosed moles, *Condylura cristata* (McIntyre et al. 2002), European water shrews, *Neomys fodiens* (Köhler 1991; Vogel 1998), and, to a more limited extent, American water shrews, *Sorex palustris* (Calder 1969; McIntyre 2000).

Weighing in at 12 to 17 g, American water shrews are the world's smallest endothermic divers. Well adapted to the cold, *S. palustris* primarily exploit riparian habitats within the boreal forests of North America, and can be found from Labrador in the east to above 60°N in the Northwest Territories, Yukon, and Alaska in the west (Whittaker et al. 2008). Semi-aquatic in nature, they forage both in and along the edges of ponds and fast-flowing streams and rivers, with up to 50–80% of their food intake coming from minnows, tadpoles, insect larvae, nymphs, crayfish, and other aquatic invertebrates (Conaway 1952; Sorenson 1962; but see Hamilton 1930). Somewhat surprisingly, vision plays a very limited role for underwater food detection (Svihla 1934) with prey capture success shown to be unaffected in total darkness (Catania et al. 2008). Instead, this flush-pursuit predator relies heavily on its remarkably acute nasal vibrissae to locate prey (even remotely through water movements), which they then attack with astonishing speed (<50 msec). The added ability to identify sedentary prey underwater via nasal olfaction

further cements American water shrews as one of the most agile and efficient of underwater hunters on the planet (Catania et al. 2008; Catania 2013).

Consistent with their aggressive predatory lifestyle, *S. palustris* are able to consume >10% of their body mass in a single sitting and, accordingly, have the highest mass-specific BMR of any eutherian diver examined to date (~3 times the mass-predicted value for a similar-sized mammal; Gusztak et al. 2005). Their strong positive buoyancy additionally requires constant paddling to remain submerged (Svihla 1934), and further suggests that the water shrew may also have an extremely high DMR and thus be limited to very short dives if it routinely stays within its calculated aerobic dive limit (cADL). Due to its small thermal inertia and large relative surface area, the water shrew should also lose heat rapidly while swimming and diving. As with other soricine (red-toothed) shrews, *S. palustris* are active year-round, though presumably are faced with greater energetic costs than terrestrial shrews given their aquatic lifestyle. This is especially true during winter while foraging under the ice, when convective heat loss is likely to pose a formidable challenge. Consequently, this diminutive insectivore is of interest not only in terms of its dive endurance, but also with respect to its thermoregulatory competence since, in theory, it should be highly susceptible to immersion hypothermia after even brief periods of aquatic activity.

The primary aim of this study was to investigate the dive performance and aquatic thermoregulatory ability of wild-caught *S. palustris* in both laboratory and semi-natural settings. Briefly, 20-min aquatic trials were completed to assess the diving behavior of this species, as well as test for the occurrence of behavioral thermoregulation in water shrews voluntarily diving over a range of water temperatures. To provide insight into the extent to which water shrews depend on aerobic metabolism while submerged, a second major objective was to determine this

species' cADL, which required measuring DMR and determining its total body oxygen storage capacity. Finally, to test whether this species takes advantage of immersion hypothermia to extend its dive duration while foraging in cold water, radio-telemetered water shrews were provided access to a semi-natural riparian environment (water temperature=3°C) where they were required to forage exclusively under water during 12- to 28-hr trials.

Methods

Animal Capture and Care

Sixty-seven water shrews were live trapped in Whiteshell (49°47'N, 95°13'W) and Nopiming (50°28'N, 95°15'W) Provincial Parks, Manitoba, Canada, using Sherman live traps (256 cm × 76 cm × 76 cm). For comparative analysis, 18 short-tailed shrews, *Blarina brevicauda*, were also captured. Trapping techniques and holding conditions in captivity are detailed elsewhere (Gusztak and Campbell 2004; Hindle et al. 2009). Briefly, traps baited with frozen minnows were set out overnight and inspected every 2 hr to minimize trap mortality. Water shrew traps were placed on the edges of streams and ponds with abundant sedge, while short-tailed shrews were trapped in grass thickets located a few meters inland from the water shrew sets. Immediately upon capture, shrews were placed individually into covered 38-l plastic containers fitted with screen lids and supplied with soil, dried grass and leaves, a layer of thick moss, rocks, logs, a nest box (100 mm × 100 mm × 125 mm), and water trays. Provisions included meal worms, *Tenebrio molitor*, hulled sunflower seeds, ground Purina cat food™, and a prepared meat mixture (see below), along with any invertebrates found while trapping. Shrews were transported to the University of Manitoba Animal Holding Facility on the morning following capture.

Vogel (1990) suggested that captive European water shrews lose the hydrophobic properties of their pelage when animals are not provided with holding conditions that permit diving and access to dry moss for burrowing following aquatic activity. Consequently, much care was taken in the design and continued maintenance of the holding tanks to ensure the fur of water shrews was always in optimal condition. Modified 264-l glass aquaria (88 cm × 50 cm × 60 cm) with screened lids served as individual long-term holding containers for water shrews (see Gusztak and Campbell 2004). Each holding tank had a discrete terrestrial (~75%) and aquatic (~25%) compartment separated by a 1-cm thick Plexiglas partition. Short-tailed shrews were individually housed in 76-l terrestrial containers and were supplied with water dishes that were refilled every 12 hr. For both species, the terrestrial area was furnished as described above for the transport containers. Both species were offered a prepared meat mixture (beef and chicken hearts, pig and beef liver, ground beef, fish fillets, and canned dog food mixed with vitamin and calcium supplements) every 12 hr. To ensure water shrews were habitually swimming and diving in the setup, they were required to swim across the tank and dive under a removable partition to access the feeding tray. Additionally, mealworm larvae and aquatic prey (leeches, dragonfly nymphs, and small crayfish) were occasionally placed in the aquatic portion of the water shrew tanks to encourage natural foraging behavior. When offered, aquatic prey was preferentially consumed over the meat ration.

To minimize stress, shrews were never handled by hand. Transfer to and from the holding tanks was accomplished by placing a short (~15 cm) blind-ending section of ABS tube (3.5 cm internal diameter) in the chamber, which the shrews readily entered. The tube was quickly capped and then opened following transfer, allowing the animal to exit freely. All wild shrews used in diving trials or for tissue processing were acclimated to holding conditions for a

minimum period of three weeks before diving trials were initiated. Water shrews were allowed a recovery period of at least 48 hr between successive experimental trials (see below). As the integrity of the air boundary in the fur was found to strongly influence DMR, T_b , and activity level, diving/metabolic trials in which the shrews fur became wetted were not included in any subsequent analyses.

Shrews are short-lived, with maximum life spans for both species in the wild being about 18 months (George et al. 1986; Beneski and Stinson 1987). Individual shrews were aged post-mortem based on the presence or absence of tooth growth rings following Hindle et al. (2009), and subsequently divided into juveniles (~1–3 months of age) or adults (~13–15 months). All study animals were captured under permission of Government of Manitoba Conservation trapping permits, and cared for in accordance with the principles and guidelines of the Canadian Council of Animal Care (University of Manitoba Animal Use Protocols: F01-025 and F05-014).

Body Temperature Recordings

Twelve water shrews were each equipped with a ~1.0-g model X-M transmitter (Mini-Mitter Inc., Sunriver, OR, USA) surgically implanted in the abdominal cavity. Five of these shrews underwent a second surgery 4 to 6 weeks later to implant replacement transmitters. Each transmitter was modified from the original packaging to decrease its overall size/mass and then calibrated following the method of Dyck and MacArthur (1992). Transmitter modification and surgical procedures are described in detail by Gusztak et al. (2005). Briefly, shrews were anesthetized with Isoflurane, first given at 3% for induction and then manually adjusted, as needed, between 2 and 3% to maintain a surgical plane of anesthesia. A midline incision through the skin and body wall was made along the *linea alba*. The sterilized transmitter was then placed

into the abdominal cavity and incisions closed with sutures; no mortalities were recorded following this surgical procedure. Post-operative surgical care included placement of the shrews in a disinfected 38-l plastic container containing a nest box and shredded paper towel. Shrews were supplied with fresh food and water every 12 hr and were transferred back to their holding tanks after 48 hr. Aquatic trials started 7 to 10 days later.

Voluntary Dive Behavior

We recorded the frequency and duration of voluntary dives by 25 water shrews (six of which were implanted with temperature-sensitive radio transmitters) over 20-min trials conducted in a 170.5 cm × 68 cm fiberglass dive tank filled with water to a depth of 60 cm (McIntyre et al. 2002). A transparent dive platform (17.5 cm × 68 cm) was situated at one end just above the waterline. A 4-cm section of ABS tube (internal diameter=3.5 cm) was fastened to the platform to provide a darkened refuge for the shrew. The center section of the tank contained open water (75 cm × 68 cm), while the remainder of the tank was covered with a sheet of 1-cm thick Plexiglas (78 cm × 68 cm) to encourage exploratory diving behavior. The Plexiglas sheet was equipped with handles so it could be removed quickly if a shrew became disorientated beneath.

Prey was absent from the dive tank, though shrews were offered a single mealworm 10 min before each trial. The mass of each animal was recorded just prior to feeding. To familiarize shrews with the dive arena, each animal completed a pre-trial training run in 30°C water. Behavioral dive trials were then conducted in randomized order in 3, 10, 20, or 30°C water. Trials were initiated when the shrew first entered the water and lasted precisely 20 min. Data were collected during the trial using a Sony Microcassette Recorder for post-trial analyses of dive durations and frequencies, inter-dive times, time in water, and grooming behavior.

For the implanted shrews, rates of body cooling and re-warming were also assessed via body temperature (T_b) data recorded during the 5-min pre-trial, 20-min dive trial, and 10-min post-trial periods, respectively. Radio signals from the Mini-mitters were continuously recorded via a Sony Cassette-Recorder that was placed outside the terrestrial section of the tank and T_b data subsequently analyzed at 1-min intervals.

In addition to the above 20-min dive trials, T_b , relative activity, swim/dive times, and foraging behavior of five of the radio-telemetered adult water shrews were continuously video recorded for a 12- to 28-hr period in a simulated riparian environment maintained at 3°C. Each shrew was only tested once in this setup. For these trials, the previously described fiberglass tank was shortened to 120 cm and furnished with river-washed rocks and an artificial riverbank set attached to one side (supplemental Fig. 1). The set had a ~10 cm wide bank that allowed the shrew to run along the length of the aquatic enclosure. A 60-cm long clear PVC tube linked the water tank to a clear plastic terrestrial chamber so that the shrew was always visible when active. The terrestrial chamber was housed with a nest box, moss, and rocks, and was placed on a motion activity detector (MAD-1; Sable Systems Inc.). *Ad libitum* invertebrate prey items (leeches, crayfish, mealworms) were placed in the water thereby requiring shrews to forage exclusively under water (~55 cm depth) for the duration of the trials. Water shrews were provided an 8-hr pretrial period to acclimate to the setup.

Metabolic Cost of Diving

The costs of bouts of diving, grooming, and re-warming of 12 radio-implanted water shrews voluntarily diving in a 208 cm × 55 cm fiberglass tank filled with water to a depth of 44 cm were measured with open-flow respirometry following McIntyre et al. (2002). The tank was covered

with three removable Plexiglas panels placed just below the waterline. A curved section of clear tubing 4 cm in diameter and 5 cm long connected the water tank to a 170-mL metabolic chamber constructed from a 6-cm length of Plexiglas tubing (internal diameter=6 cm). A removable rubber stopper at the rear of the chamber facilitated shrew transfer to and from the setup, while a removable partition was placed between the metabolic chamber and tank cover to prevent shrews from entering and exiting the water while pre- and post-dive metabolic measurements were recorded.

An outlet port was installed in the top/rear portion of the metabolic chamber while six inlet holes, each 1 mm in diameter, were drilled into the base of the chamber at the opposite end to facilitate mixing of air. Air was drawn sequentially at $\sim 500 \text{ mL min}^{-1}$ from the chamber and through a column of Drierite using a TR-SS1 gas analysis sub-sampler (Sable Systems Inc., Las Vegas, USA) calibrated against a bubble flowmeter (accurate to within $\pm 2\%$; Levy 1964). A subsample of this dry exhalent gas was analyzed using an Applied Electrochemistry S-3A O_2 analyzer. Fractional O_2 content was recorded at 1-sec intervals, while water (T_w) and chamber temperature (T_a) were recorded immediately prior to and following each trial. A respiratory quotient of 0.83 was assumed, based on an earlier study of fasted water shrews (Gusztak et al. 2005), and instantaneous $\dot{V}\text{O}_2$ derived following the method of Bartholomew et al. (1981). Mean instantaneous $\dot{V}\text{O}_2$ measurements were calculated at 20-sec intervals throughout the trial. Body temperature signals were recorded on a Sony Cassette-Recorder and subsequently analyzed at 1-min intervals throughout the pre-trial, diving, and post-trial periods, respectively.

Concrete blocks and sections of ABS tubing were placed at the bottom of the tank to encourage longer exploratory dives, since preliminary trials suggested these objects increased time under water. Each shrew completed a total of four trials presented in random order: two in

10°C water and two in 30°C water. Trials conducted at 30°C provided an estimated DMR when thermoregulatory costs are presumably minimal (MacArthur 1989), while 10°C water was chosen to assess the thermoregulatory costs associated with submersion in cold water.

The mass of each shrew (corrected for telemeter mass) was recorded immediately before placing the animal in the metabolic chamber. Each trial consisted of a 10- to 15-min pre-trial period during which time the shrew was confined to the metabolic chamber and its lowest $\dot{V}O_2$ over 5 min taken as the baseline value. After the partition was removed, 10 min was allotted for voluntary diving, which commenced upon the animal's first entry into the water. At the end of the 10-min dive session, the partition was gently slid back into place to prevent further dives. The post-dive recovery $\dot{V}O_2$ associated with re-warming and grooming was recorded until the animal's $\dot{V}O_2$ or T_b returned to within 95% of the pre-trial baseline. If this had not occurred after 15 min in the chamber, the trial was ended. Dive durations, grooming behavior, and relative activity were recorded on a Sony Microcassette-Recorder and analyzed after each trial.

Body Oxygen Stores

The total body oxygen stores of water shrews were determined after completing all diving/behavioral trials on each animal, while short-tailed shrews were allowed a 1- to 3-week acclimation period in the lab prior to determining their oxygen reserves. The mass of each shrew was recorded, after which the animal was deeply anesthetized with 3% Isoflurane inhalant anesthetic. A cardiac puncture was then performed to extract a blood sample for hemoglobin and hematocrit determinations (MacArthur 1984b; McIntyre et al. 2002), followed by euthanization via an overdose of Isoflurane. The heart, forelimb, and hindlimb muscles were then quickly removed and freeze-clamped in liquid nitrogen. Excised muscles were stored at -70°C and

myoglobin content and buffering capacity determined later following the methods of Reynafarje (1963) and Castellini et al. (1981), respectively. The lungs were carefully removed after the majority of the heart muscle had been cut away, and lung volume determined gravimetrically following the procedures described by Weibel (1970/71). This involved inserting a 3-cm section of P20 cannula 5-8 mm into the trachea, with the cannula secured in place with a 5-0 silk ligature. Vetbond™ was applied to the knot at the juncture of the cannula/trachea to ensure the preparation would not slip. The trachea/lung prep was submerged in saline (0.9 M NaCl) and then inflated at a constant pressure of 20 mmHg with humidified air for ~10–15 min before measurement. All volume measurements were corrected to STPD.

The total percentage of muscle mass, expressed as a fraction of digesta-free body mass, was also calculated for 12 water shrews and two short-tailed shrews. Skinned, eviscerated carcasses were submerged for ~24–48 hr in a detergent solution at 32°C to detach any skeletal muscle adhering to the bones. The total skeletal muscle mass was then calculated by subtracting the bone mass from the initial carcass mass (MacArthur et al. 2001; McIntyre et al. 2002).

Total blood volume of water shrews was estimated from the allometric equation of Prothero (1980): $\text{blood volume (mL)} = 76 M^{1.0}$, where M = body mass in kg. Total body O₂ stores (mL O₂ in muscle, blood and lungs, corrected to STPD) of *S. palustris* were determined following the procedures of Lenfant et al. (1970). This method assumes that water shrews dive with lungs fully inflated and with an initial lung oxygen concentration of 15%. The oxygen storage capacity of blood was calculated by assuming that 1/3 and 2/3 of the blood volume constituted the arterial and venous fractions, respectively, with the former having an oxygen saturation of 95% and the latter a 5% vol. decrease in O₂ content compared to arterial blood. Skeletal muscle myoglobin reserves were determined as the mean concentration of forelimb and hindlimb samples for each

individual, multiplied by the estimated mass of skeletal muscle in the body. Blood and myoglobin oxygen capacities were presumed to equal $1.34 \text{ mL O}_2 \text{ g pigment}^{-1}$ (Lenfant et al. 1970; Kooyman et al. 1983).

Total body oxygen stores were divided by the mean DMR ($\text{mL O}_2 \text{ sec}^{-1}$) measured in 10 and 30°C water, in order to derive a cADL for water shrews diving at each T_w . This estimate hinges on the assumption that all O_2 stores are utilized during diving, before the animal switches to anaerobic respiration (Kooyman et al. 1980). We also calculated the behavioral ADL (bADL) at each T_w , defined as the dive duration exceeded by only 5% of all voluntary dives (Kooyman et al. 1983).

Skeletal Muscle Buffering Capacity

The skeletal muscle buffering capacities of short-tailed and water shrews were tested against non-bicarbonate buffers following the procedure of Castellini et al. (1981). Buffering capacity (slykes) was standardized to represent the μmol of base needed to increase the pH of 1 g of wet muscle mass from a pH of 6 to 7. A 0.3- to 0.5-g sample of frozen skeletal muscle comprised of a mixture of forearm and hindlimb tissue was ground up in 0.9 M NaCl, following which the solution was titrated with 0.2 M NaOH using an Accumet® AB 15/15+ pH meter and an AccuTupH sensing electrode.

Statistical Analyses of Data

All statistical analyses were performed using SPSS 9.0 for Windows. Means of dive variables were compared across water temperatures using one-way ANOVA, while two-way ANOVA was employed to test for possible interaction effects between T_w and the presence or absence of an

implanted transmitter in the subject animal. When appropriate, differences between means were compared using Tukey's multiple range test. Dive profiles were compared between implanted and non-implanted adult water shrews using the log likelihood ratio test (G-test, Zar 1974). Means of variables were compared using a 1-tailed Students *t*-test. Regression lines were fitted by the method of least squares. Significance was set at 5% and means presented as ± 1 SE.

Results

Voluntary Dive Behavior

In most trials, water shrews were hesitant to dive until they had fully explored the surfaces of both the terrestrial and aquatic sections of the tank. Subsequently, predictable pre-dive behavior was routinely observed. Water shrews would approach and pause at the edge of the dive platform for 1–10 sec, during which time repetitive head nodding occurred, causing the shrew's vibrissae to repeatedly touch the water. This behavior was typically followed by the shrew diving from the platform.

Water shrews engaged in two distinct categories of dives. Dives were classified as either shallow (<10 cm) or deep (reaching the tank bottom at 60 cm). Very few dives were completed between these depths, but if they occurred, were specified as shallow. During each 20-min dive trial, the mean number of dives completed during the first 5-min was significantly greater than for any of the three remaining 5-min periods ($F_{3,92}=27.466$, $P<0.0001$), with ~50% of voluntary dives occurring during the first 5 min of the trial (supplemental Fig. 2).

During each 20-minute trial, shrews completed, on average, 14.7 dives (range=0 to 53 dives). No significant differences in diving duration or dive frequencies were found between juvenile and adult water shrews, nor between radio-implanted and non-implanted individuals (see below),

and hence all data were pooled for subsequent analyses. The mean dive time of 25 shrews (111 individual dive trials) was 5.1 ± 0.1 sec (supplemental Fig. 3), with a median of 4.5 sec. Of the 1584 recorded voluntary dives, 311 (19.6%) were deep dives with an average duration of 8.1 ± 0.2 sec. The five longest dives of each trial had a mean duration of 7.8 ± 0.3 sec, while the mean longest dive per trial was 10.3 ± 0.4 sec. Only three dives exceeded 20 sec with the longest voluntary dive recorded being 23.7 sec.

The average dive:surface ratio was 0.21 ± 0.01 sec. While short dives were often immediately followed by another dive, all dives >13 sec were accompanied by an extended (>30 sec) surface interval (supplemental Fig. 4). There was also a significant increase in the inter-dive surface time with longer dives, suggesting that longer dives require a longer recovery period than shorter dives ($F_{1,952}=20.280$, $P<0.0001$, $r^2=0.021$). This relationship held when the longest dives (>20 sec) were removed from the analysis ($F_{1,950}=14.594$, $P<0.0001$, $r^2=0.015$).

Influence of Water Temperature on Dive Behavior

A total of 18 shrews completed voluntary dive trials at all four water temperatures. For all dives combined, dive frequency was independent of T_w ($F_{3,67}=0.467$, $P=0.706$). However, T_w significantly influenced the total time water shrews spent swimming and diving ($F_{3,67}=7.892$, $P<0.001$), with shrews voluntarily spending less than half as long in 3°C compared to 30°C water (Table 1). Mean dive duration increased with T_w ($F_{3,67}=5.033$, $P=0.003$; Fig. 1) as did the dive:surface ratio ($F_{3,67}=7.146$, $P<0.001$). On average, shrews surfaced for 101 s and 66 sec before diving again in 3 and 30°C water, respectively. T_w also affected both the five longest ($F_{3,67}=5.229$, $P=0.003$) and single longest dive of each trial ($F_{3,67}=5.866$, $P=0.001$), with water

shrews diving an average of 35% longer in 30°C compared to 3°C water (Fig. 1). The bADL of water shrews progressively increased from 8.8 sec in 3°C water to 12.2 sec in 30°C water.

Influence of Transmitter Implants on Dive Performance

Dive performance was compared in twelve adult water shrews with ($N=6$) and without ($N=6$) an implanted 1.0-g abdominal temperature transmitter by first pooling the dive data for each group across all T_w 's. There were no significant differences in dive performance between the two groups in any of the examined variables (supplemental Table 1). The frequency distribution of dives by implanted and non-implanted shrews was also compared for dives times of 0 to 15 sec, using log likelihood ratios to ensure the calculated mean value was not influenced significantly by outliers (supplemental Table 2). Again, no significant difference was found between implanted and non-implanted water shrews (G value=2.78, $df=5$, $P>0.50$).

We also determined if the dive behavior of water shrews was significantly influenced by the interaction of transmitter effect and T_w , using a 2-way ANOVA. No significant interaction effects were found for any variable, including dive duration ($F_{3,40}=1.013$, $P=0.397$), dive frequency ($F_{3,40}=0.304$, $P=0.822$), dive:surface ratio ($F_{3,35}=0.378$, $P=0.770$), deep dive duration ($F_{3,30}=1.182$, $P=0.333$), deep dive frequency ($F_{3,40}=0.032$, $P=0.992$), dive:surface ratio of deep dives ($F_{3,20}=0.409$, $P=0.748$), the longest dive for each trial ($F_{3,40}=0.659$, $P=0.582$), and the total time in water ($F_{3,40}=2.321$, $P=0.0898$).

Body Temperature Patterns of Implanted Shrews During Voluntary Diving

A total of 23 T_b data sets were obtained from six water shrews that completed a total of 24 dive trials. Owing to a mechanical error with the recording device, no T_b data were obtained for one

water shrew diving in 10°C water, though behavioral dive data were collected for this animal. Water shrews usually incurred the largest drop in T_b during the first 5–7 min of each trial, but the decrease varied with T_w (Fig. 2). Following this initial curvilinear decline, T_b tended to plateau and was regulated near 38.8°C for the remainder of the dive trial. The lowest recorded T_b during the diving session did not differ significantly between T_w 's ($F_{3,19}=2.341$, $P=0.11$). Water shrews quickly elevated their T_b during the post-trial period, and attained a T_b similar to the pre-trial value regardless of T_w (Table 2; Fig. 2).

24-Hour Dive Trials

Body temperature profiles and entry and exit times to and from the water were obtained from five shrews studied in the 12- to 28-hr trials. However, reliable submergence times were only obtained for three of the trials due to differences in camera position. While substantial variability was apparent, foraging activities were typically clustered into discrete bouts that consisted of 9.9±2.5 aquatic excursions and lasted 38.1±9.1 min each, with individual bouts separated by 65.7±8.8 min. Water shrews entered the water an average of 141 times per 24-hr period (range: 92-212) with a prey capture success rate of 28.6%, or 40 prey items per day. The dive times of shrews actively foraging in the simulated riverbank enclosure were skewed towards longer duration dives (Fig. 3) than observed in the 20-min trials (Fig. 1). Consequently, the mean dive duration in the former set up (6.9±0.2 sec; $n=227$) was not only significantly longer than that of water shrews voluntary diving in 3°C water with no access to food (4.0 sec; $t=12.25$, $df=482$, $P<0.0001$), but also longer than the mean dive duration of dives completed at the four different water temperatures (5.1 sec; $t=7.92$, $df=809$, $P<0.0001$). Despite the right-shifted distribution, an abrupt drop in dive duration occurred between 9 and 11 sec yielding a bADL of 10.7 sec (Fig 3).

Body temperature profiles of free-ranging shrews exhibited several consistent patterns over the course of the trials. Principal among these was a sharp elevation in T_b by $\sim 1.2^\circ\text{C}$ in the ~ 10 min period immediately prior to initiating each aquatic foraging bout (Fig. 4). Consequently, mean T_b often averaged $>39.0^\circ\text{C}$ at the commencement of the first dive, though routinely dropped by $0.2\text{--}0.4^\circ\text{C}$ upon exiting the water which was followed by a further reduction in T_b (~ 0.5 to 1.0°C). While shrews occasionally initiated another dive at T_b 's $<37.5^\circ\text{C}$, in most cases T_b was sharply increased again prior to the next entry into water (Fig. 5 and supplemental Fig. 5). During the interbout intervals activity was minimal and core T_b was regulated near 37.5°C .

Diving Metabolic Rate

The costs of repetitive diving and re-warming were determined for 12 implanted water shrews that completed 525 dives over 48 trials. Reliable T_b measurements were not obtained for 6 of the 48 trials, due to weak/absent signals from some transmitters. During the 10-min period available for voluntary diving, shrews spent an average of 51 ± 7 sec (8.5% of total time) and 78 ± 10 sec (13% of total time) diving in 10 and 30°C water, respectively ($t=2.24$, $df=38$, $P=0.01$). As with the behavioral dive trials (see above), mean dive time in 10°C water (3.8 ± 0.2 sec; $n=256$) was significantly shorter than during the 30°C metabolic trials (5.4 ± 0.3 sec; $n=269$) ($t=2.54$, $df=32$, $P=0.007$).

Integrity of the fur air boundary layer and level of activity in the metabolic chamber were both found to significantly affect T_b and increase recorded DMR estimates (data not shown). Consequently, we only utilized DMR values from water shrews whose pelage did not show evidence of wetting during the trial and that displayed minimal terrestrial activity (Fig. 6). This resulted in estimates of DMR in only 13 of the 48 trials (five and nine from the 10°C and 30°C

trials, respectively). The mean DMR for shrews diving in 10°C water ($8.77 \pm 0.30 \text{ mL O}_2 \text{ g}^{-1} \text{ hr}^{-1}$) was 1.33× greater than that in 30°C water ($6.57 \pm 0.27 \text{ mL O}_2 \text{ g}^{-1} \text{ hr}^{-1}$).

Testing for Adaptive Hypothermia

Both regional heterothermy (peripheral hypothermy) and adaptive hypothermia have been forwarded as mechanisms that allow air breathing divers to extend their aerobic dive durations via temperature (Q_{10}) induced reductions in metabolism (Favilla and Costa 2020). While it is unlikely that small-bodied water shrews can sustain meaningful peripheral heterothermy, reductions in core T_b may potentially allow them to extend their underwater endurance. Thus, voluntary dive times and concomitant core T_b were analyzed to test for potential linkages between these variables. We first plotted dive duration against T_b to determine if shrews with lower T_b 's dove longer in cold water. To maximize the data set, dive data were combined for shrews completing voluntary dive trials in 3°C and 10°C water and for shrews completing the DMR trials in 10°C water. A statistically significant negative relationship was found between dive duration and T_b ($r^2=0.0795$, $P<0.0001$, $n=309$; Fig. 7a). Notably, shrews freely foraging in 3°C water in the semi-natural environment exhibited a similar trend ($r^2=0.0461$, $P=0.0103$; Fig. 7b). However, a true measure of adaptive hypothermia should reflect reductions in core T_b being accompanied by a greater proportion of dive times above the cADL. Limiting these regression analyses to only voluntary dives that exceeded the predicted cADL at these temperatures (~10 sec) did not return significant correlations ($r^2=0.002$, $P=0.801$, $n=27$ and $r^2=0.001$, $P=0.871$, $n=28$, respectively). Indeed, water shrews were reluctant to dive when cool, with few dives occurring at T_b 's $<37.5^\circ\text{C}$ (Figs. 7c, d).

Body Oxygen Stores and Muscle Buffering Capacity

Myoglobin concentration did not differ significantly between forelimb and hindlimb samples for either adults or juveniles in the two shrew species sampled (Table 3). The mean (forelimb and hindlimb) myoglobin concentration (mg g of wet tissue⁻¹) of adult water shrews (6.04 ± 0.25 ; $N=9$) was significantly higher than in juveniles (3.76 ± 0.12 ; $N=10$; $t=7.074$, $df=18$, $P<0.0001$) and nearly 2× greater than for adult short-tailed shrews (3.03 ± 0.12 ; $N=9$; $t=7.624$, $df=17$, $P<0.0001$). Skeletal muscle buffering capacity exhibited a similar trend, with adult water shrews having a higher value (38.22 ± 2.28 slykes; $N=13$) than either juvenile water shrews (30.67 ± 2.34 slykes; $N=11$; $t=2.395$, $df=22$, $P=0.012$) or adult short-tailed shrews (24.88 ± 1.40 slykes; $N=10$; $t=4.808$, $df=21$, $P<0.0001$).

Recorded lung volumes of adult (4.57 ± 0.28 mL STPD 100 g⁻¹; $N=5$) and juvenile (4.55 ± 0.18 mL STPD 100 g⁻¹; $N=9$) water shrews were similar (Table 3). These values are ~1.1× greater than predicted by allometry (Stahl 1967) for a mammal of 15.2 g (adult) and 14.1 g (juvenile). Lung volumes of adult short-tailed shrews (3.34 ± 0.07 mL STPD 100 g⁻¹; $N=6$) were significantly less than those recorded for adult water shrews ($t=7.401$, $df=9$, $P<0.0001$) and 6% less than that predicted by allometry for a 22.8 g mammal.

Total blood oxygen capacity of adult and juvenile water shrews was high, averaging 26.93 ± 0.58 and 26.38 ± 1.37 vol. %, respectively (Table 3), with adult water shrews having a significantly higher blood oxygen capacity than adult short-tailed shrews (23.18 ± 0.84 vol. %; $t=0.896$, $df=9$, $P<0.0001$). Hematocrit levels were high in both adult water shrews (50.23 ± 1.33) and short-tailed shrews (48.78 ± 2.50), and did not differ statistically between the two species ($t=0.735$, $df=21$, $P<0.235$). The calculated total O₂ storage capacity of summer-caught adult and juvenile water shrews was 26.31 and 24.06 mL O₂ STPD kg⁻¹, respectively (Table 4). The mass-

specific O₂ storage capacity of summer-caught adult water shrews was 1.31× greater than that of adult short-tailed shrews (Table 4). The largest single contributor to total O₂ stores in adults of both shrew species was O₂ bound to hemoglobin in the blood, accounting for 62.5% and 68.5% in water shrews and short-tailed shrews, respectively.

Calculated Aerobic Dive Limit

Assuming that diving animals fully deplete their O₂ reserves before initiating anaerobic respiration, and using the DMR estimate of 6.57 mL O₂ g⁻¹ hr⁻¹ in 30°C water, adult and juvenile American water shrews have cADLs of 14.4 and 13.2 sec, respectively (Table 4). This value is consistent with the estimated bADL at 30°C (12.2 sec), determined from voluntary dive profiles ($n=303$ dives; Fig. 1). Due to an increased DMR in 10°C water (8.77 mL O₂ g⁻¹ hr⁻¹), the cADL for adult and juvenile water shrews is 10.8 and 9.9 sec, respectively (Table 4). Again, these calculated limits closely match the bADL (10.2 sec) determined for water shrews diving in 10°C water ($n=258$ dives; Fig. 1). Only 3.9% and 2.3% of voluntary dives in 10°C and 30°C water, respectively, exceeded the cADLs for adult *S. palustris*.

Discussion

As predicted by allometry, the diminutive size of *S. palustris* in this study (mean=14.0 g) appeared to severely limit the species' aerobic dive endurance. Indeed, adult water shrews were found to exhibit the smallest total oxygen storage capacity (0.37 mL O₂), the highest mass-specific DMR (6.57–8.77 mL O₂ g⁻¹ hr⁻¹), and lowest cADL (10.8–14.4 sec) of any mammalian diver studied to date. Moreover, water shrews have the highest surface-area-to-volume ratio of any endothermic diver, which should make them extremely susceptible to immersion

hypothermia (MacArthur 1989). It is truly remarkable then, that this small insectivore can efficiently utilize aquatic foraging as a means to sustain its inherently high rate of heat production, since aquatic foraging is known to be one of the most energetically costly methods of foraging (Fish 2000). This energetic burden is highlighted by the observation that captive American water shrews in relatively warm terrestrial settings ($\sim 20^{\circ}\text{C}$) must consume at least their body weight in prey on a daily basis (Gusztak et al. 2005). This level of food consumption could conceivably be doubled or tripled in free-living individuals diving in ice-covered streams during the winter months.

Body Oxygen Stores and Muscle Buffering Capacity

Divers are known for their physiological adaptations to increase breath-hold capacity, especially those relating to their disproportionately large body oxygen stores (Butler and Jones 1997; Favilla and Costa 2020). Not surprisingly, the mass-specific oxygen stores of water shrews were $\sim 30\%$ greater than those of strictly terrestrial short-tailed shrews. Many researchers in this field consider the primary indicator of a diver's breath-hold capacity to be its muscle oxymyoglobin concentration (Kooyman and Ponganis 1998, Ponganis et al. 1999; Mirceta et al. 2013). Adult *S. palustris* appears not to follow this trend as myoglobin accounts for only 12.4% of total body O_2 stores, which is similar to that of alcids ($\sim 5\text{--}10\%$), but substantially lower than values (~ 30 to 50%) reported for various pinnipeds and toothed whales (Elliott et al. 2010; Favilla and Costa 2020). Moreover, skeletal muscle myoglobin concentration of adult water shrews ($0.60 \text{ g } 100 \text{ g}^{-1}$; Table 3) is only half that recorded for other semi-aquatic mammals: star-nosed moles ($1.36 \text{ g } 100 \text{ g}^{-1}$; McIntyre et al. 2002), muskrats ($1.21\text{--}1.38 \text{ g } 100 \text{ g}^{-1}$; MacArthur et al. 2001), and beaver ($1.2 \text{ g } 100 \text{ g}^{-1}$; McKean and Carlton 1977). Similar to other divers, however, myoglobin

stores are relatively slow to develop as values for juvenile (<4-month old) water shrews were ~62% those of adults, and only accounted for 7.7% of total body O₂ storage capacity.

The relatively low myoglobin concentrations in small-bodied alcid and semi-aquatic mammals likely arises from functional constraints imposed by muscle fiber types and relatively high mitochondrial volumes (Weibel 1985; Ordway and Garry 2004) that limit maximal attainable levels in muscle. Nonetheless, skeletal muscle myoglobin concentrations of adult water shrews are 2–4× that of strictly terrestrial short-tailed (0.16 to 0.30 g 100 g⁻¹; Stewart et al. 2005 and Table 3) and Etruscan shrews, *Suncus etruscus* (0.15 g 100 g⁻¹; Jürgens 2002), suggesting an adaptive increase associated with diving. This difference appears to be linked to the marked increase in the net surface charge of *S. palustris* myoglobin versus those of non-diving shrews, which is proposed to foster higher myoglobin concentrations by minimizing the potential for newly synthesized (apomyoglobin) and mature protein chains from aggregating/precipitating (Mirceta et al. 2013; Samuel et al. 2015; He et al. 2021). Of note, the highest myoglobin concentrations were found in the heart for both short-tailed (0.88 g 100 g⁻¹) and water shrews (1.10 g 100 g⁻¹). This finding presumably highlights a key role for this respiratory pigment in supporting the exceptionally high heart rates (>750 beats per minute) of soricids (Doremus 1965; Vornanen 1992), though this feature may also help extend aerobic metabolism of this tissue during long dives by water shrews.

The water shrew is able to compensate for its low skeletal muscle myoglobin concentration owing to the potential gain in O₂ stores in the lungs and, especially, the blood. The mean lung volume of adult water shrews (4.57 mL STPD 100 g⁻¹; Table 3) was ~1.1× greater than predicted by allometry for a 15.2 g mammal (Stahl 1967). Further, this species has a mass-specific pulmonary O₂ storage capacity that is 1.37× greater than for adult short-tailed shrews. The mass-

specific lung volume of the American water shrew is also larger than for other shrews studied, including, *N. fodiens*, *Sorex minutus*, and *Suncus etruscus* with respective lung volumes of 3.56, 3.81, and 4.02 mL STPD 100 g⁻¹ (Gehr et al. 1981). The moderately higher lung O₂ stores of *S. palustris* could serve to increase buoyancy, as well as provide an important source of O₂ while diving. It should also be noted that underwater sniffing plays an important role in food acquisition for this species (Catania 2006; Catania et al. 2008). In this context, the enlarged lungs of American water shrews may at least partially compensate for exhaled gas bubbles lost incidental to underwater sniffing (Gusztak, unpublished observations). However, lung volume of *S. palustris* is almost half that of the adult star-nosed moles (8.1 mL STPD 100 g⁻¹; McIntyre et al. 2002), which also exploits underwater olfaction (Catania 2006). Another relevant observation is that submerged water shrews would routinely inspire trapped gas released from their fur that coalesced under the Plexiglas cover during the initial training sessions. This finding implies an ability to re-breathe air bubbles trapped under the ice during winter foraging in order to extend submergence times as has been observed in muskrats (MacArthur 1992). Since water shrews commonly co-habit waterways with beavers and muskrats (Conaway 1952), it is expected that bubbles can similarly be exploited by *S. palustris* diving under the ice.

By far, the most important source of O₂ for a diving water shrew is the large reserve in the blood, which comprises 62.5–66.2% of this diver's O₂ storage capacity. This is reflected in their high hemoglobin (19.9 g 100 mL⁻¹) and hematocrit (50.4%) values, with the former resulting in a mean blood O₂ capacity of 26.9 vol. % for an adult water shrew. Many species of shrews examined to date have high recorded hemoglobin (range: 15–18 g 100 mL⁻¹) and hematocrit values (range: 45–50%), some of which are near the upper limits recorded for any mammal (Wolk 1974; Gehr et al. 1981). Even so, the blood O₂ capacity calculated for the American water

shrew is the highest value recorded of any soricid, including the European water shrew (23.9 vol. %; Wolk 1974), the Etruscan shrew (23.3 vol. %; Bartels et al. 1979) and adult short-tailed shrew (23.2 vol. %; Table 3). Adult water shrews also exhibited a higher blood O₂ capacity than other semi-aquatic divers studied to date, including adult star-nosed moles (23.0 vol. %; McIntyre et al. 2002) and muskrats (24.1 vol. %; MacArthur et al. 2001).

Buffering capacity of skeletal muscle has been shown to increase with increasing body mass and is also important for prolonging burst activity in species utilizing sprinting (Castellini et al. 1981; Hochachka and Mommsen 1983). Specifically, elevated buffering capacity is correlated with Type IIA and IIB fast-twitch muscle fibers, which are found in higher concentrations in mammals that exploit burst activity (Abe 2000; Nakagawa and Hattori 2002). In anaerobic or severely hypoxic conditions, as may be encountered during prolonged diving or for shorter periods of time during sprinting, glycolysis is the only, albeit inefficient, means of ATP production. This process is inhibited if intracellular pH drops too low, therefore intracellular buffers are critical to ensure an optimum pH for glycolysis to occur while exercising in low O₂ environments. Adult water shrews have a skeletal muscle buffering capacity (38.2 slykes) that is 1.2× greater than for sub-adult water shrews and significantly greater (1.5-fold) than for adult short-tailed shrews. Although allometric scaling of glycolytic enzymes to mass predicts that water shrews should have the poorest ability of any diver to utilize anaerobic glycogenolysis (Emmett and Hochachka 1981), adult water shrew values were similar to platypus (38.2 slykes; Evans et al. 1994), star-nosed moles (44.1 slykes; McIntyre et al. 2002) and summer-caught adult muskrats (51.5 slykes; MacArthur et al. 2001). Since few voluntary dives by water shrews exceeded the cADL, glycolytic pathways probably do not play a large role in the majority of dives completed by this species. Instead, their elevated muscle buffer capacity presumably

attenuates tissue pH changes arising from the rapid rate of CO₂ accumulation during aerobic breath hold dives, though it may also be important for their flush-pursuit aquatic foraging strategy that involves periods of brief, intense motor activity (Catania et al. 2008).

Diving Metabolic Rate

Consistent with allometric predictions, the American water shrew has the highest recorded DMR of any diver (6.57 and 8.77 mL O₂ g⁻¹ hr⁻¹ in 30°C and 10°C water, respectively). The high DMR of *S. palustris* reflects in part its inherently high mass-specific BMR (4.84 mL O₂ g⁻¹ hr⁻¹; Gusztak et al. 2005), as well as the rapid limb strokes, high buoyancy, and disproportionately high drag while diving due to the species' high mass-specific surface area. However, the DMR of *S. palustris* exposed to minimal thermal stress is only 1.4× BMR, which is less than star-nosed moles (2.1× RMR; McIntyre et al. 2002) and muskrats (2.7× BMR; MacArthur and Krause 1989), but similar to sea otters diving and capturing prey (1.6× RMR on water; Yeates et al. 2007).

Dive Performance

Prior to this study, very little was known about the voluntary breath-hold capacity of *S. palustris*. Our captive water shrews exhibited a mean voluntary dive duration of 5.1 sec during the 20-min dive trials, which is virtually identical to the value (5.2 sec) reported by McIntyre (2000) for a single captive specimen, and is consistent with the ranges reported for *S. palustris* (1–15 sec; Svihla 1934; Calder 1968) and *N. fodiens* diving in captivity (3–6 sec; Churchfield 1985; Ruthardt and Schröopfer 1985). Vogel (1998) suggested that the dive times of captive water shrews were probably shorter than those of shrews in the wild due to a small tank size and

absence of aquatic prey. However, we believe that our study provides a reasonably accurate representation of the American water shrew's dive capacity as the mean dive time of this species is slightly longer than predicted from allometry (cf. Fig. 4 of Jordaan et al. 2021). The maximum recorded dive time of 23.7 sec is also 12% greater than that predicted for a 16.3 g diver (21.2 sec; Schreer and Kovacs 1997), and is virtually identical to the maximal dive time (24 sec) for a slightly larger European water shrew diving in a 2-m deep stream (Scholetch 1980). That said, dive times in the 20-min dive trials are likely skewed towards shorter mean (but not maximum) durations than expected in nature. Indeed, the mean submergence time of water shrews foraging in 3°C water in the artificial riverbank environment (6.9 sec) was ~75% longer than that (4.0 sec) of shrews diving at the same temperature without access to food.

Even under laboratory conditions, *S. palustris* is an impressive diver compared to other small-bodied divers. Another semi-aquatic insectivore, the star-nosed mole, is ~4× larger (50–60 g) than the water shrew, yet exhibits a mean dive duration that is only 1.8 times greater (9.2 sec; McIntyre et al. 2002). Likewise, most dive times by the smallest avian diver, the dipper (*Cinclus mexicanus*; ~50 g), in the wild are even shorter at <6 sec (Calder 1968), while the much larger 325–450 g bufflehead (*Bucephala albeola*) has a mean dive time of only 12.5 sec (Gauthier 1993). Larger mink (*Mustela vison*; 850 g) and spotted neck otters (*Hydrictis maculicollis*; 4 kg) similarly have mean dive duration of just 9.9 and 8.5 sec, respectively (Dunstone and O'Connor 1979; Jordaan et al. 2021). Juvenile (300–500 g) and adult muskrats (650–900 g) exhibited only slightly longer average dive times (19.2–22.0 sec; MacArthur et al. 2001), while a mean of 31.3 sec was recorded for 1.5–2.0 kg free-ranging platypus, *Ornithorhynchus anatinus*, diving in a lake (Bethge et al. 2003).

Transmitter mass (1.00–1.16 g) ranged from 6.7–9.7% of total body mass, and did not surpass the recommended upper limit of 10% recommended by Brander and Cochran (1967). Radio equipped water shrews did not show any significant changes in dive performance compared to non-implanted animals. The large volume of air trapped in the pelage probably accounts for this lack of effect on dive performance. McIntyre (2000) measured the volume of air bound to the pelage in the water shrew and recorded a mass-specific air capacity of 0.35 mL g^{-1} , which is greater than in the star-nosed mole (0.19 mL g^{-1} ; McIntyre 2000). Not surprisingly, American water shrews also have a lower specific gravity (0.761), or stronger buoyant force in water, compared to star-nosed moles (0.826; McIntyre 2000) and muskrats (0.952; MacArthur 1992), but less than that of European water shrews (0.726; Köhler 1991). The relatively large amount of air trapped in the pelage of *S. palustris* should increase the energetic costs of submerging to depth. However, since surfacing is both passive and rapid (Svihla 1934) the energetic cost and travel time while ascending the water column is minimized, thereby lessening the probability of prey escaping. Indeed, captive *S. palustris* were routinely observed passively surfacing with 2–3 g crayfish captured at the bottom of a 30-cm tank (Gusztak, unpublished observations). Likewise, Köhler (1991) recorded that *N. fodiens* could retrieve a 12.8 g lead-filled snail shell from the tank bottom.

Aerobic Dive Limits and Diving Behavior

The relevance of determining cADL for certain diving species has been called into question since some large-bodied, benthic divers tend to have a significant number of dives exceeding their cADL. For instance, deep diving Australian and New Zealand sea lions routinely exceeded their cADLs by 1.4 and 1.5-fold, respectively (Costa et al. 2001), while beaked whales may exceed

their bADL by an astonishing 2.4 hrs (Quick et al. 2020). By contrast, small amphibious divers <2 kg submerge to relatively shallow depths with less interspecific variability in dive depth than larger species, and have been noted to rarely dive beyond their cADL. For instance, the star-nosed mole (50 g) and muskrat (680 g) only exceeded their cADLs during 2.9% and 6% of all voluntary dives, respectively (MacArthur et al. 2001; McIntyre et al. 2002). In line with these observations, only 3.9% and 2.3% of dives by adult water shrews in 10 and 30°C water exceeded their cADLs, respectively. Similarly, dive duration frequencies exhibited an abrupt drop between 9 and 11 sec during the 24-hr trials, yielding a bADL (10.7 sec) virtually identical to the cADL of adult water shrews diving in 10°C water (10.8 sec). Taken together, these observations suggest a strict adherence to an aerobic diving schedule in this species. Thus, determining the cADL of small-bodied divers seems to provide a useful estimate of their breath-hold capacity, even though this is not always the case for larger divers.

Influence of Water Temperature on Dive Behavior and Body Temperature

Presumably because it is energetically more efficient to prevent large drops in T_b than incur the costs of re-warming from a sharply depressed T_b , many small amphibious divers decrease or limit core T_b cooling behaviorally by periodically exiting the water to re-warm (MacArthur 1989; Hindle et al. 2006). Based on voluntary dive trials in 3, 10, 20, and 30°C water, shrews did not consistently show a statistically significant difference in any measured variable across adjacent T_w 's. Instead, for most variables examined, there was a gradual positive relationship between the behavioral index of dive performance and increasing T_w . However, and consistent with behavioral thermoregulation, there was a large difference in dive performance between the two extreme T_w 's. Comparing 3 and 30°C water, *S. palustris* experienced a 2-fold increase in the

total time spent in water, a 1.5-fold increase in mean dive duration, a 1.6-fold increase in the longest dive duration, a 1.4-fold increase in deep dive duration, and a $\sim 3\times$ greater increase in deep dive frequency. While many other small-bodied semi-aquatic mammals and birds display a similar positive correlation of dive duration and frequency with T_w (MacArthur 1984a; de Leeuw et al. 1999; McIntyre et al. 2002), free-diving mink and European beaver do not (Harrington et al. 2012; Graf et al. 2018).

Regardless of water temperature, water shrews experienced the greatest drop in T_b during the first 5 minutes of each diving trial, which then plateaued near $38.5\text{--}39.0^\circ\text{C}$ for the remaining 15 min. This latter interval was accompanied by a sharp reduction in aquatic activity, suggesting that water shrews principally rely on behavioral thermoregulation, likely supplemented by a fully engaged vasopressor response, to limit further declines in T_b during bouts of aquatic activity. Similarly, free-ranging muskrats and beavers were observed to maintain core T_b within $\pm 1^\circ\text{C}$ of normothermic T_b throughout most semi-aquatic activity, and also appeared to use behavioral thermoregulation to limit body cooling (MacArthur 1979; Dyck and MacArthur 1992).

As mentioned earlier, the air boundary surrounding the water shrew adds buoyancy, but more importantly, acts as a protective insulator to limit transfer of heat to the water (Vogel 1990; Köhler 1991). Some loss of this air layer is caused by rapid movements of the appendages while diving, which when combined with increased convective heat loss results in increased peripheral cooling. Rates of cooling calculated for water shrews voluntarily swimming and diving over the first 5 min of each trial ($0.92^\circ\text{C min}^{-1}$) in 10°C water were substantially lower than those previously measured by Calder (1969) for *S. palustris* at the same temperature: $2.06^\circ\text{C min}^{-1}$ (swimming) and $2.46^\circ\text{C min}^{-1}$ (diving). This difference may, in part, relate to our efforts to

maintain the pelage of the shrews in optimal condition thereby preserving the integrity of the critical air layer trapped in the fur.

The adaptive hypothermia hypothesis (Butler and Jones 1997) has been advanced to explain why some divers routinely exceed their cADL. However, the benefit of increased aerobic dive endurance comes with the mandatory energetic cost of re-warming cooled tissues after diving and, for small-bodied divers especially, the potential impairment of locomotor function and foraging efficiency accompanying immersion hypothermia. Studies examining the adaptive hypothermia tenet have yielded mixed results (Ponganis et al. 2003; Hindle et al. 2006; Niizuma et al. 2007). Owing to their low O₂ stores, high mass-specific DMR, and rapid rate of heat loss in water, water shrews might be expected to implement body cooling to prolong their underwater endurance. Consistent with this suggestion, water shrews in dive trials at 3 and 10°C, and those freely foraging in 3°C water in the semi-natural environment, showed a statistically significant inverse relationship between dive duration and core T_b. However, adaptive hypothermia should manifest as a greater proportion of dive times above the cADL in animals exhibiting the lowest core temperatures. No significant correlation was found between T_b and dive times that exceeded the cADL in either the acute or 24-hr dive trials. More telling was the finding that dive frequency significantly declined in concert with T_b, with few dives occurring at T_{b's} <37.5°C. This observation exemplifies the strict thermoregulatory control *S. palustris* exhibits during aquatic activity, and further suggests that free ranging water shrews do not utilize hypothermia to extend underwater foraging endurance.

In contrast, water shrews required to exclusively feed underwater during the 12- to 28-hr trials in 3°C water were observed to consistently elevate core body temperature by ~1.0–1.5°C immediately prior to initiating each aquatic foraging bout. This increase was often apparent

while the shrews were quietly perched near the water's edge, and thus was likely fueled by a combination of shivering and non-shivering thermogenesis. The latter mode of heat production may be especially critical during aquatic foraging bouts, and we have in fact observed large interscapular brown fat deposits in this species (Gusztak, MacArthur, and Campbell, unpublished observations). Notably, free-ranging muskrat and Eurasian otter (*Lutra lutra*) similarly increase T_b immediately before initiating foraging bouts in cold water (MacArthur 1979; Kruuk et al. 1987), suggesting this phenomenon may be relatively wide-spread among semi-aquatic mammals. However, owing to the low thermal inertia of water shrews (and the consumption of relatively large cold prey items), and potentially the release of peripheral vasoconstriction, T_b often declined by 1°C following each dive. This period was accompanied by vigorous grooming and burrowing in the moss of the chamber such that T_b generally increased back to 38.5–39.0°C over the ensuing 2 to 4 min before initiating the next aquatic excursion, as was similarly observed in the 20-min behavioral dive trials. Limited by already short foraging durations, minor increases in dive time arising from hypothermia are thus presumably outweighed by the heightened muscle power output afforded by a high T_b which likely enhances the already impressive reaction times (20 msec; Catania et al. 2008), and hence foraging efficiency of this diminutive predator while submerged. As suggested for free-ranging mink (Harrington et al. 2012), this advantage would be amplified in the physiologically challenging winter season whereby Q_{10} driven reductions in prey metabolism presumably allows these agile warm-blooded predators to capture prey items within the short submergence window (~10 sec) afforded by their inherently low cADLs. Interestingly, the relatively short (~65 min) interbout intervals closely align with the mean digesta throughput time of this species (50–55 min; Gusztak et al. 2005), thereby facilitating a dozen or more foraging bouts per day.

750

751 *Summary*

752 American water shrews are particularly intriguing from a physiological perspective as allometry
753 predicts this species has the smallest total body oxygen storage capacity, highest diving
754 metabolic rate, lowest skeletal muscle buffering capacity, and lowest anaerobic capability of any
755 endothermic diver. The water shrew must additionally balance the continual threat of immersion
756 hypothermia while submerged with the pressure to maximize its aerobic dive endurance and
757 anaerobic capacity in order to ensure adequate underwater foraging times. This challenge is
758 presumably greatest in the winter when water shrews must detect and capture prey in ice-cold
759 water, often in total darkness. Surprisingly, the world's smallest mammalian diver is able to
760 achieve this feat at least in part by elevating their core T_b by $>1^{\circ}\text{C}$ prior to diving. Despite
761 increasing the rate of O_2 use while underwater, this pre-dive hyperthermia likely enhances their
762 impressive reaction times and set of sensory modalities to efficiently capture energy-rich aquatic
763 prey within time windows as short as 5 to 10 sec. It may also enhance winter foraging efficiency
764 not only by enabling faster underwater swim speeds, but by permitting longer foraging bouts in
765 near-freezing water. This pre-dive response may thus provide a critical margin of safety that
766 reduces the risk of costly and potentially debilitating hypothermia.

767

768 **Acknowledgments**

769 We thank Chris Schneider, Dean Jeske, Allyson Hindle, and Rob Senkiw for assistance with
770 shrew trapping and husbandry, and Shane Farrow and Stefan Gusztak for lab assistance. R.W.G.
771 was supported in part by a University of Manitoba Faculty of Science Undergraduate Student
772 Research Award, a NSERC Undergraduate Research Award, and a NSERC Postgraduate

Scholarship. This research was funded by Natural Sciences and Engineering Research Council (NSERC) Discovery Grants to R.A.M. and K.L.C.

Literature Cited

Abe H. 2000. Review: role of histidine-related compounds as intracellular proton buffering constituents in vertebrate muscle. *Biochem (Moscow)* 65:757–765.

Bartels H., R. Bartels, R. Baumann, R. Fons, and K.D. Jürgens. 1979. Blood oxygen transport and organ weights of two shrew species *Suncus etruscus* and *Crocidura russula*. *Am J Physiol* 236:R221–R224.

Bartholomew G.A., D. Vleck, and C.M. Vleck. 1981. Instantaneous measurements of oxygen consumption during pre-flight warm-up and post-flight cooling in sphingid and saturniid moths. *J Exp Biol* 90:17–32.

Beneski Jr J.T. and D.W. Stinson 1987. Mammalian species: *Sorex palustris*. *Am Soc Mammal* 296:1–6.

Bethge P., S. Munks, H. Otley, and S. Nicol. 2003. Diving behaviour, dive cycles and aerobic dive limit in the platypus, *Ornithorhynchus anatinus*. *Comp Biochem Physiol A* 136:799–809.

Brander R.B. and W.W. Cochran. 1967. Radio-location telemetry. In R.G. Giles (ed.), *Wildlife Management Techniques*. The Wildlife Society, Washington, D.C.

Butler P.J. and D.R. Jones. 1997. Physiology of diving birds and mammals. *Physiol Revs* 77:837–899.

Calder W.A. 1967. Physiology of diving and breath-holding in small birds and mammals. The Jackson Hole Research Station Annual Report. 1967:5–7.

- 796 Calder W.A. 1969. Temperature relations and under water endurance of the smallest
 797 homeothermic diver, the water shrew. *Comp Biochem Physiol A* 30:1075–1082.
- 798 Calder W.A. 1996. *Size, Function, and Life History*. Courier Corporation.
- 799 Castellini M.A., G.N. Somero, and G.L. Kooyman. 1981. Glycolytic enzyme activities in tissues
 800 of marine and terrestrial mammals. *Physiol Zool* 54:242–252.
- 801 Catania K.C. 2006. Olfaction: underwater ‘sniffing’ by semi-aquatic mammals. *Nature*
 802 444:1024–1025.
- 803 Catania K.C. 2013. The neurobiology and behavior of the American water shrew (*Sorex*
 804 *palustris*). *J Comp Physiol A* 199: 545–554.
- 805 Catania K.C., J.F. Hare, and K.L. Campbell. 2008. Water shrews detect movement, shape, and
 806 smell to find prey underwater. *Proc Natl Acad Sci* 105:571–576.
- 807 Churchfield S. 1985. The feeding ecology of the European water shrew. *Mammal Rev* 1:13–21.
- 808 Conaway C.H. 1952. Life history of the water shrew (*Sorex palustris navigator*). *Amer Midland*
 809 *Nat* 48:219–248.
- 810 Costa D.P., N.J. Gales, and M.E. Goebel. 2001. Aerobic dive limit: how often does it occur in
 811 nature? *Comp Biochem Physiol A* 129:771–783.
- 812 Dawson T.J. and D. Fanning. 1981. Thermal and energetic problems of semiaquatic mammals: a
 813 study of the Australian water rat, including comparisons with the platypus. *Physiol Zool*
 814 54:285–296.
- 815 Doremus H.M. 1965. Heart rate, temperature and respiration rate of the short-tailed shrew in
 816 captivity. *J Mammal* 46:424–425.

- 817 de Leeuw J.J., M.R. Erden, and G.H. Visser. 1999. Wintering tufted ducks, *Aythya fuligula*,
818 diving for zebra mussels, *Dreissena polymorpha*, balance feeding costs within narrow
819 margins of their budget. J Avian Biol 30:182–192.
- 820 Dunstone N. and R.J. O’Conner. 1979. Optimal foraging in an amphibious mammal. I. The aqua-
821 lung effect. Anim Behav 27:1182–1194.
- 822 Dyck A.P. and R.A. MacArthur. 1992. Seasonal body temperature and activity in free-ranging
823 beaver (*Castor canadensis*). Can J Zool 70:1668–1672.
- 824 Elliott K.H., A. Shoji, K.L. Campbell, and A.J. Gaston. 2010. Oxygen stores and foraging
825 behavior of two sympatric, planktivorous auks. Aquatic Biol 8:221–235.
- 826 Emmett B. and P.W. Hochachka. 1981. Scaling of oxidative and glycolytic enzymes in
827 mammals. Respir Physiol 45:261–72.
- 828 Evans B.K., D.R. Jones, J. Baldwin, and G.R.J. Gabbott. 1994. Diving ability of the platypus.
829 Aust J Zool 42:17–27.
- 830 Fish F.E. 2000. Biomechanics and energetics in aquatic and semiaquatic mammals: platypus to
831 whale. Physiol Biochem Zool 73:683–698.
- 832 Gauthier G. 1993. Bufflehead (*Bucephala albeola*). In *Birds of North America*, No. 67. A. Poole
833 and F. Gill, eds. The Academy of Natural Sciences, Philadelphia, PA and The American
834 Ornithologists’ Union, Washington, D.C.
- 835 Gehr P., D.K. Mwangi, A. Amman, G.M.O. Maloiy, C.R. Taylor, and E.R. Weibel. 1981. Design
836 of the mammalian respiratory system. V. Scaling morphometric pulmonary diffusion capacity
837 to body mass: wild and domestic mammals. Respir Physiol 44:61–86.
- 838 George S.B., J.R. Choate, and H.H. Genoways 1986. Mammalian species: *Blarina brevicauda*.
839 Am Soc Mammal 261:1–9.

- 840 Graf P.M., R.P. Wilson, L.C. Sanchez, K. Hackländer, and F. Rosell. 2018. Diving behavior in a
 841 free-living, semi-aquatic herbivore, the Eurasian beaver *Castor fiber*. *Ecol Evol* 8:997–1008.
- 842 Gusztak R.W. and K.L. Campbell. 2004. Growth, development and maintenance of American
 843 water shrews (*Sorex palustris*) in captivity. *Mammal Study* 29:65-72.
- 844 Gusztak R.W., R.A. MacArthur, and K.L. Campbell. 2005. Bioenergetics and thermal
 845 physiology of American water shrews (*Sorex palustris*). *J Comp Physiol B* 175:87–95.
- 846 Halsey L.G., P.J. Butler, and T.M. Blackburn. 2006. A phylogenetic analysis of the allometry of
 847 diving. *Am Nat* 167:276–287.
- 848 Hamilton W.J. 1930. The food of the Soricidae. *J Mammal* 11:26–39.
- 849 Harrington L.A. G.C. Hays, L. Fasola, A.L. Harrington, D. Righton, and D.W. Macdonald. 2012.
 850 Dive performance in a small-bodied, semi-aquatic mammal in the wild. *J Mammal* 93:198–
 851 210.
- 852 He K., T.G. Eastman, H. Czolacz, S. Li, A. Shinohara, S.-I. Kawada, M.S. Springer, M.
 853 Berenbrink, and K.L. Campbell. 2021. Myoglobin primary structure reveals multiple
 854 convergent transitions to semi-aquatic life in the world's smallest mammalian divers. *eLife*
 855 10:e66797.
- 856 Hindle A.G., R.W. Senkiw, and R.A. MacArthur. 2006. Body cooling and the diving capabilities
 857 of muskrats (*Ondatra zibethicus*): A test of the adaptive hypothermia hypothesis. *Comp*
 858 *Biochem Physiol A* 144:232–241.
- 859 Hindle A.G, J.M. Lawler, K.L. Campbell, and M. Horning. 2009. Muscle senescence in short-
 860 lived wild mammals, the soricine shrews *Blarina brevicauda* and *Sorex palustris*. *J Exp Zool*
 861 *A Ecol Genet Physiol* 311A:358–367.
- 862 Hochachka P.W. and T.P. Mommsen. 1983. Protons and anaerobiosis. *Science* 4591:1391–1397.

- 863 Jordaan R.K., M.J. Somers, and T. McIntyre. 2021. The diving behavior of African clawless and
 864 spotted-necked otters in freshwater environments. *J Mammal* gyab031.
- 865 Jürgens K.D. 2002. Etruscan shrew muscle: the consequences of being small. *J Exp Biol*
 866 205:2161–2166.
- 867 Kleiber M. 1975. *The Fire of Life: An Introduction to Animal Energetics*. Huntington, NY: R.E.
 868 Krieger.
- 869 Köhler D. 1991. Notes on the diving behaviour of the water shrew, *Neomys fodiens* (Mammalia,
 870 Soricidae). *Zool Anz* 227:218–228.
- 871 Kooyman G.L., E.A. Wahrenbrock, M.A. Castellini, R.W. Davis, and E.E. Sinnett. 1980.
 872 Aerobic and anaerobic metabolism during voluntary diving in Weddell seals: evidence of
 873 preferred pathways from blood chemistry and behavior. *J Comp Physiol* 138:335–346.
- 874 Kooyman G.L., M.A. Castellini, R.W. Davis, and R.A. Maue. 1983. Aerobic dive limits in
 875 immature Weddell seals. *J Comp Physiol A* 151:171–174.
- 876 Kooyman G.L. and P.J. Ponganis. 1998. The physiological basis of diving to depth: birds and
 877 mammals. *Annu Rev Physiol* 60:19–32.
- 878 Kruuk H., P.T. Taylor, and G.A.T. Mom. 1997. Body temperature and foraging behaviour of the
 879 Eurasian otter (*Lutra lutra*), in relation to water temperature. *J Zool Lond* 241:689–697.
- 880 Levy A. 1964. The accuracy of the bubble meter method for gas flow measurements. *Sci*
 881 *Instrument* 41:449–453.
- 882 Lenfant C., R. Elsner, G.L. Kooyman, and C.M. Drabek. 1970. Gas transport and oxygen storage
 883 capacity in some pinnipeds and the sea otter. *Respir Physiol* 9:277–286.
- 884 MacArthur R.A. 1979. Seasonal patterns of body temperature and activity in free-ranging
 885 muskrats (*Ondatra zibethicus*). *Can J Zool* 57:25–33.

- 886 MacArthur R.A. 1984a. Aquatic thermoregulation in the muskrat (*Ondatra zibethicus*): energy
887 demands of swimming and diving. Can J Zool 62:241–248.
- 888 MacArthur R.A. 1984b. Seasonal changes in hematological and respiratory properties of muskrat
889 (*Ondatra zibethicus*) blood. Can J Zool 62:241–248.
- 890 MacArthur R.A. 1989. Aquatic mammals in cold. In *Advances in Comparative and*
891 *Environmental Physiology*, vol. 4, *Animal Adaptations to Cold* (ed. L.C.H. Wang) pp. 289–
892 325. New York: Springer-Verlag.
- 893 MacArthur R.A. and R.E. Krause. 1989. Energy requirements of freely diving muskrats
894 (*Ondatra zibethicus*). Can J Zool 67:2194–2200.
- 895 MacArthur R.A. 1992. Gas bubble release by muskrats diving under ice lost gas or a potential
896 oxygen pool. J Zool Lond 226:151–164.
- 897 MacArthur R.A., M.M. Humphries, G.A. Fines, and K.L. Campbell. 2001. Body oxygen stores,
898 aerobic dive limits, and the diving abilities of adult and juvenile muskrats (*Ondatra*
899 *zibethicus*). Physiol Biochem Zool 74:178–190.
- 900 McIntyre I.W. 2000. Diving energetics and temperature regulation of the star-nosed mole
901 (*Condylura cristata*) with comparisons to non-aquatic talpids and the water shrew (*Sorex*
902 *palustris*). MSc Thesis, University of Manitoba, Winnipeg.
- 903 McIntyre I.W., K.L. Campbell, and R.A. MacArthur. 2002. Body oxygen stores, aerobic dive
904 limits and diving behaviour of the star-nosed mole (*Condylura cristata*) and comparisons
905 with non-aquatic talpids. J Exp Biol 205:45–54.
- 906 McKean T. and C. Carlton. 1977. Oxygen storage in beavers. J Appl Physiol 42:545–547.

- 907 Mirceta S., A.V. Signore, J.M. Burns, A.R. Cossins, K.L. Campbell, and M. Berenbrink. 2013.
 908 Evolution of mammalian diving capacity traced by myoglobin net surface charge. *Science*
 909 340:1234192.
- 910 Nakagawa Y. and M. Hattori. 2002. Relationship between muscle buffering capacity and fiber
 911 type during anaerobic exercise in human. *J Physiol Anthropol* 21:129–131.
- 912 Niizuma Y., G.W. Gabrielsen, K. Sato, Y. Watanuki, and Y. Naito. 2007) Brünnich's guillemots
 913 (*Uria lomvia*) maintain high temperature in the body core during dives. *Comp Biochem*
 914 *Physiol A* 147:438–444.
- 915 Ordway G.A. and D.J. Garry. 2004. Myoglobin: an essential hemoprotein in striated muscle. *J*
 916 *Exp Biol* 207:3441–3446.
- 917 Ponganis P.J., L.N. Starke, M. Horning, and G.L. Kooyman. 1999. Development of diving
 918 capacity in emperor penguins. *J Exp Biol* 202:781–786.
- 919 Ponganis P.J., R.P. Van Dam, D.H. Levenson, R. Knowler, K.V. Ponganis, and G. Marshall.
 920 2003. Regional heterothermy and conservation of core temperature in emperor penguins
 921 diving under sea ice. *Comp Biochem Physiol A* 135:477–487.
- 922 Prothero J.W. 1980. Scaling of blood parameters in mammals. *Comp Biochem Physiol A*
 923 67:649–657.
- 924 Quick N.J., W.R. Cioffi, J.M. Shearer, A. Fahlman, and A.J. Read. 2020. Extreme diving in
 925 mammals: first estimates of behavioural aerobic dive limits in Cuvier's beaked whales. *J Exp*
 926 *Biol* 223: jeb222109.
- 927 Reynafarje B. 1963. Simplified method for the determination of myoglobin. *J Lab Clin Med*
 928 61:138–145.

- 929 Ruthardt M. and R. Schröpfer. 1985. Zum Verhalten der wasserspitzmaus *Neomys fodiens*
 930 (Pennant, 1771) unter wasser. Z Angew Zool 72:49–57.
- 931 Samuel P.P., L.P. Smith, G.N. Phillips Jr., and J.S. Olson. 2015. Apoglobin stability is the major
 932 factor governing both cell-free and in vivo expression of holomyoglobin. J Biol Chem
 933 290:23479–23495.
- 934 Scholetch R. 1980. Freilandbeobachtungen an der waserspitzmaus, *Neomys fodiens* (Pennant,
 935 1771) im Schweizerischen Nationalpark. Revue Suisse Zool 87:937–939.
- 936 Schreer J.F. and K.M. Kovacs. 1997. Allometry of diving capacity in air-breathing vertebrates.
 937 Can J Zool 75:339–358.
- 938 Svihla, A. 1934. The mountain water shrew. Murrelet 15:44–45.
- 939 Sorenson M.W. 1962. Some aspects of water shrew behavior. Am Midl Nat 68:445–462.
- 940 Stewart J.M., A.K. Woods, and J.A. Blakely. 2005. Maximal enzyme activities, and myoglobin
 941 and glutathione concentrations in heart, liver and skeletal muscle of the Northern short-tailed
 942 shrew (*Blarina brevicauda*; Insectivora: Soricidae). Comp Biochem Physiol B 141:267–273.
- 943 Stahl W.R. 1967. Scaling of respiratory variables in mammals. J Appl Physiol 22:453–460.
- 944 Vogel P. 1990. Body temperature and fur quality in swimming water-shrews, *Neomys fodiens*
 945 (Mammalia, Insectivora). Z Saeugetierkd 55:73–80.
- 946 Vogel P. 1998. Diving capacity and foraging behaviour of the water shrew (*Neomys fodiens*).
 947 Symp Zool Soc Lond 71:31–47.
- 948 Vornanen M.A. 1992. Maximum heart rate of soricine shrews: correlation with contractile
 949 properties and myosin composition. Am J Physiol Reg Integr Comp Physiol 262:R842–R851.
- 950 Weibel E.R. 1970/71. Morphometric estimation of pulmonary diffusing capacity. I. Model and
 951 method. Resp Physiol 11:54–75.

- 952 Weibel E.R. 1985. Design and performance of muscular systems: an overview. J Exp Biol
953 115:405–412.
- 954 White C.R. and R.S. Seymour. 2003. Mammalian basal metabolic rate is proportional to body
955 mass^{2/3}. Proc Natl Acad Sci 100:4046–4049.
- 956 Whittaker J.C., G. Hammerson, and S.J. Norris. 2008. *Sorex palustris*. In: IUCN 2009. IUCN
957 Red list of threatened species. Version 2009.2. www.iucnredlist.org.
- 958 Wolk E. 1974. Variations in hematological parameters of shrews. Acta Theriol 19:315–346.
- 959 Yeates L.C., T.M. Williams, and T.L. Fink. 2007. Diving and foraging energetics of the smallest
960 marine mammal, the sea otter (*Enhydra lutris*). J Exp Biol 210:1960–1970.
- 961 Zar J.H. 1974. Biostatistical analysis. Prentice-Hall. Englewood Cliff, NJ.

Table 1. The effect of water temperature on the voluntary dive behavior of 18 captive American water shrews.

Variable	Water temperature (°C)			
	3	10	20	30
Total time in water (sec)	111±14 ^a	134±15 ^a	184±21 ^{a,b}	235±25 ^b
All dives (<i>n</i>)	258	257	279	303
Dive duration (sec)	4.0±0.2 ^a	4.9±0.2 ^b	5.2±0.2 ^b	5.5±0.2 ^b
Dive frequency (dives • min ⁻¹)	0.71±0.08 ^a	0.71±0.09 ^a	0.77±0.07 ^a	0.84±0.11 ^a
Dive:surface ratio	0.12±0.02 ^a	0.20±0.02 ^{a,b}	0.25±0.09 ^b	0.27±0.03 ^b
Duration of five longest dives (sec)	6.4±0.5 ^a	7.6±0.5 ^{a,b}	8.5±0.5 ^{b,c}	9.3±0.6 ^c
Duration of longest dive (sec)	8.1±0.7 ^a	9.7±0.6 ^{a,b}	11.3±0.8 ^{b,c}	12.4±0.9 ^c

Values presented are means ± 1 SE.

Within each row, means sharing the same letter are not statistically different ($P>0.05$).

Table 2. Telemetered body temperatures (T_b) recorded from six American water shrews during 20-min voluntary dive trials in 3–30°C water.

Variable	Water temperature (°C)			
	3	10	20	30
Time in water (sec)	106.9±22.4 ^a	152.5±34.0 ^a	180.4±29.4 ^a	162.1±18.7 ^a
Rate of cooling during first 5 min of trial (°C min ⁻¹)	1.61±0.63 ^a	0.92±0.24 ^a	0.99±0.31 ^a	0.43±0.15 ^a
T_b measurements (°C)				
Pre-trial T_b	40.0±0.2 ^{a,1}	40.1±0.1 ^{a,1}	39.9±0.1 ^{a,1}	40.1±0.2 ^{a,1}
Mean T_b during diving	38.9±0.3 ^{a,2,3}	39.1±0.2 ^{a,2}	38.7±0.4 ^{a,2,3}	39.4±0.2 ^{a,2}
Lowest T_b in trial	38.0±0.3 ^{a,3}	38.4±0.2 ^{a,3}	37.7±0.5 ^{a,3}	38.8±0.2 ^{a,3}
Highest post-trial T_b	39.5±0.2 ^{a,1,2}	39.8±0.2 ^{a,1}	39.8±0.2 ^{a,1,2}	39.5±0.1 ^{a,1,2}

Values presented are means ± 1 SE.

Values within a row sharing the same letter do not differ significantly ($P>0.05$).

Values within a column sharing the same number do not differ significantly ($P>0.05$).

Table 3. Respiratory characteristics of short-tailed and American water shrews. Values are presented as means $\pm$ 1 SE, with number of animals sampled indicated in parentheses.

Variable	American water shrew		Short-tailed shrew	
	Juvenile	Adult	Adult	Juvenile
Body mass (g)	13.00 $\pm$ 0.45 (20)	14.57 $\pm$ 0.36 (21) ^a	24.26 $\pm$ 1.41 (10) ^b	21.78 $\pm$ 0.13 (7)
% total body muscle mass	36.72 $\pm$ 2.34 (2)	40.51 $\pm$ 3.38 (9)	24.13 $\pm$ 1.80 (2) ^b	No data
Myoglobin concentration (mg g wet tissue ⁻¹)				
Forelimb	3.75 $\pm$ 0.22 (10)	6.18 $\pm$ 0.31 (10) ^a	2.98 $\pm$ 0.32 (7) ^b	No data
Hindlimb	3.76 $\pm$ 0.16 (10)	5.89 $\pm$ 0.45 (9) ^a	3.01 $\pm$ 0.40 (6) ^b	No data
Mean skeletal muscle concentration ^x	3.76 $\pm$ 0.12 (10)	6.04 $\pm$ 0.25 (9) ^a	3.03 $\pm$ 0.26 (9) ^b	2.70 $\pm$ 0.49 (7)
Heart	10.97 $\pm$ 1.22 (5)	9.47 $\pm$ 0.63 (6)	8.77 $\pm$ 0.55 (5)	8.77 $\pm$ 0.79 (5)
Buffering capacity (slykes) ^c	30.67 $\pm$ 2.34 (11)	38.22 $\pm$ 2.28 (13) ^a	24.88 $\pm$ 1.40 (10) ^b	25.05 $\pm$ 1.92 (6)
Total lung volume (mL STPD 100 g ⁻¹)	4.55 $\pm$ 0.18 (9)	4.57 $\pm$ 0.28 (5)	3.34 $\pm$ 0.07 (6) ^b	3.43 $\pm$ 0.11 (5)
Hematocrit (%)	50.57 $\pm$ 1.33 (10)	50.23 $\pm$ 1.33 (15)	48.78 $\pm$ 2.50 (8)	50.17 $\pm$ 2.50 (7)
Hemoglobin content (g 100 mL ⁻¹)	19.69 $\pm$ 1.02 (8)	20.09 $\pm$ 0.44 (13)	17.29 $\pm$ 0.62 (7) ^b	16.87 $\pm$ 1.54 (6)
Blood O ₂ capacity (vol. %)	26.38 $\pm$ 1.37 (8)	26.93 $\pm$ 0.58 (13)	23.18 $\pm$ 0.84 (7) ^b	22.61 $\pm$ 0.14 (6)

^aValues for adult and juvenile water shrews are significantly different (P<0.05).

^bValues for adult water shrews and adult short-tailed shrews are significantly different (P<0.05).

^cμmoles of base required to titrate the pH of 1 g of wet muscle by 1 pH unit.

Table 4. Oxygen storage capacities of the lungs, blood, and skeletal muscle of adult and juvenile American water shrews (*Sorex palustris*) and adult short-tailed shrews (*Blarina brevicauda*).

Species	N	Oxygen stores (mL O ₂ STPD kg ⁻¹)					Calculated ADL (sec) ^a	
		Lung	Arterial blood	Venous blood	Muscle	Total	Water temperature (°C) 10	30
Adult <i>S. palustris</i>	8	6.60±0.08	6.33±0.04	10.12±0.07	3.26±0.21	26.31±0.24	10.8	14.4
Juvenile <i>S. palustris</i>	4	6.29±0.22	6.15±0.06	9.77±0.11	1.84±0.09	24.06±0.16	9.9	13.2
Adult <i>B. brevicauda</i>	2	5.01 ^b	5.45±0.37	8.37±0.75	1.32±0.04	20.14±1.07	N/A	N/A

^aThe aerobic dive limit (ADL) was calculated for a 14.0 g water shrew assuming a diving metabolic rate of 8.77 mL O₂ g⁻¹ hr⁻¹ in 10°C water and 6.57 mL O₂ g⁻¹ hr⁻¹ in 30°C water (see text for details).

^bValue estimated from data in Table 3.

Supplemental Table 1. Contingency table using the method of log-likelihood ratio to compare the frequency distribution of voluntary dives by twelve adult American water shrews with ($N=6$) and without ($N=6$) an implanted abdominal temperature transmitter.

Variable	Dive duration (sec)						Total
	0-2	3-5	6-8	9-11	12-14	>15	
Non-implanted water shrews	109	96	69	29	13	3	319
Implanted water shrews	92	104	75	36	10	3	320
Total	201	200	144	65	23	6	639

df=5, critical value=11.07.

Calculated G-value=2.78, $P>0.50$.

Supplemental Table 2. Comparison of the dive performance of adult American water shrews with and without a surgically implanted abdominal temperature transmitter. Dive data for each animal are pooled values from 20-min dive trials completed in 3, 10, 20, and 30°C water.

	Mass (g) (without transmitter)	Mean dive duration (sec)	Mean dive frequency (dives min ⁻¹)	Mean deep dive duration (sec)	Mean deep dive frequency (dives min ⁻¹)	Longest dive (sec)	Dive:surface ratio	Total time in water per trial (sec)
Non-implanted								
Shrew 1	16.3	5.1	0.39	8.8	0.05	12.2	0.21	132.0
Shrew 2	13.6	5.8	0.48	13.9	0.04	23.7	0.21	160.0
Shrew 5	14.0	6.4	0.85	8.5	0.41	13.2	0.26	170.5
Shrew 8	17.8	4.2	0.76	7.8	0.08	13.9	0.13	193.0
Shrew 9 ^a	13.3	5.3	0.94	8.5	0.20	11.7	No data	163.9
Shrew 10	15.6	4.1	0.53	7.4	0.10	12.6	0.31	83.9
Mean	15.1±0.8	5.2±0.4	0.66±0.10	9.2±1.1	0.15±0.06	14.6±2.0	0.22±0.03	150.5±17.0
Transmitter implanted								
Shrew 12	15.1	4.3	0.40	8.1	0.14	16.7	0.10	108.2
Shrew 13	14.1	5.9	0.85	7.8	0.13	14.8	0.11	198.0
Shrew 14	15.2	4.3	0.70	5.9	0.14	9.8	0.21	151.0
Shrew 15	14.0	6.3	1.00	8.8	0.29	20.1	0.33	185.8
Shrew 16	14.0	4.1	0.44	6.1	0.05	10.2	0.19	80.7
Shrew 17	14.2	7.0	0.58	9.7	0.14	14.5	0.22	167.8
Mean	14.4±0.3	5.3±0.6	0.66±0.11	7.7±0.7	0.15±0.03	14.4±1.8	0.19±0.04	148.6±20.4
<i>P</i> -value ^b	0.199	0.397	0.490	0.124	0.487	0.468	0.263	0.468

^aDive data taken from McIntyre 2000; note that dive:surface ratio was not reported for this animal.

^bCalculated using 1-tailed Student's *t*-test.

Figure Legends

Figure 1. Influence of water temperature on the diving behavior of 18 American water shrews in 30, 20, 10, and 3°C water. Shallow dives (<10 cm deep) are denoted by open bars, while deep dives (60 cm) are denoted by stippled bars. Behavioral (bADL) and calculated aerobic dive limits (cADL) are denoted by dotted and dashed lines, respectively, while solid circles indicate mean dive time at each water temperature.

Figure 2. Telemetered body temperatures (T_b) $\pm$ 1 SE of six American water shrews voluntarily diving during 20-min trials in 3–30°C water. Pre- and post-trial measurements were taken from shrews in temporary holding containers immediately before and after the trials (see text for details).

Figure 3. The distribution of dive times for three American shrews diving in a semi-natural riparian environment in 3°C water over 12- to 28-hr periods. The calculated behavioral aerobic dive limit (bADL) of 10.7 sec is indicated by the dotted line.

Figure. 4. Mean body temperature $\pm$ 1 SE of five American water shrews in the 35 min prior to initiating foraging bouts in a semi-natural riparian environment.

Figure 5. Representative body temperature (dashed lines) and relative activity traces (grey bars) of two adult American water shrews during 12- to 28-hr trials in a natural riparian environment (water temperature=3°C). Open boxes represent body temperature at the start of each water excursion, while red circles denote body temperature immediately following exit from the water.

Figure 6. The relationship of diving metabolic rate to total submergence time of American water shrews voluntarily diving in 10°C (open circles, dash line) and 30°C (closed squares, solid line). The dotted line denotes the basal metabolic rate (BMR; 4.84 mL O₂ g⁻¹ hr⁻¹) of water shrews (Gusztak et al. 2005).

Figure 7. Pooled behavioral and body temperature (T_b) data for American water shrews implanted with a 1.0-g intra-abdominal transmitter and voluntarily diving in a) 3°C and 10°C water and b) during the 12- to 28-hr trials in 3°C water. Although the relationship between dive duration and core T_b is significant in both instances, no significant relationship was observed when the analysis was limited dives greater than the calculated aerobic dive limit (~10 sec; denoted by filled circles). The frequency distribution of T_b recordings in c) 3°C and 10°C water and d) during the 12- to 28-hr trials in 3°C water are also presented.

Supplemental Figure 1. Photograph of the semi-natural riparian set up for observing the voluntary aquatic behavior, body temperature, and activity of five adult water shrews over periods ranging from 12 to 28 hr.

Supplemental Figure 2. Percentage of a) total voluntary dives completed and b) corresponding mean dive times of 6 transmitter-implanted American water shrews during successive 5-min periods of each dive trial ($n=317$). Bars sharing the same letters are not significantly different ($P<0.05$).

Supplemental Figure 3. Frequency distribution of voluntary dive times of 25 American water shrews diving in 3–30°C water ($n=1584$ dives). Shallow dives (<10 cm deep) are denoted by open bars, while deep dives (60 cm) are denoted by stippled bars. The mean dive time of all dives combined was 5.1 ± 0.1 sec, while the mean duration of deep dives was 8.1 ± 0.2 sec ($n=311$).

Supplemental Figure 4. The relationship of inter-dive surface time to duration of preceding dive in water shrews completing consecutive dives during voluntary dive trials. The regression line was fitted by the method of least squares.

Supplemental Figure 5. Body temperature (dashed lines) and relative activity traces (grey bars) of three adult American water shrews during 12- to 28-hr trials in a natural riparian environment (water temperature=3°C). Open boxes represent body temperature at the start of each water excursion, while red circles denote body temperature immediately following exit from the water.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Shrew #13: 24-hr trial (average mass = 14.5 g)

Shrew #17: 12-hr trial (average mass = 13.3 g)

Figure 6

Figure 7

Supplemental figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental figure 4

Supplemental figure 5

Shrew #16: 28-hr trial (average mass = 16.2 g)

Shrew #14: 12-hr trial (average mass = 13.6 g)

Shrew #12: 24.5-hr trial (average mass = 17.2 g)