**The genome sequence of the avian vampire fly (*Philornis downsi*), an invasive nest
parasite of Darwin's finches in Galápagos**

Melia Romine¹, Sarah A. Knutie^{2,3}, Carly M. Crow⁴, Grace J. Vaziri², Jaime Chaves^{5,6},
Jennifer A.H. Koop^{4*} & Sangeet Lamichhaney^{1,7*#}

¹School of Biomedical Sciences, Kent State University, Kent, OH, USA

²Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT USA

³Institute for Systems Genomics, University of Connecticut, Storrs, CT USA

⁴Department of Biological Sciences, Northern Illinois University, DeKalb, IL, USA

⁵Department of Biology, San Francisco State University, San Francisco, CA, USA

⁶Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito,
Ecuador

⁷Department of Biological Sciences, Kent State University, Kent, OH, USA

*co-senior authors

#corresponding author (slamichh@kent.edu)

Abstract

The invasive avian vampire fly (*Philornis downsi*) is considered one of the greatest threats to the unique and endemic avifauna of the Galápagos Islands, Ecuador. The fly parasitizes nearly every passerine species, including Darwin's finches, in the Galápagos. The fly is thought to have been introduced from mainland Ecuador, although the full pathway of invasion is not yet known. The majority of research to date has focused on the effects of the fly on the fitness of avian hosts and explorations of mitigation methods. A lag in research related to the genetics of this invasion demonstrates, in part, a need to develop full-scale genomic resources with which to address further questions within this system. In this study, an adult *P. downsi* collected from San Cristobal Island within the Galápagos archipelago was sequenced to generate a high-quality genome assembly. We examined various features of the genome (e.g., coding regions, non-coding transposable elements) and carried out comparative genomics analysis against other dipteran genomes. We identified lists of gene families that are significantly expanding/contracting in *P. downsi* that are related to insecticide resistance, detoxification, and potential feeding ecology and counter defense against host immune responses. The *P. downsi* genome assembly provides an important foundational resource for studying the molecular basis of its successful invasion in the Galápagos and the dynamics of its population across multiple islands. The findings of significantly changing gene families associated with insecticide resistance and immune responses highlight the need for further investigations into the role of different gene families in aiding the fly's successful invasion. Furthermore, this genomic resource will also better help inform future research studies and mitigation strategies aimed at minimizing the fly's impact on the birds of the Galápagos.

Keywords: Avian vampire fly genome, parasitic invasion, Galápagos, insecticide resistance, Darwin's finches

Introduction

The invasive avian vampire fly (*Philornis downsi*) (Figure 1a) is considered among the greatest threats to the unique and endemic avifauna of the Galápagos Islands, Ecuador (Causton *et al.*, 2013). The fly parasitizes eleven species of Darwin's finches (Figure 1b) as well as nearly every other Galápagos passerine species (Causton *et al.*, 2013; Fessl *et al.*, 2018). The earliest record of adult flies from the Galápagos occurred in 1964, but the fly was not observed in the nests of birds until 1997 (Fessl and Tebbich, 2002; Causton *et al.*, 2006). The fly has now been found on all major islands within the archipelago except Española, Genovesa, Darwin, and Wolf (Causton *et al.*, 2013). The impact of parasitism by the fly has been severe in some populations of birds in Galápagos and while the effects are variable, some studies have reported near-total nest failure rates due to parasitism (Dudaniec *et al.*, 2007; Koop *et al.*, 2011; Koop, Le Bohec, *et al.*, 2013; O'Connor *et al.*, 2014; Knutie *et al.*, 2016a; Heimpel *et al.*, 2017; Adesso *et al.*, 2020). The fly has also been implicated in the decline of the medium tree finch (*Camarhynchus pauper*), the warbler finch (*Certhidia olivacea*), and the mangrove finch (*Camarhynchus heliobates*) (Dvorak *et al.*, 2004; Grant *et al.*, 2005; Cunninghame *et al.*, 2017; Peters and Kleindorfer, 2018; Bulgarella *et al.*, 2019). Furthermore, the potential for population-level declines of even relatively prominent bird species, e.g., the medium ground finch, have also been demonstrated using predictive models (Koop *et al.*, 2015).

The genus *Philornis* includes approximately 50 species found primarily in the Neotropics and into North America (Dodge, 1955, 1963; Spalding *et al.*, 2002; Dudaniec and Kleindorfer, 2006; Couri *et al.*, 2007). Of the described species within *Philornis*, only two species (including *P. downsi*) are free-living ectoparasites within the nests of their hosts (Fessl *et al.*, 2006), while most others are subcutaneous on their hosts. Several *Philornis* species are found in mainland

Ecuador (Bulgarella *et al.*, 2015, 2017), but *P. downsi* remains the only recorded species of the genus present, to date, in the Galápagos. *P. downsi* is thought to have been introduced from mainland Ecuador, though the full pathway of invasion is not yet known (Fessl *et al.*, 2018). Preliminary population genetics studies show that flies within the archipelago have a high degree of relatedness relative to those on the mainland, which supports the hypothesis of a relatively recent invasion and also the possibility of continued gene flow between populations in the Galápagos (Dudaniec *et al.*, 2008; Koop *et al.*, 2021).

A collaborative research effort has been made to continue monitoring the effects of the avian vampire fly on bird populations in the Galápagos. These efforts aim to identify source population(s), the pathway and mechanism of invasion, and possible long-term mitigation methods, including the Sterile Insect Technique and biocontrol (Causton *et al.*, 2013, 2019). Current stopgap mitigation efforts in the Galápagos include the direct application of insecticide to bird nests. Newly hatched nestlings are removed, and a permethrin solution is sprayed inside the nest and allowed to dry, at which point the nestlings are placed back in the nest. This method has been shown to effectively reduce parasites in the nest and increase fledging success (Koop *et al.*, 2011). A continuation of this method relies on birds incorporating permethrin-treated cotton into their nests, achieving similar increases in fledging success (Knutie *et al.*, 2014). Both methods, with subtle variations, are being used in attempts to increase the nesting success of the critically endangered mangrove finch, currently one of the most endangered birds in the world (Fessl *et al.*, 2018). The same method is also being used successfully to protect endangered bird species against their respective parasites in Australia (Alves *et al.*, 2020).

Despite the robust number of studies that explored the effects of these flies on hosts in the Galápagos, questions remain about the underlying ecological and evolutionary mechanisms of

their successful invasion in the Galápagos. This knowledge gap demonstrates the need to develop a full-scale genomic resource of the fly as it provides a critical knowledge base from which to explore these questions, similar to other parasites of concern (Scott *et al.*, 2020). In this study, we generated a high-quality draft genome of the avian vampire fly, which is expected to become an important resource of future molecular studies in this system. We further carried out comparative genomics analysis with additional published dipteran genomes that identified evidence of significantly changing gene families associated with insecticide resistance, detoxification, and possible counter defense against host immune responses. These results serve as the first step toward investigations of this fly's ability to rapidly evolve traits associated with its successful invasion in the Galápagos. From an applied aspect, these genomic resources will also help inform future research studies and mitigation strategies aimed at minimizing the fly's impact on the birds of the Galápagos.

Materials and Methods

Sampling and DNA extraction for genome sequencing

Flies were collected in Jardín de las Opuntias on San Cristobal Island, Galápagos, Ecuador (-0.9491651°, -89.5528454°) in March-April of 2019. Adult flies were reared from pupae collected in the nests of small ground finches (*Geospiza fuliginosa*). When the nests were empty because nestlings died or fledged, all larvae and pupae were extracted from the nest and placed in a falcon tube with a modified lid that allowed airflow. After flies eclosed from their pupal case, they were placed in the freezer, then preserved in 95% ethanol. Preserved flies were transported to the University of Connecticut, then shipped to Northern Illinois University for further processing. DNA was extracted using DNeasy Blood and Tissue Kit (Qiagen, Valencia,

California, USA) on whole fly samples after wings and legs were removed using forceps. All samples were treated with Monarch® RNase A (New England Biolabs) to remove RNA from genomic DNA samples. Multiple samples were extracted and those with low yields of DNA or contamination were discarded from further processing. Quantification of double-stranded DNA was done using QuBit® and samples were run on a gel to assess fragmentation.

Estimate of genome size and ploidy

We prepared a 10X Chromium GEM library from the extracted DNA above according to the manufacturer's recommended protocols. The library was sequenced using the Illumina 10X platform to generate paired-end 150 bp reads. We used a k-mer based approach to estimate genome size, heterozygosity, and repeat content from unprocessed short sequencing reads (Vurture *et al.*, 2017). We also used the k-mer distribution to extract heterozygous k-mer pairs to assess the ploidy level in *P. downsi* (Ranallo-Benavidez *et al.*, 2020). These estimates of genome size and ploidy were then used to choose parameters for the downstream genome assembly pipeline.

Genome assembly and annotation

The 10x Chromium linked reads generated above were used to generate a reference genome using Supernova.v.2.1.1 (Weisenfeld *et al.*, 2017) with default parameters. The genome contiguity statistics such as scaffold N₅₀ and the total number of scaffolds etc. were calculated using custom Perl scripts. We further compared the draft genome assembly against a set of conserved genes in Diptera using BUSCO (Waterhouse *et al.*, 2017) to assess gene-space completeness. We also mapped the short sequencing reads back to the draft genome assembly

using BWA (Li and Durbin, 2009) to estimate how many short-sequence reads were used to build the reference genome and infer genome completeness. The annotation of the *P. downsi* genome was done using the MAKER annotation pipeline (Cantarel *et al.*, 2008). We combined protein homology evidence (using publicly available protein datasets from major dipteran lineages, downloaded from the Ensemble database (Cunningham *et al.*, 2019) and *ab-initio* gene predictions models to annotate the genome.

Transposable elements (TE) in *P. downsi*

We combined homology-based and de novo approaches using Repeatmasker (Smit Hubley P., 2013; Flynn *et al.*, 2019) to characterize TEs in *P. downsi*. We used two different sets of repeat libraries, (a) reference repeat library downloaded from the Repbase database (Bao *et al.*, 2015) and (b) *de novo* custom-built species-specific repeat library generated for *P. downsi* using RepeatModeler (Flynn *et al.*, 2019). The usage of the *de novo* species-specific repeat library increased the accuracy of detection and annotation of transposable elements. For an unbiased comparison of repeat landscapes among all species, we used similar approaches to detect TEs in the Housefly (*M. domestica*), Stable fly (*S. calcitrans*), Tsetse fly (*G. morsitans*), Mediterranean fruit fly (*C. capitata*), Fruit fly (*D. melanogaster*) and Yellow fever mosquito (*A. aegypti*), together with an outgroup, Postman butterfly (*H. melpomene*).

Orthologs to other dipteran genomes

We inferred orthogroups and orthologs by comparative analysis of proteomes in *P. downsi* and six additional dipteran species (*M. domestica*, *S. calcitrans*, *G. morsitans*, *C. capitata*, *D. melanogaster*, and *A. aegypti*), together with an outgroup, *H. melpomene*. This

analysis used the draft proteome of *P. downsi* generated from our genome annotation pipeline described above. Complete proteomes of each of the remaining six species were downloaded from the Ensemble database (Cunningham *et al.*, 2019). The orthologs inference was carried out using Orthofinder (Emms and Kelly, 2015).

Gene family evolution

To explore the evolution of gene families in *P. downsi* to other Dipterans, we first constructed a maximum likelihood phylogeny using 3,070 single-copy orthologs among these eight species used for inferring orthologues (Figure 3a). We then used this phylogeny to analyze the changes in gene family size across lineages leading to each of these eight species using a maximum likelihood approach (CAFÉ) (De Bie *et al.*, 2006) that uses a birth and death process to model gene gain and loss across a phylogenetic tree.

Results

Genome sequencing and assembly

We sampled an adult *P. downsi* fly in Jardín de las Opuntias on San Cristobal Island, Galápagos. DNA library was prepared using 10X Chromium linked-read approach (Zheng *et al.*, 2016) and sequenced using Illumina 10X platform to generate paired-end 150 bp reads which resulted in ~479 million read pairs (~72 Gb of raw sequence data). We first generated a k-mer distribution based on these short-sequencing reads for preliminary characterization of genome structure in *P. downsi* (e.g., genome size, an abundance of repetitive elements, rate of heterozygosity) (Supplementary Table 1, Supplementary Figure 1) that further allowed us to make informed decisions on parameters needed for building a reference genome. We also

utilized the k-mer distribution to estimate the ploidy level in *P. downsi*, which indicated it to be a diploid species (Supplementary Figure 2).

Based on genome size estimates using k-mer distributions from short sequence reads (Supplementary Table 1), we had ~ 91X sequence coverage for generating the draft *de-novo* genome assembly. The total estimated length of the draft assembly was 971.6 Mb. The assembly contained 41,176 total scaffolds (minimum 1000 bp to maximum 8.6 Mb) with scaffold N₅₀ of 1.3 Mb. We further assessed genome contiguity and gene-space completeness using benchmarking universal single-copy orthologs (BUSCO) (Waterhouse *et al.*, 2017). Among 3,258 genes highly conserved across Diptera, 3,147 (95.8%) full-length genes were detected in the *P. downsi* draft genome assembly. Partial sequences of 71 genes were identified (2.2%) and only 67 (2.0%) were missing, indicating a high degree of contiguity in the *P. downsi* genome. In addition, 99.16% of total short sequencing reads aligned back to the draft genome indicating a high degree of genome completeness as well.

Genome annotation

We combined protein homology-based evidence and *ab-initio* gene prediction models using the MAKER genome annotation pipeline (Cantarel *et al.*, 2008) which identified 15,774 protein-coding genes in the *P. downsi* genome. We compared the genome annotation of *P. downsi* with two other published and highly curated dipteran genomes, *Musca domestica* (Scott *et al.*, 2014) and *D. melanogaster* (Adams *et al.*, 2000). We annotated fewer genes in the *P. downsi* genome (15,774) compared to that of *M. domestica* (17,283) or *D. melanogaster* (17,468) (Table 1). The average length of genes is similar among *P. downsi* and *D. melanogaster*, whereas

M. domestica typically has longer genes. The mean number of exons per gene is fewer in *P. downsi* compared to the other two species.

Table 1: Statistics of genomic features among three fly genomes

Genome statistics	<i>P. downsi</i>	<i>M. domestica</i> (Scott et al. 2014)	<i>D. melanogaster</i> (Adams et al. 2000)
Genome size (Gb)	0.97	0.69	0.18
Total genes	15,774	17,283	17,468
Avg. gene length (bp)	4,789	13,747	5,830
Total exons	56,595	123,831	188,405
Avg. exon length (bp)	367	453	485
Number of Exons per gene	3.59	7.16	10.79

Transposable elements in *P. downsi*

Mobile transposable elements (TE) are key features of eukaryotic genomes, being major determinants of genome size variation (Kapusta et al., 2017; Lamichhaney et al., 2021), and important contributors to the evolutionary potential of an organism (Pourrajab and Hekmatimoghaddam, 2021). We characterized the transposable elements in the *P. downsi* genome using homology-based (Smit Hubley P., 2013) and *de-novo* approaches (Flynn et al., 2019). More than half of the genome (51.7%) of *P. downsi* consists of transposable elements (Supplementary Table 2). Among these sequences, 9.3% of the genome are retroelements (7.7% LINEs and 1.6% LTRs) and 23.4% DNA transposons. Short interspersed nuclear elements (SINEs), a major category of retroelements, were not detected in the *P. downsi* genome.

We also used similar methods to detect, annotate, and compare the repeat content across several other species including the house fly (*Musca domestica*), stable fly (*Stomoxys calcitrans*), tsetse fly (*Glossina morsitans*), Mediterranean fruit fly (*Ceratitis capitata*), fruit fly (*Drosophila melanogaster*) and yellow fever mosquito (*Aedes aegypti*). The postman butterfly (*Heliconius melpomene*) was used as an outgroup. Transposable elements are known to be highly correlated

with genome size across the tree of life (Kidwell, 2002; Lynch, 2007), and our results across various dipteran genomes are consistent with this pattern (Figure 2a).

We also compared different categories of transposable elements across these seven genomes (Figure 2b). *P. downsi* had the highest proportion of DNA transposons (23.4%) among all taxa analyzed. In comparison, only 6.36% of the *S. calcitrans* genome and 15.96% of the *M. domestica* genome consisted of DNA transposons. Although *P. downsi* and *S. calcitrans* have similar genome sizes, *S. calcitrans* had slightly higher amounts of repeat content in its genome (58.3%), compared to *P. downsi* (51.7%) (Figure 2b). *S. calcitrans* had higher amounts of long interspersed nuclear elements (LINEs) (23.8%), compared to *P. downsi* (7.7%). We observed that most Diptera (except *Aedes*) have only a few SINE elements, one of the major classes of transposable elements (Figure 2b). LTR elements were relatively common in *D. melanogaster*, whereas other dipterans had a low amount of LTR elements (including *P. downsi*).

Orthologs to other dipteran genomes and outgroup *Heliconius*

Comparative genomics analysis of 113,047 protein sequences from *P. downsi* and seven other species (listed above) identified 11,112 orthogroups. A total of 95,567 proteins (out of 113,047, 86.3%) were assigned to these orthogroups. The mean size of an orthogroup was 8.8 genes/species, and 3,069 orthogroups had singly copy genes in each species. A total of 5,754 orthogroups were shared among all eight species. Only a few orthogroups (minimum 496 in *A. aegypti* and a maximum of 1,445 in *M. domestica*) were present in fewer than four species (Figure 3a). The number of unique orthologs in each species was consistent with their phylogenetic relationships (e.g., *A. aegypti* and the outgroup *H. melpomene* had the highest number of unique orthogroups). The number of shared orthologs among each pair of species is

presented in Supplementary Table 3. The number of orthologs identified in *P. downsi* is consistent with other published dipteran genomes (Figure 3a, Supplementary Table 3), indicating no major bias in the genome assembly and annotation pipeline used in this study.

Within *P. downsi*, 13,706 out of 15,774 (86.9%) annotated genes were assigned to orthogroups. We expect the remaining missing genes to either be the most recently evolved orphan genes in the branch leading to the *P. downsi* lineage or the consequence of a lack of inclusion of enough closely related species of *P. downsi* in the analysis. The distribution of these unique genes in the *P. downsi* genome is random and does not show specific clustering patterns across various locations of the genome.

We also compared the number of pairwise orthogroups that are uniquely shared among all eight species (Figure 3b). A total of 993 orthogroups were unique to Diptera (after excluding the outlier *H. melpomene*), 79 orthogroups were shared only between *M. domestica* and *P. downsi* and 12 putative gene families were unique to only *P. downsi*. These 12 gene families consisted of 27 genes and the great majority had the best BLAST hits against “uncharacterized” or “hypothetical” proteins in other related species (Supplementary Table 4). This result indicates that these gene families that appeared “unique” in *P. downsi* are likely due to the lack of proper gene annotation in other species.

We further examined the gene ontology terms of the 993 orthogroups unique to Diptera using the PANTHER gene ontology database (Mi *et al.*, 2021). The common biological processes of these genes included localization, locomotion, immune system processes, response to stimulus, and reproduction (Supplementary Figure 3), many of which are likely key genes for the overall development and function of dipterans.

Gene family evolution

We used a maximum likelihood approach to analyze changes in gene family size among the same eight species used for inferring orthologs (De Bie *et al.*, 2006). This method uses a statistical approach to model gene gain and loss, accounting for phylogenetic history and assess the significance of the observed gene family size differences among taxa. Across 11,112 orthogroups we identified before among eight taxa; 101 gene families were significantly expanding/contracting ($p < 0.01$) across the phylogenetic tree. 25 out of these 101 gene families were identified in the branch leading to *P. downsi*. The list of these gene families in *P. downsi* included those associated with insecticide resistance or detoxification and host defense or immunity proteins (Table 2).

Table 2: List of significantly expanding/contracting ($p < 0.01$) gene families in *P. downsi*

Functional Categories	Gene Family
Insecticide Resistance and Detoxification	Cytochrome P450
	Glutathione S Transferase
	Cuticular Protein
Defense against host Immunity	Fibrinogen C-terminal Domain-Containing Protein
	Scp Domain-Containing Protein

Gene families associated with insecticide resistance and detoxification

Cytochrome P450 gene family

We examined the number of genes in the cytochrome P450 (CYP450) family across the seven Diptera and the *H. melpomene* outgroup. *P. downsi* and *M. domestica* have an expanded CYP450 gene family in comparison to their most recent common ancestor with *G. morsitans* (family Glossinidae) (Figure 4a). For example, The *P. downsi* CYP450 family is composed of 102 genes in comparison to 66 in *G. morsitans*. An even greater level of expansion was observed in two other members of Muscidae (*M. domestica*: 143 genes and *S. calcitrans*: 193 genes).

Compared to *G. morsitans*, the expansion of CYP450 genes is mainly found in CYP4, CYP6, and CYP28 genes (Figure 4b). The CYP6 subfamily in *P. downsi* is composed of 25 genes, almost doubling the number from the 14 genes present in *G. morsitans*. We also found an expansion in the CYP4 subfamily from 15 genes in *G. morsitans* to 31 genes in *P. downsi*. A similar expansion was seen in the CYP28 subfamily (4 genes in *G. morsitans* to 8 genes in *P. downsi*).

Glutathione S-transferase gene family

The *P. downsi* Glutathione S-transferase family consists of 25 genes and the pattern across the dipterans is similar to observations of the Cytochrome P450 gene family expansion (Figure 4c). For example, in comparison to *G. morsitans* (15 genes), *P. downsi* and *M. domestica* have an expanded number of genes (25 genes in *P. downsi*, 31 genes in *M. domestica*). The GSTs are grouped into six subclasses (Delta, Epsilon, Omega, Sigma, Theta, and Zeta). The expansion of GSTs in *P. downsi* occurred mainly in the Delta, Epsilon, and Zeta subclasses (Figure 4d).

Cuticular Protein

The *P. downsi* cuticular gene family consists of 214 genes and the pattern across the dipterans is similar to observations of the Cytochrome P450 and GST gene family expansions (Supplementary Figure 4). For example, the *Muscidae* family (represented here by *P. downsi*, *M. domestica*, and *S. calcitrans*) has an expanded gene family in comparison to their most recent common ancestor with *G. morsitans* (115 genes). An even greater level of expansion was

observed by two other members of *Muscidae* (*M. domestica* – 357 genes and *S. calcitrans* – 270 genes).

Gene families associated with immunity

Two gene families (Fibrinogen C-terminal Domain-Containing Protein and Scp Domain-Containing Protein) associated with immunity were also identified as significantly changing in *P. downsi* (Table 2). Opposite of genes associated with insecticide resistance and detoxification (Cytochrome P450 and Glutathione S-transferase), these gene families showed a reduction in the number of genes in *P. downsi* compared to the most recent common ancestor with *G. morsitans* (Supplementary Table 5). They also show contraction in comparison with two included members of the Family *Muscidae* (*M. domestica* and *S. calcitrans*).

Discussion

In this study, we report the first genome sequence of the avian vampire fly, a highly invasive parasitic nest fly that threatens endemic avifauna of the Galápagos Islands (Fessl *et al.*, 2018; Causton *et al.*, 2019). This genome is meant to serve as an important resource to research efforts aimed at characterizing the molecular mechanisms of the fly's successful invasion in the Galápagos. The genome size of *P. downsi* is 971.6 Mb and its high quality is reflected both in terms of genome contiguity (scaffold N₅₀ of 1.3 Mb) and completeness (98% BUSCO gene-space score). Interestingly, the total number of annotated genes in *P. downsi* is slightly lower compared to other published fly genomes (*M. domestica* and *D. melanogaster*) (Table 1). The low gene count likely reflects the lack of *P. downsi* specific transcriptome data in this study, which perhaps led to reduced gene predictions. We aim to improve the *P. downsi* genome annotations

in the future using additional transcriptomic resources using additional flies collected from the Galápagos.

Transposable elements (TE) are typically non-coding sequences that can insert themselves in various places of the genome, often with neutral or deleterious phenotypic consequences (Bourque *et al.*, 2018). The role of TEs, as well as their evolution across insect genomes, is still an area of major research, but they are thought to be important drivers of genomic architecture depending on the location of the genome to which they insert themselves (i.e., coding versus non-coding regions). Furthermore, TE may also be a critical mechanism of adaptive evolution, as has been shown in an invasive ant species (Schrader *et al.*, 2014). Analysis of transposable elements in *P. downsi* and across other dipteran genomes showed a strong positive correlation between genome size and repeat content (Figure 2a), consistent with similar findings across other taxa (Lynch, 2007; Lamichhaney *et al.*, 2021). Interestingly, *P. downsi* had a higher number of DNA transposons (Class II TEs) than any other compared genome, including *M. domestica*, *S. calcitrans*, and *G. morsitans*. While long terminal repeat (LTR) transposons, LINEs, and SINEs were present in the species studied, *P. downsi* had no SINEs, a finding consistent with a study by Petersen and colleagues (Petersen *et al.*, 2019) showing that SINEs contribute less than 1% to the TE content of dipterans. However, it is important to note that some SINEs may be present in *P. downsi* but are currently masked as unclassified. Future research should explore the role of TEs, especially DNA transposons, in aiding the invasion of *P. downsi* to the Galápagos.

Comparative genomics analysis of *P. downsi* and other additional dipteran genomes allowed us to identify gene families that were significantly expanded/contracted ($p < 0.01$) in *P. downsi*. The list of these gene families in *P. downsi* included those associated with insecticide resistance or detoxification and host defense or immunity proteins, and we predict that these

gene families are associated with a successful invasion of *P. downsi* in the Galápagos (Table 1). E.g. CYP450 mono-oxygenases are a diverse superfamily of proteins, including enzymes, associated with catabolism and anabolism of xenobiotics and endogenous compounds. These monooxygenase-mediated metabolisms have allowed numerous insect species to develop insecticide resistance and detoxification (Scott, 1999; Wen *et al.*, 2011). The expansion of CYP450 genes in *P. downsi* is mainly found in CYP4 and CYP6 gene subfamilies (Figure 4b), which has also been shown in other species from the family Muscidae (Scott *et al.*, 2014; Olafson *et al.*, 2021). Interestingly, in higher Diptera, many of the genes within the CYP6 subfamily (e.g., CYP6A, CYP6G, CYP6D) are associated with insecticide resistance (Feyereisen, 2012). For example, the *Cyp6g1* gene is involved in resistance to the insecticide Dichlorodiphenyltrichloroethane (DDT) in *D. melanogaster* (Festucci-Buselli *et al.*, 2005). The CYP4 subfamily can also influence the breakdown of synthetic insecticides (Iga and Kataoka, 2012). *Cyp4d4v2*, *Cyp4g2*, and *Cyp6a38* can be co-up-regulated in house flies that are resistant to the insecticide permethrin, which is used to control *P. downsi* in the Galapagos (Zhu *et al.*, 2008). Overall, the expansion of CYP4 and CYP6 subfamilies may indicate the evolution of insecticide resistance in *P. downsi* over macroevolutionary time (i.e., before arriving in the Galápagos), which could have facilitated its invasion to the Galápagos and might affect its management on the islands.

Glutathione S-transferase (GSTs) was another major expanding gene family in *P. downsi*. GSTs are a highly conserved, large family of dimeric enzymes associated with detoxification of endogenous and/or xenobiotic compounds, such as insecticides (Ketterman *et al.*, 2011). The GST family is further grouped into six subclasses (Delta, Epsilon, Omega, Sigma, Theta, and Zeta), with Delta and Epsilon being specific subclasses found in the class Insecta. We observed

an expansion of GST genes in *P. downsi* relative to *G. morsitans*, but fewer than were found in *M. domestica* and *S. calcitrans*. The major expansions of the GST family in *P. downsi* were observed in Delta and Epsilon subclasses (Figure 4d). Insecticides, such as permethrin, are used to experimentally manipulate *P. downsi* abundance in bird nests to study the effects of the parasite on the health of the birds (Fessler *et al.*, 2010; Koop *et al.*, 2011; Koop, Le Bohec, *et al.*, 2013; Koop, Owen, *et al.*, 2013; Knutie *et al.*, 2014, 2016b; O'Connor *et al.*, 2014; McNew *et al.*, 2020; Adesso *et al.*, 2020). Increased expression of GSTs following permethrin exposure has been documented in several insect species including oriental fruit flies (*Bactrocera dorsalis*) (Hu *et al.*, 2008). Expansion of the GST gene family in *P. downsi* may be a result of such permethrin exposure. However, gene expression studies that further explore the role of the GST gene family in insecticide resistance and detoxification are needed across insect taxa including *P. downsi*. It is important to note that the expansion of CYP450 and GST families observed in *P. downsi* may be an artifact of phylogenetic relationships rather than ecological adaptations. Still, given the observed and predicted impacts of *P. downsi* on native endemic host populations, it is important to consider the implications of expanded gene families related to detoxification and insecticide resistance.

Previous studies have shown that Darwin's finch species can increase specific antibody responses to parasitism (Koop, Owen, *et al.*, 2013). This response is most prominent in adult females that are likely parasitized while brooding nestlings or eggs on the nest. However, little is known about the ability of host bird immunological responses to effectively reduce fly fitness, in part, because so little is known about the fly itself. One of the most prominent questions is whether the *P. downsi* possesses the ability to counter defend against host immune responses. We identified a reduction in the size of two additional gene families (Fibrinogen C-terminal

Domain-Containing and SCP domain-containing gene family), both with immune function properties in *P. downsi* (Table 1, Supplementary Table 5). Fibrinogen plays a key role in blood clot formation through the conversion of fibrinogen to insoluble fibrin (Weisel and Litvinov, 2017) and the C-terminal domain of fibrinogen is the primary binding site of platelets (Hanington and Zhang, 2011). The Sperm-coating glycoprotein (Scp) family contains, among other proteins, antigen 5 (Ag5), which is associated with the venom secretory ducts of stinging insects (Gibbs *et al.*, 2008). However, the interpretation of these results is difficult without further investigation into whether these genes are associated with innate immune responses of *P. downsi* toward their hosts or whether they might be important components of their feeding ecology. The goal of our study was not to make such inferences, but rather to highlight promising avenues of future research.

The invasion of *P. downsi* has had dramatic negative effects on the endemic avifauna of the Galápagos, including Darwin's finches. As researchers work to better understand the pathway of invasion and the ecological and evolutionary processes that may have facilitated its invasion to the Galápagos, the need for a high-quality whole genome sequence has grown. The addition of this resource is therefore meant to provide the foundation for further investigations using genomics tools in this system. These genomic resources will further allow us to understand the evolution of the *P. downsi* defense in response to host defenses or disease resistance itself. Gene expression studies could shed light on the development of larvae and at what stage they are most vulnerable to host defenses. Further population-scale resequencing of various populations of *P. downsi* will also allow us to explore mechanisms of local adaptation of the parasite to the environment across islands.

Data availability

All raw data generated in this study (raw short sequence reads and draft genome assembly) has been deposited to NCBI, under accession number PRJNAXXX. The final genome assembly and annotation can be found under the accession number GCA_XXX. Supplementary material is available on figshare.

Acknowledgment

We would like to thank Lauren Albert, Taylor Verrett, and Corrine Arthur for field assistance and the Galápagos Science Center and the Galápagos National Park for logistical support. We also thank Noah K. Whiteman for his helpful comments on the manuscript. The sampling was done under Galápagos National Parks permits PC 28-19 and Genetic Access permit MAE-DNB-CM-2016-0041. This project was supported by the Department of Biological Sciences, Kent State University to SL, and the University of Connecticut to SAK.

Competing interest statement

The authors declare no competing interests.

Author contributions

SL and JAHK conceived the idea. SAK, GJV, and JC collected the specimens. CMC and JAHK carried out the laboratory work. MR and SL designed and performed the bioinformatic analyses with the support of CMC and JAHK. SL and JAHK prepared the manuscript, and all authors edited and approved the final version.

List of Figures

Figure 1: The avian vampire fly, *Philornis downsi* **(a)**, parasitizes many endemic bird species of the Galapagos Islands, including the medium ground finch, *Geospiza fortis* **(b)**. The fly is parasitic in its larval forms **(c, bottom three)** when it feeds on the blood and other fluids of its avian hosts. The larva then pupates **(c, second from top)** and ecloses **(c, top)** as adult flies.

Figure 2: Landscape of Transposable elements in *P. downsi* **(a)** Comparison of repeat content and genome size across Diptera and its outgroup **(b)** Repeat statistics on various classes of transposable elements across dipteran genomes.

Figure 3: Orthogroups in *P. downsi* **(a)** Phylogenetic relationship between *P. downsi* and other seven published Diptera genomes, estimated using alignments from 3,069 orthogroups had singly copy orthogroups in each species. Horizontal bars for each species show number of orthogroups that are single-copy orthologs in all species, present in all species, present in the majority of species, present in few species, and unique to the species **(b)** Number of shared orthologs among all species.

Figure 4: Gene family evolution **(a)** Number of genes in Cytochrome P450 gene family across Diptera **(b)** Number of genes in various subfamilies of Cytochromes P450 gene family in *G. morsitans* and *P. downsi*, *M. domestica* and *D. melanogaster* **(c)** Number of genes in Glutathione S-transferase gene family across Diptera **(d)** Number of genes in various subfamilies of

Glutathione S-transferase gene family in *G. morsitans* and *P. downsi*, *M. domestica* and *D. melanogaster*.

References

- Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, *et al.* (2000). The genome sequence of *Drosophila melanogaster*. *Science* **287**: 2185–95.
- Addesso AM, Harvey JA, Vaziri GJ, Verrett TB, Albert L, Arthur C, *et al.* (2020). Effect of introduced parasites on the survival and microbiota of nestling cactus finches (*Geospiza scandens*) in the Galápagos Islands. *Journal of Ornithology* **161**: 1011–1019.
- Alves F, Langmore N, Heinsohn R, Stojanovic D (2020). ‘Self-fumigation’ of nests by an endangered avian host using insecticide-treated feathers increases reproductive success more than tenfold. *Animal Conservation*.
- Bao W, Kojima KK, Kohany O (2015). Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mobile DNA* **6**: 11.
- Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, *et al.* (2018). Ten things you should know about transposable elements. *Genome Biol* **19**: 199–199.
- Bulgarella M, Quiroga MA, Boulton RA, Ramirez IE, Moon RD, Causton CE, *et al.* (2017). Life Cycle and Host Specificity of the Parasitoid *Conura annulifera* (Hymenoptera: Chalcididae), a Potential Biological Control Agent of *Philornis downsi* (Diptera: Muscidae) in the Galapagos Islands. *Ann Entomol Soc Am* **110**: 317–328.
- Bulgarella M, Quiroga MA, Brito vera GA, Dregni JS, Cunningham F, Mosquera Munoz DA, *et al.* (2015). *Philornis downsi* (Diptera: Muscidae), an avian nest parasite invasive to the Galapagos islands, in mainland Ecuador. *Annals of the Entomological Society of America*.
- Bulgarella M, Quiroga MA, Heimpel GE (2019). Additive negative effects of *Philornis* nest parasitism on small and declining Neotropical bird populations. *Bird Conservation International* **29**: 339–360.
- Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, *et al.* (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Res* **18**: 188–196.
- Causton CE, Cunningham F, Tapia W (2013). *Management of the avian parasite Philornis downsi in the Galapagos Islands: a collaborative and strategic action plan*. GNPS, GCREG, CDF, and GC: Puerto Ayora, Galapagos, Ecuador.

- 510 Causton CE, Moon RD, Cimadom A, Boulton RA, Cedeño D, Lincango MP, *et al.* (2019).
511 Population dynamics of an invasive bird parasite, *Philornis downsi* (Diptera: Muscidae),
512 in the Galapagos Islands. *PLoS One* **14**: e0224125.
- 513 Causton CE, Peck SB, Sinclair BJ, Roque-Albelo L, Hodgson CJ, Landry B (2006). Alien
514 insects: Threats and implications for conservation of Galápagos Islands. *Annals of the*
515 *Entomological Society of America* **99**: 121–143.
- 516 Couri MS, de Carvalho CJB, Lowenberg-Neto P (2007). Phylogeny of *Philornis* Meinert species
517 (Diptera: Muscidae). *Zootaxa*: 19–26.
- 518 Cunningham F, Achuthan P, Akanni W, Allen J, Amode MR, Armean IM, *et al.* (2019).
519 Ensembl 2019. *Nucleic Acids Research* **47**: D745–D751.
- 520 Cunnigham F, Fessl B, Sevilla CR, Young GR, La Greco N (2017). *Manejo de la*
521 *conservacion a largo plazo para salvar al pinzon de manglar (Camarhynchus heliobates)*
522 *en peligro critico de extincion*. DPNG, CGREG, FCD and GC: Puerto Ayora, Galapagos,
523 Ecuador.
- 524 De Bie T, Cristianini N, Demuth JP, Hahn MW (2006). CAFE: a computational tool for the
525 study of gene family evolution. *Bioinformatics (Oxford, England)* **22**: 1269–1271.
- 526 Dodge HR (1955). New Muscid flies from Florida and the West Indies (Diptera: Muscidae). *The*
527 *Florida Entomologist* **38**: 147–151.
- 528 Dodge HR (1963). A new *Philornis* with coprophagous larva, and some related species (Diptera:
529 Muscidae). *Journal of the Kansas Entomological Society* **36**: 239–247.
- 530 Dudaniec RY, Fessl B, Kleindorfer S (2007). Interannual and interspecific variation in intensity
531 of the parasitic fly, *Philornis downsi*, in Darwin’s finches. *Biological Conservation* **139**:
532 325–332.
- 533 Dudaniec RY, Gardner MG, Donnellan S, Kleindorfer S (2008). Genetic variation in the invasive
534 avian parasite, *Philornis downsi* (Diptera, Muscidae) on the Galapagos archipelago. *BMC*
535 *Ecology* **8**: 13.
- 536 Dudaniec RY, Kleindorfer S (2006). Effects of the parasitic flies of the genus *Philornis* (Diptera :
537 Muscidae) on birds. *Emu* **106**: 13–20.
- 538 Dvorak M, Vargas H, Fessl B, Tebbich S (2004). On the verge of extinction: a survey of the
539 mangrove finch *Cactospiza heliobates* and its habitat on the Galápagos Islands. *Oryx* **38**:
540 1–9.
- 541 Emms DM, Kelly S (2015). OrthoFinder: solving fundamental biases in whole genome
542 comparisons dramatically improves orthogroup inference accuracy. *Genome Biology* **16**:
543 157.

- 544 Fessl B, Heimpel G, Causton C (2018). Invasion of an avian nest parasite, *Philornis downsi*, to
545 the Galapagos Islands: colonization history, adaptations to novel ecosystems, and
546 conservation challenges. In: Parker P (ed) *Disease Ecology, Social and Ecological*
547 *Interactions in the Galapagos Islands*, Springer International Publishing.
- 548 Fessl B, Sinclair BJ, Kleindorfer S (2006). The life-cycle of *Philornis downsi* (Diptera :
549 Muscidae) parasitizing Darwin's finches and its impacts on nestling survival.
550 *Parasitology* **133**: 739–747.
- 551 Fessl B, Tebbich S (2002). *Philornis downsi* - a recently discovered parasite on the Galápagos
552 archipelago - a threat for Darwin's finches? *Ibis* **144**: 445–451.
- 553 Fessl B, Young GH, Young RP, Rodríguez-Matamoros J, Dvorak M, Tebbich S, *et al.* (2010).
554 How to save the rarest Darwin's finch from extinction: the mangrove finch on Isabela
555 Island. *Philos Trans R Soc Lond B Biol Sci* **365**: 1019–1030.
- 556 Festucci-Buselli RA, Carvalho-Dias AS, de Oliveira-Andrade M, Caixeta-Nunes C, Li HM,
557 Stuart JJ, *et al.* (2005). Expression of Cyp6g1 and Cyp12d1 in DDT resistant and
558 susceptible strains of *Drosophila melanogaster*. *Insect Mol Biol* **14**: 69–77.
- 559 Feyereisen R (2012). Insect CYP genes and P450 enzymes. In: Gilbert LI, editor. *Insect*
560 *Molecular Biology and Biochemistry*: 236–316.
- 561 Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, *et al.* (2019).
562 RepeatModeler2: automated genomic discovery of transposable element families.
563 *bioRxiv*: 856591.
- 564 Gibbs GM, Roelants K, O'Bryan MK (2008). The CAP Superfamily: Cysteine-Rich Secretory
565 Proteins, Antigen 5, and Pathogenesis-Related 1 Proteins—Roles in Reproduction,
566 Cancer, and Immune Defense. *Endocrine Reviews* **29**: 865–897.
- 567 Grant PR, Grant BR, Petren K, Keller LF (2005). Extinction behind our backs: the possible fate
568 of one of the Darwin's finch species on Isla Floreana, Galápagos. *Biological*
569 *Conservation* **122**: 499–503.
- 570 Hanington PC, Zhang S-M (2011). The primary role of fibrinogen-related proteins in
571 invertebrates is defense, not coagulation. *J Innate Immun* **3**: 17–27.
- 572 Heimpel GE, Hillstrom A, Freund D, Knutie SA, Clayton DH (2017). Invasive Parasites and the
573 Fate of Darwin's Finches in the Galapagos Islands: The Case of the Vegetarian Finch
574 (*Platyspiza crassirostris*). *Wilson J Ornithol* **129**: 345–349.
- 575 Hu J, Zhang JL, Nardi F, Zhang RJ (2008). Population genetic structure of the melon fly,
576 *Bactrocera cucurbitae* (Diptera: Tephritidae), from China and Southeast Asia. *Genetica*
577 **134**: 319–24.
- 578 Iga M, Kataoka H (2012). Recent Studies on Insect Hormone Metabolic Pathways Mediated by
579 Cytochrome P450 Enzymes. *Biological and Pharmaceutical Bulletin* **35**: 838–843.

580 Kapusta A, Suh A, Feschotte C (2017). Dynamics of genome size evolution in birds and
581 mammals. *Proceedings of the National Academy of Sciences* **114**: E1460 LP-E1469.

582 Ketterman AJ, Saisawang C, Wongsantichon J (2011). Insect glutathione transferases. *Drug*
583 *Metabolism Reviews* **43**: 253–265.

584 Kidwell MG (2002). Transposable elements and the evolution of genome size in eukaryotes.
585 *Genetica* **115**: 49–63.

586 Knutie SA, McNew SM, Bartlow AW, Vargas DA, Clayton DH (2014). Darwin’s finches
587 combat introduced nest parasites with fumigated cotton. *Current Biology* **24**: R355–
588 R356.

589 Knutie SA, Owen JP, McNew SM, Bartlow AW, Arriero E, Herman JM, *et al.* (2016a).
590 Galapagos mockingbirds tolerate introduced parasites that affect Darwin’s finches.
591 *Ecology* **97**: 940–950.

592 Koop JAH, Causton CE, Bulgarella M, Cooper E, Heimpel GE (2021). Population structure of a
593 nest parasite of Darwin’s finches within its native and invasive ranges. *Conservation*
594 *Genetics* **22**: 11–22.

595 Koop JAH, Huber SK, Lavery SM, Clayton DH (2011). Experimental demonstration of the
596 fitness consequences of an introduced parasite of Darwin’s finches. *PLoS ONE* **6**: e19706
597 doi:10.1371/journal.pone.0019706.

598 Koop JAH, Kim PS, Knutie SA, Adler F, Clayton DH (2015). Introduced parasitic fly may lead
599 to local extinction of Darwin’s finch populations. *Journal of Applied Ecology*.

600 Koop JAH, Le Bohec C, Clayton DH (2013). Dry year does not reduce invasive parasitic fly
601 prevalence or abundance in Darwin’s finch nests. *Reports in Parasitology* **3**: 11–17.

602 Koop JAH, Owen JP, Knutie SA, Aguilar MA, Clayton DH (2013). Experimental demonstration
603 of a parasite-induced response in wild birds: Darwin’s finches and introduced nest flies.
604 *Ecology and Evolution* **3**: 2514–2523.

605 Lamichhaney S, Catullo R, Keogh JS, Clulow S, Edwards SV, Ezaz T (2021). A bird-like
606 genome from a frog: Mechanisms of genome size reduction in the ornate burrowing frog,
607 *Platyplectrum ornatum*. *Proceedings of the National Academy of Sciences* **118**:
608 e2011649118.

609 Li H, Durbin R (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.
610 *Bioinformatics* **25**: 1754–1760.

611 Lynch M (2007). *The Origins of Genome Architecture*, 1 edition. Sinauer Associates Inc:
612 Sunderland.

613 McNew SM, Knutie SA, Clayton DH (2020). No evidence of sex ratio manipulation by
614 Galápagos mockingbirds in response to environment. *Journal of Avian Biology* **51**.

615 Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, *et al.* (2021). PANTHER
616 version 16: a revised family classification, tree-based classification tool, enhancer regions
617 and extensive API. *Nucleic Acids Research* **49**: 394–403.

618 O'Connor JA, Robertson J, Kleindorfer S (2014). Darwin's finch begging intensity does not
619 honestly signal need in parasitised nests. *Ethology* **120**: 228–237.

620 Olafson PU, Aksoy S, Attardo GM, Buckmeier G, Chen X, Coates CJ, *et al.* (2021). The genome
621 of the stable fly, *Stomoxys calcitrans*, reveals potential mechanisms underlying
622 reproduction, host interactions, and novel targets for pest control. *BMC Biology* **19**: 41.

623 Peters KJ, Kleindorfer S (2018). Avian population trends in Scalesia forest on Floreana Island
624 (2004–2013): Acoustical surveys cannot detect hybrids of Darwin's tree finches
625 *Camarhynchus* spp. *Bird Conservation International* **28**: 319–335.

626 Petersen M, Armisen D, Gibbs RA, Hering L, Khila A, Mayer G, *et al.* (2019). Diversity and
627 evolution of the transposable element repertoire in arthropods with particular reference to
628 insects. *BMC Evol Biol* **19**: 11–11.

629 Pourrajab F, Hekmatimoghaddam S (2021). Transposable elements, contributors in the evolution
630 of organisms (from an arms race to a source of raw materials). *Heliyon* **7**: e06029–
631 e06029.

632 Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020). GenomeScope 2.0 and Smudgeplot for
633 reference-free profiling of polyploid genomes. *Nature Communications* **11**: 1432.

634 Schrader L, Kim JW, Ence D, Zimin A, Klein A, Wyschetzki K, *et al.* (2014). Transposable
635 element islands facilitate adaptation to novel environments in an invasive species. *Nature*
636 *Communications* **5**: 5495.

637 Scott JG (1999). Cytochromes P450 and insecticide resistance. *Insect Biochem Mol Biol* **29**:
638 757–77.

639 Scott MJ, Benoit JB, Davis RJ, Bailey ST, Varga V, Martinson EO, *et al.* (2020). Genomic
640 analyses of a livestock pest, the New World screwworm, find potential targets for genetic
641 control programs. *Communications Biology* **3**: 424–424.

642 Scott JG, Warren Wc Fau - Beukeboom LW, Beukeboom Lw Fau - Bopp D, Bopp D Fau - Clark
643 AG, Clark Ag Fau - Giers SD, Giers Sd Fau - Hediger M, *et al.* (2014). Genome of the
644 house fly, *Musca domestica* L., a global vector of diseases with adaptations to a septic
645 environment.

646 Smit Hubley P. AFA R& Green (2013). RepeatMasker Open-4.0,.

647 Spalding MG, Mertins JW, Walsh PB, Morin KC, Dunmore DE, Forrester DJ (2002). Burrowing
648 fly larvae (*Philornis porteri*) associated with mortality of eastern bluebirds in Florida.
649 *Journal of Wildlife Diseases* **38**: 776–783.

- Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, *et al.* (2017). GenomeScope: fast reference-free genome profiling from short reads. *Bioinformatics* **33**: 2202–2204.
- Waterhouse RM, Seppey M, Simao FA, Manni M, Ioannidis P, Klioutchnikov G, *et al.* (2017). BUSCO applications from quality assessments to gene prediction and phylogenomics. *Molecular biology and evolution*.
- Weisel JW, Litvinov RI (2017). Fibrin Formation, Structure and Properties. *Subcell Biochem* **82**: 405–456.
- Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB (2017). Direct determination of diploid genome sequences. *Genome Research* **27**: 757–767.
- Wen Z, Zhang X, Zhang Y (2011). P450—mediated Insecticide Detoxification and Its Implication in Insecticide Efficacy. In: Liu T, Kang L (eds) *Recent Advances in Entomological Research: From Molecular Biology to Pest Management*, Springer Berlin Heidelberg: Berlin, Heidelberg, pp 229–245.
- Zheng GXY, Lau BT, Schnall-Levin M, Jarosz M, Bell JM, Hindson CM, *et al.* (2016). Haplotyping germline and cancer genomes with high-throughput linked-read sequencing. *Nature Biotechnology* **34**: 303–311.
- Zhu F, Li T, Zhang L, Liu N (2008). Co-up-regulation of three P450 genes in response to permethrin exposure in permethrin resistant house flies, *Musca domestica*. *BMC Physiol* **8**: 18.

b

Figure 1

a

b

Figure 2

Figure 4

Supplementary Materials

The genome sequence of the avian vampire fly (Philornis downsi), an invasive nest parasite of Darwin's finches

Supplementary Table 1: Genome Statistics of *P. downsi* estimated from short-sequencing reads

<i>Property</i>	<i>min</i>	<i>max</i>
Heterozygosity	1.40879%	1.42194%
Genome Haploid Length	796,797,589 bp	799,050,413 bp
Genome Repeat Length	232,405,721 bp	233,062,812 bp
Genome Unique Length	564,391,869 bp	565,987,600 bp
Model Fit	95.7713%	98.7012%
Read Error Rate	1.228%	1.228%

Supplementary Table 2: Repeat statistics in *P. downsi* genome

sequences: 41,176
total length: 971,6346,46 bp
GC level: 35.10 %
bases masked: 502,200,627 bp (51.69 %)

	number of elements*	length occupied	percentage of sequence
Retroelements	224657	90253146 bp	9.29 %
SINEs:	58	6761 bp	0.00 %
Penelope	3187	661022 bp	0.07 %
LINEs:	196268	74550945 bp	7.67 %
CRE/SLACS	0	0 bp	0.00 %
L2/CR1/Rex	43733	18550153 bp	1.91 %
R1/LOA/Jockey	15852	7502835 bp	0.77 %
R2/R4/NeSL	834	673762 bp	0.07 %
RTE/Bov-B	102395	29493616 bp	3.04 %

L1/CIN4	1415	94740 bp	0.01 %
LTR elements:	28331	15695440 bp	1.62 %
BEL/Pao	12565	7084057 bp	0.73 %
Ty1/Copia	1561	775084 bp	0.08 %
Gypsy/DIRS1	9951	7481940 bp	0.77 %
Retroviral	2481	143387 bp	0.01 %
DNA transposons	621139	227262774 bp	23.39 %
hobo-Activator	8749	1542960 bp	0.16 %
Tc1-IS630-Pogo	576750	212960682 bp	21.92 %
En-Spm	0	0 bp	0.00 %
MuDR-IS905	0	0 bp	0.00 %
PiggyBac	1280	395680 bp	0.04 %
Tourist/Harbinger	4060	1924044 bp	0.20 %
Other	275	11451 bp	0.00 %
Rolling-circles	19653	4597029 bp	0.47 %
Unclassified:	828801	166508339 bp	17.14 %
Total interspersed repeats:		484024259 bp	49.82 %
Small RNA:	7148	2907047 bp	0.30 %
Satellites:	796	124775 bp	0.01 %
Simple repeats:	213281	8909635 bp	0.92 %
Low complexity:	34278	1638267 bp	0.17 %

Supplementary Table 3: Number of shared orthogroups among various dipteran species and their outgroup

	<i>H.melpomene</i>	<i>A.aegypti</i>	<i>C.capitata</i>	<i>S.calcitrans</i>	<i>M.domestica</i>	<i>P.downsi</i>	<i>G.morsitans</i>	<i>D.melanogaster</i>
<i>H.melpomene</i>		6,978	6,874	6,883	6,910	6,800	6,337	6,859
<i>A.aegypti</i>	6,978		8,015	8,008	8,018	7,796	7,228	7,995
<i>C.capitata</i>	6,874	8,015		9,123	9,186	8,855	8,014	9,096
<i>S.calcitrans</i>	6,883	8,008	9,123		9,650	9,197	8,087	9,041
<i>M.domestica</i>	6,910	8,018	9,186	9,650		9,295	8,144	9,117
<i>P.downsi</i>	6,800	7,796	8,855	9,197	9,295		8,103	8,770
<i>G.morsitans</i>	6,337	7,228	8,014	8,087	8,144	8,103		7,940
<i>D.melanogaster</i>	6,859	7,995	9,096	9,041	9,117	8,770	7,940	

Supplementary Table 4: *Gene families unique only to P. downsi*

EGT47331	hypothetical protein CAEBREN_08836 [Caenorhabditis brenneri]
GBO13109	hypothetical protein AVEN_233885-1 [Araneus ventricosus]
GBP44057	PiggyBac transposable element-derived protein 4 [Eumeta japonica]
KMQ83311	transposable element tc3 transposase [Lasius niger]
KNC20639	hypothetical protein FF38_06891, partial [Lucilia cuprina]
KNC20966	hypothetical protein FF38_09018 [Lucilia cuprina]
KNC26906	hypothetical protein FF38_10930 [Lucilia cuprina]
KXJ68891	hypothetical protein RP20_CCG001210 [Aedes albopictus]
OAF68034	hypothetical protein A3Q56_04227 [Intoshia linei]
PCG68510	hypothetical protein B5V51_5157, partial [Heliothis virescens]
XP_011295145	PREDICTED: tigger transposable element-derived protein 6-like isoform X2 [Musca domestica]
XP_011295374	PREDICTED: tigger transposable element-derived protein 6 [Musca domestica]
XP_013109704	PREDICTED: uncharacterized protein LOC106088638 [Stomoxys calcitrans]
XP_017475229	PREDICTED: uncharacterized protein LOC108365650 [Rhagoletis zephyria]
XP_017478109	PREDICTED: uncharacterized protein LOC108367917 [Rhagoletis zephyria]
XP_017478991	PREDICTED: uncharacterized protein LOC108368617 [Rhagoletis zephyria]
XP_017479715	PREDICTED: uncharacterized protein LOC108369194 [Rhagoletis zephyria]
XP_017481195	PREDICTED: RNA-directed DNA polymerase from mobile element jockey-like [Rhagoletis zephyria]
XP_019891578	PREDICTED: ATP-binding cassette sub-family A member 3-like [Musca domestica]
XP_019894716	PREDICTED: uncharacterized protein LOC105262305 isoform X1 [Musca domestica]
XP_021704105	protein ALP1-like [Aedes aegypti]
XP_022823959	piggyBac transposable element-derived protein 4-like [Spodoptera litura]
XP_022834134	uncharacterized protein LOC111361914 [Spodoptera litura]
XP_033325321	uncharacterized protein LOC117219890 [Megalopta genalis]
XP_036214104	trypsin zeta-like [Bactrocera oleae]
XP_036337749	uncharacterized protein LOC118747737 isoform X3 [Rhagoletis pomonella]
XP_036342696	uncharacterized protein LOC118751975 [Rhagoletis pomonella]

Supplementary Table 5: Number of genes in *Fibrinogen C-Terminal Domain-Containing* gene family and *SCP domain-containing* gene family

Species	Fibrinogen C-Terminal Domain-Containing gene family	SCP domain-containing gene family
<i>H.melpomene</i>	1	2
<i>A.aegypti</i>	33	20
<i>C.capitata</i>	10	6
<i>S.calcitrans</i>	45	17
<i>M.domestica</i>	32	25
<i>P.downsi</i>	5	7
<i>G.morsitans</i>	7	7
<i>D.melanogaster</i>	11	15

Supplementary Figure 1: *k*-mer spectrum and fitted modelling used for estimating genome parameters of *P. downsi* from short sequencing reads

Supplementary Figure 2: Distribution of the total coverage of the *k*-mer pair (y-axis) against relative minor *k*-mer coverage (x-axis) providing evidence of diploidy in *P. downsi*.

Supplementary Figure 3: *Gene ontology terms associated with 993 gene families unique to diptera*

Supplementary Figure 4: *Number of genes in Cuticular gene family across dipterans*