Control of Arabidopsis shoot stem cell homeostasis by two antagonistic CLE peptide signalling pathways

Jenia Schlegel, Grégoire Denay, Karine Gustavo Pinto, Yvonne Stahl, Julia Schmid, Patrick Blümke, and Rüdiger Simon*

Institute for Developmental Genetics and Cluster of Excellence on Plant Sciences, Heinrich Heine University, Universitätsstraße 1, 40225 Düsseldorf, Germany

*Corresponding author; email: ruediger.simon@hhu.de

Keywords: shoot meristem, CLAVATA signalling, stem cells, CLE40, BAM1, WUS, shape

Abstract

Stem cell homeostasis in plant shoot meristems requires tight coordination between stem cell proliferation and cell differentiation. In *Arabidopsis*, stem cells express the secreted dodecapeptide CLAVATA3 (CLV3), which signals through the leucine-rich repeat (LRR)–receptor kinase CLAVATA1 (CLV1) and related CLV1-family members to downregulate expression of the homeodomain transcription factor *WUSCHEL* (*WUS*). *WUS* protein moves from cells below the stem cell domain to the meristem tip and promotes stem cell identity, together with *CLV3* expression, generating a negative feedback loop. How stem cell activity in the meristem centre is coordinated with organ initiation and cell differentiation at the periphery is unknown.

We show here that the *CLE40* gene, encoding a secreted peptide closely related to *CLV3*, is expressed in the SAM in differentiating cells in a pattern complementary to that of *CLV3*. *CLE40* promotes *WUS* expression via *BAM1*, a CLV1-family receptor, and *CLE40* expression is in turn repressed in a *WUS*-dependent manner. Together, *CLE40-BAM1-WUS* establish a second negative feedback loop. We propose that stem cell homeostasis is achieved through two intertwined pathways that adjust *WUS* activity and incorporate information on the size of the stem cell domain, via *CLV3-CLV1*, and on cell differentiation via *CLE40-BAM1*.

Introduction

In angiosperms, the stem cell domain in shoot meristem is controlled by the directional interplay of two adjacent groups of cells. These are the central zone (CZ) at the tip of the dome-shaped meristem, comprising slowly dividing stem cells, and the underlying cells of the organising centre (OC). Upon stem cell division, daughter cells are displaced laterally into the peripheral zone (PZ), where they can enter differentiation pathways (Fletcher et al., 1999; Hall & Watt, 1989; Reddy et al., 2004; Schnablová et al., 2020; Stahl & Simon, 2005; Steeves & Sussex, 1989). Cells in the OC express the homeodomain transcription factor *WUSCHEL* (*WUS*), which moves through plasmodesmata to CZ cells to maintain stem cell fate and promote expression of the secreted signalling peptide *CLAVATA3* (*CLV3*) (Brand et al., 2000; Daum et al., 2014; Müller et al., 2006; Schoof et al., 2000; Yadav et al., 2011). Perception of *CLV3* by plasma-membrane localised receptors in the OC cells triggers a signal transduction cascade and downregulates *WUS* activity, thus establishing a negative feedback loop (Mayer et al., 1998; Ogawa et al., 2008; Yadav et al., 2011). Mutants of *CLV3* or its receptors (see below) fail to confine *WUS* expression and cause stem cell proliferation, while *WUS* mutants cannot maintain an active stem cell population (Brand et al., 2002; Clark et al., 1993, 1995; Endrizzi et al., 1996; Laux et al., 1996; Schoof et al., 2000). *WUS* function in the OC is negative regulated by *HAM* transcription factors, and only *WUS* protein that moves upwards to the stem cell zone, which lacks *HAM* expression, can activate *CLV3* expression (Han et al., 2020; Zhou et al., 2018). The *CLV3-WUS* interaction can serve to maintain the relative sizes of the CZ and OC, and thereby meristem growth along the apical-basal axis. However, cell loss from the PZ due to production of lateral organs requires a compensatory size increase of the stem cell domain.

The CLV3 signalling pathway, which acts along the apical-basal axis of the meristem, has been widely studied in several plant species and shown to be crucial for stem cell homeostasis in shoot and floral meristems (Somssich et al., 2016). The CLV3 peptide is perceived by a leucine-rich-repeat (LRR) receptor kinase, CLAVATA1 (CLV1), which interacts with coreceptors of the CLAVATA3 INSENSITIVE RECEPTOR KINASES (CIK) 1-4 family (Clark et al., 1997; Cui et al., 2018). CLV1 activation involves autophosphorylation, interaction with membrane-associated and cytosolic kinases and phosphatases (Blümke et al., 2021; Defalco et al., 2021). Furthermore, heterotrimeric G-proteins and MAPKs have been implicated in this signal transduction cascade in maize and Arabidopsis (Betsuyaku et al., 2011; Bommert et al., 2013; Ishida et al., 2014; Lee et al., 2019). Besides CLV1, several other receptors contribute to WUS regulation, among them RECEPTOR-LIKE PROTEIN KINASE2 (RPK2), the CLAVATA2-CORYNE heteromer (CLV2-CRN) and BARELY ANY MERISTEM1-3 (BAM1-3) (Bleckmann et al., 2010; DeYoung & Clark, 2008; Hord et al., 2006; Jeong et al., 1999; Kinoshita et al., 2010; Müller et al., 2008). The BAM receptors share high sequence similarity with CLV1, and perform diverse functions throughout plant development. Double mutants of *BAM1* and *BAM2* maintain smaller shoot and floral meristems, thus displaying the opposite phenotype to mutants of CLV1 (DeYoung et al., 2006; DeYoung & Clark, 2008; Hord et al., 2006). Interestingly, ectopic expression experiments showed that CLV1 and BAM1 can perform similar functions in stem cell control (Nimchuk et al., 2015). In addition, one study showed that CLV3 could interact with CLV1 and BAM1 in cell extracts (Shinohara & Matsubayashi, 2015), although another *in vitro* study did not detect *BAM1-CLV3* interaction at physiological levels of CLV3 (Crook et al., 2020). Furthermore, CLV1 was shown to act as a negative regulator of *BAM1* expression, which was interpreted as a genetic buffering system, whereby a loss of CLV1 is compensated by upregulation of BAM1 in the meristem

84 centre (Nimchuk, 2017; Nimchuk et al., 2015). Comparable genetic compensation
 85 models for CLE peptide signalling in stem cell homeostasis were established for other
 86 species, such as tomato and maize (Rodriguez-Leal et al., 2019).
 87 Maintaining the overall architecture of the shoot apical meristem during the entire life
 88 cycle of the plant requires replenishment of differentiating stem cell descendants in the
 89 PZ, indicating that cell division rates and cell fate changes in both regions are closely
 90 connected (Stahl & Simon, 2005). Overall meristem size is restricted by the ERECTA-
 91 family signalling pathway, which is activated by EPIDERMAL PATTERNING FACTOR
 92 (EPF)-LIKE (EPFL) ligands from the meristem periphery and confines both *CLV3* and
 93 *WUS* expression (Mandel et al., 2014; Shpak, 2013; Shpak et al., 2004; Torii et al.,
 94 1996; Zhang et al., 2021). In the land plant lineage, the shoot meristems of bryophytes
 95 such as the moss *Physcomitrium patens* appear less complex than those of
 96 angiosperms, and carry only a single apical stem cell which ensures organ initiation by
 97 continuous asymmetric cell divisions (de Keijzer et al., 2021; Harrison et al., 2009).
 98 Broadly expressed CLE peptides were here found to restrict stem cell identity, and act
 99 in division plane control (Whitewoods et al., 2018). Proliferation of the apical notch cell
 100 in the liverwort *Marchantia polymorpha* is promoted by MpCLE2 peptide which acts
 101 from outside the stem cell domain via the receptor MpCLV1, while cell proliferation is
 102 confined by MpCLE1 peptide through a different receptor (Hata & Kyoizuka, 2021;
 103 Hirakawa et al., 2019, 2020; Takahashi et al., 2021). Thus, antagonistic control of stem
 104 cell activities through diverse CLE peptides is conserved between distantly related land
 105 plants. In the grasses, several CLEs were found to control the stem cell domain. In
 106 maize, *ZmCLE7* is expressed from the meristem tip, while *ZmFCP1* is expressed in
 107 the meristem periphery and its centre. Both peptides restrict stem cell fate via
 108 independent receptor signalling pathways (Liu et al., 2021; Rodriguez-Leal et al.,
 109 2019). In rice, overexpression of the CLE peptides OsFCP1 and OsFCP2

downregulates the homeobox gene *OSH1* and arrests meristem function (Ohmori et al., 2013; Suzuki et al., 2008). Common for rice and maize, CLE peptide signalling restricts stem cell activities in the shoot meristem, but a stem cell promoting pathway were not been identified so far.

Importantly, how stem cell activities in the CZ and OC are coordinated to regulate organ initiation and cell differentiation in the PZ, which is crucial to maintain an active meristem, is not yet known. In maize, the CLV3-related peptide ZmFCP1 was suggested to be expressed in primordia, and convey a repressive signal on the stem cell domain (Je et al., 2016). In Arabidopsis, the most closely related peptide to CLV3 is CLE40, which was shown to act in the root meristem to restrict columella stem cell fate and regulate the expression of the *WUS* paralog *WOX5* (Berckmans et al., 2020; Hobe et al., 2003; Pallakies & Simon, 2014; Stahl et al., 2013; Stahl & Simon, 2010). Functions of CLE40 in the SAM have not previously been described. Overexpression of *CLE40* causes shoot stem cell termination, while *CLE40* expression from the *CLV3* promoter fully complements the shoot and floral meristem defects of *clv3* mutants (Hobe et al., 2003). We therefore hypothesized that *CLE40* could act in a *CLV3*-related pathway in shoot stem cell control.

Here, we show that the expression level of *WUS* in the OC is subject to feedback regulation from the PZ, which is mediated by the secreted peptide CLE40. In the shoot meristem, *CLE40* is expressed in a complementary pattern to *CLV3*, and excluded from the CZ and OC. In *cle40* loss of function mutants, *WUS* expression is reduced, and shoot meristems remain small and flat, indicating that *CLE40* signalling is required to maintain *WUS* expression in the OC. Ectopic expression of *WUS* represses *CLE40* expression, while in *wus* loss-of-function mutants *CLE40* is expressed in the meristem centre, indicating that CLE40, in contrast to CLV3, is subject to negative feedback

135 regulation by WUS. CLE40 likely acts as an autocrine signal that is perceived by BAM1
 136 in a domain flanking the OC.
 137 Based on our findings, we propose a new model for the regulation of the stem cell
 138 domain in the shoot meristem in which signals and information from both, the CZ and
 139 the PZ are integrated through two interconnected negative feedback loops that sculpt
 140 the dome-shaped shoot meristems of angiosperms.

Results

CLE40 signalling promotes IFM growth from the peripheral zone

Previous studies showed that *CLE40* expression from the *CLV3* promoter can fully complement a *clv3-2* mutant, indicating that CLE40 can substitute CLV3 function in the shoot meristem to control stem cell homeostasis, if expressed from the stem cell domain. Furthermore, while all other *CLE* genes in Arabidopsis lack introns, the *CLE40* and *CLV3* genes carry two introns at very similar positions (Hobe et al., 2003). Phylogenetic analysis revealed that CLV3 and CLE40 locate in the same cluster together with CLV3 orthologues from rice, maize and tomato (Goad et al., 2017) (Fig. 1A).

Mutations in *CLE40* were previously found to affect distal stem cell maintenance in the root meristem, revealing that a CLV3 related signalling pathway also operates in the root stem cell niche. To uncover a potential role of *CLE40* in shoot development, we analysed seedling and flower development, and inflorescence meristem (IFM) sizes of the wild type *Col-0*, and *clv3-9* and *cle40-2* loss-of-function mutants. At 4 weeks after germination (WAG), leaves of *clv3-9* mutants remained shorter than those of *Col-0* or *cle40-2* (Fig1-SupplFig.1). After floral induction, the inflorescences of *clv3-9* mutants were compact with many more flowers than the wild type, while *cle40-2* mutant inflorescences appeared smaller than the control (Fig. 1B-D). To first investigate effects on meristem development in detail, longitudinal optical sections through the inflorescence meristem (IFM) at 6 WAG were obtained by confocal microscopy and meristem areas were analysed (Fig. 1B-E). In *clv3-9* mutants, meristem areas increased to approx. 450% of wild type (*Col-0*) levels, while shoot meristems from 4 independent *cle40* mutant alleles in a *Col-0* background (*cle40-2*, *cle40-cr1*, *cle40-cr2*, *cle40-cr3*) reached only up to 65% of wild type (Fig. 1E, Fig1-SupplFig.2C) (Yamaguchi et al., 2017). Next, we used carpel number as a rough proxy for flower meristem (FM)

size, which was 2 ± 0.0 (N=290) in *Col-0* and *cle40-2* (N=290) but 3.7 ± 0.4 (N=340) in *clv3-9* (Fig1-SupplFig.3). Hence, we concluded that CLE40 mainly promotes IFM growth, whereas CLV3 serves to restrict both IFM and FM sizes.

Fig. 1: CLV3 and CLE40 exert opposite effects on meristem size

(A) The amino acid (AA) sequences of the mature CLV3 and CLE40 peptides differ in four AAs (differences marked in red). (B) *Col-0* inflorescence at 6 WAG with flowers. (B') IFM at 6 WAG, maximum intensity projection (MIP) of a z-stack taken by confocal microscopy. (C) *clv3-9* inflorescence at 6 WAG (C') MIP of a *clv3-9* IFM at 6 WAG. (D) Inflorescence of *cle40-2* at 6 WAG (D') MIP of a *cle40-2* IFM. (E) Box and whisker plot of IFM sizes of *Col-0* (N=59), *clv3-9* (N=22), and *cle40-2* (N=27) plants.

Scale bars: 10mm (B, C, D), 50 μm (B', C', D'), Statistical groups were assigned after calculating p-values by ANOVA and Turkey's multiple comparison test (differential grouping from $p \leq 0.01$). Yellow dotted lines in B' to D' enclose the IFM, red line in the inset meristem in E indicates the area that was used for the quantifications in E.

We next analysed the precise *CLE40* expression pattern using a transcriptional reporter line, *CLE40:Venus-H2B* (Wink, 2013). We first concentrated on the IFMs and FMs. *CLE40* is expressed in IFMs and in FMs, starting at P5 to P6 onwards (Fig. 2A-C). We found stronger expression in the PZ than in the CZ, and no expression in young primordia. Using MorphoGraphX software, we extracted the fluorescence signal originating from the outermost cell layer (L1) of the IFM, and noted reduced *CLE40* expression in the CZ (Fig. 2B). Optical longitudinal sections through the IFM showed that *CLE40* is not expressed in the CZ, and only occasionally in the OC region (Fig. 2C). Expression of *CLE40* changed dynamically during development: expression was concentrated in the IFM, but lacking at sites of primordia initiation (P0 to P4/5, Fig. 2C). In older primordia from P5/6 onwards, *CLE40* expression is detectable from the centre of the young FM and expands towards the FM periphery. In the FMs, *CLE40* is lacking in young sepal primordia (P6), but starts to be expressed on the adaxial sides of petals at P7 (Fig. 2A, P1-P7).

To compare the *CLE40* pattern with that of *CLV3*, we introgressed a *CLV3:NLS-3xmCherry* transcriptional reporter into the *CLE40:Venus-H2B* background. *CLV3* and *CLE40* are expressed in almost mutually exclusive domains of the IFM, with *CLV3* in the CZ surrounded by *CLE40* expressing cells (Fig. 2D-F"). In the deeper region of the IFM, where the OC is located, both *CLV3* and *CLE40* are not expressed (Fig. 2F).

We noted that *CLE40* is downregulated where *WUS* is expressed, or where *WUS* protein localises, such as the OC and CZ. Furthermore, *CLE40* is also lacking in very early flower primordia and in incipient organs.

Fig. 2: *CLE40* and *CLV3* show complementary expression patterns in the IFM

(A) MIP of an inflorescence at 5 WAG expressing the transcriptional reporter *CLE40:Venus-H2B//Col-0* showing *CLE40* expression in the IFM, older primordia and sepals

(N=23). **(B)** The L1 projection shows high expression in the epidermis of the periphery of the IFM and only weak expression in the CZ. **(C)** Longitudinal section through the IFM shows expression of *CLE40* in the periphery, but lack of expression in the CZ. **(P1 –P6)** Longitudinal section through primordia show no *CLE40* expression in young primordia (P1-P4), but in the centre of older primordia (P5-P6). **(D)** The MIP of the double reporter line of *CLE40* and *CLV3* (*CLE40:Venus-H2B;CLV3:NLS-3xmCherry//Col-0*) shows *CLV3* expression in the CZ surrounded by *CLE40* expression in the periphery (N=12). **(E-E’)** The L1 projection shows *CLV3* **(E’)** expression in the centre of the IFM and *CLE40* **(E’)** expression in a distinct complementary pattern in the periphery of the IFM. **(F)** The longitudinal section through the centre of the IFM shows *CLV3* expression in the CZ while *CLE40* **(F’)** is mostly expressed in the surrounding cells. **(F’)** *CLE40* and *CLV3* are expressed in complementary patterns. Dashed blue lines indicate magnified areas, dashed white and orange lines indicate planes of optical sections, dashed yellow line in B marks CZ and in F’ the OC. Scale bars: 50µm (A, D), 20µm (B, C, E, E’, F’), 10µm (P0 to P6), MIP = Maximum intensity projection, PI = Propidium iodide, L1 = visualisation of layer 1 only, P1 to P7 = primordia at consecutive stages.

***CLE40* expression is repressed by *WUS* activity**

To further analyse the regulation of *CLE40* expression, we introduced the *CLE40* transcriptional reporter into the *clv3-9* mutant background (Fig. 3A-B, Fig3-SupplFig.1). In *clv3-9* mutants, *WUS* is no longer repressed by the *CLV* signalling pathway, and the CZ of the meristem increases in size as described previously (Clark et al., 1995). In the *clv3-9* mutant meristems, both *CLV3* and *WUS* promoter activity is now found in an expanded domain (Fig3-SupplFig.1). *CLE40* is not expressed in the tip and centre of the IFM but is rather confined to the peripheral domain, where neither *CLV3* nor *WUS* are expressed (Fig. 3B’, Fig3-SupplFig.1B’). To further explore the expression dynamics of *CLE40* in connection with regulation of stem cell fate and *WUS*, we misexpressed *WUS* from the *CLV3* promoter and introgressed it into plants carrying

the *CLE40:Venus-H2B* construct. Since *WUS* activates the *CLV3* promoter, *CLV3:WUS* misexpression triggers a positive feedback loop. This results in a continuous enlargement of the CZ (Brand et al., 2002). Young seedlings carrying the *CLV3:WUS* transgene at 10 DAG displayed a drastically enlarged SAM, compared to wild type seedlings of the same age (Fig. 3C-D'). Wild type seedlings at this stage express *CLE40* in older leaf primordia and in deeper regions of the vegetative SAM (Fig. 3E-E'). The *CLV3:WUS* transgenic seedlings do not initiate lateral organs from the expanded meristem, and *CLE40* expression is confined to the cotyledons (Fig. 3F-F'). *CLE40* is also lacking in the deeper regions of the vegetative SAM (Fig. 3F'). Thus, we conclude that either *WUS* itself, or a *WUS*-dependent regulatory pathway represses *CLE40* gene expression.

We next determined if *CLE40* repression in the CZ can be alleviated in mutants with reduced *WUS* activity. Since *wus* loss-of-function mutants fail to maintain an active CZ and shoot meristem, we used the hypomorphic *wus-7* allele (Graf et al., 2010; Ma et al., 2019). *wus-7* mutants are developmentally delayed. Furthermore, *wus-7* mutants generate an IFM, but the FMs give rise to sterile flowers that lack inner organs (Fig3-SupplFig.2). We introgressed the *CLE40* reporter into *wus-7*, and found that at 5WAG, all *wus-7* mutants expressed *CLE40* in both the CZ and the OC of the IFM (Fig. 3G-H', Fig3-SupplFig.2). Similar to wild type, *CLE40* is only weakly expressed in the young primordia of *wus-7*. Therefore, we conclude that a *WUS*-dependent pathway downregulates *CLE40* in the centre of the IFM during normal development.

Fig. 3: WUS-dependent repression of CLE40 expression in the shoot meristem

(A) MIP of CLE40 expression (CLE40:Venus-H2B//Col-0) at 5 WAG, **(A')** Optical section through the centre of the IFM (indicated by orange line in (A)) reveals no CLE40 expression in

the CZ and in the centre of the meristem. Cells in the L2 layer also show less *CLE40* expression. High *CLE40* expression is found in the PZ (N=23). **(B)** MIP of *CLE40* expression in a *clv3-9* mutant (*CLE40:Venus-H2B//clv3-9*) shows expression only in the PZ of the meristem, in FMs and in sepals (N=6). **(B')** Optical section through the IFM depicts no *CLE40* expression at the tip and the centre of the meristem. *CLE40* expression is only detected in cells at the flanks of the IFM and in sepals. **(C)** Arabidopsis seedling at 10 DAG. **(D)** Seedling expressing *WUS* from the *CLV3* promoter, 10 DAG. **(D')** Magnification of seedling in (D). The meristem fasciates without forming flowers. **(E)** L1 projection, vegetative seedling with *CLE40* expression in the PZ and in leaf primordia starting from LP4, at 10 DAG (N=5). **(E')** Optical section of (E) with *CLE40* expression primordia and rib meristem or periphery. **(F)** MIP of fasciated meristem as in (D). *CLE40* expression can only be found in the cotyledons (C1 and C2) next to the meristem (N=5). **(F')** Optical section shows *CLE40* expression only in the epidermis of cotyledons. **(G and G')** MIP **(G)** and optical section **(G')** of *CLE40* expression (*CLE40:Venus-H2B//Ler*) in a wild type (*L.er*) background at 5WAG shows no signal in the CZ or OC. *CLE40* is confined to the PZ and the centre of older flower primordia, and to sepals (N=8). **(H and H')** MIP of *CLE40* in a *wus-7* background shows expression through the entire IFM and in the centre of flower primordia. The optical section **(H')** reveals that *CLE40* is also expressed in the CZ as well as in the OC of the IFM (N= 12).

Dashed orange lines indicate the planes of optical sections, dashed blue lines in E and F enclose the meristem, the dashed white line in E' marks the CZ. Scale bars: 50µm (A, B, G, H), 20µm (A', B', E, E', F, F', G', H'), 1mm (C, D), 500µm (D'), MIP = Maximum intensity projection, PI = propidium iodide, L1 = layer 1 projection, C = cotyledon, LP = leaf primordium

CLE40 signals through BAM1

Given that *CLV1* and *BAM1* perform partially redundant functions to perceive *CLV3* in shoot and floral meristems, we asked if these receptors also contribute in a *CLE40* signalling pathway. We therefore generated the translational reporter lines *CLV1:CLV1-GFP* and *BAM1:BAM1-GFP*, and analysed their expression patterns in

detail. We observed dynamic changes of *CLV1* expression during the different stages of flower primordia initiation. *CLV1:CLV1-GFP* is continuously expressed in deeper regions of the IFM comprising the OC, and in the meristem periphery where new FMs are initiating (Fig. 4A). *CLV1* is expressed strongly in cells of the L1 and L2 of incipient organ primordia (P-1, P0), and only in L2 at P1. P2 and P3 show only very faint expression in the L1, but in stages from P4 to P6, *CLV1* expression expands from the L3 into the L2 and L1 (Fig. 4, P1-P6).

Fig. 4: *CLV1* is expressed in the OC and in cells of incipient organ primordia

(A) MIP of *CLV1* under its endogenous promoter (*CLV1:CLV1-GFP/Col-0*) at 5 WAG shows *CLV1* expression in the OC of the meristems, IFM and FMs, in incipient organ primordia (P-1

to P1) and in sepals (N=15). **(B)** In the L1 projection *CLV1* expression is detected in cells of incipient organs. **(C)** Optical section through the IFM shows *CLV1* expression in the OC and in P0. **(P-1-P6)** *CLV1* expression is detected in incipient organ primordia in L1 and L2 (-P1, P0), in the L2 of P1, and in the OC of the IFM and FMs from P4 to P6.

Dashed white and blue lines indicate the planes of optical sections, yellow dashed line in (A) and (B) mark incipient organ primordia (P-1 to P1), yellow lines (P-1 to P6) indicate the IFM region, white lines mark the primordium. Scale bars: 50µm (A), 20µm (B, C), 10µm (P1 to P6), MIP = maximum intensity projection, PI = propidium iodide, L1 = layer 1, P = primordium

The translational *BAM1:BAM1-GFP* reporter is expressed in the IFM, the FMs and in floral organs (Fig. 5A). In the IFM, expression is found throughout the L1 layer of the meristem, and, at an elevated level, in L2 and L3 cells of the PZ, but not in the meristem centre around the OC, where *CLV1* expression is detected (Fig. 5B,C, compare to Fig. 4C). *BAM1* is less expressed in the deeper regions of primordia from P6 onwards (Fig. 5C). *BAM1* transcription was reported to be upregulated in the meristem centre in the absence of *CLV3* or *CLV1* signalling (Nimchuk, 2017). Using our translational *BAM1* reporter in the *clv1-20* mutant background, we confirmed that *BAM1* is expressed in the meristem centre, similar to the pattern of *CLV1* in the wild type, and that *BAM1* is upregulated in the L1 of the meristem. Importantly, in a *clv1-20* background *BAM1* is absent in the peripheral region of the IFM and the L2 (Fig. 5D-F).

Fig. 5: *BAM1* expression is elevated in the flanks of the IFM and not detectable in the OC

(A) MIP of *BAM1* under its endogenous promoter (*BAM1:BAM1-GFP//bam1-3*) at 5 WAG. *BAM1* expression is detected nearly throughout the entire inflorescence (IFM, FM, sepals) with

weak expression in the CZ of IFM and FMs (N=15). **(B)** The L1 projection of the IFM shows ubiquitous expression of *BAM1*. **(C)** Optical section through the IFM shows elevated *BAM1* expression in the flanks (yellow arrows) and a lack of *BAM1* expression in the OC. **(P1 - P6)** *BAM1* expression is found in all primordia cells. **(D)** MIP of *BAM1* in a *clv1-20* mutant (*BAM1:BAM1-GFP//bam1-3;clv1-20*). *BAM1* expression is detected in most parts of the inflorescence, especially in the centre of the IFM and FMs (N=9). **(E)** Cross section (XY) through of the IFM (from D) shows *BAM1* expression in a *clv1-20* mutant in the CZ (IFM and FMs) and in the L1/L2. **(F)** Optical section through the meristem (from D) shows *BAM1* expression in the OC and in the L1, while no *BAM1* expression is detected in the PZ (yellow arrows).

Dashed white and orange lines indicate longitudinal sections; yellow lines (P1 to P6) indicate the IFM region, white lines (P1 to P6) mark the primordium, yellow arrows indicate high (C) or no (F) *BAM1* expression in the PZ. Scale bars: 50µm (A, D), 20µm (B, C, E, F), 10µm (P1 to P6), MIP = maximum intensity projection, PI = propidium iodide, L1 = layer 1, P = primordium

In longitudinal and optical cross sections through the IFM, we found that complementarity of *CLE40* and *CLV3* is reflected in the complementary expression patterns of *BAM1* and *CLV1* (Fig. 6A-D'). Therefore, we conclude that expression patterns of *CLV1* and *BAM1* are mostly complementary in the meristem itself and during primordia development. When comparing *CLE40* and *BAM1* expression patterns, we found a strong overlap in the peripheral zone of the meristem, during incipient primordia formation, in older primordia, and in L3 cells surrounding the OC (Fig. 6A'B', Fig6-SupplFig. 1). Similarly, *CLV3* and *CLV1* are confined to the CZ and OC, respectively.

To analyse if *CLE40*-dependent signalling requires *CLV1* or *BAM1*, we measured the sizes of IFMs in the respective single and double mutants (Fig. 6E, Fig6-SupplFig. 2). While *cle40-2* mutant IFMs reached 65% of the wild type size, *clv1-101*

plants develop IFMs that were 140% wild type size, whereas *bam1-3;clv1-101* double mutant meristems reached 450% wild type size, similar to those of *clv3-9* mutants. This supports the notion that BAM1 can partially compensate for CLV1 function in the CLV3 signalling pathway when expressed in the meristem centre (Fig. 5F) (Nimchuk et al., 2015). The relationship between CLV1 and BAM1 is not symmetrical, since CLV1 is expressed in a wildtypic pattern in *bam1-3* mutants (Fig8-SupplFig. 4). Meristem sizes of *bam1-3* mutants reached 70% of the wild type, and double mutants of *cle40-2;bam1-3* did not differ significantly. However, double mutants of *cle40-2;clv1-101* developed like the *clv1-101* single mutant, indicating an epistatic relationship. Importantly, both *clv1-101* and *bam1-3* mutants lack BAM1 function in the meristem periphery (Fig. 5F), where also *CLE40* is highly expressed, which could explain the observed epistatic relationships of *cle40-2* with both *clv1-101* and *bam1-3*. Similar genetic relationships for *CLV3*, *CLE40*, *CLV1* and *BAM1* were noticed when analysing carpel number as a proxy for FM sizes. We also noted that generation of larger IFMs and FMs in different mutants was negatively correlated with leaf size, which we cannot explain so far (Fig1-SupplFig.1).

We hypothesize that *CLE40* signals from the meristem periphery via BAM1 to promote meristem growth. Next, we aimed to determine if the commonalities between *cle40-2* and *bam1-3* mutants extend beyond their effects on meristem size.

Fig. 6: BAM1 and CLV1 are receptors for CLE40 and CLV3, respectively

(A and A') Longitudinal and cross sections of *CLE40* (*CLE40:Venus-H2B//Col-0*) through the IFM show *CLE40* expression in the PZ while no *CLE40* expression is detected in the CZ or the

OC (dashed yellow line). **(B and B')** In optical sections of *BAM1* (*BAM1:BAM1-GFP//bam1-3*) through the IFM elevated *BAM1* expression in the PZ and in young primordia can be detected, while low expression is found in the CZ and no expression is observed in the OC (dashed yellow line). **(C and C')** Optical and cross section of *CLV3* through the IFM (*CLV3:NLS-3xmCherry//Col-0*) show *CLV3* expression in the CZ (dashed yellow line). **(D and D')** The native expression of *CLV1* (*CLV1:CLV1-GFP//Col-0*) in an optical and cross section through the IFM is depicted in the OC (dashed yellow line) and in cells of the L1 and L2 close to emerging primordia. **(E)** Box and whisker plot of the IFM area size of *Col-0* (N=82), various single (*clv3-9* (N=22), *cle40-2* (N=42), *clv1-101* (N=32), *bam1-3* (N=68)) and double mutants (*cle40-2;clv1-101* (N=37), *cle40-2;bam1-3* (N=25) and *bam1-3;clv1-101* (N=36)) at 6 WAG. Scale bars: 20µm (A – D'), yellow dashed lines indicate the OC (in A', B', D') or the CZ (C'), Statistical groups were assigned after calculating p-values by ANOVA and Turkey's multiple comparison test (differential grouping from $p \leq 0.01$). Red line in the inset meristem in (E) indicates the area that was used for the quantifications in (E).

A CLE40 and BAM1 signalling pathway promotes *WUS* expression in the meristem periphery

We next analysed the number of *WUS*-expressing cells in wild type and mutant meristems using a *WUS:NLS-GFP* transcriptional reporter. Compared to wild type, the *WUS* expression domain was laterally strongly expanded in both *clv3-9* and *clv1-101*. Interestingly, *WUS* signal extended also into the L1 layer of *clv1-101*, albeit in a patchy pattern (Fig. 7A-C',F). Also noteworthy is that *BAM1* was expressed at a higher level in the L1 layer of *clv1* mutants. *cle40-2* mutants showed a reduction in the number of *WUS* expressing cells down to approx. 50% wild-type levels (Fig. 7D-D',F). Importantly, *WUS* remained expressed in the centre of the meristem, but was there found in a narrow domain. In *bam1-3* mutants, the *WUS* domain was similarly reduced as in *cle40-2*, and *WUS* expression focussed in the meristem centre (Fig. 7E,E',F). In

contrast, both *clv3-9* and *clv1-101* mutants express *WUS* in a laterally expanded domain (Fig. 7B',C').

To integrate our finding that *CLE40* expression is repressed by *WUS* activity with the observation that *WUS*, in turn, is promoted by *CLE40* signalling, we hypothesize that the *CLE40-BAM1-WUS* interaction establishes a new negative feedback loop. The *CLE40-BAM1-WUS* negative feedback loop acts in the meristem periphery, while the *CLV3-CLV1-WUS* negative feedback loop acts in the meristem centre along the apical-basal axis. Both pathways act in parallel during development to regulate the size of the *WUS* expression domain in the meristem, possibly by perceiving input signals from two different regions, the CZ and the PZ, of the meristem.

We then asked how the two signalling pathways, converge on the regulation of *WUS* expression, control meristem growth and development. So far, we showed that both *CLV3-CLV1* and *CLE40-BAM1* signalling control meristem size, but in an antagonistic manner. However, we noticed that the different mutations in peptides and receptors affected distinct aspects of meristem shape. We therefore analysed meristem shape by measuring meristem height (the apical-basal axis) at its centre, and meristem diameter (the radial axis) at the base in longitudinal sections. The ratio of height to width then gives a shape parameter " σ " (from the greek word σχήμα = shape). In young inflorescence meristems at 4-5 WAG, when inflorescence stems were approximately 5-8 cm long, meristems of *cle40-2* and *bam1-3* mutants were slightly reduced in width, and strongly reduced in height, resulting in reduced σ in comparison to *Col-0* (Fig. 7G, Fig7-SupplFig. 1A). Meristems of *clv1-101* and *clv3-9* mutants were similar in width to wild type, but strongly increased in height, giving high σ values (Fig. 7, Fig7-SupplFig. 1A-C). This indicates that *CLV3-CLV1* signalling mostly restricts meristem growth along the apical-basal axis, while *CLE40-BAM1* signalling promotes meristem growth along both axes.

Fig. 7: CLE40 and BAM1 promote *WUS* expression

(A – E') MIP and optical section of inflorescences at 5WAG expressing the transcriptional reporter *WUS:NLS-GFP* in a (A and A') *Col-0*, (B and B') *clv1-101*, (C and C') *clv3-9*, (D and D') *cle40-2* and (E and E') *bam1-3* background. In (A) wild type plants the *WUS* domain is smaller compared to the expanded *WUS* domain in (B) *clv1-101* and (C) *clv3-9* mutants. The *WUS* domain of (D) *cle40-2* and (E) *bam1-3* mutants is decreased compared to wild type plants. Optical sections of (B') *clv1-101* and (C') *clv3-9* mutants expand along the basal-apical axis while the meristem shape of (D') *cle40-2* and (E') *bam1-3* mutants are flatter compared to (A') wild type plants. (F) Box and whisker plot shows the number of *WUS*-expressing cells in the OC of IFMs of *Col-0* (N=9), *cle40-2* (N=9), *bam1-3* (N=9), *clv1-101* (N=8), and *clv3-9* (N=5). (G) After 5 WAG *bam1-3* (N=11) and *cle40-2* (N=11) mutants have flatter meristems than wild type plants (decreased σ value compared to *Col-0* (N=11)), while *clv1-101* (N=9) and *clv3-9* (N=6) mutants increase in their IFM height showing a higher σ value.

Scale bars: 50µm (A – E), 20µm (A'-J), Statistical groups and stars were assigned after calculating p-values by ANOVA and Turkey's multiple comparison test (differential grouping from $p \leq 0.01$). WAG = weeks after germination, yellow numbers = *WUS* expressing cells in the CZ, σ value = height/width of IFMs

Our data expand the current model of shoot meristem homeostasis by taking into account that stem cells are lost from the OC during organ initiation in the PZ (Fig. 8). *CLV3* signals from the CZ via *CLV1* in the meristem centre to confine *WUS* expression to the OC. The diffusion of *WUS* protein along the apical-basal axis towards the meristem tip establishes the CZ and activates *CLV3* expression as a feedback signal. During plant growth, rapid cell division activity and organ initiation requires the replenishment of PZ cells from the CZ, which can be mediated by increased *WUS* activity. We now propose that the PZ generates *CLE40* as a short range or autocrine signal that acts through *BAM1* in the meristem periphery. Since *BAM1* and *WUS* expression do not overlap, we postulate the generation of a diffusible factor that relies on *CLE40-BAM1*, and acts from the PZ to promote *WUS* expression. *WUS*, in turn, represses *CLE40* expression from the OC, thus establishing a second negative feedback regulation. Together, the two intertwined pathways serve to adjust *WUS* activity in the OC and incorporate information on the actual size of the stem cell domain, via *CLV3-CLV1*, and the growth requirements from the PZ via *CLE40-BAM1*.

Fig. 8: Schematic model of two intertwined signalling pathways in the shoot meristem

(A and B) Schematic representation of two negative feedback loops in the IFM of *Arabidopsis thaliana*. CLV3 in the CZ binds to the LRR receptor CLV1 to activate a downstream signalling cascade which leads to the repression of the transcription factor WUS. In a negative feedback loop WUS protein moves to the stem cells to activate *CLV3* gene expression. In the PZ of the IFM a second negative feedback loop controls meristem growth by CLE40 and its receptor BAM1. CLE40 binds to BAM1 in an autocrine manner, leading to the activation of a downstream signal “X” which promotes WUS activity. WUS protein in turn represses the expression of the *CLE40* gene.

CZ = central zone, PZ = peripheral zone, arrows indicate a promoting effect and the blocked line indicates a repressing signal

Discussion

Shoot meristems are the centres of growth and organ production throughout the life of a plant. Meristems fulfil two main tasks, which are the maintenance of a non-differentiating stem cell pool, and the assignment of stem cell daughters to lateral organ primordia and differentiation pathways (Hall & Watt, 1989). Shoot meristem homeostasis requires extensive communication between the CZ, the OC and the PZ. The discovery of CLV3 as a signalling peptide, which is secreted exclusively from stem cells in the CZ, and its interaction with WUS in a negative feedback loop was fundamental for our understanding of such communication pathways (Fletcher, 2020). Here, we analysed the function of *CLE40* in shoot development of Arabidopsis, and found that *WUS* expression in the OC is under positive control from the PZ due to the activity of a CLE40-BAM1 signalling pathway. IFM size is reduced in *cle40* mutants, indicating that *CLE40* signalling promotes meristem size. Importantly, *CLE40* is expressed in the PZ, in late stage FMs and in differentiating organs. A common denominator for the complex and dynamic expression pattern is that *CLE40* expression is confined to meristematic tissues, but not in organ founder sites or in regions with high *WUS* activity, such as the OC and the CZ. Both misexpression of *WUS* in the *CLV3* domain (Fig. 3F), studies of *clv3* mutants with expanded stem cell domains (Fig. 3B, Fig3-SupplFig.1) and analysis of *wus* mutants (Fig. 3, Fig3-SupplFig.2) underpinned the notion that *CLE40* expression, in contrast to *CLV3*, is negatively controlled in a *WUS*-dependent manner. Furthermore, we found that the number of *WUS* expressing cells in *cle40* mutant IFMs is strongly reduced, indicating that *CLE40* exerts its positive effects on IFM size by expanding the *WUS* expression domain.

So far, the antagonistic effects of Arabidopsis *CLV3* and *CLE40* on meristem size can only be compared to the antagonistic functions of *MpCLE1* and *MpCLE2* on the gametophytic meristems of *M. polymorpha*, which signal through two distinct

receptors, MpTDR and MpCLV1, respectively (Hata & Kyozyuka, 2021). By the complementation of *clv3* mutants through expression of *CLE40* from the *CLV3* promoter it was shown previously that *CLE40* and *CLV3* are able to activate the same downstream receptors (Hobe et al., 2003). Our detailed analysis of candidate receptor expression patterns showed that *CLV3* and *CLV1* are expressed in partially overlapping domains in the meristem centre, while *CLE40* and *BAM1* are confined to the meristem periphery. Like *cle40* mutants, *bam1* mutant IFMs are smaller and maintain a smaller *WUS* expression domain, supporting the notion that *CLE40* and *BAM1* comprise a signalling unit that increases meristem size by promoting *WUS* expression. The antagonistic functions of the *CLV3-CLV1* and *CLE40-BAM1* pathways in the regulation of *WUS* are reflected in their complementary expression patterns. There is cross-regulation between these two signalling pathways at two levels: (1) *WUS* has been previously shown to promote *CLV3* levels in the CZ, and we here show that *WUS* represses (directly or indirectly) *CLE40* expression in the OC and in the CZ (Fig. 3B, Fig3-SupplFig.1); (2) *CLV1* represses *BAM1* expression in the OC, and thereby restricts *BAM1* to the meristem periphery (Fig. 5, Fig. 6). In *clv1* mutants, *BAM1* shifts from the meristem periphery to the OC, and the *WUS* domain laterally expands in the meristem centre (Fig. 5F, Fig. 7B'). Furthermore, *BAM1* expression increases also in the L1, which could cause the observed irregular expression of *WUS* in the outermost cell layer of *clv1* mutants. The role of *BAM1* in the OC is not entirely clear: despite the high sequence similarity between *CLV1* and *BAM1*, the expression of *BAM1* in the OC is not sufficient to compensate for the loss of *CLV1* (Fig.5D-F, Nimchuk et al., 2015). In the OC, *BAM1* appears to restrict *WUS* expression to some extent, since *clv1;bam1* double mutants reveal a drastically expanded IFM (DeYoung & Clark, 2008). However, it is possible that *BAM1* in the absence of *CLV1* executes a

dual function: to repress *WUS* in response to *CLV3* in the OC as a substitute for *CLV1*, and simultaneously to promote *WUS* expression in the L1 in response to *CLE40*.

The expression domains of *CLE40* and its receptor *BAM1* largely coincide, suggesting that *CLE40* acts as an autocrine signal. Similarly, protophloem sieve element differentiation in roots is inhibited by *CLE45*, which acts as an autocrine signal via *BAM3* (Kang & Hardtke, 2016). Since *WUS* is not expressed in the same cells as *BAM1*, we have to postulate a non-cell autonomous signal X that is generated in the peripheral zone due to *CLE40-BAM1* signalling, and diffuses towards the meristem centre to promote *WUS* expression (Hohm et al., 2010). As a result, *CLE40-BAM1* signalling from the PZ will provide the necessary feedback signal that stimulates stem cell activity and thereby serves to replenish cells in the meristem for the initiation of new organs. The *CLV3-CLV1* signalling pathway then adopts the role of a necessary feedback signal that avoids an excessive stem cell production.

The two intertwined, antagonistically acting signalling pathways that we described here allow us to better understand the regulation of shoot meristem growth, development and shape. The previous model, which focussed mainly on the interaction of the CZ and the OC via the *CLV3-CLV1-WUS* negative feedback regulation, lacked any direct regulatory contribution from the PZ. *EPFL* peptides were shown to be expressed in the periphery and to restrict both *CLV3* and *WUS* expression via *ER* (Zhang et al., 2021). However, *EPFL* peptide expression is not reported to be feedback regulated from the OC or CZ, and the main function of the *EPFL-ER* pathway is therefore to restrict overall meristem size (Zhang et al., 2021). The second negative feedback loop controlled by *CLE40*, which we uncovered here, enables the meristem to fine-tune stem cell activities in response to fluctuating requirements for new cells during organ initiation. Due to the combined activities of *CLV3* and *CLE40*, the OC (with *WUS* as a key player) can now record and compute information from both, the

CZ and PZ. Weaker *CLV3* signalling, indicating a reduction in the size of the CZ, induces preferential growth of the meristem along the apical-basal axis (increasing σ), while weaker *CLE40* signals, reporting a smaller PZ, would decrease σ and flatten meristem shape. It will be intriguing to investigate if different levels of *CLV3* and *CLE40* also contribute to the shape changes that are observed during early vegetative development, or upon floral transition in *Arabidopsis*.

Many shoot-expressed CLE peptides are encoded in the genomes of maize, rice and barley, which could act analogously to *CLV3* and *CLE40* of *Arabidopsis*. It is tempting to speculate that in grasses, a *CLE40*-like, stem cell promoting signalling pathway is more active than a *CLV3*-like, stem cell restricting pathway. This could contribute to the typical shape of cereal SAMs, which are, compared to the dome-shaped SAM of dicotyledonous plants, extended along the apical-basal axis.

Material and Methods

All chemicals used for the experiments are listed in Tab. 1.

Plant material and growth conditions

All wild type *Arabidopsis thaliana* (L.) Heynh. plants used in this study are ecotype Columbia-0 (*Col-0*), except for *wus-7* mutants which are in Landsberg *erecta* (L.er.) background. Details about *Arabidopsis thaliana* plants carrying mutations in the following alleles: *bam1-3*, *cle40-2*, *cle40-cr1*, *cle40-cr2*, *cle40-cr3*, *clv1-101*, *clv3-9* and *wus-7* are described in Tab. 2. All mutants are in *Col-0* background and are assumed to be null-mutants, except for *wus-7* mutants. *cle40* mutants (*cle40-2*, *cle40-cr1*, *cle40-cr2*, *cle40-cr3*) have either a stop codon, a T-DNA insertion or deletion in or before the crucial CLE box domain (Fig1-SupplFig.2B'). *clv3-9* mutants were generated in 2003 by the lab of R. Simon. *clv3-9* mutants were created by EMS resulting in a W62STOP mutation before the critical CLE domain region. *bam1-3* and *clv1-101* mutants have been described as null mutants before (DeYoung et al., 2006; Kinoshita et al., 2010), while *clv1-20* is a weak allele which contains a insertion within the 5'-UTR of CLV1 and results in a reduced mRNA level (Durbak & Tax, 2011). *wus-7* is a weak allele and mutants were described in previous publications (Graf et al., 2010). Double mutants were obtained by crossing the single mutant plants until both mutations were proven to be homozygous for both alleles. Genotyping of the plants was performed either by PCR or dCAPS method with the primers and restrictions enzymes listed in Tab. 3. Before sowing, seeds were either sterilized for 10min in an ethanol solution (80% v/v ethanol, 1,3% w/v sodium hypochloride, 0,02% w/v SDS) or for 1h in a desiccator in a chloric gas atmosphere (50mL of 13% w/v sodium hypochlorite with 1mL 37% HCL). Afterwards, seeds were stratified for 48h at 4°C in darkness. Seeds on soil were then cultivated in phytochambers under long day (LD) conditions (16h light/ 8h dark) at

21°C. For selection of seeds or imaging of vegetative meristems seeds were sowed on ½ Murashige & Skoog (MS) media (1% w/v sucrose, 0.22% w/v MS salts + B5 vitamins, 0.05% w/v MES, 12g/L plant agar, adjusted to pH 5.7 with KOH) in squared petri dishes. Seeds in petri dishes were kept in phytocabinets under continuous light conditions at 21°C and 60% humidity.

Cloning of reporter lines

CLV1 (*CLV1:CLV1-GFP*), BAM1 (*BAM1:BAM1-GFP*) and CLV3 (*CLV3:NLS-3xmCherry*) reporter lines were cloned using the GreenGate method (Lampropoulos et al., 2013). Entry and destination plasmids are listed in Tab. 4 and Tab. 5. Promoter and coding sequences were PCR amplified from genomic *Col-0* DNA which was extracted from rosette leaves of *Col-0* plants. Primers used for amplification of promoters and coding sequences can be found in Tab. 6 with the specific overhangs used for the GreenGate cloning system. Coding sequences were amplified without the stop codon to allow transcription of fluorophores at the C-terminus. BsaI restriction sites were removed by site-directed mutagenesis using the “QuickChange II Kit” following the manufacturer’s instructions (Agilent Technologies). Plasmid DNA amplification was performed by heat-shock transformation into *Escherichia coli* DH5α cells (10min on ice, 1min at 42°C, 1min on ice, 1h shaking at 37°C), which were subsequently plated on selective LB medium (1% w/v tryptone, 0.5% w/v yeast extract, 0.5% w/v NaCl) and cultivated overnight at 37°C. All entry and destination plasmids were validated by restriction digest and Sanger sequencing.

Generation of stable *A. thaliana* lines

Generation of stable *Arabidopsis thaliana* lines was done by using the floral dip method (Clough & Bent, 1998).

619 Translational *CLV1* (*CLV1:CLV1-GFP*) and transcriptional *CLV3*
620 (*CLV3:NLS-3xmCherry*) reporter carry the BASTA plant resistance cassette. T1 seeds
621 were sown on soil and sprayed with Basta ® (120mg/mL) after 5 and 10 DAG. Seeds
622 of ~10 independent Basta-resistant lines were harvested. The translational *BAM1*
623 (*BAM1:BAM1-GFP*) reporter line carries a D-Alanin resistance cassette and T1 seeds
624 were sown on ½ MS media containing 3-4mM D-Alanin. T2 seeds were then selected
625 on ½ MS media supplied with either 3-4mM D-Alanin or 10µg/mL of DL-
626 phosphinothricin (PPT) as a BASTA alternative. Only plants from lines showing about
627 ~75% viability were kept and cultivated under normal plant conditions (21°C, LD). Last,
628 T3 seeds were plated on ½ MS media supplied with 3-4mM D-Alanin or PPT again
629 and plant lines showing 100% viability were kept as homozygous lines. The
630 *CLV3:NLS-3xmCherry* and *CLV1:CLV1-GFP* constructs were transformed into *Col-0*
631 wild type plants and after a stable T3 line was achieved, plants carrying the
632 *CLV1:CLV1-GFP* construct were crossed into *bam1-3*, *cle40-2*, *clv3-9* and *clv1-101*
633 mutants until a homozygous mutant background was reached. *BAM1:BAM1-GFP* lines
634 were floral dipped into *bam1-3* mutants and subsequently crossed into the *clv1-20*
635 mutant background which rescued the extremely fasciated meristem phenotype of
636 *bam1-3;clv1-20* double mutants (Fig. 5D-F). *BAM1:BAM1-GFP//bam1-3* plants were
637 also crossed into *cle40-2* and *clv3-9* mutants until a homozygous mutant background
638 was achieved. The *CLE40:Venus-H2B* reporter line was created and described in
639 Wink, 2013 and the *WUS:NLS-GFP;CLV3:NLS-mCherry* reporter line was a gift from
640 the Lohmann lab (Wink, 2013). *CLE40:Venus-H2B* reporter line was crossed into
641 homozygous *clv3-9* and heterozygous *wus-7* mutants. Homozygous *clv3-9* mutants
642 were detected by its obvious phenotype and were brought into a stable F3 generation.
643 Homozygous *wus-7* mutants were genotyped. Seeds were kept in the F2 generation,
644 since homozygous *wus-7* plants do not develop seeds. The *CLE40:Venus-H2B*

reporter line was also crossed with the *CLV3:NLS-3xmCherry* reporter line and was brought into a stable F3 generation. To generate the *CLE40:Venus-H2B//CLV3:WUS* line, plants carrying the *CLE40:Venus-H2B* line were floral dipped with the *CLV3:WUS* construct. T1 seeds were sown on 10µg/mL of DL-phosphinothricin (PPT) and the viable seedlings were imaged. *WUS:NLS-GFP/CLV3:NLS-mCherry//Col-0* reporter line was crossed into *clv3-9*, *cle40-2*, *clv1-101* and *bam1-3* mutants until a stable homozygous F3 generation was reached respectively. Detailed information of all used *A. thaliana* lines can be found in Tab. 7.

Confocal imaging of IFMs

To image IFMs *in vivo*, plants were grown under LD (16h light/ 8h dark) conditions and inflorescences were cut off at 5 or 6 WAG. Inflorescences were stuck on double sided adhesive tape on an objective slide and dissected until only the meristem and primordia from P0 to maximum P10 were visible. Next, inflorescences were stained with either propidium iodide (PI 5mM) or 4',6-Diamidin-2-phenylindol (DAPI 1µg/mL) for 2 to 5min. Inflorescences were then washed three times with water and subsequently covered with water and a cover slide and placed under the microscope. Imaging was performed with a Zeiss LSM780 or LSM880 using a W Plan-Apochromat 40x/1.2 objective. Laser excitation, emission detection range and detector information for fluorophores and staining can be found in Tab. 8. All IFMs were imaged from the top taking XY images along the Z axis, resulting in a Z-stack through the inflorescence. The vegetative meristems were imaged as described for IFMs. Live imaging of the reporter lines in *A. thaliana* plants was performed by dissecting primary inflorescences (except for *clv3-9* mutants) at 5 WAG under LD conditions. For imaging of the reporter lines in the mutant backgrounds of *clv3-9* secondary IFMs were dissected, since the primary meristems are highly fasciated. Vegetative meristems were cultivated in

continuous light conditions at 21°C on ½ MS media plates and were imaged at 10 DAG. For each reporter line at least 3 independent experiments were performed and at least 5 IFMs were imaged.

Phenotyping of *CLV* mutants

For meristem measurements (area size, width and height) primary and secondary IFMs of wild type (*Col-0*) and mutant plants (*cle40-2*, *cle40-cr1-3*, *bam1-3*, *cle40-2;bam1-3*, *clv1-101*) were dissected at 6 WAG under LD conditions. For *clv3-9* and *clv1-101;bam1-3* only secondary IFMs were imaged and analyzed, due to the highly fasciated primary meristems. Optical sections of the Z-stacks were performed through the middle of the meristem starting in the centre of primordia P5 and ending in the centre of primordia P4. Based on the optical sections (XZ), meristem height and area size were measured as indicated in Fig6-SupplFig. 2. IFM sizes from Fig. 1E are also used in Fig. 6E for *Col-0*, *cle40-2* and *clv3-9* plants.

Same procedure was used to count the cells expressing *WUS* in different mutant backgrounds (Fig. 7A-E). Optical sections of IFMs at 5 WAG were performed from P4 to P5 and only nuclei within the meristem area were counted and plotted. For analyses of carpel numbers, the oldest 10 - 15 siliques per plant at 5 WAG were used. Each carpel was counted as one, independent of its size. N number depicts number of siliques. Leaf measurements were performed at 4 WAG and four leaves of each plant were measured and plotted. Data was obtained from at least 3 independent experiments.

Data analysis

For visualization of images the open-source software ImageJ v 1.53c (Schneider et al., 2012) was used. All images were adjusted in “Brightness and Contrast”. IFMs in Fig. 7

were imaged with identical microscopy settings (except for *c/v3-9* mutants) and were all changed in “Brightness and Contrast” with the same parameters to ensure comparability. *c/v3-9* mutants were imaged with a higher laser power since meristems are highly fasciated. MIPs were created by using the “Z-Projection” function and optical sections were performed with the “Reslice...” function resulting in the XZ view of the image. Meristem width, height and area size were measured with the “Straight line” for width and height and the “Polygon selection” for area size. The shape parameter σ was calculated by the quotient of height and width from each IFM. For L1 visualization the open-source software MorphoGraphX (<https://www.mpipz.mpg.de/MorphoGraphX/>) was used that was developed by Richard Smith. 2½ D images were created by following the steps in the MorphoGraphX manual (de Reuille et al., 2015). After both channels (PI and fluorophore signal) were projected to the created mesh, both images were merged using ImageJ v 1.53c. For all statistical analyses, GraphPad Prism v8.0.0.224 was used. Statistical groups were assigned after calculating p-values by ANOVA and Turkey’s or Dunett’s multiple comparison test (differential grouping from $p \leq 0.01$) as indicated under each figure. Same letters indicate no statistical differences. All plasmid maps and cloning strategies were created and planned using the software VectorNTI®.

716 Tab. 1: Chemicals and substances used in this study.

Name	Producer	Product no.	CAS no.
BASTA® non-selective herbicide	Bayer CropScience	84442615	N/A
bacto™ agar	gibco	214010	9105960
bacto™ yeast extract	gibco	212750	9070604
Carbenicillin disodium salt	Carl Roth	6344.2	4800-94-6
DAPI (4,6-diamidino-2-phenylindole)	N/A	N/A	28718-90-3
DL-phosphinothricin (PPT)	Duchefa Biochemie bv	P0159	77182-82-2
D-Alanin	Sigma-Aldrich (Merck)	A7377	338-69-2
D(+)-Saccharose	Carl Roth	4661.1	N/A
Gentamicin sulfate	Sigma-Aldrich (Merck)	G1264	1405-41-0
Hygromycin B	Duchefa Biochemie bv	H0192	31282-04-9
Hypochloride acid (~37%)	Thermo Fischer Scientific	H/120/PB15	1884567
Kanamycin monosulfate	Duchefa Biochemie bv	K0126.0005	25389-94-0
Magnesium chloride x 6H ₂ O	Grüssig GmbH	12087	205
MES hydrate	Sigma-Aldrich (Merck)	10240885	1266615-59-1
Murashige & Skoog (+ Gamborg B5 vitamins)	Duchefa Biochemie bv	M0231.0050	N/A
Phusion High-Fidelity PCR polymerase	Thermo Fischer Scientific	F530S	N/A
Plant agar	Duchefa Biochemie bv	P1001.1000	9002-18-0
Potassium hydroxide	Sigma-Aldrich (Merck)	9643807	N/A
Propidium iodide	Thermo Fischer Scientific	P1304MP	25535-16-4
Rifampicin	TCI	R0079	13292-46-1
Spectinomycin HCl pentahydrate	Duchefa Biochemie bv	S0188	22189-32-8
Sodium dodecyl sulfate (SDS)	Sigma-Aldrich (Merck)	L3771	151-21-3

Sodium chloride	Carl Roth	3957.1	N/A
Sodium hypochloride (13%)	Zentrale Chemikalienlager (ZCL)	2N370	N/A
Tetracycline	Sigma-Aldrich (Merck)	87128	60-54-8
Tryptone	gibco	N/A	57091

Tab. 2: Mutants analysed in this study.

Allele	Gene	Mutation	Reference	Background
<i>bam1-3</i>	AT5G65700	T-DNA	Alonso et al., 2003; SALK_015302	<i>Col-0</i>
<i>cle40-2</i>	AT5G12990	Transposon mutation	Stahl et al., 2009	<i>Col-0</i>
<i>cle40-cr1</i> <i>cle40-cr2</i> <i>cle40-cr3</i>	AT5G12990	CRISPR	Yamaguchi et al., 2017	<i>Col-0</i>
<i>clv3-9</i>	AT2G27250	EMS	Rüdiger Simon, 2003	<i>Col-0</i>
<i>clv1-20</i>	AT1G75820	T-DNA	SALK_008670	<i>Col-0</i>
<i>clv1-101</i>	AT1G75820	T-DNA	Kinoshita et al., 2010; CS858348	<i>Col-0</i>
<i>wus-7</i>	AT2G17950	EMS	Graf et al., 2010	<i>L.er.</i>

Tab. 3: Primers and methods used for genotyping.

Allele	Method	Primer	PCR product
<i>bam1-3</i>	PCR	bam1-3_F: GGAGCTAATTGCGGATTAACC bam1-3_R: GGAACCTAAACCGGAGAGGTTG Lbb1.3_R: ATTTTGCCGATTTCGGAAC	WT amp. : 1208 bp mutant amp. : 998 bp
<i>cle40-2</i>	dCAPS	cle40-2_F: GGAGAAACACAAGATACGAAAGCCATG cle40-2_R: ATTGTGATTTGATACCAACTTAAAA	Restriction enzyme: AseI WT amp. : 460 + 200 bp mutant amp. : 410 + 200 + 60 bp
<i>cle40-cr1</i> <i>cle40-cr2</i> <i>cle40-cr3</i>	dCAPS	cle40-cr_F: ATGGCGGCGATGAAATACAA cle40-cr_R: GTTACGCTTTGGCATCTTTCC	Restriction enzyme: BamHI WT amp.: 750 bp mutant amp. : 491 + 259 bp
<i>clv1-20</i>	PCR	clv1-20_F: TTTGAATAGTGTGTGACCAAATTTGA clv1-20_R: TCCAATGGTAATTCACCGGTG LBa.1: TGGTTCACGTAGTGGGCCATCG	WT amp.: 860bp mutant amp: 1200bp
<i>clv1-101</i>	PCR	clv1-101_F: TTCTCCAAATTCACCAACAGG clv1-101_R: CAACGGAGAAATCCCTAAAGG WiscLox_LT6_R: AATAGCCTTTACTTGAGTTGGCGTAAAAG	WT amp. : 1158 bp mutant amp. : 896 bp
<i>wus-7</i>	dCAPS	wus-7_F: CCGACCAAGAAAGCGGCAACA wus-7_R: AGACGTTCTTGCCCTGAATCTTT	Restriction enzyme: XmnI WT amplification: 216 bp mutant amp. : 193 + 23 bp

725 Tab. 4: Entry vectors used for cloning.

Name	Description	Bacterial resistance	Backbone	Reference/ Origin
proBAM1 (pGD288)	BAM1 promoter 3522bp upstream from transcription start	Ampicillin	pGGA000	Grégoire Denay
proCLV3	CLV3 promoter 1480bp upstream from transcription start	Ampicillin	pGGA000	Jenia Schlegel
proCLV1	CLV1 promoter 5759bp upstream from transcription start	Ampicillin	pGGA000	Patrick Schultz
omega-element (pGGB002)	Omega- element	Ampicillin	pGGB000	Lampropoulos et al., 2013
SV40 NLS (pGGB005)	SV40 NLS (SIMIAN VIRUS 40 NUCLEAR LOCALIZATION SIGNAL)	Ampicillin	pGGB000	Lampropoulos et al., 2013
BAM1_CDS (pGD351)	BAM1 coding region genomic region of BAM1 START to one codon before STOP, including introns, internal BsaI sites removed	Ampicillin	pGGC000	Grégoire Denay
CLV1_CDS	CLV1 coding region 2946 bp coding region amplified from genomic Col-0 DNA without STOP codon and internal BsaI site removed	Ampicillin	pGGC000	Jenia Schlegel
3x-mCherry (pGGC026)	3x mCherry	Ampicillin	pGGC000	Lampropoulos et al., 2013
linker-GFP (pGD165)	linker(10aa)-eGFP	Ampicillin	pGGD000	Grégoire Denay
d-dummy (pGGD002)	d-dummy	Ampicillin	pGGD000	Lampropoulos et al., 2013
tCLV3	CLV3 terminator 1257 bp downstream of transcription stop	Ampicillin	pGGE000	Jenia Schlegel
tUBQ10 (pGGE009)	UBQ10 terminator	Ampicillin	pGGE000	Lampropoulos et al., 2013
BastaR (pGGF008)	pNOS:BastaR (chi sequence removed):tNOS	Ampicillin	pGGF000	Lampropoulos et al., 2013
D-AlaR (pGGF003)	pMAS:D-AlaR:tMAS	Ampicillin	pGGF000	Lampropoulos et al., 2013

726

727

Tab. 5: Destination vectors used to generate transgenic *A.thaliana* reporter lines.

Name	Backbone	Promoter	N-tag	CDS	C-tag	Terminator	Resistance
<i>BAM1:BAM1-GFP</i>	pGGZ001	proBAM1	Ω- element (pGGB002)	BAM1-CDS	linker-GFP (pGD165)	tUBQ10 (pGGE009)	D-Alanin (pGGF003)
<i>CLV1:CLV1-GFP</i>	pGGZ001	proCLV1	Ω- element (pGGB002)	CLV1-CDS	linker-GFP (pGD165)	tUBQ10 (pGGE009)	BastaR (pGGF008)
<i>CLV3:NLS-3xmCherry</i>	pGGZ001	proCLV3	SV40 NLS	3x-mCherry (pGGC026)	d-dummy (pGGD002)	tCLV3	BastaR (pGGF008)

Tab. 6: Primers used for cloning the entry vectors.

Name	Primer
proBAM1 (pGD288)	F: AAAGGTCTCAACCTATGATCCGATCCTCAAAAGTATGTA R: AAAGGTCTCATGTTTCTCTCTATCTCTCTTGTGTG
BAM1_CDS (pGD351)	F: TTTGGTCTCAGGCTCTATGAACTTTTCTTCTCCTTC R: TTTGGTCTCACTGATAGATTGAGTAGATCCGGC Bsal-site_#1_F: CTTGATCTCTCCGGA CTCAACCTCTCCGG Bsal-site_#1_R: CCGGAGAGGTTGAGTCCGGAGAGATCAAG Bsal-site_#2_F: CTCATGTTGCTGACTTTGGACTCGCTAAATTCCTTCAAG Bsal-site_#2_R: CTTGAAGGAATTTAGCGAGTCCAAAGTCAGCAACATGAG
proCLV1	F: AAAGGTCTCAACCTGACTATTGTTTATACTTAGTTG R: TTTGGTCTCATGTTTCAATTTTTTAGTGTCTC
CLV1_CDS	F: AAAGGTCTCAGGCTTAATGGCGATGAGAC R: TTTGGTCTCACTGAACGCGATCAAGTTC Bsl-site_#1_F: CTAAAGGACACGGACTGCACGACTG Bsl-site_#1_R: CAGTCGTGCAGTCCGTGTCCTTTAG Bsl-site_#2_F: CTTAGAGTATCTTGGACTGAACGGAGCTGG Bsl-site_#2_R: CCAGCTCCGTTTCACTCAAGATACTCTAAG
proCLV3	F: AAAGGTCTCAACCTCGGATTATCCATAATAAAAAAC R: AAAGGTCTCATGTTTTTTAGAGAGAAAGTGACTGAG
tCLV3	F: TTTGGTCTCTCTGCCGCCCTAATCTCTTGT R: TTTGGTCTCGTGATATGTGTGTTTTTCTAAACAATC

Tab. 7: Arabidopsis lines that were analysed in this study.

Name/Construct	Background	Plant resistance	Generation	Reference
<i>BAM1:BAM1-GFP</i>	<i>bam1-3</i>	D-Ala	T4	this study
<i>BAM1:BAM1-GFP</i>	<i>bam1-3;clv1-20</i>	D-Ala	F3	this study
<i>BAM1:BAM1-GFP</i>	<i>bam1-3;clv3-9</i>	D-Ala	F3	this study
<i>BAM1:BAM1-GFP</i>	<i>bam1-3;cle40-2</i>	D-Ala	F3	this study
<i>CLE40:Venus-H2B</i>	<i>Col-0</i>	Hygromycin	T5	Rene Wink
<i>CLE40:Venus-H2B</i>	<i>clv3-9</i>	Hygromycin	F3	this study
<i>CLE40:Venus-H2B</i>	<i>wus-7</i>	Hygromycin	F2	this study
<i>CLE40:Venus-H2B</i>	<i>CLV3:WUS//Col-0</i>	Hygromycin/ Basta	T1*	this study
<i>CLV1:CLV1-GFP</i>	<i>Col-0</i>	Basta	T4	this study
<i>CLV1:CLV1-GFP</i>	<i>bam1-3</i>	Basta	F3	this study
<i>CLV1:CLV1-GFP</i>	<i>clv3-9</i>	Basta	F3	this study
<i>CLV1:CLV1-GFP</i>	<i>cle40-2</i>	Basta	F3	this study
<i>CLV3:NLS-3xmCherry</i>	<i>CLE40:Venus-H2B//Col-0</i>	Basta/ Hygromycin	F3	this study
<i>CLV3:NLS-mCherry</i> <i>WUS:NLS-GFP</i>	<i>Col-0</i>	Kanamycin	N/A	Anne Pfeiffer
<i>CLV3:NLS-mCherry</i> <i>WUS:NLS-GFP</i>	<i>cle40-2</i>	Kanamycin	F3	this study
<i>CLV3:NLS-mCherry</i> <i>WUS:NLS-GFP</i>	<i>bam1-3</i>	Kanamycin	F3	this study
<i>CLV3:NLS-mCherry</i> <i>WUS:NLS-GFP</i>	<i>clv1-101</i>	Kanamycin	F3	this study
<i>CLV3:NLS-mCherry</i> <i>WUS:NLS-GFP</i>	<i>clv3-9</i>	Kanamycin	F3	this study

* Plants do not overcome seedling stage

Tab. 8: Microscopy settings used for imaging.

Fluorophore/ Staining	Excitation	Emission	MBS	Detector	Light source
DAPI	405 nm	410 - 490 nm	405	PMT*	Diode
GFP	488 nm	500 - 545 nm	488/561	GaAsP	Argon laser
Venus	514 nm	518 - 540 nm	458/514	GaAsP	Argon laser
mCherry	561 nm	570 - 640 nm	458/561	PMT*	DPSS laser**
PI	561 nm	595 - 650 nm	488/561	PMT*	DPSS laser**

* Photomultiplier tubes

** Diode-pumped solid state

Acknowledgements

This study was funded by DFG through iGrad-Plant (IRTG 2466), CRC 1208 and CEPLAS (EXC 2048). We thank Cornelia Gieseler, Silke Winters, and Carin Theres for technical support and Yasuka L. Yamaguchi (Sawa lab), Anne Pfeiffer (Lohmann lab) and Rene Wink (Simon lab) for sharing Arabidopsis seeds. We also thank Vicky Howe for proof reading the manuscript, the Center for Advanced imaging (CAi) at HHU for microscopy support and Aleksandra Sapala for support with MorphoGraphX.

Author contributions

J.S., G.D., Y.S. and R.S. designed and planned the experiments. J.S. performed experiments and data analysis, besides counting carpels (Fig1-SupplFig.3), which was performed by J. Schmidt. G.D., K.G.P. and J.S. generated stable Arabidopsis lines. Y.S. provided material. G.D., P.B. and J.S. performed the cloning. J.S. and R.S wrote the manuscript with input from all authors.

Declaration of Interests

The authors declare no competing interests.

References

- Berckmans, B., Kirschner, G., Gerlitz, N., Stadler, R., & Simon, R. (2020). CLE40 signaling regulates root stem cell fate. *Plant Physiology*, 182(4), 1776–1792. <https://doi.org/10.1104/PP.19.00914>
- Betsuyaku, S., Takahashi, F., Kinoshita, A., Miwa, H., Shinozaki, K., Fukuda, H., & Sawa, S. (2011). Mitogen-activated protein kinase regulated by the CLAVATA receptors contributes to shoot apical meristem homeostasis. *Plant and Cell Physiology*, 52(1), 14–29. <https://doi.org/10.1093/pcp/pcq157>
- Bleckmann, A., Weidtkamp-Peters, S., Seidel, C. a M., & Simon, R. (2010). Stem cell signaling in Arabidopsis requires CRN to localize CLV2 to the plasma membrane. *Plant Physiology*, 152(1), 166–176. <https://doi.org/10.1104/pp.109.149930>
- Blümke, P., Schlegel, J., Gonzalez-Ferrer, Carmen Becher, S., Gustavo Pinto, K., Monaghan, J., & Simon, R. (2021). Receptor-like cytoplasmic kinase MAZZA mediates developmental processes with CLAVATA1-family receptors in Arabidopsis. *Journal of Experimental Botany*.
- Bommert, P., Je, B. Il, Goldshmidt, A., & Jackson, D. (2013). The maize Gα gene COMPACT PLANT2 functions in CLAVATA signalling to control shoot meristem size. *Nature*, 502(7472), 555–558. <https://doi.org/10.1038/nature12583>
- Brand, U., Fletcher, J. C., Hobe, M., Meyerowitz, E. M., & Simon, R. (2000). Dependence of stem cell fate in Arabidopsis on a feedback loop regulated by CLV3 activity. *Science*, 289(5479), 617–619. <https://doi.org/10.1126/science.289.5479.617>
- Brand, U., Grünewald, M., Hobe, M., & Simon, R. (2002). Regulation of CLV3 expression by two homeobox genes in Arabidopsis. *Plant Physiology*, 129(2), 565–575. <https://doi.org/10.1104/pp.001867>
- Clark, S. E., Running, M. P., & Meyerowitz, E. M. (1993). CLAVATA1, a regulator of meristem and flower development in Arabidopsis. *Development*, 119, 397–418.
- Clark, S. E., Running, M. P., & Meyerowitz, E. M. (1995). CLAVATA3 is a specific regulator of shoot and floral meristem development affecting the same processes as CLAVATA1. *Development*, 121(7), 2057–2067.
- Clark, S. E., Williams, R. W., & Meyerowitz, E. M. (1997). The CLAVATA1 Gene Encodes a Putative Receptor Kinase That Controls Shoot and Floral Meristem Size in Arabidopsis. *Cell*, 89, 575–585.

- 793 Clough, S. J., & Bent, A. F. (1998). Floral dip: A simplified method for *Agrobacterium*-
794 mediated transformation of *Arabidopsis thaliana*. *Plant Journal*, 16(6), 735–743.
795 <https://doi.org/10.1046/j.1365-3113X.1998.00343.x>
- 796 Crook, A. D., Willoughby, A. C., Hazak, O., Okuda, S., Van Der Molen, K. R., Soyars,
797 C. L., Cattaneo, P., Clark, N. M., Sozzani, R., Hothorn, M., Hardtke, C. S., &
798 Nimchuk, Z. L. (2020). BAM1/2 receptor kinase signaling drives CLE peptide-
799 mediated formative cell divisions in *Arabidopsis* roots. *Proceedings of the*
800 *National Academy of Sciences of the United States of America*, 117(51), 32750–
801 32756. <https://doi.org/10.1073/pnas.2018565117>
- 802 Cui, Y., Hu, C., Zhu, Y., Cheng, K., Li, X., Wei, Z., Xue, L., Lin, F., Shi, H., Yi, J.,
803 Hou, S., He, K., Li, J., & Gou, X. (2018). Cik receptor kinases determine cell fate
804 specification during early anther development in *Arabidopsis*. *Plant Cell*, 30(10),
805 2383–2401. <https://doi.org/10.1105/tpc.17.00586>
- 806 Daum, G., Medzihradsky, A., Suzuki, T., & Lohmann, J. U. (2014). A mechanistic
807 framework for noncell autonomous stem cell induction in *Arabidopsis*.
808 *Proceedings of the National Academy of Sciences of the United States of*
809 *America*, 111(40), 14619–14624. <https://doi.org/10.1073/pnas.1406446111>
- 810 de Keijzer, J., Rios, A. F., & Willemsen, V. (2021). *Physcomitrium patens*: A single
811 model to study oriented cell divisions in 1d to 3d patterning. *International Journal*
812 *of Molecular Sciences*, 22(5), 1–16. <https://doi.org/10.3390/ijms22052626>
- 813 de Reuille, P. B., Routier-Kierzkowska, A. L., Kierzkowski, D., Bassel, G. W.,
814 Schüpbach, T., Tauriello, G., Bajpai, N., Strauss, S., Weber, A., Kiss, A., Burian,
815 A., Hofhuis, H., Sapala, A., Lipowczan, M., Heimlicher, M. B., Robinson, S.,
816 Bayer, E. M., Basler, K., Koumoutsakos, P., ... Smith, R. S. (2015).
817 MorphoGraphX: A platform for quantifying morphogenesis in 4D. *ELife*, 4, 1–20.
818 <https://doi.org/10.7554/eLife.05864>
- 819 Defalco, T. A., Anne, P., James, S. R., Willoughby, A., Johannndrees, O., Genolet, Y.,
820 Pullen, A.-M., Zipfel, C., Hardtke, C. S., & Nimchuk, Z. L. (2021). A conserved
821 regulatory module regulates receptor kinase signaling in immunity and
822 development. *BioRxiv*, 2021.01.19.427293.
823 <https://doi.org/10.1101/2021.01.19.427293>
- 824 DeYoung, B. J., Bickle, K. L., Schrage, K. J., Muskett, P., Patel, K., & Clark, S. E.
825 (2006). The CLAVATA1-related BAM1, BAM2 and BAM3 receptor kinase-like
826 proteins are required for meristem function in *Arabidopsis*. *Plant Journal*, 45(1),

- 1–16. <https://doi.org/10.1111/j.1365-313X.2005.02592.x>
- DeYoung, B. J., & Clark, S. E. (2008). BAM receptors regulate stem cell specification and organ development through complex interactions with CLAVATA signaling. *Genetics*, 180(2), 895–904. <https://doi.org/10.1534/genetics.108.091108>
- Durbak, A. R., & Tax, F. E. (2011). CLAVATA signaling pathway receptors of arabidopsis regulate cell proliferation in fruit organ formation as well as in meristems. *Genetics*, 189(1), 177–194. <https://doi.org/10.1534/genetics.111.130930>
- Endrizzi, K., Moussian, B., Haecker, A., Levin, J. Z., & Laux, T. (1996). The SHOOT MERISTEMLESS gene is required for maintenance of undifferentiated cells in Arabidopsis shoot and floral meristems and acts at a different regulatory level than the meristem genes WUSCHEL and ZWILLE. *The Plant Journal*, 10, 967–979.
- Fletcher, J. C. (2020). Recent Advances in Arabidopsis CLE Peptide Signaling. *Trends in Plant Science*, 25(10), 1005–1016. <https://doi.org/10.1016/j.tplants.2020.04.014>
- Fletcher, J. C., Brand, U., Running, M. P., Simon, R., & Meyerowitz, E. M. (1999). Signaling of cell fate decisions by CLAVATA3 in Arabidopsis shoot meristems. *Science*, 283(5409), 1911–1914. <https://doi.org/10.1126/science.283.5409.1911>
- Goad, D. M., Zhu, C., & Kellogg, E. A. (2017). Comprehensive identification and clustering of CLV3/ESR-related (CLE) genes in plants finds groups with potentially shared function. *New Phytologist*, 216(2), 605–616. <https://doi.org/10.1111/nph.14348>
- Graf, P., Dolzblasz, A., Würschum, T., Lenhard, M., Pfreundt, U., & Laux, T. (2010). MGOUN1 encodes an Arabidopsis type IB DNA topoisomerase required in stem cell regulation and to maintain developmentally regulated gene silencing. *Plant Cell*, 22(3), 716–728. <https://doi.org/10.1105/tpc.109.068296>
- Hall, P., & Watt, F. (1989). Stem cells: the generation and maintenance of cellular diversity. *Development*, 106(4), 619–633. <http://dev.biologists.org/content/106/4/619.abstract>
- Han, H., Geng, Y., Guo, L., Yan, A., Meyerowitz, E. M., Liu, X., & Zhou, Y. (2020). The Overlapping and Distinct Roles of HAM Family Genes in Arabidopsis Shoot Meristems. *Frontiers in Plant Science*, 11(541968). <https://doi.org/10.3389/fpls.2020.541968>

- 861 Harrison, C. J., Roeder, A. H. K., Meyerowitz, E. M., & Langdale, J. A. (2009). Local
862 Cues and Asymmetric Cell Divisions Underpin Body Plan Transitions in the
863 Moss *Physcomitrella patens*. *Current Biology*, 19(6), 461–471.
864 <https://doi.org/10.1016/j.cub.2009.02.050>
- 865 Hata, Y., & Kyoizuka, J. (2021). Fundamental mechanisms of the stem cell regulation
866 in land plants: lesson from shoot apical cells in bryophytes. *Plant Molecular*
867 *Biology*. <https://doi.org/10.1007/s11103-021-01126-y>
- 868 Hirakawa, Y., Fujimoto, T., Ishida, S., Uchida, N., Sawa, S., Kiyosue, T., Ishizaki, K.,
869 Nishihama, R., Kohchi, T., & Bowman, J. L. (2020). Induction of Multichotomous
870 Branching by CLAVATA Peptide in *Marchantia polymorpha*. *Current Biology*,
871 30(19), 3833–3840.e4. <https://doi.org/10.1016/j.cub.2020.07.016>
- 872 Hirakawa, Y., Uchida, N., Yamaguchi, Y. L., Tabata, R., Ishida, S., Ishizaki, K.,
873 Nishihama, R., Kohchi, T., Sawa, S., & Bowman, J. L. (2019). Control of
874 proliferation in the haploid meristem by CLE peptide signaling in *marchantia*
875 *polymorpha*. *PLoS Genetics*, 15(3), 1–20.
876 <https://doi.org/10.1371/journal.pgen.1007997>
- 877 Hobe, M., Müller, R., Grünewald, M., Brand, U., & Simon, R. (2003). Loss of CLE40,
878 a protein functionally equivalent to the stem cell restricting signal CLV3,
879 enhances root waving in *Arabidopsis*. *Development Genes and Evolution*,
880 213(8), 371–381. <https://doi.org/10.1007/s00427-003-0329-5>
- 881 Hohm, T., Zitzler, E., & Simon, R. (2010). A dynamic model for stem cell homeostasis
882 and patterning in *Arabidopsis* meristems. *PLoS ONE*, 5(2), 1–9.
883 <https://doi.org/10.1371/journal.pone.0009189>
- 884 Hord, C. L. H., Chen, C., DeYoung, B. J., Clark, S. E., & Ma, H. (2006). The
885 BAM1/BAM2 receptor-like kinases are important regulators of *Arabidopsis* early
886 anther development. *Plant Cell*, 18(7), 1667–1680.
887 <https://doi.org/10.1105/tpc.105.036871>
- 888 Ishida, T., Tabata, R., Yamada, M., Aida, M., Mitsumasu, K., Fujiwara, M.,
889 Yamaguchi, K., Shigenobu, S., Higuchi, M., Tsuji, H., Shimamoto, K., Hasebe,
890 M., Fukuda, H., & Sawa, S. (2014). Heterotrimeric G proteins control stem cell
891 proliferation through CLAVATA signaling in *Arabidopsis*. *EMBO Reports*, 15(11),
892 1202–1209. <https://doi.org/10.15252/embr.201678010>
- 893 Je, B. II, Gruel, J., Lee, Y. K., Bommert, P., Arevalo, E. D., Eveland, A. L., Wu, Q.,
894 Goldshmidt, A., Meeley, R., Bartlett, M., Komatsu, M., Sakai, H., Jönsson, H., &

- Jackson, D. (2016). Signaling from maize organ primordia via FASCIATED
EAR3 regulates stem cell proliferation and yield traits. *Nature Genetics*, 48(7),
785–791. <https://doi.org/10.1038/ng.3567>
- Jeong, S., Trotochaud, A. E., & Clark, S. E. (1999). The Arabidopsis CLAVATA2
gene encodes a receptor-like protein required for the stability of the CLAVATA1
receptor-like kinase. *Plant Cell*, 11(10), 1925–1933.
<https://doi.org/10.1105/tpc.11.10.1925>
- Kang, Y. H., & Hardtke, C. S. (2016). Arabidopsis MAKR 5 is a positive effector of
BAM3-dependent CLE45 signaling. *EMBO Reports*, 17(8), 1145–1154.
<https://doi.org/10.15252/embr.201642450>
- Kinoshita, A., Betsuyaku, S., Osakabe, Y., Mizuno, S., Nagawa, S., Stahl, Y., Simon,
R., Yamaguchi-Shinozaki, K., Fukuda, H., & Sawa, S. (2010). RPK2 is an
essential receptor-like kinase that transmits the CLV3 signal in Arabidopsis.
Development, 137(24), 4327–4327. <https://doi.org/10.1242/dev.061747>
- Lampropoulos, A., Sutikovic, Z., Wenzl, C., Maegele, I., Lohmann, J. U., & Forner, J.
(2013). GreenGate - A novel, versatile, and efficient cloning system for plant
transgenesis. *PLoS ONE*, 8(12). <https://doi.org/10.1371/journal.pone.0083043>
- Laux, T., Mayer, K. F. X., Berger, J., & Jürgens, G. (1996). The WUSCHEL gene is
required for shoot and floral meristem integrity in Arabidopsis. *Development*,
122, 87–96.
- Lee, H., Jun, Y. S., Cha, O. K., & Sheen, J. (2019). Mitogen-activated protein kinases
MPK3 and MPK6 are required for stem cell maintenance in the Arabidopsis
shoot apical meristem. *Plant Cell Reports*, 38(3), 311–319.
<https://doi.org/10.1007/s00299-018-2367-5>
- Liu, L., Gallagher, J., Arevalo, E. D., Chen, R., Skopelitis, T., Wu, Q., Bartlett, M., &
Jackson, D. (2021). Enhancing grain-yield-related traits by CRISPR–Cas9
promoter editing of maize CLE genes. *Nature Plants*, 7(3), 287–294.
<https://doi.org/10.1038/s41477-021-00858-5>
- Ma, Y., Miotk, A., Šutiković, Z., Ermakova, O., Wenzl, C., Medzihradszky, A.,
Gaillochet, C., Forner, J., Utan, G., Brackmann, K., Galván-Ampudia, C. S.,
Vernoux, T., Greb, T., & Lohmann, J. U. (2019). WUSCHEL acts as an auxin
response rheostat to maintain apical stem cells in Arabidopsis. *Nature*
Communications, 10(5093), 1–11. <https://doi.org/10.1038/s41467-019-13074-9>
- Mandel, T., Moreau, F., Kutsher, Y., Fletcher, J. C., Carles, C. C., & Williams, L. E.

- (2014). The ERECTA receptor kinase regulates Arabidopsis shoot apical meristem size, phyllotaxy and floral meristem identity. *Development*, 141(4), 830–841. <https://doi.org/10.1242/dev.104687>
- Mayer, K. F. X., Schoof, H., Haecker, A., Lenhard, M., Jürgens, G., & Laux, T. (1998). Role of WUSCHEL in Regulating Stem Cell Fate in the Arabidopsis Shoot Meristem. *Cell Press*, 95, 805–815.
- Müller, R., Bleckmann, A., & Simon, R. (2008). The receptor kinase CORYNE of Arabidopsis transmits the stem cell-limiting signal CLAVATA3 independently of CLAVATA1. *Plant Cell*, 20(4), 934–946. <https://doi.org/10.1105/tpc.107.057547>
- Müller, R., Borghi, L., Kwiatkowska, D., Laufs, P., & Simon, R. (2006). Dynamic and Compensatory Responses of Arabidopsis Shoot and Floral Meristems to CLV3 Signaling. *The Plant Cell*, 18(5), 1188–1198. <https://doi.org/10.1105/tpc.105.040444>
- Nimchuk, Z. L. (2017). CLAVATA1 controls distinct signaling outputs that buffer shoot stem cell proliferation through a two-step transcriptional compensation loop. *PLoS Genetics*, 13(3), 1–19. <https://doi.org/10.1371/journal.pgen.1006681>
- Nimchuk, Z. L., Zhou, Y., Tarr, P. T., Peterson, B. a, & Meyerowitz, E. M. (2015). Plant stem cell maintenance by transcriptional cross-regulation of related receptor kinases. *Development*, 142(6), 1043–1049. <https://doi.org/10.1242/dev.119677>
- Ogawa, M., Shinohara, H., Sakagami, Y., & Matsubayash, Y. (2008). Arabidopsis CLV3 peptide directly binds CLV1 ectodomain. *Science*, 319(5861), 294. <https://doi.org/10.1126/science.1150083>
- Ohmori, Y., Tanaka, W., Kojima, M., Sakakibara, H., & Hirano, H. Y. (2013). WUSCHEL-RELATED HOMEODOMAIN4 Is involved in meristem maintenance and is negatively regulated by the CLE gene FCP1 in rice. *The Plant Cell*, 25(1), 229–241. <https://doi.org/10.1105/tpc.112.103432>
- Pallakies, H., & Simon, R. (2014). The CLE40 and CRN/CLV2 signaling pathways antagonistically control root meristem growth in arabidopsis. *Molecular Plant*, 7(11), 1619–1636. <https://doi.org/10.1093/mp/ssu094>
- Reddy, V. G., Heisler, M. G., Ehrhardt, D. W., & Meyerowitz, E. M. (2004). Real-time lineage analysis reveals oriented cell divisions associated with morphogenesis at the shoot apex of Arabidopsis thaliana. *Development*, 131(17), 4225–4237. <https://doi.org/10.1242/dev.01261>

- Rodriguez-Leal, D., Xu, C., Kwon, C. T., Soyars, C., Demesa-Arevalo, E., Man, J., Liu, L., Lemmon, Z. H., Jones, D. S., Van Eck, J., Jackson, D. P., Bartlett, M. E., Nimchuk, Z. L., & Lippman, Z. B. (2019). Evolution of buffering in a genetic circuit controlling plant stem cell proliferation. *Nature Genetics*, 51(5), 786–792. <https://doi.org/10.1038/s41588-019-0389-8>
- Schnablová, R., Neustupa, J., Woodard, K., Klimešová, J., & Herben, T. (2020). Disentangling phylogenetic and functional components of shape variation among shoot apical meristems of a wide range of herbaceous angiosperms. *American Journal of Botany*, 107(1), 20–30. <https://doi.org/10.1002/ajb2.1407>
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9(7), 671–675. <https://doi.org/10.1038/nmeth.2089>
- Schoof, H., Lenhard, M., Haecker, A., Mayer, K. F. X., Jürgens, G., & Laux, T. (2000). The Stem Cell Population of Arabidopsis Shoot Meristems Is Maintained by a Regulatory Loop between the CLAVATA and WUSCHEL Genes. *Cell*, 100, 635–644.
- Shinohara, H., & Matsubayashi, Y. (2015). Reevaluation of the CLV3-receptor interaction in the shoot apical meristem: Dissection of the CLV3 signaling pathway from a direct ligand-binding point of view. *Plant Journal*, 82(2), 328–336. <https://doi.org/10.1111/tpj.12817>
- Shpak, E. D. (2013). Diverse roles of ERECTA family genes in plant development. *Journal of Integrative Plant Biology*, 55(12), 1238–1250. <https://doi.org/10.1111/jipb.12108>
- Shpak, E. D., Berthiaume, C. T., Hill, E. J., & Torii, K. U. (2004). Synergistic interaction of three ERECTA-family receptor-like kinases controls Arabidopsis organ growth and flower development by promoting cell proliferation. *Development*, 131(7), 1491–1501. <https://doi.org/10.1242/dev.01028>
- Somssich, M., Je, B. II, Simon, R., & Jackson, D. (2016). CLAVATA-WUSCHEL signaling in the shoot meristem. *Development*, 143(18), 3238–3248. <https://doi.org/10.1242/dev.133645>
- Stahl, Y., Grabowski, S., Bleckmann, A., Kühnemuth, R., Weidtkamp-Peters, S., Pinto, K. G., Kirschner, G. K., Schmid, J. B., Wink, R. H., Hülsewede, A., Felekyan, S., Seidel, C. A. M., & Simon, R. (2013). Moderation of arabidopsis root stemness by CLAVATA1 and ARABIDOPSIS CRINKLY4 receptor kinase

complexes. *Current Biology*, 23(5), 362–371.
<https://doi.org/10.1016/j.cub.2013.01.045>

Stahl, Y., & Simon, R. (2005). Plant stem cell niches. *International Journal of Developmental Biology*, 49, 479–489. <https://doi.org/10.1387/ijdb.041929ys>

Stahl, Y., & Simon, R. (2010). Plant primary meristems: shared functions and regulatory mechanisms. *Current Opinion in Plant Biology*, 13, 53–58.
<https://doi.org/10.1016/j.pbi.2009.09.008>

Stahl, Y., Wink, R. H., Ingram, G. C., & Simon, R. (2009). A Signaling Module Controlling the Stem Cell Niche in Arabidopsis Root Meristems. *Current Biology*, 19(11), 909–914. <https://doi.org/10.1016/j.cub.2009.03.060>

Steeves, T. A., & Sussex, I. M. (1989). Patterns in Plant Development. *Cambridge University Press*, 2nd edn.

Suzaki, T., Yoshida, A., & Hirano, H. Y. (2008). Functional diversification of CLAVATA3-related CLE proteins in meristem maintenance in rice. *Plant Cell*, 20(8), 2049–2058. <https://doi.org/10.1105/tpc.107.057257>

Takahashi, G., Betsuyaku, S., Okuzumi, N., Kiyosue, T., & Hirakawa, Y. (2021). An Evolutionarily Conserved Coreceptor Gene Is Essential for CLAVATA Signaling in Marchantia polymorpha. *Frontiers in Plant Science*, 12(657548).
<https://doi.org/10.3389/fpls.2021.657548>

Torii, K. U., Mitsukawa, N., Oosumi, T., Matsuura, Y., Yokoyama, R., Whittier, R. F., & Komeda, Y. (1996). The Arabidopsis ERECTA Gene Encodes a Putative Receptor Protein Kinase with Extracellular Leucine-Rich Repeats. *The Plant Cell*, 8(4), 735–746. <https://doi.org/10.1105/tpc.8.4.735>

Whitewoods, C. D., Cammarata, J., Nemec Venza, Z., Sang, S., Crook, A. D., Aoyama, T., Wang, X. Y., Waller, M., Kamisugi, Y., Cuming, A. C., Szövényi, P., Nimchuk, Z. L., Roeder, A. H. K., Scanlon, M. J., & Harrison, C. J. (2018). CLAVATA Was a Genetic Novelty for the Morphological Innovation of 3D Growth in Land Plants. *Current Biology*, 28(15), 2365–2376.
<https://doi.org/10.1016/j.cub.2018.05.068>

Wink, R. (2013). *On the function of peptide signaling pathways in the root meristem of Arabidopsis thaliana*.

Yadav, R. K., Perales, M., Gruel, J., Girke, T., Jönsson, H., & Reddy, V. G. (2011). WUSCHEL protein movement mediates stem cell homeostasis in the Arabidopsis shoot apex. *Genes and Development*, 25(19), 2025–2030.

1031 <https://doi.org/10.1101/gad.17258511>

1032 Yamaguchi, Y. L., Ishida, T., Yoshimura, M., Imamura, Y., Shimaoka, C., & Sawa, S.

1033 (2017). A Collection of Mutants for CLE-Peptide-Encoding Genes in Arabidopsis

1034 Generated by CRISPR/Cas9-Mediated Gene Targeting. *Plant and Cell*

1035 *Physiology*, 58(11), 1848–1856. <https://doi.org/10.1093/pcp/pcx139>

1036 Zhang, L., DeGennaro, D., Lin, G., Chai, J., & Shpak, E. D. (2021). ERECTA family

1037 signaling constrains CLAVATA3 and WUSCHEL to the center of the shoot apical

1038 meristem. *Development*, 148(5), 1–10. <https://doi.org/10.1242/dev.189753>

1039 Zhou, Y., Yan, A., Han, H., Li, T., Geng, Y., Liu, X., & Meyerowitz, E. M. (2018). Hairy

1040 meristem with wuschel confines clavata3 expression to the outer apical

1041 meristem layers. *Science*, 361(6401), 502–506.

1042 <https://doi.org/10.1126/science.aar8638>

1043

Supplementary Figures

Fig1-SupplFig.1: Mutants from the CLV pathway show differences in their leaf lengths

(A) Wild type (*Col-0*) and different single and double mutants (*clv3-9*, *cle40-2*, *clv1-101*, *clv1-101;cle40-2*, *bam1-3*, *bam1-3;cle40-2*, *bam1-3;clv1-101*) at 4 WAG. **(B)** Leaf lengths were measured and plotted. Wild type (*Col-0* N=47), *cle40-2* (N=32) and *bam1-3;cle40-2* (N=29) mutant plants do not show a significant difference in leaf length to each other. While *bam1-3* (N=32) mutants exhibit in average significantly longer leaves than wild type plants, the single mutants *clv3-9* (N=33) and *clv1-101* (N=33) and the double mutants *clv1-101;cle40-2* (N=32) and *bam1-3;clv1-101* (N=45) show significantly shorter leaves. Statistical groups were assigned after calculating p-values by ANOVA and Turkey's multiple comparison test (differential grouping from $p \leq 0.01$).

Fig1-SupplFig.2: *cle40* mutants have smaller meristems

(A) Wild type *A. thaliana* plants (*Col-0*) and *cle40* mutants (*cle40-2*, *cle40-cr1*, *cle40-cr2*, *cle40-cr3*) at 6 WAG. All plants show a similar height ranging from 17.5cm to 21.2cm and do not have an obvious plant phenotype. **(B)** Schematic representation of the *CLE40* gene, consisting of three exons (green arrows) and two introns. Exon 3 carries the crucial CLE box (dashed red line). **(B')** Schematic representation of all four *cle40* mutations. All four lines have mutations in or before the CLE box domain in Exon 3. *cle40-2* mutants were created by transposon mutagenesis resulting in a stop codon in front of the CLE box (Stahl et al., 2009). *cle40-cr1*, *cle40-cr2* and *cle40-cr3* mutants were created using the CRISPR-Cas9 method (Yamaguchi et al., 2017). *cle40-cr1* has an 11bp insertion inside the CLE box domain while *cle40-cr2* and *cle40-cr3* have a deletion of -34bp and -28bp within the CLE box. **(C)** At 6 WAG, IFMs of wild type (*Col-0* N=16) and *cle40* mutant plants were dissected and the area of each meristem was imaged and measured. All four *cle40* mutants show significantly reduced IFM sizes compared

to *Col-0* plants (*cle40-2* N=17, *cle40-cr1* N=24, *cle40-cr2* N=20, *cle40-cr3* N=19). Statistical stars were assigned after calculating p-values by ANOVA and Dunnett's multiple comparison test (differential grouping from $p \leq 0.01$).

Fig1-SupplFig.3: Siliques of various mutants differ in their carpel number

(A) Carpels of *Arabidopsis thaliana* plants at 6 WAG in wild type (*Col-0*) or different mutant backgrounds: *clv3-9*, *cle40-2*, *clv1-20*, *clv1-20;cle40-2*, *bam1-3*, *bam1-3;cle40-2*, *bam1-3;clv1-101*. **(B)** Carpel number was counted and plotted. Wild type (*Col-0* N=290), *cle40-2* (N=300), *bam1-3* (N=300) and *bam1-3;cle40-2* (N=280) mutant plants always develop two carpels, while *clv3-9* (N=340) plants exhibit 3 to 5 carpels and *clv1-20* (N=350) and *clv1-20;cle40-2* (N=320) mutants show an average of 2 to 3 carpels. The double mutant *bam1-3;clv1-101* (N=280) develops 6 carpels on average. N number depicts number of siliques. Statistical

groups were assigned after calculating p-values by ANOVA and Turkey's multiple comparison test (differential grouping from $p \leq 0.01$).

Fig3-SupplFig.1: *CLE40* expression is lacking in the CZ and OC

(A) MIP of *CLV3* and *WUS* expression (*CLV3:NLS-mCherry;WUS:NLS-GFP//clv3-9*) in a *clv3-9* mutant IFM (N=5). *CLV3* expression is detected at the tip of the meristem, while *WUS* expression is predominantly found in young primordia surrounding the meristem. **(A')** Optical section through the IFM shows an extended expression domain of *CLV3* in the CZ and *WUS* expressing cells in OC of the IFM. **(B)** MIP of a *clv3-9* mutant IFM expressing *CLE40:Venus-H2B*. *CLE40* is expressed in the PZ of the IFM, in flower primordia and in mature sepals (N=6). **(B')** Optical section through the IFM shows *CLE40* expression in the outer layers of the PZ while it is lacking in the CZ and OC, where *CLV3* and *WUS* are expressed. Dashed orange line indicates the planes of optical sections; Scale bars: 50 μ m (C, D), 10 μ m (C', D'), MIP = maximum intensity projection, PI = propidium iodide

Fig3-SupplFig.2: *CLE40* expression is extended in *wus-7* mutants

(A) *L. er.* wild type plant at 5 WAG shows normal plant growth, while *wus-7* mutants at 5 WAG are delayed in their development (dashed white line). **(B)** *wus-7* mutant at 8 WAG. *wus-7* mutants develop IFMs but give rise to sterile flowers that lack inner organs. **(C and D)** MIP of *wus-7* IFMs at 5 WAG expressing *CLE40:Venus-H2B*. *CLE40* expression is detected through the entire meristem and in the centre of primordia (N=12). **(D' and D')** Optical sections through the meristem show *CLE40* expression in an extended pattern in the PZ and the OC. Dashed white line in B encloses homozygous *wus-7* mutants, dashed orange line indicates the planes of optical sections; Scale bars: 20mm (A, B), 20µm (C-D')

Fig6-SupplFig. 1 Expression patterns of *CLE40* and *BAM1* overlap in the IFM

Optical sections through an IFM and its developing primordia P1 to P6 expressing either **(A)** *CLE40* (*CLE40:Venus-H2B*) (N=23) or **(B)** *BAM1* (*BAM1:BAM1-GFP*) (N=15). In the IFM, *CLE40* and *BAM1* expression patterns overlap in the PZ, while both genes are lacking in the OC. No *CLE40* expression is detected in young primordia in P1 to P3. From P4 on a faint signal in CZ of the primordia express *CLE40*. Its expression expands in P5 and can be found in almost all cells of P6. *BAM1* is expressed ubiquitously in all primordia from P2 to P6. Yellow lines (P1 to P6) indicate the IFM region, white lines (P1 to P6) mark the primordium, Scale bar: 10µm, P = primordium

Fig6-SupplFig. 2: Meristem measurements in mutant backgrounds

IFMs of (A) *Col-0*, (B) *clv3-9*, (C) *cle40-2*, (D) *clv1-101*, (E) *clv1-101; cle40-2*, (F) *bam1-3*, (G) *bam1-3; cle40-2* and (H) *bam1-3; clv1-101* were imaged after 6 WAG. Z-stacks were taken from

the top of the IFMs with a confocal microscope. A MIP and an optical section from P4 to P5 was performed for each meristem. The yellow dashed line depicts the area of the meristem that was measured and the dashed red line indicates the height of the meristems. Scale bar: 20 μ m

Fig7-SupplFig. 1: IFM height, width and shape after 5 WAG

(A) The width of *Col-0* (N=11), *cle40-2* (N=10), *bam1-3* (N=11), *clv1-101* (N=9) and *clv3-9* (N=5) mutants at 5 WAG does not significantly differ from each other. The average width lays between 85 to 100 μ m. Wild type plants are in average 90 μ m wide while *clv3-9* mutants depict the widest meristem average of 100 μ m. Only *bam1-3* mutants have with an average of 79 μ m a significantly smaller meristem wide compared to wild type plants. **(B)** The height of *cle40-2* (~22 μ m) and *bam1-3* (~23 μ m) mutants is significantly shorter after 5 compared to wild type plants (~28 μ m). In contrast, *clv1-101* and *clv3-9* have significantly higher meristems than *Col-0*, *cle40-2* and *bam1-3* mutants. **(C)** The σ -value represents the shape of the meristem and is defined by the quotient of height and width. *cle40-2* and *bam1-3* mutants have a significantly smaller σ -value compared to wild type plants, resulting in flatter meristems. *clv1-101* and *clv3-9* have with an average of 0.55 a significantly higher σ -value and thus have more dome-shaped meristems.

Green line in the inset meristem in (A) indicates the width that was used for the quantifications in (A); magenta line in the inset meristem in (B) indicates the height that was used for the quantifications in (B); green and magenta line in the inset meristem in (C) indicates the width and height that was used for the quantifications in (C),

Fig8-SupplFig. 1: Schematic model of the two intertwined signaling pathways in a *clv1-101* mutant background

(A) Optical sections of through IFMs show the expression patterns of *CLV3* (N=8), *WUS* (N=8), *BAM1* (N=9) and *CLV1* (N=5) in a *clv1* mutant. Compared to wild type plants, the expression of *CLV3* and *WUS* is expanded and *WUS* is found in a patchy pattern in the L1. *BAM1* expression shifts to the CZ and is found in an elevated expression in the L1. **(B and C)** Schematic representation of two intertwined negative feedback loops in the IFM of a *clv1-101* mutant. The lack of *CLV1* leads to a shift of *BAM1* expression to the OC and to an elevated expression in the L1. In the L3, *BAM1* can partly substitute for *CLV1* and thus *CLV3* can act

via *BAM1* in order to repress *WUS* activity. The elevated expression of *BAM1* in the L1 overlaps in very few cells with *CLE40* expression in the periphery and leads to a weak activation of the downstream signal “X” that promotes *WUS* activity. Since, *WUS* expression is only partly repressed by the *CLV3-BAM1* signalling pathway, the *WUS* domain is extended and leads to an increase in stem cells (expanded *CLV3* expression). *WUS* is now also detected in the L1 of the meristem, together with *BAM1* expression. Scale bars: 20µm (A), CZ =central zone, L1 =layer 1

Fig8-SupplFig. 2: Schematic model of the intertwined signalling pathways in a *clv3-9* mutant background

(A) Optical sections of through IFMs show the expression patterns of *CLV3* (N=5), *WUS* (N=5), *CLE40* (N=6), *BAM1* (N=5) and *CLV1* (N=5) in a *clv3-9* mutant. Compared to wild type plants, the meristem is highly increased in its size along the apical-basal axis and the expression of

CLV3 and *WUS* is expanded in the CZ and OC. *CLE40* expression is limited to the outer layers of the meristems' periphery and excluded from the CZ and OC, while *BAM1* expression shifts towards the inner layers of the PZ. *CLV1* expression is found at the tip and not in the centre of the fasciated meristem. **(B and C)** Schematic representation of two intertwined negative feedback loops in the IFM of a *clv3-9* mutant. The lack of *CLV3* leads to a fasciated meristem with increased number of stem cells and thus an expanded CZ and a decreased PZ. Since no *CLV3* peptide is available, *CLV1* is not activated and expression of *CLV1* shifts from the OC to the tip of the CZ, where it represses *BAM1* expression. *BAM1* is expressed in the inner layers of the PZ, while *CLE40* expression is found in the outer layers of the PZ since it is repressed by the expanded *WUS* domain in the centre of the meristem. Thus only very few cells express both, *BAM1* and *CLE40* and hence, nearly no *WUS* promoting factor "X" is produced and the *CLV3-CLV1* signaling pathway does not repress *WUS* activity. Scale bars: 20µm (A), CZ = central zone, PZ = peripheral zone

Fig8-SupplFig. 4: Schematic model of the intertwined signalling pathways in a *bam1-3* mutant background

(A) Optical sections of through IFMs show the expression patterns of *CLV3* (N=9), *WUS* (N=9), *BAM1* (N=15) and *CLV1* (N=7) in a *bam1-3* mutant. *CLV3* expression is similar to wild type plants, at the tip of the meristem in a cone shaped domain. *WUS* expression is found in the OC, but in less cells than in *Col-0* plants. *CLV1* expression seems to increased in its intensity compared to wild type plants. **(B and C)** Schematic representation of two intertwined negative feedback loops in the IFM of a *bam1-3* mutant. In *bam1-3* mutants, *CLV1* expression appears to be increased. Since *BAM1* is lacking in the periphery, the *WUS* promoting diffusion factor “X” is not being produced and thus *WUS* expression is decreased and confined to the centre of the OC, similar to *cle40-2* plants. With the loss of *BAM1*, the main receptor for *CLE40* is missing, and thus *CLE40* peptide now might signal through *CLV1* leading to a stronger repression of *WUS* from the centre of the meristem.

Scale bars: 20µm (A), CZ = central zone, PZ = peripheral zone

List of Figures

Fig. 1: CLV3 and CLE40 exert opposite effects on meristem size	9
Fig1-SupplFig.1: Mutants from the CLV pathway show differences in their leaf lengths	54
Fig1-SupplFig.2: <i>cle40</i> mutants have smaller meristems	55
Fig1-SupplFig.3: Siliques of various mutants differ in their carpel number	56
Fig. 2: CLE40 and CLV3 show complementary expression patterns in the IFM	11
Fig. 3: WUS-dependent repression of CLE40 expression in the shoot meristem	14
Fig3-SupplFig.1: CLE40 expression is lacking in the CZ and OC	57
Fig3-SupplFig.2: CLE40 expression is extended in <i>wus-7</i> mutants	58
Fig. 4: CLV1 is expressed in the OC and in cells of incipient organ primordia	16
Fig. 5: BAM1 expression is elevated in the flanks of the IFM and not detectable in the OC	18
Fig. 6: BAM1 and CLV1 are receptors for CLE40 and CLV3, respectively	21
Fig6-SupplFig. 1: Expression patterns of CLE40 and BAM1 overlap in the IFM	59
Fig6-SupplFig. 2: Meristem measurements in mutant backgrounds	60
Fig. 7: CLE40 and BAM1 promote WUS expression	24
Fig7-SupplFig. 1: IFM height, width and shape after 5 WAG	61
Fig. 8: Schematic model of two intertwined signalling pathways in the shoot meristem	26
Fig8-SupplFig. 1: Schematic model of the two intertwined signaling pathways in a <i>clv1-101</i> mutant background	62
Fig8-SupplFig. 2: Schematic model of the intertwined signalling pathways in a <i>clv3-9</i> mutant background	63

1253	Fig8-SupplFig. 3: Schematic model of the intertwined signalling pathways in a <i>cle40-2</i>	
1254	mutant background	65
1255	Fig8-SupplFig. 4: Schematic model of the intertwined signalling pathways in a	
1256	<i>bam1-3</i> mutant background	66
1257		

1258 **List of Tables**

1259	Tab. 1: Chemicals and substances used in this study.	37
1260	Tab. 2: Mutants analysed in this study.	38
1261	Tab. 3: Primers and methods used for genotyping.	39
1262	Tab. 4: Entry vectors used for cloning.	40
1263	Tab. 5: Destination vectors used to generate transgenic <i>A.thaliana</i> reporter lines.	41
1264	Tab. 6: Primers used for cloning the entry vectors.	41
1265	Tab. 7: Arabidopsis lines that were analysed in this study.	42
1266	Tab. 8: Microscopy settings used for imaging.	43
1267		