Neurofunctional basis underlying audiovisual integration of print and speech sound in Chinese children

Zhichao Xia ^{a, b * †}, Ting Yang ^{c, d *}, Xin Cui ^{a, e}, Fumiko Hoeft ^{e, f, g, h}, Hong Liu ^{c, d, f},
Xianglin Zhang ^a, Hua Shu ^{a †}, Xiangping Liu ^{c, d †}

^a State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research, Beijing Normal University, China

^b School of Systems Science, Beijing Normal University, China

^c Faculty of Psychology, Beijing Normal University, China

^d Beijing Key Laboratory of Applied Experimental Psychology, National Demonstration Center for Experimental Psychology Education, Faculty of Psychology, Beijing Normal University, China

^e Haskins Laboratories, USA

^f Department of Psychological Sciences and Brain Imaging Research Center, University of Connecticut, USA

^g Department of Psychiatry and Weill Institute for Neurosciences and Dyslexia Center, University of California, San Francisco, USA

^h Department of Neuropsychiatry, Keio University School of Medicine, Japan

* These authors contributed equally to this work

†

† Corresponding authors: Dr. Hua Shu, State Key Lab of Cognitive Neuroscience and Learning, Beijing Normal University, China. Email: shuhua@bnu.edu.cn; Dr. Xiangping Liu, Faculty of Psychology, Beijing Normal University, China. Email: lxp599@163.com; Dr. Zhichao Xia, State Key Lab of Cognitive Neuroscience and Learning, Beijing Normal University, China. Email: xiazc.psy@gmail.com

Acknowledgments: The authors would like to thank all the children and their families for participating in this study and all the examiners for helping with data collection. This work was supported by the Key Program of the National Social Science Foundation of China (14ZDB157), National Key Basic Research Program of China (2014CB846103), National Natural Science Foundation of China grants (31271082, 31671126, 31611130107, 61374165, 81801782), Beijing Municipal Science & Technology Commission (Z151100003915122), Fundamental Research Fund for the Central Universities, China Postdoctoral Science Foundation (2019T120062, 2018M641235), and Social Science Fund of Beijing (17YYA004).

Conflict of Interest: The authors declare no competing financial interests.

Author Contributions: Conceptualization: ZX, HS, XL; Investigation: ZX, TY, XC, HL, XZ; Formal Analysis: ZX; Data Curation: ZX; Writing – Original Draft Preparation: ZX; Writing – Review & Editing: ZX, TY, XC, FH, HL, XZ, HS, XL; Funding Acquisition: ZX, HS, XL; Supervision: ZX, HS, XL

43 **Data Availability Statement:** The data that support the findings of this study are
44 available from the corresponding authors upon reasonable request. The data are not
45 publicly available due to privacy or ethical restrictions.

46 **Running Title:** Brain basis of audiovisual integration

47

Abstract

Effortless print-sound integration is essential to reading development, and the superior temporal cortex (STC) is one most critical brain regions in alphabetic languages. To examine the neural basis in a non-alphabetic language and its relationship with reading ability, we conducted a functional magnetic resonance imaging study in typically developing Chinese children. Audiovisual additive enhancement and congruency effect in two scripts with different orthographic depths—characters (opaque) and pinyin (a transparent alphabetic transcription of characters)—were investigated. In terms of commonality, bilateral STCs displayed additive enhancement in processing both character and pinyin stimulations. Moreover, the magnitudes of congruency effect in the left STC for the two scripts were highly correlated. On the other hand, differences were also observed. First, areas in addition to the STC showed additive enhancement in processing pinyin-sound associations. Second, while the bilateral superior temporal, left inferior/middle frontal and parietal regions manifested a striking incongruency effect (incongruent > congruent) for character-sound pairs, no significant clusters were revealed for pinyin-sounds. Finally, the magnitude of congruency effect in the left middle frontal gyrus for characters was specifically associated with silent reading comprehension proficiency, indicating automatic semantic access and integration during implicit character-sound processing. In contrast, the congruency effect in the left STC for pinyin was specifically associated with oral reading fluency that relies mainly on grapho-phonological mapping. To summarize, this study

revealed both universal and script-specific neurofunctional substrates underlying audiovisual integration as well as their processing- and region-dependent associations with reading abilities in typical Chinese children.

Keywords

Audiovisual integration, character, fMRI, pinyin, Chinese, reading development

Highlights

- The superior temporal cortex is involved in print-sound integration regardless of orthographies.
- The neural manifestation of audiovisual integration is modulated by orthographic transparency.
- The congruency effect in the left middle frontal gyrus for characters is correlated to reading comprehension.
- The congruency effect in the left superior temporal gyrus for pinyin is correlated with oral reading fluency.
- This study provides neural evidence on the role of pinyin in Chinese reading acquisition.

1. Introduction

Efficient print-speech sound mapping is a prerequisite for and a hallmark of successful reading acquisition (Blomert & Froyen, 2010; Fabio Richlan, 2019). Behavioral and neuroimaging studies have revealed that the level of automatic audiovisual integration uniquely accounts for reading outcomes in both shallow (e.g., Dutch, German) and opaque orthographies (e.g., English) (Bakos, Landerl, Bartling, Schulte-Körne, & Moll, 2017; Blau et al., 2010; Leo Blomert & Willems, 2010; Kronschnabel, Brem, Maurer, & Brandeis, 2014; McNorgan, Randazzo-Wagner, & Booth, 2013).

Better knowledge about the neural basis underlying audiovisual integration can further our understanding of how this process contributes to reading development and provide objective training targets for children with reading disorders (RD). In the past two decades, functional magnetic resonance imaging (fMRI) has been widely used to answer the ‘where’ question. Two analytic approaches are the most common methods to identify brain regions involved in processing multimodal information. One is based on the additive model that initially examined in animal electrophysiological studies where researchers found audiovisual signals induced stronger (i.e., super-additive) or lower (i.e., sub-additive) neural response than the summation of responses to unimodal auditory and visual inputs in areas such as the STC (Stein & Stanford, 2008). This super-additive enhancement criterion was then applied to fMRI data but proven too strict

about detecting multisensory integration regions. In contrast, with liberal variants such as mean additivity enhancement criterion (i.e., congruent condition > mean of unimodal auditory and visual conditions), areas including bilateral STCs can be successfully identified (Beauchamp, 2005; Kronschnabel et al., 2014; McNorgan & Booth, 2015). In terms of functional significance, the super- (and mean) additive enhancement occurs early and reflects general aspects of processing (Kronschnabel et al., 2014; Xu, Kolozsvari, Oostenveld, Leppanen, & Hamalainen, 2019). No associations between additive enhancement with reading performance were found (Kronschnabel et al., 2014). In this case, we expected that the bilateral STCs would also show an additive enhancement in processing Chinese print-sound pairs.

Distinguishable brain response to congruent versus incongruent audiovisual pairs is another reliable measure of multisensory integration (a.k.a. congruency effect or incongruency effect if in the opposite direction) (Blau et al., 2010; Blau, van Atteveldt, Ekkebus, Goebel, & Blomert, 2009; Holloway, van Atteveldt, Blomert, & Ansari, 2015; van Atteveldt, Formisano, Goebel, & Blomert, 2004; van Atteveldt, Formisano, Goebel, & Blomert, 2007). Compared with additive enhancement, the congruency effect can reflect deep processes on the information carried by the stimuli. In the pioneering study by van Atteveldt et al. (2004), the STC showed stronger activation in the congruent condition compared with incongruent condition. Although the involvement of STC in audiovisual integration of print and speech sound has been replicated regardless of the level of processing (e.g., phoneme, syllable, or whole word) (Kronschnabel et al., 2014; McNorgan et al.,

2013; Wang, Karipidis, Pleisch, Fraga-Gonzalez, & Brem, 2020), the precise manifestation of the effect (e.g., direction, strength) and whether other areas display such an effect are associated with multiple factors, such as orthographic depth (Holloway et al., 2015; Weiyong Xu, Kolozsvari, Monto, & Hämäläinen, 2018) and reading ability (Blau et al., 2010; Blau et al., 2009; Kronschnabel et al., 2014). In addition, task paradigm (McNorgan & Booth, 2015; van Atteveldt et al., 2007), stimuli properties (Kronschnabel et al., 2014), and developmental stage (Plewko et al., 2018; Wang et al., 2020) also affect the congruency effect but are beyond the scope of this study.

Orthographic depth describes the consistency of mapping between orthographic and phonological forms (e.g., grapheme-phoneme correspondence in alphabetic languages, GPC hereafter), which significantly influences behavioral and neural profiles of reading development and impairments in a given writing system (Borleffs, Maassen, Lyytinen, & Zwarts, 2018; Richlan, 2014). The neural processing accompanying audiovisual integration is also modulated by orthographic transparency. For example, native speakers of English (deep) show an incongruency effect in the STC during processing letter-sound pairs, which differs from the congruency effect observed in Dutch (shallow) (Holloway et al., 2015). More importantly, when the same group of English-speaking participants processed numbers and number names, where the mapping was one-on-one, a strong congruency effect appeared. However, the number of studies directly comparing different orthographies is limited, and most existing evidence is restricted to

alphabetic languages. The question of whether the same neural substrate exists in non-alphabetic languages with dramatically different linguistic characteristics, such as having no GPC rules, remains largely open.

Chinese is particularly suited for answering cross-linguistic questions given its logographic nature and extremely deep orthography (Perfetti, Cao, & Booth, 2013). Since that characters directly map to syllables and there are no GPC rules, the print-sound mapping is quite arbitrary. Moreover, in many cases, different characters share the same pronunciation (i.e., homophones), whereas a single character may also have multiple sounds (i.e., polyphone). Therefore, the link between print (character) and speech sound (syllable) is even more opaque. Based on previous findings in alphabetic languages, an incongruity effect during audiovisual integration was expected. In addition, grapho-semantic correspondence is more systematic in Chinese characters, deeply involved in character recognition, and plays a role equal to or even more important than grapho-phonological processing in reading acquisition (Liu et al., 2017; Ruan, Georgiou, Song, Li, & Shu, 2018; Yang, Shu, McCandliss, & Zevin, 2013; Yang, Zevin, Shu, McCandliss, & Li, 2006). Thus, it is also reasonable to expect that additional regions, such as the inferior and middle frontal cortex known to be involved in semantic processing, would be recruited (Wu, Ho, & Chen, 2012; Zhao et al., 2014). Recently, Xu et al. (2019) conducted a magnetoencephalography (MEG) study in Chinese adults with a one-back task and identified an incongruity effect across the left frontal cortex and STC. While the finding is interesting, some aspects need to be considered,

including the small sample size ($n = 12$), flat tone speech sounds that do not correspond to any character as stimuli, and source reconstruction accuracy of MEG data. Moreover, for experienced adult readers, audiovisual integration of characters and corresponding sound is fully automated. Thus, in addition to cross-linguistic issues, another important but unanswered question involves the neurofunctional features of this process in the developing brain. Against this background, the current study was to investigate the neural basis of integrating characters and speech sounds in typically developing Chinese children.

In particular, we focused on the early-acquired single-element characters given their fundamental role in learning-to-read in Chinese (Shu, Chen, Anderson, Wu, & Xuan, 2003). At the very beginning stage, a small set of pictographic characters are taught first. These simple characters will later be used as radicals in complex compound characters (i.e., phonograms), which comprise greater than 80% of Chinese characters (Chan & Siegel, 2001). Importantly, as a pictographic character does not contain any components (i.e., radicals), there is no clue on its pronunciation or meaning. Thus, the mappings between its visual form with sound (syllable) and meaning (morpheme) are entirely arbitrary. In addition, Mandarin Chinese offers a valuable opportunity to examine the influence of orthographic depth since children learn pronunciations of characters with Hanyu pinyin (pinyin hereafter). In brief, pinyin is an official Roman alphabetic system used to teach the sounds of characters in Mainland China and Singapore (Yin et al., 2011). A child learns pinyin first once entering school and then uses it to assist in conquering the

pronunciation of new characters (Guan, Liu, Chan, Ye, & Perfetti, 2011; Shu, Peng, & McBride-Chang, 2008; Yan, Miller, Li, & Shu, 2008). In other words, pinyin acts as a scaffold during reading acquisition. Given that pinyin borrows letters from alphabets and the majority have consistent mappings with speech sounds (phonemes), it is pretty transparent. Therefore, we included pinyin in this study to directly examine the effect of orthographic depth and expected to observe a similar activation pattern in processing pinyin-sounds as in shallow languages.

In addition to investigating the neural basis of audiovisual integration at the group level, understanding how it correlates to individual heterogeneity in reading abilities is important. On the one hand, children with RD showed a reduced congruency effect compared with their typical peers (e.g., Blau et al., 2010; Blau et al., 2009; Plewko et al., 2018). On the other hand, large individual differences in reading ability are noted even within each group and can be accounted for by variances at the brain level (Kast, Bezzola, Jancke, & Meyer, 2011; McNorgan, Awati, Desroches, & Booth, 2014; McNorgan et al., 2013; Plewko et al., 2018; Wang et al., 2020). Understanding such individual differences in linguistic processing, which is often masked by averaging at the group level, is valuable theoretically and practically (Kanai & Rees, 2011; Kidd, Donnelly, & Christiansen, 2018; Seghier & Price, 2018). Therefore, we also adopted an individual differences approach to investigate whether neural circuitries link congruency effects in processing character/pinyin-sound pairs with typical reading acquisition.

2. Materials and Methods

2.1. Participants

The initial sample consisted of 54 typically developing children according to the following inclusion criteria: (a) native speaker of Mandarin Chinese, (b) right-handed, (c) normal audition, (d) normal or corrected-to-normal vision, (e) full-scale intelligence quotient (IQ) ≥ 80 , (f) no history of neurological or psychiatric disorders, and (g) no reading difficulties ($> 16^{\text{th}}$ percentile on the standardized reading screening task *Character Recognition*) (Xue, Shu, Li, Li, & Tian, 2013). All the children received formal instruction on pinyin in the first half of the semester in the 1st grade. Before the experiment, the children and their parents/guardians were informed about the aim and procedure in detail and signed written consent. All behavioral and neuroimaging data were collected during 2018. This study received ethical approval from the Institutional Review Board of State Key Laboratory of Cognitive Neuroscience and Learning at Beijing Normal University.

The quality of each functional run was evaluated based on in-scanner task performance and the number of outliers (i.e., 'bad' data points) in blood-oxygen-level-dependent signal caused by severe head motion or spikes. Acceptance depended on the following criteria: (i) no more than 15% of the volumes (10 data points) were marked as outliers; (ii) no less than 75% in overall accuracy (combining all the conditions) and accuracy for each condition should be no less than 50%. The run failing to meet either criterion was removed from the analysis. As a result, 36

children had acceptable imaging data for the character experiment (23 girls; age 110-141 months; mean age = 127, grades 3-5). Eighteen children were excluded due to incomplete data collection ($n = 9$), poor data quality ($n = 8$), or the combination of poor data quality and task performance ($n = 1$). For pinyin, the final sample consisted of 41 children (27 girls; age 110-141 months; mean age = 126, grades 3-5). Thirteen children were excluded due to incomplete data collection ($n = 6$), poor data quality ($n = 6$), or the combination of poor data quality and task performance ($n = 1$).

2.2. Behavior measures

Each child underwent testing individually in a silent booth. IQ was measured with the abbreviated version of the *Chinese Wechsler Intelligence Scale for Children (WISC-CR)* (Wechsler, 1974). Specifically, *Information*, *Similarities*, and *Digit Span* subtests were used to estimate verbal IQ; *Picture Completion*, *Block Design*, and *Coding* subtests were used to estimate performance IQ.

A set of neuropsychological tasks was used to measure children's reading and reading-related cognitive-linguistic skills. In brief, *Character Recognition* that had been used for identifying Chinese dyslexic children (Cui, Xia, Su, Shu, & Gong, 2016; Xia, Hoefft, Zhang, & Shu, 2016) was used here for the screening aim. *Word List Reading* and *Silent Reading Comprehension* were used to measure oral reading fluency and reading comprehension proficiency, respectively. The former relies more on grapho-phonological mapping, whereas the latter relies more on grapho-semantic

processing (Xia et al., 2018). Finally, *Phoneme Deletion*, *RAN*, and *Morphological Production* tasks were used to measure PA, RAN, and morphological awareness (MA), the three critical cognitive-linguistic skills in Chinese reading development (Lei et al., 2011).

Character Recognition is a standardized test for estimating the number of characters children have learned. The test consisted of 150 characters selected from grades 1-6 textbooks (Xue et al., 2013). The characters were arranged in the order of difficulty on ten sheets of A4 paper. During the task, the participant was asked to name the characters in sequence until failing in all 15 items on one page. There was no time limit. Each correct answer is worth 1 point, and the full mark is 150.

Word List Reading is a standardized test for measuring the level of oral reading fluency (Zhang et al., 2012). The test consisted of 180 high-frequency two-character words arranged in a 9-column by 20-row matrix in a sheet of A4 paper. The participant was required to read the words aloud in left-to-right, up-to-down order as accurately and quickly as possible. The time used to complete the task was recorded, and an index referring to the number of words the participant read correctly per minute was calculated.

Silent Reading Comprehension is a standardized test for assessing children's reading comprehension (Lei et al., 2011). The test included 100 items arranged based on the number of characters (i.e., from a simple sentence to a short paragraph). The participant was requested to read each sentence/paragraph silently

and decide the correctness of the meaning with a mark of ✓ or ✗ as fast as possible within 3 minutes. The total number of characters in sentences with correct responses was calculated and then transformed to the number of characters the participant read per minute.

Phoneme Deletion was used to assess PA (Li, Shu, McBride-Chang, Liu, & Peng, 2012). This task included three sections, and the participant was asked to delete the initial, middle, or final sound from an orally presented syllable and pronounce the remaining portion. Each section consisted of 3 practice trials and 8-10 testing items. One correct response is worth 1 point for a total possible score of 28.

Digit RAN required the participant to name 50 one-digit numbers on a card as accurately and rapidly as possible (Liu et al., 2017). Numbers 1, 2, 3, 5, and 8 were repeated ten times each and arranged randomly in a 5-column by 10-row matrix. The task was administered twice, and the average time the participant completed the task was used as the score.

Morphological Production was used to measure MA at the character level (Shu, McBride-Chang, Wu, & Liu, 2006). In each trial, a character was orally presented to the participant in a high-frequency word. The participant had to respond with two new words. In one word, the given character retains the same meaning as in the original case, whereas in the other word, the meaning of the

character should be different. This task contained 15 characters for a total possible score of 30.

2.3. Stimuli and experimental design

In this study, we investigated neurofunctional correlates of audiovisual integration in processing character-sound and pinyin-sound pairs separately. In the character experiment, the materials were characters and corresponding speech sounds (syllables) (**Figure 1A**). Fifty-six high frequency pictographic characters were selected (Chinese Single-Character Word Database; <https://blclab.org/psycholinguistic-norms-database/>). All the characters are visually simple, taught early, and used as radicals in complex compound characters. During the experiment, visual stimuli were displayed in white, ‘KaiTi’ font, and 96 pt at the center of a black background. A native Chinese male recorded the speech sounds with a sampling rate of 44.1 kHz and 16-bit quantization. The audio files were normalized to 85 dB and then edited with a bandpass (100-4000 Hz) filter with Audacity (<https://www.audacityteam.org/>). The average duration of all sounds was 476.3 (± 87.5) ms without pre- or post-silence. All the speech sounds can be recognized easily. In the pinyin experiment, the visual stimuli were syllables written in pinyin (e.g., sound /sheng 1/ can be written as ‘shēng’, **Figure 1B**), in white, ‘Century Schoolbook’ font, and 90 pt. The auditory stimuli were the same as those used in character conditions.

The fMRI procedure was adapted from Blau et al. (2010) (**Figure 1C**). A block design was used with four conditions in each experiment: unimodal auditory (Aud), unimodal visual (Vis), audiovisual congruent (avC), and audiovisual incongruent (avI). The visual and auditory stimuli were presented simultaneously in the avC and avI conditions. The entire study contained 4 functional runs, half of which were for character and the other half for pinyin. To avoid priming from character processing on pinyin processing, the pinyin experiment was always conducted first. Each run included 8 experimental blocks (duration 20.8 s), with each condition replicated twice and interleaved with 9 rest blocks (duration 20.8 s). In each experimental block, there were 4 mini-blocks. Each mini-block consisted of a 1.5-s brain volume acquisition period and a 3.7-s silence period (i.e., sparse-sampling) during which 4 trials of stimuli were presented. The order of the stimuli and conditions were pseudorandomized. Considering the age of the participants, a simple target detection task that does not require active monitoring of congruency status was used to help maintain attention during scanning. Compared to the active matching task, the passive paradigm will not change the automatic response pattern during audiovisual integration (Blau et al., 2010; van Atteveldt et al., 2007). The participant was asked to press the button with the right index finger accurately and quickly to the auditory (440 Hz pure tone), visual (an unpronounceable symbol) targets, or their combinations that appeared twice randomly per experimental block (**Figure 1D**). To ensure that the child performed the task correctly, two practice

sessions were administered, including one outside the scanner and the other before the first fMRI session inside the scanner.

2.4. Image acquisition

The 3-Tesla Siemens MAGNETOM Trio Tim scanner in Imaging Center for Brain Research at Beijing Normal University was used to collect all the images with a 12-channel head coil. For each child, 4 T2* sensitive gradient echo planar imaging sequences were collected with the following parameters: repetition time (TR) = 5200 ms (1500 ms image acquisition plus 3700 ms delay), echo time = 32 ms, flip angle = 90 degrees, slice thickness = 4.5 mm, voxel size = $3.0 \times 3.0 \times 4.5 \text{ mm}^3$, interscan gap = 0.675 mm, number of slices = 24, number of volumes = 68, and time of acquisition = 5 minutes and 54 s. The sparse-sampling method provided a 3.7-s period of silence for stimuli presentation. Such a design is recommended for fMRI experiments investigating processing with auditory inputs (Talavage & Hall, 2012). Three dummy scans were administered before image collection for each functional run to avoid scanner equilibration effects. In addition, a high-resolution whole-brain T1-weighted structural image (magnetization-prepared rapid acquisition with gradient echo, TR = 2530 ms, echo time = 3.39 ms, inversion time = 1100 ms, flip angle = 7 degrees, slice thickness = 1.33 mm, voxel size = $1.33 \times 1.0 \times 1.33 \text{ mm}^3$, number of axial slices = 144, and time of acquisition = 8 minutes and 7 s) was collected. Prior to the formal scan, children were informed about the experimental procedure, familiarized with the noise of MRI sequences, and trained to hold still. Earplugs and noise-proof headphones were used to help reduce scanner noise, and foam pads

were used to help reduce head motion during scanning. A radiologist blinded to the details of the study reviewed all the images to ascertain any pathological deviations.

2.5. fMRI preprocessing

Preprocessing of brain images was conducted with SPM12 (Wellcome Department of Cognitive Neurology, London, UK, <http://www.fil.ion.ucl.ac.uk/spm/>). It included the following steps: (a) head motion correction (realign), (b) ART-based outlier detection (the ‘bad’ time point was defined as intensity > global mean $\pm$ 9 SD or frame-to-frame head motion [framewise displacement] > 2 mm; https://www.nitrc.org/projects/artifact_detect), (c) T1 image segmentation, (d) normalization to the standard template in Montreal Neurological Institute (MNI) space, and (e) smoothing with an 8-mm full-width at half-maximum Gaussian kernel. No slice timing correction was performed given the discontinuous nature of the signal in fMRI data collected using sparse-sampling (Perrachione & Ghosh, 2013). At the individual level, experimental conditions were modeled with a generalized linear model. Parameters referring to outliers in the time course and head motion (3 translation, 3 rotation, and 1 framewise displacement) were included in the model to exclude the effect of nuisance covariates. Then, we calculated two brain maps—avC vs. (Aud + Vis) / 2 and avC vs. avI, which were used in the subsequent analyses on additive enhancement and congruency effect, respectively.

2.6. Statistical analyses

Descriptive statistics of demographic and behavioral measures were conducted first, followed by correlations between reading abilities and cognitive-linguistic skills. The Bonferroni method was used to deal with multiple comparisons (corrected $p < 0.05 / 15$). Next, we evaluated in-scanner task performance. In this study, we defined the accurate response as a button press to the target with a reaction time (RT) between 200 and 2000 ms. The average reaction time was calculated for correct responses.

The two aforementioned approaches were used in the whole brain analyses to identify areas involved in audiovisual processing of character and pinyin stimulations separately. The first approach was to detect mean additive enhancement. It consisted of two criteria: (i) stronger activation in each unimodal condition against the baseline, i.e., $Aud > 0$, $Vis > 0$. This criterion was used to exclude areas that only responded to one input modality; (ii) stronger activation in the audiovisual congruent condition compared with the mean of unimodal conditions, i.e., $avC > (Aud + Vis) / 2$. For this, three one-sample t -tests were administered, and intersections were identified. For each t -test, an uncorrected threshold of p -voxel < 0.05 was applied, resulting in a conjoint $p < 0.05^3 = 0.000125$ for the overlapping voxel. To be conservative, only clusters containing more than 200 continuous voxels were considered to be significant. The second approach was used to identify regions showing congruency effects. A one-sample t -test was conducted with individual contrast maps of congruency (i.e., congruent against incongruent). We focused on brain areas showing activation (uncorrected p -

voxel < 0.05) in at least one multimodal condition to avoid results produced by differences in de-activation. A threshold of p -voxel < 0.005 , FWE corrected p -cluster < 0.05 was used to identify significant clusters while dealing with multiple comparisons. Previous studies have repeatedly reported that the congruency effect is associated with reading ability. To avoid missing the effect masked at the group level, we conducted two regression analyses where the standard score of oral reading fluency or silent reading comprehension proficiency was included as the variate of interest, respectively. The same threshold of FWE corrected $p < 0.05$ (cluster-forming p -voxel < 0.005) was used to address multiple comparisons. Nuisance variables of age and sex were controlled statistically in all the analyses.

With the results of whole brain analyses, we defined areas with different combinations of additive enhancement, congruency effect, and congruency-reading correlation as the regions-of-interest (ROIs). Such differences may indicate different roles the areas play in audiovisual integration. Given that domain-specific (i.e., reading-related) processes were of particular interest, we did not include areas that only showed additive enhancement. Those regions could be more involved in domain-general processing. The aims of the ROI analysis were (1) to explicitly present the effects and (2) to exclude the possibility that the absence of significance on a specific effect was due to the strict whole brain threshold. For this, in the region only showing congruency-related effects but no additive enhancement at the whole brain level, we further examined its activation in unimodal conditions and the contrast $avC - (Aud + Vis) / 2$. In the region showing additive enhancement and

congruency-behavior correlation but no congruency effect across the entire group at the whole brain level, we examined the congruency effect with the average values within the cluster. We also examined brain-behavior associations with the other reading ability and cognitive-linguistic skills, while the nuisance variables of age and sex were controlled statistically. Bonferroni correction was used to control the number of behavior variables (corrected $p < 0.05 / 4 = 0.0125$). Finally, given that the left STC was identified in both the character (showing additive enhancement and incongruency effect) and pinyin experiments (showing additive enhancement and congruency-reading correlation), we performed conjunction analysis to examine the spatial relationship between the two clusters. We also calculated the correlation between the magnitudes of congruency effect in processing character-sounds and processing pinyin-sounds.

Significant clusters identified in the whole brain analyses were presented over a FreeSurfer surface template with BrainNet Viewer (Xia, Wang, & He, 2013). The peak location was reported in MNI coordinates, and corresponding brain regions were localized and reported with the AAL atlas implemented in DPABI (<http://rfmri.org/dpabi>). Behavioral analysis and ROI analysis were performed with SPSS Statistics version 24 (SPSS Inc., Chicago, IL, USA).

3. Results

3.1. Behavioral profiles and in-scanner task performance

Descriptive statistics of demographics, cognitive-linguistic and reading skills for all 41 children are presented in **Table 1**. Significantly positive correlations were observed between reading measures (all p 's < 0.05 / 15, Bonferroni correction; see **Table 2** for the correlation coefficients and raw p -values). Regarding the relationship with cognitive-linguistic skills, *Word List Reading* was significantly associated with *Digit RAN* ($r = -0.597$, $p < 0.001 < 0.05 / 15$), but not with *Phoneme Deletion* ($r = 0.288$, $p = 0.068 > 0.05 / 15$) or *Morphological Production* ($r = 0.344$, $p = 0.028 > 0.05 / 15$) with Bonferroni correction. In contrast, *Silent Reading Comprehension* was significantly correlated with morphological production ($r = 0.487$, $p = 0.001 < 0.05 / 15$), but not with *Phoneme Deletion* ($r = 0.135$, $p = 0.400 > 0.05 / 15$) or *Digit RAN* ($r = -0.332$, $p = 0.034 > 0.05 / 15$) with Bonferroni correction. This pattern indicates that in typically developing Chinese children who have received formal reading instruction for 3-5 years, oral reading necessitates more print-to-sound mapping, whereas reading comprehension relies more on semantic processing.

Regarding the in-scanner task, all the children performed well (character accuracy: $M [SD] = 96.5\% [4.8]$, RT $M [SD] = 512 [81]$ ms; pinyin accuracy $M [SD] = 96.4\% [3.7]$, RT $M [SD] = 530 [80]$ ms; **Table S1**), and no significant differences were found between runs (all p 's > 0.1). These results indicate that participants maintained attention to incoming stimuli during the entire scan session. The results of ANOVA with condition as the repeated measure factor revealed no significant effect on accuracy ($F = 1.00$, $p = 0.395$) and a significant effect on RT in

the character experiment ($F = 35.8, p < 0.001$). Post-hoc tests revealed the longest RT for the unimodal visual condition (p 's ≤ 0.002 in all comparisons, Bonferroni correction), followed by the auditory condition (Aud > avC: corrected $p < 0.001$; Aud > avI: corrected $p < 0.001$). There was no difference between the two crossmodal conditions. In the pinyin experiment, the effect of condition was significant on both accuracy ($F = 10.1, p < 0.001$) and RT ($F = 54.1, p < 0.001$). Post-hoc tests revealed that while children had the lowest accuracy in the visual condition (p 's ≤ 0.005 in all comparison, Bonferroni correction), no differences were found between any other pairs of conditions. In the unimodal visual condition, participants also showed the longest RT (all corrected p 's ≤ 0.05), followed by the auditory condition (Aud > avC: corrected $p < 0.001$; Aud > avI: corrected $p < 0.001$). No differences were observed between the two crossmodal conditions.

3.2. Brain results: the character experiment

The bilateral STCs were identified as showing additive enhancement (**Figure 2A; Table 3**; see **Figure S1** for activation in two unimodal conditions and the brain map for avC > (Aud + Vis) / 2). On the other hand, a significantly incongruency effect (i.e., avC < avI) was found in the left frontal (peak MNI -40, 14, 32), parietal (peak MNI -36, -62, 48) and bilateral STCs (peak MNI, left: -54, -22, -2; right: 62, -34, 10; **Figure 2B, Table 4**) at the FWE corrected threshold of p -cluster < 0.05. In addition, whole-brain regression analysis identified a cluster in the left middle frontal cortex (peak MNI -32, 10, 42) showing a positive correlation between the congruency effect and silent reading comprehension proficiency at the FWE

corrected threshold of p -cluster < 0.05 (**Figure 2C, Table 4**). No regions were identified as significant in whole-brain regression for oral reading fluency.

Based on the whole brain results, we defined three types of ROI and further examined their response properties and relationships with reading abilities. The first type of region (bilateral STCs) showed both additive enhancement and incongruency effect (**Figure 2D**). No congruency-reading correlations were found even at uncorrected p -voxel < 0.05 . The second type of region (left IPL) showed only incongruency effect in the whole brain analyses (**Figure 2E**). We found no activation in either the unimodal condition or the audiovisual congruent condition. No brain-behavior correlations were found in this area neither. The third type of region (left MFG) showed a significant incongruency effect, of which the magnitude was further associated with silent reading comprehension proficiency (**Figure 2F**). In the ROI analysis, we found significant activation for unimodal auditory and visual conditions in this area but observed no activation in the audiovisual congruent condition. For the brain-behavior relationship, the magnitude of the congruency also correlated with oral reading fluency ($r = 0.382$, $p = 0.026$) at an uncorrected threshold, and showed correlated trends for digit RAN ($r = -0.319$, $p = 0.066$) and MA ($r = 0.300$, $p = 0.084$). Importantly, the positive correlation between congruency and silent reading comprehension remained significant ($r = 0.494$, $p = 0.004$) even after additionally controlling for oral reading fluency. Finally, we found that the positive correlation between congruency and reading comprehension was

driven by a negative correlation in the incongruent condition (avC: $r = 0.084$, $p = 0.637$; avI: $r = -0.487$, $p = 0.003$).

3.3. Brain results: the pinyin experiment

The audiovisual additive enhancement was detected in the left superior temporal, inferior frontal, right superior, inferior temporal, and subcortical cortices (**Figure 3A; Table 3**; see **Figure S1** for activation in the unimodal conditions and brain map for $\text{avC} > (\text{Aud} + \text{Vis}) / 2$). No regions showed a significant congruency effect in either direction in the whole brain analysis, which is unexpected. However, voxel-wise regression analysis revealed that oral reading fluency was positively correlated with the magnitude of congruency effect in the left STC (peak MNI -56 -36 14) (FWE corrected p -cluster < 0.05 ; **Figure 3B, Table 4**). No regions were identified as significant in whole-brain regression for silent reading comprehension proficiency. Therefore, we defined the area (left STC) showing both additive enhancement and significant congruency-reading correlation as the ROI (**Figure 3C**). In the subsequent analyses, we found no congruency or incongruency effect in this region even at the uncorrected threshold of $p < 0.05$. In terms of brain-behavior correlation, the magnitude of congruency effect was correlated with silent reading comprehension without multiple comparisons correction ($r = 0.371$, $p = 0.020$, uncorrected). No correlations or correlated trends were found with PA ($r = -0.087$, $p = 0.600$), digit RAN ($r = -0.248$, $p = 0.127$), or MA ($r = 0.195$, $p = 0.235$). Of importance, the correlation between the congruency effect and oral reading fluency remained significant ($r = 0.437$, $p = 0.006$) even after additionally controlling for

reading comprehension. Finally, different from the left MFG observed in the character experiment, the positive congruency-reading correlation in the left STC was driven by a positive correlation in the congruent condition (avC: $r = 0.381$, $p = 0.017$; avI: $r = -0.039$, $p = 0.815$), suggesting distinct mechanisms.

Given that the left STC was identified as functionally important (significant in both additive enhancement and congruency effect related analyses) in audiovisual processing in both experiments, we conducted follow-up conjunction analysis and correlation analysis to examine their relationship. First, the two ROIs spatially overlapped (**Figure 4A**). Second, the magnitudes of the congruency effect in processing character-sound pairs and pinyin-sound pairs were significantly correlated ($r = 0.512$, $p = 0.002$; **Figure 4B**), while nuisance variables of age and sex were statistically controlled.

4. Discussion

The present study investigated the brain bases underlying audiovisual integration in characters (deep orthography) and pinyin (a transparent alphabetic coding system) in typically developing Chinese children. The results revealed both script-shared and script-unique neural substrates, supporting a universal principle underlying the neural mechanism of reading-related processes across languages. Furthermore, the evidence implies that the left MFG links audiovisual integration of character-sounds with silent reading comprehension proficiency by supporting automatic semantic access—reflecting a language-specific manifestation, while the

left STG links audiovisual integration of pinyin-sounds with oral reading fluency by underpinning the shared grapho-phonological mapping component—providing a possible mechanism with which pinyin helps with Chinese reading development.

4.1. Neural processes accompanying print-sound integration in scripts with contrasting transparencies

Several neuroimaging studies have investigated the neural mechanism underlying print-sound integration and its role in reading development with different techniques. On the one hand, electroencephalogram (EEG) and MEG were used to reveal the temporal properties of neural activities (Froyen, van Atteveldt, & Blomert, 2010; Froyen, Van Atteveldt, Bonte, & Blomert, 2008; Froyen, Bonte, van Atteveldt, & Blomert, 2009; Raij, Uutela, & Hari, 2000; Xu et al., 2019). These studies demonstrated that integration occurs preconsciously in expert reading and is reflected in the component of mismatch negativity/response. On the other hand, fMRI research provided more precise and detailed information about the spatial localization of the processing. Among others, the bilateral STCs have been identified as the most relevant areas. Like other stimuli categories, processing audiovisual information of scripts induces super-additivity (or mean additive) enhancement in STC (Beauchamp, 2005; McNorgan & Booth, 2015; van Atteveldt et al., 2004). Strikingly, differences in brain responses to audiovisual congruent against incongruent print-sound pairs have been repeatedly reported in these areas (van Atteveldt et al., 2004; van Atteveldt, Formisano, Blomert, & Goebel, 2007).

Importantly, such effects were observed in passive paradigms, pinpointing the automatic nature of this effect (Blau et al., 2010; van Atteveldt et al., 2007).

Additive enhancement is more domain-general and has been reported in the left STC in both transparent (van Atteveldt et al., 2004) and opaque languages (McNorgan & Booth, 2015). Consistent with previous studies, we observed audiovisual additive enhancement in the bilateral STCs in processing both character-sound (deep) and pinyin-sound (shallow) associations. Compared with additive enhancement, the congruency effect reflects more domain-specific processing and is affected by factors, such as linguistic characteristics (Holloway et al., 2015) and reading abilities (Blau et al., 2010; Blau et al., 2009; Karipidis et al., 2018). Consistency of grapheme-phoneme mapping (orthographic depth or transparency) is an essential feature in alphabetic languages. Differences in the direction of the congruency effect and brain regions showing this effect have been reported between languages with different orthographic depths (Blomert & Froyen, 2010; Holloway et al., 2015). In particular, greater activation in the congruent condition than incongruent condition is more consistently reported in skilled readers when processing letter-sound associations in transparent scripts. In contrast, the opposite pattern is observed in opaque scripts (Holloway et al., 2015). Chinese is a logographic language with an extremely deep orthography. Therefore, we expected to observe an incongruency effect. In support of this hypothesis, the results of the character experiment revealed that four regions (including the bilateral STC) displayed significant incongruency effects. In contrast, in the absence

of a congruency or incongruency effect in processing pinyin-sound pairs at the group level, we observed a positive correlation between the magnitude of congruency effect and oral reading fluency. Higher skill readers showed a congruency effect, whereas lower skill readers showed an incongruency effect. This pattern is consistent with previous studies that revealed the lack of a congruency effect in dyslexia in transparent languages (Blau et al., 2010; Blau et al., 2009), as well as these investigated brain-behavior correlations at both the phoneme (Karipidis et al., 2016; Karipidis et al., 2018) and syllable levels (McNorgan et al., 2014; Wang et al., 2020).

4.2. Left frontal and parietal cortex in implicitly integrating audiovisual information of Chinese characters

Opaque orthography is not the only linguistic feature that affects the neural processing of character-sound processing. Chinese characters are morpheme-syllabic, which means that semantic information is deeply involved. Therefore, in addition to grapho-phonological mapping, semantic access is an equally or even more important aspect in reading (Bi, Han, Weekes, & Shu, 2007; Dang, Zhang, Wang, & Yang, 2018; Liu et al., 2017; Ruan et al., 2018; Yang et al., 2013; Yang et al., 2006). In this study, the left MFG displayed a significant incongruency effect and showed a unique association between the magnitude of the effect and silent reading comprehension proficiency. Considering that this region did not show additive enhancement, we speculate that the observed incongruency effect is more likely to be downstream and reflects semantic processing. In previous research, frontal areas have also been reported along with a broad bilateral fronto-parietal

network. Its function was proposed as task-related top-down modulation (van Atteveldt et al., 2007) or implicit domain-general conflict detection and resolution (Holloway et al., 2015). Recently, the incongruency effect measured with MEG was sourced to the frontal cortex in experienced Chinese adult readers (Xu et al., 2019). Here, we replicated this finding by revealing the incongruency effect in the left MFG in a developmental population in Chinese with fMRI and further demonstrated that the role of this region in the implicit audiovisual integration of characters is domain-specific and that the information automatically processed in this region corresponds to semantics. This finding is also consistent with the opinion that while the fundamental neural circuit for reading is universal, linguistic features in a given language introduce different demands for specific cognitive components (Perfetti et al., 2013; Rueckl et al., 2015).

In Chinese, neither grapho-semantic mapping nor phono-semantic mapping is reliable at the character level. Given that no activation was found in the audiovisual congruent condition for characters, the incongruency effect in the left frontal cortex is more likely to reflect a resolution of the conflict between different meanings accessed via visual and auditory inputs. Interestingly, the positive correlation between the magnitude of congruency effect and reading comprehension proficiency was driven by a negative correlation in the incongruent condition. In other words, individuals with better comprehension performance showed lower activation for non-matching print-sound pairs. This pattern could be caused by higher neural efficiency in relevant processing, such as implicit suppression of

conflicting semantic information. It has previously been found that in the initial stage of establishing a binding between grapheme and phoneme in Swiss-German, typical readers show higher activation in incongruent conditions, and such activation decreases as automation is achieved (Plewko et al., 2018).

In addition to the bilateral temporal and left frontal cortices, a left parietal area also displayed a significant incongruency effect. Like the frontal region, this region did not show additive enhancement, implying that higher activation in the incongruent condition is more likely to be a downstream effect. Unlike the left MFG, no activation was observed in either the unimodal auditory or visual condition, and there was no congruency-reading correlation. Therefore, although previous studies regarded the left parietal cortex as a part of the semantic network (Binder & Desai, 2011; Wang, Zhao, Zevin, & Yang, 2016), there was an absence of evidence indicating that the incongruency effect in this area reflects semantic processing. In another line of research, compared with English, the left parietal region was more recruited in Chinese reading for visual orthographic analysis given the visual complexity of characters (Perfetti et al., 2013). In this case, the incongruency effect might reflect a conflict of orthographic information. However, here we found no activation in the visual condition, going against this explanation. Third, although the left parietal cortex has not been associated with print-sound integration previously in one's native language, a recent training study revealed that this region was involved in acquiring novel grapho-phonological mapping (Xu,

Kolozsvari, Oostenveld, & Hamalainen, 2020). As its function in audiovisual processing remains unclear, future studies are needed.

4.3. The left STG links implicit integration of pinyin-sound pairs with oral reading fluency

While conquering the sounds of letters to an automatic level is one of the most critical prerequisites in alphabetic languages (Blomert, 2011; Richlan, 2019), learning pinyin—a transparent phonological coding system—plays a scaffolding role at the initial stage of Chinese reading acquisition in Mainland China. Therefore, the present study not only examined whether and how the neural features of audiovisual integration are associated with orthographic transparency but also provided information that helps illustrate the role of pinyin in developing fluent reading ability in Chinese.

First, widespread brain regions, including the bilateral STCs, showed additive enhancement in processing pinyin-sound associations, in line with previous findings in alphabetic languages (Kronschabel et al., 2014; McNorgan & Booth, 2015). For the congruency effect, we expected to observe a stronger activation in the congruent condition than in the incongruent condition in the STC based on previous studies in transparent scripts (Blau et al., 2010; van Atteveldt et al., 2004). At first glance, no significant congruency or incongruency effect was observed. There could be several reasons. First, the stimuli we used were at the syllable level. Differences between the letter-phoneme integration and string-syllable integration have been

reported in previous studies (Kronshabel et al., 2014). However, the authors also found that neural response to word-like stimuli is more sensitive to reading abilities. Thus, the alternative possibility is that the direction and strength of the congruency effect are associated with reading abilities and masked by group averaging. In line with the latter, we did observe a positive correlation between the magnitude of the congruency effect in the left STG and oral reading fluency. Specifically, children with higher oral reading scores are more likely to show stronger congruency effects. In contrast, those with lower scores are more likely to show weaker congruency effects or even incongruency effects. This finding is consistent with studies, where a reduced congruency effect was found in individuals with RD in shallow languages (Blau et al., 2010; Blau et al., 2009), studies exploring brain-behavior correlation in transparent (Wang et al., 2020), and opaque languages (McNorgan et al., 2014), as well as studies using a training paradigm (Karipidis et al., 2016; Karipidis et al., 2018). The left STG is the central area representing phonological information (Boets et al., 2013; Glezer et al., 2016) and is involved in visual phonological processing in Chinese children and adults (Cao et al., 2010). Its activation and connectivity change after learning visual-sound mappings (Dehaene et al., 2010; Li, Xu, Luo, Zeng, & Han, 2020; Thiebaut de Schotten, Cohen, Amemiya, Braga, & Dehaene, 2014), which can further scaffold later reading development (Wang, Joanisse, & Booth, 2020). Noteworthy, the positive correlation between the congruency effect in the left STG for pinyin-sound pairs and oral reading fluency was driven by a positive correlation in the congruent condition.

This pattern is different from that in the left MFG in processing character-sound pairs, indicating different mechanisms. It should also be noted that the current findings were revealed with typical readers. Whether children with RD have deficits or show other processing patterns requires further investigation.

The significant correlation between the magnitude of congruency effect in the left STG to pinyin-sound pairs with oral reading fluency implies that learning pinyin helps children develop the related cognitive components and underlying neural circuits for fluent oral reading (though causal claims cannot be made from this study). Compared with cognitive-linguistic skills such as PA and RAN, less attention has been given to pinyin processing in understanding Chinese reading development, probably because of its auxiliary role as a scaffold in reading acquisition but not reading itself. However, although the number of studies is limited, the pinyin-reading relationship has been demonstrated (Lin et al., 2010; Pan et al., 2011; Siok & Fletcher, 2001; Zhang, Georgiou, Inoue, Zhong, & Shu, 2020). Pan et al. found that invented pinyin spelling at age 6 independently predicted reading performance at ages 8 and 10 (Pan et al., 2011). Zhang and colleagues' recent finding suggests a bidirectional relationship (Zhang et al., 2020). Several explanations were proposed based on behavioral evidence. One is that pinyin can help to establish better phonological awareness, the cognitive processing shared by pinyin and character reading (Ding, Liu, McBride, & Zhang, 2015; Li et al., 2020; McBride, Wang, & Cheang, 2018; Shu et al., 2008; Yin et al., 2011). Above and beyond PA, there is also a unique contribution from pinyin processing skills to

character reading (Ju, Zhou, & delMas, 2021). It is reasonable to hypothesize that pinyin may also help establish the correspondence between print and sound. In support of this notion, a study revealed that training on pinyin uniquely improved character-to-syllable mapping in Chinese as a second language learner (Guan et al., 2011). In the present study, a spatially overlapping region also showed additive enhancement and incongruency effect in processing characters. Furthermore, we found that the magnitudes of congruency effect in processing two types of stimuli were highly correlated. As the left posterior STC is involved in visual rhyming judgment of Chinese characters and proposed to play a role in mapping between orthography and phonology (Cao et al., 2010), these findings collectively suggest that learning pinyin may shape the neural circuit for print-sound integration that will later be recruited in learning characters.

4.4. Caveats and future directions

While this study investigated the neurofunctional basis of integrating audiovisual information of characters and pinyin, representing one of the first efforts, caution should be considered when interpreting the results. First, we focused on print-sound integration at the syllable level, whereas previous studies mainly explored letter-speech sound associations (but see Kronschnabel et al., 2014; Wang et al., 2020). This is because Chinese has no GPC rules, and print-sound mapping occurs at the syllable level. But similar to GPC rules in alphabetic languages, simple characters are taught at the earliest stages and play fundamental roles in reading development (Shu et al., 2003; Xing, Shu, & Li, 2004). Second, because our

participants were in grades 3-5, pinyin may play a different role than it does in earlier stages of reading development. Siok and Fletcher (2001) unveiled significant correlations between pinyin knowledge and reading in grades 2-5, but not in grade 1. Thus, it could be the particular age range that enabled us to observe the relationship between print-sound processing in pinyin and oral reading fluency. To further address this issue, it is necessary to conduct studies following preliterate children until they achieve fluent reading. Finally, in this study, we uncovered the brain bases of print-sound integrations in two contrasting scripts and their relationship with reading performance in typically developing children. One next question that warrants investigation in the future is whether neural processes are altered in children with RD.

5. Conclusion

In the current study, we investigated regions that are involved in audiovisual integration of character-sounds and pinyin-sounds. The results suggest the bilateral STCs as the universal brain substrates underlying audiovisual integration across scripts. At the same time, the precise manifestation of specific effects is influenced by linguistic features in the given language. In particular, the left MFG is strongly involved in the audiovisual integration of characters and correlated with reading comprehension, suggesting that semantic representations were more efficiently accessed and processed in readers with higher comprehension proficiency. In contrast, the brain-behavior correlation in the left STC implies that pinyin may

help children develop better reading skills by shaping the shared component of grapho-phonological mapping and shaping the underlying brain circuits.

References

- Bakos, S., Landerl, K., Bartling, J., Schulte-Körne, G., & Moll, K. (2017). Deficits in Letter-Speech Sound Associations but Intact Visual Conflict Processing in Dyslexia: Results from a Novel ERP-Paradigm. *Front Hum Neurosci*, 11(116).
- Beauchamp, M. S. (2005). Statistical criteria in fMRI studies of multisensory integration. *Neuroinformatics*, 3(2), 93-113.
- Bi, Y., Han, Z., Weekes, B., & Shu, H. (2007). The interaction between semantic and the nonsemantic systems in reading: Evidence from Chinese. *Neuropsychologia*, 45(12), 2660-2673.
- Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. *Trends Cogn Sci*, 15(11), 527-536.
- Blau, V., Reithler, J., van Atteveldt, N., Seitz, J., Gerretsen, P., Goebel, R., & Blomert, L. (2010). Deviant processing of letters and speech sounds as proximate cause of reading failure: a functional magnetic resonance imaging study of dyslexic children. *Brain*, 133(Pt 3), 868-879.
- Blau, V., van Atteveldt, N., Ekkebus, M., Goebel, R., & Blomert, L. (2009). Reduced neural integration of letters and speech sounds links phonological and reading deficits in adult dyslexia. *Current Biology*, 19(6), 503-508.

- 792 Blomert, L. (2011). The neural signature of orthographic-phonological binding in
793 successful and failing reading development. *Neuroimage*, 57(3), 695-703.
- 794 Blomert, L., & Froyen, D. (2010). Multi-sensory learning and learning to read. *Int J*
795 *Psychophysiol*, 77(3), 195-204.
- 796 Blomert, L., & Willems, G. (2010). Is there a causal link from a phonological
797 awareness deficit to reading failure in children at familial risk for dyslexia?
798 *Dyslexia*, 16(4), 300-317.
- 799 Boets, B., Op de Beeck, H. P., Vandermosten, M., Scott, S. K., Gillebert, C. R.,
800 Mantini, D., . . . Ghesquiere, P. (2013). Intact but less accessible phonetic
801 representations in adults with dyslexia. *Science*, 342(6163), 1251-1254.
- 802 Borleffs, E., Maassen, B. A. M., Lyytinen, H., & Zwarts, F. (2018). Cracking the
803 Code: The Impact of Orthographic Transparency and Morphological-Syllabic
804 Complexity on Reading and Developmental Dyslexia. *Front Psychol*, 9(2534),
805 2534.
- 806 Cao, F., Lee, R., Shu, H., Yang, Y. H., Xu, G. Q., Li, K. C., & Booth, J. R. (2010).
807 Cultural Constraints on Brain Development: Evidence from a Developmental
808 Study of Visual Word Processing in Mandarin Chinese. *Cereb Cortex*, 20(5),
809 1223-1233.
- 810 Chan, C. K. K., & Siegel, L. S. (2001). Phonological Processing in Reading Chinese
811 among Normally Achieving and Poor Readers. *J Exp Child Psychol*, 80(1), 23-
812 43.

- 813 Cui, Z., Xia, Z., Su, M., Shu, H., & Gong, G. (2016). Disrupted white matter
814 connectivity underlying developmental dyslexia: A machine learning
815 approach. *Human Brain Mapping*, 37(4), 1443-1458.
- 816 Dang, M., Zhang, R., Wang, X., & Yang, J. (2018). The Interaction Between
817 Phonological and Semantic Processing in Reading Chinese Characters. *Front*
818 *Psychol*, 9, 2748.
- 819 Dehaene, S., Pegado, F., Braga, L. W., Ventura, P., Nunes, G., Jobert, A., . . . Cohen,
820 L. (2010). How Learning to Read Changes the Cortical Networks for Vision
821 and Language. *Science*, 330(6009), 1359-1364.
- 822 Ding, Y., Liu, R. D., McBride, C., & Zhang, D. (2015). Pinyin Invented Spelling in
823 Mandarin Chinese-Speaking Children With and Without Reading
824 Difficulties. *J Learn Disabil*, 48(6), 635-645.
- 825 Froyen, D., van Atteveldt, N., & Blomert, L. (2010). Exploring the Role of Low Level
826 Visual Processing in Letter-Speech Sound Integration: A Visual MMN Study.
827 *Front Integr Neurosci*, 4, 9.
- 828 Froyen, D., Van Atteveldt, N., Bonte, M., & Blomert, L. (2008). Cross-modal
829 enhancement of the MMN to speech-sounds indicates early and automatic
830 integration of letters and speech-sounds. *Neurosci Lett*, 430(1), 23-28.
- 831 Froyen, D. J. W., Bonte, M. L., van Atteveldt, N., & Blomert, L. (2009). The Long
832 Road to Automation: Neurocognitive Development of Letter-Speech Sound
833 Processing. *J Cogn Neurosci*, 21(3), 567-580.

- 834 Glezer, L. S., Eden, G., Jiang, X., Luetje, M., Napoliello, E., Kim, J., & Riesenhuber,
835 M. (2016). Uncovering phonological and orthographic selectivity across the
836 reading network using fMRI-RA. *Neuroimage*, 138, 248-256.
- 837 Guan, C. Q., Liu, Y., Chan, D. H. L., Ye, F., & Perfetti, C. A. (2011). Writing
838 strengthens orthography and alphabetic-coding strengthens phonology in
839 learning to read Chinese. *Journal of Educational Psychology*, 103(3), 509-522.
- 840 Holloway, I. D., van Atteveldt, N., Blomert, L., & Ansari, D. (2015). Orthographic
841 Dependency in the Neural Correlates of Reading: Evidence from Audiovisual
842 Integration in English Readers. *Cereb Cortex*, 25(6), 1544-1553.
- 843 I Karipidis, I., Pleisch, G., Röthlisberger, M., Hofstetter, C., Dornbierer, D.,
844 Stämpfli, P., & Brem, S. (2016). Neural initialization of audiovisual
845 integration in prereaders at varying risk for developmental dyslexia. *Human*
846 *Brain Mapping*.
- 847 Ju, Z., Zhou, Y., & delMas, R. (2021). The contributions of separate pinyin skills and
848 oral vocabulary to Chinese word reading of U.S. Mandarin immersion third
849 graders. *Reading and Writing*.
- 850 Kanai, R., & Rees, G. (2011). The structural basis of inter-individual differences in
851 human behaviour and cognition. *Nature Reviews Neuroscience*, 12(4), 231-
852 242.
- 853 Karipidis, I. I., Pleisch, G., Brandeis, D., Roth, A., Röthlisberger, M., Schneebeli, M.,
854 . . . Brem, S. (2018). Simulating reading acquisition: The link between

- 855 reading outcome and multimodal brain signatures of letter–speech sound
- 856 learning in prereaders. *Scientific Reports*, 8(1), 7121.
- 857 Kast, M., Bezzola, L., Jancke, L., & Meyer, M. (2011). Multi- and unisensory
- 858 decoding of words and nonwords result in differential brain responses in
- 859 dyslexic and nondyslexic adults. *Brain and Language*, 119(3), 136-148.
- 860 Kidd, E., Donnelly, S., & Christiansen, M. H. (2018). Individual Differences in
- 861 Language Acquisition and Processing. *Trends Cogn Sci*, 22(2), 154-169.
- 862 Kronschnabel, J., Brem, S., Maurer, U., & Brandeis, D. (2014). The level of
- 863 audiovisual print-speech integration deficits in dyslexia. *Neuropsychologia*,
- 864 62, 245-261.
- 865 Lei, L., Pan, J., Liu, H., McBride-Chang, C., Li, H., Zhang, Y., . . . Shu, H. (2011).
- 866 Developmental trajectories of reading development and impairment from
- 867 ages 3 to 8 years in Chinese children. *J Child Psychol Psychiatry*, 52(2), 212-
- 868 220.
- 869 Li, H., Shu, H., McBride-Chang, C., Liu, H., & Peng, H. (2012). Chinese children's
- 870 character recognition: Visuo-orthographic, phonological processing and
- 871 morphological skills. *Journal of Research in Reading*, 35(3), 287-307.
- 872 Li, M., Xu, Y., Luo, X., Zeng, J., & Han, Z. (2020). Linguistic experience acquisition
- 873 for novel stimuli selectively activates the neural network of the visual word
- 874 form area. *Neuroimage*, 215, 116838.
- 875 Li, Y., Chen, X., Li, H., Sheng, X., Chen, L., Richardson, U., & Lyytinen, H. (2020).
- 876 A computer-based Pinyin intervention for disadvantaged children in China:

- 877 Effects on Pinyin skills, phonological awareness, and character reading.
878 *Dyslexia*, 26(4), 377-393.
- 879 Lin, D., McBride-Chang, C., Shu, H., Zhang, Y., Li, H., Zhang, J., . . . Levin, I.
880 (2010). Small Wins Big: Analytic Pinyin Skills Promote Chinese Word
881 Reading. *Psychol Sci*, 21(8), 1117-1122.
- 882 Liu, Y., Georgiou, G. K., Zhang, Y., Li, H., Liu, H., Song, S., . . . Shu, H. (2017).
883 Contribution of cognitive and linguistic skills to word-reading accuracy and
884 fluency in Chinese. *International Journal of Educational Research*, 82, 75-90.
- 885 McBride, C., Wang, Y., & Cheang, L. M.-L. (2018). Dyslexia in Chinese. *Current*
886 *Developmental Disorders Reports*, 5(4), 217-225.
- 887 McNorgan, C., Awati, N., Desroches, A. S., & Booth, J. R. (2014). Multimodal
888 Lexical Processing in Auditory Cortex Is Literacy Skill Dependent. *Cereb*
889 *Cortex*, 24(9), 2464-2475.
- 890 McNorgan, C., & Booth, J. R. (2015). Skill dependent audiovisual integration in the
891 fusiform induces repetition suppression. *Brain and Language*, 141, 110-123.
- 892 McNorgan, C., Randazzo-Wagner, M., & Booth, J. R. (2013). Cross-modal
893 integration in the brain is related to phonological awareness only in typical
894 readers, not in those with reading difficulty. *Front Hum Neurosci*, 7(388),
895 388.
- 896 Pan, J., McBride-Chang, C., Shu, H., Liu, H., Zhang, Y., & Li, H. (2011). What is in
897 the naming? A 5-year longitudinal study of early rapid naming and
898 phonological sensitivity in relation to subsequent reading skills in both native

- 899 Chinese and English as a second language. *Journal of Educational*
- 900 *Psychology*, 103(4), 897-908.
- 901 Perfetti, C., Cao, F., & Booth, J. (2013). Specialization and Universals in the
- 902 Development of Reading Skill: How Chinese Research Informs a Universal
- 903 Science of Reading. *Sci Stud Read*, 17(1), 5-21.
- 904 Perrachione, T. K., & Ghosh, S. S. (2013). Optimized Design and Analysis of Sparse-
- 905 Sampling fMRI Experiments. *Frontiers in Neuroscience*, 7(55).
- 906 Plewko, J., Chyl, K., Bola, Ł., Łuniewska, M., Dębska, A., Banaszkiewicz, A., . . .
- 907 Jednoróg, K. (2018). Letter and Speech Sound Association in Emerging
- 908 Readers With Familial Risk of Dyslexia. *Front Hum Neurosci*, 12(393).
- 909 Raij, T., Uutela, K., & Hari, R. (2000). Audiovisual integration of letters in the
- 910 human brain. *Neuron*, 28(2), 617-625.
- 911 Richlan, F. (2014). Functional neuroanatomy of developmental dyslexia: the role of
- 912 orthographic depth. *Front Hum Neurosci*, 8, 347.
- 913 Richlan, F. (2019). The Functional Neuroanatomy of Letter-Speech Sound
- 914 Integration and Its Relation to Brain Abnormalities in Developmental
- 915 Dyslexia. *Front Hum Neurosci*, 13(21).
- 916 Ruan, Y., Georgiou, G. K., Song, S., Li, Y., & Shu, H. (2018). Does writing system
- 917 influence the associations between phonological awareness, morphological
- 918 awareness, and reading? A meta-analysis. *Journal of Educational*
- 919 *Psychology*, 110(2), 180.

- 920 Rueckl, J. G., Paz-Alonso, P. M., Molfese, P. J., Kuo, W. J., Bick, A., Frost, S. J., . . .
- 921 Frost, R. (2015). Universal brain signature of proficient reading: Evidence
- 922 from four contrasting languages. *Proc Natl Acad Sci U S A*, 112(50), 15510-
- 923 15515.
- 924 Seghier, M. L., & Price, C. J. (2018). Interpreting and Utilising Intersubject
- 925 Variability in Brain Function. *Trends Cogn Sci*, 22(6), 517-530.
- 926 Shu, H., Chen, X., Anderson, R. C., Wu, N., & Xuan, Y. (2003). Properties of school
- 927 Chinese: Implications for learning to read. *Child development*, 74(1), 27-47.
- 928 Shu, H., McBride-Chang, C., Wu, S., & Liu, H. Y. (2006). Understanding Chinese
- 929 developmental dyslexia: Morphological awareness as a core cognitive
- 930 construct. *Journal of Educational Psychology*, 98(1), 122-133.
- 931 Shu, H., Peng, H., & McBride-Chang, C. (2008). Phonological awareness in young
- 932 Chinese children. *Dev Sci*, 11(1), 171-181.
- 933 Siok, W. T., & Fletcher, P. (2001). The role of phonological awareness and visual-
- 934 orthographic skills in Chinese reading acquisition. *Dev Psychol*, 37(6), 886-
- 935 899.
- 936 Stein, B. E., & Stanford, T. R. (2008). Multisensory integration: current issues from
- 937 the perspective of the single neuron. *Nat Rev Neurosci*, 9(4), 255-266.
- 938 Talavage, T. M., & Hall, D. A. (2012). How challenges in auditory fMRI led to
- 939 general advancements for the field. *Neuroimage*, 62(2), 641-647.

- 940 Thiebaut de Schotten, M., Cohen, L., Amemiya, E., Braga, L. W., & Dehaene, S.
941 (2014). Learning to read improves the structure of the arcuate fasciculus.
942 *Cereb Cortex*, 24(4), 989-995.
- 943 van Atteveldt, N., Formisano, E., Goebel, R., & Blomert, L. (2004). Integration of
944 letters and speech sounds in the human brain. *Neuron*, 43(2), 271-282.
- 945 van Atteveldt, N. M., Formisano, E., Blomert, L., & Goebel, R. (2007). The effect of
946 temporal asynchrony on the multisensory integration of letters and speech
947 sounds. *Cereb Cortex*, 17(4), 962-974.
- 948 van Atteveldt, N. M., Formisano, E., Goebel, R., & Blomert, L. (2007). Top-down
949 task effects overrule automatic multisensory responses to letter-sound pairs
950 in auditory association cortex. *Neuroimage*, 36(4), 1345-1360.
- 951 Wang, F., Karipidis, II, Pleisch, G., Fraga-Gonzalez, G., & Brem, S. (2020).
952 Development of Print-Speech Integration in the Brain of Beginning Readers
953 With Varying Reading Skills. *Front Hum Neurosci*, 14(289), 289.
- 954 Wang, J., Joanisse, M. F., & Booth, J. R. (2020). Neural representations of
955 phonology in temporal cortex scaffold longitudinal reading gains in 5- to 7-
956 year-old children. *Neuroimage*, 207, 116359.
- 957 Wang, X., Zhao, R., Zevin, J. D., & Yang, J. (2016). The Neural Correlates of the
958 Interaction between Semantic and Phonological Processing for Chinese
959 Character Reading. *Front Psychol*, 7.
- 960 Wechsler, D. (1974). *WISC-R, Wechsler intelligence scale for children, revised:*
961 Psychological Corporation.

- 962 Wu, C. Y., Ho, M. H., & Chen, S. H. (2012). A meta-analysis of fMRI studies on
963 Chinese orthographic, phonological, and semantic processing. *Neuroimage*,
964 63(1), 381-391.
- 965 Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: a network visualization tool
966 for human brain connectomics. *PLoS One*, 8(7), e68910.
- 967 Xia, Z., Hoeft, F., Zhang, L., & Shu, H. (2016). Neuroanatomical anomalies of
968 dyslexia: Disambiguating the effects of disorder, performance, and
969 maturation. *Neuropsychologia*, 81, 68-78.
- 970 Xia, Z., Zhang, L., Hoeft, F., Gu, B., Gong, G., & Shu, H. (2018). Neural correlates of
971 oral word reading, silent reading comprehension, and cognitive
972 subcomponents. *International Journal of Behavioral Development*, 0(0),
973 0165025417727872.
- 974 Xing, H., Shu, H., & Li, P. (2004). The acquisition of Chinese characters: Corpus
975 analyses and connectionist simulations. *Journal of Cognitive Science*, 5(1), 1-
976 49.
- 977 Xu, W., Kolozsvari, O. B., Monto, S. P., & Hämäläinen, J. A. (2018). Brain
978 Responses to Letters and Speech Sounds and Their Correlations With
979 Cognitive Skills Related to Reading in Children. *Front Hum Neurosci*,
980 12(304).
- 981 Xu, W., Kolozsvari, O. B., Oostenveld, R., & Hamalainen, J. A. (2020). Rapid
982 changes in brain activity during learning of grapheme-phoneme associations
983 in adults. *Neuroimage*, 220, 117058.

- 984 Xu, W., Kolozsvari, O. B., Oostenveld, R., Leppanen, P. H. T., & Hamalainen, J. A.
985 (2019). Audiovisual Processing of Chinese Characters Elicits Suppression and
986 Congruency Effects in MEG. *Front Hum Neurosci*, 13(18), 18.
- 987 Xue, J., Shu, H., Li, H., Li, W., & Tian, X. J. J. o. P. R. (2013). The Stability of
988 Literacy-Related Cognitive Contributions to Chinese Character Naming and
989 Reading Fluency. 42(5), 433-450.
- 990 Yan, M., Miller, K. F., Li, H., & Shu, H. (2008). What is the place for Pinyin in
991 beginning Chinese reading? Evidence from eye movements. In *Cognitive and*
992 *cultural influences on eye movements*. (pp. 343-360). Tianjin, China: Tianjin
993 People's Publishing House.
- 994 Yang, J., Shu, H., McCandliss, B. D., & Zevin, J. D. (2013). Orthographic influences
995 on division of labor in learning to read Chinese and English: Insights from
996 computational modeling. *Biling (Camb Engl)*, 16(Spec Iss 2), 354-366.
- 997 Yang, J., Zevin, J. D., Shu, H., McCandliss, B. D., & Li, P. (2006). A “triangle model”
998 of Chinese reading. Paper presented at the 28th annual conference of the
999 cognitive science society, Mahwah, NJ.
- 1000 Yin, L., Li, W., Chen, X., Anderson, R. C., Zhang, J., Shu, H., & Jiang, W. (2011).
1001 The role of tone awareness and pinyin knowledge in Chinese reading. *Writing*
1002 *Systems Research*, 3(1), 59-68.
- 1003 Zhang, S., Georgiou, G. K., Inoue, T., Zhong, W., & Shu, H. (2020). Do pinyin and
1004 character recognition help each other grow? *Early Childhood Research*
1005 *Quarterly*, 53, 476-483.

1006 Zhang, Y., Zhang, L., Shu, H., Xi, J., Wu, H., Zhang, Y., & Li, P. (2012). Universality
 1007 of categorical perception deficit in developmental dyslexia: an investigation of
 1008 Mandarin Chinese tones. *Journal of Child Psychology and Psychiatry*, 53(8),
 1009 874-882.

1010 Zhao, J., Wang, X., Frost, S. J., Sun, W., Fang, S.-Y., Mencl, W. E., . . . Rueckl, J. G.
 1011 (2014). Neural division of labor in reading is constrained by culture:
 1012 A training study of reading Chinese characters. *Cortex*, 53, 90-106.
 1013

Table 1 Demographics and behavior profiles (n = 41)

Measure	<i>Mean</i>	<i>SD</i>	<i>Range</i>
Age (month)	126	8	110-141
Grade (3 rd / 4 th / 5 th)		4 / 23 / 14	
Sex (female / male)		27 / 14	
Verbal IQ (standard score)	107	13	85-145
Performance IQ (standard score)	114	13	75-137
Full-scale IQ (standard score)	111	12	80-143
Character Recognition (corrected item)	119	12	86-146
Word List Reading (word / minute)	90	19	55-140
Silent Reading Comprehension (character / minute)	327	119	132-565
Phoneme Deletion (corrected item)	21	4	7-25
Rapid Naming (second)	18	3	11-24
Morphological Production (item)	24	4	13-30
Number of 'bad' time points per fMRI run			
Character Experiment ^a	2.4	3.37	0-10
Pinyin Experiment	0.74	1.54	0-9

Note. ^a n = 36

Table 2 Correlation between reading and cognitive-linguistic skills (n = 41)

Measures	1	2	3	4	5	6
1 Character Recognition	---					
2 Word List Reading	$r = 0.559$ $p < 0.001$	---				
3 Silent Reading Comprehension	$r = 0.647$ $p < 0.001$	$r = 0.701$ $p < 0.001$	---			
4 Phoneme Deletion	$r = 0.286$ $p = 0.070$	$r = 0.288$ $p = 0.068$	$r = 0.135$ $p = 0.400$	---		
5 Rapid Naming	$r = -0.357$ $p = 0.022$	$r = -0.597$ $p < 0.001$	$r = -0.332$ $p = 0.034$	$r = -0.160$ $p = 0.319$	---	
6 Morphological Production	$r = 0.566$ $p < 0.001$	$r = 0.344$ $p = 0.028$	$r = 0.487$ $p = 0.001$	$r = 0.122$ $p = 0.446$	$r = -0.117$ $p = 0.466$	---

Table 3 Cluster identified as showing audiovisual additive enhancement in each experiment

Experiment	# Voxel	Brain area
<i>Character</i>	822	Temporal_Sup_R, Temporal_Mid_R
	354	Temporal_Sup_L, Temporal_Mid_L
<i>Pinyin</i>	2752	Temporal_Mid_L, Temporal_Sup_L, Rolandic_Oper_L, Occipital_Mid_L, Temporal_Inf_L, SupraMarginal_L, Fusiform_L, Postcentral_L, Heschl_L, Temporal_Pole_Sup_L, Occipital_Inf_L
	1532	Temporal_Sup_R, Temporal_Mid_R, SupraMarginal_R
	967	Temporal_Mid_R, Temporal_Inf_R, Fusiform_R, Cerebelum_6_R, Occipital_Inf_R, Occipital_Mid_R
	943	Precentral_L, Frontal_Inf_Oper_L, Frontal_Inf_Tri_L, Rolandic_Oper_L
	936	Putamen_L, Thalamus_L, Insula_L, Vermis_3

Note. Brain area labeling is based on the AAL atlas (> 20 overlapping voxels).

Table 4 Results of the voxel-wise whole brain analyses on congruency effect and congruency-reading relationship

Experiment and contrast	Brain area (AAL)	Direction	# Voxel	Peak T	X	Y	Z
<i>Character</i>							
Congruency / Incongruency	Temporal_Mid_L, Temporal_Sup_L, Temporal_Pole_Sup_L, Rolandic_Oper_L, Hippocampus_L Precuneus_R, Parietal_Inf_L, Parietal_Sup_L, Precuneus_L, Cuneus_R, Angular_L, Precentral_L, Occipital_Sup_L, Paracentral_Lobule_L, Occipital_Sup_R, occipital_Mid_L Temporal_Sup_R, Temporal_Mid_R, Temporal_Pole_Sup_R, Temporal_Inf_R Precentral_L, Frontal_Inf_Tri_L, Frontal_Mid_L, Frontal_Inf_Oper_L	avC < avI avC < avI avC < avI	3310 3308 2319 2894	-7.92 -5.79	-54 -36	-22 -62	-2 48
Congruency-ORF Correlation		n.s.					
Congruency-SRC Correlation	Frontal_Mid_L, Precentral_L	positive	558	4.07	-32	10	42
<i>Pinyin</i>							
Congruency / Incongruency		n.s.					
Congruency-ORF correlation	Temporal_Sup_L, Temporal_Mid_L, Thalamus_L	positive	468	4.32	-40	-42	6

Congruency-SRC correlation	<i>n.s.</i>
----------------------------	-------------

Note: Brain area labeling is based on the AAL atlas (> 20 overlapping voxels). Abbreviations: avC = audiovisual congruent, avI = audiovisual incongruent, ORF = oral reading fluency, SRC = silent reading comprehension.

Figure 1 Materials and experimental design. Example stimuli used in the character (A) and pinyin (B) experiments. (C) Schematic illustration of the fMRI procedure. (D) Auditory and visual targets in the in-scanner task.

1022

1023 **Figure 2** Results of the character experiment. (A) Clusters (yellow) showing

audiovisual additive enhancement are identified with the mean rule (Beauchamp, 2005) consisting of two criteria: (1) The region should activate in both unisensory conditions (i.e., $Aud > 0$, $Vis > 0$); (2) The audiovisual congruent stimuli should induce stronger activation than the mean activation of the unisensory conditions, i.e., $avC > (Aud + Vis) / 2$. An uncorrected threshold of $p\text{-voxel} < 0.05$ is used for each contrast, resulting in a conjoint probability of $p\text{-voxel} < 0.05^3 = 0.000125$. A cluster containing no less than 200 continuous voxels was considered to show significant additive enhancement. (B) Areas showing a significant congruency effect (blue = incongruency, i.e., $avC < avI$). Threshold: cluster forming $p\text{-voxel} < 0.005$, Family-Wise Error (FWE) corrected $p\text{-cluster} < 0.05$. (C) The region showing a significant correlation between the magnitude of congruency effect and reading scores. Threshold: cluster forming $p\text{-voxel} < 0.005$, FWE corrected $p\text{-cluster} < 0.05$. (D, E, F) The areas showing both additive enhancement and congruency effect (red cluster), congruency effect only (green cluster), and both congruency effect and congruency-reading correlation (violet cluster) were defined as the regions-of-interest (ROIs). For each ROI, a bar plot displaying activation for each condition and a boxplot displaying the contrast $avC - (Aud + Vis) / 2$ are presented. A scatter plot is presented in the ROI showing a significant correlation between the magnitude of congruency effect and reading (residuals were obtained by regressing out nuisance variables of age and sex with a general linear model). *Abbreviations*: Aud = unisensory auditory, Vis = unisensory visual, avC = audiovisual congruent, avI = audiovisual incongruent, L = left hemisphere, R = right hemisphere; *** $p < 0.001$, *n.s.* = non-significant.

Figure 3 Results of the pinyin experiment. (A) Clusters (yellow) showing audiovisual additive enhancement are identified with the mean rule (Beauchamp, 2005), which consists of two criteria: (1) The region should activate in both unisensory conditions (i.e., $Aud > 0$, $Vis > 0$); (2) The audiovisual congruent stimuli pairs should induce stronger activation than the average activation of the auditory and visual conditions, i.e., $avC > (Aud + Vis) / 2$. An uncorrected threshold p -voxel < 0.05 is used for each contrast, resulting in a conjoint probability of p -voxel $< 0.05^3 = 0.000125$. A cluster consisting of no less than 200 continuous voxels under the threshold is considered to show additive enhancement. (B) A cluster displaying a significant correlation between the magnitude of congruency effect and oral reading fluency. Threshold: cluster forming p -voxel < 0.005 , Family-Wise Error (FWE) corrected p -cluster < 0.05 . (C) The intersection between areas showing mean additive audiovisual enhancement

1059 and areas showing association with reading is defined as the region-of-interest (ROI).
 1060 The bar plot shows activation in each condition. There was no significant difference
 1061 between the audiovisual congruent condition (avC) and incongruent condition (avI).
 1062 Boxplot displays the contrast $\text{avC} - (\text{Aud}, \text{Vis}) / 2$. The scatter plot shows the
 1063 correlation between the magnitude of congruency effect and oral reading fluency
 1064 (residuals were obtained by regressing out nuisance variables of age and sex with a
 1065 general linear model). *Abbreviations*: Aud = unisensory auditory, Vis = unisensory
 1066 visual, avC = audiovisual congruent, avI = audiovisual incongruent, L = left
 1067 hemisphere, R = right hemisphere; *n.s.* = non-significant.

Figure 4 (A) The regions-of-interest (ROIs) in the left superior temporal cortex defined in the character experiment (green) and pinyin experiment (red), as well as their intersection (yellow), are presented on a template brain. (B) The scatter plot shows the significant correlation between the magnitude of congruency effect in the character ROI and the pinyin ROI (residuals were obtained by regressing out nuisance variables of age and sex with a general linear model).