Lipid membranes modulate the activity of RNA through sequence-dependent interactions

Authors: Tomasz Czerniak^a and James P Saenz^{a*}

^a Technische Universität Dresden, B CUBE Center for Molecular Bioengineering, 01307 Dresden, Germany

*corresponding author and lead contact: james.saenz@tu-dresden.de

Abstract

RNA is a ubiquitous biomolecule that can serve as both catalyst and information carrier. Understanding how RNA activity is controlled and how it in turn regulates bioactivity is crucial for elucidating its physiological roles and potential applications in synthetic biology. Here we show that lipid membranes can act as RNA organization platforms, introducing a novel mechanism for ribo-regulation. The activity of R3C ribozyme can be modified by the presence of lipid membranes, with direct RNA-lipid interactions dependent on RNA nucleotide content, base pairing and length. In particular, the presence of guanine in short RNAs is crucial for RNA-lipid interactions, while double-stranded RNAs further increase lipid-binding affinity. Lastly, by artificially modifying the R3C substrate sequence to enhance membrane binding we generated a lipid-sensitive ribozyme reaction with riboswitch-like behavior. These findings introduce RNA-lipid interactions as a tool for developing synthetic riboswitches and novel RNA-based lipid biosensors, and bear significant implications for RNA World scenarios for the origin of life.

Introduction

RNA performs diverse functions ranging from information storage to regulation of other biomolecules and direct catalysis of biochemical reactions. The functional versatility of RNA has implications for understanding plausible scenarios for the origin of life¹⁻³, and for developing tools in synthetic biology⁴⁻⁶. Research aimed at understanding how an RNA World could have emerged has motivated development of ribozymes with functions including RNA ligation⁷⁻¹⁰, replication^{11,12} and other activities¹³⁻¹⁶. The experimental development of functional RNAs raises the possibility of RNA-based synthetic systems. For both synthetic life and understanding the origin of self-replicating organisms, RNA has intrinsic appeal: it can serve functions of both DNA (information storage) and proteins (enzymes), obviating the need for translation machineries and protein chaperones. Furthermore, RNA does not undergo irreversible denaturation like proteins, lending robustness against a broad range of physical and chemical conditions. In order to design a system based on RNA, however, it is essential to be able to coordinate RNA activity in space and time.

Key to harnessing the functional versatility of RNA is understanding how to spatially and temporally modulate its properties and to selectively modulate the activity of different RNAs within one system. The physicochemical environment surrounding an RNA molecule is a central determinant of its structure, stability and activity. Spontaneous RNA hydrolysis and ligation, as well as catalytic RNA activity are sensitive to pH¹⁷, ionic strength^{18,19} and RNA concentration changes¹¹, among other parameters. Similarly, RNA activity can be modulated by interactions with molecules such as ions,

proteins, and other nucleic acids²⁰. Thus, one approach to regulating RNA activity could be via tunable interactions with binding partners that affect RNA structure, concentration or chemical microenvironment.

One mechanism for modulating RNA activity could be through direct RNA-lipid interactions^{19,21,22}. Because of their amphiphilic nature, lipids spontaneously self-assemble into membranous structures that can encapsulate RNA into protected and selective microcompartments^{23–25}. Alternatively, direct RNA-lipid interactions could localize RNAs to membrane surfaces, increasing its local concentration and reducing dimensionality for intermolecular interactions²¹. Lastly, localization to a lipid surface brings RNA into a physicochemically unique microenvironment with sharp gradients of hydrophobicity, electrical permittivity, and water activity. Through these effects, RNA-lipid interactions could provide a powerful mechanism for modulating RNA activity.

The first functional RNA-lipid interaction was described more than 40 years ago²⁶, with subsequent research revealing various factors that facilitate nucleic acid-lipid binding^{27–37}. More recently, specific RNA sequences have been generated through SELEX with affinity for fluid membranes comprised of phospholipids and cholesterol^{38–40}. Interestingly, mixtures of RNAs have also been shown to bind to membranes that are in a solid crystalline (gel) phase^{41,42}. These studies revealed that, while most randomized mixtures of RNA sequences can bind to gel membranes, there is a relatively small chemical space of oligomers that have affinity for fluid membranes. Thus, conceptually, gel membranes could provide a platform for modulating the activity of a diverse range of RNAs. However, the effects of gel membranes on RNA activity and the sequence selectivity of such interactions are relatively unexplored.

This study reports the effect of lipid membranes on RNA catalytic activity. We show that RNA-lipid binding depends on the primary sequence, secondary structure, and length of RNA. Using the trans-acting R3C ligase ribozyme, we observed that R3C-lipid binding changes ribozyme activity in a concentration-dependent manner. Lipid binding assays show that the interaction of short RNA sequences with gel membranes depends on guanine content and the presence of double-stranded structures. Lastly, modification of R3C's substrate sequence increased the tunability of R3C-based reactions through a lipid-dependent mechanism. Our findings demonstrate that membranes can serve as platforms for ribo-regulation, which could contribute to the development of RNA-based lipid biosensors and lipid-sensitive riboswitches. This approach introduces new tools for molecular and synthetic biology and raises the prospect of previously unrecognized roles for RNA-lipid interactions in the origin and evolution of life.

Results

The discovery that RNA can catalyse reactions in addition to encoding information^{18,43}, opened new directions for engineering life and the possibility of protocells emerging from an RNA world². But, a key missing ingredient for RNA-based systems is a mechanism to organize RNAs and regulate their activity. We hypothesized that RNA-membrane interactions could influence ribozyme activity by changing local RNA concentrations at the membrane surface or influencing RNA conformations.

To test the hypothesis that lipid membranes can serve as platforms for ribo-regulation, we took advantage of the observation that RNA can interact with solid crystalline (gel) phase membranes composed of phosphatidylcholine lipids^{41,42}. To visibly demonstrate this phenomenon we prepared giant unilamellar vesicles (GUVs) that are phase separated into gel and liquid domains and we then

observed the distribution of a random mixture of RNA oligomers stained with SybrGold by fluorescence microscopy (**Fig. 1a**). A fluorescent lipid probe (DiD) was enriched in membrane liquid domains and was excluded from gel phase domains. In contrast, RNA was enriched within the gel phase domains. RNA enrichment on GUVs was reversed by heating the system above the gel phase melting temperature, resulting in an entirely liquid phase vesicle (**Fig. 1b**). Binding of RNA oligonucleotides to gel-phase small unilamellar vesicles (SUVs) can also lead to aggregation into large visible RNA-lipid assemblies (**Fig. 2a**)^{27,35,41,42}. Depletion of divalent cations or increasing temperature above the lipid melting temperature (i.e. producing fluid instead of gel membranes) reduced RNA-membrane binding and aggregation (**Fig. 2a**). Taken together, reversible RNA-lipid binding and lipid-dependent aggregation could provide a tunable mechanism to concentrate and regulate RNAs in simple bottom up synthetic systems, or in a prebiotic environment.

RNA-lipid binding changes the probability of RNA-RNA interactions

To determine whether local RNA concentration is influenced by interaction with lipid membranes, we relied on UV-mediated crosslinking. All small vesicle membranes in this study were composed of dipalmitoylphosphatidylcholine (DPPC), which is in a solid crystalline (gel) phase at the experimental temperature of 24°C. Nucleic acid bases absorb UV light, producing chemical changes that yield base-base covalent bonds in a distance-dependent manner, yielding insights into the structure and interactions of nucleic acids^{44–46}. We first observed the effect of gel membranes on the crosslinking of a single defined RNA sequence, the R3C ligase, which predominately forms one structure (R3C ligase, **Fig. 2d**). When subjected to UV, the R3C ligase shows slightly increased crosslinking in the presence of low lipid concentrations. A further increase in lipid concentration led to decreased crosslinking efficiency, possibly through dilution of the RNA species on the surface of the lipid membranes. This effect was inversely correlated with RNA-lipid binding efficiency (**Fig. 2b**). In contrast, a mixture of RNA oligomers with randomized sequences which is able to create a more diverse range of intermolecular structures (**Fig. 2d**) showed continuously increasing and overall higher UV-crosslinking efficiency in the presence of lipids (**Fig. 2c**). The contrasting effect of crosslinking for R3C and randomized oligomers might be explained by the influence of lipid membranes on RNA interactions: membrane binding could potentially stabilize base-pairing⁴⁷ and thereby enhance crosslinking efficiency, whereas the decrease of crosslinking might be caused by a dilution effect at higher membrane surface areas or by lipid-based steric hindrance. These observations demonstrate that RNA-membrane interactions can influence RNA-RNA interactions in a manner that is dependent on lipid concentration and RNA sequence diversity. This further suggests that membrane binding could have an effect on trans-acting ribozyme activity derived from base pairing.

RNA-lipid interactions influence RNA catalytic activity

To investigate the functional consequence of RNA-lipid binding, we tested the effect of lipids on the activity of the trans-acting R3C ligase ribozyme. R3C ligase is a relatively short ribozyme that catalyzes ligation of substrate strands to the ribozyme¹⁰(**Fig. 3a**). Since R3C is also part of an RNA self-replication system^{11,12}, the effects of lipids on R3C are also interesting with regard to the emergence, evolution and artificial synthesis of autonomous self-replicating systems. The R3C reaction rate in the absence of membranes was 11 pM/min, with a reaction constant rate of $k = 6.4 \times 10^{-4} \text{ min}^{-1}$. Addition of lipid vesicles led to an increased reaction rate (14 pM/min, +29%), which could plausibly be due to increased ligase-substrate interaction on the membrane either through increased concentrations at the membrane surface or through stabilization of base pairing⁴⁷. At the highest lipid concentration, the reaction rate dropped to 8 pM/min (-27%) (**Fig. 3b**). A decrease in reaction rate was correlated with

further R3C ligase binding to the lipid membranes (>100 lipid/R3C, **Fig. 2b**) and we speculate that this may be caused by steric hindrance from the lipid vesicles. These results show that gel membranes can influence the reaction rate of a trans-acting ribozyme in a lipid concentration-dependent manner.

R3C ligase and its substrate bind differently to the membrane

To further understand how R3C and substrate membrane binding might contribute to changes in ligation rates, we analyzed lipid-binding affinities of the ligase and its substrate. To quantify binding to the membranes, we calculated lipid–buffer partition coefficients (K). The partition coefficient of the short 12 nt substrate was more than 3 orders of magnitude lower than that of the 84 nt R3C ligase (**Fig. 3c**). Thus, in this system, substrate-membrane binding is essentially negligible compared with R3C-membrane binding. Combined with our observation that RNA-RNA interactions are stabilized by membrane binding (**Fig. 2c**), this indicates that the increased activity observed in **Figure 3b** was due to an effect of the membrane on the R3C ligase-substrate complex stability, rather than through substrate-lipid interactions on the membrane surface. Thus, membranes can enhance ribozyme activity by promoting the stability of reaction intermediates.

RNA sequence influences binding to gel membranes

Understanding the features that influence RNA-lipid interactions could allow us to engineer RNAs with higher membrane affinity, and thereby potentially enhance activity by concentrating RNAs at the membrane surface. Interactions of RNA with gel-phase membranes have been previously shown to exhibit low sequence-dependence, based on binding of oligos with randomized sequences^{41,42}. However, the large difference in binding for R3C and its substrate suggests that RNA-gel membrane interactions might in fact be dependent on RNA composition or structure.

To investigate which features of RNA influence its interaction with lipid membranes we first evaluated how base content influences membrane binding of short RNA species. 19nt oligomers with only one type of nucleotide were tested together with a random sequence as a control. To quantify relative changes of RNA-membrane binding, we calculated the RNA binding efficiency as a percentage of total RNA bound to membranes at a fixed oligo-to-lipid ratio. Remarkably, whereas the 19xG oligomer bound very efficiently to the gel-phase membrane (84% binding efficiency), oligomers of A, C or U showed practically no binding. A mixture of 19-mer RNA with randomized sequences showed moderate binding (33% binding efficiency), as did one with AG-repeats (**Fig. 4a**), most probably due to the introduction of guanine. Guanine could conceivably enhance binding either through direct interactions with the membrane, or alternatively by promoting G≡C base pairing or presence of G-quadruplexes, thereby influencing structure. The enhanced binding of an oligomer with only AG-repeats suggests that membrane binding can be influenced by direct guanine-membrane interactions⁴⁸.

We next examined the effect of deleting a specific base from mixtures of 40-mer oligos with randomized sequences. Depletion of A, C or U led to a slight decrease in membrane binding, whereas removal of G showed a significant binding decrease. The decrease in binding affinity in all of the base-depleted oligos (**Fig. 4b**) indicates that base-pairing may influence membrane binding. However, the larger effect of G depletion indicates that either depletion of G causes highest bias on RNA base-pairing or base pairing is not the only factor responsible for membrane binding, consistent with our previous observation that G-content alone can influence binding (**Fig. 4a**). Finally, by comparing 19-mer and 40-

mer randomized oligos, we observed that increasing RNA length also led to enhanced binding, possibly due to increased length and G content (**Fig. 4 a,b**).

To further determine how guanine content affects membrane binding, lipid-buffer partition coefficients were determined for short oligomers with varying G-content. RNA membrane-buffer partition coefficients indicate how well a particular molecule binds to the membrane by comparing the relative amount of RNA in buffer and on the lipid membrane. The RNA oligos measured in this study had partition coefficients around 10^5 , which is in the range of partition coefficients determined for hydrophobic peptides⁴⁹. Remarkably, addition of 2 G's per oligo led to a 2.7-fold increase of the partition coefficient relative to guanine-depleted RNA. Oligos with 4 and more guanine residues showed a further increase of partition coefficient values with highest value for 9 G's. This result shows that the presence of guanine affects RNA-lipid binding and even low guanine content significantly increases membrane binding efficiency (**Fig. 4c**).

RNA-membrane binding is dependent on RNA base pairing

Our observations of the effect of base depletion on oligomer binding suggested that base-pairing may be a key factor in membrane binding. To investigate the effect of base-pairing on membrane interactions, we estimated binding efficiency for ssRNA and dsRNA (composed of 100% complementary sequences) species. We observed that not only was the membrane binding efficiency higher for dsRNA compared with ssRNA species, but also that only dsRNA promoted vesicle aggregation (**Fig. 5a**). It is interesting to note that the difference in binding efficiency of the two ssRNAs, which have identical base composition but different sequence, is consistent with the importance of sequence as a key determinant of membrane binding. We further observed that dsRNA interacts with fewer lipids than ssRNA, implying different models of binding (**Fig. 5b**). Thus, membrane binding efficiency can be dependent on the propensity of an RNA molecule to form intra- or inter-molecular structures through base pairing. Since base pairing is a fundamental element of RNA secondary structure, our results indicate that structure in general can influence the selectivity of RNA-lipid interactions.

To gain further insight into the effect of base pairing-dependent intra- and inter-molecular interactions, we estimated the lipid-to-nucleotide binding ratios of ssRNA and dsRNA (**Fig. 5c**). By calculating the molecular ratio of lipid/nucleotide binding, we introduce a normalization factor that allows us to directly compare membrane binding for oligos of varying length. Furthermore, varying lipid to nucleotide stoichiometry reflects changes in the average lipid/nucleotide interaction strength. With this estimate, we compared lipid/nucleotide binding for ssRNA and dsRNA alongside oligos with varying guanine content from our previous analysis (**Fig. 4c**). The differences between ssRNA and dsRNA, all containing 9 guanines per oligo, confirmed our previous observation that sequence variation (with constant base content) and base pairing can both influence binding. However, the reason for a change in lipid/nucleotide ratio for oligos with varying guanine content was not readily apparent. Analysis of oligos with varying guanine content by native non-denaturing gel revealed that the presence of guanine promotes a larger spread in migration (**Fig. 5c, inset**), indicating that guanines promote intermolecular interactions. These observations show that guanine can enhance binding by promoting intermolecular interactions, and further confirm the importance of base pairing in RNA-lipid interactions.

Modifying R3C substrate sequence enhances membrane binding and modifies reaction rates

In previous experiments, we observed that the binding affinity of the short substrate was >3 orders of magnitude lower than for the R3C ligase and that reaction rates correlated strongly with R3C-membrane binding (**Fig. 3c, d**). We therefore reasoned that by modifying the R3C ligase substrate to enhance membrane binding, we might be able to further modulate ligase activity through increased RNA density at the membrane surface. Having determined that guanine content is a key factor influencing RNA-lipid binding (**Fig. 4**), we modified the R3C substrate sequence to increase lipid membrane binding efficiency by addition of a 5'-overhang with varying amounts of AG repeats (**Fig. 6a**).

We first assessed the effect of modifying the R3C ligase substrate on ligation rate in the absence of membranes (**Fig. 6a**). We observed that increasing the guanine content decreased ligase activity, suggesting that 5' modification of the substrate was inhibiting the reaction either through steric hindrance or promotion of inhibitory inter- or intra-molecular interactions. To determine whether AG-rich substrates form intra-molecular structures⁵⁰ or exhibit reduced binding with the R3C ligase, we performed electrophoresis in native conditions with and without ligase (**Fig. 6b**). Firstly, the non-modified short substrate migrated as one band and was fully bound to the R3C ligase. Secondly, a 4x(AG) 20 nt substrate variant migrated as two bands, which suggests that part of the substrate molecules have some folding that might bias binding to the R3C ligase. In the presence of ligase, 30% of the substrate was non-bound, which correlated with a 22% reduction in reaction rate. For the longest substrate variant (31 nt), electrophoretic mobility suggested higher folding, since only one band was present and migrated half-way between the non-folded 12 nt and 20 nt substrates. 70% of the 9x(AG) substrate was not bound to the ligase, which correlates with a reduced R3C reaction rate. Lastly, the AG-rich oligomer did not co-migrate with the R3C ligase, indicating that the inhibition effect is not based on non-specific substrate-ligase interactions (not shown). Therefore, increasing guanine content reduces R3C ligase-substrate binding, likely accounting for decreased activity.

We next investigated whether varying guanine content through 5' modification of the substrate influences membrane binding and, consequently, ligation activity in the presence of membranes. As expected, the longer guanine-containing R3C substrates showed significantly higher membrane binding compared with the shorter variant (**Fig. 6c**). To determine whether binding of the substrates to the membrane enhances ligase activity, we determined R3C reaction rates in the presence of the gel lipid membranes (**Fig. 6d**). For modified substrates, the reaction rates increased significantly to the level of the unmodified substrate. Interestingly, none of the modified substrates exhibited higher activity than the unmodified substrate, indicating that the rescue of activity of the other guanine variants is not due to increased substrate-ligase interactions through enhanced binding to the membrane, but rather due to interaction of the membrane with 5' overhangs. Thus, substrate-membrane interaction increases activity by rescuing the inhibitory effect of the guanine-rich 5' overhangs (**Fig. 6e**).

In summary, by modifying the R3C ribozyme substrate with an additional short sequence containing guanine residues, we increased membrane affinity. The addition of guanines, however, also introduced an inhibitory effect on ligation activity that was reversed through the introduction of gel-membranes to the reaction. Thus, we unexpectedly produced a lipid-sensitive riboswitch-like system that could be tuned in a lipid concentration-dependent manner.

Discussion

We have demonstrated that RNA-gel membrane interactions are dependent on the nucleotide content and base pairing of ribonucleic acids. Varying the guanosine content of RNAs changed membrane-buffer partitioning by several orders of magnitude. Additionally, double-stranded RNAs exhibited higher membrane partitioning than single-stranded RNAs, indicating that secondary structure is a critical factor in RNA-membrane interactions. Although gel-membranes are not widely observed in living cells, RNA-gel membrane interactions could be employed for ribo-regulation in synthetic biological systems. Gel membranes also plausibly accumulated in ancient prebiotic scenarios, potentially serving as an organizational scaffold, as has been proposed for “ribofilms” on mineral surfaces⁵¹. Furthermore, understanding the factors involved in RNA-gel membrane interactions should provide insights that can be used to design RNA sequences with affinity for physiologically relevant fluid membranes. Indeed, our finding that guanosine is a key factor in promoting RNA-gel membrane partitioning is consistent with previous speculation that guanosine residues might play a role in the binding of RNA aptamers to fluid membranes³⁸. Taken together, the sequence and structural dependence of RNA-gel membrane interactions shown here provides the necessary framework to design synthetic RNAs with varying membrane affinities.

Our observation of riboswitch-like behavior suggests a novel mechanism for regulation of RNA through membrane interactions. Riboswitches are RNAs with regulatory effects that are derived from sensitivity to physicochemical conditions or through binding with an interaction partner. Riboswitch regulation through temperature changes, interaction with nucleic acids and small molecules have been described previously^{52–56}. However, lipid sensitive riboswitches have not been documented. It is now plausible that spontaneously self-assembling lipid membranes in ancient environments could have served as an organizational scaffold regulating the activity of RNAs in a sequence- and structure-dependent manner. Furthermore, the importance of guanosine in membrane binding suggests a possible mechanism for its selection as one of the canonical nucleic acids. In extant organisms, synthetic or naturally occurring RNAs exhibiting sensitivity to lipids could potentially act as biosensors or riboswitches. By demonstrating that RNA-gel membrane interactions can induce riboswitch-like behaviors, we raise the possibility that such functionalities could be an ancient mechanism of ribo-regulation and may be present among the myriad uncharacterized small RNAs in modern cells.

RNA-membrane interactions provide a means to select RNAs by sequence and structure. For example, diverse RNA species can be segregated or co-localized based on their differential membrane affinities. Furthermore, selective RNA-membrane localization can be controlled by shifting the temperature above and below the membrane gel-liquid transition temperature, thereby turning on and off RNA-membrane binding. Such lipid-dependent sorting of RNAs could provide a mechanism to selectively control the concentration of reactants in primordial scenarios and in synthetic RNA-based systems. Selective RNA-membrane interactions would also influence the sequence space explored by evolving ribozymes. Ribozymes generated through natural or artificial selection in the presence of lipid membranes may, therefore, evolve enhanced or novel functions. The sequence and structural selectivity of RNAs for membranes can also be implemented in the design of novel RNA membrane biosensors. Looking forward, RNA-membrane interactions can provide tools for selective ribo-regulation in synthetic systems, development of membrane biosensors, and might facilitate the emergence of novel functions through artificial and natural evolution of ribozymes in the presence of membranes.

In conclusion, our findings reveal that lipids, which are present in every modern cell and were plausibly part of a prebiotic world^{57–59}, can interact with RNA and change its activity. These findings have significant applications in fields such as synthetic biology, where merging the selective affinities of aptamers with ribozyme activity (aptazymes) is currently a developing field^{60–62}. Our results provide a foundation for exploring the basis of membrane-RNA interactions with physiologically relevant fluid membranes exhibited by synthetic aptamers^{16,40,63} thereby providing new regulatory mechanisms in synthetic biology through the specific and anchor-free localization of RNA molecules within lipid compartments. Furthermore, insights from the present study can already be implemented in bioengineering applications such as in improving mRNA drug delivery mechanisms and introducing lipid sensitive ribo-regulation to synthetic ribozyme networks. Finally, the interaction between RNA and lipid membranes could have provided an evolutionary pressure in ancient times through selection for base composition or secondary structural features.

Acknowledgements:

We would like to thank Mario Mörl, Ilya Levental, Robert Ernst, Andre Nadler, Dora Tang, Grzegorz Chwastek, Michał Grzybek and Mike Thompson for helpful discussions and feedback. We would also like to thank Anatol Fritsch for home-made microscopy stage-top temperature-control device. This work was supported by the B CUBE of the TU Dresden, a Simons Foundation Fellowship (to J.S.), a German Federal Ministry of Education and Research BMBF grant (to J.S., project 03Z22EN12), and a VW Foundation “Life” grant (to J.S., project 93090).

The authors declare no conflict of interests.

References

1. Pace, N. R. & Marsh, T. L. RNA catalysis and the origin of life. *Origins Life Evol B* **16**, 97–116 (1985).
2. Gilbert, W. Origin of life: The RNA world. *Nature* **319**, 618–618 (1986).
3. Rich, A. On the problems of evolution and biochemical information transfer. *Horizons in biochemistry* 103–126 (1962).
4. Breaker, R. R. Engineered allosteric ribozymes as biosensor components. *Curr Opin Biotech* **13**, 31–39 (2002).
5. Wittmann, A. & Suess, B. Engineered riboswitches: Expanding researchers’ toolbox with synthetic RNA regulators. *Febs Lett* **586**, 2076–2083 (2012).
6. Isaacs, F. J., Dwyer, D. J. & Collins, J. J. RNA synthetic biology. *Nat Biotechnol* **24**, 545–554 (2006).
7. Bartel, D. & Szostak, J. Isolation of new ribozymes from a large pool of random sequences. *Science* **261**, 1411–1418 (1993).

- 359 8. Eklund, E., Szostak, J. & Bartel, D. Structurally complex and highly active RNA ligases
360 derived from random RNA sequences. *Science* **269**, 364–370 (1995).
- 361 9. Robertson, M. P., Hesselberth, J. R. & Ellington, A. D. Optimization and optimality of a
362 short ribozyme ligase that joins non-Watson–Crick base pairings. *Rna* **7**, 513–523 (2001).
- 363 10. Rogers, J. & Joyce, G. F. The effect of cytidine on the structure and function of an RNA
364 ligase ribozyme. *Rna* **7**, 395–404 (2001).
- 365 11. Paul, N. & Joyce, G. F. A self-replicating ligase ribozyme. *Proc National Acad Sci* **99**,
366 12733–12740 (2002).
- 367 12. Lincoln, T. A. & Joyce, G. F. Self-Sustained Replication of an RNA Enzyme. *Science* **323**,
368 1229–1232 (2009).
- 369 13. Saragliadis, A., Krajewski, S. S., Rehm, C., Narberhaus, F. & Hartig, J. S. Thermozyms:
370 synthetic RNA thermometers based on ribozyme activity. *Rna Biol* **10**, 1009–1016 (2013).
- 371 14. Ferretti, A. C. & Joyce, G. F. Kinetic Properties of an RNA Enzyme That Undergoes Self-
372 Sustained Exponential Amplification. *Biochemistry-us* **52**, 1227–1235 (2013).
- 373 15. Lam, B. J. & Joyce, G. F. Autocatalytic aptazymes enable ligand-dependent exponential
374 amplification of RNA. *Nat Biotechnol* **27**, 288–292 (2009).
- 375 16. Janas, T., Janas, T. & Yarus, M. A membrane transporter for tryptophan composed of
376 RNA. *Rna* **10**, 1541–1549 (2004).
- 377 17. Pino, S., Ciciriello, F., Costanzo, G. & Mauro, E. D. Nonenzymatic RNA Ligation in Water*.
378 *J Biol Chem* **283**, 36494–36503 (2008).
- 379 18. Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N. & Altman, S. The RNA moiety of
380 ribonuclease P is the catalytic subunit of the enzyme. *Cell* **35**, 849–857 (1983).
- 381 19. Anella, F. & Danelon, C. Prebiotic Factors Influencing the Activity of a Ligase Ribozyme.
382 *Life* **7**, 17 (2017).
- 383 20. Dai, X., Zhang, S. & Zaleta-Rivera, K. RNA: interactions drive functionalities. *Mol Biol Rep*
384 **47**, 1413–1434 (2020).
- 385 21. Müller, U. F. & Bartel, D. P. Improved polymerase ribozyme efficiency on hydrophobic
386 assemblies. *Rna* **14**, 552–562 (2008).
- 387 22. Anella, F. & Danelon, C. Reconciling Ligase Ribozyme Activity with Fatty Acid Vesicle
388 Stability. *Life* **4**, 929–943 (2014).
- 389 23. Chen, I. A., Salehi-Ashtiani, K. & Szostak, J. W. RNA Catalysis in Model Protocell Vesicles. *J*
390 *Am Chem Soc* **127**, 13213–13219 (2005).

- 391 24. Mansy, S. S., Schrum, J. P., Krishnamurthy, M., Tobé, S., Treco, D. A. & Szostak, J. W.
392 Template-directed synthesis of a genetic polymer in a model protocell. *Nature* **454**, 122–125
393 (2008).
- 394 25. Bianconi, G., Zhao, K., Chen, I. A. & Nowak, M. A. Selection for Replicases in Protocells.
395 *Plos Comput Biol* **9**, e1003051 (2013).
- 396 26. Budker, V. G., Kazatchkov, Yu. A. & Naumova, L. P. Polynucleotides adsorb on
397 mitochondrial and model lipid membranes in the presence of bivalent cations. *Febs Lett* **95**,
398 143–146 (1978).
- 399 27. McManus, J. J., Rädler, J. O. & Dawson, K. A. Does Calcium Turn a Zwitterionic Lipid
400 Cationic? *J Phys Chem B* **107**, 9869–9875 (2003).
- 401 28. Kharakoz, D. P., Khusainova, R. S., Gorelov, A. V. & Dawson, K. A. Stoichiometry of
402 dipalmitoylphosphatidylcholine-DNA interaction in the presence of Ca²⁺: a temperature-
403 scanning ultrasonic study. *Febs Lett* **446**, 27–29 (1999).
- 404 29. Gromelski, S. & Brezesinski, G. Adsorption of DNA to zwitterionic DMPE monolayers
405 mediated by magnesium ions. *Phys Chem Chem Phys* **6**, 5551–5556 (2004).
- 406 30. Uhríková, D., Hanulová, M., Funari, S. S., Khusainova, R. S., Šeršeň, F. & Balgavý, P. The
407 structure of DNA–DOPC aggregates formed in presence of calcium and magnesium ions: A
408 small-angle synchrotron X-ray diffraction study. *Biochimica Et Biophysica Acta Bba -*
409 *Biomembr* **1713**, 15–28 (2005).
- 410 31. Gromelski, S. & Brezesinski, G. DNA Condensation and Interaction with Zwitterionic
411 Phospholipids Mediated by Divalent Cations. *Langmuir* **22**, 6293–6301 (2006).
- 412 32. McLoughlin, D., Dias, R., Lindman, B., Cardenas, M., Nylander, T., Dawson, K., Miguel, M.
413 & Langevin, D. Surface Complexation of DNA with Insoluble Monolayers. Influence of
414 Divalent Counterions. *Langmuir* **21**, 1900–1907 (2005).
- 415 33. Michanek, A., Yanez, M., Wacklin, H., Hughes, A., Nylander, T. & Sparr, E. RNA and DNA
416 Association to Zwitterionic and Charged Monolayers at the Air–Liquid Interface. *Langmuir*
417 **28**, 9621–9633 (2012).
- 418 34. Michanek, A., Kristen, N., Höök, F., Nylander, T. & Sparr, E. RNA and DNA interactions
419 with zwitterionic and charged lipid membranes — A DSC and QCM-D study. *Biochimica Et*
420 *Biophysica Acta Bba - Biomembr* **1798**, 829–838 (2010).
- 421 35. Marty, R., N’soukpoé-Kossi, C. N., Charbonneau, D. M., Kreplak, L. & Tajmir-Riahi, H.-A.
422 Structural characterization of cationic lipid–tRNA complexes. *Nucleic Acids Res* **37**, 5197–
423 5207 (2009).
- 424 36. Suga, K., Tanabe, T., Tomita, H., Shimanouchi, T. & Umakoshi, H. Conformational change
425 of single-stranded RNAs induced by liposome binding. *Nucleic Acids Res* **39**, 8891–8900
426 (2011).

37. Lu, D. & Rhodes, D. G. Binding of phosphorothioate oligonucleotides to zwitterionic liposomes. *Biochimica Et Biophysica Acta Bba - Biomembr* **1563**, 45–52 (2002).
38. Khvorova, A., Kwak, Y.-G., Tamkun, M., Majerfeld, I. & Yarus, M. RNAs that bind and change the permeability of phospholipid membranes. *Proc National Acad Sci* **96**, 10649–10654 (1999).
39. Vlassov, A. & Yarus, M. Interaction of RNA with Phospholipid Membranes. *Mol Biol+* **36**, 389–393 (2002).
40. Vlassov, A., Khvorova, A. & Yarus, M. Binding and disruption of phospholipid bilayers by supramolecular RNA complexes. *Proc National Acad Sci* **98**, 7706–7711 (2001).
41. Janas, T., Janas, T. & Yarus, M. Specific RNA binding to ordered phospholipid bilayers. *Nucleic Acids Res* **34**, 2128–2136 (2006).
42. Pannwitt, S., Slama, K., Depoix, F., Helm, M. & Schneider, D. Against Expectations: Unassisted RNA Adsorption onto Negatively Charged Lipid Bilayers. *Langmuir* **35**, 14704–14711 (2019).
43. Kruger, K., Grabowski, P. J., Zaug, A. J., Sands, J., Gottschling, D. E. & Cech, T. R. Self-splicing RNA: Autoexcision and autocyclization of the ribosomal RNA intervening sequence of tetrahymena. *Cell* **31**, 147–157 (1982).
44. Quarless, S. A. & Cantor, C. R. Analysis of RNA structure by ultraviolet crosslinking and denaturation gel electrophoresis. *Anal Biochem* **147**, 296–300 (1985).
45. Behlen, L. S., Sampson, J. R. & Uhlenbeck, O. C. An ultraviolet light-induced crosslink in yeast tRNA phe. *Nucleic Acids Res* **20**, 4055–4059 (1992).
46. Nejedlý, K., Kittner, R., Pospíšilová, Š. & Kypr, J. Crosslinking of the complementary strands of DNA by UV light: dependence on the oligonucleotide composition of the UV irradiated DNA. *Biochimica Et Biophysica Acta Bba - Gene Struct Expr* **1517**, 365–375 (2001).
47. Lengyel, A., Uhríková, D., Klacsová, M. & Balgavý, P. DNA condensation and its thermal stability influenced by phospholipid bilayer and divalent cations. *Colloids Surfaces B Biointerfaces* **86**, 212–217 (2011).
48. Black, R. A., Blosser, M. C., Stottrup, B. L., Tavakley, R., Deamer, D. W. & Keller, S. L. Nucleobases bind to and stabilize aggregates of a prebiotic amphiphile, providing a viable mechanism for the emergence of protocells. *Proc National Acad Sci* **110**, 13272–13276 (2013).
49. White, S. H., Wimley, W. C., Ladokhin, A. S. & Hristova, K. [4] Protein folding in membranes: Determining energetics of peptide-bilayer interactions. *Methods Enzymol* **295**, 62–87 (1998).
50. Lee, J. S., Evans, D. H. & Morgan, A. R. Polypurine DNAs and RNAs form secondary structures which may be tetra-stranded. *Nucleic Acids Res* **8**, 4305–4320 (1980).

- 463 51. Baross, J. A. & Martin, W. F. The Ribofilm as a Concept for Life's Origins. *Cell* **162**, 13–15
464 (2015).
- 465 52. Johansson, J., Mandin, P., Renzoni, A., Chiaruttini, C., Springer, M. & Cossart, P. An RNA
466 Thermosensor Controls Expression of Virulence Genes in *Listeria monocytogenes*. *Cell* **110**,
467 551–561 (2002).
- 468 53. Winkler, W., Nahvi, A. & Breaker, R. R. Thiamine derivatives bind messenger RNAs
469 directly to regulate bacterial gene expression. *Nature* **419**, 952–956 (2002).
- 470 54. Morita, M. T., Tanaka, Y., Kodama, T. S., Kyogoku, Y., Yanagi, H. & Yura, T. Translational
471 induction of heat shock transcription factor ζ 32: evidence for a built-in RNA thermosensor.
472 *Gene Dev* **13**, 655–665 (1999).
- 473 55. Schilling, O., Langbein, I., Müller, M., Schmalisch, M. H. & Stülke, J. A protein-dependent
474 riboswitch controlling ptsGHI operon expression in *Bacillus subtilis*: RNA structure rather
475 than sequence provides interaction specificity. *Nucleic Acids Res* **32**, 2853–2864 (2004).
- 476 56. Nahvi, A., Sudarsan, N., Ebert, M. S., Zou, X., Brown, K. L. & Breaker, R. R. Genetic Control
477 by a Metabolite Binding mRNA. *Chem Biol* **9**, 1043–1049 (2002).
- 478 57. Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D. & Sutherland, J. D. Common origins of
479 RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nat Chem* **7**, 301–307
480 (2015).
- 481 58. Rushdi, A. I. & Simoneit, B. R. T. Lipid Formation by Aqueous Fischer-Tropsch-Type
482 Synthesis over a Temperature Range of 100 to 400 °C. *Origins Life Evol B* **31**, 103–118 (2001).
- 483 59. Brady, R. C. & Pettit, R. Mechanism of the Fischer-Tropsch reaction. The chain
484 propagation step. *J Am Chem Soc* **103**, 1287–1289 (1981).
- 485 60. Robertson, M. P. & Ellington, A. D. Design and optimization of effector-activated
486 ribozyme ligases. *Nucleic Acids Res* **28**, 1751–1759 (2000).
- 487 61. Debiais, M., Lelievre, A., Smietana, M. & Müller, S. Splitting aptamers and nucleic acid
488 enzymes for the development of advanced biosensors. *Nucleic Acids Res* **48**, 3400–3422
489 (2020).
- 490 62. Stifel, J., Spöring, M. & Hartig, J. S. Expanding the toolbox of synthetic riboswitches with
491 guanine-dependent aptazymes. *Synthetic Biology* **4**, ysy022- (2019).
- 492 63. Janas, T. & Yarus, M. Visualization of membrane RNAs. *Rna* **9**, 1353–1361 (2003).
- 493 64. Walker, S. C., Avis, J. M. & Conn, G. L. General plasmids for producing RNA in vitro
494 transcripts with homogeneous ends. *Nucleic Acids Res* **31**, e82–e82 (2003).
- 495 65. Schmidt, M. L., Ziani, L., Boudreau, M. & Davis, J. H. Phase equilibria in DOPC/DPPC:
496 Conversion from gel to subgel in two component mixtures. *J Chem Phys* **131**, 175103 (2009).

For the confocal image acquisition, 488 nm (SybrGold) and 635 nm (DiD) lasers were used at low power (<0.5%) to avoid photobleaching and the signal was collected using hybrid detectors (500 - 550 nm for SybrGold and 700 – 800 nm for DiD). Z-scans were acquired as 6-8 averaged images per layer and Z-stacked as a sum of the slides using FIJI software⁶⁶.

Temperature ramps were captured using a camera with FITC dichroics for DiD detection. 1.5 sec of exposure without binning was used to acquire each channel and measurement was repeated every 5 seconds. Temperature changes were registered and calculated to be 3.4 °C/min. Image analysis was conducted in FIJI software.

Small lipid vesicle preparation

DPPC chloroform stock was pipetted into a glass vial and briefly evaporated under a steady flow of nitrogen gas. To remove organic solvent residues, a lipid film was dried under vacuum overnight. To obtain multi-layer DPPC vesicles (MLV DPPC), lipid films were hydrated in reaction buffer, with a final lipid concentration of 10 mM in 1 mL. Buffer-lipid mixtures were shaken at 60 °C, above the melting temperature of DPPC ($T_m = 42$ °C), for one hour. A cloudy liposome suspension was freeze-and-thawed 10x and extruded 17x through 100 nm polycarbonate filters (Merck Millipore, USA) to reduce multi-lamellarity and achieve a consistent size distribution of vesicles. Lipid vesicle stocks were kept at 4 °C. Stock lipid concentrations were confirmed using a phosphate assay⁶⁷.

Dynamic light scattering

The size distribution of DPPC vesicles was estimated using a ZetaSizer Nano in 173° backscatter with multiple narrow mode (high resolution) analysis. Final concentration of lipids was 10 μM and the amount of RNA was fixed at a ratio of 10 lipids / nucleotide. Results were plotted as the size distribution in the number of detected species (NumberPSD).

UV-crosslinking

RNA was incubated for 10 minutes in the reaction buffer with or without lipids followed by 10 minutes incubation under UV-B light (300 mW LEUVA77N50KKU00 LED 305 nm) from a distance of 1.5 – 2 cm. Samples were analysed on denaturing PAGE with SybrGold poststaining.

R3C activity assays

5 pmol of R3C ligase was mixed with 0.5 pmol of the substrate to ensure saturation of the system (pseudo 1st order reaction) and decrease any batch-to-batch differences coming from R3C purification. Final reaction volume was adjusted to 30 μl. Incubation was conducted at various concentrations of DPPC vesicles and samples were ethanol precipitated after 30, 60, 90 and 120 minutes. Reactions were quantified on denaturing PAGE using FIJI software. Product and substrate intensities (I_P and I_S , respectively) were quantified and the concentration of product at time t was calculated:

$$[P]_t = [S]_0 * \left(\frac{I_P}{I_P + I_S} \right)$$

To determine reaction rates (M/min), concentration of the product was plotted as a function of time and then fitted with a linear function (fixed [0;0] point).

RNA-lipid binding assays

Binding assays for the results in Fig. 3c (R3C ligase), Fig. 4a, b, c and Fig. 5a were conducted using liposome-coated magnetic beads. Binding assays for the results in Fig. 2b, c and 5b, c were conducted using an ultracentrifugation method.

Magnetic beads binding assay

Magnetic beads (DynaBeads streptavidin T1, Invitrogen) were coupled with DPPC liposomes doped with 0.5 mol% biotinoyl cap PE. Lipid concentration on non-diluted beads was estimated using a phosphate assay⁶⁷. Liposome conjugated beads final working concentration was of 800 μM lipid.

25 pmol of 19 nt or 10 pmol of 40/36 nt RNAs were incubated with 2 – 1.5x diluted magnetic beads for 30'. After incubation the supernatant was separated from the beads using a magnet. The amount of RNA left in solution was estimated either using absorbance or Qubit miRNA quantification kit (ThermoScientific, USA). The reported binding efficiencies (Fig. 4a, b) correspond to 15 μL of liposome-coated beads in a 30 μL reaction volume. The partition coefficient K (Fig. 4c, Fig. 5a) was calculated according to following formula:

$$K = \frac{\frac{[R_L]}{[R_L] + [L]}}{\frac{[R_w]}{[R_w] + [W]}} \quad \text{Eq. 1}$$

Assuming high excess of [L] and [W] we simplified the equation to:

$$K = \frac{\frac{[R_L]}{[L]}}{\frac{[R_w]}{[W]}} \quad \text{Eq. 2}$$

Where:

[R_L] – RNA bound to the membranes

[R_w] – non-bound RNA

[L] – outer lipid concentration

[W] – water concentration

Ultracentrifugation binding assay

RNA was incubated with various concentrations of DPPC vesicles for 30'. After incubation ultracentrifugation was conducted to separate vesicles from the solution (125 000 x G, 40', 24 °C). 9 μL of supernatant (out of 30 μL reaction mix) was pulled and incubated with 1.5 μL of 0.5 M EDTA (90 °C, 5'). After incubation 150 μL of 1xQubit dye was added to the solution and the amount of RNA was measured using a plate reader (SPARK 20M, TECAN).

Measured fluorescence intensity values were plotted as a function of lipid/oligomer or lipid/nucleotide ratios (we assumed that RNA interacts only with the outer membrane leaflet, so final outer lipid concentration is equal half of the total lipid amount) and fitted using Hill's fit (OriginLab software):

Eq. 3

$$f(x) = Start + (End - Start) \frac{x^n}{x^n + k^n}$$

Where Start and End are function plateau values, k is inflection point of curve and n is cooperativity index. Inflection point of the curve (k) is a parameter which we used for further data analysis.

RNA-lipid binding partition coefficient determination for R3C substrates

5 pmol of RNA was incubated with an excess of DPPC vesicles (5 mM, 2.5 mM outer leaflet) for 30 minutes. After incubation ultracentrifugation was conducted to separate vesicles from solution (125 000 x G, 40', 24 °C). 10 µL out of 30 µL was taken as a supernatant sample, and the rest of the supernatant was discarded to leave only the pellet. RNA from supernatant and pellet samples was ethanol precipitated and analyzed using denaturing PAGE. Partition coefficient values (Fig. 3c "substrate", Fig. 6c) were calculated using Eq. 2.

Statistical analysis

All of the binding efficiency, partition coefficient, crosslinking efficiency and R3C activity assays were repeated at least 3 times. Presented error bars are SEM values. Fits to the lipid/RNA binding assay ratio determinations were included with $R^2 > 0.9$.

a) In gel-liquid phase separated membranes, RNA partitions into gel domains

b) RNA-lipid binding is abolished when gel phase domains are melted

Fig. 1 RNA selectively binds to gel phase membrane domains. (a) gel-liquid phase separated giant unilamellar vesicles (GUVs) were prepared from a mixture of DOPC and DPPC with a molar ratio of 1:1 and labeled with 0.5 mol % of the fluorescent lipophilic probe DiD. DiD is excluded from gel phase domains, which are observed as non-stained regions on the surface of the vesicle. A mixture of random RNA oligomers (40xN) stained with SybrGold is enriched within the gel phase domains. A control without RNA shows that SybrGold itself does not stain the GUVs. (b) When gel phase domains are melted by increasing temperature (3.4 °C/min – see supplementary movie 1) RNA no longer enriches at the GUV surface. Decreasing the temperature again restores gel-liquid phase separation and, consequently, RNA-membrane binding (~28 °C, white arrows – see supplementary movie 2). All of the scale bars are 5 μ m.

Fig. 2 RNA-lipid interactions induce reversible aggregation of vesicles and enhance intermolecular RNA interactions. (a) RNA-induced vesicular aggregation can be observed through changes in vesicle size distribution measured by dynamic light scattering (left), or visually as depicted in the cuvette images (right). The introduction of a randomized pool of RNA oligos to gel lipid membranes (gel phase = **G**) induces vesicle aggregation, which can be reversed either through chelation of divalent cations with EDTA or by increasing the temperature to achieve membranes in a liquid phase (liquid phase = **L**). **(b)** R3C ligase (84 nt) shows a small increase in UV-crosslinking at low lipid concentration, whereas crosslinking efficiency decreases at higher lipid concentration. Binding and UV crosslinking are inversely correlated and show similar inflection points with respect to lipid concentration. **(c)** Randomized RNA oligos (75-85 nt) show increasing crosslinking efficiency in the presence of lipids with a slight decrease at the highest lipid concentration; crosslinking and lipid binding are closely correlated. **(d)** Analysis of RNAs by native PAGE gel demonstrates that the R3C ligase forms primarily one structure, whereas randomized RNA forms diverse intermolecular complexes.

a) Determination of R3C ligase activity

b) Effect of lipid vesicles on R3C ligase activity

c) Lipid binding and nucleotide characteristics of R3C reactants

Oligo	log(K)	Nucleotide %				Length
		G	A	C	U	
R3C	7.22 ± 0.08	27	28	13	31	84 nt
Substrate	3.39 ± 0.15	8	33	33	25	12 nt

Fig. 3 Lipid membranes influence R3C ligase activity through lipid-ligase interactions. (a) R3C ligase reaction rates were quantified by measuring the amount of product within a 2 hour incubation time. **(b)** R3C reaction rate is dependent on the presence of gel membranes - increased activity is observed for lower lipid concentration followed by drop in activity at higher lipid concentration. **(c)** The 12 nt R3C substrate shows several orders of magnitude lower membrane-buffer partition coefficient $\log(K)$ compared with the R3C ligase, which could be derived from different compositional or structural characteristics of both RNA species.

Fig. 4 RNA-lipid binding is RNA sequence-dependent. (a) Short 19 nucleotide oligomers show sequence-dependent binding to the gel membranes, with highest binding observed for the pure G oligomer followed by oligos with randomized sequence or AG repeats. Oligomers containing only A, C or U bases show negligible binding to the lipid membranes. (b) Oligonucleotide length also influences binding efficiency as seen in the comparison of (a) 19xN and (b) 40xN. Depletion of one type of nucleotide within oligomers lowers binding efficiency. Lack of guanine decreased binding by 57%, whereas lack of adenine, uracil or cytidine showed a less pronounced effect. (c) Oligonucleotides with increasing G content showed higher lipid – buffer partition coefficients compared to a G-deficient oligo with the highest value observed for the 9xG oligomer.

a) Effect of RNA base-pairing on vesicle aggregation and lipid membrane binding

b) Lipid:RNA binding ratios for ssRNA and dsRNA

c) Effect of G-content on lipid:nucleotide binding ratio

Fig. 5 RNA-gel membrane binding depends on RNA structure. (a) The effect of RNA base-pairing structure on lipid vesicle aggregation (left) and membrane binding (right) was evaluated by comparing RNA's with limited ability to create inter- and intramolecular structures (ssRNA: 36 nucleotide ACUG or CAGU repeats) with 100% complementary double stranded RNAs (dsRNA: mixed ACUG and CAGU oligos). Single stranded RNAs do not cause lipid vesicle aggregation despite moderate membrane binding, whereas double stranded RNA species show both high binding and vesicle aggregation. **(b)** RNA:lipid binding ratios (shown in table) estimated from binding curves of ssRNA and dsRNA (plotted) are affected by base pairing and sequence. **(c)** Oligomers with different nucleotide composition show different lipid/nucleotide binding stoichiometry. Analysis of oligomers by native PAGE gel (**c, inset**) indicates that the presence of guanine promotes inter-molecular interactions, suggesting, that base-pairing enhances the average strength of nucleotide-lipid interactions.

Fig. 6 Modification of the R3C substrate sequences enables lipid-dependent tunability of ligation rate. **(a)** Modified R3C reaction substrates were synthesized in order to increase binding to the membrane. The catalytic 3' end of the molecule was enriched with 5' tails with varying G content. Reactions for modified substrates showed lower reaction rate constants compared with the unmodified substrate. **(b)** The mechanism for inhibition of activity was evaluated by analysis of R3C reactants using native PAGE. The unmodified short substrate is entirely bound to the ligase, whereas AG-modified substrates show mobility abnormalities, which suggest substrate folding (magenta points). **(c)** Modification of the substrate sequence through addition of a 5' overhang increases binding to the membrane. **(d)** Addition of lipid membranes to reactions with the G-rich substrates increases reaction rates to the level of the unmodified substrate. **(e)** We propose that the decrease of R3C reaction rates in the presence of G-rich substrates is based on the inhibition of the catalytic part of the substrate by the 5' G-rich overhang. The presence of lipid membranes not only increases R3C-substrate stability, but also screens the 5'-rich substrate part enabling the R3C reaction to proceed.