1 Progesterone receptor-A isoform interaction with RUNX transcription factors controls
2 chromatin remodelling at promoters during ovulation

3 Authors: Dinh DT ¹(*), Breen J ², Nicol B ³, Smith KM ¹, Nicholls M ¹, Emery A ¹,
4 Wong YY ¹, Barry SC ¹, Yao HHC ³, Robker RL ¹, Russell DL ¹(*)

5 Affiliations:

6 1. Robinson Research Institute, Adelaide Medical School, University of Adelaide, Australia

7 2. South Australian Genomics Centre (SAGC), South Australian Health and Medical

8 Research Institute (SAHMRI), Adelaide, South Australia, Australia

9 3. Reproductive Developmental Biology Group, National Institute of Environmental Health

10 Sciences, Research Triangle Park, NC 27709, USA

11

12 Summary:

13 Progesterone receptor (PGR) plays diverse roles in reproductive tissues and thus coordinates
14 mammalian fertility. In the ovary, acutely induced PGR is the key determinant of ovulation
15 through transcriptional control of a unique set of genes that culminates in follicle rupture.

16 However, the molecular mechanisms for PGR's specialised function in ovulation is poorly
17 understood. To address this, we assembled a detailed genomic profile of PGR action through
18 combined ATAC-seq, RNA-seq and ChIP-seq analysis in wildtype and isoform-specific PGR
19 null mice. We demonstrated the unique action of PGR-A isoform in the ovary through a
20 transcriptional complex involving physical interaction with RUNX and JUN/FOS
21 transcription factors. The assembly of this unique complex directs targeting of PGR binding
22 to proximal promoter regions and enables chromatin accessibility, leading to ovulatory gene

induction. This PGR signalling mechanism is specific to ovulation and provides potential targets for infertility treatments as well as new contraceptives that block ovulation.

25

1. INTRODUCTION

Ovulation is a highly controlled process that ensures the release of appropriately mature oocytes, timed to coincide with the receptivity of the uterus for embryo implantation. The mechanism of ovulation begins with the activation of cAMP-dependent kinase (PKA) and extracellular regulated kinase (ERK) signalling cascades in granulosa cells of the follicle in response to the pituitary LH surge¹. These signals result in the induction of an array of transcriptional regulators that collectively mediate complete structural remodelling of the ovarian follicle and the resumption of oocyte meiotic maturation². Meanwhile, follicular granulosa cells undergo pervasive differentiation and rapid neoangiogenesis to form the corpus luteum, a transient endocrine organ that produces high levels of progesterone to promote uterine receptivity³ and mammary gland maturation⁴ in preparation for pregnancy and lactation.

Progesterone signalling also has an obligatory role in granulosa cells where it is essential for ovulation. In the ovary, PGR is a key transcription factor that determines ovulation^{5,6}, as demonstrated in mutant animal models as well as in primates and human. Female mice with either total or granulosa-specific null mutation of the nuclear progesterone receptor (PGR) have a complete anovulatory phenotype without affecting oocyte meiosis or luteinisation⁵⁻⁷. Likewise, in primates and humans, administration of PGR antagonists (Mifepristone or Ulipristal Acetate) blocks ovulation^{8,9} and thus have been explored as potential acute contraceptive agents. However, disrupting systemic progesterone action has unacceptable side effects across many target organs, necessitating further research into the unique ovarian

47 PGR action and its downstream target genes in order to discover highly selective targets for
48 ovulation-blocking contraceptives.

49 As a steroid receptor, PGR is a nuclear transcription factor that is activated by its ligand,
50 progesterone. In the uterus and mammary gland, PGR expression is maintained by estrogen
51 action. However, in the ovary PGR is not estrogen dependent¹⁰ but is highly induced in
52 granulosa cells in response to the LH surge¹¹. In general, PGR is believed to regulate
53 transcription through binding the PGR response element (PRE) motif that can be recognised
54 by all NR3C steroid receptors. However, PGR function can be highly intricate and involves
55 binding to target genes in cooperation with other transcription factors^{12,13}. Such
56 transcriptional complexes can recruit histone and chromatin remodellers including steroid
57 receptor coactivators (SRC), CBP/p300 and PRMT1¹⁴⁻¹⁶ to facilitate accessibility for the
58 basal transcriptional machinery and RNA Pol II to induce expression of target genes¹⁷. The
59 two major isoforms of PGR – PGR-A and PGR-B – possess specific properties and functions,
60 including distinct protein-protein interaction capabilities¹⁸ and cross-isoform regulation, as
61 demonstrated by the widely reported repression of PGR-B transcriptional activity by PGR-A
62 in the uterus¹⁹ and breast cancer cells²⁰. In the ovary, both PGR isoforms are induced by the
63 LH surge^{21,22}; however, PGR-A is credited as the more essential isoform in ovulation, as
64 shown through PGR-A specific KO mouse model (AKO)²³. In contrast, female mice lacking
65 only PGR-B (BKO) exhibit normal ovulation²⁴. The relative importance of PGR-A and
66 PGR-B throughout reproductive physiology remains an active topic of investigation^{3,25,26},
67 and their molecular mechanisms in granulosa cells that control ovulation remain unknown.

68 Our previous characterisation of the granulosa cell PGR cistrome through ChIP-seq suggests
69 that the classic pathway involving the canonical PRE motif is not favoured by PGR in
70 granulosa cells and that the cooperation between PGR and context-specific transcription
71 factors, such as JUN/FOS and RUNX transcription factor families, establishes unique

transcriptional regulation during ovulation²². The interaction between transcription modulators and PGR has been shown to be important in other biological contexts^{12,13}; however, whether such interactions are crucial for PGR-mediated ovarian functions remains unexplored. Among the potential binding partners for PGR, the RUNX family was uniquely identified in granulosa cells²². Included in this family is RUNX1, which acts as a regulator of gene expression in peri-ovulatory granulosa cells^{27,28}. The ablation of CBF β , the canonical protein partner of RUNX1 and the closely related RUNX2, results in the reduction of important ovulatory genes and consequentially impaired fertility in female mice^{29,30}. However, the nature of PGR/RUNX1 relationships in the context of ovulation remains unknown.

This study provides novel insights into the molecular mechanisms employed by PGR in granulosa cells to cause ovulation. The scope of global changes in the open chromatin landscape and transcriptome induced by the LH surge and PGR in mouse granulosa cells was defined using ATAC-seq and RNA-seq in wildtype and PGR-null mice. The formation of molecular complexes between PGR and other transcription factors was confirmed. For RUNX1, comparative ChIP-seq analysis pointed to a high level of PGR/RUNX1 cooperativity at mutual chromatin sites and especially at proximal promoters of ovulatory genes. Isoform-specific granulosa cell transcriptomes also identified the dominant role of PGR-A in ovulation, through interaction with RUNX transcription factors. The results demonstrate a unique molecular mechanism that involves PGR-A and RUNX1 co-operativity to shape granulosa cell chromatin epigenetic reprogramming and transcriptional response to the LH surge. This novel progesterone signalling pathway is critical for normal fertility and will enable the identification of causes of unexplained infertility as well as the discovery of molecular complexes or protein targets for future contraceptive discovery.

97 **2. RESULTS**

98 **Rapid global reprogramming of chromatin state and transcription by LH-stimulus** 99 **through the induction and repression of distinct transcription factor binding**

100 Ovulation induction by hCG injection to simulate the LH surge in mice provides an *in vivo*
101 model for the rapid response to ovulatory cues in granulosa cells. ATAC-seq in granulosa
102 cells isolated before or 6 h after *in vivo* LH-stimulus revealed dynamic changes in chromatin
103 accessibility in response to the induction of ovulatory transcription factors, including PGR.
104 Stringent ATAC-seq peak calling criteria identified a total of 71,287 peaks across both
105 conditions, of which 28,729 sites (40.3%) were LH-induced, 21,257 sites (29.8%) were
106 significantly repressed and 21,301 sites (29.8%) with no change in accessibility after LH-
107 stimulus (Figure 1A). LH-mediated differentially accessible sites, including those more
108 accessible before or after LH-stimulus, indicated that chromatin remodelling occurred not
109 only in promoters but also at distal enhancers and throughout gene bodies (Figure 1B). LH-
110 stimulus induced the accessibility of promoters corresponding to genes involved in
111 cytoskeletal reorganisation, mitochondrial functions, and signalling pathway activation which
112 are important processes leading up to ovulation (Supplementary figure 1). Significantly, LH-
113 stimulus also repressed access to genes that were responsible for gene silencing and
114 nucleosome modification. Genes that were involved in normal cellular processes, such as
115 RNA processing, biosynthesis and vesicle transportation, showed no change in promoter
116 accessibility by the treatment.

117 As ATAC-seq provides DNA accessibility data at footprint resolution, it can be used to
118 analyse transcription factor binding sites in an unbiased manner. To predict the activity of
119 key transcription factors involved in changes in the accessibility of LH-mediated open
120 chromatin sites, differential enrichment analysis of transcription factor-binding motifs was

performed (Figure 1C, Supplementary table 1). The motif for many ovulatory transcription factors, such as for JUN/FOS, RUNX, CREB and CEBP β , showed significantly increased binding score in peaks for open chromatin after LH-stimulus, whereas those for CTCF, NR5A and ESRR were enriched at pre-LH open chromatin sites. Interestingly, the canonical motifs for the NR3C family members, such as glucocorticoid receptor (NR3C1/GR), mineralocorticoid receptor (NR3C2/MR), androgen receptor (NR3C4/AR) as well as PGR, were significantly more enriched in open chromatin before LH-stimulus. In support of these findings, *de novo* motif enrichment analysis using HOMER also identified JUN/FOS, RUNT and CEBP motifs in LH-induced, but not LH-repressed open chromatin regions (Figure 1D). Motifs for NR5A and GATA families were similarly enriched in all ATAC-seq peak subsets. Importantly, a motif best matched to that bound by NR3C members was identified only in the LH-repressed subset, consistent with the differential transcription factor binding analysis pattern above. Direct binding of transcription factors to these motifs was confirmed through DNA motif footprinting (Figure 1E). Footprint patterns indicative of transcription factors bound to these sites could be observed with medium to deep indentation (JUN/FOS, CTCF, NR5A) and shallower patterns (RUNT, NR3C), alluding to differences in the dwell time of these transcription factors. Again JUN/FOS footprints were only evident after LH-stimulus, while evidence of NR3C footprints was more obvious in unstimulated cells.

In order to correlate chromatin accessibility profiles with gene expression changes, the complete transcriptional response of peri-ovulatory granulosa cells to LH-stimulus was profiled through RNA-seq of granulosa cells collected before or 8 h after *in vivo* LH-stimulus. This timepoint was chosen as it is approximately 2 h after the peak induction of key ovulatory transcription factors. A total of 2,088 genes were differentially expressed (DEG), of which 52% were upregulated by LH-stimulus (Figure 2A). This included many genes previously associated with ovulation, such as *Ptgs2*, *Pgr*, *Runx1*, *Runx2* and *Adamts1*

(Supplementary Table 2). Prediction of the potential upstream regulators of LH DEGs using tools from the Ingenuity Pathway Analysis (IPA) software identified a suite of transcription factors, many of which were also among the DEGs in response to the LH surge, including PGR (logFC = 6.4), RUNX1 (logFC = 4.6), CEBP β (logFC = 2.5) and JUNB (logFC = 2.1) (Supplementary Table 2, 3). Notably, the binding motifs for these same transcription factors also showed increased accessibility following LH-stimulus (Figure 1C, D). The global relationship between promoter accessibility and downstream gene expression in peri-ovulatory granulosa cells was illustrated through a comparison between LH DEG and LH-dependent chromatin accessibility at the promoter (within 3 kb upstream of TSS) of these genes. Significant changes in promoter accessibility corresponded with transcriptional change in 69.3% of LH-driven DEGs (1,447 out of 2,088 genes) (Figure 2B, red symbols), in which promoter accessibility largely increased in LH-induced genes and was less accessible in LH-repressed genes (Spearman correlation coefficient = 0.467). For example, in *Cxcr4* and *Inhbb*, two genes that were respectively upregulated and downregulated post-LH, clear association between expression pattern and ATAC-seq signal at the gene promoter could be observed (Figure 2C). In some instances, however, genes that were differentially regulated had constitutively accessible promoters, as seen in *Cited1*.

PGR and RUNX1 mutually bind chromatin in peri-ovulatory granulosa cells

Since RUNX1 is functionally important in granulosa cells³¹ and RUNT motifs were enriched in granulosa cell PGR ChIP-seq peaks²² as well as regions which gained chromatin accessibility after LH-stimulus (Figure 1C), we asked whether RUNX1 is key to the transcriptional regulatory mechanism that mediates ovulatory gene expression. Having confirmed RUNX1 expression to be induced in granulosa cells in response to the LH surge (Supplementary figure 2A, B), we sought to biochemically determine its targets on a genome-wide scale. RUNX1 ChIP-seq was performed on granulosa cells before or 6 h after *in vivo*

hCG treatment to induce ovulation. Global RUNX1-chromatin binding was detected prior to the LH surge but was also induced 7-fold after the LH surge (16,065 LH-induced peaks) (Figure 3A). Such LH-activated RUNX1 binding was strongly associated with LH-induced open chromatin as defined through ATAC-seq (Supplementary figure 3A). A clear relationship between RUNX1 and PGR chromatin interaction in peri-ovulatory granulosa cells was demonstrated through comparison between RUNX1 ChIP-seq and PGR ChIP-seq at 6h post-LH stimulation. PGR and RUNX1 shared a remarkable number of mutual binding sites within 100 bp, with 9,704 chromatin sites (or 70% of total PGR and 52% of total RUNX1 binding sites) having both PGR and RUNX1 binding (Figure 3B). Among these shared PGR/RUNX1 binding sites, 9,288 were located in ATAC-seq open chromatin sites identified at the same time point. Dividing transcriptionally active RUNX1 and PGR cistromes into those that were shared or uniquely bound by each transcription factor showed a relatively low level of promoter binding in uniquely PGR-bound sites, yet a strong preference for proximal promoter binding (within 1 kb of TSS) was a predominant characteristic of PGR/RUNX1 co-binding sites (Figure 3C). The remarkably high overlap of PGR and RUNX1 binding sites, coupled with our previous report of enriched RUNX1-binding motifs in granulosa cell PGR ChIP-seq²² suggests that the previously reported granulosa-specific ability of PGR to target proximal promoter relies on co-binding with RUNX1. Such dependency was not reciprocal, as RUNX1 showed a prevalence for proximal promoter occupancy that was independent of PGR co-binding. To test whether LH-induced RUNX1 chromatin binding was dependent on PGR, we assessed whether LH-induced RUNX1 binding intensity was greater at PGR-overlapped sites versus PGR-independent sites. No difference was observed in the level of RUNX1 enrichment, suggesting that PGR was unlikely to play a role in facilitating the tethering and recruitment of RUNX1 to these shared sites (Supplementary figure 3B).

The importance of PGR and RUNX1 cooperation on ovulatory gene regulation at a genome-wide level was demonstrated by comparing genes with PGR and/or RUNX1 binding at their promoters against all DEG responding to LH in granulosa cell (Figure 3D). Remarkably, 62.4% of all LH-mediated DEGs contained at least PGR or RUNX1 binding. Of these, the majority of RUNX1- (55%) and PGR-bound (87%) LH DEGs were in fact co-bound by both transcription factors, implying that simultaneous interaction of both PGR and RUNX1 at target promoters was a major mechanism for each of these transcription factors in gene regulation. Examples of PGR/RUNX1 mutual chromatin occupancy were evident in the promoters of many known PGR and RUNX1 target genes in peri-ovulatory granulosa cells (*Adamts1*, *Rgcc* and *Zbtb16*) (Figure 3E).

DNA motifs preferentially bound by RUNX1 in granulosa cells were identified using motif enrichment analysis. As expected, the canonical RUNT motif was significantly highly enriched in the RUNX1 cistrome (3.5-fold over background, p-value 1e-787) as well as PGR cistrome (2.8-fold, p-value 1e-370) (Figure 3F). The NR3C motif was also enriched in RUNX1-bound peaks (2-fold over background, p-value 1e-25); interestingly, such enrichment was restricted to only PGR/RUNX1 co-binding sites (Supplementary figure 3C). Other enriched non-canonical motifs included those corresponding to JUN/FOS, CEBP and GATA transcription factors.

The potential physical interaction between PGR and RUNX1 was supported by positive proximity ligation assay (PLA) signals between PGR/RUNX1 in the nucleus of mouse granulosa cells treated with hCG and R5020 for up to 8 hours to mimic the *in vivo* ovulatory stimuli (Figure 3G). PLA signal showed a rapid transient induction of PGR/RUNX1 complexes by LH-stimulus, indicating an acute temporal PGR/RUNX1 interaction that was largely absent before stimulus but increased within 4 h after stimulation.

In addition to RUNX1, other members of the RUNT transcription factor family are capable of binding the RUNT motif. Notably, RUNX2 was induced by the LH surge in granulosa cells of antral follicles and had a transient temporal expression pattern during the peri-ovulatory window similar to PGR and RUNX1 (Supplementary figure 2A, B), consistent with previous reports³². The interaction between PGR and RUNX2 was also demonstrated through PLA of granulosa cells exposed to ovulatory stimuli (Figure 4A). Similar to the interaction between PGR and RUNX1, PGR/RUNX2 PLA signal was induced within 4 h of stimulus. Additional potential candidate protein partners from the NR5A and JUN/FOS transcription factor families were selected based on their known involvement in granulosa cell functions during ovulation. Interactions of PGR with LRH1 and JUND were observed in granulosa cells after 6 h hCG, but not with cJUN or JUNB (Figure 4B). The interaction between PGR and ovulatory transcription factors was predominantly in the nuclear compartment of the cell, indicating that these transcription factors concurrently interact in chromatin-bound complexes in response to ovulatory cues.

PGR mediates chromatin accessibility in granulosa cells

PGR-bound regions predominantly overlap with transcriptionally active chromatin, as shown by 75% overlap in PGR- and H3K27ac ChIP-seq peaks²²; likewise 90% of PGR binding sites were located in 6 h hCG ATAC-seq peaks (Supplementary figure 4A). In addition, an interaction between PGR and the histone acetyltransferase CBP/p300 was illustrated in hormone-stimulated granulosa cells (Supplementary figure 4B). A functional relationship between PGR binding and chromatin accessibility was further investigated through ATAC-seq in granulosa cells isolated from PGRWT and PGRKO mice. Of a total of 73,764 identified ATAC peaks, 1,499 sites were found to have significantly altered peak intensity in the absence of PGR (Figure 5A); interestingly, only 6 of the 1,499 sites had increased ATAC peak intensity in PGRKO compared to PGRWT. Whereas PGR-chromatin binding has been

shown to favour proximal promoter regions in granulosa cells, the PGR-dependent ATAC-seq peaks were more evenly distributed throughout all genomic features (Figure 5B). Through RNA-seq in granulosa cells from PGRWT and PGRKO mice 8 h post-hCG treatment, 2 h after the peak of PGR abundance, 286 PGR-dependent DEGs were identified (Supplementary table 2). Comparison of these PGR-regulated genes and PGRKO ATAC signal showed a consistent reduction in ATAC-seq intensity at promoters of genes with PGR-dependent expression (Spearman correlation coefficient = 0.5116, p-value < 0.0001) (Figure 5C). 28 DEGs showed a significant difference in ATAC-seq peak intensity as determined through DiffBind (Figure 5C, light green), indicating that promoter accessibility for these genes is significantly influenced by PGR. Examples of the difference in PGR influence on promoters are shown for *Gas7*, whose promoter was differentially accessible in PGRKO, and *Abhd2*, the promoter accessibility of which was not significantly altered in the absence of PGR (Figure 5D).

Transcription factor motif enrichment analysis was performed through HOMER on PGR-driven ATAC peaks and compared with motifs enriched in PGR ChIP-seq. As with ChIP-seq, the most highly enriched and most significant motif identified in PGR-driven ATAC peak sequences was the canonical NR3C motif (12.4-fold over background). Again, several non-canonical motifs were also significantly enriched in ATAC-seq as with PGR ChIP-seq (Figure 5E). To examine the possibility that differential chromatin remodelling in PGRKO was mainly at sites with direct PGR-NR3C motif binding, we compared PGR and RUNX1 ChIP-seq cistromes with PGR-driven ATAC sites. Interestingly, 53% (795 of 1,499) of PGR-dependent ATAC sites were bound by PGR but only 17.8% (267 sites) were bound by RUNX1, almost all of which (250/267 sites) were PGR/RUNX1 co-bound (Figure 5F). This indeed suggests that PGR-dependent accessible chromatin sites were mainly bound by only PGR and not RUNX1.

PGR-A is the key mediator of RUNX1 interaction and ovulatory gene regulation

We confirmed that PGR-A was the most important isoform in ovulation by verifying the disrupted ovulatory phenotype in both total PGRKO and AKO mice but not BKO mice, as previously documented^{23,24} (Supplementary figure 5A). RNA-seq in granulosa cells obtained from PGRKO, AKO or BKO and WT littermates from each strain was analysed and hierarchical clustering analysis showed that the global gene expression pattern in AKO granulosa cells resembled that of total PGRKO, while BKO more closely resembled the WTs of each strain (Figure 6A). The loss of PGR-A resulted in the differential expression of 310 genes, 153 (or 49.4%) of which were also identified in the absence of both PGR isoforms (Figure 6B, Supplementary table 2). Among these, 48.6% and 58.5% of AKO and PGRKO DEG respectively were also differentially expressed in response to the LH surge. The loss of PGR-B, on the other hand, resulted in no significant transcriptional changes (Figure 6B), confirming that PGR-A plays a predominant role in the overall effect of PGR during ovulation. Examples of RNA-seq signals can be observed in PGR target genes such as *Pparg* and *Stard5* (Supplementary figure 5B, C), which exhibits dependence on PGR-A but not PGR-B.

Upstream analysis of the DEG datasets using IPA analysis predicted a number of transcription factors as potential upstream regulators of the PGRKO and AKO responses (Figure 6C). As expected, PGR was among the most significant predicted upstream regulators, as well as other transcription factors we and others have shown to be targets of PGR action, such as PPARG³³ and HIF1A³⁴. Importantly, this independent approach also implicates RUNX1, CBF β , JUN, FOS and CBP/p300 as mediators of the PGR-dependent transcriptional response. The lack of PGR-B effect on transcription regulation is suggestive of PGR isoform-specific interactomes in granulosa cells. To confirm whether the direct PGR interaction with RUNX was specific for PGR-A, we performed PLA for PGR/RUNX1 and

PGR/RUNX2 in hCG/progestin-stimulated granulosa cells obtained from WT, het and KO animals from the AKO and BKO strains. A decrease in protein-protein interaction with both RUNX1 and RUNX2 was observed exclusively in AKO but not BKO granulosa cells (Figure 6D).

3. DISCUSSION

Safer, more accessible contraception as well as new treatments for anovulatory infertility remain key unmet needs in fertility management. In the era of high specificity, low side-effect drug development, this can be delivered once the molecular mechanisms driving the biological process are fully elaborated and validated. In this study we applied unbiased genome-wide approaches to characterise the global chromatin remodelling in granulosa cells responding to the *in vivo* ovulatory stimulus and identified the cell-specific gene regulatory network controlled by the action of PGR, the critical transcription factor required for ovulation (Figure 7). Our data confirm a specific role for PGR-A in granulosa cells through a unique interaction with RUNX transcription factors at non-canonical DNA motifs. We verified the cooperative binding of PGR with RUNX1/2 by demonstrating substantial overlap between PGR and RUNX1 cistromes as well as showing LH-dependent formation of nuclear complexes containing PGR and RUNX1/2 that depend on the presence of PGR-A but not PGR-B. Together, these findings support our hypothesis that the unique ovarian action of PGR involves the formation of PGR-A/RUNX transcriptional complexes with specific DNA sequence binding activity.

The LH surge leads to critical shifts in the chromatin landscape and the ovarian gene expression profile, eventually resulting in the activation of biological pathways that are important for follicle rupture. Markedly increased chromatin accessibility around promoters

of ovulatory genes and also at many intergenic regions indicates global epigenetic changes in histone modifications throughout the genome within 6 h of the ovulatory stimulus, leading to the rapid and pervasive reprogramming of gene expression. Our approach also allowed comprehensive identification of the transcription factors involved. *De novo* motif identification and footprinting analysis confirm RUNX and JUN/FOS as well as NR5A and CEBP transcription factor binding at open chromatin sites. This is consistent with the reported induced expression of JUN and FOS³⁵, RUNX1/2^{32,36} and CEBP β ³⁷ by ovulatory stimulus, as well as an activation of the PKA-CREB1 signalling pathway^{38,39}.

Surprisingly, the canonical sequence motif used by NR3C steroid receptors, including PGR, shows the largest repression in chromatin accessibility in response to the ovulatory stimulus. Considering that PGR expression is highly induced in the peri-ovulatory period²² and that it is a key determinant of ovulation^{6,40}, the loss of NR3C response element representation in open chromatin regions suggests that PGR-chromatin interaction is less dependent on this consensus sequence and supports our previous report that PGR binding sites in granulosa cells are exceptionally enriched in RUNX and JUN/FOS sequence motifs²². Tissue-specific PGR interactomes have been shown to be important for the diversity of PGR transactivation functions in various contexts⁴¹. Here, an interaction between PGR and RUNX1 is described for the first time, in which PGR and RUNX1/2 form a very close physical association specifically within the nucleus of granulosa cells in response to ovulatory cues, strongly suggesting that they reside in a cooperative transcriptional complex. Further evidence that PGR/RUNX1 interact in a transcriptional complex comes from the remarkable overlap in PGR/RUNX1 co-occupancy on granulosa cell chromatin (70% of PGR binding sites overlapping with RUNX1), especially at ovulatory genes. The very high prevalence of promoter-centric RUNX1 binding and RUNX1/PGR co-binding mimics the promoter-centric enrichment of the PGR cistrome we previously reported to be highly specific in granulosa

cells and not other PGR target tissues^{12,22}. We now show that this selective PGR-promoter interaction is dependent on RUNX1 and is also not reciprocal, since RUNX1 preferably targets proximal promoters both in the presence or absence of PGR. The predominance of PGR and RUNX1 binding identified at LH-regulated genes is equally striking. Together the results imply that RUNX1 directs PGR binding to promoter regions and they cooperatively regulate a significant proportion of the ovulatory gene expression profile. Consistent with this conclusion, RUNX1 has been previously reported to behave as a pioneer factor for the SWI/SNF chromatin remodelling complex⁴² and has a role in programming chromatin conformation⁴³. Reports of the interaction between members of the RUNX family and NR3C members, AR and GR, leading either to the repression⁴⁴ or transactivation of these transcription factors⁴⁴⁻⁴⁷, as well as RUNX1 tethering effect on ER α ⁴⁸, further substantiate the conclusion that RUNX transcription factors tether PGR to granulosa-specific chromatin targets. Utilising isoform-specific knockout mouse models, we also show that PGR-A is the predominant isoform obligatory for ovulation. The unique granulosa cell function of PGR-A is likely due to isoform-specific interaction with other transcription mediators, exemplified by the specific interaction between PGR-A and RUNX1. PGR is also known to have an active role in mediating chromatin accessibility through the recruitment of chromatin remodellers^{14,15}. The interaction between PGR and the histone acetyltransferase CBP/p300, identified through PLA, and altered accessibility at 1,499 chromatin sites in PGRKO granulosa cells shows that PGR is involved in chromatin reprogramming in ovulation. Together this indicates that tissue-specific PGR response is driven by proximal promoter-centric PGR/RUNX interaction as well as RUNX1-independent PGR binding outside of proximal regulatory regions that coordinately facilitates chromatin remodelling.

The physical interaction of PGR with other transcription factors, such as NR5A2 (LRH1) and JUND, is also identified. Although a low level of PLA signal suggests that PGR and LRH1

are closely co-localised, a functional interaction is less likely as they possess different windows of expression in granulosa cells (PGR is completely absent in granulosa cells prior to the LH surge whereas LRH1 is expressed from primordial stage up to corpus luteum⁴⁹. In support of this, less than 20% of PGR-regulated genes were also found to be LRH1 targets in peri-ovulatory granulosa cells². On the other hand, JUN/FOS transcription factors are highly induced by the LH surge in granulosa cells, similar to PGR^{35,50}. PGR/JUND PLA signal was abundant and substantial direct occupancy of the JUN/FOS motif was exclusive to LH-induced open chromatin sites. We suggest that JUN/FOS activation and chromatin binding is a parallel pathway to PGR/RUNX1/2 complexes and these transcription factors cooperatively mediate transcriptional activation of ovulatory genes. Further studies of JUN/FOS cistromes and transcriptomes in peri-ovulatory granulosa cells are necessary to define their contribution to PGR action and ovulatory function.

In summary, our results have clearly defined a unique mechanistic model of PGR action in granulosa cells which involves PGR-A interaction with RUNX1 to drive the specialised ovulatory PGR-mediated gene regulatory network. This ovarian-specific molecular mechanism of PGR action is distinct from the mechanisms in other PGR-responsive organs. These new insights are critical for understanding and addressing causes of anovulatory infertility and for the development of ovulation-specific contraceptives with minimal side-effects that often arise from progestin therapies.

4. METHODS

Reagents and antibodies

Unless otherwise stated, reagents were purchased from Sigma-Aldrich (St. Louis, MO, USA).

Animals

21-day old CBA x C57BL/6 F1 (CBAF1) mice were obtained from the Laboratory Animal Services (University of Adelaide). Null mutant mouse strains with ablation of either all PGR isoforms, or specific ablation of PGR-A or PGR-B^{7,23,24} were obtained from the Jackson Laboratory (Bar Harbor, USA). PGRKO mice are a targeted mutation strain with the Jackson Laboratory designation Pgr^{tm1Bwo}. PGR-A KO (AKO) mice are a targeted mutation strain with the Jackson Laboratory designation Pgr^{tm1Omc}. PGR-B KO (BKO) mice are a targeted mutation strain with the Jackson Laboratory designation Pgr^{tm2Omc}. Mice from KO strains are routinely genotyped from ear or tail biopsies before allocation to experiments and confirmed from replicate biopsies collected at the time of experiments. All mice were maintained in 12 h light / 12 h dark conditions and given water and rodent chow *ad libitum*. Littermate females of each genotype were allocated for experiments at 21 days of age. All experiments were approved by The University of Adelaide Animal Ethics Committee and were conducted in accordance with the Australian Code of Practice for the Care and Use of Animals for Scientific Purposes (ethics number m/2015/075, m/2018/100, m/2018/122, m/2018/117).

Granulosa cell culture and hormone treatment

CBAF1 female mice were stimulated with 5 IU eCG (Lee BioSolutions, Maryland Heights, USA) and culled at 44 hours post-eCG, upon which ovaries were dissected and granulosa cells isolated. Cumulus-oocyte complexes were removed and granulosa cells were counted before being seeded into an 8-well fibronectin-coated chamber slide (minimum 100,000 cells/well). Cells were cultured in DMEM:F12 media (Thermo Fisher) at 37°C, 5% CO₂ for 90 minutes then washed with PBS to remove debris. Cells were incubated overnight and were treated with 2 IU/ml hCG (Merck Sharp & Dohme B.V., Haarlem, Netherlands) and 100 nM R5020 (PerkinElmer, Waltham, USA) for indicated durations at 37°C, 5% CO₂.

416 **ATAC-seq**

417 ATAC-seq was performed as previously described⁵¹. Briefly, granulosa cells were collected
418 from superovulated female CBAF1 or PGRKO mice at 0 h or 6 h post-hCG as above. Cells
419 from 2-3 animals were pooled together for one biological replicate, two replicates were
420 obtained per condition. Granulosa cells were lysed and tagmentation was performed on the
421 nuclei using the Illumina Tagment DNA TDE1 Enzyme and Buffer Kits (Illumina, San
422 Diego, USA). DNA purification was performed using the Monarch PCR & DNA Cleanup Kit
423 (NEB, Ipswich, USA). Libraries were then amplified and purified before being size selected
424 using magnetic AMPure XP Beads (Beckman Coulter, Brea, USA). Library fragmentation
425 was confirmed using the LabChIP GX Bioanalyser (PerkinElmer) and sequencing was
426 performed on the HiSeq2500 Sequencing System (Illumina). For all datasets, adapters were
427 trimmed from 150-base sequences using Cutadapt⁵² and aligned to the mm10 mouse genome
428 using Bowtie2 algorithm⁵³. Alignment was filtered using samtools for non-mitochondrial
429 reads with MAPQ ≥ 30 cut-off and was de-duplicated using Picard tools. Peak calling from
430 read count followed the algorithm for Model-based Analysis for ChIP-Seq (MACS2) with a
431 q-value cut-off = 10^{-10} and a mouse genome size of 1.87×10^9 ⁵⁴. Differential binding analysis
432 for LH ATAC-seq and PGR ATAC-seq was performed using DiffBind⁵⁵ with DESeq2 linear
433 model, with differentially enriched sites determined to have FDR ≤ 0.05 . Genomic
434 distribution of peak subsets were determined using the ChIPseeker package⁵⁶. Gene ontology
435 enrichment analysis was performed using GREAT⁵⁷. Differential enrichment of transcription
436 factor binding motif and DNA footprinting analysis was performed using TOBIAS⁵⁸. Motif
437 analysis for *de novo* sequence motifs was performed using HOMER motif finding algorithm
438 with random 200 bp-long sequences from the mouse genome used to estimate motif
439 frequency in random sequence⁵⁹. Visual representation of ATAC signal was through the
440 UCSC Genome Browser (Santa Cruz, USA).

RNA-seq

Three-week old female mice, either CBAF1 (for LH RNA-seq) or WT and KO from PGRKO, AKO and BKO strains (for PGR RNA-seq) were hormonally stimulated as above, before being culled at 8 h post-hCG injection for granulosa cell collection. For LH RNA-seq, eCG-treated mice that did not receive hCG injection served as negative control. For each condition, four individual replicates were collected, each containing pooled cells from three animals. RNA was extracted from granulosa cells using RNeasy Mini kit (Qiagen, Chadstone, VIC, Australia). RNA quality was assessed using the RNA ScreenTape System (Agilent, Santa Clara, CA, USA).

For LH response RNA-seq, RNA was subjected to rRNA depletion using a modified Ribozero method and library preparation using Clontech SMARTer Stranded RNA-Seq Kits (Takara Bio, Shiga, Japan). Sequencing was performed on the Illumina HiSeq2500 100bp PE Rapid run (Ramaciotti Centre for Genomics, Kensington, Australia). For PGR RNA-seq, the library was prepared using the Universal RNA-Seq library kit with NuQuant Mouse AnyDeplete (Nugen, Redwood City, USA) and sequencing was performed on the NovaSeq 6000 S1 Sequencing System (South Australian Health & Medical Research Institute, Adelaide, Australia). Overrepresented adapters were checked using FastQC (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) and adapters were trimmed using AdapterRemoval⁶⁰ when required. Quasi-alignment of each dataset to the GENCODE mouse transcriptome (GRCm38.p6, M25 release) and transcript quantification were through Salmon⁶¹. Expression profiles and differential gene expression were assessed with limma⁶² and edgeR⁶³ using mean-variance estimates from log counts as described in the limma voom method. Differential gene expression for each experiment was defined as a fold-change ≥ 2 ($|\log FC| \geq 1$) and Benjamini-Hochberg adjusted p-value ≤ 0.01 . Upstream regulator analysis

of DEGs was through IPA software (QIAGEN). For visualisation of RNA-seq on the UCSC Genome Browser, read coverage was processed using deepTools ⁶⁴.

ChIP-seq

Granulosa cells from CBAF1 female mice were collected by puncturing ovarian follicles 44 h after eCG (0 h hCG) and 6 h after hCG stimulation. Two biological replicates were obtained for each sample from 10 mice, each with at least 1×10^7 cells. ChIP-seq for RUNX1 was performed by Active Motif (Carlsbad, USA) as previously described ²² using in-house RUNX1 antibody ⁶⁵. For all datasets, sequences were aligned to the mm10 mouse genome using Bowtie2 algorithm. Peak calling from read count followed the algorithm for MACS2 with a p-value cut-off = 10^{-10} . Overlapped peaks determined using ChIPseeker were used as the consensus data in all subsequent comparisons. Differential binding analysis for RUNX1 ChIP-seq was performed using DiffBind as described above. Peak characterisation was determined using the ChIPseeker package. Motif analysis for known and *de novo* sequence motifs was performed using HOMER as described above. Visualisation of ChIP-seq signal was through the UCSC Genome Browser.

Proximity ligation assay (PLA)

PLA was performed on cultured granulosa cells using the Duolink PLA Probes and PLA Fluorescence in situ Detection Kit Red (Sigma) following the manufacturer's protocol. Briefly, cultured granulosa cells were fixed in 4% paraformaldehyde and permeabilised with PBS + 0.01% Triton X-100 for 1 hour at room temperature. Cells were blocked with Blocking Buffer for 1 hour at 37°C and incubated with primary antibody couples (PGR –Cell Signalling Technology (CST); RUNX1 – Santa Cruz Biotechnology (SCB); RUNX2 – SCB; c-JUN – SCB; JUNB – SCB; JUND – SCB; LRH1 – Perseus Proteomics, CBP/p300 – CST) diluted in Antibody Diluent for 2 hours at room temperature or overnight at 4°C. Then, cells

were incubated with PLA probes of appropriate species for 1 hour at 37°C, oligo probes were ligated for 30 minutes at 37°C and the amplification reaction was at 37°C for a minimum of 100 minutes. Between steps, cells were washed using the provided wash buffers. Slides were mounted with Prolong Gold Mounting Media with DAPI (Thermo Fisher), cured for at least 1 hour in the dark and were stored at -20°C prior to imaging by Olympus confocal microscope. The nuclear boundary for each cell was determined through DAPI staining. PLA signals were identified as fluorescent puncta in the nucleus and quantified using the ‘Count Maxima’ function in ImageJ for each cellular compartment. Significant differences between time points were determined through one-way ANOVA with Tukey test for multiple comparison.

Data availability

The datasets generated and/or analysed during the current study are available in the NCBI Gene Expression Omnibus repository: GSE166878 (LH and PGRKO ATAC-seq), GSE178314 (LH RNA-seq), GSE168213 (PGRKO RNA-seq), GSE152941 (RUNX1 ChIP-seq).

Acknowledgement

The RUNX1 antibody for ChIP-seq was a kind gift from Dr Yoram Groner and Dr Ditsa Levanon, the Weizmann Institute of Science, Israel. This work was partially supported by the Intramural Research Program of National Institute of Environmental Health Sciences in the USA (Z01-ES102965 for HHCY).

Author contribution

D.L.R conceived and designed the study. T.D.D, J.B, R.L.R and D.L.R conceived and designed the experiments. T.D.D, B.N, K.M.S, M.N, A.E, Y.Y.W performed experiments. T.D.D and J.B analysed data. T.D.D, S.C.B, H.H.C.Y, R.L.R and D.L.R interpreted results.

T.D.D prepared figures. T.D.D and D.L.R drafted manuscript. All authors contributed to manuscript editing and revision.

Competing interests

The authors declare no competing interest.

Materials & Correspondence

Correspondence to Thao D Dinh and Darryl L Russell.

5. REFERENCES

- 1 Russell, D. L. & Robker, R. L. in *The Ovary (Third Edition)* (eds Peter C. K. Leung & Eli Y. Adashi) 217-234 (Academic Press, 2019).
- 2 Hughes, C. H. K. & Murphy, B. D. Nuclear receptors: Key regulators of somatic cell functions in the ovulatory process. *Molecular Aspects of Medicine*, 100937, doi:<https://doi.org/10.1016/j.mam.2020.100937> (2020).
- 3 Wu, S.-P., Li, R. & DeMayo, F. J. Progesterone Receptor Regulation of Uterine Adaptation for Pregnancy. *Trends in Endocrinology & Metabolism* **29**, 481-491, doi:<https://doi.org/10.1016/j.tem.2018.04.001> (2018).
- 4 Conneely, O. M., Mulac-Jericevic, B. & Arnett-Mansfield, R. in *Progestins and the Mammary Gland*. (eds O. Conneely & C. Otto) 175-185 (Springer Berlin Heidelberg).
- 5 Robker, R. L., Russell, D. L., Espey, L. L., Lydon, J. P., O'Malley, B. W. & Richards, J. S. Progesterone-regulated genes in the ovulation process: ADAMTS-1 and cathepsin L proteases. *Proc Natl Acad Sci U S A* **97**, 4689-4694 (2000).

- 533 6 Park, C. J., Lin, P.-C., Zhou, S., Barakat, R., Bashir, S. T., Choi, J. M., Cacioppo, J. A.,
534 Oakley, O. R., Duffy, D. M., Lydon, J. P. & Ko, C. J. Progesterone Receptor Serves the
535 Ovary as a Trigger of Ovulation and a Terminator of Inflammation. *Cell Reports* **31**,
536 107496, doi:<https://doi.org/10.1016/j.celrep.2020.03.060> (2020).
- 537 7 Lydon, J. P., DeMayo, F. J., Funk, C. R., Mani, S. K., Hughes, A. R., Montgomery, C. A.,
538 Jr., Shyamala, G., Conneely, O. M. & O'Malley, B. W. Mice lacking progesterone
539 receptor exhibit pleiotropic reproductive abnormalities. *Genes Dev* **9**, 2266-2278
540 (1995).
- 541 8 Ledger, W. L., Sweeting, V. M., Hillier, H. & Baird, D. T. Inhibition of ovulation by low-
542 dose mifepristone (RU 486). *Human Reproduction* **7**, 945-950,
543 doi:10.1093/oxfordjournals.humrep.a137776 (1992).
- 544 9 Brache, V., Cochon, L., Jesam, C., Maldonado, R., Salvatierra, A. M., Levy, D. P.,
545 Gainer, E. & Croxatto, H. B. Immediate pre-ovulatory administration of 30 mg
546 ulipristal acetate significantly delays follicular rupture. *Human Reproduction* **25**,
547 2256-2263, doi:10.1093/humrep/deq157 (2010).
- 548 10 Clemens, J. W., Robker, R. L., Kraus, W. L., Katzenellenbogen, B. S. & Richards, J. S.
549 Hormone induction of progesterone receptor (PR) messenger ribonucleic acid and
550 activation of PR promoter regions in ovarian granulosa cells: evidence for a role of
551 cyclic adenosine 3',5'-monophosphate but not estradiol. *Mol Endocrinol* **12**, 1201-
552 1214, doi:10.1210/mend.12.8.0157 (1998).
- 553 11 Ismail, P. M., Li, J., DeMayo, F. J., O'Malley, B. W. & Lydon, J. P. A Novel LacZ
554 Reporter Mouse Reveals Complex Regulation of the Progesterone Receptor
555 Promoter During Mammary Gland Development. *Molecular Endocrinology* **16**, 2475-
556 2489, doi:10.1210/me.2002-0169 (2002).

557 12 Yin, P., Roqueiro, D., Huang, L., Owen, J. K., Xie, A., Navarro, A., Monsivais, D., Coon
558 V, J. S., Kim, J. J., Dai, Y. & Bulun, S. E. Genome-Wide Progesterone Receptor Binding:
559 Cell Type-Specific and Shared Mechanisms in T47D Breast Cancer Cells and Primary
560 Leiomyoma Cells. *PLOS ONE* **7**, e29021, doi:10.1371/journal.pone.0029021 (2012).

561 13 Li, X. & O'Malley, B. W. Unfolding the Action of Progesterone Receptors. *Journal of*
562 *Biological Chemistry* **278**, 39261-39264, doi:10.1074/jbc.R300024200 (2003).

563 14 Han, S. J., DeMayo, F. J., Xu, J., Tsai, S. Y., Tsai, M.-J. & O'Malley, B. W. Steroid
564 Receptor Coactivator (SRC)-1 and SRC-3 Differentially Modulate Tissue-Specific
565 Activation Functions of the Progesterone Receptor. *Molecular Endocrinology* **20**, 45-
566 55, doi:10.1210/me.2005-0310 (2006).

567 15 Béguelin, W., Díaz Flaqué, M. C., Proietti, C. J., Cayrol, F., Rivas, M. A., Tkach, M.,
568 Rosembli, C., Tocci, J. M., Charreau, E. H., Schillaci, R. & Elizalde, P. V. Progesterone
569 receptor induces ErbB-2 nuclear translocation to promote breast cancer growth via a
570 novel transcriptional effect: ErbB-2 function as a coactivator of Stat3. *Molecular and*
571 *cellular biology* **30**, 5456-5472, doi:10.1128/MCB.00012-10 (2010).

572 16 Malbeteau, L., Poulard, C., Languilaire, C., Mikaelian, I., Flamant, F., Le Romancer, M.
573 & Corbo, L. PRMT1 Is Critical for the Transcriptional Activity and the Stability of the
574 Progesterone Receptor. *iScience* **23**, 101236,
575 doi:<https://doi.org/10.1016/j.isci.2020.101236> (2020).

576 17 Cho, H., Orphanides, G., Sun, X., Yang, X. J., Ogryzko, V., Lees, E., Nakatani, Y. &
577 Reinberg, D. A human RNA polymerase II complex containing factors that modify
578 chromatin structure. *Molecular and cellular biology* **18**, 5355-5363,
579 doi:10.1128/mcb.18.9.5355 (1998).

580 18 Dong, X., Challis, J. R. & Lye, S. J. Intramolecular interactions between the AF3
581 domain and the C-terminus of the human progesterone receptor are mediated
582 through two LXXLL motifs. *J Mol Endocrinol* **32**, 843-857, doi:10.1677/jme.0.0320843
583 (2004).

584 19 Merlino, A. A., Welsh, T. N., Tan, H., Yi, L. J., Cannon, V., Mercer, B. M. & Mesiano, S.
585 Nuclear Progesterone Receptors in the Human Pregnancy Myometrium: Evidence
586 that Parturition Involves Functional Progesterone Withdrawal Mediated by Increased
587 Expression of Progesterone Receptor-A. *The Journal of Clinical Endocrinology &*
588 *Metabolism* **92**, 1927-1933, doi:10.1210/jc.2007-0077 (2007).

589 20 Menendez, J. A., Peirce, S. K., Papadimitropoulou, A., Cuyàs, E., Steen, T. V., Verdura,
590 S., Vellon, L., Chen, W. Y. & Lupu, R. Progesterone receptor isoform-dependent
591 cross-talk between prolactin and fatty acid synthase in breast cancer. *Aging (Albany*
592 *NY)* **12**, 24671-24692, doi:10.18632/aging.202289 (2020).

593 21 Teilmann, S. C., Clement, C. A., Thorup, J., Byskov, A. G. & Christensen, S. T.
594 Expression and localization of the progesterone receptor in mouse and human
595 reproductive organs. *J Endocrinol* **191**, 525-535, doi:10.1677/joe.1.06565 (2006).

596 22 Dinh, D. T., Breen, J., Akison, L. K., DeMayo, F. J., Brown, H. M., Robker, R. L. &
597 Russell, D. L. Tissue-specific progesterone receptor-chromatin binding and the
598 regulation of progesterone-dependent gene expression. *Scientific reports* **9**, 11966-
599 11966, doi:10.1038/s41598-019-48333-8 (2019).

600 23 Mulac-Jericevic, B., Mullinax, R. A., DeMayo, F. J., Lydon, J. P. & Conneely, O. M.
601 Subgroup of Reproductive Functions of Progesterone Mediated by Progesterone
602 Receptor-B Isoform. *Science* **289**, 1751-1754, doi:10.1126/science.289.5485.1751
603 (2000).

- 604 24 Mulac-Jericevic, B., Lydon, J. P., DeMayo, F. J. & Conneely, O. M. Defective mammary
605 gland morphogenesis in mice lacking the progesterone receptor B isoform.
606 *Proceedings of the National Academy of Sciences* **100**, 9744-9749,
607 doi:10.1073/pnas.1732707100 (2003).
- 608 25 Hu, M., Li, J., Zhang, Y., Li, X., Brännström, M., Shao, L. R. & Billig, H. Endometrial
609 progesterone receptor isoforms in women with polycystic ovary syndrome. *Am J*
610 *Transl Res* **10**, 2696-2705 (2018).
- 611 26 Wetendorf, M., Li, R., Wu, S.-P., Liu, J., Creighton, C. J., Wang, T., Janardhan, K. S.,
612 Willson, C. J., Lanz, R. B., Murphy, B. D., Lydon, J. P. & DeMayo, F. J. Constitutive
613 expression of progesterone receptor isoforms promotes the development of
614 hormone-dependent ovarian neoplasms. *Science Signaling* **13**, eaaz9646,
615 doi:10.1126/scisignal.aaz9646 (2020).
- 616 27 Liu, J., Park, E.-S. & Jo, M. Runt-related transcription factor 1 regulates luteinized
617 hormone-induced prostaglandin-endoperoxide synthase 2 expression in rat
618 periovulatory granulosa cells. *Endocrinology* **150**, 3291-3300 (2009).
- 619 28 Liu, J., Park, E.-S., Curry, T. E., Jr. & Jo, M. Periovulatory expression of hyaluronan and
620 proteoglycan link protein 1 (Hapln1) in the rat ovary: hormonal regulation and
621 potential function. *Mol Endocrinol* **24**, 1203-1217, doi:10.1210/me.2009-0325
622 (2010).
- 623 29 Lee-Thacker, S., Choi, Y., Taniuchi, I., Takarada, T., Yoneda, Y., Ko, C. & Jo, M. Core
624 binding factor β expression in ovarian granulosa cells is essential for female fertility.
625 *Endocrinology* **159**, 2094-2109 (2018).

- 626 30 Lee-Thacker, S., Jeon, H., Choi, Y., Taniuchi, I., Takarada, T., Yoneda, Y., Ko, C. & Jo,
627 M. Core Binding Factors are essential for ovulation, luteinization, and female fertility
628 in mice. *Scientific Reports* **10**, 9921, doi:10.1038/s41598-020-64257-0 (2020).
- 629 31 Nicol, B., Grimm, S. A., Chalmel, F., Lecluze, E., Pannetier, M., Pailhoux, E., Dupin-De-
630 Beyssat, E., Guiguen, Y., Capel, B. & Yao, H. H. C. RUNX1 maintains the identity of the
631 fetal ovary through an interplay with FOXL2. *Nature communications* **10**, 5116
632 (2019). <<https://doi.org/10.1038/s41467-019-13060-1>>.
- 633 32 Park, E.-S., Park, J., Franceschi, R. T. & Jo, M. The role for runt related transcription
634 factor 2 (RUNX2) as a transcriptional repressor in luteinizing granulosa cells.
635 *Molecular and Cellular Endocrinology* **362**, 165-175,
636 doi:<https://doi.org/10.1016/j.mce.2012.06.005> (2012).
- 637 33 Kim, J., Sato, M., Li, Q., Lydon, J. P., DeMayo, F. J., Bagchi, I. C. & Bagchi, M. K.
638 Peroxisome Proliferator-Activated Receptor γ Is a Target of Progesterone Regulation
639 in the Preovulatory Follicles and Controls Ovulation in Mice. *Molecular and cellular*
640 *biology* **28**, 1770-1782, doi:10.1128/mcb.01556-07 (2008).
- 641 34 Kim, J., Bagchi, I. C. & Bagchi, M. K. Signaling by hypoxia-inducible factors is critical
642 for ovulation in mice. *Endocrinology* **150**, 3392-3400, doi:10.1210/en.2008-0948
643 (2009).
- 644 35 Sharma, S. C. & Richards, J. S. Regulation of AP1 (Jun/Fos) factor expression and
645 activation in ovarian granulosa cells. Relation of JunD and Fra2 to terminal
646 differentiation. *J Biol Chem* **275**, 33718-33728, doi:10.1074/jbc.M003555200 (2000).
- 647 36 Jo, M. & Curry, T. E., Jr. Luteinizing Hormone-Induced RUNX1 Regulates the
648 Expression of Genes in Granulosa Cells of Rat Periovarian Follicles. *Molecular*
649 *Endocrinology* **20**, 2156-2172, doi:10.1210/me.2005-0512 (2006).

650 37 Fan, H.-Y., Liu, Z., Johnson, P. F. & Richards, J. S. CCAAT/Enhancer-Binding Proteins
651 (C/EBP)- α and - β Are Essential for Ovulation, Luteinization, and the Expression of Key
652 Target Genes. *Molecular Endocrinology* **25**, 253-268, doi:10.1210/me.2010-0318
653 (2011).

654 38 Russell, D. L., Doyle, K. M. H., Gonzales-Robayna, I., Pipaon, C. & Richards, J. S. Egr-1
655 Induction in Rat Granulosa Cells by Follicle-Stimulating Hormone and Luteinizing
656 Hormone: Combinatorial Regulation By Transcription Factors Cyclic Adenosine 3',5'-
657 Monophosphate Regulatory Element Binding Protein, Serum Response Factor, Sp1,
658 and Early Growth Response Factor-1. *Molecular Endocrinology* **17**, 520-533,
659 doi:10.1210/me.2002-0066 (2003).

660 39 Puri, P., Little-Ihrig, L., Chandran, U., Law, N. C., Hunzicker-Dunn, M. & Zeleznik, A. J.
661 Protein Kinase A: A Master Kinase of Granulosa Cell Differentiation. *Scientific Reports*
662 **6**, 28132, doi:10.1038/srep28132 (2016).

663 40 Robker, R. L., Russell, D. L., Espey, L. L., Lydon, J. P., O'Malley, B. W. & Richards, J. S.
664 Progesterone-regulated genes in the ovulation process: ADAMTS-1 and cathepsin L
665 proteases. *Proceedings of the National Academy of Sciences* **97**, 4689-4694,
666 doi:10.1073/pnas.080073497 (2000).

667 41 Grimm, S. L., Hartig, S. M. & Edwards, D. P. Progesterone Receptor Signaling
668 Mechanisms. *Journal of Molecular Biology* **428**, 3831-3849,
669 doi:<https://doi.org/10.1016/j.jmb.2016.06.020> (2016).

670 42 Chuang, L. S. H., Ito, K. & Ito, Y. RUNX family: Regulation and diversification of roles
671 through interacting proteins. *International Journal of Cancer* **132**, 1260-1271,
672 doi:10.1002/ijc.27964 (2013).

673 43 Barutcu, A. R., Hong, D., Lajoie, B. R., McCord, R. P., van Wijnen, A. J., Lian, J. B.,
674 Stein, J. L., Dekker, J., Imbalzano, A. N. & Stein, G. S. RUNX1 contributes to higher-
675 order chromatin organization and gene regulation in breast cancer cells. *Biochimica*
676 *et Biophysica Acta (BBA) - Lipids and Lipid Metabolism* **1859**, 1389-1397,
677 doi:10.1016/j.bbagrm.2016.08.003 (2016).

678 44 Kawate, H., Wu, Y., Ohnaka, K. & Takayanagi, R. Mutual transactivational repression
679 of Runx2 and the androgen receptor by an impairment of their normal
680 compartmentalization. *The Journal of Steroid Biochemistry and Molecular Biology*
681 **105**, 46-56, doi:<https://doi.org/10.1016/j.jsbmb.2006.11.020> (2007).

682 45 Yang, R., Browne, J. A., Eggener, S. E., Leir, S.-H. & Harris, A. A novel transcriptional
683 network for the androgen receptor in human epididymis epithelial cells. *Molecular*
684 *Human Reproduction* **24**, 433-443, doi:10.1093/molehr/gay029 (2018).

685 46 Ning, Y.-M. & Robins, D. M. AML3/CBF α 1 Is Required for Androgen-specific
686 Activation of the Enhancer of the Mouse Sex-limited Protein (Slp) Gene. *Journal of*
687 *Biological Chemistry* **274**, 30624-30630, doi:10.1074/jbc.274.43.30624 (1999).

688 47 Takayama, K.-i., Suzuki, T., Tsutsumi, S., Fujimura, T., Urano, T., Takahashi, S.,
689 Homma, Y., Aburatani, H. & Inoue, S. RUNX1, an androgen- and EZH2-regulated
690 gene, has differential roles in AR-dependent and -independent prostate cancer.
691 *Oncotarget* **6** (2014).

692 48 Stender, J. D., Kim, K., Charn, T. H., Komm, B., Chang, K. C. N., Kraus, W. L., Benner,
693 C., Glass, C. K. & Katzenellenbogen, B. S. Genome-Wide Analysis of Estrogen
694 Receptor α DNA Binding and Tethering Mechanisms Identifies Runx1 as a Novel
695 Tethering Factor in Receptor-Mediated Transcriptional Activation. *Molecular and*
696 *cellular biology* **30**, 3943-3955, doi:10.1128/mcb.00118-10 (2010).

697 49 Falender, A. E., Lanz, R., Malenfant, D., Belanger, L. & Richards, J. S. Differential
698 Expression of Steroidogenic Factor-1 and FTF/LRH-1 in the Rodent Ovary.
699 *Endocrinology* **144**, 3598-3610, doi:10.1210/en.2002-0137 (2003).

700 50 Choi, Y., Rosewell, K. L., Brännström, M., Akin, J. W., Curry Jr, T. E. & Jo, M. FOS, a
701 Critical Downstream Mediator of PGR and EGF Signaling Necessary for Ovulatory
702 Prostaglandins in the Human Ovary. *The Journal of Clinical Endocrinology &*
703 *Metabolism* **103**, 4241-4252 (2018).

704 51 Corces, M. R., Trevino, A. E., Hamilton, E. G., Greenside, P. G., Sinnott-Armstrong, N.
705 A., Vesuna, S., Satpathy, A. T., Rubin, A. J., Montine, K. S., Wu, B., Kathiria, A., Cho, S.
706 W., Mumbach, M. R., Carter, A. C., Kasowski, M., Orloff, L. A., Risca, V. I., Kundaje, A.,
707 Khavari, P. A., Montine, T. J., Greenleaf, W. J. & Chang, H. Y. An improved ATAC-seq
708 protocol reduces background and enables interrogation of frozen tissues. *Nature*
709 *Methods* **14**, 959-962, doi:10.1038/nmeth.4396 (2017).

710 52 Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
711 reads. *EMBnet. journal* **17**, 10-12 (2011).

712 53 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature*
713 *Methods* **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

714 54 Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E.,
715 Nusbaum, C., Myers, R. M., Brown, M., Li, W. & Liu, X. S. Model-based analysis of
716 ChIP-Seq (MACS). *Genome Biol* **9**, R137, doi:10.1186/gb-2008-9-9-r137 (2008).

717 55 Stark R, B. G. DiffBind: differential binding analysis of ChIP-Seq peak data. (2011).

718 56 Yu, G., Wang, L. G. & He, Q. Y. ChIPseeker: an R/Bioconductor package for ChIP peak
719 annotation, comparison and visualization. *Bioinformatics* **31**, 2382-2383,
720 doi:10.1093/bioinformatics/btv145 (2015).

721 57 McLean, C. Y., Bristor, D., Hiller, M., Clarke, S. L., Schaar, B. T., Lowe, C. B., Wenger,
722 A. M. & Bejerano, G. GREAT improves functional interpretation of cis-regulatory
723 regions. *Nature biotechnology* **28**, 495 (2010).

724 58 Bentsen, M., Goymann, P., Schultheis, H., Klee, K., Petrova, A., Wiegandt, R., Fust, A.,
725 Preussner, J., Kuenne, C., Braun, T., Kim, J. & Looso, M. ATAC-seq footprinting
726 unravels kinetics of transcription factor binding during zygotic genome activation.
727 *Nature Communications* **11**, 4267, doi:10.1038/s41467-020-18035-1 (2020).

728 59 Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre,
729 C., Singh, H. & Glass, C. K. Simple combinations of lineage-determining transcription
730 factors prime cis-regulatory elements required for macrophage and B cell identities.
731 *Mol Cell* **38**, 576-589, doi:10.1016/j.molcel.2010.05.004 (2010).

732 60 Lindgreen, S. AdapterRemoval: easy cleaning of next-generation sequencing reads.
733 *BMC Research Notes* **5**, 337, doi:10.1186/1756-0500-5-337 (2012).

734 61 Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast
735 and bias-aware quantification of transcript expression. *Nature methods* **14**, 417
736 (2017).

737 62 Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W. & Smyth, G. K. limma
738 powers differential expression analyses for RNA-sequencing and microarray studies.
739 *Nucleic Acids Research* **43**, e47-e47, doi:10.1093/nar/gkv007 (2015).

740 63 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
741 differential expression analysis of digital gene expression data. *Bioinformatics*
742 (Oxford, England) **26**, 139-140, doi:10.1093/bioinformatics/btp616 (2010).

743 64 Ramírez, F., Ryan, D. P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., Heyne, S.,
744 Dündar, F. & Manke, T. deepTools2: a next generation web server for deep-

745 sequencing data analysis. *Nucleic Acids Research* **44**, W160-W165,
746 doi:10.1093/nar/gkw257 (2016).
747 65 Umansky, K. B., Gruenbaum-Cohen, Y., Tsoory, M., Feldmesser, E., Goldenberg, D.,
748 Brenner, O. & Groner, Y. Runx1 Transcription Factor Is Required for Myoblasts
749 Proliferation during Muscle Regeneration. *PLOS Genetics* **11**, e1005457,
750 doi:10.1371/journal.pgen.1005457 (2015).

751

752

753

754

Figure 1: Global changes in the chromatin landscape of granulosa cells after hCG treatment. (A) Heatmap of LH ATAC-peak signals (left), with peaks subset into those that were unchanged (top), LH-induced (middle) or LH-repressed (bottom). Representative profile plots of each condition are on the right. (B) Genomic distribution of ATAC-peaks that were unchanged by LH, LH-induced or LH-repressed. (C) Differential enrichment analysis of transcription factor-binding motifs in ATAC-seq peaks found in 6h-hCG and 0h-hCG. Top 5% motifs that were differentially enriched in either condition are highlighted. (D) *de novo* motif enrichment analysis of LH-induced, LH-repressed or unchanged ATAC peaks as identified through HOMER. For each dataset, top 5 motifs are shown with p-value and fold change to background (FC). (E) DNA footprints of selected motifs in ATAC-peaks that were LH-induced (red) or LH-repressed (pink).

Figure 2: Global changes in the transcriptome of granulosa cells after hCG treatment.

(A) Heatmap of the expression level of ovulatory DEG identified in LH RNA-seq. (B) Correlation between expression level and promoter accessibility at RNA-seq identified ovulatory genes, shown as LH DEG RNA-seq log fold change vs ATAC-seq log fold change at their respective promoter (within 3 kb of TSS). DEGs with LH-dependent promoter are labelled red, DEGs with LH-independent promoter are labelled pink. Correlation between RNA-seq and ATAC-seq intensity is determined through Spearman's correlation coefficient ($r = 0.4297$, $p < 0.0001$) (C) Examples of ATAC-seq (track 1, 2) and RNA-seq (track 3, 4) signals for 0h-hCG (light) and 6h/8h-hCG (dark) at *Cxcr4*, *Inhbb* and *Cited1*. Differentially accessible ATAC peaks are highlighted in yellow.

779

780

Figure 3: PGR and RUNX1 share mutual chromatin targets in peri-ovulatory granulosa

cells. (A) Heatmap of RUNX1 ChIP-seq peak signals, with peaks subset into those that were unchanged (top) or LH-induced (bottom). Only 1 RUNX1 peak is LH-repressed and is thus not plotted (B) Venn diagram showing shared and factor-unique peak counts for RUNX1 and PGR in relation to 6h-hCG LH ATAC. (C) Genomic distribution of transcriptionally active PGR-specific (top), PGR/RUNX1 shared (middle) and RUNX1-specific (bottom) binding sites. (D) Venn diagram showing the relationship between LH DEG and genes with RUNX1 and/or PGR binding in their promoters. (E) Example of LH-dependent RUNX1 binding (track 1, 2) in conjunction with PGR (track 3) at *Rgcc*, *Adamts1* and *Zbtb16*. ATAC-seq peaks at 0h, 6h and H3K27ac ChIP-seq peaks at 6h (track 4-6) indicate open chromatin regions. (F) Top most common known sequence motifs found to be enriched at RUNX1 (pink) and PGR (orange) binding sites. Bars indicate fold enrichment of motif to background (bottom x axis). Circles indicate $-\log(p\text{-value})$ (top x axis). (G) PLA for PGR/RUNX1 in cultured granulosa cells under hCG/R5020 time course treatment. One representative image is shown for each treatment. PGR/IgG antibody pair was used as negative control. PLA was performed in 3 biological replicates (4 mice per replicate). Red indicates protein-protein interaction. Blue is DAPI nuclear stain. Scale bar = 10 μm . Quantification of PLA signal is displayed as the number of nuclear foci per cell, displayed as mean $\pm$ SEM, p-value = 0.1044.

Figure 4: PGR interacts with other ovulatory transcription factors in granulosa cells in response to the LH surge. (A) PLA for PGR/RUNX2 in cultured granulosa cells under hCG/R5020 time course treatment. One representative image is shown for each treatment. PGR/IgG antibody pair was used as negative control. PLA was performed in 3 biological replicates (4 mice per replicate). Quantification of PLA signal is displayed as the number of nuclear foci per cell, displayed as mean $\pm$ SEM, p-value = 0.2294. (B) PLA showing the interaction between PGR and LRH1, JUND, cJUN and JUNB. Red indicates protein-protein interaction. Blue is DAPI nuclear stain. Scale bar = 10 μ m.

810

811

Figure 5: PGR role in mediating chromatin accessibility in granulosa cells (A) Heatmap of PGR ATAC-peaks (left) that were unchanged (top) or differentially accessible between PGRKO WT and KO (bottom). Representative profile plots of each group are on the right. (B) Genomic distribution of peaks that were differentially enriched in PGRKO. (C) Correlation between expression level and promoter accessibility at PGR DEGs, shown as PGR DEG fold change vs ATAC fold change at their respective promoter. DEGs with PGR-dependent promoter accessibility are labelled light green, DEGs with promoter accessibility unchanged by PGR are labelled dark green. Correlation between RNA-seq and ATAC-seq intensity is determined through Spearman's correlation coefficient ($r = 0.5116$, $p < 0.0001$) (D) Examples of LH ATAC-seq (track 1, 2), PGR ATAC-seq (track 3, 4), PGR and H3K27ac ChIP-seq (track 5, 6) and PGR RNA-seq (track 7, 8) signals at *Gas7* and *Abhd2*. ATAC peaks at promoters are highlighted in yellow. (E) Top most common known sequence motifs found to be enriched at PGR-dependent ATAC peaks (green), with the level of enrichment for the same motif at PGR binding sites as reference (orange). Bars indicate fold enrichment of motif to background (bottom x axis). Circles indicate $-\log(p\text{-value})$ (top x axis). (F) Upset plot showing the number of overlapped peaks between PGR-dependent ATAC sites, PGR and RUNX1 binding sites.

Figure 6: PGR isoform-driven transcriptome in peri-ovulatory granulosa cells through interaction with specific transcription factor partners. (A) Heatmap showing changes in gene expression in granulosa cells that are WT or KO for total PGR, PGR-A or PGR-B, each shown with the collective PGRKO, AKO and BKO DEGs. (B) Upset plot showing the number of overlapped genes between PGRKO/AKO/BKO DEG and LH DEG. (C) Top 50 upstream regulators of PGRKO and AKO DEGs as identified through IPA. Circle colour indicates the number of downstream DEG regulated by the upstream regulator, circle size indicates $-\log(p\text{-value})$ of regulator enrichment. (D) PLA of PGR/RUNX1 (top) and PGR/RUNX2 (bottom) in *in vitro* WT/het/KO granulosa cells from AKO or BKO mice, treated for 6 h with hCG and R5020. Red indicates protein-protein interaction. Blue is DAPI nuclear stain. Scale bar = 10 μm . Quantification of PLA signal is displayed as the number of nuclear foci per cell, displayed as mean $\pm$ SEM, * = $p\text{-value} < 0.05$, N = 3 biological replicates (2 mice per replicate per genotype).

846

847 **Figure 7: Model of PGR action in peri-ovulatory granulosa cells.**