The Rise of West Nile Virus in Southern and Southeastern Europe: a spatial-temporal analysis investigating the combined effects of climate, land use and economic changes

Matthew J. Watts^a, Victor Sarto i Monteys^{a,b}, P. Graham Mortyn^{a,d},
Panagiota Kotsila^{a,c}

^a*Institute of Environmental Science and Technology (ICTA) , Autonomous University of Barcelona (UAB), Bellaterra, Spain*

^b*Departament d'Agricultura, Ramaderia, Pesca, Alimentació i Medi Natural, Generalitat de Catalunya, Avinguda Meridiana, Barcelona, Spain*

^c*Barcelona Laboratory for Urban Environmental Justice and Sustainability (BCNEJ), Institute of Environmental Science and Technology (ICTA) , Autonomous University of Barcelona (UAB), Bellaterra, Spain*

^d*Department of Geography, Autonomous University of Barcelona (UAB), Bellaterra, Spain*

Abstract

West Nile Virus (WNV) has recently emerged as a major public health concern in Europe, its recent expansion also coincided with some remarkable socio-economic and environmental changes, including an economic crisis and some of the warmest temperatures on record. Here we empirically investigate the drivers of this phenomenon at a European wide scale by constructing and analyzing a unique spatial-temporal data-set, that includes data on climate, land-use, the economy, and government spending on environmental related sectors. Drivers and risk factors of WNV were identified by building a conceptual framework, and relationships were tested using a Generalized Additive Model (GAM), which could capture complex non-linear relationships and also account for the spatial and temporal auto-correlation. Some of the key risk factors identified in our conceptual framework, such as a higher percentage of wetlands and arable land, climate factors (higher summer rainfall and higher summer temperatures) were positive predictors of WNV infections. Interestingly, winter temperatures of

*Corresponding author

Email address: matthewjohnwatts@gmail.com (Matthew J. Watts)

between 2 °C and 6 °C were among some of the strongest predictors of annual WNV infections, one possible explanation for this result is that successful overwintering of infected adult mosquitoes (likely *Culex pipiens*) is key to the intensity of outbreaks for a given year. Furthermore, lower surface water extent over the summer is also associated with more intense outbreaks, suggesting that drought, which is known to induce positive changes in WNV prevalence in mosquitoes is also contributing to the upward trend in WNV cases in affected regions. Our indicators representing the economic crisis were also strong predictors of WNV infections, suggesting there is an association between austerity and cuts to key sectors, which could have benefited vector species and the virus during this crucial period. These results, taken in the context of recent winter warming due to climate change, and more frequent droughts, may offer an explanation of why the virus has become so prevalent in Europe.

Keywords: West-Nile-virus, climate-change, economic-crisis, mosquito, austerity, vector-borne-disease, drought.

1. Introduction

Over the past few decades, new health risks have been emerging in Europe, particularly with the recent appearance of neglected tropical diseases (NTDs) such as Chikungunya, West Nile Virus (WNV), Dengue (DENV-1) and Crimean-
5 Congo haemorrhagic fever [1, 2, 3]. Rising temperatures are likely increasing the transmission potential of NTDs in Europe, by affecting the geographic spread, abundance, survival and feeding activity of vector species and benefiting pathogen development in infected vectors [4, 5, 6, 7, 8, 9]. This, combined with other factors such as human population growth, intensive animal rearing,
10 global commerce, air travel, urbanization and land-use changes, is increasing the chances of novel diseases to enter and emerge in Europe [10, 11, 12, 13].

In this study, we focus on WNV, a single-stranded RNA *Flavivirus* closely related to other *Flaviviridae* pathogens such as dengue, Japanese encephalitis and yellow fever viruses [14]. Although WNV is a zoonotic pathogen, infecting

15 mammals, particularly humans and horses, the transmission cycle is believed to be driven mainly by mosquitoes and birds [15], although some wild mammals may serve as intermediate hosts for West Nile virus [16]. Generally, WNV distribution is determined by the presence of suitable mosquito vectors and avian hosts, such as terrestrial and wetland birds. The spring migration of birds
20 from infected regions of Sub-Saharan Africa to temperate regions of Europe is considered to be one of the main drivers of the disease in Europe [17, 18]. In Europe, the main vectors are *Culex pipiens*, *Culex modestus*, and *Coquilleltidia richiardii*, although *Aedes* species can also transmit the disease [18] (see figures S1-S3 in additional information for vector distribution maps).

25 Although WNV infections are more common of late, sporadic West Nile virus outbreaks have occurred in humans and equines in southern and eastern European countries over the last century; the majority of which have occurred in wetland areas and densely inhabited urban areas [18].

Since 2010, WNV has been reported in 13 EU countries including Austria,
30 Bulgaria, Croatia, Cyprus, Czechia, France, Greece, Hungary, Italy, Portugal, Romania, Slovenia and Spain. It has also been reported in and five EU candidate countries including Albania, Montenegro, Serbia, Turkey and Kosovo [19]. In 2010, major outbreaks hit Greece, Hungary, Romania, and Turkey. Since then, outbreaks have occurred annually in multiple regions, including more northerly
35 regions that had not previously reported cases, including Germany, Slovakia, Slovenia, and the Czech Republic (see Figure 1). This culminated in another major outbreak in 2018, that affected more regions than had been recorded in previous years.

We should also bear in mind that the actual number of cases in Europe is
40 likely to be much higher than reported, since most people infected with WNV will not develop symptoms (are asymptomatic). We can therefore assume that only serious cases are reported. Around 20% of those infected with WNV will develop West Nile fever, a flu like illness, or severe West Nile disease. The most severe manifestations of the disease include acute aseptic meningitis or
45 encephalitis, anterior myelitis, hepatosplenomegaly, hepatitis, pancreatitis, and

myocarditis. On rare occasions, infection can lead to Guillain–Barré syndrome and other demyelinating neuropathies [15, 14].

1.1. Motivation for study

West Nile Virus (WNV) has recently emerged as a major public health concern in Europe, its recent expansion also coincided with some remarkable socio-economic and environmental changes, including an economic crisis and some of the warmest temperatures on record. To date, there is current very little research investigating this phenomenon at a European wide scale and more work is required to reveal the key drivers of the disease. A better understanding of this phenomenon can help public health officials design health prevention measures and develop better predictive models for public health risk management. Furthermore, little work has been done to explore the association between the rise of WNV in Europe and the economic crisis. Although the study of physical factors is key in understanding disease transmission and distribution, few articles have considered how anthropogenic factors, like economic factors and policy making [20, 21], which can have wider and unintended effects on natural ecosystems and, eventually, on disease [22]. Although examining such factors presents certain challenges and uncertainties given the scales involved and lack of data. We feel there is a need to examine the statistical relationships more closely, at the very least to open up the scholarly debate and instigate further research.

To examine the recent rise of WNV infections in Europe in more depth, we empirically investigate the combined effect of three sets of factors: (1) Climate/environmental factors including temperature, rainfall and surface water; (2) land-use factors including continuous/discontinuous urban fabric, regional coverage of wetlands and arable land; (3) socio-economic factors that capture the associations of the economic crisis and are proxied by GDP growth, central government spending on areas of the environment including agriculture, forestry and fisheries and waste water management. Our analysis focuses on regions in the seven European countries where WNV has been regularly reported - Austria,

Bulgaria, Croatia, Greece, Hungary Italy, and Romania. The time series data set captures the time period before and after the economic crisis (2007-2019).

Since WNV infection data is only available at the European NUTS 3 level (aggregated areal health data), our empirical strategy relies on aggregated areal
80 health data. We recognize that the use of aggregate data presents some challenges since we cannot account for individual heterogeneity, which may lead to confounding bias i.e., relationships observed at the group level does not necessarily hold for individuals. Additionally, we may not be able to draw causal inference due to the likely presence of endogeneity bias (e.g., omitted variable
85 bias or reverse causality). Nevertheless, we think this type of empirical investigation maintains high merit, as it allows us to quickly explore geographic associations between the disease and the predictor variables, which in turn allows us to further the debate on this topic and instigate more refined research.

1.2. Conceptual framework

90 WNV transmission requires the presence of competent vectors, a suitable climate and a susceptible host population. Studying WNV transmission at a macro scale presents significant challenges, since key data on the seasonal and annual abundance of competent mosquitoes and birds are not available, this is further complicated given the hundreds of potential host bird species in Europe
95 [23]. To explain human infections, we therefore use environmental risk factors known to attract vector, and host. Furthermore, vector abundance for a given season is modulated by physical environmental factors, such as temperature, rainfall and water resource availability [18, 24, 14]. We therefore use proxies that can predict mosquito abundance.

100

1.2.1. Modulating Factors

Typically, with most tropical and temperate mosquito species, elevated temperatures allow vector populations to increase their growth and reproduction rates, which in turn decreases blood meals intervals, accelerating transmission

105 and virus evolution rates [25]. Furthermore, increasingly warmer winters allow mosquito vectors to expand their breeding seasons and survive during winter, either as eggs or as overwintering female mosquitoes. Weather conditions and climatic factors can also affect vector competence [14]. Viral replication rates and transmission of WNV are modulated by ambient temperature, affecting the
110 length of the extrinsic incubation period (EIP), seasonal phenology of mosquito host populations and also times at which humans and mosquitoes come into contact [26]. Generally, higher rainfall in warmer weather can lead to higher mosquito abundance and disease transmission by increasing the potential habitat suitable for mosquito reproduction, e.g. standing water [27, 18, 28]. Con-
115 versely, sometimes drought and shrinking water resources can also bring some species into closer contact, facilitating transmission and amplification of WNV within these locations [14]. To represent these points in our data-set / model, we selected mean winter temperature, mean summer temperature, number of days of rainfall in summer and summer surface water extent (the number
120 of satellite surface water observations per region).

1.2.2. Risk factors

West Nile virus circulation in Europe is usually confined to two different cycles and ecosystems: the sylvatic and the urban synanthropic cycle. Rural locations, including river deltas and floodplain areas, help create a sylvatic
125 cycle, where wild, usually nesting wetland birds and ornithophilic mosquitoes (*Culex pipiens*, *Culex modestus*, *Coquillettidia richiardii*) create the conditions for maintaining WNV transmission. In urban synanthropic cycles mosquitoes, such as *Culex pipiens* or *Culex modestus*, feed on domestic birds and humans.
130 However, these two cycles can overlap, so areas with wetlands close to human population can be particularly at risk to the disease [18]. Irrigation from agriculture is also heavily linked to a greater incidence of human and veterinary WNV infections [29]. In order represent these factors in our data set and final models, we selected land use variables (% cover) representing urban areas (metro areas),

135 semi urban areas (lower density human settlements), wetlands and arable land.

1.2.3. *Economic crisis*

We would expect the repercussions of an economic crisis to affect WNV transmission in several ways, at the individual level (bottom-up) and govern-
 140 ment level (top-down). Previous studies have shown that socio-economic factors tend to influence the distribution and intensity of mosquito-borne diseases both pre-infection and post-infection [30]. Poorer communities are less likely to have air-conditioned homes, tap water and adequate drainage, and therefore may be more exposed to biting mosquitoes. Several studies have demonstrated the link
 145 between WNV infections and a range of local-level socio-economic and demographic factors such as income, sanitation, and population density [17, 31, 32]. In general, we would expect to see a drop in living standards in regions experiencing an economic shock followed by sluggish economic growth. Those people most affected would find it more difficult to prevent mosquito infections through
 150 direct measures i.e. sprays and repellents and less likely to pay for things that indirectly influence WNV transmission, like the up keep of homes and use of air conditioning. Factors associated with higher economic status can also bring humans into closer contact with mosquitoes; for example, home owners with gardens and potted plants, swimming pools and ponds or having good access
 155 to recreational space where mosquitoes can breed [33, 34]. However, neglect of such things through economic decline can have further unintended effects, even in wealthy neighborhoods [35, 36, 37].

Currently, the most effective way to prevent transmission of WNV from mosquito to human is through mosquito control (mosquito abatement). This
 160 is crucial to prevent transmission and prevent an epidemic once transmission has begun. It is well documented, for example, that during the European debt crisis, the Greek government cut mosquito abatement budgets which may have led to a rise in vector borne disease outbreaks such as malaria and WNV [38]. WNV transmission is most likely to occur in places that favor the larval devel-

165 opment of *Culex pipiens*, such as poorly drained low-lying areas, urban storm-water catch basins and manhole chambers, roadside ditches, sewage treatment lagoons, and man-made containers around houses, or other aquatic environments where mosquitoes deposit their eggs [14]. Additionally, during periods of austerity, governments can neglect hazard prevention efforts, such as spending
170 on flood defenses, as well as essential works like sanitation and up-keep and improvement of infrastructure [39]. Such degradation can lead to the creation of mosquito habitats [14, 18, 39]. Another critical component of preventing disease transmission is through public education programs and health promotion, educating the public on measures which can prevent being bitten can reduce
175 risk of exposure. In general, we would expect to see a general deterioration in a government ability to run such programs during crisis and austerity. Other consequences of austerity can be expected in decreased disease detection because of cuts in public health services, prolonged periods between initial infection and treatment seeking due to dysfunctional healthcare systems, and reduced
180 treatment of disease; all of which can lead to more intense outbreaks [40].

In order to represent the economic crisis in our model, we selected regional GDP and central government on spending on health and environment factors including agriculture, forest & fisheries and waste water management. Rather than using actual annual values, or year on year growth, we look at increases or
185 decreases in growth using 2007 baseline levels, just before the crisis hit Europe. As a priori, we would expect WNV incidence to be associated with negative growth or very low growth in these sectors.

2. Materials and methods

In this study, we compiled a unique spatial temporal data-set that captures
190 the main drivers and risk factors of WNV infections in Europe, based on findings from the conceptual framework. We selected regions for the study if autochthonous virus transmission had occurred at least once over the reporting period and if vectors were present in a region (see figures S1-3). We assumed

that all regions included in the study could be influenced by migratory birds
 195 that form part of the African and European flyways [41].

2.1. Aggregation

All data were aggregated annually to produce the final yearly panel data-set and aggregated at the NUTS 3 country subdivision level, apart from economic data which was sourced at the country level (central government). The Nomenclature of territorial units for statistics (NUTS) is a classification system used
 200 to divide economic territories of the EU into three hierarchical sub categories for the purpose of data collection and and statistical analysis:

- NUTS 1: Major socio-economic regions with a population ranging from 3 to 7 million.
- 205 • NUTS 2: Basic regions generally used for the application of regional policies with a population ranging from 800,000 to 3 million.
- NUTS 3: Small regions for specific diagnoses with a population ranging from 150,000 to 800,000.

For further details see <https://ec.europa.eu/eurostat/web/nuts/background>.

210 2.1.1. WNF Case Data

WNV case data were provided at request by the European Centre for Disease Prevention and Control (www.ecdc.europa.eu). Case data are collected weekly by EU member states and affiliates. Data were aggregated at NUTS 3 country subdivisions [42]. Positive cases were confirmed by at least one of the following
 215 techniques: 1) isolating WNV or WNV nucleic acid from blood or cerebrospinal fluid (CSF); 2) inducing a WNV-specific antibody response (either IgG / IgM) in a serological test. All cases were aggregated yearly to create the annual panel data-set.

2.2. Economic, Socio-Economic and Demographic

220 Economic data were extracted from the Eurostat database (<https://ec.europa.eu/eurostat/data/database>), which provides comparable statistics and indicators and is presented in yearly time series. To capture factors determining the economic crisis, austerity and cuts to public spending we selected NUTS3 regional level Gross Domestic Product (GDP); and country level agri-
225 culture, forestry, fisheries spending, waste water spending Health spending. The “Agriculture, forestry, fisheries spending” variable captures spending in rural areas that help to improve the environment and agricultural development, that can benefit agricultural workers and/or mechanize production [43]. In order to represent spending before and after the economic crisis, we created a baseline
230 index for each variable set at 2007 levels, which represented negative or positive growth from the point just before the economic crisis hit Europe.

Population count data to predict the number of people at risk in a region were sourced from the Socio-economic Data and Applications Center’s Gridded Population of the World data set [44]. This data-set estimates population count
235 for the years 2000, 2005, 2010, 2015, and 2020, consistent with national censuses and population registers. R’s Zoo package was used to replace values for missing years, by implementing a linear interpolation method that would predict trends between years. This way, increases or decreases in human population were controlled for in the final model.

240 2.3. Climate Data

Climate data were sourced from the E-OBS Gridded Data-set [45]. The data-set was created using a series of daily climate observations at meteorological stations throughout Europe and the Mediterranean. Files were processed in R with the Tidyverse, NetCDF and Raster packages to create regional seasonal
245 variables: “Mean temp winter (°C)”, “Mean temp spring (°C)”, “Mean temp summer (°C)”, “Winter precipitation (total mm per region)”, “Spring precipitation (total mm per region)”, “summer precipitation (total mm per region)”. Although, to standardize the data across regions we created variables which

counted rain occurrence per season i.e. “Days of rain in winter, “Days of rain
 250 in spring” and ”Days of rain in summer”. Winter was designated as December
 to March, Spring as March to June, and summer June to September.

2.4. Land-use data

Land use statistics were captured at the NUTS3 level using the CORINE
 Land Cover (CLC) 2006, 2012 and 2018 data-sets [46]. These data-sets provide
 255 information on the biophysical characteristics of the Earth’s surface in the form
 of categorical raster data. For each region, we calculated percentage land cover
 for each of the land-use risk factors identified in our conceptual framework
 i.e., “Continuous urban fabric”, “Discontinuous urban fabric’, “Wetlands (fresh
 water)” and “Arable land”. R’s SF and Raster packages were used to extract
 260 information for each available year (2006, 2012, 2018). R’s Zoo package was
 used to calculate values for missing years, by implementing a linear interpolation
 method that would predict trends between years, apart from 2019 where 2018
 values were used.

2.5. Surface Water data

265 Regional surface water data was sourced using the JRC Monthly Water His-
 tory, v1.2 data set [47] via Google Earth Engine at a 30 meter pixel resolution.
 This data set contains maps of the location and temporal distribution of surface
 water from 1984 to 2019 and provides statistics on the extent and change of wa-
 ter surfaces. Data were generated using scenes from Landsat 5, 7, and 8. Each
 270 pixel was individually classified into water / non-water using an expert system
 and the results were collated into a monthly history. Surface water observations
 were standardized by region and season by converting them to Z-scores. This
 would help determine if the seasonal water extent was average, below the mean
 (low), or above the mean (high) for a given year.

275 2.6. Statistical Methods

2.6.1. General additive regression model to assess associations of independent variables on WNV case data at regional level

One of the main issues with our data-set is that it did not meet some basic assumptions for statistical inference, and specifically the data are not independent and identically distributed random variables (iid). More specifically, the data-set captured repeated measurements over the same regions, and observations were not independent because of spill over effects from neighboring regions, therefore we needed to implement an appropriate statistical design to control for both temporal and spatial pseudo replication (lack of independence). We could deal with this in two ways, 1) either using a generalized linear mixed model (GLMM) approach, relaxing the assumption of independence and estimating the spatial/temporal correlation between residuals, or 2) model the spatial and temporal dependence in the systematic part of the model [48]. We opted to use a Generalized Additive Model (GAM) using R's MgcV statistical package because of its versatility and ability to fit complex models that would converge even with low numbers of observations and could capture potential complex non-linear relationships. One of the advantages of GAMs is that we do not need to determine the functional form of the relationship beforehand. In general, such models transform the mean response to an additive form so that additive components are smooth functions (e.g., splines) of the covariates, in which functions themselves are expressed as basis-function expansions. The spatial auto-correlation in the GAM model was approximated by a Markov random field (MRF) smoother, defined by the geographic areas and their neighborhood structure. We used R's Spdep package to create a queen neighbors list (adjacency matrix) based on regions with contiguous boundaries i.e. those sharing one or more boundary point. We used a full rank MRF, which represented roughly one coefficient for each area. The local Markov property assumes that a region is conditionally independent of all other regions unless regions share a boundary. This feature allowed us to model the correlation between geographi-

cal neighbors and smooth over contiguous spatial areas, summarizing the trend
of the response variable as a function of the predictors see section 5.4.2 of [49].
In order to account for variation in the response variable over time, not at-
tributed to the other explanatory variables in our model, we used a saturated
time effect for years, where a separate effect per time point is estimated.

We first tried to fit our model using a Poisson distribution. However, the
mean of our dependent variable (dengue cases by region and year) was lower
than its variance - $E(Y) < \text{Var}(Y)$, suggesting that the data are over-dispersed.
We also tried to fit our models using the negative binomial, quasi-Poisson and
Tweedie distribution, all particularly suited when the variance is much larger
than the mean. After several tests, we concluded that the Tweedie distribution
worked well with our data since it can handle excess zeros [50], and allows us
to model the incident rate, although results were comparative across all dis-
tributions (note that WNV infection count data, offset by a log of population
at risk was used for the neg bin and quasi-Poisson models). Analysis of model
diagnostic tests did not reveal any major issues; in general residuals appeared to
be randomly distributed (see additional information - figures S8-S15 and Table
S1 for diagnostics).

Tweedie distributions are defined as subfamily of (reproductive) exponential
dispersion models (ED), with a special mean-variance relationship . A random
variable Y is Tweedie distributed if:

$TW_p(\mu, \sigma^2)$ if $Y \sim ED(\mu, \sigma^2)$, with mean $= \mu = E(Y)$, positive dispersion
parameter σ^2 and $\text{Var}(Y) = \mu\sigma^2$.

The empirical model can then be written as:

$$E(Y) = f_1(X_{it}) + f_n(\text{Year}_t) + f_m(\text{Region}_i)$$

Where the $f(\cdot)$ stands for smooth functions; $E(Y)_{it}$ is equal to the WNV
infection incidence per 100,000 in region i at time t , which we assume to be
Tweedie distributed; X_{it} - is a vector of economic, demographic, environmental

and climate variables. $Year_t$ is a function of the time intercept and $Region_i$ represents neighborhood structure of region.

3. Results

Table 1 provides descriptive statistics of the final data-set. We did not
 335 include “Mean temp spring (°C)” in our final data set as it was correlated with
 winter and summer temperature variables, we concluded that we would capture
 more variation using the winter and summer variables which were not highly
 correlated. Public health spending was also not included in the final analysis
 as it was highly correlated with GDP (see Additional file 1 for data and model
 340 diagnostics)

To carry out our empirical analysis, we compiled a spatial-temporal data-
 set that captures some of the main drivers and risk factors of WNV infections
 in Europe. Figure 1 characterizes the climate in the study regions. As we
 can observe, WNV infections occurred in regions with climates that can be
 345 described as “Hot-summer Mediterranean”, “Humid subtropical”, “Tempera-
 ture oceanic” and “Warm-summer humid continental” or “Temperate oceanic”.
 Figure 2 shows the WNV infection incidence rates over the study period. From
 2007 to 2009, very few regions were affected by WNV, however, 2010 saw an
 outbreak that spread far and wide. Since 2010, the number of regions that have
 350 been affected by WNV increased. In 2018, a massive outbreak affected almost
 all of the regions in our study.

The relationship between the incidence of WNV infections (per 100,000)
 and the climate, land-use, and economic factors was modeled via a Generalized
 355 Additive Model (GAM), which also accounted for the spatial and temporal
 auto-correlation. Table 2 shows the results of our statistical analysis. We built
 the statistical model in a stepwise fashion using the lowest Akaike Information
 Criterion (AIC) to help us assess the different specifications. The AIC allows
 us to measure model performance accounting for model complexity and reflects

360 how well the model fits the data.

We selected relevant variables in each specification according to their category, i.e. climate, land-use and economic. All variables were included in the final specification to ascertain the contribution of each driver, all else equal. Table 2 also summarizes the relevant statistics (AIC, BIC and Deviance explained and
365 so on) to compare the different specifications. We find that our final model (Full model) has the best fit in terms of the AIC, followed by the economic model, the climate model and land use model, as shown in Table 2.

Note that as we are not estimating a standard regression model, the figures reported should not be read as coefficients, but degrees of freedom of the smooth
370 terms. Given that we cannot interpret the coefficients to infer the sign and magnitude of the relationship, we visualize it by plot. Figures (3-5) plot the partial effects—the relationship between a change in each of the covariates and a change in the fitted values in the full model.

All climate variables in both model specifications are highly statistically
375 significant (“Clim model” & “Full model”). Mean summer temperatures of above 22°C are (Figure 3, top left) positively associated with WNV infections, the relationship is linear and strong.

Mean winter temperature has a quadratic relationship with WNV. Temperatures of between 2 °C and 6°C have a positive association with WNV infections,
380 whereas colder and warmer temperatures outside of this range are negatively associated with WNV infections. The number of rain days per summer is also a strong predictor of WNV infections and has a linear positive relationship. Surface water in the summer is negatively correlated with WNV infections. The relationship is fairly strong considering its complexity i.e. the variable com-
385 plexity has been reduced to standard deviation scores to standardize it across regions and seasons.

As for the land use variables, the percentage of arable land and wetlands in a region (“Land-use model”) is positively correlated with the incidence of WNV and highly significant. However, the wetlands variable loses significance in the
390 final model. The percentage of “Continuous” and “Discontinuous urban fabric”

variables, which represents metropolitan and built up areas (residential suburbs, villages) are not statistically significant in any of the specifications. Although the partial effect plot for “Continuous urban fabric” reveals that it has slightly negative relationship with WNV infections and “Discontinuous urban fabric” has a positive association with WNV infections.

The economic indicators are negatively correlated with WNV infections. These variables represent 2007 baseline regional GDP and central government spending growth. Regional GDP has a gentle negative association with WNV infections, but loses significance in the final model. The two indicators that directly represent central government spending on areas of the environment, such as agriculture, forest & fisheries spending and waste water management, have very strong negative associations with WNV infections.

4. Discussion

To help us explain the rise in WNV infections in Europe over the last 13 years (2007-2019). We compiled a unique spatial-temporal data-set that merged a range of regional abiotic factors associated with human WNV infections, along with a set of economic indicators that captured the economic crisis. This allowed for the identification of changes, in both environmental and economic factors that possibly led to the recent rise in infections over the past decade or so.

4.1. Meteorological factors

Over the past 70 years, the countries analyzed in this study have been experiencing increasingly warmer temperatures throughout the year (see Additional information: figures S4-7). According to our initial analysis, the last decade has been the warmest. The results of our final model (Figures 3-5) show that average summer temperatures above 22 °C are positively associated with an increase in WNV incidence. It is likely that rising temperatures have created more favorable conditions for the virus and this is why it is occurring in more northerly regions than in previous decades. Higher temperature, especially in summer months,

is a key driver of WNV transmission, since warmer temperatures influence the
420 hatching rate and development time of mosquitoes, and shorten the extrinsic
incubation period (EIP) of WNV and related viruses [51, 52, 53, 54, 55, 26].

Our analysis of the mean winter temperature (Dec-Feb) reveals a quadratic
relationship with WNV and is one of the strongest predictors of annual WNV
infections. Temperatures below 2 °C and above 6 °C have a negative association
425 on WNV infections. These results are consistent with findings by Koenraadt et
al., 2019 [56]. The authors looked at *Culex pipiens* overwintering potential in
mosquitoes in sheds and houses. Temperatures in sheds ranged from 2.7–17.0°C,
and in houses 8.8–23.7°C. Survival rates were much better in sheds that had
cooler temperatures. Diapausing *Culex pipiens* mosquitoes do not necessarily do
430 better under warmer conditions and there is a temperature range in which they
can successfully diapause. Although our results may be slightly less accurate
than results derived through an experimental study (since we analyzed infection
data at a regional scale using mean monthly temperatures), as with the study by
Koenraadt et al. [56], the results do adhere to the same quadratic relationship.
435 This can be interpreted as WNV infections for the months following winter are
higher, if the specific temperature range of between 2 °C and to 6 °C is met.
This is likely because WNV can overwinter with the mosquitoes when conditions
are right, and then can be transmitted early on in the year, and throughout
the year. This result, taken in the context of recent winter warming due to
440 climate change, may explain why the virus has become so prevalent in Europe.
Given current trends, we can also expect to see regions that have previously
been too cold for *Culex pipiens* to survive over winter become viable locations
and cause further havoc in regions that are currently experiencing just a few
annual cases. On the contrary, regions which currently have optimal conditions
445 for overwintering mosquitoes may become too warm. It is also important to
note that it is not currently clear at what temperatures *Culex modestus* and
Coquillettidia richiardii overwinter as adults, since our literature search didn't
yield any findings.

The number of rain days per summer is also positively correlated with WNV

infections. This result is consistent with the literature, i.e., a steady flow of aquatic resources for mosquitoes has a positive association on their abundance, and therefore an increase in disease transmission [28, 27]. Rainfall patterns have also been shifting since the 1950s (see Additional information: figures S4-7) although unlike climate, there is no a clear trend and results are more difficult to interpret. In general, Austria, Croatia and Italy are seeing less intense rainfall in the summer months, but higher in autumn, whereas Bulgaria, Greece, Hungary and Romania are receiving more rainfall in summer months.

Our results also show that higher regional summer surface water extent for a given year, is negatively correlated with WNV and is one of the strongest predictors of WNV incidence. This was not an expected finding, since we would expected higher levels of surface water to be positively correlated with WNV incidence because of the extra water resources available to mosquitoes. However, the result is consistent with the literature as sometimes desiccation of water resources can bring mosquito and bird hosts closer together, increasing transmission potential and therefore the prevalence of the virus. [18, 14]. This was also a major finding in a recent study by Paull et al., 2017 [57], who reported that drought was closely linked to the intensity of outbreaks for a given year in the United States. It may be the case that this phenomenon is also acting at a macro-scale in Europe and is a significant driver of recent outbreaks, especially given that meteorological and hydrological droughts are becoming more frequent and extreme [58]. Another explanation for this result is that, with higher surface water extent may be associated with flooding and fast water movement, which may wash away mosquito eggs and larvae, and also may inhibit contact between birds, mosquitoes and humans [18, 14]. It is important to note that this variable probably does not capture the creation of short-term water resources created by rainfall (e.g. pools, puddles), which can be used as breeding habitat by mosquitoes. It rather captures long term and large water surface such as deltas, lakes and flood plains.

4.1.1. *Land-use*

480 As for the land-use variables, as expected, regions with a larger proportion of arable land (including rice paddies and irrigated agriculture) and wetlands are associated with WNV incidence. This is consistent with the literature as such land-use patterns attract bird hosts and mosquito species capable of transmitting WNV, which can then have knock-on effects on the human population. 485 The percentage of discontinuous urban fabric, that represents populated areas of low to medium density such as residential suburbs, villages etc. [59] is positively associated with WNV incidence, which is consistent with the literature, however, this variable is not statistically significant.

4.1.2. *Economic-factors*

490 In terms of economic factors associated with WNV infections, lower GDP growth, lower spending growth on environmental factors such as agriculture, forest, fisheries, and waste water management are positively associated with WNV incidence. Although GDP is a very broad measure, it does act as a proxy, albeit a weak one for population health and well-being as laid out in the 495 conceptual framework. It may be the case that populations living in locations that were harder hit were more exposed to mosquitoes for reasons outlined in the conceptual framework i.e. direct and indirect interventions i.e., drops in income make it difficult to afford mosquito repellents, air conditioning and upkeep of homes leading to the creation of mosquito habitat. Waste water management 500 spending may be a reliable causative factor, due to, for instance, the neglect of hazard prevention efforts, such as spending on flood defenses, essential works like sanitation and upkeep of infrastructure, that can lead to the creation of mosquito breeding habitats, e.g. potholes, blocked drains [14, 18, 39]. Lack of central government spending in this area may have led to a general degradation 505 of infrastructure and benefited mosquito species. Agriculture, forest + fisheries sectors, which represents investment or subsidies, that can go into improving environmental care, landscapes and biodiversity in rural locations, as well as investments in agricultural technology that improve aspects of the environment,

worker safety or mechanize agriculture, also had a negative association with
 510 WNV infections. Many studies report strong associations between agriculture
 [60, 29, 61, 62] and WNV incidence, in general, cuts and lower spending in
 this sector may have led to degradation on farms and the wider environment
 which may have benefited mosquitoes through the creation of habitat or lack of
 measures to control their abundance. Although it is difficult to lay out actual
 515 mechanisms when dealing with data at this scale, the overall take home message
 here is lower annual growth in the areas i.e. economic slowdown and austerity,
 is positively associated with WNV incidence.

4.1.3. Conclusions

In this study we set out to investigate why WNV outbreaks have become
 520 so frequent in Europe over the past decade. Our findings suggest that this
 phenomenon occurred because of a combination of environmental changes: 1)
 Rising winter temperatures, or rather the creation of optimal temperature con-
 ditions allowed the virus to overwinter with *Culex pipiens*; 2) Warmer summer
 temperatures are benefiting mosquitoes, influencing their hatching rate and de-
 525 velopment time, and shortening the extrinsic incubation period (EIP) of WNV;
 and 3) Shrinking water resources are increasing in WNV prevalence in birds
 and mosquitoes during some seasons. These changes also occurred during an
 economic crisis and subsequent austerity, where government institutions were
 severely weakened and had to limit spending on key sectors, and segments of
 530 the human population were exposed to increased financial hardship.

4.1.4. Limitations

Some of the limitations of the study are as follows. Since we used data-driven
 approaches to investigate how aspects of the economy, environment and climate
 drive WNV incidence at a macro-scale, we were limited to using aggregated data
 535 at the NUTS-3 regional level. With any analysis dealing with population-level
 areal data, issues can arise due to individual heterogeneity, which may lead to
 confounding bias, and we are not able to draw causal inference due to the likely

presence of endogeneity bias (e.g., omitted variable bias or reverse causality). Therefore, we carefully evaluated results from this study with individual-level
540 and clinical based studies to draw conclusions. We would have also liked to include further explanatory variables on avian host and mosquito abundance but were restricted by the availability of data.

It is also important to note that data quality issues arise owing to the under-reporting of cases i.e., through under-diagnosis, lack of diagnostic tests and a
545 lack of resources/time to carry out and implement mass testing. Another factor we did not consider is bird immunity, which may influence WNV incidence following a major outbreak, although this was not considered an important factor in explaining the rise in WNV infections in Europe, but may have influenced the results. Furthermore, a growing body of literature reports that mammals can
550 serve as intermediate hosts for West Nile virus [16] and more research needs to be done on to determine if wild mammals act as reservoirs and contribute significantly to the transmission cycle. We also realize that the economic analysis is slightly limited, in part because of a lack of refined data and in part because scale issues i.e. the amount of work required to look at individual local level
555 policies and spending was no feasible for 166 regions. Nevertheless, we hope this study would spur further research into this topic i.e., the impacts of the economic crisis on health; and the long term trade-offs and unintended consequences austerity can they have on the environment and human health. This is an especially important topic when considering we are facing multiple threats
560 brought about by global warming and other anthropogenic induced changes that can benefit emerging diseases, i.e., global trade in wild animals, intensive agriculture / animal rearing and land use conversion.

References

References

- 565 [1] P. J. Hotez, Southern europe's coming plagues: Vector-borne neglected tropical diseases, PLoS Negl Trop Dis 10 (6) (2016) e0004243. doi:10.

1371/journal.pntd.0004243.

URL <https://www.ncbi.nlm.nih.gov/pubmed/27362640>

- [2] O. Olesen, M. Ackermann, Increasing european support for neglected infectious disease research, Computational and Structural Biotechnology Journal 15 (2017) 180–184. doi:10.1016/j.csbj.2017.01.007.

URL <http://dx.doi.org/10.1016/j.csbj.2017.01.007>

- [3] G. Calleri, A. Angheben, M. Albonico, Neglected tropical diseases in europe: rare diseases and orphan drugs?, Infection 47 (1) (2018) 3–5. doi:10.1007/s15010-018-1241-2.

URL <http://dx.doi.org/10.1007/s15010-018-1241-2>

- [4] J. A. Patz, P. R. Epstein, T. A. Burke, J. M. Balbus, Global climate change and emerging infectious diseases, JAMA 275.

- [5] M. Bouzid, F. J. Colón-González, T. Lung, I. R. Lake, P. R. Hunter, Climate change and the emergence of vector-borne diseases in europe: case study of dengue fever, BMC Public Health 14 (1) (2014) 781. doi:10.1186/1471-2458-14-781.

URL <https://doi.org/10.1186/1471-2458-14-781>

- [6] W. J. Tabachnick, Climate change and the arboviruses: Lessons from the evolution of the dengue and yellow fever viruses, Annual review of Virologydoi:10.1146/annurev-virology-110615-035630.

URL <GotoISI>://WOS:000389586100006<https://www.annualreviews.org/doi/pdf/10.1146/annurev-virology-110615-035630>

- [7] C. N. Mweya, S. I. Kimera, G. Stanley, G. Misinzo, L. E. G. Mboera, Climate change influences potential distribution of infected aedes aegypti co-occurrence with dengue epidemics risk areas in tanzania, Plos One 11 (9). doi:10.1371/journal.pone.0162649.

URL <GotoISI>://WOS:000384171400014<http://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0162649&type=printable>

- [8] R. Shope, Global climate change and infectious diseases, *Environ Health Perspect* 96. doi:10.1289/ehp.9196171.
- [9] A. Morrison, H. Astete, C. Rocha, V. Lopez, J. Olson, T. Kochel, M. Si-huincha, J. Stancil, Impact of the dengue vector control system (dvcs) on
600 aedes aegypti populations in iquitos, peru 2004-2005, *American Journal of Tropical Medicine and Hygiene* 73 (6) (2005) 326–326.
URL <GotoISI>://WOS:000202990001415
- [10] J. A. Patz, W. J. Martens, D. A. Focks, T. H. Jetten, Dengue fever epi-
demic potential as projected by general circulation models of global climate
605 change, *Environ Health Perspect* 106. doi:10.1289/ehp.98106147.
URL <https://doi.org/10.1289/ehp.98106147><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1533051/pdf/envhper00526-0073.pdf>
- [11] S. Naish, P. Dale, J. S. Mackenzie, J. McBride, K. Mengersen, S. L. Tong, Climate change and dengue: a critical and systematic review
610 of quantitative modelling approaches, *Bmc Infectious Diseases* 14. doi:10.1186/1471-2334-14-167.
URL <GotoISI>://WOS:000333603300001<https://bmcinfectdis.biomedcentral.com/track/pdf/10.1186/1471-2334-14-167>
- [12] D. J. Gubler, Dengue, urbanization and globalization: The unholy trinity
615 of the 21(st) century, *Tropical Medicine and Health* 39 (4 Suppl) (2011) 3–11. doi:10.2149/tmh.2011-S05.
URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3317603/>
- [13] K. Magori, J. M. Drake. The population dynamics of vector-borne diseases [online] (2013).
- [14] S. Paz, J. C. Semenza, Environmental drivers of west nile fever epidemi-
620 ology in europe and western asia—a review, *International journal of environmental research and public health* 10 (8) (2013) 3543–3562. doi:10.3390/ijerph10083543.

- URL <https://pubmed.ncbi.nlm.nih.gov/23939389><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3774453/>
- [15] ECDC. Factsheet about west nile virus infection [online] (2018).
- [16] J. Root, A. Bosco-Lauth, West nile virus associations in wild mammals: An update, *Viruses* 11 (5) (2019) 459. doi:10.3390/v11050459.
URL <http://dx.doi.org/10.3390/v11050459>
- [17] W. J. Landesman, B. F. Allan, R. B. Langerhans, T. M. Knight, J. M. Chase, Inter-annual associations between precipitation and human incidence of west nile virus in the united states, *Vector-Borne and Zoonotic Diseases* 7 (3) (2007) 337–343. doi:10.1089/vbz.2006.0590.
URL <https://doi.org/10.1089/vbz.2006.0590>
- [18] Z. Hubálek, J. Halouzka, West nile fever a reemerging mosquito borne viral disease in europe, *Emerging infectious diseases* 5 (5) (1999) 643–650. doi:10.3201/eid0505.990505.
- [19] J. Young, J. Haussig, S. Aberle, D. Pervanidou, F. Riccardo, N. Sekulić, T. Bakonyi, C. Gossner, Epidemiology of human west nile virus infections in the european union and european union enlargement countries, 2010 to 2018, *Eurosurveillance* 26 (19). doi:10.2807/1560-7917.es.2021.26.19.2001095.
URL <http://dx.doi.org/10.2807/1560-7917.ES.2021.26.19.2001095>
- [20] M. Karanikolos, P. Mladovsky, J. Cylus, S. Thomson, S. Basu, D. Stuckler, J. P. Mackenbach, M. McKee, Financial crisis, austerity, and health in europe, *Lancet* 381 (9874) (2013) 1323–31. doi:10.1016/s0140-6736(13)60102-6.
- [21] A. Baumbach, G. Gulis, Impact of financial crisis on selected health outcomes in europe, *The European Journal of Public Health* 24 (3) (2014) 399–403. doi:10.1093/eurpub/cku042.
URL <http://dx.doi.org/10.1093/eurpub/cku042>

- [22] S. Quinn, S. Kumar, Health inequalities and infectious disease epidemics: A challenge for global health security, *Biosecurity and Bioterrorism: Biodefense Strategy, Practice, and Science* 12 (5) (2014) 263–273. doi:10.1089/bsp.2014.0032.
655 URL <http://dx.doi.org/10.1089/bsp.2014.0032>
- [23] B. Durand, A. Tran, G. Balança, V. Chevalier, Geographic variations of the bird-borne structural risk of west nile virus circulation in europe, *PLOS ONE* 12 (10) (2017) e0185962. doi:10.1371/journal.pone.0185962.
660 URL <http://dx.doi.org/10.1371/journal.pone.0185962>
- [24] ECDC. West nile virus infection [online] (2020).
- [25] R. P. Meyer, J. L. Hardy, W. K. Reisen, Diel changes in adult mosquito microhabitat temperatures and their relationship to the extrinsic incubation of arboviruses in mosquitoes in kern county, california, *J Med Entomol* 27 (4) (1990) 607–14. doi:10.1093/jmedent/27.4.607.
665
- [26] W. K. Reisen, Y. Fang, V. M. Martinez, Effects of temperature on the transmission of west nile virus by culex tarsalis (diptera: Culicidae), *J Med Entomol* 43 (2) (2006) 309–17. doi:10.1603/0022-2585(2006)043[0309:Eotott]2.0.Co;2.
- [27] J. E. Soverow, G. A. Wellenius, D. N. Fisman, M. A. Mittleman, Infectious disease in a warming world: how weather influenced west nile virus in the united states (2001-2005), *Environ Health Perspect* 117 (7) (2009) 1049–52. doi:10.1289/ehp.0800487.
670
- [28] T. Takeda, C. A. Whitehouse, M. Brewer, A. D. Gettman, T. N. Mather, Arbovirus surveillance in rhode island: assessing potential ecologic and climatic correlates, *J Am Mosq Control Assoc* 19 (3) (2003) 179–89.
675
- [29] M. C. Gates, R. C. Boston, Irrigation linked to a greater incidence of human and veterinary west nile virus cases in the united states from 2004 to 2006,

Prev Vet Med 89 (1-2) (2009) 134–7. doi:10.1016/j.prevetmed.2008.
12.004.

[30] J. M. Brunkard, J. L. R. Lopez, J. Ramirez, E. Cifuentes, S. J. Rothenberg, E. A. Hunsperger, C. G. Moore, R. M. Brussolo, N. A. Villarreal, B. M. Haddad, Dengue fever seroprevalence and risk factors, texas-mexico border, 2004, *Emerging Infectious Diseases* 13 (10) (2007) 1477–1483. doi:10.3201/eid1310.061586.
URL <GotoISI>://WOS:000249962800006https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2851499/pdf/06-1586_finalR.pdf

[31] J. P. DeGroote, R. Sugumaran, National and regional associations between human west nile virus incidence and demographic, landscape, and land use conditions in the coterminous united states, *Vector Borne Zoonotic Dis* 12 (8) (2012) 657–65. doi:10.1089/vbz.2011.0786.

[32] J. Tackett, R. Charnigo, G. Caldwell, Relating west nile virus case fatality rates to demographic and surveillance variables, *Public health reports* (Washington, D.C. : 1974) 121 (6) (2006) 666–673. doi:10.1177/003335490612100606.
URL <https://pubmed.ncbi.nlm.nih.gov/17278401><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1781908/>

[33] Z. Dowling, S. L. Ladeau, P. Armbruster, D. Biehler, P. T. Leisnham, Socioeconomic status affects mosquito (diptera: Culicidae) larval habitat type availability and infestation level, *J Med Entomol* 50 (4) (2013) 764–72. doi:10.1603/me12250.

[34] I. Unlu, A. Farajollahi, D. Strickman, D. M. Fonseca, Crouching tiger, hidden trouble: urban sources of aedes albopictus (diptera: Culicidae) refractory to source-reduction, *PloS one* 8 (10) (2013) e77999–e77999. doi:10.1371/journal.pone.0077999.
URL <https://pubmed.ncbi.nlm.nih.gov/24167593><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3805523/>

- [35] W. Reisen, R. Takahashi, B. Carroll, R. Quiring, Delinquent mortgages, neglected swimming pools, and west nile virus, california, *Emerging Infectious Diseases* 14 (11) (2008) 1747–1749. doi:10.3201/eid1411.080719. URL <http://dx.doi.org/10.3201/eid1411.080719>
- [36] M. Kim, J. Holt, R. Eisen, K. Padgett, W. Reisen, J. Croft, Detection of swimming pools by geographic object-based image analysis to support west nile virus control efforts, *Photogrammetric Engineering and Remote Sensing* 77 (11) (2011) 1169–1179. doi:10.14358/pers.77.11.1169. URL <http://dx.doi.org/10.14358/PERS.77.11.1169>
- [37] W. Reisen, B. Carroll, R. Takahashi, Y. Fang, S. Garcia, V. Martinez, R. Quiring, Repeated west nile virus epidemic transmission in kern county, california, 2004–2007, *Journal of Medical Entomology* 46 (1) (2009) 139–157. doi:10.1603/033.046.0118. URL <http://dx.doi.org/10.1603/033.046.0118>
- [38] E. Kondilis, S. Giannakopoulos, M. Gavana, I. Ierodiakonou, H. Waitzkin, A. Benos, Economic crisis, restrictive policies, and the population’s health and health care: the greek case, *American journal of public health* 103 (6) (2013) 973–979. doi:10.2105/AJPH.2012.301126. URL <https://pubmed.ncbi.nlm.nih.gov/23597358><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3698730/>
- [39] R. Wallace, L. F. Chaves, L. R. Bergmann, C. Ayres, L. Hogerwerf, R. Kock, R. G. Wallace, *Clear-cutting disease control: capital-led deforestation, public health austerity, and vector-borne infection*, Springer, 2018.
- [40] L. Steele, E. Orefuwa, P. Dickmann, Drivers of earlier infectious disease outbreak detection: a systematic literature review, *International Journal of Infectious Diseases* 53 (2016) 15–20. doi:<https://doi.org/10.1016/j.ijid.2016.10.005>. URL <http://www.sciencedirect.com/science/article/pii/S1201971216311882>

- [41] UNEP, A review of migratory bird flyways and priorities for management, CMS Technical Series.
URL https://www.cms.int/sites/default/files/publication/CMS_Flyways_Reviews_Web.pdf
- [42] Eurostat. Nuts - nomenclature of territorial units for statistics [online] (2020).
- [43] Eurostat, Agriculture, forestry and fishery statistics, Report, Eurostat (2019). doi:10.2785/798761.
URL <https://ec.europa.eu/eurostat/web/products-statistical-books/-/KS-FK-19-001>
- [44] SEDAC. Gridded population of the world, version 4 (gpwv4): Population count, revision 11 [online] (2018).
- [45] R. C. Cornes, G. van der Schrier, E. J. M. van den Besselaar, P. D. Jones, An ensemble version of the e-obs temperature and precipitation data sets, Journal of Geophysical Research: Atmospheres 123 (17) (2018) 9391–9409. doi:<https://doi.org/10.1029/2017JD028200>.
URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2017JD028200>
- [46] EU. Copernicus land monitoring service 2018 [online] (2018).
- [47] J.-F. Pekel, A. Cottam, N. Gorelick, A. S. Belward, High-resolution mapping of global surface water and its long-term changes, Nature 540 (7633) (2016) 418–422. doi:10.1038/nature20584.
URL <https://doi.org/10.1038/nature20584>
- [48] A. Aswi, S. M. Cramb, P. Moraga, K. Mengersen, Bayesian spatial and spatio-temporal approaches to modelling dengue fever: a systematic review, Epidemiology and infection 147 (2018) 1–14. doi:10.1017/S0950268818002807.

URL <https://pubmed.ncbi.nlm.nih.gov/30369335><https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6518570/>

[49] S. N. Wood, Generalized additive models: an introduction with R, CRC press, 2017.

[50] C. Kurz, Tweedie distributions for fitting semicontinuous health care utilization cost data, BMC Medical Research Methodology 17 (1). doi:10.1186/s12874-017-0445-y.

URL <http://dx.doi.org/10.1186/s12874-017-0445-y>

[51] S. Chen, J. Blanford, S. Fleischer, M. Hutchinson, M. Saunders, M. Thomas, Estimating west nile virus transmission period in pennsylvania using an optimized degree-day model, Vector-Borne and Zoonotic Diseases 13 (7) (2013) 489–497. doi:10.1089/vbz.2012.1094.

URL <http://dx.doi.org/10.1089/vbz.2012.1094>

[52] T. Kamiya, M. Greischar, K. Wadhawan, B. Gilbert, K. Paaijmans, N. Mideo, Temperature-dependent variation in the extrinsic incubation period elevates the risk of vector-borne disease emergence, Epidemics 30 (2020) 100382. doi:10.1016/j.epidem.2019.100382.

URL <http://dx.doi.org/10.1016/j.epidem.2019.100382>

[53] A. Mohammed, D. D. Chadee, Effects of different temperature regimens on the development of aedes aegypti (l.)(diptera: Culicidae) mosquitoes, Acta tropica 119 (1) (2011) 38–43.

[54] W. Tun-Lin, A. Lenhart, V. S. Nam, E. Rebollar-Tellez, A. C. Morrison, P. Barbazan, M. Cote, J. Midega, F. Sanchez, P. Manrique-Saide, A. Kroeger, M. B. Nathan, F. Meheus, M. Petzold, Reducing costs and operational constraints of dengue vector control by targeting productive breeding places: a multi-country non-inferiority cluster randomized trial, Tropical Medicine and International Health 14 (9) (2009) 1143–1153. doi:10.1111/j.1365-3156.2009.02341.x.

URL <GotoISI>://WOS:000269263800024<https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-3156.2009.02341.x>

- [55] D. M. Watts, D. S. Burke, B. A. Harrison, R. E. Whitmire, A. Nisalak, Effect of temperature on the vector efficiency of aedes aegypti for dengue 2 virus, The American journal of tropical medicine and hygiene 36 (1) (1987) 143–152.

- [56] C. J. M. Koenraadt, T. W. R. Mohlmann, N. O. Verhulst, J. Spitzen, C. B. F. Vogels, Effect of overwintering on survival and vector competence of the west nile virus vector culex pipiens, Parasites and Vectors 12 (1) (2019) 147. doi:10.1186/s13071-019-3400-4. URL <https://doi.org/10.1186/s13071-019-3400-4>

- [57] S. Paull, D. Horton, M. Ashfaq, D. Rastogi, L. Kramer, N. Diffenbaugh, A. Kilpatrick, Drought and immunity determine the intensity of west nile virus epidemics and climate change impacts, Proceedings of the Royal Society B: Biological Sciences 284 (1848) (2017) 20162078. doi:10.1098/rspb.2016.2078. URL <http://dx.doi.org/10.1098/rspb.2016.2078>

- [58] M. Hanel, O. Rakovec, Y. Markonis, P. Máca, L. Samaniego, J. Kyselý, R. Kumar, Revisiting the recent european droughts from a long-term perspective, Scientific Reports 8 (1). doi:10.1038/s41598-018-27464-4. URL <http://dx.doi.org/10.1038/s41598-018-27464-4>

- [59] ETC/ULS, Updated clc illustrated nomenclature guidelines, Report, European Environment Agency (2019). URL <https://land.copernicus.eu/user-corner/technical-library/corine-land-cover-nomenclature-guidelines/html/index-clc-112.html>

- [60] D. Crowder, E. Dykstra, J. Brauner, A. Duffy, C. Reed, E. Martin, W. Peterson, Y. Carrière, P. Dutilleul, J. Owen, West nile virus prevalence across

- 820 landscapes is mediated by local effects of agriculture on vector and host
communities, PLoS ONE 8 (1) (2013) e55006. doi:10.1371/journal.
pone.0055006.
URL <http://dx.doi.org/10.1371/journal.pone.0055006>
- [61] L. Eisen, C. Barker, C. Moore, W. Pape, A. Winters, N. Cheronis, Irrigated
825 agriculture is an important risk factor for west nile virus disease in the
hyperendemic larimer-boulder-weld area of north central colorado, Journal
of Medical Entomology 47 (5) (2010) 939–951. doi:10.1093/jmedent/47.
5.939.
URL <http://dx.doi.org/10.1093/jmedent/47.5.939>
- 830 [62] R. Miramontes, W. Lafferty, B. Lind, M. Oberle, Is agricultural activity
linked to the incidence of human west nile virus?, American Journal of
Preventive Medicine 30 (2) (2006) 160–163. doi:10.1016/j.amepre.2005.
10.008.
URL <http://dx.doi.org/10.1016/j.amepre.2005.10.008>

835 **Acknowledgments**

This research was funded by ICTA's Maria de Maeztu Unit of Excellence, awarded by the Spanish Ministry of Economy and Competitiveness. The award is the highest institutional recognition of scientific research in Spain. Thanks also to Patrizia Ziveri and Pedro Manuel Gonzalez Hernandez for supporting
840 the project. Thanks also to Cesira Urzi Brancati for statistical help and advice.

Author contributions statement

MW led the work and was responsible for the conceptualization of the project, data collection, data processing, formulation of the methodology, statistical analysis, modeling, analysis and writing the original draft and interpreting
845 the results. All authors made contributions to developing the original paper outline. PK gave critical advice throughout the project. All contributors revised the manuscript and copy-edited the final submission version. All the authors were also involved in revising the manuscript critically for important intellectual content. All authors read and approved the final manuscript. PK, PGM
850 VSM supervised the project and are responsible for formulating ideas for the umbrella project Impacts of Climate Change (CC) on Human Health (HH) at ICTA-UAB: Integrating socio-economic and policy studies with natural science studies to enhance consilience of climate policy science.

Data availability

855 An R project containing all the data that supports the findings of this study are available in .Rdata format from <https://doi.org/10.5281/zenodo.4656902>).

Code availability

An R project containing all the code used to set-up the models is available
860 here is available from <https://doi.org/10.5281/zenodo.4656902>.

5. Tables

Table 1: Final dataset - 2007-2019

Statistic	N	Min	Max	Mean	St. Dev.
WNV cases	2,158	0	100	1.649	6.012
Human population	2,158	6,254.326	10,534,640.000	443,384.400	513,000.000
Mean temp winter (C)	2,158	-6.072	14.564	3.190	3.623
Mean temp spring (C)	2,158	4.092	18.684	12.433	2.157
Mean temp summer (C)	2,158	13.109	28.011	22.465	2.285
Days of rain in winter	2,158	0	68	30.156	12.546
Days of rain in spring	2,158	0	71	31.723	12.918
Days of rain in summer	2,158	0	65	26.021	14.374
Spring surface water extent Z-score (30m2)	2,158	-2.876	2.404	0.000	0.958
Summer surface water extent Z-score (30m2)	2,158	-3.301	3.080	-0.000	0.958
Continuous urban fabric % cover	2,158	0.000	45.056	1.336	6.754
Discontinuous urban_fabric (% cover)	2,158	0.534	60.457	5.511	7.044
Wetlands (% cover)	2,158	0.000	25.460	0.569	2.026
Arable land (% cover)	2,158	0.000	86.307	33.893	22.172
Regional GDP growth (2007=100%)	2,158	57	217	106.752	24.587
Agri, forest + fish spending (2007=100%)	2,158	27	251	80.202	35.165
Waste water mngmnt spending (2007=100%)	2,158	5	352	93.476	53.933
Health spending (2007=100%)	2,158	60	212	114.802	33.307

Table 2: Final model results

	Clim model	Land-use model	Econ model	Full model
Intercept	-2.21*** (0.47)	-2.13*** (0.45)	-2.25*** (0.33)	-2.35*** (0.40)
Mean temp summer (C)	1.00*** (1.00)			1.00* (1.00)
Mean temp winter (C)	1.96*** (1.99)			1.94*** (1.99)
Days of rain in summer	1.00** (1.00)			1.00** (1.00)
Summer surface water extent (30m2)	1.60** (1.84)			1.02*** (1.03)
Continuous urban fabric %		1.00 (1.00)		1.00 (1.00)
Discontinuous urban fabric %		1.00 (1.00)		1.00 (1.00)
Wetlands %		1.00** (1.00)		1.00 (1.00)
Arable land %		1.81*** (1.89)		1.74** (1.84)
Regional GDP index (2007=100%)			1.00* (1.00)	1.00 (1.00)
Agri, forest + fish spending (2007=100%)			1.95*** (1.99)	1.93*** (1.99)
Waste water managment spending (2007=100%)			1.60*** (1.83)	1.10*** (1.19)
Spatial lag	78.44*** (109.60)	80.54*** (111.42)	88.90*** (121.08)	76.19*** (106.52)
Year	11.73*** (12.00)	11.75*** (12.00)	11.48*** (12.00)	11.56*** (12.00)
AIC	3952.99	3992.59	3931.25	3907.56
BIC	4526.68	4569.26	4560.01	4538.85
Log Likelihood	-1875.44	-1894.71	-1854.87	-1842.57
Deviance	3747.15	3895.52	3592.25	3520.85
Deviance explained	0.63	0.62	0.64	0.65
Dispersion	2.85	2.95	2.76	2.73
R ²	0.26	0.36	0.32	0.25
GCV score	1900.89	1927.27	1889.38	1871.90
Num. obs.	2158	2158	2158	2158
Num. smooth terms	6	6	5	13

*** $p < 0.01$; ** $p < 0.05$; * $p < 0.1$

Figure 1: Koppen-Geiger Climate Classification in Study Regions (Data source: koeppen-geiger.vu-wien.ac.at).

Figure 2: Regional West Nile Virus distribution 2006 to 2019: Confirmed cases per 100,000 (Data source: ECDC).

6. Figures

Figure 3: Partial effects plots of main model results 1: represents the relationship between a change in each of the covariates and a change in the fitted values in the full model.

Figure 4: Partial effects plots of main model results 2: represents the relationship between a change in each of the covariates and a change in the fitted values in the full model.

Figure 5: Partial effects plots of main model results 3: represents the relationship between a change in each of the covariates and a change in the fitted values in the full model.

Köppen-Geiger Climate Classification