AtPQT11, a P450 enzyme, detoxifies paraquat via N-demethylation

Yi-Jie Huang¹, Yue-Ping Huang^{1,2}, Jin-Qiu Xia¹, Zhou-Ping Fu³, Yi-Fan Chen³,
Yi-Peng Huang³, Aimin Ma⁴, Wen-Tao Hou¹, Yu-Xing Chen¹, Xiaoquan Qi⁴, Li-Ping
Gao³, Cheng-Bin Xiang¹

¹School of Life Sciences, Division of Molecular & Cell Biophysics, Hefei National
Science Center for Physical Sciences at the Microscale, University of Science and
Technology of China, The Innovation Academy of Seed Design, Chinese Academy of
Sciences, Hefei, Anhui Province 230027, China

²School of Life Sciences, Sun Yat Sen University, No. 135, Xingang Xi Road,
Guangzhou, 510275, P. R. China

³School of Life Science, Anhui Agricultural University, 130 West Changjiang Rd,
Hefei 230036, Anhui, China

⁴ **Key Laboratory of Plant Molecular Physiology**, Institute of Botany, The Innovation
Academy of Seed Design, Chinese Academy of Sciences, Beijing, China

Correspondence:

Cheng-Bin Xiang

Email: xiangcb@ustc.edu.cn

orcid.org/0000-0002-7152-1458

Li-Ping Gao

E-mail: gaolp62@126.com

Abstract

Paraquat is one of the most widely used nonselective herbicides in agriculture. Due to its wide use, paraquat resistant weeds have emerged and is becoming a potential threat to agriculture. The molecular mechanisms of paraquat resistance in weeds remain largely unknown. Physiological studies indicated that the impaired translocation of paraquat and enhanced antioxidation could improve paraquat resistance in plants. However, the detoxification of paraquat via active metabolism by plants has not been reported to date. Here we report that an activated expression of *At1g01600* encoding the P450 protein CYP86A4 confers paraquat resistance as revealed by the gain-of-function mutant *paraquat tolerance 11D* (*pqt11D*), in which a T-DNA with four 35S enhancers was inserted at 1646 bp upstream the ATG of *At1g01600*. The paraquat resistance can be recapitulated in Arabidopsis wild type by overexpressing *AtPQT11* (*At1g01600*), while its knockout mutant is hypersensitive to paraquat. Moreover, *AtPQT11* also confers paraquat resistance in *E. coli* when overexpressed. We further demonstrate that AtPQT11 has P450 enzyme activity that converts paraquat to N-demethyl paraquat nontoxic to Arabidopsis, therefore detoxifying paraquat in plants. Taken together, our results unequivocally demonstrate that AtPQT11/ CYP86A4 detoxifies paraquat via active metabolism, thus revealing a novel molecular mechanism of paraquat resistance in plants and providing a means potentially enabling crops to resist paraquat.

Key words: paraquat, paraquat resistance, *AtPQT11*, *pqt11*, CYP86A4, P450, herbicide, demethylation.

Introduction

Paraquat (1,1'-dimethyl-4,4'-bipyridinium dichloride) has been widely used as a broad spectrum nonselective herbicide in agriculture for decades¹. Plants absorb paraquat from their environment and transport to chloroplasts, where it competes for electrons on the PSI and generates superoxide that is converted to H₂O₂ by superoxide dismutases (SODs), resultantly accumulating a large amount of ROS, which may lead to cell death².

Paraquat resistant weeds have emerged after the wide use of paraquat in agriculture. The underlying mechanisms of paraquat resistance in weeds have been studied mostly at physiological and biochemical levels³⁻⁶. From these studies, two major mechanisms of paraquat resistance were suggested, including defects in paraquat translocation restricting the transport of paraquat to chloroplasts^{4,7}, and enhanced antioxidation scavenging ROS more efficiently^{8,9}. However, no genetic or molecular evidence for the proposed mechanisms has been revealed in weeds. With the power of the model plant *Arabidopsis*, several research groups have isolated paraquat resistant mutants, originally aiming at identifying novel components in oxidative stress response or cell death^{10,11}. These studies with *Arabidopsis* paraquat resistant mutants helped reveal the molecular mechanisms of paraquat resistance proposed by weed scientists based on their studies on paraquat resistant weeds. The unraveled molecular mechanisms from these studies fall into two categories; one involves paraquat uptake and transport¹²⁻¹⁵, and the other is the enhanced ability scavenging ROS^{10,11,16,17}.

However, no evidence of paraquat degradation by plant active metabolism has been reported. Herbicide degradation by plants via active metabolism is a desirable mechanism for engineering crop resistance to herbicides. CYP450s (cytochrome P450s) constitute the biggest plant protein family and can mediate redox reaction in plant. They usually function as demethylase and/or hydroxylase in herbicide metabolism¹⁸. Some P450s metabolize herbicides to lower toxic products which are transferred to vacuole as glucosyl conjugates¹⁹. Two plant cytochrome P450 enzymes,

CYP71A10 from *Helianthus tuberosus* and CYP76B1 from soybean, were reported having urea herbicide metabolism activity²⁰. In addition, tobacco CYP81B2 and CYP71A11 catabolize chlortoluron by ring methyl hydroxylation and N-methylation²¹.

We previously isolated a number of paraquat tolerant Arabidopsis mutants and reported *pqt24* and *pqt3*^{15,16}. Here we continue to characterize those paraquat tolerance mutants by focusing on *pqt11D* in this study. We isolated *pqt11D* as a gain-of-function mutant with enhanced paraquat tolerance and identified that the *At1G01600* (*AtPQT11*) was activated by the enhancers of the T-DNA insertion, which caused paraquat resistance in the mutant. The mutant phenotype could be recapitulated by overexpressing *AtPQT11*, encoding the protein CYP86A4, a member of CYP450 super family, in the wild type Arabidopsis. Further analyses revealed that AtPQT11/CYP86A4 could convert paraquat to N-demethyl paraquat *in vitro* by the mean of N-demethylation, and demethylated paraquat was found nontoxic to Arabidopsis, clearly demonstrating paraquat degradation by a plant P450 enzyme. Our study revealed a novel molecular mechanism of paraquat resistance that has not been reported in plants. Our findings should provide a new means for developing paraquat resistance crops using *AtPQT11/CYP86A4*.

Result

Activated expression of *At1G01600* confers paraquat resistance in *pqt11D*

From an Arabidopsis activation-tagging library of approximately 55000 lines, we isolated a number of paraquat tolerance mutants. One of them, *pqt11D*, displayed enhanced resistance to paraquat (Fig. S1A). We found that the T-DNA with four 35S enhancers next to the right border was inserted at 1646 bp upstream the ATG codon of *At1g01600* (Fig. S1B). As a result, the gene *At1g01600* was activated (Fig. S1C), indicating that the activated expression of *At1g01600* may be responsible for observed paraquat resistance in *pqt11D*. We therefore named *At1g01600* as *AtPQT11*.

108 *AtPQT11* is predicted to encode CYP86A4, a cytochrome P450 enzyme with 554
109 amino acids. It is expressed in all tissues examined with relatively higher levels in
110 roots, flowers and siliques (Fig. S1D), and induced by paraquat treatment for the first
111 3 hours, then declined (Fig. 1E).

112 **The *pqt11D* phenotype can be recapitulated by overexpressing *AtPQT11* in the** 113 **wild type**

114 To confirm whether the paraquat resistance of *pqt11D* was caused by an activated
115 expression of *AtPQT11*, we generated two *AtPQT11* overexpression lines (OX1-4 and
116 OX3-1) and obtained a knockout mutant *pqt11* (Salk_073078), in which the T-DNA
117 was inserted in the first exon of *AtPQT11* from ABRC (Fig. S1B), and confirmed all
118 the genetic materials by 3-primers PCR (Fig. S1F) and qRT-PCR analyses (Fig. S1G).

119 Seed germination assay in response to paraquat treatment showed that two OX
120 lines had significantly higher survival after 7 days on MS medium with 2 μ M
121 paraquat compared with the wild type, while the knockout mutant *pqt11* became
122 hypersensitive (Fig. 1A and B).

123 We also conducted paraquat resistance assay on soil-grown plants of different
124 *AtPQT11* genotypes. When the plants grew to the stage with 8 rosette leaves, they
125 were sprayed with 10 μ M paraquat every 3 days. 10 days after the first spray, The OX
126 lines showed a significantly higher survival than wild type control, whereas the
127 knockout mutant exhibited sicker phenotype and lower survival (Fig. 1C and D).

128 Together, these results demonstrate that an elevated expression of *AtPQT11*
129 confers paraquat resistance in Arabidopsis.

130 ***AtPQT11* renders *E. coli* resistant to paraquat**

131 To further confirm *AtPQT11*-conferred paraquat resistance, we cloned *AtPQT11*
132 cDNA into pET28a vector (pET28a-P450) and expressed in *E. coli*. Based on 3-D
133 simulation of AtPQT11 structure (Fig. S2), we created single mutations of three
134 residues G309A, S461A, G463A around the predicted substrate binding pocket and
135 further expressed these three mutants in *E. coli*, respectively. In the absence of
136 paraquat in liquid culture medium, there was no difference was observed between the
137 bacteria expressing pET28a-P450 and the empty pET28a. In contrast, the bacteria

expressing pET28a-P450 displayed a remarkable resistance to 1 mM paraquat than those harboring the empty pET28a, whereas the bacteria transformed with the constructs carrying 3 different point mutations in *AtPQT11* exhibited no paraquat resistance as the empty vector control (Fig. 1E). When grew on solid agar medium containing 1 mM paraquat, only bacteria expressing pET28a-P450 formed colony plaques (Fig. 1F). These results further confirm the role of *AtPQT11* in paraquat resistance and indicate that the amino acid residues are critical for its function.

AtPQT11 catalyzes the conversion of paraquat to N-demethyl paraquat *in vitro*

Considering that AtPQT11 encodes a P450 protein, we reasoned AtPQT11 might catabolize paraquat to a nontoxic product. To demonstrate this, we first established an *in vitro* enzyme assay using *E. coli* expressed AtPQT11 protein, paraquat as substrate, and NADPH as electron donor as described in Materials and Methods. The consumption of NADPH provides a convenient spectrophotometric monitor at OD_{340nm} of the reaction. As shown in Fig. 2A, the consumption of NADPH was AtPQT11-dependent in the reaction and increased with incubation time. Therefore, we established the *in vitro* enzyme assay for AtPQT11 and used this assay to determine the enzyme kinetic properties (Fig. 2B-D).

To identify the product of the reaction, the paraquat metabolite of the *in vitro* reaction was analyzed by LC-MS/MS. A product peak was found at 2.7 min with m/z 171, which was AtPQT11- and NADPH-dependent (Fig. 3A-D, left panels). To confirm that this product is derived from paraquat, we used the ²H-labeled paraquat as the substrate, whose eight hydrogen atoms on paraquat pyridine ring were replaced by deuterium so that the m/z of paraquat changed from 186 to 194. As expected, we found the product peak with m/z 179, which was also AtPQT11- and NADPH-dependent (Fig. 3A-D, right panels). The reaction product has the same mass as N-demethyl paraquat (m/z 171).

To resolve the identity of the product, we analyzed the MS/MS spectrum and found that the product showed the same fragment ions as N-demethyl paraquat standard (Fig. 3F, left panel and G, right panel), which was further confirmed by the same experiments with ²H-labeled paraquat (Fig. 3E and F, right panels). These

168 results unequivocally show that the product is derived from the substrate and the
169 product matches N-demethyl paraquat based on their fragment ion pattern (Fig. 3G).
170 Thus, we have demonstrated that AtPQT11/ CYP86A4 functions as an N-demethylase
171 capable of converting paraquat to N-demethyl paraquat.

172 **N-demethyl paraquat is nontoxic to Arabidopsis**

173 Have identified N-demethyl paraquat as the product, we decided to find out
174 whether N-demethyl paraquat has herbicide activity. To demonstrate this, we simply
175 tested whether the N-demethyl paraquat is toxic to Arabidopsis. Seed germination
176 assay in the presence of N-demethyl paraquat showed that two OX lines, WT and
177 knockout mutant *pqt11* all survived and grew well on MS medium with 2 μ M
178 N-demethyl paraquat or 2 μ M bipyridine (double demethylated paraquat) compared
179 with paraquat (Fig. 4A and B). This result clear shows that N-demethyl paraquat and
180 bipyridine are nontoxic to Arabidopsis. Therefore, we have uncovered the molecular
181 mechanism underlying the paraquat resistance of *pqt11D*.

182

183 **Discussion**

184 In this study we reported the paraquat resistance mutant *pqt11D* and the
185 underlying molecular mechanism. The phenotype of the mutant was caused by an
186 activated expression of *AtPQT11*, which was confirmed by recapitulation analyses in
187 Arabidopsis wild type as well as in *E. coli* (Fig. 1 and Fig. S1A-C). The paraquat
188 hypersensitive phenotype of the *AtPQT11* knockout mutant also supports that
189 *AtPQT11* is responsible for the observed paraquat resistance (Fig. 1 and 4). Further
190 analyses revealed the AtPQT11 catalyzed biochemical reaction converting paraquat to
191 single demethylated product (Fig. 2 and 3), and we also found that N-demethyl
192 paraquat is nontoxic to Arabidopsis (Fig. 4).

193 CYP450 enzymes is widely found in biosynthesis or catabolic processes
194 through the whole life in all living organisms²². Several members of CYP450 family
195 have been reported to detoxify herbicides^{23,24}. As a xenobiotic, paraquat is not a
196 natural substrate of the CYP450 enzyme AtPQT11. The natural substrates of

AtPQT11 remain unknown at present. AtPQT11 can catalyze the conversion of paraquat to N-demethyl paraquat perhaps because paraquat may have similar structural resemblance to its natural substrates.

Since paraquat is a widely used herbicide in agriculture, paraquat resistant weeds have emerged, which is a potential threat to agriculture. Generally, plants could gain paraquat resistance in three ways. First, paraquat absorption and transport are limited, thus preventing paraquat from reaching chloroplasts. Second, an enhanced intracellular antioxidant capacity would help plants to scavenge ROS more efficiently. Third, paraquat may be degraded to nontoxic metabolites. Plants may have these strategies in single or in combination to produce paraquat resistance²⁵. There are many reports for the first two strategies^{4,10,13,15,16}. However, paraquat degradation by plants has not been reported thus far. Our finding with *pqt11D* fills up the gap for the third strategy.

In conclusion, our findings with the paraquat resistant mutant *pqt11D* unequivocally demonstrate that AtPQT11, a member of the P450 super family detoxifies paraquat and confers paraquat resistance in Arabidopsis when overexpressed, therefore providing a much needed means to engineering paraquat resistance crops.

Methods and materials

Plant material and growth conditions

Arabidopsis Col-0 was used as wild type in this study. We also used Col-0 as the genetic background for all the mutants and transgenic plants. Salk_073078 was ordered from the Arabidopsis Biological Resource Center (ABRC). Seeds were sterilized in 10% bleach for 10 minutes and washed with sterile water 5 times. The washed seeds were kept in the dark at 4°C for 2 days before germination on MS medium at 22°C under 14 h light/10 h dark.

Construction of *AtPQT11* overexpression lines

In order to get overexpression lines, we used forward primer 5'-GGGGACAAGTTTGTACAAAAAAGCAGGCT ATGGAAATATCCAATGCCATGC-3 and reverse primer 5'-GGGGACCACTTTGTACAAGAAAGCTGGGT TTAAACCACTGCAACTCCCGTA-3' to obtain the CDS of CYP86A4. By means of GATEWAY system²⁶, the CDS was cloned into vector pCB2004 via the shuttle vector of pDONR207.

The constructed plasmid was transformed into *Agrobacterium tumefaciens* C58C1 by electroporation. The floral-dip method was used to transfer the construct into plants²⁷. The positive lines were screened as glufosinate resistant plants and homozygous lines were obtained from F2 population.

RNA extraction and qRT-PCR

We extracted the RNA from one-week-old seedlings by Trizol (Invitrogen, Carlsbad, California, USA) and reverse transcribed RNA to cDNA by TransScript Kit (TaKaRa). The expression of CYP86A4 was detected by RT-PCR or quantitative RT-PCR using forward primer 5'-CCCCAAGGGTTTCACTGAATTC-3' and reverse primer 5'-AAGTAAATGCGAAGCCTGCTTG -3' and Applied Biosystem Step One real-time PCR system or TaKaRa SYBR Premix Ex Taq II reagent kit. *UBQ5* was used as the internal reference.

Identification of the *AtPQT11* knockout mutant

3 primers including forward primer 5'- TTCACCACATACAGCTGCATC -3', reverse primer 5'- AAATGTTGTCGAATGTGAGCC-3' and intermediate primer LBb1.3 5'-ATTTTGCCGATTTTCGGAAC-3' were used to identify homozygotes of SALK_073078. The homozygotes showed only one band around 750 bp while the wild type showed one band around 1000 bp. RT-PCR was performed to confirm null expression of the *Atlg01600* in SALK_073078 by using forward primer 5'- ATGGAAATATCCAATGCCATGC-3' and reverse primer 5'- TTAAACCACTGCAACTCCCGTA-3'. *TUBULIN* was used as the internal control.

254 **Seed germination assay**

255 Sterilized seeds for each line including WT, knock out mutant SALK_073078
256 (*pqt11/cyp86a4*) and two overexpression lines OX1-4 and OX3-1 were treated at 4°C
257 in dark condition for 2 days. Then we put these seeds on MS medium with 2 μM
258 paraquat, MS medium was used as control. The plates were placed at 22°C under 14
259 h light/10 h dark cycles. We considered the appearance of two green cotyledons as
260 successful germination, 1 week later we recorded germination %.

261 **Paraquat resistance assay on soil-grown plants**

262 One-week-old seedlings grown on MS medium were transferred to soil. We
263 planted 5 seedlings per pot, grew to the stage with 8 rosette leaves, and sprayed with
264 10 μM paraquat and sprayed again 3 days later. Photos and survival rate were
265 recorded 10 days after the first spray.

266 **3-D model of At PQT11 structure**

267 The 3-D structure model of CYP86A4 was produced by Phyre2 and paraquat
268 was docked into the structure model. Residues within the distance of 4 Å around
269 paraquat are colored in red. Conservative residues likely responsible for the binding
270 of paraquat are labeled and chosen as mutation sites.

271 **Site-directed mutagenesis of *AtPQT11***

272 Specific primers for site-directed mutagenesis were designed, for changing
273 309Gly to Ala, 461Cys to Ala, 463Gly to Ala as listed in Table S1. pET28-CYP86A4
274 or pCB2004-CYP86A4 were set as template and the PCR procedure was set for 20
275 cycles. After the PCR products were recycled, 1 μl DpnI was added to digest these
276 products. Then transferred these products to DH5α and extracted plasmids from
277 overnight culture.

278 **Paraquat resistance assay in *E. coli***

279 We used Rosetta or BL21 as host, transferred the constructed pET28-P450 and
280 the three plasmids with point mutation: pET28-P450A (Gly309Ala), pET28-P450B
281 (Cys461Ala) and pET28-P450C (Gly463Ala) into the host bacteria, and empty vector
282 pET28a as control. Single colonies grown on the plate were picked and cultured

overnight. Then the overnight culture was diluted into fresh LB medium with a starting OD₆₀₀ value of 0.1, and paraquat was added to make its final concentration of 1 mM. Another set of culture with 0 mM paraquat was used as control. The OD₆₀₀ value of the culture was monitored at regular intervals for 36 hours. Each culture was triplicated and the growth curves were obtained. When tested on solid LB plate with 1 mM paraquat, the overnight liquid bacteria culture was inoculated on the plate with 0 or 1 mM paraquat and incubated overnight at 37°C.

Bacterial expression of AtPQT11

The CDS of *AtPQT11* was ligated to the vector pET28a by T4 ligase. The constructed plasmid was transformed into Rosetta strain with kanamycin selection. A clone was picked from medium and was added with liquid LB, when the OD₆₀₀ of medium reached 0.6, IPTG was added with a final concentration of 0.3 mM. The culture was shaken in 16°C with the speed of 120 rpm for 12-16 h. The culture was centrifuged for 10 min at 4500 rpm to harvest bacteria, then the bacterial pellet was resuspended with protein isolation buffer (20 mM Tris-Cl pH7.5, 0.2 M NaCl, 5% glycerin, 1 mM EDTA, 1 mM PMSF) + 1% Dodecyl-beta-D-maltoside (m/v), and ultrasonicated to break bacterial cell walls. The cell lysate was rotated at 4°C for 2h and then centrifuged at 45000 rpm for 30 min. The supernatant was collected for AtPQT11 protein purification using Ni-IDA-sefinose resin kit.

***In vitro* enzyme assay of AtPQT11**

The reaction mixtures contained 0.1-10 µM paraquat as substrate, 50-100 µg AtPQT11/ CYP86A4 protein, 1 mM NADPH as electron donor, 25 mM phosphate buffer (pH7.0) in a total volume of 50 µl. The reaction was initiated by adding NADPH. The reaction mixtures without NADPH or AtPQT11/CYP86A4 were set as controls. The mixtures were incubated for 30 min at 27°C and terminated by adding 50 µl acetonitrile. To determine the enzyme kinetic properties, the reaction was spectrophotometrically monitored at 340 nm for NADPH consumption. To identify the reaction product, the mixtures was centrifuged at 14000g for 20 min, and the

311 supernatant was analyzed by HPLC-MS/MS as described below.

312 **HPLC-MS/MS**

313 HPLC-MS/MS analysis was performed using Agilent Q-TOF-LC/MS 6545
314 device (Agilent Technologies, Palo Alto, CA, USA) . A XBridge BEH HILIC 2.5 μ m
315 Column (2.1*100 mm) from Waters was used for chromatographic separation. The
316 mobile phase was a mixture of solvent A (10 mM ammonium acetate added with 0.1%
317 V/V formic acid) and solvent B (10 mM ammonium acetate dissolved in 95%
318 acetonitrile) and was delivered at a flow rate of 0.3 mL/min. The column temperature
319 was set at 40°C and the injection volume was 2 μ l. The sample was followed the
320 gradient: 90% B (0 min), 10% B (3 min), 1% B (5 min), 90% B (5 min), with a total
321 run time of 12 min.

322 Mass acquisition was performed in the positive ionization mode at a
323 fragmentation voltage of 175 V. The following parameters were used: drying gas flow,
324 8 L/min; temperature, 325 °C; sheath gas flow, 11 L/min; temperature, 350 °C;
325 nebulizer pressure, 45 psi; and capillary voltage, 3500 V. The collision energy was set
326 at 15–40 V, and the mass range was recorded from m/z 50–1700. The acquired MS
327 and MS/MS data files were analyzed using Agilent MassHunter Qualitative Analysis
328 B.07.00.

329 **Toxicity assay of N-demethyl paraquat**

330 Seeds of different genotypes were washed and treated at 4°C in dark condition
331 for 2 days, then plated on MS medium with 2 μ M N-demethyl paraquat, 2 μ M
332 bipyridine, 2 μ M paraquat, respectively. MS medium was used as mock. Germination
333 and survival was recorded after 1 week.

334 **Statistical analyses**

335 Statistical significance was evaluated at the 0.05 probability level using Student's
336 t-test.

337 **Supplemental information**

338 Figure S1. Paraquat resistance phenotype of *pqt11D* and confirmation of

339 *AtPQT11* expression in *pqt11*, *pqt11D*, and overexpression lines.

340 Fig. S2. 3-D model of AtPQT11 structure.

341 Table S1. Primers used in this study.

342 **Acknowledgements**

343 This work was supported by grants from the National Natural Science Foundation
344 of China (grant no. 31770273). We thank ABRC for providing Arabidopsis mutant
345 seeds.

346 **Author's Contributions**

347 CBX and YJH designed the experiments. YJH. performed the major experiments.
348 YPH isolated *pqt11D* and identified the T-DNA insertion site. JQX, WTH, AMM,
349 XQQ, YXC, participated in experiments and data analyses. ZPF, YFC, YiPH, LPG
350 contributed to LC-MS/MS analyses. YJH wrote the manuscript. CBX and LPG
351 revised the manuscript and supervised the project.

352 **Conflict Interests**

353 The authors declare no conflict of interest.

354

355

References

- 1 Baldwin, B., Clarke, C. & Wilson, I. Paraquat in chloroplasts. *Biochimica et Biophysica Acta (BBA)-Bioenergetics* **162**, 614-617 (1968).
- 2 Farrington, J., Ebert, M., Land, E. & Fletcher, K. Pulse radiolysis studies of the reaction of paraquat radical with oxygen. Implication for the mode of action of bipuridyl herbicides. *Biochimica et Biophysica Acta* **314**, 372-381 (1973).
- 3 Lewthwaite, S. & Triggs, C. Identification of paraquat-resistant *Solanum nigrum* and *S. americanum* biotypes. *New Zealand Plant Protection* **62**, 349-355 (2009).
- 4 Jóri, B. *et al.* Role of transporters in paraquat resistance of horseweed *Conyza canadensis* (L.) Cronq. *Pesticide Biochemistry and Physiology* **88**, 57-65, doi:10.1016/j.pestbp.2006.08.013 (2007).
- 5 Lasat, M. M., DiTomaso, J. M., Hart, J. J. & Kochian, L. V. Evidence for vacuolar sequestration of paraquat in roots of a paraquat-resistant *Hordeum glaucum* biotype. *Physiologia Plantarum* **99**, 255-262 (1997).
- 6 Hawkes, T. R. Mechanisms of resistance to paraquat in plants. *Pest Manag Sci* **70**, 1316-1323, doi:10.1002/ps.3699 (2014).
- 7 Shaaltiel, Y. & Gressel, J. Kinetic analysis of resistance to paraquat in *Conyza*: evidence that paraquat transiently inhibits leaf chloroplast reactions in resistant plants. *Plant physiology* **85**, 869-871 (1987).
- 8 Ye, B. & Gressel, J. Transient, oxidant-induced antioxidant transcript and enzyme levels correlate with greater oxidant-resistance in paraquat-resistant *Conyza bonariensis*. *Planta* **211**, 50-61 (2000).
- 9 Tsuji, K., Hosokawa, M., Morita, S., Miura, R. & Tominaga, T. Resistance to paraquat in *M. azu pumilus*. *Weed Research* **53**, 176-182 (2013).
- 10 Chen, R. *et al.* The Arabidopsis PARAQUAT RESISTANT2 gene encodes an S-nitrosoglutathione reductase that is a key regulator of cell death. *Cell Research* **19**, 1377-1387, doi:10.1038/cr.2009.117 (2009).
- 11 Kurepa, J., Smalle, J., Va, M., Montagu, N. & Inzé, D. Oxidative stress tolerance and longevity in Arabidopsis: the late-flowering mutant *gigantea* is tolerant to paraquat. *The Plant Journal* **14**, 759-764 (1998).
- 12 Fujita, M. *et al.* Natural variation in a polyamine transporter determines paraquat tolerance in Arabidopsis. *Proceedings of the national academy of sciences of the united states of america* **109**, 6343-6347, doi:10.1073/pnas.1121406109 (2012).
- 13 Li, J. *et al.* Paraquat Resistant1, a Golgi-localized putative transporter protein, is involved in intracellular transport of paraquat. *Plant Physiology* **162**, 470-483, doi:10.1104/pp.113.213892 (2013).
- 14 Soar, C., Preston, C., Karotam, J. & Powles, S. Polyamines can inhibit paraquat toxicity and translocation in the broadleaf weed *Arctotheca calendula*. *Pesticide Biochemistry and Physiology* **80**, 94-105 (2004).

- 15 Xi, J., Xu, P. & Xiang, C. B. Loss of AtPDR11, a plasma membrane-localized ABC transporter, confers paraquat tolerance in *Arabidopsis thaliana*. *Plant Journal* **69**, 782-791, doi:10.1111/j.1365-3113X.2011.04830.x (2012).
- 16 Luo, C. *et al.* PARAQUAT TOLERANCE3 Is an E3 Ligase That Switches off Activated Oxidative Response by Targeting Histone-Modifying PROTEIN METHYLTRANSFERASE4b. *PLOS Genetics* **12** (2016).
- 17 Sipari, N., Lihavainen, J., Shapiguzov, A., Kangasjärvi, J. & Keinänen, M. Primary Metabolite Responses to Oxidative Stress in Early-Senescing and Paraquat Resistant *Arabidopsis thaliana* rcd1 (Radical-Induced Cell Death1). *Frontiers in plant science* **11**, 194 (2020).
- 18 Schuler, M. A. & Werck-Reichhart, D. Functional genomics of P450s. *Annual review of plant biology* **54**, 629-667 (2003).
- 19 Kreuz, K., Tommasini, R. & Martinoia, E. Old enzymes for a new job (herbicide detoxification in plants). *Plant physiology* **111**, 349 (1996).
- 20 Didierjean, L. *et al.* Engineering herbicide metabolism in tobacco and *Arabidopsis* with CYP76B1, a cytochrome P450 enzyme from Jerusalem artichoke. *Plant Physiology* **130**, 179-189 (2002).
- 21 Yamada, T., Kambara, Y., Imaishi, H. & Ohkawa, H. Molecular Cloning of Novel Cytochrome P450 Species Induced by Chemical Treatments in Cultured Tobacco Cells. *Pesticide Biochemistry and Physiology* **68**, 11-25 (2000).
- 22 Mansuy, D. The great diversity of reactions catalyzed by cytochromes P450. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol* **121**, 5-14, doi:10.1016/s0742-8413(98)10026-9 (1998).
- 23 Xiang, W., Wang, X. & Ren, T. Expression of a wheat cytochrome P450 monooxygenase cDNA in yeast catalyzes the metabolism of sulfonylurea herbicides. *Pesticide biochemistry and physiology* **85**, 1-6 (2006).
- 24 Siminszky, B., Corbin, F. T., Ward, E., Fleischmann, T. J. & Dewey, R. E. Expression of a soybean cytochrome P450 monooxygenase cDNA in yeast and tobacco enhances the metabolism of phenylurea herbicides. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 1750-1755 (1999).
- 25 Baucom, R. S. The remarkable repeated evolution of herbicide resistance. *Am J Bot* **103**, 181-183, doi:10.3732/ajb.1500510 (2016).
- 26 Lei, Z. *et al.* High-throughput Binary Vectors for Plant Gene Function Analysis. *Journal of Integrative Plant Biology* **49**, 556-567 (2007).
- 27 Clough, S. J. & Bent, A. F. Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* **16**, 735-743 (1998).

AtPQT11 by YJ Huang et al.

437

438

439

Figure Legends

Figure 1. *AtPQT11* confers paraquat resistance in *Arabidopsis* and *E. coli* when overexpressed.

(A) The seeds of wild type (WT), *AtPQT11* overexpression lines (OX1-4, OX3-1), and *pqt11* were germinated on MS medium with 0 or 2 μ M paraquat (PQ) for 1 week before survival was recorded. 100 seeds were used for each genotype. 3 replicates were performed for each treatment. Bar = 1 cm.

(B) Survival % in germination as in (A). Values are mean $\pm$ SD (n=3). The low case letters indicate significant differences ($P < 0.05$).

(C) Phenotypes of wild type (WT), *AtPQT11* overexpression lines (OX1-4, OX3-1), and *pqt11* grown in the soil. When plants grew to 8-rosette-leaf stage, they were sprayed with 0 (control) or 10 μ M paraquat and sprayed again 3 days later. Photos and survival % were recorded 10 days after the first spray. 40 plants were used for each genotype. 3 replicates were performed for each treatment.

(D) Survival % as in (C). Values are mean $\pm$ SD (n=3). The low case letters indicate significant differences ($P < 0.05$).

(E) Bacterial growth curves. Plasmids pET28a-*AtPQT11* (pET28a-P450) with different changed residue: pET28-P450A (Gly309Ala), pET28-P450B (Cys461Ala) and pET28-P450C (Gly463Ala) were transferred into Rosetta, and cultured in LB medium with 0 or 1 mM paraquat. The growth of the cultures with initial OD₆₀₀ adjusted to 0.1 was monitored for 36 hours. 3 replicates were performed for each treatment. Values are mean $\pm$ SD (n=3).

(F) Bacterial growth on solid agar plates. Bacterial strains as in (E) were inoculated on solid LB medium with 0 or 1 mM paraquat and incubated overnight at 37°C.

Figure 2. *In vitro* kinetic parameters of *AtPQT11*

(A) *AtPQT11*-dependent reaction. The enzyme assay was carried out as described in Materials and Methods with 2 controls, no *AtPQT11* (CYP684A) and inactivated *AtPQT11*. NADPH consumption was only observed in the reaction containing

470 AtPQT11. Values are mean $\pm$ SD (n=3).
 471 (B) Michaelis-Menton plot for paraquat. 2 mM NADPH was added to ensure it is in
 472 excess in the reaction while varying the concentration of paraquat. Values are mean
 473 $\pm$ SD (n=3).
 474 (C) Michaelis-Menton plot for NADPH. 2 mM Paraquat was added to ensure it is in
 475 excess in the reaction while varying the concentration of NADPH. Values are mean
 476 $\pm$ SD (n=3).
 477 (D) The enzyme kinetics parameters. V_{max} and K_m were obtained from (B) and (C), K_m
 478 and V_{max} values were estimated by using non-linear regression analysis.

479

480 **Figure 3. Identification of the reaction product by LC-MS/MS.**

481 (A-D) The reaction was carried out with paraquat or ^2H -paraquat as substrate as
 482 described in Materials and Methods for 30 min. The reaction supernatant was
 483 subjected to LC-MS/MS analysis by scanning product at M/Z 171 or 179. (A) and (B)
 484 are the reaction for 0 or 30 min, (C) and (D) are the reaction for 30 min but without
 485 NADPH or P450 enzyme.
 486 (E-G) The MS/MS spectrum of substrate and product. (E) The MS/MS spectrum of
 487 paraquat when the reaction was just started. (F) The MS/MS spectrum of product
 488 when the reaction was completed (30 min). (G) The MS/MS spectrum of paraquat
 489 standard and N-demethyl paraquat standard. MS/MS spectrum of the product closely
 490 matches with that of N-demethyl paraquat (M/Z 171).

491

492 **Figure 4. N-demethyl paraquat is nontoxic to Arabidopsis.**

493 (A) Germination assay. The seeds of wild type (WT), *AtPQT11* overexpression lines
 494 (OX1-4, OX3-1), and *pqt11* were germinated on MS medium with 0 (Mock), 2 μM
 495 paraquat (PQ), 2 μM N-demethyl paraquat (N-methyl PQ), or 2 μM bipyridine,
 496 respectively for 1 week before survival was recorded. 100 seeds were used for each
 497 genotype. 3 replicates were performed for each treatment. Bar = 1 cm.
 498 (B) Survival % in germination as in (A). Values are mean $\pm$ SD (n=3). The low case
 499 letters indicate significant differences ($P < 0.05$).

Figure 1. *AtPQT11* confers paraquat resistance in *Arabidopsis* and *E. coli* when overexpressed.

(A) The seeds of wild type (WT), *AtPQT11* overexpression lines (OX1-4, OX3-1), and *pqt11* were germinated on MS medium with 0 or 2 μ M paraquat (PQ) for 1 week before survival was recorded. 100 seeds were used for each genotype. 3 replicates were performed for each treatment. Bar = 1 cm.

(B) Survival % in germination as in (A). Values are mean $\pm$ SD (n=3). The low case letters indicate significant differences ($P < 0.05$).

(C) Phenotypes of wild type (WT), *AtPQT11* overexpression lines (OX1-4, OX3-1), and *pqt11* grown in the soil. When plants grew to 8-rosette-leaf stage, they were sprayed with 0 (control) or 10 μ M paraquat and sprayed again 3 days later. Photos and survival % were recorded 10 days after the first spray. 40 plants were used for each genotype. 3 replicates were performed for each treatment.

(D) Survival % as in (C). Values are mean $\pm$ SD (n=3). The low case letters indicate significant differences ($P < 0.05$).

(E) Bacterial growth curves. Plasmids pET28a-*AtPQT11* (pET28a-P450) with different changed residue (pET28a-P450A, pET28a-P450B, pET28a-P450C) were transferred into Rosetta, and cultured in LB medium with 0 or 1 mM paraquat. The growth of the cultures with initial OD₆₀₀ adjusted to 0.1 was monitored for 36 hours. 3 replicates were performed for each treatment. Values are mean $\pm$ SD (n=3).

(F) Bacterial growth on solid agar plates. Bacterial strains as in (E) were inoculated on solid LB medium with 0 or 1 mM paraquat and incubated overnight at 37°C.

Figure 2. *In vitro* kinetic parameters of AtPQT11

(A) AtPQT11-dependent reaction. The enzyme assay was carried out as described in Materials and Methods with 2 controls, no AtPQT11 (CYP684A) and inactivated AtPQT11. NADPH consumption was only observed in the reaction containing AtPQT11. Values are mean $\pm$ SD (n=3).

(B) Michaelis-Menton plot for paraquat. 2 mM NADPH was added to ensure it is in excess in the reaction while varying the concentration of paraquat. Values are mean $\pm$ SD (n=3).

(C) Michaelis-Menton plot for NADPH. 2 mM Paraquat was added to ensure it is in excess in the reaction while varying the concentration of NADPH. Values are mean $\pm$ SD (n=3).

(D) The enzyme kinetics parameters. V_{max} and K_m were obtained from (B) and (C), K_m and V_{max} values were estimated by using non-linear regression analysis.

Figure 3. Identification of the reaction product by LC-MS/MS.

(A-D) The reaction was carried out with paraquat or ^2H -paraquat as substrate as described in Materials and Methods for 30 min. The reaction supernatant was subjected to LC-MS/MS analysis by scanning product at M/Z 171 or 179. (A) and (B) are the reaction for 0 or 30 min, (C) and (D) are the reaction for 30 min but without NADPH or P450 enzyme.

(E-G) The MS/MS spectrum of substrate and product. (E) The MS/MS spectrum of paraquat when the reaction was just started. (F) The MS/MS spectrum of product when the reaction was completed (30 min). (G) The MS/MS spectrum of paraquat standard and N-demethyl paraquat standard. MS/MS spectrum of the product closely matches with that of N-demethyl paraquat (M/Z 171).

A

WT OX1-4 OX3-1 *pqt11*

bioRxiv preprint doi: <https://doi.org/10.1101/2021.06.23.449549>; this version posted June 23, 2021. The copyright holder for this preprint (which was not certified by peer review) is the author/funder. All rights reserved. No reuse allowed without permission.

Mock

PQ

**N-demethyl
PQ**

bipyridine

B

Figure 4. N-demethyl paraquat is nontoxic to Arabidopsis.

(A) Germination assay. The seeds of wild type (WT), *AtPQT11* overexpression lines (OX1-4, OX3-1), and *pqt11* were germinated on MS medium with 0 (Mock), 2 μ M paraquat (PQ), 2 μ M N-demethyl paraquat (N-methyl PQ), or 2 μ M bipyridine, respectively for 1 week before survival was recorded. 100 seeds were used for each genotype. 3 replicates were performed for each treatment. Bar = 1 cm.

(B) Survival % in germination as in (A). Values are mean $\pm$ SD (n=3). The low case letters indicate significant differences ($P < 0.05$).