Danish dementia and excitatory transmission

▪ Familial Danish dementia young Knock-in rats expressing humanized APP and human A β show impaired pre and postsynaptic glutamatergic transmission

▪ Tao Yin¹, Wen Yao^{1,2}, Kelly A. Norris¹ & Luciano D'Adamio^{1*}

¹Department of Pharmacology, Physiology & Neuroscience, Brain Health Institute, New Jersey Medical School, Rutgers, The State University of New Jersey, 205 South Orange Ave, Newark, NJ, 07103, USA

²Present address: Biomedical Research Center, Southern Medical University, 1023 South Shatai Road, Guangzhou, Guangdong, 510515, China.

*To whom correspondence should be addressed: luciano.dadamio@rutgers.edu

Running title: Danish dementia and excitatory transmission

Keywords: Familial Danish Dementia, Amyloid precursor protein (APP), Amyloid β , neurodegeneration, synaptic plasticity, rat, animal model, glutamate, Familial Danish Dementia, BRI2, *Integral membrane protein 2B (ITM2b)*

ABSTRACT

Familial British and Danish dementia (FBD and FDD) are two neurodegenerative disorders caused by mutations in the *Integral membrane protein 2B (ITM2b)*. BRI2, the protein encoded by *ITM2b*, tunes excitatory synaptic transmission at both pre- and post-synaptic terminus. Too, BRI2 interacts with and modulates proteolytic processing of Amyloid- β precursor Protein (APP), whose mutations cause familial forms of Alzheimer disease (FAD). To study pathogenic mechanism triggered by the Danish mutation we generated rats carrying the Danish mutation into the rat *Itm2b* gene (*Itm2b^D* rats). Given the BRI2/APP interaction and the widely accepted relevance of human A β , a proteolytic product of APP, to AD, *Itm2b^D* rats were engineered to express two humanized *App* alleles, to produce human A β . Here, we studied young *Itm2b^D* rats to investigate early pathogenic changes. We found that peri-adolescent *Itm2b^D* rats present subtle changes in human A β levels along with decreased spontaneous glutamate release and AMPAR-mediated responses but increased short-term

synaptic facilitation in the hippocampal Schaeffer-collateral pathway. These changes are like those observed in adult mice producing rodent A β and carrying either the Danish or British mutations into the mouse *Itm2b* gene. Collectively, the data show that the pathogenic Danish mutation alters the physiological function of BRI2 at glutamatergic synapses; these functional alterations are detected across species and occur early in life. Future studies will be needed to determine whether this phenomenon represents an early pathogenic event in human dementia.

INTRODUCTION

Model organisms that reproduce the pathogenesis of human diseases are useful to dissect disease mechanisms, identify therapeutic targets and test therapeutic strategies. Because genetic manipulation has been easier in mice, mice have overtaken rats as the major rodent-based model organism in neurodegeneration research. Thus, to study FDD and FBD, fifteen years ago we generated mice carrying the pathogenic Danish and British dementia mutations (*Itm2b^D* and *Itm2b^B*

Danish dementia and excitatory transmission

mice) into the *Itm2b* mouse gene (1-3). We choose a knock in (KI) approach rather than the more common transgenic overexpression approach for several reasons. KIs mimic the genetic of FDD and FBD and make no assumption about pathogenic mechanisms (except the unbiased genetic one), while the transgenic approach aims to reproduce pathology (plaques, Neurofibrillary tangles (NFTs), etc.), under the assumption that this “pathology” is pathogenic. In KI models, expression of mutant genes is controlled by endogenous regulatory elements, avoiding issues related to over-expression of disease-proteins in a non-physiological quantitative-spatial-temporal manner. Finally, potential confounding “insertion” effects of transgenes are avoided.

Because rats are better suited to study neurodegenerative diseases, we took advantage of recent developments in gene-editing technologies and introduced the familial Danish mutation into the genomic *Itm2b* rat locus (*Itm2b^D* rats). The rat was the organism of choice for most behavioral, memory and cognitive research -which is critical when studying neurodegenerative diseases- because physiological processes are similar in rats and humans and the rat is an intelligent and quick learner (4-7).

Several procedures that are important in dementia research are more easily performed in rats as compared to mice due to the larger size of the rat brain. Cannulas -to administer drugs, biologics, viruses etc.- and microdialysis probes -for sampling extracellular brain levels of neurotransmitters, A β , soluble tau etc.- can be accurately directed to individual brain regions, causing less damage and increasing specificity. *In vivo* brain imaging techniques, such as MRI (8) and PET (9-11), can assess the extent and course of neurodegeneration with better spatial resolution in rats. Moreover, rats are large enough for convenient *in vivo* electrophysiological recordings or serial sampling of cerebrospinal fluid for detection of biomarkers.

Finally, gene-expression differences suggest that rats may be advantageous model

of neurodegenerative diseases over mice. For example, alternative splicing of *tau* (12-15), which forms NFTs and is mutated in Frontotemporal Dementia (16-23), leads to expression of tau isoforms with three or four microtubule binding domains (3R and 4R, respectively). Adult human and rat brains express both 3R and 4R tau isoforms (24): in contrast, adult mouse brains express only 4R tau(25), suggesting that the rat may be a better model organism for dementias with tauopathy, such as FDD and FBD.

BRI2 physically interacts with and modulates processing of APP, which bears relevance to AD pathogenesis (26-30). In addition, APP processing mediates LTP and memory deficits of Danish and British KI mice (31-36). Aggregated forms of A β , a product of APP processing, are by and large considered the main pathogenic molecule in AD. Rat and human APP differ by 3 amino-acids in the A β region: given that human A β are believed to have higher propensity to form toxic A β species as compared to rodent A β , we produced rats carrying the humanized A β sequence (*App^h* rats) (37,38). Thus, to study possible interactions between the Danish mutation and human A β , *Itm2b^D* rats were backcrossed to *App^h* rats. Hence, all rats used in this study produce human and not rodent A β species.

Here, we studied peri-adolescent *Itm2b^D* rats, with the purpose of investigating early dysfunctions that may underlie initial pathogenic mechanisms leading to dementia later in life.

RESULTS

Generation of *Itm2b^D* KI rats carrying humanized *App^h* alleles.

The knock-in founder F0-*Itm2b^D* rat, which is carrying FDD mutation on *Itm2b* rat gene, was generated by CRISPR/Cas-mediated genome engineering as described in Experimental Procedures and Supporting Information. The F0-*Itm2b^D* rat, which is a chimera for the *Itm2b* gene, was crossed to WT (*Itm2b^{w/w}*) Long-Evans rats to generate F1-*Itm2b^{D/w}* rats. F1-*Itm2b^{D/w}* rats were crossed to WT Long-Evans to generate F2-*Itm2b^{D/w}* rats. These crossing were repeated three more

Danish dementia and excitatory transmission

times to obtain F5-*Itm2b*^{D/w} rats. The probability that F5 rats carry unidentified off-target mutations (except those, if present, on Chr. 15) is ~1.5625%. Male and female F5-*Itm2b*^{D/w} rats were crossed to obtain *Itm2b*^{D/w}, *Itm2b*^{D/D} and *Itm2b*^{w/w} rats.

The FDD mutation consist of a 10 nucleotides duplication one codon before the normal stop codon (39). This produces a frameshift in the BRI2 sequence generating a precursor protein 11 amino acids larger-than-normal (Figure 1A). To verify that the Danish mutation was correctly inserted into *Itm2b* exon 6, we amplified by PCR the *Itm2b* gene exon 6 *Itm2b*^{D/w}, *Itm2b*^{D/D} and *Itm2b*^{w/w} rats. Sequencing of the PCR products shows that the Danish mutation was correctly inserted in the *Itm2b* gene exon 6 (Figure 1B) and encoded for the COOH-terminus of the Danish BRI2 mutant. When we generated FDD KI mice, we humanized the mouse COOH-terminal region of BRI2 by introducing an alanine (A) was substituted for threonine (T) at codon 250 (3). Since that humanization did not result into deposition of ADan peptides in amyloid plaques in KI mice (3), that modification was not repeated in rats.

To generate *Itm2b*^{D/w}, *Itm2b*^{D/D} and *Itm2b*^{w/w} rats on a background in which rat *App* has a humanized Aβ region, *Itm2b*^{D/w} and *App*^{h/h} rats were crossed to generate *Itm2b*^{D/w}; *App*^{h/h} rats. The *App*^w allele was removed in subsequent crosses. Henceforth, *Itm2b*^{D/D}, *Itm2b*^{D/w} and *Itm2b*^{w/w} rats used in this study have an *App*^{h/h} background and produce human and not rodent Aβ species.

To determine whether *Itm2b* expression is disrupted by the introduced mutations, we examined *Itm2b* mRNA levels in p21 *Itm2b*^{D/D} and *Itm2b*^{w/w} rats by standard RNA-Seq analysis on total brain RNA. The mRNA expression of *Itm2b* shows no significant difference between *Itm2b*^{D/D} and *Itm2b*^{w/w} rats (Figure 1C).

The *Itm2b*^D allele encodes for a longer Bri2 precursor protein (Bri2-ADan) that accumulates in primary neurons.

BRI2 is a type II membrane protein that is synthesized as an immature precursor (imBRI2). imBRI2 is cleaved at the COOH-

terminus by proprotein convertase to produce the NH₂-terminal mature BRI2 (mBRI2) and the 23 amino acid-long COOH-terminal peptide called Bri23 (40). As noted above, in the Danish patients, a frameshift caused by a 10 nucleotides duplication 5' to the stop codon leads to the synthesis of a BRI2 precursor protein 11 amino acids larger-than-normal (39). Convertase-mediated cleavage of immature Danish BRI2 generates a WT-like mBRI2 and a 34 amino acid long peptide called ADan, which co-deposits with Aβ species in amyloid fibrils in patients. For clarity, we will refer to the wild type imBri2 as Bri2-Bri23, and to the Danish mutant imBri2 as Bri2-ADan.

To determine whether the *Itm2b*^D allele codes for Bri2-ADan we examined Bri2 expression in total neuronal lysates isolated from male and female 2 months old *Itm2b*^{D/w}, *Itm2b*^{D/D} and *Itm2b*^{w/w} rats. However, the Bri2 antibody tested identified many non-specific bands (Figure S1), making a rigorous assessment of Bri2 expression in rat brains challenging.

Analysis of mouse *Itm2b*^{w/w} and *Itm2b*^{D/D} primary neurons showed that the mBri2/Bri2-Bri23 ratio in *Itm2b*^{w/w} primary neurons was significantly higher than the mBri2/Bri2-ADan ratio in *Itm2b*^{D/D} primary neurons (41). In addition, lysosomal inhibition caused accumulation of mBri2 but not Bri2-Bri23 in *Itm2b*^{w/w} primary neurons; in contrast, both mBri2 and Bri2-ADan accumulated in *Itm2b*^{D/D} primary neurons (41). These observations indicated that the Danish mutation reduced maturation of the mutant precursor Bri2 in mouse neurons. Based on these observations, we probed whether primary neurons could be used to assess mBri2, Bri2-Bri23 and Bri2-ADan expression in KI rats. Primary neurons are a simpler system compared to total brain; this, *per se*, may reduce the number of non-specific bands identified by anti-Bri2 antibodies. Moreover, inhibition of lysosome-mediated degradation of Bri2 species in primary neurons may help identify specific Bri2 molecules. Thus, primary neurons derived from *Itm2b*^{w/w} and *Itm2b*^{D/D} rat were treated with the lysosomal inhibitor chloroquine and analyzed by Western

Danish dementia and excitatory transmission

blot. The anti-Bri2 antibody identified a band of ~34 kDa in all samples, which was increased by chloroquine (Figure 2A and 2C). These observations are consistent with the ~34 kDa corresponding to mBri2. A second band of ~36 kDa was detected in *Itm2b^{w/w}* primary neurons (Figure 2A). In contrast, a slightly larger second band (~37 kDa) that was increased by chloroquine treatment, was detected in *Itm2b^{D/D}* primary neurons (Figure 2A and 2C). These observations are consistent with the ~36 kDa and ~37 kDa bands corresponding to Bri2-Bri23 and Bri2-ADan, respectively.

Without treatment, the levels of Bri2-ADan in *Itm2b^{D/D}* primary neurons were significantly higher than the levels of Bri2-Bri23 in *Itm2b^{w/w}* primary neurons (Figure 2A and 2C) and the mBri2/Bri2-Bri23 ratio in *Itm2b^{w/w}* primary neurons was significantly higher than the mBri2/Bri2-ADan ratio in *Itm2b^{D/D}* primary neurons (Figure 2A and 2C). Chloroquine significantly reduced the LC3A I/LC3A II and LC3B I/LC3B II ratios (Figure 2B and 2C), confirming inhibition of lysosome-mediated degradation.

Subtle increase in Aβ42 levels in young *Itm2b^D* KI rats.

Sequential processing of APP by α-/γ-secretase and β-/γ-secretase generate the following APP metabolites: sAPPβ, sAPPα, β-CTF, α-CTF, AID/AICD, P3 and Aβ. Since BRI2 interacts with APP and modulates APP processing by α-, β- and γ-secretase (26-30), we determined the steady-state levels of several of these APP metabolites in the central nervous system (CNS) of young male and female *Itm2b^D* KI rats. Full length APP, α-CTF and β-CTF were measured by Western blot: soluble APPs (sAPPα/sAPPβ) were detected by ELISA, and human Aβ species (Aβ38, Aβ40, Aβ42 and Aβ43) were detected by human Aβ specific-ELISA. These measurements have previously been used for other KI rats generated in our lab (38,42-44).

Levels of full-length APP, CTFs, Aβ38, Aβ40, Aβ43, sAPPα and sAPPβ were unchanged in 8 weeks old *Itm2b^{D/w}*, *Itm2b^{D/D}* and *Itm2b^{w/w}* rats (Figure 3A-C), nor was the Aβ43/Aβ42 ratio altered (Figure 3C). In

contrast, there was a slight but significant increase in Aβ42 as well as the Aβ42/Aβ40 ratio in *Itm2b^{D/D}* compared to *Itm2b^{w/w}* rats (Figure 3C). Small but statistically significant decreases in both Aβ43 and Aβ43/Aβ42 ratio were evident in *Itm2b^{D/D}* as compared to *Itm2b^{w/w}* rats (Figure 3C). Overall, these data indicate a gene dosage-dependent minor increase in steady-state levels of Aβ42, and decrease in Aβ43, in peri-adolescent *Itm2b^D* rats. Analysis of older rats will be needed to determine whether the Danish mutation in *Itm2b* alters APP processing in KI rats and whether these alterations may more robustly change the steady-state levels of APP metabolites with aging.

It has been postulated that toxic forms of Aβ are oligomeric (45). Thus, we tested whether toxic oligomers are augmented in peri-adolescent *Itm2b^D* rats. To this end, we used the prefibrillar oligomer-specific antibody A11 to perform dot blots (46). We found no evidence supporting an increase in neurotoxic brain oligomer levels in peri-adolescent *Itm2b^{D/w}* and *Itm2b^{D/D}* rats as compared with *Itm2b^{w/w}* rats (Figure 4). However, Aβ oligomers appeared to be significantly increased in *Itm2b^{D/w}* rats compared to *Itm2b^{D/D}* animals (Figure 4). Analysis of older rats will be needed to clarify the relevance of this odd observation.

Glutamatergic synaptic transmission at hippocampal SC-CA3>CA1 synapses is impaired in peri-adolescent *Itm2b^D* rats.

Bri2 modulates glutamatergic synaptic transmission at both pre- and post-synaptic termini of Schaeffer-collateral pathway (SC)-CA3>CA1 synapses (47). This function is compromised in both adult *Itm2b^D* and *Itm2b^B* KI mice (41). Here, analyzed glutamatergic transmission at SC-CA3>CA1 synapses in young peri-adolescent *Itm2b^D* male and female rats. First, we analyzed miniature excitatory postsynaptic currents (mEPSC), the frequency of which is determined, in part, by the probability of release (Pr) of glutamatergic synaptic vesicles release (48). Thus, mEPSC frequency is regulated mostly by pre-synaptic mechanisms: As shown in Figure 5A, B,C, the Danish *Itm2b* mutation

Danish dementia and excitatory transmission

caused a significant reduction in the frequency of mEPSC: this reduction is gene-dosage dependent (*Itm2b^{w/w}* vs. *Itm2b^{D/w}*, $P=0.003$; *Itm2b^{w/w}* vs. *Itm2b^{D/D}*, $P<0.0001$; *Itm2b^{D/w}* vs. *Itm2b^{D/D}*, $P=0.0051$) and suggests a decrease in Pr of glutamatergic synaptic vesicles.

The amplitude of mEPSC is instead dependent on post-synaptic AMPA receptor (α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid-receptor, AMPAR) responses. AMPAR-mediated mEPSC responses amplitude was also significantly decreased in *Itm2b^D* rats (Figure 5A, D, E, G). Also in this case, the reduction is gene-dosage dependent (*Itm2b^{w/w}* vs. *Itm2b^{D/w}*, $P=0.0016$; *Itm2b^{w/w}* vs. *Itm2b^{D/D}*, $P=0.0005$). Decay time of mEPSC was not significantly affected in *Itm2b^D* rats compared to littermate controls (Figure 5A, F, G).

Since mEPSCs AMPAR-mediated responses are reduced in amplitude, we measured the AMPA/NMDA ratio in evoked responses. Consistent with the hypothesis that the Danish *Itm2b* mutation impairs AMPAR-mediated responses, the AMPA/NMDA ratio was reduced in Danish KI rats (Figure 5H). This difference was statistically different only between *Itm2b^{w/w}* and *Itm2b^{D/D}* rats, with *Itm2b^{D/w}* rats showing an intermedia phenotype.

Finally, we examined the effect of the pathogenic Danish mutation on paired-pulse facilitation (PPF). PPF is a form of short-term synaptic plasticity that is in part determined by changes in Pr of glutamatergic synaptic vesicles (48)(41). Facilitation at both 50ms and 200ms interstimulus interval (ISI), was significantly increased in *Itm2b^{D/D}* (Figure 5I). Even in this case the changes were gene-dosage-dependent (50ms ISI: *Itm2b^{w/w}* vs. *Itm2b^{D/D}*, $P<0.0001$; *Itm2b^{D/w}* vs. *Itm2b^{D/D}*, $P=0.0031$; 200ms ISI: *Itm2b^{w/w}* vs. *Itm2b^{D/D}*, $P=0.0205$). Interestingly, also an increase in PPF is consistent with a decrease in Pr, just like a decrease in mEPSC frequency. Overall, our data indicate that the pathogenic Danish *Itm2b* mutation alters glutamatergic synaptic transmission at excitatory hippocampal SC-CA3>CA1 synapses in peri-adolescent KI

rats. These alterations are like those seen in *Itm2b* KO and *Itm2b^D/Itm2b^B* KI adult mice.

DISCUSSION

The choice of the genetic approach and the model organisms used to model human diseases have major implications on the phenotypic expression of disease-associated genetic mutations. For the last 13 years, our laboratory has modeled AD and AD-like neurodegenerative disorders in mice, using a KI approach (3,49-52). The KI approach was preferred because it generates models genetically faithful to human diseases and make no preconceived assumption about pathogenic mechanisms (except the unbiased genetic one). We have recently extended our KI modeling of familial and sporadic forms of AD and AD-related disorders to rats (38,42,44,53,54), because the rat is better suited for behavioral tests and other procedures that are important when studying neurodegenerative diseases. In addition, gene-expression differences suggest that rats may be advantageous model of neurodegenerative diseases over mice. Alternative splicing of *Mapt* (12-15), which forms NFTs and is mutated in Frontotemporal Dementia (16-23), leads to expression of tau isoforms with three or four microtubule binding domains (3R and 4R, respectively). Adult human and rat brains express both 3R and 4R tau isoforms (24): in contrast, adult mouse brains express only 4R tau (25), suggesting that the rat may be a better model organism for dementias with tauopathy.

To explore early dysfunctions that may underlie initial mechanisms leading to dementia, we studied young KI rats carrying the *Itm2b^D* Familial Danish dementia mutation. Consistent with the findings in *Itm2b^{D/D}* mouse KIs (41), we found that Bri2-ADan maturation is altered and accumulates in *Itm2b^{D/D}* primary neurons (Figure 2). Analysis of APP metabolism in peri-adolescent *Itm2b^D* Ki rats (Figure 3 and 4) only showed subtle but significant changes in A β 42 and A β 43 steady-state levels, which were slightly increased and decreased, respectively (Figure 3).

Danish dementia and excitatory transmission

We have previously shown that the Danish and British *ITM2b* mutations lead to reduced glutamatergic neurotransmitter release and AMPAR-mediated responses in adult *Itm2b^B* and *Itm2b^D* mice (41). These reductions are like those seen in adult *Itm2b* knock-out mice (41,47). Interestingly, we detected identical, gene dosage-dependent, pre-and post-synaptic glutamatergic transmission changes in the SC pathway of peri-adolescent *Itm2b^D* rats (Figure 5). More specifically, the frequency of mEPSC and the PPF are significantly decreased and increased, respectively, in *Itm2b^D* rats, suggesting a pre-synaptic reduction of the Pr of glutamatergic synaptic vesicles. In addition, mEPSCs amplitude and the AMPA/NMDA ratio were both decreased in *Itm2b^D* rats, suggesting a post-synaptic reduction of AMPAR-mediated responses. Collectively, these data together with our previously published observations, indicate that the synaptic transmission alteration caused by Danish mutation occur early in life, and are neither species nor gene-editing technology-specific. These studies underlie the potential relevance of our studies to functional changes caused by the pathogenic *ITM2b* mutations in humans.

Given the functional and pathological interaction between APP and BRI2 (26-36), it is possible that the presence of human A β in the rat model may lead to an earlier manifestation of synaptic plasticity deficit in rat as compared to mice, which express rodent A β . Moreover, the evidence that both APP and BRI2 tune excitatory synaptic transmission, and that these functions are altered by pathogenic mutations in both APP and BRI2 (37,41,47,55-57) suggest that early alterations in glutamatergic transmission may underlie initial pathogenic mechanisms in dementia. Future studies will be needed to test these hypotheses.

EXPERIMENTAL PROCEDURES

Rats and ethics statement- Rats were handled according to the NIH Ethical Guidelines for Treatment of Laboratory Animals. The procedures were described and approved by the Institutional Animal Care and

Use Committee (IACUC) at Rutgers (IACUC, protocol number PROTO201702513).

Generation of rats expressing the Danish *Itm2b* mutation (*Itm2b^D* rats). The rat *Itm2b* gene (GenBank accession number: NM_001006963.1; Ensembl: ENSRNOG00000016271) is located on rat chromosome 15. It comprises 6 exons, with ATP start codon in exon 1 and TGA stop codon in exon 6. The FDD mutation (TTTAATTGTTCTTGAACAGTCAAGAA AACATTAT) KI site in oligo donor was introduced into exon 6, which is the target site by homology-directed repair. A silent mutation (GTG to GTC) was also introduced to prevent the binding and re-cutting of the sequence by Cas9 after homology-directed repair. The detailed procedures are reported in the Supporting Information file.

Standard RNA-Seq analysis- Total brain RNA from 21 days old *Itm2b^{D/D}* and *Itm2b^{ww/}* rats (2 male and 2 females per each genotype) was extracted with RNeasy RNA Isolation kit (Qiagen). Standard RNA-Seq procedures and data analysis was performed by Genewiz following proprietary methods (https://cdn2.hubspot.net/hubfs/3478602/NGS/RNA-Seq/GENEWIZ_RNA-Seq_Technical_Specifications_US.pdf). Student's t-test was used for all analyses, with data presented as mean $\pm$ SD.

Rats brain proteins preparation, Western blots and ELISA- These procedures were performed as previously described (42,54). Briefly, rats were anesthetized with isoflurane and perfused via intracardiac catheterization with ice-cold PBS. Brains were extracted and homogenized with a glass-teflon homogenizer in 250 mM Sucrose, 20 mM Tris-base pH 7.4, 1 mM EDTA, 1mM EGTA plus protease and phosphatase inhibitors (ThermoScientific). All steps were carried out on ice. Homogenates were solubilized with 1% NP-40 for 30 min rotating and spun at 20,000 g for 10 minutes. Supernatants were collected and protein content was quantified by Bradford.

For Western blot analyses, proteins were diluted with PBS and LDS Sample buffer-10% β -mercaptoethanol (Invitrogen

Danish dementia and excitatory transmission

NP0007) and 4.5M urea to 1µg/µl, loaded on a 4-12% Bis-Tris polyacrylamide gel (Biorad 3450125), and transferred onto nitrocellulose at 25V for 7min using the Trans-blot Turbo system (Biorad). Blotting efficiency was visualized by red Ponceau staining on membranes. For Dot-blot analysis 2.5 µg of material was directly spotted with a p20 pipette on a nitrocellulose membrane. Dot membrane was also visualized by red Ponceau after it was totally dried. Membranes were blocked in 5%-milk (Biorad 1706404) for 30 minutes and washed in PBS/Tween20-0.05%. Primary antibodies were applied dilution in blocking solution (Thermo 37573). The following antibodies were used: Polyclonal anti-Bri2 serum test bleeds provided by Cell Signaling Technology was used at 1:500 with overnight shaking at 4°C. APP-Y188 (Abcam32136), Oligomer Aβ A11 (shared by Rakez Kaye's Lab), LC3A (CST 4599) and LC3B (CST 2775) were used at 1:1000 with same other condition. Secondary antibodies [either anti-mouse (Southern Biotech 1031-05) or a 1:1 mix of anti-rabbit (Southern Biotech, OB405005) and anti-rabbit (Cell Signaling, 7074)], were diluted 1:1000 in 5% milk and used against either mouse or rabbit primary antibodies for 1 hour at RT, with shaking. Membranes were washed with PBS/Tween20-0.05% (3 times, 10 minutes each time), developed with West Dura ECL reagent (Thermo, PI34076) and visualized on a ChemiDoc MP Imaging System (Biorad). Signal intensity was quantified with Image Lab software (Biorad). Data were analyzed using Prism software and represented as mean ± SD.

For analysis of human Aβ peptides and sAPPα/sAPPβ, brain lysates were diluted at 4µg/µl. Aβ38, Aβ40, and Aβ42 were measured with V-PLEX Plus Aβ Peptide Panel 1 6E10 (K15200G, Meso Scale Discovery) and sAPPα/sAPPβ were measured with sAPPα/sAPPβ (K15120E, Meso Scale Discovery). Plates were read on a MESO QuickPlex SQ 120. Aβ43 was quantified using the IBL human Aβ43 Assay Kit #27710.

Primary hippocampal neuron culture- Rat hippocampal neurons were prepared from *Itm2b^{w/w}* and *Itm2b^{D/D}* post-natal day 1 pups. Briefly, after removal of meninges, the hippocampi were collected in HBSS without magnesium and calcium, 1mM sodium pyruvate, 0.1% glucose, 10mM HEPES. Hippocampi were dissected into single cell by trituration followed by 15 minutes incubation at 37°C in 0.25% trypsin. Cells were subsequently treated with 0.1% DNase (Sigma, dn25) in plating media (BME, 10% FBS, 0.09% Glucose, 1mM Sodium Pyruvate, 2mM Glutamine, 1x Pen/Strep). Cells were filtered through a Falcon 70µm nylon cell strainer and were plated in Poly L lysine pretreated 12-well-plate (300,000 cells/well) in Neurobasal media, 1x B-27, 2mM glutamine, 1x Pen/Strep. Half of the culture media was changed every 2 days.

Pharmacological treatment and sample preparation- After 9 days in culture, primary neurons were treated with 50µM Chloroquine (Cell signaling, 14774s) or PBS (Veh) for 18 hours. After treatment, cells were washed with PBS and lysed in RIPA buffer with protease/phosphatase inhibitor for 15 minutes on ice. Lysed cells were centrifuged at full speed for 15 minutes. Cell lysates were quantified and analyzed by Western blot as described earlier for brain lysates.

Electrophysiological recording- These procedures were performed as previously described (44). Briefly, rats were anesthetized with isoflurane and perfused intracardially with an ice-cold cutting solution containing (in mM) 120 choline chloride, 2.6 KCl, 26 NaHCO₃, 1.25 NaH₂PO₄, 0.5 CaCl₂, 7 MgCl₂, 1.3 ascorbic acid, 15 glucose, prebubbled with 95% O₂/5% CO₂ for 15 min. The brains were rapidly removed from the skull and coronal brain slices containing the hippocampal formation (350µm thick) were prepared in the ice-cold cutting solution bubbled with 95% O₂/5% CO₂ using Vibratome VT1200S (Leica Microsystems) and then incubated in an interface chamber in ACSF containing (in mM): 126 NaCl, 3 KCl, 1.2 NaH₂PO₄, 1.3 MgCl₂, 2.4 CaCl₂, 26 NaHCO₃, and 10 glucose (at pH 7.3), bubbled with 95% O₂ and 5% CO₂

Danish dementia and excitatory transmission

at 30°C for 1hr and then kept at room temperature. The hemi-slices were transferred to a recording chamber perfused with ACSF at a flow rate of ~2ml/min using a peristaltic pump. Experiments were performed at 28.0 ± 0.1°C.

Whole-cell recordings in the voltage-clamp mode(-70 mV) were made with patch pipettes containing (in mM): 132.5 Cs-gluconate, 17.5 CsCl, 2 MgCl₂, 0.5 EGTA, 10 HEPES, 4 ATP, and 5 QX-314, with pH adjusted to 7.3 by CsOH. Patch pipettes (resistance, 8–10 MΩ) were pulled from 1.5 mm thin-walled borosilicate glass (Sutter Instruments, Novato, CA) on a horizontal puller (model P-97; Sutter Instruments, Novato, CA). Basal synaptic responses were evoked at 0.05 Hz by electrical stimulation of the Schaffer collateral afferents using concentric bipolar electrodes. CA1 neurons were viewed under upright microscopy (FN-1, Nikon Instruments, Melville, NY) and recorded with Axopatch-700B amplifier (Molecular Devices, San Jose, CA). Data were low-pass filtered at 2 kHz and acquired at 5–10 kHz. The series resistance (Rs) was consistently monitored during recording in case of reseal of ruptured membrane. Cells with Rs >20 MΩ or with Rs deviated by >20% from initial values were excluded from analysis. Excitatory postsynaptic currents (EPSCs) were recorded in ACSF containing the GABA-A receptors inhibitor bicuculline methiodide (15μM). The stimulation intensity was adjusted to evoke EPSCs that were 40% of the maximal evoked amplitudes (“test intensity”). 5–10 min after membrane rupture, EPSCs were recorded for 7 minutes at a test stimulation intensity that produced currents of ~40% maximum. For recording of paired-pulse ratio (PPR), paired-pulse stimuli with 50ms or 200ms inter-pulse interval were given. The PPR was calculated as the ratio of the second EPSC amplitude to the first. For recording of AMPA/NMDA ratio, the membrane potential was firstly held at -70 mV to record only AMPAR current for 20 sweeps with 20s intervals. Then the membrane potential was turned to +40 mV to record NMDAR current for 20 sweeps with perfusion

of 5μM NBQX to block AMPAR. Mini EPSCs were recorded by maintaining neurons at -70 mV with ACSF containing action potentials blocker (1μM TTX) and GABA-A receptors inhibitors (15μM bicuculline methiodide). mEPSCs were recorded for ~10 mins. Data were collected with Axopatch 700B amplifiers and analyzed with pCLAMP10 software (Molecular Devices). mEPSCs are analyzed using mini-Analysis Program.

Statistics- All the experiments mentioned in the paper were analyzed by one-way ANOVA or two-way ANOVA as indicated. Data showing statistical significance by one-way ANOVA or two-way ANOVA were subsequently analyzed by either Tukey's multiple comparisons test or Sidak's multiple comparisons. All statistical analyses were performed using Prism 9 (GraphPad) software.

Acknowledgement- All authors read and approved the final manuscript.

Declarations-The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Conflicts of interest-The authors declare that they have no competing interests.

Author Contributions- LD generated the animals; KAN set up breeding and genotyped animals; TY performed the biochemical and molecular experiments; WY performed the electrophysiology experiments; All authors designed the experiments; LD and TY wrote the paper.

L.D. was funded by the NIH/NIA R01AG063407, R01AG064821, 1R01AG033007

References

1. Tamayev, R., Matsuda, S., Fa, M., Arancio, O., and D'Adamio, L. (2010) Danish dementia mice suggest that loss of function and not the amyloid cascade causes synaptic plasticity and memory deficits. *Proc Natl Acad Sci U S A* **107**, 20822-20827

Danish dementia and excitatory transmission

2. Tamayev, R., Giliberto, L., Li, W., d'Abramo, C., Arancio, O., Vidal, R., and D'Adamio, L. (2010) Memory deficits due to familial British dementia BRI2 mutation are caused by loss of BRI2 function rather than amyloidosis. *J Neurosci* **30**, 14915-14924
3. Giliberto, L., Matsuda, S., Vidal, R., and D'Adamio, L. (2009) Generation and initial characterization of FDD knock in mice. *PLoS One* **4**, e7900
4. Deacon, R. M. (2006) Housing, husbandry and handling of rodents for behavioral experiments. *Nat Protoc* **1**, 936-946
5. Whishaw, I. Q., Metz, G. A., Kolb, B., and Pellis, S. M. (2001) Accelerated nervous system development contributes to behavioral efficiency in the laboratory mouse: a behavioral review and theoretical proposal. *Dev Psychobiol* **39**, 151-170
6. Kepecs, A., Uchida, N., Zariwala, H. A., and Mainen, Z. F. (2008) Neural correlates, computation and behavioural impact of decision confidence. *Nature* **455**, 227-231
7. Foote, A. L., and Crystal, J. D. (2007) Metacognition in the rat. *Curr Biol* **17**, 551-555
8. Bartelle, B. B., Barandov, A., and Jasanoff, A. (2016) Molecular fMRI. *J Neurosci* **36**, 4139-4148
9. Zimmer, E. R., Leuzy, A., Bhat, V., Gauthier, S., and Rosa-Neto, P. (2014) In vivo tracking of tau pathology using positron emission tomography (PET) molecular imaging in small animals. *Transl Neurodegener* **3**, 6
10. Leuzy, A., Zimmer, E. R., Heurling, K., Rosa-Neto, P., and Gauthier, S. (2014) Use of amyloid PET across the spectrum of Alzheimer's disease: clinical utility and associated ethical issues. *Amyloid* **21**, 143-148
11. Zimmer, E. R., Leuzy, A., Gauthier, S., and Rosa-Neto, P. (2014) Developments in Tau PET Imaging. *Can J Neurol Sci* **41**, 547-553
12. Andreadis, A. (2005) Tau gene alternative splicing: expression patterns, regulation and modulation of function in normal brain and neurodegenerative diseases. *Biochim Biophys Acta* **1739**, 91-103
13. Janke, C., Beck, M., Stahl, T., Holzer, M., Brauer, K., Bigl, V., and Arendt, T. (1999) Phylogenetic diversity of the expression of the microtubule-associated protein tau: implications for neurodegenerative disorders. *Brain Res Mol Brain Res* **68**, 119-128
14. Hong, M., Zhukareva, V., Vogelsberg-Ragaglia, V., Wszolek, Z., Reed, L., Miller, B. I., Geschwind, D. H., Bird, T. D., McKeel, D., Goate, A., Morris, J. C., Wilhelmsen, K. C., Schellenberg, G. D., Trojanowski, J. Q., and Lee, V. M. (1998) Mutation-specific functional impairments in distinct tau isoforms of hereditary FTDP-17. *Science* **282**, 1914-1917
15. Roberson, E. D., Scarce-Levie, K., Palop, J. J., Yan, F., Cheng, I. H., Wu, T., Gerstein, H., Yu, G. Q., and Mucke, L. (2007) Reducing endogenous tau ameliorates amyloid beta-induced deficits in an

Danish dementia and excitatory transmission

- 863 Alzheimer's disease mouse model.
- 864 *Science* **316**, 750-754
- 865 16. Spillantini, M. G., and Goedert, M.
- 866 (1998) Tau protein pathology in
- 867 neurodegenerative diseases.
- 868 *Trends Neurosci* **21**, 428-433
- 869 17. Goedert, M., Crowther, R. A., and
- 870 Spillantini, M. G. (1998) Tau
- 871 mutations cause frontotemporal
- 872 dementias. *Neuron* **21**, 955-958
- 873 18. Grundke-Iqbal, I., Iqbal, K., Tung, Y.
- 874 C., Quinlan, M., Wisniewski, H. M.,
- 875 and Binder, L. I. (1986) Abnormal
- 876 phosphorylation of the
- 877 microtubule-associated protein
- 878 tau (tau) in Alzheimer cytoskeletal
- 879 pathology. *Proc Natl Acad Sci U S A*
- 880 **83**, 4913-4917
- 881 19. Hutton, M., Lendon, C. L., Rizzu, P.,
- 882 Baker, M., Froelich, S., Houlden,
- 883 H., Pickering-Brown, S.,
- 884 Chakraborty, S., Isaacs, A., Grover,
- 885 A., Hackett, J., Adamson, J.,
- 886 Lincoln, S., Dickson, D., Davies, P.,
- 887 Petersen, R. C., Stevens, M., de
- 888 Graaff, E., Wauters, E., van Baren,
- 889 J., Hillebrand, M., Joosse, M.,
- 890 Kwon, J. M., Nowotny, P., Che, L.
- 891 K., Norton, J., Morris, J. C., Reed, L.
- 892 A., Trojanowski, J., Basun, H.,
- 893 Lannfelt, L., Neystat, M., Fahn, S.,
- 894 Dark, F., Tannenberg, T., Dodd, P.
- 895 R., Hayward, N., Kwok, J. B.,
- 896 Schofield, P. R., Andreadis, A.,
- 897 Snowden, J., Craufurd, D., Neary,
- 898 D., Owen, F., Oostra, B. A., Hardy,
- 899 J., Goate, A., van Swieten, J.,
- 900 Mann, D., Lynch, T., and Heutink,
- 901 P. (1998) Association of missense
- 902 and 5'-splice-site mutations in tau
- 903 with the inherited dementia FTDP-
- 904 17. *Nature* **393**, 702-705
- 905 20. Stanford, P. M., Shepherd, C. E.,
- 906 Halliday, G. M., Brooks, W. S.,
- 907 Schofield, P. W., Brodaty, H.,
- 908 Martins, R. N., Kwok, J. B., and
- 909 Schofield, P. R. (2003) Mutations
- 910 in the tau gene that cause an
- 911 increase in three repeat tau and
- 912 frontotemporal dementia. *Brain*
- 913 **126**, 814-826
- 914 21. Yasuda, M., Takamatsu, J.,
- 915 D'Souza, I., Crowther, R. A.,
- 916 Kawamata, T., Hasegawa, M.,
- 917 Hasegawa, H., Spillantini, M. G.,
- 918 Tanimukai, S., Poorkaj, P., Varani,
- 919 L., Varani, G., Iwatsubo, T.,
- 920 Goedert, M., Schellenberg, D. G.,
- 921 and Tanaka, C. (2000) A novel
- 922 mutation at position +12 in the
- 923 intron following exon 10 of the tau
- 924 gene in familial frontotemporal
- 925 dementia (FTD-Kumamoto). *Ann*
- 926 *Neurol* **47**, 422-429
- 927 22. Kowalska, A., Hasegawa, M.,
- 928 Miyamoto, K., Akiguchi, I.,
- 929 Ikemoto, A., Takahashi, K., Araki,
- 930 W., and Tabira, T. (2002) A novel
- 931 mutation at position +11 in the
- 932 intron following exon 10 of the tau
- 933 gene in FTDP-17. *J Appl Genet* **43**,
- 934 535-543
- 935 23. Grover, A., England, E., Baker, M.,
- 936 Sahara, N., Adamson, J., Granger,
- 937 B., Houlden, H., Passant, U., Yen, S.
- 938 H., DeTure, M., and Hutton, M.
- 939 (2003) A novel tau mutation in
- 940 exon 9 (1260V) causes a four-
- 941 repeat tauopathy. *Exp Neurol* **184**,
- 942 131-140
- 943 24. Hanes, J., Zilka, N., Bartkova, M.,
- 944 Caletkova, M., Dobrota, D., and
- 945 Novak, M. (2009) Rat tau
- 946 proteome consists of six tau
- 947 isoforms: implication for animal
- 948 models of human tauopathies. *J*
- 949 *Neurochem* **108**, 1167-1176

Danish dementia and excitatory transmission

25. McMillan, P., Korvatska, E., Poorkaj, P., Evstafjeva, Z., Robinson, L., Greenup, L., Leverenz, J., Schellenberg, G. D., and D'Souza, I. (2008) Tau isoform regulation is region- and cell-specific in mouse brain. *J Comp Neurol* **511**, 788-803
26. Matsuda, S., Giliberto, L., Matsuda, Y., Davies, P., McGowan, E., Pickford, F., Ghiso, J., Frangione, B., and D'Adamio, L. (2005) The familial dementia BRI2 gene binds the Alzheimer gene amyloid-beta precursor protein and inhibits amyloid-beta production. *J Biol Chem* **280**, 28912-28916
27. Fotinopoulou, A., Tsachaki, M., Vlavaki, M., Pouloupoulos, A., Rostagno, A., Frangione, B., Ghiso, J., and Efthimiopoulos, S. (2005) BRI2 interacts with amyloid precursor protein (APP) and regulates amyloid beta (Abeta) production. *J Biol Chem* **280**, 30768-30772
28. Matsuda, S., Giliberto, L., Matsuda, Y., McGowan, E. M., and D'Adamio, L. (2008) BRI2 inhibits amyloid beta-peptide precursor protein processing by interfering with the docking of secretases to the substrate. *J Neurosci* **28**, 8668-8676
29. Matsuda, S., Matsuda, Y., Snapp, E. L., and D'Adamio, L. (2011) Maturation of BRI2 generates a specific inhibitor that reduces APP processing at the plasma membrane and in endocytic vesicles. *Neurobiol Aging* **32**, 1400-1408
30. Matsuda, S., Tamayev, R., and D'Adamio, L. (2011) Increased AβPP processing in familial Danish dementia patients. *J Alzheimers Dis* **27**, 385-391
31. Tamayev, R., Matsuda, S., Giliberto, L., Arancio, O., and D'Adamio, L. (2011) APP heterozygosity averts memory deficit in knockin mice expressing the Danish dementia BRI2 mutant. *EMBO J* **30**, 2501-2509
32. Tamayev, R., and D'Adamio, L. (2012) Inhibition of γ-secretase worsens memory deficits in a genetically congruous mouse model of Danish dementia. *Mol Neurodegener* **7**, 19
33. Tamayev, R., and D'Adamio, L. (2012) Memory deficits of British dementia knock-in mice are prevented by Aβ-precursor protein haploinsufficiency. *J Neurosci* **32**, 5481-5485
34. Tamayev, R., Matsuda, S., Arancio, O., and D'Adamio, L. (2012) β- but not γ-secretase proteolysis of APP causes synaptic and memory deficits in a mouse model of dementia. *EMBO Mol Med* **4**, 171-179
35. Lombino, F., Biundo, F., Tamayev, R., Arancio, O., and D'Adamio, L. (2013) An intracellular threonine of amyloid-β precursor protein mediates synaptic plasticity deficits and memory loss. *PLoS One* **8**, e57120
36. Biundo, F., Ishiwari, K., Del Prete, D., and D'Adamio, L. (2016) Deletion of the γ-secretase subunits Aph1B/C impairs memory and worsens the deficits of knock-in mice modeling the Alzheimer-

Danish dementia and excitatory transmission

- 1037 like familial Danish dementia.
- 1038 *Oncotarget* **7**, 11923-11944
- 1039 37. Tambini, M. D., Yao, W., and
- 1040 D'Adamio, L. (2019) Facilitation of
- 1041 glutamate, but not GABA, release
- 1042 in Familial Alzheimer's APP mutant
- 1043 Knock-in rats with increased β -
- 1044 cleavage of APP. *Aging Cell* **18**,
- 1045 e13033
- 1046 38. Tambini, M. D., Norris, K. A., and
- 1047 D'Adamio, L. (2020) Opposite
- 1048 changes in APP processing and
- 1049 human A β levels in rats carrying
- 1050 either a protective or a pathogenic
- 1051 APP mutation. *Elife* **9**
- 1052 39. Vidal, R., Revesz, T., Rostagno, A.,
- 1053 Kim, E., Holton, J. L., Bek, T.,
- 1054 Bojsen-Møller, M., Braendgaard,
- 1055 H., Plant, G., Ghiso, J., and
- 1056 Frangione, B. (2000) A decamer
- 1057 duplication in the 3' region of the
- 1058 BRI gene originates an amyloid
- 1059 peptide that is associated with
- 1060 dementia in a Danish kindred. *Proc*
- 1061 *Natl Acad Sci U S A* **97**, 4920-4925
- 1062 40. Choi, S. I., Vidal, R., Frangione, B.,
- 1063 and Levy, E. (2004) Axonal
- 1064 transport of British and Danish
- 1065 amyloid peptides via secretory
- 1066 vesicles. *FASEB J* **18**, 373-375
- 1067 41. Yin, T., Yao, W., Lemenze, A. D.,
- 1068 and D'Adamio, L. (2020) Danish
- 1069 and British dementia ITM2b/BRI2
- 1070 mutations reduce BRI2 protein
- 1071 stability and impair glutamatergic
- 1072 synaptic transmission. *J Biol Chem*
- 1073 42. Tambini, M. D., and D'Adamio, L.
- 1074 (2020) Trem2 Splicing and
- 1075 Expression are Preserved in a
- 1076 Human Abeta-producing, Rat
- 1077 Knock-in Model of Trem2-R47H
- 1078 Alzheimer's Risk Variant. *Sci Rep*
- 1079 **10**, 4122
- 1080 43. Ren, S., Breuillaud, L., Yao, W., Yin,
- 1081 T., Norris, K. A., Zehntner, S. P.,
- 1082 and D'Adamio, L. (2020) TNF- α -
- 1083 mediated reduction in inhibitory
- 1084 neurotransmission precedes
- 1085 sporadic Alzheimer's disease
- 1086 pathology in young. *J Biol Chem*
- 1087 44. Ren, S., Yao, W., Tambini, M. D.,
- 1088 Yin, T., Norris, K. A., and D'Adamio,
- 1089 L. (2020) Microglia *TREM2*^{R47H}
- 1090 Alzheimer-linked variant enhances
- 1091 excitatory transmission and
- 1092 reduces LTP via increased TNF- α
- 1093 levels. *Elife* **9**
- 1094 45. Shankar, G. M., Li, S., Mehta, T. H.,
- 1095 Garcia-Munoz, A., Shepardson, N.
- 1096 E., Smith, I., Brett, F. M., Farrell,
- 1097 M. A., Rowan, M. J., Lemere, C. A.,
- 1098 Regan, C. M., Walsh, D. M.,
- 1099 Sabatini, B. L., and Selkoe, D. J.
- 1100 (2008) Amyloid-beta protein
- 1101 dimers isolated directly from
- 1102 Alzheimer's brains impair synaptic
- 1103 plasticity and memory. *Nat Med*
- 1104 **14**, 837-842
- 1105 46. Kaye, R., Head, E., Thompson, J.
- 1106 L., McIntire, T. M., Milton, S. C.,
- 1107 Cotman, C. W., and Glabe, C. G.
- 1108 (2003) Common structure of
- 1109 soluble amyloid oligomers implies
- 1110 common mechanism of
- 1111 pathogenesis. *Science* **300**, 486-
- 1112 489
- 1113 47. Yao, W., Yin, T., Tambini, M. D.,
- 1114 and D'Adamio, L. (2019) The
- 1115 Familial dementia gene
- 1116 ITM2b/BRI2 facilitates glutamate
- 1117 transmission via both presynaptic
- 1118 and postsynaptic mechanisms. *Sci*
- 1119 *Rep* **9**, 4862
- 1120 48. Zucker, R. S., and Regehr, W. G.
- 1121 (2002) Short-term synaptic
- 1122 plasticity. *Annu Rev Physiol* **64**,
- 1123 355-405

Danish dementia and excitatory transmission

- 1124 49. Barbagallo, A. P., Wang, Z., Zheng,
1125 H., and D'Adamio, L. (2011) The
1126 intracellular threonine of amyloid
1127 precursor protein that is essential
1128 for docking of Pin1 is dispensable
1129 for developmental function. *PLoS*
1130 *One* **6**, e18006
- 1131 50. Barbagallo, A. P., Wang, Z., Zheng,
1132 H., and D'Adamio, L. (2011) A
1133 single tyrosine residue in the
1134 amyloid precursor protein
1135 intracellular domain is essential for
1136 developmental function. *J Biol*
1137 *Chem* **286**, 8717-8721
- 1138 51. Barbagallo, A. P., Weldon, R.,
1139 Tamayev, R., Zhou, D., Giliberto, L.,
1140 Foreman, O., and D'Adamio, L.
1141 (2010) Tyr(682) in the intracellular
1142 domain of APP regulates
1143 amyloidogenic APP processing in
1144 vivo. *PLoS One* **5**, e15503
- 1145 52. Garringer, H. J., Murrell, J.,
1146 D'Adamio, L., Ghetti, B., and Vidal,
1147 R. (2010) Modeling familial British
1148 and Danish dementia. *Brain Struct*
1149 *Funct* **214**, 235-244
- 1150 53. Ren, S., Breuillaud, L., Yao, W., Yin,
1151 T., Norris, K. A., Zehntner, S. P.,
1152 and D'Adamio, L. (2020) TNF- α -
1153 mediated reduction in inhibitory
1154 neurotransmission precedes
1155 sporadic Alzheimer's disease
1156 pathology in young Trem2 R^{47H}
1157 rats. *J Biol Chem* **296**, 100089
- 1158 54. Tambini, M. D., and D'Adamio, L.
1159 (2020) Knock-in rats with
1160 homozygous PSEN1 L^{435F} Alzheimer
1161 mutation are viable and show
1162 selective γ -secretase activity loss
1163 causing low A β 40/42 and high
1164 A β 43. *J Biol Chem* **295**, 7442-7451
- 1165 55. Yao, W., Tambini, M. D., Liu, X.,
1166 and D'Adamio, L. (2019) Tuning of
1167 Glutamate, But Not GABA, Release
1168 by an Intrasyntaptic Vesicle APP
1169 Domain Whose Function Can Be
1170 Modulated by β - or α -Secretase
1171 Cleavage. *J Neurosci* **39**, 6992-
1172 7005
- 1173 56. Fanutza, T., Del Prete, D., Ford, M.
1174 J., Castillo, P. E., and D'Adamio, L.
1175 (2015) APP and APLP2 interact
1176 with the synaptic release
1177 machinery and facilitate
1178 transmitter release at
1179 hippocampal synapses. *Elife* **4**,
1180 e09743
- 1181 57. Del Prete, D., Lombino, F., Liu, X.,
1182 and D'Adamio, L. (2014) APP is
1183 cleaved by Bace1 in pre-synaptic
1184 vesicles and establishes a pre-
1185 synaptic interactome, via its
1186 intracellular domain, with
1187 molecular complexes that regulate
1188 pre-synaptic vesicles functions.
1189 *PLoS One* **9**, e108576
1190

1191

1192

Danish dementia and excitatory transmission

Figure legends

Figure 1. Characterization of *Itm2b*^D KI rats. (A) Sequences of the COOH-terminus of Bri2-23 (WT) and Bri2-ADan (Danish). (B) PCR amplification and sequencing of the *Itm2b* gene exon 6 from *Itm2b*^{w/w} and *Itm2b*^{D/D} rats shows that the Danish mutation was correctly inserted in the *Itm2b* exon 6 of *Itm2b*^{D/D} rats. This mutation causes the predicted frameshift in the BRI2 sequence generating a precursor protein 11 amino acids larger-than-normal coding for the Bri2-ADan mutant protein (partial DNA sequences of WT and Danish exon 6 are shown. Inserted nucleotides are highlighted in black, and the amino-acid sequences are indicated above -for the Danish mutant allele- and below -for the WT allele- the DNA sequences). (C) Levels of *Itm2b* mRNA in brains of 21 days old *Itm2b*^{w/w} and *Itm2b*^{D/D} rats were determined by Standard-RNAseq analysis. No significant differences between *Itm2b*^{w/w} and *Itm2b*^{D/D} rats were evident. Data are represented as mean ± SD. Data were analyzed by Student's t-test. N=4 rats per genotype.

Danish dementia and excitatory transmission

Danish dementia and excitatory transmission

Figure 4. Levels of human Aβ oligomeric species in the brain of peri-adolescent *Itm2b^{D/D}*, *Itm2b^{D/w}* and *Itm2b^{w/w}* male and female rats. (A) We analyzed material from the same rats analyzed in Figure 3. Quantitation of dot-blot using the oligomer-specific antibody A11. Before immunoblot analysis, membranes were stained with Ponceau red. Quantitative analysis of A11 blot was normalized to the Ponceau red quantitative analysis. Data are represented as mean ± SD and were analyzed by ordinary one-way ANOVA followed by post-hoc Tukey's multiple comparisons test when ANOVA showed statistically significant differences. ANOVA summary: $F_{(2,27)} = 4.593$, $P=0.0192^*$; post-hoc Tukey's multiple comparisons test: *Itm2b^{w/w}* vs *Itm2b^{D/w}*, $P=0.6406$, *Itm2b^{w/w}* vs *Itm2b^{D/D}*, $P=0.1195$; *Itm2b^{D/w}* vs *Itm2b^{D/D}*, $P=0.0170^*$.

Danish dementia and excitatory transmission

Figure 5. Glutamatergic synaptic transmission is reduced at hippocampal SC-CA3>CA1 synapses of *Itm2b^D* KI rats. Data are represented as mean $\pm$ SD and were analyzed by ordinary one-way ANOVA followed by post-hoc Tukey's multiple comparisons test when ANOVA showed significant differences. We used the following animals: *Itm2b^{w/w}* N=11 (4M/6, 3F/5, indicating that 6 recordings were obtained from the 4 males and 5 recordings from the 3 females), *Itm2b^{D/w}* N=15 (3M/7, 4F/8), *Itm2b^{D/D}* N=10 (3M/5, 3F/5). (A) Representative recording traces of mEPSC at SC-CA3>CA1 synapses. (B) The Danish mutation causes a significant decrease in mEPSC frequency [ANOVA summary, $F_{(2,33)} = 20.66$, $P < 0.0001$ ****; post-hoc Tukey's multiple comparisons test: *Itm2b^{w/w}* vs *Itm2b^{D/w}*, $P = 0.003$ ***, *Itm2b^{w/w}* vs *Itm2b^{D/D}*, $P < 0.0001$ ****; *Itm2b^{D/w}* vs *Itm2b^{D/D}*, $P = 0.0051$ **]. (C) Cumulative probability of AMPAR-mediated mEPSC frequency inter event intervals. (D) [ANOVA summary, $F_{(2,33)} = 10.78$, $P = 0.0002$ ***; post-hoc Tukey's multiple comparisons test: *Itm2b^{w/w}* vs *Itm2b^{D/w}*, $P = 0.0016$ ***, *Itm2b^{w/w}* vs *Itm2b^{D/D}*, $P = 0.0005$ ****; *Itm2b^{D/w}* vs *Itm2b^{D/D}*, $P = 0.6691$]. (E) Cumulative probability of AMPAR-mediated mEPSC amplitude. (F) In contrast, decay time of mEPSC was not significantly changed [ANOVA summary, $F_{(2,33)} = 1.292$, $P = 0.2882$]. (G) Average mEPSC of *Itm2b^{D/D}*, *Itm2b^{D/w}* and *Itm2b^{w/w}* rats. (H) AMPA/NMDA ratio is significantly decreased in *Itm2b^{D/D}* rats [ANOVA summary, $F_{(2,16)} = 8.417$, $P = 0.0032$ ***, post-hoc Tukey's multiple comparisons test: *Itm2b^{w/w}* vs *Itm2b^{D/w}*, $P = 0.2393$; *Itm2b^{w/w}* vs *Itm2b^{D/D}*, $P = 0.0023$ ***, *Itm2b^{D/w}* vs *Itm2b^{D/D}*, $P = 0.0818$]. Representative traces are shown on top of the graph (traces are averaged from 20 sweeps). Animals used: *Itm2b^{w/w}* N=7 (4M/4, 3F/3), *Itm2b^{D/w}* N=6 (3M/3, 3F/3), *Itm2b^{D/D}* N=6 (3M/3, 3F/3). (I) Average PPF at 50ms (left panel) and 200ms (right panel) Inter stimulus Interval (ISI) shows that PPF is increased in *Itm2b^{D/D}* rats ISI [50ms ISI PPF ANOVA summary: $F_{(2,60)} = 11.89$, $P < 0.0001$ ****; post-hoc Tukey's multiple comparisons test: *Itm2b^{w/w}* vs *Itm2b^{D/w}*, $P = 0.2315$; *Itm2b^{w/w}* vs *Itm2b^{D/D}*, $P < 0.0001$ ****; *Itm2b^{D/w}* vs *Itm2b^{D/D}*, $P = 0.0031$ **]. 200ms ISI PPF ANOVA summary: $F_{(2,64)} = 3.802$, $P = 0.0275$ *; post-hoc Tukey's multiple comparisons test: *Itm2b^{w/w}* vs *Itm2b^{D/w}*, $P = 0.4504$; *Itm2b^{w/w}* vs *Itm2b^{D/D}*, $P = 0.0205$ *; *Itm2b^{D/w}* vs *Itm2b^{D/D}*, $P = 0.2462$]. Representative traces are shown on top of the panels. $P < 0.05$ *; $P < 0.01$ **; $P < 0.001$ ***; $P < 0.0001$ ****. Animals used: *Itm2b^{w/w}* N=18 (4M/10, 3F/8), *Itm2b^{D/w}* N=20 (3M/10, 4F/10), *Itm2b^{D/D}* N=17 (3M/8, 3F/9).