1 **Title: High photosynthesis rate in the selected wild rice is driven by leaf anatomy**
2 **mediating high Rubisco activity and electron transport rate**

3

4 **Running Head:** Efficient photosynthesis in the selected wild rice

5

6 Jyotirmaya Mathan[#], Anuradha Singh[#], Vikram Jathar[#], Aashish Ranjan*

7 National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-110067,
8 India.

9 [#]These authors contributed equally to the work.

10

11

12 ***Corresponding Author**

13 email: aranjan@nipgr.ac.in

14 Phone no. 91-11-26735117

15

16 Authors' email

17 Jyotirmaya Mathan: jyotirmaya@nipgr.ac.in

18 Anuradha Singh: annusingh1206@gmail.com

19 Vikram Jathar: vikram204@nipgr.ac.in

20

21

22

23 Number of figures and tables: 6 figures and 4 tables

24

25

26 The word count: 5,794

27

28

29 Supplementary data: 11 supplemental figures and 6 supplemental tables

30

31

32

33

34

35

36

37

Title: High photosynthesis rate in the selected wild rice is driven by leaf anatomy mediating high Rubisco activity and electron transport rate

Running Head: Efficient photosynthesis in the selected wild rice

Highlight

Distinct leaf biochemical, photochemical, and developmental features contribute to efficient photosynthesis in the selected wild rice accessions that could potentially be exploited to increase rice leaf photosynthesis.

Abstract

The importance of increasing photosynthetic efficiency for sustainable crop yield increases to feed the growing world population is well recognized. The natural genetic variation for leaf photosynthesis in crop plants is largely unexploited for increasing genetic yield potential. The genus *Oryza*, including cultivated rice and wild relatives, offers tremendous genetic variability to explore photosynthetic differences, and underlying biochemical, photochemical, and developmental bases. We quantified leaf photosynthesis and related physiological parameters for six cultivated and three wild rice genotypes, and identified photosynthetically efficient wild rice accessions. Fitting A/C_i curves and biochemical analyses showed that the leaf photosynthesis in cultivated rice varieties, IR64 and Nipponbare, was limited due to leaf nitrogen content, Rubisco activity, and electron transport rate compared to photosynthetically efficient accessions of wild rice *Oryza australiensis* and *Oryza latifolia*. The selected wild rice accessions with high leaf photosynthesis per unit area had striking anatomical features, such as larger mesophyll cells with more chloroplasts, fewer mesophyll cells between two consecutive veins, and higher mesophyll cell and chloroplast surface area exposed to intercellular space. Our results show the existence of desirable variations in Rubisco activity, electron transport rate, and leaf anatomical features in the rice system itself that could be targeted for increasing the photosynthetic efficiency of cultivated rice varieties.

Keywords: A/C_i curve, Chlorophyll fluorescence, Electron transport rate, Leaf anatomy, Photosynthesis, Photosystem II (PSII), Rubisco activity, Wild rice.

72

73 **Introduction**

74 Photosynthesis, the key biochemical process fixing atmospheric CO₂ in the form of
75 carbohydrates, is a multifaceted process that has contributions from leaf physiological,
76 biochemical, and developmental features (Long *et al.*, 2006; Mathan *et al.*, 2016). Increasing
77 photosynthetic efficiency of crop plants is suggested to be an important strategy for
78 sustainable increases in crop yield potential and productivity (Zhu *et al.*, 2010; Lawson *et al.*,
79 2012; Ort *et al.*, 2015; Simkin *et al.*, 2019). The extensive biochemical understanding of
80 photosynthesis over the years has highlighted the potential biochemical targets, such as
81 decreasing Rubisco oxygenase activity, increasing RuBP regeneration, and suppressing
82 photorespiration, to improve photosynthetic efficiency (Horton, 2000; Long *et al.*, 2006;
83 Murchie and Niyogi, 2011). *in vivo* Rubisco activity (V_{max}), maximum electron transport rate
84 (J_{max}) determining the regeneration of ribulose phosphate (RuBP) for Rubisco, and the
85 maximum rate of triose phosphate utilization (TPU) as the three major biochemical
86 limitations to leaf photosynthesis (Farquhar *et al.*, 1980; Sharkey, 1985; von Caemmerer,
87 2000). The response of net CO₂ assimilation under saturating light to intercellular CO₂
88 concentrations (A/C_i curve) is extensively used to deduce these biochemical limitations to
89 efficient leaf photosynthesis (Long and Bernacchi, 2003; Sharkey *et al.*, 2007). However, the
90 limited characterization of genetic variability of these biochemical limitations has restricted
91 the potential of increasing leaf photosynthesis in crop improvement.

92 Leaf photosynthesis limitation due to Rubisco activity, which primarily occurs at low
93 intercellular CO₂ concentration ($< 200 \mu\text{mol mol}^{-1}$), is due to the low activity of Rubisco
94 along with competing carboxylation and oxygenation reactions mediated by the enzyme (von
95 Caemmerer, 2000; Parry *et al.*, 2007; Sharkey *et al.*, 2007). Exploration of natural variation
96 in Rubisco activity, as well as developmental and metabolic features mediating increased
97 concentration of CO₂ and RuBP at the active site of Rubisco, are possible strategies to
98 overcome this limitation of photosynthesis (Parry *et al.*, 2007; Parry *et al.*, 2013; Lin *et al.*,
99 2014; Sharwood, 2017). Organization of LHCs, the quantum efficiency of the photosystem,
100 and photochemical and non-photochemical quenching are key factors for determining
101 photosynthetic limitation due to electron transport rate and RuBP regeneration that operates
102 at CO₂ concentration $> 300 \mu\text{mol mol}^{-1}$ (Murchie and Lawson, 2013; Wu *et al.*, 2019). TPU
103 drives sucrose synthesis from primary photosynthates, and TPU -limited photosynthesis
104 primarily operates at high CO₂ concentrations that often determine A_{max} (Sharkey *et al.*, 2007;
105 Fabre *et al.*, 2019). The genotypic variations for these limitations in crop plants have been

attributed to photosynthetic differences (Acevedo-Siaca *et al.*, 2020; De Souza *et al.*, 2020). For example, Acevedo-Siaca *et al.* (2020) recently investigated the variation among rice accessions to show V_{cmax} as the primary limitation to photosynthetic induction. Therefore, understanding these biochemical limitations in cultivated crop varieties will provide the potential ways to improve photosynthetic efficiency.

In addition to the biochemical limitations, stomatal conductance (g_s) and mesophyll conductance (g_m) that affect leaf gaseous exchange are critical determinants of leaf photosynthesis (Farquhar and Sharkey, 1982; Adachi *et al.*, 2013; Gago *et al.*, 2016). Leaf developmental features, besides affecting the leaf gaseous exchange, influence light harvesting and distribution for photosynthesis (Terashima *et al.*, 2011; Mathan *et al.*, 2016). Flag leaf width (FLW), flag leaf thickness (FLT), and specific leaf area showed a positive correlation with crop photosynthetic efficiency (Liu *et al.*, 2015). Anatomical features such as size, lobbing, and surface area of mesophyll cells are critical for proper light distribution and CO₂ diffusion within a leaf (Flexas *et al.*, 2008; Sage and Sage, 2009; Tholen *et al.*, 2012). Furthermore, the surface area of mesophyll cell (S_{mes}) and chloroplast (S_c) exposed to intercellular spaces are the key determinants of g_m (Scafaro *et al.*, 2011). Similarly, vein density and interveinal distance influence CO₂ fixation efficiency (Amiard *et al.*, 2005; Brodribb *et al.*, 2007). Thus, identifying the exploitable desirable variations in leaf morphological and anatomical features could help in optimizing crop photosynthetic efficiency (Evans, 2013).

Rice, with two cultivated species: *Oryza sativa* (Asian rice with two subspecies: *indica* and *japonica*) and *Oryza glaberrima* (African rice), and many wild rice species, provides an excellent system to investigate the genetic diversity for crop improvement. Rice domestication, which primarily targeted grain features and plant architectural traits, significantly reduced the genetic diversity for many favourable traits in cultivated rice varieties (Huang *et al.*, 2012; Chen *et al.*, 2021). Wild rice species may harbour more natural allelic diversity than cultivated varieties, and hence mining hidden useful variants from wild species could be an important strategy in rice improvement. Wild rice species are already shown to be reservoirs of many useful traits, particularly for biotic and abiotic stress resistance, which have been used in rice breeding programs (Sanchez *et al.*, 2013; Atwell *et al.*, 2014; Wing *et al.*, 2018). Interestingly, some of the wild rice accessions of *O. rufipogon*, *O. latifolia*, and *O. australiensis* are reported to accumulate higher biomass than cultivated varieties, and introgression from a wild *Oryza* species to cultivated rice has also been shown to increase biomass (Yeo *et al.*, 1994; Zhao *et al.*, 2008). Moreover, a few wild rice

accessions are also reported to have high leaf photosynthesis rate than the cultivated varieties (Zhao *et al.*, 2010; Kiran *et al.*, 2013). Thus, wild relatives of rice could be a valuable resource for increasing leaf photosynthetic efficiency of cultivated varieties. However, a systematic effort to identify photosynthetically efficient wild rice accessions along with biochemical, photochemical, and developmental traits promoting high photosynthesis rate in those accessions is warranted.

We examined a set of cultivated and wild rice accessions to find out photosynthetically efficient wild accessions. We identified that the Rubisco activity and electron transport rate limit leaf photosynthesis in the cultivated varieties compared to the selected wild accessions. Photochemical and developmental differences, further, corroborated the leaf photosynthesis differences across the selected cultivated and wild rice.

Materials and methods

Plant material and growth conditions

Nine rice genotypes, consisting of four varieties of *Oryza sativa* ssp. *indica* (IR 64, Swarna, Nagina 22, and Vandana), a variety of *Oryza sativa* ssp. *japonica* (Nipponbare), African cultivated rice *Oryza glaberrima*, and three wild rice species, *Oryza rufipogon*, *Oryza latifolia*, and *Oryza australiensis* were used for the study (Supplementary Table S1). Seeds were germinated on germination paper for seven days at room temperature and then transferred to Hoagland solution in test tubes for two weeks of hydroponic growth. Next, fifteen to twenty individual plants of each genotype were transplanted in the research field of the National Institute of Plant Genome Research, New Delhi (average air temperature > 25 °C, 70-80% humidity, and more than 100 cm annual rainfall). In addition, five plants of each genotype were transplanted in the plastic pots, one plant per pot (15 cm diameter × 14 cm height), and were grown in randomized complete block design under controlled growth chambers with re-randomization every seventh day to control the confounder. Plants were supplied with the same amount of Hoagland solution weekly. The growth conditions were maintained as 28/22°C day and night air temperature, 70-80% relative humidity, a photoperiodic cycle of 14h:10h (light:dark), and 1000 $\mu\text{mol m}^{-2}\text{s}^{-1}$ light intensity. The transition from dark to day light and temperature conditions was incremental. The supply of nutrients, including nitrogen, was controlled for each genotype in both field and controlled growth chamber experiments.

Photosynthesis rate and related physiological trait measurements

Both field-grown plants, as well as chamber-grown plants, were used for detailed quantification of photosynthesis rate and related physiological traits. Leaf photosynthesis of field-grown plants was measured at ten days after heading from the middle widest portion of fully expanded flag leaves using LI-6400XT portable photosynthesis system using 2×3 cm standard leaf chamber (LI-6400, LI-COR, Inc). For the accessions with leaf width less than 2 cm, the leaf chamber was filled using two flag leaves carefully without any overlap (Busch, 2018). For all measurements, the leaf chamber was maintained at ambient atmospheric CO_2 concentration (C_a , $400 \mu\text{mol mol}^{-1}$), natural light irradiance ($1400\text{-}1500 \mu\text{mol m}^{-2}\text{s}^{-1}$), constant airflow ($300 \mu\text{mol s}^{-1}$), 70-80% relative humidity, leaf and air temperature ($28\text{-}30^\circ\text{C}$), and leaf-to-air maximum vapor pressure deficit in the range of $1.0\text{-}1.5$ kPa. Gas exchange measurements were performed between 9.00 h to 11.00 h on clear sunny days. Fifteen flag leaves from different plants were used to quantify net photosynthesis rate (A), intercellular CO_2 concentration (C_i), stomatal conductance to water (g_s), and carboxylation efficiency (CE , the ratio of A to C_i). Values were recorded when A , g_s , and C_i became stable.

A , g_s , C_i and CE were also quantified from the middle widest part of fully-grown eighth leaves of chamber-grown plants using a 2 cm^2 leaf chamber with a LED light source (LI-6400-02B; LI-COR, Inc). Leaf chamber was maintained at $1500 \mu\text{mol m}^{-2}\text{s}^{-1}$ light intensity, $400 \mu\text{mol mol}^{-1}$ CO_2 concentration, $300 \mu\text{mol s}^{-1}$ constant airflow, 70-80% relative humidity, leaf and air temperature of 28 and 30°C , respectively, and leaf-to-air maximum vapor pressure deficit in the range of $1.0\text{-}1.5$ kPa.

Estimation of leaf chlorophyll content

Total chlorophyll content (TCC) was measured in aqueous 80% acetone following the procedure described in Porra *et al.* (1989). Briefly, flag leaves (2.0×1.0 cm) of field-grown plants and fully-grown eighth leaves of chamber-grown plants (five biological replicates per genotypes) were harvested and incubated in 5 ml of ice-cold 80% acetone for 48 h, centrifuged at $5,000 \times g$ for 5 min, and absorbance was recorded at 663 nm (Chl a) and 645 nm (Chl b). The pigment concentration (Chl a + b) was calculated using the method described by Arnon, (1949) and expressed in $\mu\text{mol m}^{-2}$.

Quantification of leaf photosynthesis (A) and stomatal conductance (g_s) at different light intensities

Plants grown under controlled growth conditions were used to examine the effect of light intensity on A and g_s . The measurements were performed on the same middle widest part of fully-grown eighth leaves of chamber-grown plants (five biological replicates per genotypes) using a 2 cm² standard leaf chamber with a LED light source (LI-6400-02B; LI-COR, Inc). The leaf chamber was maintained at a constant temperature and relative humidity for all measurements as explained above. The CO₂ concentration (C_a) was set at 400 μmol mol⁻¹. Leaves were initially stabilized at the highest irradiance (i.e. 2000 μmol m⁻² s⁻¹), and then PPFD was gradually decreased in a step-wise manner (1800, 1500, 1200, 1000, 700, 500, 300, 100 μmol m⁻² s⁻¹). Data were recorded at each point once A and g_s values were stable.

Quantification of A , g_s , and chlorophyll fluorescence at different CO₂ concentrations

The effect of varying CO₂ concentration on A and g_s , as well as on leaf chlorophyll fluorescence, was quantified simultaneously using plants grown under controlled growth conditions. Values were recorded from the same middle widest part of fully-grown eighth leaves of chamber-grown plants (five biological replicates per genotypes) under 1500 μmol m⁻² s⁻¹ light conditions intensity using 2 cm² standard leaf chamber fluorometer (LI-6400-40 LI-COR, Inc) according to the method described in Li *et al.* (2009) and Fabre *et al.* (2019). Before measurements, leaves were stabilized at ambient CO₂ concentration (400 μmol mol⁻¹) and saturating light intensity for at least 10 minutes. The concentration of CO₂ in the leaf chamber (C_a) was decreased from 400 μmol mol⁻¹ concentration to 300, 200, 150, 100, 80, μmol mol⁻¹ CO₂, and then increased in a step-wise manner to 400, 600, 800, 1000, 1200, and 1400 μmol mol⁻¹ CO₂. The values of C_i at each CO₂ concentration for the selected accessions are shown in Fig. S4B. The CO₂ and H₂O diffusion through the gasket was corrected according to the manufacturer's instructions (LI-COR 6400XT manual version 6). The leaf and air temperature, relative humidity, airflow, and vapor pressure deficit were maintained as mentioned above.

For photochemical traits, steady-state fluorescence (F_s) was quantified at different CO₂ concentrations. Light-adapted maximum fluorescence (F_m') was determined at each CO₂ concentration by illuminating 2.0 cm² leaf surface inside the chamber with a saturated pulse of light (about 8000 μmol m⁻² s⁻¹ for 0.8 sec), provided by an array of light-emitting diodes (a mixture of 90% red and 10% blue lights at 640 and 460 nm peak wavelength, respectively). Minimal fluorescence level in the light-adapted state (F_o') was determined after removal of the actinic light, and illuminating the leaf with 1 sec of far-red light (735 nm). Using F_s , F_m' and F_o' , we calculated (a) the efficiency of PSII (Φ_{PSII}) (b) the coefficient of photochemical

quenching (qP), and (c) the photosynthetic electron transport rate (J_{PSII}) in a light-adapted state using following formulae as described in Murchie and Lawson, (2013).

$$\square PSII = (F_m' - F_s) / F_m'$$

$$qP = (F_m' - F_s) / (F_m' - F_o')$$

$$J_{PSII} = \alpha \times \beta \times \square PSII \times PPFD$$

where PPFD is incident photosynthetic photon flux density (i.e., $1500 \mu\text{mol m}^{-2} \text{s}^{-1}$), β is the photosystem partitioning factor between PSI and PSII for the distribution of light (0.5), and α is the leaf absorbance calculated from total chlorophyll content per unit leaf area (TCC , $\mu\text{mol m}^{-2}$) for each rice genotype. TCC was quantified from fully grown leaves of growth-chamber grown plants at the same stage when photosynthesis was measured (Supplementary Table S2). α was calculated from TCC using the following formula as described in Evans and Poorter, (2001):

$$\alpha = [TCC] / ([TCC] + 76)$$

Quantification of mitochondrial respiration rate in the light (R_d) and the CO_2 compensation point related to C_i (Γ^*)

R_d and Γ^* were quantified at the intersection point of multiple A/C_i response curves, conducted at a series of CO_2 concentrations and multiple light intensities using plants grown under controlled growth conditions (Laisk, 1977; Li *et al.*, 2009). A was recorded from the same middle widest part of fully-grown eighth leaves of chamber-grown plants using a 2 cm^2 standard leaf chamber fluorometer (LI-6400-40 LI-COR, Inc). Light intensity inside the leaf chamber was controlled as a series of 100, 200, and $500 \mu\text{mol m}^{-2} \text{s}^{-1}$. At each light intensity, CO_2 concentration was adjusted in the series of 75, 100, 125, and $150 \mu\text{mol mol}^{-1}$ inside the chamber. The leaf and air temperature, relative humidity, airflow, and vapor pressure deficit were maintained as explained above. CO_2 and H_2O diffusion through the gasket was corrected according to the manufacturer's instructions (LI-COR 6400XT manual version 6). The value of A at the point of intersection of the three A/C_i curves represented $-R_d$, and C_i at the point represented Γ^* (Li *et al.*, 2009). The representative graphs for the quantification of R_d and Γ^* of one plant of each genotype are shown in Supplementary Fig. S5. The average values of R_d and Γ^* were obtained from the values of three individual plants of each genotype.

Quantification of mesophyll conductance (g_m)

Mesophyll conductance (g_m) at ambient CO₂ concentration (400 μmol mol⁻¹) was quantified using the formula $g_m = A/(C_i - C_c)$, where C_c is the chloroplastic CO₂ concentration. C_c was calculated according to the variable J method (Harley *et al.*, 1992) as:

$$C_c = \frac{\Gamma^* \times [J + 8(A + R_d)]}{J - 4(A + R_d)}$$

where Γ^* is CO₂ compensation point (μmol mol⁻¹), R_d is mitochondrial respiration rate in the light (μmol m⁻² s⁻¹), and J is the rate of photosynthetic electron transport, which was estimated using chlorophyll fluorescence analysis as J_{PSII} (described above). Genotype-specific values of variables were used for the calculation of g_m for each genotype.

Fitting the FvCB model to estimate biochemical limitations

We used the non-linear fitting method explained by Sharkey (2016), version 2 to fit the FvCB model. Three major photosynthetic limitations, the maximum velocity of Rubisco carboxylation (V_{cmax}), maximum electron transport rate for RuBP regeneration (J_{max}), and maximum rate of triose phosphate utilization (TPU) were estimated simultaneously by minimizing the sum of squares of the residuals. Genotype-specific values of g_m , R_d , and Γ^* (as explained above) were used for fitting the model. Rice Rubisco catalytic constants were taken from Perdomo *et al.* (2017). All parameters were scaled to a constant temperature of 25°C. Before fitting the curves, values for each curve were corrected for diffusive leaks between the cuvette and the external environment (Bernacchi *et al.*, 2001). Rubisco limitation (A_c), RuBP-regeneration limitation (A_j), and TPU limitation (A_t) to photosynthesis for different accessions were estimated using the following formulae:

$$A_c = V_{cmax} \left[\frac{C_c - \Gamma^*}{C_c + K_c \left(1 + \frac{O}{K_o}\right)} \right] - R_d$$

$$A_j = J_{max} \frac{C_c - \Gamma^*}{4C_c + 8\Gamma^*} - R_d$$

$$A_t = 3TPU - R_d$$

where O is partial pressure for oxygen, and K_c and K_o are Rubisco catalytic constants.

Rubisco activity assay

Rubisco activity assay was performed following the procedure described in Kubien *et al.* (2011) and Sharwood *et al.* (2016). Leaf tissues (3.6 × 1.0 cm) were excised from the same middle widest part of the same leaves of field-grown plants, from where photosynthesis data

were recorded, and homogenized in extraction buffer [EB, 5 mM DTT, 10 mg ml⁻¹ PVPP, 2 mg ml⁻¹ BSA, 2 mg ml⁻¹ PEG, 2% (v/v) Tween-80, 12 mM 6-aminocaproic acid, 2.4 mM benzamidine in HEPES stock) at 4 °C. The homogenate was centrifuged for 1 min at ~16,000 × g and the pellet was discarded. To activate rubisco enzyme, leaf extracts were incubated with BMEC (100 mM Bicine-NaOH, pH 8.2, 20 mM MgCl₂, 1 mM Na₂-EDTA and 10 mM NaHCO₃) in a 9:1 ratio (e.g. 900 µL leaf extract to 100 µL BMEC) at 25 °C for 25 min. To assess the activity of Rubisco from leaf extract, 50 µl of leaf extract was incubated with 1,550 µl of the EME buffer, 40 µl 10 mM of NADH (final concentration 200 µM), 80 µl of 250 mM NaH¹²CO₃ (10 mM), 200 µl of ATP/phosphocreatine (1 and 5 mM, respectively), 40 µl of coupling enzymes (Carbonic anhydrase 500U, Glyceraldehyde-3-P dehydrogenase 50U, 3-PGA kinase 50U, Triose-P Isomerase/Glycerol-3-P dehydrogenase 400/40U), 40 µl of 25 mM RuBP (500 µM), mixed gently, and decrease of absorbance were monitored at 340 nm continuously for 0 to 1800 sec using a UV–Vis spectrophotometer.

The Rubisco carboxylation rate was determined using the equation:

$$RuBP - use (\mu mol s^{-1}) = \frac{\Delta OD_{340} * Vol}{6.18 \times 4}$$

assuming an extinction coefficient of 6.18 µmol ml⁻¹ cm⁻¹ for NADH oxidation at 340 nm. ΔOD₃₄₀ is the rate of change of absorbance at 340 nm per second, and Vol is assay volume (1 ml). The equation accounts for the four NADH molecules oxidized per RuBP carboxylated by Rubisco.

The activity of Rubisco (V_{max}) is calculated as:

$$V_{max} (\mu mol m^{-2} s^{-1}) = \frac{(RuBP - use; \mu mol s^{-1})(EV; \mu l)(10000 cm^2/m^2)}{(LA; cm^2) (Sample volume; \mu l)}$$

where EV is extraction volume and LA is leaf area. All the chemicals used for the Rubisco activity assay, including RuBP, were commercially procured from Sigma Aldrich (Kubien *et al.*, 2011; Sharwood *et al.*, 2016).

Quantification of leaf nitrogen content

Nitrogen content in leaf tissues was calculated by excising 2.0 × 1.0 cm leaf area from the same portion of the leaves where Rubisco activity was performed. The fresh weight of the leaf tissues was determined followed by drying of the tissues for three days at 60 °C. The dried tissues were weighed and ground to a fine powder. The leaf mass per area (LMA) was calculated by dividing the leaf dry weight by leaf area. The nitrogen content of leaves per unit

dry weight (N_{mass}) was then quantified using a Kjeldahl auto-analyzer (Technicon, Tarrytown, New York) as described in Traw and Ackerly (1995). The leaf nitrogen content per unit area (N_{area}) was derived from N_{mass} and LMA (Pinto *et al.*, 2016).

RNA isolation and expression analysis

Total RNA was isolated from flag leaves in three biological replicates for each genotype, and each biological replicate consisted of tissues from multiple individual plants. Tissues were harvested and immediately frozen in liquid nitrogen. Total RNA extraction (TRIzol reagent), DNase treatment, and cDNA synthesis (RevertAid First-strand cDNA synthesis kit) were performed according to the manufacturer protocol (Thermo Fisher Scientific). The PowerUp™ SYBR® Green fluorescence dye was used for quantitative gene expression analysis on CFX connect Real-time PCR detection system (Bio-Rad). The details of genes used for expression analysis and the primers are presented in Supplementary Table S3 and S4. The specificity of amplified gene products across the selected rice genotypes using the listed primers was viewed through dissociation curve analysis. Before quantitative real-time PCR, the efficiency of each primers pair was evaluated using a standard curve method where four cDNA quantities (1, 10, 50, and 100 ng) from one of the cultivated rice (Nipponbare) was used to construct a standard curve, and efficiency was calculated for each genotype using the formula $E = (10^{[-1/\text{slope}]} - 1) * 100$. Expression analysis of each gene was performed in three biological replicates for each genotype and repeated twice for the reproducibility of the results. We normalized the transcript levels of genes independently to two stable internal controls actin (LOC_Os03g50885.1) and ubiquitin (LOC_Os03g03920) with similar results. Relative expressions of the target genes are presented using both actin and ubiquitin as the internal control following the method by Livak and Schmittgen (2001) (Fig. 4, Supplementary Fig. S9).

Visualization and quantification of leaf developmental traits

The widest middle part of the flag leaves of 10 -15 field-grown plants was used to measure flag leaf width (FLW) and leaf thickness (FLT). For anatomical traits visualization, transverse sections were cut from the mid-portion of flag leaves (one leaf per plant, five biological replicates per genotype), and stained with toluidine blue with some modification of the method as described in Lux *et al.* (2005). Briefly, chlorophyll pigments of the thin sections were cleared using 85 % (w/v) lactic acid saturated with chloral hydrate, heated on a water bath at 70 °C for 1 h, and washed several times with distilled water. To visualize

mesophyll cells, sections were incubated in absolute alcohol overnight and stained with 1% toluidine blue in 1% (w/v) disodium tetraborate for 15 sec. Finally, thin sections were mounted with 50 % glycerol and visualized using a bright field microscope (LMI, UK). The quantification of leaf anatomical features related to the minor vein, mesophyll cell and, bundle sheath cells were performed under a light microscope at 10X and 100X magnification (one leaf per plant, four images per leaf, two on each side of the midrib that consisted of about 3-6 complete cells per image, five biological replicates per genotype) using Fiji-Image J software (Wayne Rasband, National Institute of Health) as described in Chatterjee *et al.* (2016) and Ivanova *et al.* (2018). The mesophyll cell number per unit leaf area, mesophyll cell area, and the mesophyll surface area exposed to the intercellular space (S_{mes}) were calculated from transverse sections under bright field microscopy as described in Scafaro *et al.* (2011). Total 30 to 40 mesophyll cells per genotype were quantified for the calculation of S_{mes} .

Chloroplast ultrastructure

Freshly harvested flag leaf segments (1 × 1 mm; one leaf each from three individual plants per genotype) were fixed under buffer (3% glutaraldehyde in 0.1 M cacodylate buffer, pH 7.2) for 2 h, and then treated with 1% osmium tetroxide overnight at 4 °C. The fixed segments were dehydrated in a graded acetone series and embedded in epoxy resin. Ultrathin sections were cut with diamond knives, stained with uranyl acetate and lead citrate double staining procedure as described by Reynolds, (1963). Chloroplasts of the uppermost part of the leaf sections were viewed using a digital camera (BH-2, Olympus) equipped with transmission electron microscopy at 1000X magnification (JEM 1230; JEOL). The transverse sections prepared for electron microscopy were used to calculate chloroplast area, chloroplast number per unit mesophyll cell area, and the surface area of the chloroplast exposed to intercellular space (S_c) as described in Scafaro *et al.* (2011). 15 to 20 mesophyll cells with clear images of chloroplast cells were quantified for the calculation of S_c .

Statistical analysis

Differences among the genotypes were tested by one-way analysis of variance (ANOVA) (JMP Pro, version 12.0.1; SAS Institute, Cary, NC, USA). Statistically similar genotypes were grouped according to post-hoc Tukey HSD calculation. Significance test for the genotypes at varying light intensities and CO₂ concentrations, as well as for Rubisco activity differences was performed using one-way Analysis of Variance (ANOVA). Pearson

correlation matrix was calculated based on mean values of each trait for each genotype to evaluate the trait-to-trait associations. The threshold for statistical significance was $P \leq 0.05$. Linear regression analyses between V_{cmax}/J_{max} , V_{cmax}/TPU , and J_{max}/TPU were performed using Microsoft Excel.

Results

Quantification of leaf photosynthesis and related physiological features

We observed remarkable differences in leaf photosynthesis (A) and related physiological features among the selected cultivated and wild rice accessions (Fig. 1). The mean A value of the wild rice species, *O. latifolia*, *O. australiensis*, and *O. rufipogon*, and African cultivated rice *O. glaberrima* were found to be significantly higher than that of cultivated rice varieties, such as IR64 and Nipponbare (Fig. 1A). *O. australiensis* and *O. latifolia* showed the highest A , 23.5 and 24.2 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, respectively, which were 29% to 57% higher compared with the different Asian cultivated varieties. The African cultivated rice *O. glaberrima* showed A value intermediate between the wild accessions and Asian cultivated varieties. We also observed significant variations for A among the cultivated varieties. For instance, one of the high-yielding *indica* variety Swarna and *japonica* variety Nipponbare displayed 19% and 14% higher A , respectively, compared with IR64 (Fig. 1A).

We further examined important physiological traits of the selected accessions that are known to influence leaf photosynthesis. As expected, wild rice accessions of *O. latifolia* and *O. australiensis* with the highest A showed higher stomatal conductance (g_s), and intercellular CO_2 concentration (C_i) (Fig. 1B, C). For example, *O. latifolia* and *O. australiensis* accessions showed 1.3 to 3-fold higher g_s compared with different Asian cultivated varieties. In addition, these species along with *O. rufipogon* displayed higher carboxylation efficiency (CE) than those of cultivated varieties (Fig. 1D). Among the cultivated varieties, Nipponbare that displayed higher A also showed reasonably higher g_s and C_i compared with the other cultivated varieties. We also observed significantly higher total chlorophyll content (TCC) in flag leaves of the selected wild rice accessions compared with the cultivated varieties. Consistent with the high A , wild accessions of *O. latifolia* and *O. australiensis* had 33% and 43% higher TCC per unit leaf area, respectively, than IR64. *O. glaberrima*, Swarna, and Nipponbare also had reasonably higher TCC than IR 64, Nagina 22, and Vandana (Fig. 1E). Pearson correlation analysis showed a significant positive correlation of A with g_s ($r = 0.629$), C_i ($r = 0.570$), CE ($r = 0.819$), and TCC ($r = 0.912$), g_s with C_i ($r = 0.965$) and CE with TCC ($r = 0.906$) at $P \leq 0.05$ (Supplementary Fig. S1).

We then selected representative cultivated rice accessions (*O. sativa* ssp. *indica* cv. IR 64, *O. sativa* ssp. *japonica* cv. Nipponbare, and African cultivated rice *O. glaberrima*) and wild rice accessions (*O. latifolia* and *O. australiensis*), which captured the entire variation range for the leaf photosynthesis per unit area in the field experiment, for the quantification of leaf photosynthesis and related physiological traits at the vegetative stage under controlled conditions (Fig. 1F; Supplementary Fig. S2). Similar to our observations at the full heading stage under field conditions, the selected wild rice accessions together with African cultivated rice *O. glaberrima* showed significantly higher A , g_s , and CE compared to cultivated rice varieties, IR64 and Nipponbare, at the vegetative stage (Table 1). Moreover, we have recently reported consistently higher photosynthesis per unit leaf area in the *O. australiensis* accession compared with Nipponbare at different developmental stages in field conditions (Mathan *et al.* 2021). Taken together, these results showed significant differences in photosynthetic traits across the selected cultivated and wild rice accessions.

Next, we examined the effect of different light intensities and CO_2 concentrations on A and g_s of the selected wild and cultivated rice accessions grown under controlled conditions (Fig. S3). *O. latifolia* and *O. australiensis* accessions exhibited higher increases in A in response to increasing PPFD and maintained higher g_s at different light intensities compared with the cultivated varieties (Supplementary Fig. S3A and B). *O. latifolia* and *O. australiensis* had significantly higher light-saturated A_{max} , 20.94 and 21.68 $\mu mol CO_2 m^{-2} s^{-1}$, respectively than *O. sativa* cv. IR 64 and *O. sativa* cv. Nipponbare (11.89 and 13.02 $\mu mol CO_2 m^{-2} s^{-1}$, respectively). Since the differences in the C_i values at different PPFD may confound the photosynthesis rate in different accessions, we quantified the C_i values at each PPFD for the selected accessions. Changes in C_i in response to increasing PPFD during the light response curves were mostly similar across all the genotypes (Fig. S4). The effect of increasing CO_2 concentration on A segregated the selected genotypes under three distinct groups: *O. australiensis* and *O. latifolia* with higher A , African cultivated rice *O. glaberrima* with intermediate A , and cultivated varieties Nipponbare and IR 64 with lower A at different CO_2 concentration (Fig. S3C). The g_s , however, showed a decreasing trend at increasing CO_2 concentration, likely due to the closure of stomata (Fig. S3D). Altogether, the selected wild rice accessions of *O. australiensis* and *O. latifolia* were able to maintain higher leaf photosynthesis per unit area at different light intensities and CO_2 concentrations compared with the selected cultivated varieties.

Fitting CO_2 response curves to investigate biochemical limitations V_{max} , J_{max} , and TPU

To investigate the biochemical limitations to photosynthesis across the selected accessions using the FvCB model, we first estimated mesophyll conductance (g_m) at ambient CO_2 concentration. g_m significantly varied for the selected cultivated and wild rice species (Table 2). *O. australiensis* and *O. latifolia*, with highest A , also showed higher g_m than the cultivated varieties, suggesting the key importance of g_m for photosynthetic variations among the genotypes. We also quantified the mitochondrial respiration rate in the light (R_d) and CO_2 compensation point related to C_i (Γ^*) for each genotype to fit the FvCB model (Table 2 and Supplementary Fig. S5). R_d was found to be significantly higher for *O. australiensis* and *O. latifolia* compared to the selected cultivated accessions, whereas no significant difference was observed for Γ^* across the accessions.

We, then, used carbon dioxide response (A/C_i) curves to estimate maxima of Rubisco carboxylation activity (V_{cmax}), regeneration of ribulose-1,5-biphosphate expressed as electron transport rate (J_{max}), and triose phosphate utilization (TPU) for the selected accessions (Farquhar *et al.*, 1980; von Caemmerer, 2000) (Supplementary Fig. S6). CO_2 -saturated A_{max} for the selected accessions ranged from $21.37 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ for cultivated variety IR 64 to $36.81 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ for the wild species *O. australiensis* (Table 3). We found significantly higher values V_{cmax} , J_{max} , and TPU for the wild species *O. australiensis* and *O. latifolia* compared to cultivated varieties IR64 and Nipponbare, suggesting V_{cmax} , J_{max} , and TPU together might contribute to differences in A_{max} . Consistent with the intermediate value of A_{max} for *O. glaberrima*, the values of biochemical limitations were also intermediate between the wild species and cultivated *indica* and *japonica* varieties. We investigated the relationship among the modelled biochemical limitations to photosynthesis across the selected accessions. A distinct linear relationship was observed between V_{cmax} and J_{max} ($R^2 = 0.973$), J_{max} and TPU ($R^2 = 0.987$), and V_{cmax} and TPU ($R^2 = 0.968$) at $P < 0.001$, suggesting the robustness of the modelled limitations (Supplementary Fig. S7). These results, expectedly, indicated the important contribution of biochemical attributes to photosynthetic differences among the selected cultivated and wild rice accessions.

Next, we calculated the photosynthetic limitations due to V_{cmax} (A_c), J_{max} (A_j), and TPU (A_t). The transition from Rubisco-limited (A_c) to RuBP-regeneration limited (A_j) photosynthesis in Asian cultivated rice, IR 64 and Nipponbare, was observed at $C_i < 250 \mu\text{mol mol}^{-1}$ (Supplementary Table S5). In contrast, wild rice species *O. australiensis* and *O. latifolia* along with African cultivated rice *O. glaberrima* showed A_c to A_j transition at $C_i > 300 \mu\text{mol mol}^{-1}$. Similarly, the transition from RuBP-regeneration limited (A_j) to TPU-limited (A_t) photosynthesis in Nipponbare and IR 64 was observed at C_i 524 and $608 \mu\text{mol mol}^{-1}$,

respectively, which was lesser compared to *O. glaberrima* (657 $\mu\text{mol mol}^{-1}$), *O. australiensis* (700 $\mu\text{mol mol}^{-1}$), and *O. latifolia* (713 $\mu\text{mol mol}^{-1}$) (Supplementary Table S5). CO_2 -saturated A_{max} that ranged from 21.37 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in IR 64 to 36.81 $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ in *O. australiensis*, further, indicated TPU-limitation (A_t) in IR64 and Nipponbare compared with the wild rice accessions. However, the TPU-limitation is observed at C_i much higher than the physiological conditions as reported earlier (Kumarathunge *et al.*, 2019). These results suggested that leaf photosynthesis per unit area in Asian cultivated varieties, IR 64 and Nipponbare, was limited due to Rubisco activity (V_{cmax}) and RuBP-regeneration/Electron Transport Rate (J_{max}) compared to the selected wild rice genotypes under ambient CO_2 concentrations.

Photochemical differences among the selected rice accessions

Since the estimation of biochemical limitations to photosynthetic differences across the selected accessions suggested J_{max} as one of the limiting factors, we used leaf chlorophyll fluorescence parameters to quantify the photochemical features of the selected rice accessions. We observed strong differences in electron transport rate (J_{PSII}), the efficiency of PSII ($\square\text{PSII}$), and photochemical quenching (qP) among the selected cultivated and wild rice accessions across different CO_2 concentrations, with distinct differences at CO_2 concentration higher than 200 $\mu\text{mol mol}^{-1}$. *O. latifolia* and *O. australiensis* showed higher values of J_{PSII} than the cultivated Asian rice varieties IR 64 and Nipponbare at different CO_2 concentrations, with 19% to 53% higher J_{max} in the two wild rice accession compared with the two cultivated varieties (Fig. 2A). Similar to J_{PSII} , values of $\square\text{PSII}$ and qP were also significantly higher for *O. latifolia* and *O. australiensis* compared with the cultivated varieties (Fig. 2B, C). *O. glaberrima*, the African cultivated rice with reasonably higher A , also showed relatively higher J_{PSII} , $\square\text{PSII}$, and qP than *O. sativa* cultivated varieties at different CO_2 concentrations. Similar response patterns of both A and J_{PSII} were observed at different CO_2 concentrations, and were found to be strongly correlated with each other for all the selected genotypes (Supplementary Fig. S8). Taken together, these results showed that photochemical features of *O. australiensis* and *O. latifolia* contributed to the higher leaf photosynthesis rate, and lower J_{PSII} , $\square\text{PSII}$, and qP of IR 64 and Nipponbare accounted for RuBP-regeneration-limited photosynthesis in those cultivated varieties.

Rubisco activity differences across the selected rice genotypes

Our analyses on biochemical limitations to photosynthesis suggested V_{cmax} as a major limitation to leaf photosynthesis per unit area in the cultivated varieties IR64 and Nipponbare. Therefore, we quantified the *RuBP-use* and *in vitro* Rubisco activity (V_{cmax}) over 30 minutes across selected cultivated and wild rice species (Fig. 3A, B). The CO_2 -saturated *RuBP-use* gradually increased immediately from 10 sec to 10 min, followed by saturation for all the accessions. The *RuBP-use* was significantly higher in photosynthetically efficient wild rice accessions (*O. latifolia* and *O. australiensis*) than that of Asian cultivated varieties (*O. sativa* cv. IR 64, and *O. sativa* cv. Nipponbare) (Fig. 3A). Consistent with the *RuBP-use*, V_{cmax} was maximum for the wild rice *O. australiensis* and minimum for the cultivated variety IR64 at saturation (Fig. 3B). Table 4 shows significantly higher values of *RuBP-use* per sec and V_{cmax} for *O. latifolia* and *O. australiensis* compared to IR 64 and Nipponbare at different time points during the assay. The results not only supported the estimated photosynthetic limitations due to V_{cmax} in the selected cultivated varieties but also suggested the existence of variation in Rubisco carboxylation activity per unit leaf area within the rice system.

The differences in Rubisco activity per unit leaf area across the selected accessions may be attributed either to Rubisco amount or to enzyme kinetics differences. Since the leaf nitrogen content would inform about Rubisco amount per unit leaf area, we quantified the leaf nitrogen content of the selected accessions. Photosynthetically efficient *O. australiensis* and *O. latifolia* accessions showed significantly higher nitrogen content per unit leaf area (N_{area}) compared with IR64 and Nipponbare (Fig. 3C). The selected *O. australiensis* accession had 22% and 14% higher N_{area} , whereas *O. latifolia* accession had 12% and 5% higher N_{area} than IR64 and Nipponbare, respectively, suggesting an important contribution of leaf nitrogen content to higher Rubisco activity per unit area in the selected wild accessions.

Expression pattern of genes encoding core photosynthetic and chlorophyll biosynthesis enzymes

Since the differences in the Rubisco amount per unit leaf area may also contribute to the Rubisco-limitation of the photosynthesis in IR 64 and Nipponbare, we checked the expression pattern genes encoding Rubisco small and large subunits. In addition, we checked the expression of LHC1 (Light-harvesting chlorophyll protein complex) and TPT (Triose phosphate translocator) that may contribute to RuBP-regeneration- and TPU- limitations of the photosynthesis, respectively (Supplementary Table S3). The expression of the selected photosynthetic genes was significantly higher in *O. latifolia* and *O. australiensis* compared to the selected cultivated varieties (Fig. 4A, Supplementary Fig. S9A), suggesting higher

photosynthesis in those species could, at least in part, be due to the higher abundance of these gene products. These genes could increase net photosynthetic efficiency by increasing the efficiency of light capture, CO₂ assimilation, and transport of triose phosphate from chloroplast to cytoplasm. Since wild rice species with higher photosynthesis also showed higher chlorophyll content, we checked the expression level of chlorophyll biosynthesis genes. The expression of chlorophyll biosynthesis genes, such as *GluTR*, *PPOX*, *Mg-CHLD*, *CHLSYN*, and *CAO* were found to be significantly higher in *O. latifolia* and *O. australiensis* compared to the selected cultivated varieties (Fig. 4B, Supplementary Fig. S9B).

Leaf developmental traits underpinning photosynthetic variations

Leaves are the prime photosynthetic organs, and expectedly the selected cultivated and wild rice accessions exhibited considerable variation for flag leaf width (FLW) and thickness (FLT) (Supplementary Fig. S10). Wild rice species had thicker and wider leaves as compared to cultivated rice. Strong significant positive correlations of leaf photosynthesis were found with FLW ($r = 0.850$) and FLT ($r = 0.938$) (Fig. 5A). At the anatomical level, remarkable differences in the vascular architecture and mesophyll cell features were observed for the selected rice accessions that differ in leaf photosynthesis (Supplementary Table S6). Wild rice species with higher *A* exhibited a larger vein height and width. *O. sativa* cv. Nipponbare that showed relatively higher *A* compared to other cultivated varieties also showed larger vein height and width (Supplementary Table S6). Significant positive correlations were observed for *A* with vein features: minor vein width (MiV-W, $r = 0.752$), minor vein height (MiV-H, $r = 0.899$), and total number of veins (TV, $r = 0.848$) (Fig. 5A). Similar strong correlations of *A* were observed for mesophyll cell features: mesophyll cell lobbing (M-Lo, $r = 0.760$), mesophyll cell length (M-L, $r = 0.863$), mesophyll cell width (M-W, $r = 0.928$), and mesophyll cell area (M-A, $r = 0.897$) at $P \leq 0.05$. However, no significant correlations of *A* and related physiological traits were observed with bundle sheath features: bundle-sheath number (BS-No), and bundle-sheath area (BS-A) (Fig. 5A).

Photosynthetically efficient wild rice species, *O. latifolia*, and *O. australiensis*, had larger and closely spaced veins with a fewer number of mesophyll cells between two consecutive veins (Fig. 5B). Those wild species also had larger mesophyll cell size, and thus a lesser number of mesophyll cells per unit leaf area (Fig. 5C, D). We, then, quantified attributes related to the mesophyll cell surface area of the selected accessions. Wild rice accessions of *O. australiensis* and *O. latifolia*, showed remarkably higher average mesophyll cell area (2.2 to 2.6-fold larger mesophyll cells) as well as significantly higher mesophyll cell

surface area exposed to intercellular space (S_{mes}) compared with IR64 and Nipponbare (Fig. 5E, F). We also investigated the organization of chloroplasts in mesophyll cells of the representative wild and cultivated accessions. Photosynthetically efficient wild rice exhibited larger and numerous chloroplasts, which were systematically distributed along the mesophyll cell walls (Fig. 5G, H; Supplementary Fig. S11). The surface area of chloroplasts exposed to intercellular space (S_c) was also found to be significantly higher in the photosynthetically efficient wild rice accessions compared with the cultivated varieties IR64 and Nipponbare (Fig. 5I). These results underline the strong influence of mesophyll and chloroplast features in driving high leaf photosynthesis per unit area in the selected wild rice accessions.

Discussion

The importance of investigating variations in leaf photosynthesis within and across species to be exploited for crop improvement has been emphasized (Ort *et al.*, 2015; Adachi *et al.* 2019; Simkin *et al.*, 2019; van Bezouw *et al.*, 2019). Exploiting the photosynthetic variations within the rice system itself, including cultivated and wild rice, could be one of the potential strategies to increase genetic yield potential. In this study, we not only quantified leaf photosynthetic differences across the selected cultivated and wild rice accessions but also identified the biochemical limitations to leaf photosynthesis per unit area in the selected cultivated varieties, IR64 and Nipponbare. In addition, we also identified photochemical and leaf developmental traits associated with higher leaf photosynthesis rates in the selected wild rice accessions of *O. australiensis* and *O. latifolia*.

High leaf photosynthesis rate in the selected wild rice accessions, and related physiological traits

Wild relatives of rice are shown to have more genetic variation compared with cultivated varieties as domestication was primarily targeted for optimal plant architectural and grain features (Huang *et al.*, 2012; Chen *et al.*, 2021). Therefore, the wild relatives are suggested to have useful variants for various desirable agronomic traits. Indeed, wild relatives of rice have been used for breeding stress tolerance in cultivated rice varieties (Sanchez *et al.*, 2013; Atwell *et al.*, 2014; Wing *et al.*, 2018). We found higher leaf photosynthesis per unit area (A) in the selected accessions of wild rice *O. australiensis*, *O. latifolia*, and *O. rufipogon*, as well as African domesticated rice *O. glaberrima* compared with the cultivated varieties of *Oryza sativa* (Fig. 1A). A few previous studies have reported high leaf photosynthesis rate in some of the wild relatives of rice, including *O. australiensis*

and *O. rufipogon*, than cultivated species *O. sativa* (Zhao *et al.*, 2010; Kiran *et al.*, 2013; Kondamudi *et al.*, 2016; Haritha *et al.*, 2017). However, a few other studies found no obvious difference or lower *A* in wild relatives of rice compared with *O. sativa*, suggesting that high *A* is not a general feature of wild rice relatives, rather an accession-specific feature (Yeo *et al.*, 1994; Scafaro *et al.*, 2011; Giuliani *et al.*, 2013). Higher leaf photosynthesis rate could be attributed to differences in the life strategies of the selected wild rice accessions compared with the cultivated varieties. The perennial nature of the wild accessions with remarkably higher biomass accumulation than cultivated varieties might require more photosynthates, and thus increasing leaf photosynthesis rate. We have recently shown preferential utilization of a higher amount of photosynthates in the vegetative tissues of the specific *O. australiensis* accession for the synthesis of structural carbohydrates (Mathan *et al.*, 2021). Furthermore, a higher amount of photosynthates would also support more organ initiation and larger organ size, as well as the general maintenance of the higher biomass in the selected wild rice accessions. Thus, the requirement of photosynthates for general growth and biomass as well as specific biochemical, photochemical, and developmental features would determine the accession-specific value of *A*. Therefore, understanding the underlying traits that drive high *A* in specific accessions is crucial.

Consistent with leaf photosynthesis rate, photosynthetically efficient wild rice accessions of *O. australiensis* and *O. latifolia* showed higher values of stomatal conductance (g_s) and mesophyll conductance (g_m) that would facilitate higher CO_2 concentration in intercellular spaces (C_i) as well as inside mesophyll cells of those species (Fig. 1B, C; Table 2). Further, the wild rice accession also had reasonably higher *CE* than cultivated varieties. Higher leaf nitrogen content, indicating more Rubisco per unit leaf area, supports higher *CE* and *A* in the selected wild rice accessions than the cultivated varieties (Fig. 3C). The *A* has been shown to be positively associated with *CE* in rice (Yeo *et al.*, 1994). Indeed, quantification of photosynthesis and related physiological parameters for introgressions from *O. rufipogon* in a cultivated background showed the influence of both *CE* as well as g_s on higher leaf photosynthesis (Haritha *et al.*, 2017). Comparison of several cultivated and wild rice genotypes had shown a close positive relation between *A* and g_m , and g_m could increase leaf photosynthesis per unit area without affecting transpiration (Flexas *et al.*, 2008; Zhu *et al.*, 2010; Giuliani *et al.*, 2013). Therefore, efforts to increase g_s , g_m , and *CE* in the cultivated rice varieties comparable to the high photosynthetic wild accessions will help in increasing photosynthetic efficiency.

Rubisco carboxylation rate (V_{cmax}) and electron transport rate (J_{max}) promote leaf photosynthesis rate in the selected wild rice accessions

Rubisco is the rate-limiting enzyme for photoassimilation of CO₂, and Rubisco abundance and activity strongly determine the photosynthetic performance of a plant (Parry *et al.*, 2013; Sage, 2002). Higher Rubisco activity per unit leaf area in the selected wild accessions compared with the cultivated varieties could be due to either more Rubisco per unit area or the presence of an efficient Rubisco in the wild species. High photosynthetic wild accessions had higher expression of genes encoding Rubisco large subunit and small subunit, indicating the more abundance of Rubisco in those species (Fig. 4A). Thick leaves of the photosynthetically efficient wild species would support the first alternative where increasing leaf thickness would lead to more Rubisco per unit area. More Rubisco per unit area, in addition, could also be achieved by more nitrogen allocation to Rubisco. Higher nitrogen content per unit area in the leaves of the photosynthetically efficient *O. australiensis* and *O. latifolia* accessions than the cultivated varieties would also support higher Rubisco content per unit area in the wild accession (Fig. 3C). Therefore, possible ways to increase Rubisco content per unit leaf area may overcome the Rubisco limitation of leaf photosynthesis in the cultivated varieties. It would, further, be interesting to investigate if there are differences in Rubisco kinetic properties across photosynthetically contrasting cultivated and wild rice that might contribute to differences in Rubisco carboxylation activity.

The comparative higher values for the quantum efficiency of PSII (Φ_{PSII}) and photochemical quenching (qP) along with J_{PSII} in *O. australiensis* and *O. latifolia* accessions indicate more receptive PSII reaction centres for electron transport from PSII to PSI in those species, making an efficient link between light and light-independent reactions to increase A (Fig. 2A-C) (Kramer and Evans, 2011). Gene expression studies also showed higher expression of the *LHCI* gene, which is involved in the organization of reaction centre, in photosynthetically efficient wild species. Results, further, showed higher expression of chlorophyll biosynthetic genes and higher chlorophyll content in those wild species, aiding the efficient light harvesting (Fig. 1E, 4B). Understanding the differential regulation of genes encoding components of photosystem as well as chlorophyll biosynthesis in the wild species could help in improving photochemical features in the cultivated varieties.

Targeting leaf thickness and width to overcome the biochemical limitations to leaf photosynthesis in the cultivated varieties

We observed a strong positive association of A with leaf width and thickness (Fig.

5A). Leaf thickness has been suggested to influence photosynthesis through efficient absorption and utilization of the light by homogenizing the leaf internal light distribution as well as maintaining leaf temperature (Tholen *et al.*, 2012). Increasing leaf thickness, and thus more tissue per unit area would be instrumental for more nitrogen and Rubisco content, chlorophyll content, and electron transport machinery per unit area. Higher chlorophyll and electron transport machinery per unit area would facilitate efficient light harvesting for RuBP regeneration, whereas more Rubisco content per unit leaf area would increase carboxylation efficiency. Thus, targeting leaf thickness, similar to the photosynthetically efficient wild rice, may result in high net photosynthesis per unit leaf area in the cultivated varieties by overcoming Rubisco and electron transport rate limitations. Such leaves would indeed be beneficial under high light intensity, but may not be optimal for light harvesting and photosynthesis under low light conditions. Nonetheless, increasing leaf thickness, resulting in higher nitrogen and Rubisco content per unit area, could be a potential way to overcome the biochemical limitation of photosynthesis in the cultivated rice varieties under ambient light conditions. The association of leaf width with photosynthetic performance and yield in rice has been observed earlier. *WUSCHEL-RELATED HOMEODOMAIN* (*WOX*) genes *NAL1*, *NAL2*, and *NAL3* are associated with increased leaf width, which in turn results in increased photosynthetic efficiency and yield (Fujita *et al.*, 2013; Ishiwata *et al.*, 2013). Introgression of a natural variant of *NAL1* from *japonica* to *indica* cultivar led to thicker and wider leaves with higher leaf photosynthesis rate and yield (Takai *et al.*, 2013).

Distinct leaf anatomical features of the photosynthetically efficient wild rice accessions

Photosynthetically efficient wild rice accessions have mesophyll cell features that promote both higher accesses to available CO₂ as well as efficient light harvesting. Larger sized and lobed mesophyll cells along with more surface area exposed to intercellular spaces in wild rice accessions likely provide increased surface area to access more internal CO₂ (Fig. 5C-F), whereas the higher number of larger and systematically distributed chloroplasts in the mesophyll cells of those species may promote efficient light capture to enhance photosynthesis (Fig. 5G, H; S11). In addition, higher values of S_{mes} and S_c would support the higher g_m of the photosynthetically efficient wild rice accessions, leading to efficient CO₂ diffusion inside the mesophyll cells (Adachi *et al.* 2013; Pathare *et al.*, 2020). These, together with thicker leaves having more nitrogen and Rubisco content per unit area, would provide a higher amount of photosynthetic inputs as well as machinery in the photosynthetically efficient wild rice accessions. Lesser number of mesophyll cells between two consecutive

veins in photosynthetically efficient wild species, a feature similar to C_4 plants, would be promoting efficient transport of photoassimilate (Amiard *et al.*, 2005; Feldman *et al.*, 2017). Larger vein size in the wild rice species would, further, be adding to the efficient transport of photoassimilates, thus increasing the leaf photosynthesis (Sack and Scoffoni, 2013). Thus, photosynthetically efficient wild rice accessions with desirable anatomical features provide a potential genetic resource towards the identification of genes for desirable manipulation in mesophyll, vein, and chloroplast features.

In summary, we identified biochemical limitations to leaf photosynthesis per unit area in the selected cultivated rice varieties, IR64 and Nipponbare, compared to photosynthetically efficient wild rice accessions of *O. australiensis* and *O. latifolia*. Rubisco carboxylation activity and electron transport rate along with photochemical features (quantum efficiency of PSII and Photochemical quenching) and leaf developmental features (thicker and wider leaves, larger and lobbed mesophyll cells, larger veins, and lesser number of mesophyll cells between two consecutive veins) contribute to higher leaf photosynthesis in the selected *O. australiensis* and *O. latifolia* accessions (Fig. 6). The distinct leaf anatomy in the selected wild accessions would provide structure to accommodate more nitrogen and photosynthetic machinery per unit leaf area than the cultivated varieties. Similarly, higher S_{mes} and S_c would lead to more CO_2 diffusion inside the mesophyll cells and chloroplasts of those wild accessions. Together, these anatomical features would mediate higher Rubisco activity and electron transport rate driving higher leaf photosynthesis rate in the selected wild accessions. Future efforts to dissect the genetic and molecular basis of the underlying biochemical, photochemical, and developmental traits shall be instrumental in tailoring photosynthetically efficient rice towards the possible increases in genetic yield potential.

Supplementary data

Fig. S1. Pair-wise Pearson correlation coefficient (r) analysis among the leaf photosynthesis and physiological traits.

Fig. S2. Representative images of growth-chambers-grown plants of the selected cultivated and wild rice accessions.

Fig. S3. Effect of varying light intensity and CO_2 concentration on leaf photosynthesis and stomatal conductance.

Fig. S4. C_i values for light-response curves and CO_2 -response curves for each genotype.

Fig. S5. Representative graphs for the quantification of mitochondrial respiration rate in the light (R_d) and the CO_2 compensation point related to C_i (Γ^*) for each genotype.

Fig. S6. Relationship between A and C_i with A/C_i curve fitting for the selected cultivated and wild rice accessions.

Fig. S7. Linear regression analysis between V_{cmax} and J_{max} (A), TPU and J_{max} (B), and V_{cmax} and TPU (C).

Fig. S8. Response and correlation of leaf photosynthesis (A) and electron transport rate (J_{PSII}) at varying CO_2 concentrations for the selected rice accessions.

Fig. S9. Expression pattern of photosynthetic and chlorophyll biosynthetic genes across the selected accessions normalized to rice ubiquitin internal control.

Fig. S10. Flag leaf width and thickness of the selected cultivated and wild rice accessions.

Fig. S11. TEM images for the organization of chloroplasts in mesophyll cells.

Table S1. Details of rice accessions used in the present study.

Table S2. Total chlorophyll content (TCC , $\mu\text{mol m}^{-2}$) and leaf absorbance (α) of fully-grown eighth leaves of plants grown under controlled conditions.

Table S3. Details of genes used for expression analysis.

Table S4. List of primers for qRT-PCR analysis.

Table S5. A_c-A_j and A_j-A_t transitions for the selected genotypes.

Table S6. Quantification of leaf developmental traits of the fully expanded flag leaves of the selected cultivated rice varieties and wild rice species.

Acknowledgments

This work was supported by the core funding from the National Institute of Plant Genome Research as well as Ramalingaswamy Re-entry Fellowship (BT/RLF/Re-entry/05/2013) and Innovative Young Biotechnologist Award (BT/09/IYBA/2015/01) from the Department of Biotechnology, Ministry of Science and Technology, India. JM, AS, and VJ acknowledge their CSIR-JRF, SERB-NPDF, and UGC-JRF, fellowships, respectively. Seeds of wild rice species and *Oryza glaberrima* were kindly provided by Dr. Kuldeep Singh and Dr. Kumari Neelam, Punjab Agricultural University, Ludhiana, India. We thank Dr. Subodh K Sinha for his help in leaf nitrogen content estimation.

Author contributions

JM and AR conceptualized the study. JM, AS, VJ, and AR designed experiments. JM, AS, and VJ performed experiments, analysed data, and compiled figures. AS and AR wrote the manuscript. All authors contributed to and edited the final manuscript.

Data availability statement

Data sharing is not applicable to this article as all created data is already contained within this article or in the supplementary material.

References

- Acevedo** Siaca LG, Coe R, Wang Y, Kromdijk W, Quick WP, Long SP. 2020. Variation in photosynthetic induction between rice accessions and its potential for improving productivity. *New Phytologist* **227**, 1097-1108.
- Adachi S, Nakae T, Uchida M, Soda K, Takai T, Oi T, Yamamoto T, Ookawa T, Miyake H, Yano M, Hirasawa T.** 2013. The mesophyll anatomy enhancing CO₂ diffusion is a key trait for improving rice photosynthesis. *Journal of Experimental Botany* **64**, 1061-72.
- Adachi S, Yamamoto T, Nakae T, Yamashita M, Uchida M, Karimata R Ichihara N, Soda K, Ochiai T, Ao R, Otsuka C, Nakano R, Takai T, Ikka T, Kondo K, Ueda T, Ookawa T, Hirasawa T.** 2019. Genetic architecture of leaf photosynthesis in rice revealed by different types of reciprocal mapping populations. *Journal of Experimental Botany* **70**, 5131–5144.
- Amiard V, Mueh KE, Demmig-Adams B, Ebbert V, Turgeon R, Adams WW.** 2005. Anatomical and photosynthetic acclimation to the light environment in species with differing mechanisms of phloem loading. *Proceedings of the National Academy of Sciences of the USA* **102**, 12968–12973.
- Arnon DI.** 1949. Copper enzymes in isolated chloroplasts, polyphenol oxidase in *Beta vulgaris*. *Plant Physiology* **24**, 1-15.
- Atwell BJ, Wang H, Scafaro AP.** 2014. Could abiotic stress tolerance in wild relatives of rice be used to improve *Oryza sativa*? *Plant Science* **215–216**, 48–58.
- Bernacchi CJ, Singaas EL, Pimentel C, Portis Jr AR, Long SP.** 2001. Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell and Environment* **24**, 253–259.
- van Bezouw RFHM, Keurentjes JJB, Harbinson J, Aarts MGM.** 2019. Converging phenomics and genomics to study natural variation in plant photosynthetic efficiency. *Plant Journal* **97**, 112–133.
- Brodribb TJ, Feild TS, Jordan GJ.** 2007. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiology* **144**, 1890–1898.
- Busch FA.** 2018. Photosynthetic gas exchange in land plants at the leaf level. In Covshoff S ed., *Photosynthesis: Methods and Protocols, Methods in Molecular Biology*, vol. 1770, 25 - 44.
- von Caemmerer S.** 2000. *Biochemical Models of Leaf Photosynthesis*, pp. 1–165. CSIRO Publishing, Collingwood, Victoria, Australia.
- Chatterjee J, Dionora J, Elmido-Mabilangan A, Wanchana S, Thakur V, Bandyopadhyay A, Brar DS, Quick WP.** 2016. The evolutionary basis of naturally

- diverse rice leaves anatomy. *PLoS ONE* **11**, e0164532.
- Chen Q, Li W, Tan L, Tian F.** 2021. Harnessing Knowledge from Maize and Rice Domestication for New Crop Breeding. *Molecular Plant* **14**(1), 9-26.
- De Souza AP, Wang Y, Orr DJ, Carmo-Silva E, Long SP.** 2020. Photosynthesis across African cassava germplasm is limited by Rubisco and mesophyll conductance at steady state, but by stomatal conductance in fluctuating light. *New Phytologist* **225**, 2498–2513.
- Evans JR, Poorter H.** 2001. Photosynthetic acclimation of plants to growth irradiance: the relative importance of SLA and nitrogen partitioning in maximising carbon gain. *Plant, Cell and Environment* **24**, 755–768.
- Evans JR.** 2013. Improving photosynthesis. *Plant Physiology* **162**, 1780–1793.
- Fabre D, Yin X, Dingkuhn M, Clément-Vidal A, Roques S, Rouan L, Soutiras A, Delphine L.** 2019. Is triose phosphate utilization involved in the feedback inhibition of photosynthesis in rice under conditions of sink limitation? *Journal of Experimental Botany* **70**, 5773–5785.
- Farquhar GD, von Caemmerer S, Berry JA.** 1980. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ species. *Planta* **149**, 78–90.
- Farquhar GD, Sharkey TD.** 1982. Stomatal conductance and photosynthesis. *Annual Review of Plant Physiology* **33**, 317–345.
- Feldman AB, Leung H, Baraoidan M, Elmido-mabilangan A.** 2017. Increasing leaf vein density via mutagenesis in rice results in an enhanced rate of photosynthesis, smaller cell sizes and can reduce interveinal mesophyll cell number. *Frontiers in Plant Science* **8**, 1883.
- Flexas J, Ribas-Carbó M, Diaz-Espejo A, Galmés J, Medrano H.** 2008. Mesophyll conductance to CO₂: Current knowledge and future prospects. *Plant, Cell and Environment* **31**, 602–621.
- Fujita D, Trijatmiko KR, Tagle AG, Sapasap MV, Koide Y, Sasaki K, et al.** 2013. NAL1 allele from a rice landrace greatly increases yield in modern indica cultivars. *Proceedings of the National Academy of Sciences of the USA* **110**, 20431–20436.
- Gago J, Daloso DM, Figueroa CM, Flexas J, Fernie AR, Nikoloski Z.** 2016. Relationships of leaf net photosynthesis, stomatal conductance, and mesophyll conductance to primary metabolism: A multispecies meta-analysis approach. *Plant Physiology* **171**, 265–279.
- Giuliani R, Koteyeva N, Voznesenskaya E, Evans MA, Cousins AB, Edwards GE.** 2013. Coordination of leaf photosynthesis, transpiration, and structural traits in rice and wild relatives (Genus *Oryza*). *Plant Physiology* **162**, 1632–1651.
- Haritha G, Vishnukiran T, Yugandhar P, Sarla N, Subrahmanyam D.** 2017.

- Introgressions from *Oryza rufipogon* increase photosynthetic efficiency of KMR3 Rice Lines. *Rice Science* **24**, 85–96.
- Harley PC, Loreto F, Marco GDi, Sharkey TD.** 1992. Theoretical considerations when estimating the mesophyll conductance to CO₂ flux by analysis of the response of photosynthesis to CO₂. *Plant Physiology* **98**, 1429–1436.
- Horton P.** 2000. Prospects for crop improvement through the genetic manipulation of photosynthesis: morphological and biochemical aspects of light capture, *Journal of Experimental Botany* **51**, 475–485.
- Huang X, Kurata N, Wei X, Wang ZX, Wang A, Zhao Q, Zhao Y, Liu K, Lu H, Li W, et al.** 2012. A map of rice genome variation reveals the origin of cultivated rice. *Nature* **490**, 497–501.
- Ishiwata A, Ozawa M, Nagasaki H, Kato M, Noda Y, Yamaguchi T, Nosaka M, Shimizu-Sato S, Nagasaki A, Maekawa M, Hirano H-Y, Sato Y.** 2013. Two WUSCHEL-related homeobox genes, narrow leaf2 and narrow leaf3, control leaf width in rice. *Plant and Cell Physiology* **54**, 779–792.
- Ivanova LA, Yudina PK, Ronzhina DA, Ivanov LA, Hölzel N.** 2018. Quantitative mesophyll parameters rather than whole-leaf traits predict response of C₃ steppe plants to aridity. *New Phytologist* **217**, 558–570.
- Kiran TV, Rao YV, Subrahmanyam D, Rani NS, Bhadana VP, Rao PR.** 2013. Variation in leaf photosynthetic characteristics in wild rice species. *Photosynthetica* **51**, 350–358.
- Kondamudi R, Swamy KN, Rao YV, et al.** 2016. Gas exchange, carbon balance and stomatal traits in wild and cultivated rice (*Oryza sativa* L.) genotypes. *Acta Physiologiae Plantarum* **38**, 160.
- Kramer DM, Evans JR.** 2011. The Importance of energy balance in improving photosynthetic productivity. *Plant Physiology* **155**, 70–78.
- Kubien DS, Brown CM, Kane HJ.** 2011. Quantifying the amount and activity of rubisco in leaves. *Photosynthesis Research Protocol* **684**, 349–362.
- Kumarathunge DP, Medlyn BE, Drake JE, Rogers A, Tjoelker MG.** 2019. No evidence for triose phosphate limitation of light-saturated leaf photosynthesis under current atmospheric CO₂ concentration. *Plant, Cell and Environment* **42**, 3241–3252.
- Laisk AK.** 1977. Kinetics of photosynthesis and photorespiration in C₃-Plants. Nauka, Moscow
- Lawson T, Kramer DM, Raines CA.** 2012. Improving yield by exploiting mechanisms underlying natural variation of photosynthesis. *Current Opinion in Biotechnology* **23**, 215–220.
- Li Y, Gao Y, Xu X, Shen Q, Guo S.** 2009. Light-saturated photosynthetic rate in

- high-nitrogen rice (*Oryza sativa* L.) leaves is related to chloroplastic CO₂ concentration. *Journal of Experimental Botany* **60**, 2351–2360.
- Lin MT, Occhialini A, Andralojc PJ, Parry MAJ & Hanson MR.** 2014. A faster Rubisco with potential to increase photosynthesis in crops. *Nature* **513**, 547–550.
- Liu L, Sun G, Ren X, Li C, Sun D.** 2015. Identification of QTL underlying physiological and morphological traits of flag leaf in barley. *BMC Genetics* **16**, 29.
- Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCT} method. *Methods* **25**, 402–408.
- Long SP, Bernacchi CJ.** 2003. Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *Journal of Experimental Botany* **54**, 2393–2401.
- Long SP, Zhu XG, Naidu SL, Ort DR.** 2006. Can improvement in photosynthesis increase crop yields? *Plant, Cell and Environment* **29**, 315–330.
- Lux A, Morita S, Abe J, Ito K.** 2005. An improved method for clearing and staining free-hand sections and whole-mount samples. *Annals of Botany* **96**, 989–996.
- Mathan J, Bhattacharya J, Ranjan A.** 2016. Enhancing crop yield by optimizing plant developmental features. *Development* **143**, 3283–3294.
- Mathan J, Singh A, Ranjan A.** 2021. Sucrose transport and metabolism control carbon partitioning between stem and grain in rice. *Journal of Experimental Botany* <https://doi.org/10.1093/jxb/erab066>.
- Murchie EH, Lawson T.** 2013. Chlorophyll fluorescence analysis: A guide to good practice and understanding some new applications. *Journal of Experimental Botany* **64**, 3983–3998.
- Murchie EH, Niyogi KK.** 2011. Manipulation of photoprotection to improve plant photosynthesis. *Plant Physiology* **155**, 86–92.
- Ort DR, Merchant SS, Alric J, Barkan A, Blankenship RE, et al.** 2015. Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proceedings of the National Academy of Sciences of the USA* **112**, 8529–8536.
- Parry MAJ, Madgwick PJ, Carvahlo JFC, Andralojc PJ.** 2007. Prospects for increasing photosynthesis by overcoming the limitations of Rubisco. *Journal of Agricultural Science* **145**, 31–43.
- Parry MAJ, Andralojc PJ, Scales JC, Salvucci ME, Carmo-Silva AE, Alonso H, Whitney SM.** 2013. Rubisco activity and regulation as targets for crop improvement. *Journal of Experimental Botany* **64**, 717–730.
- Pathare VS, Koteyeva N, Cousins AB.** 2020. Increased adaxial stomatal density is associated with greater mesophyll surface area exposed to intercellular air spaces and

- mesophyll conductance in diverse C₄ grasses. *New Phytologist* **225**,169-182.
- Perdomo JA, Capó-Bauçà S, Carmo-Silva E and Galmés J.** 2017. Rubisco and rubisco activase play an important role in the biochemical limitations of photosynthesis in rice, wheat, and maize under high temperature and water deficit. *Frontiers in plant science* **8**, 490.
- Pinto H, Powell JR, Sharwood RE, Tissue DT, Ghannoum O.** 2016. Variations in nitrogen use efficiency reflect the biochemical subtype while variations in water use efficiency reflect the evolutionary lineage of C₄ grasses at inter-glacial CO₂. *Plant, Cell and Environment* **39**, 514-26.
- Porra RJ, Thompson WA, Kriedemann PE.** 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: Verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **975**, 384–394.
- Reynolds ES.** 1963. The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *The Journal of Cell Biology* **17**, 208–212.
- Sack L, Scoffoni C.** 2013. Leaf venation: structure, function, development, evolution, ecology and applications in the past, present and future. *New Phytologist* **198**, 983-1000.
- Sage RF.** 2002. Variation in the kcat of Rubisco in C₃ and C₄ plants and some implications for photosynthetic performance at high and low temperature. *Journal of Experimental Botany* **53**, 609–620.
- Sage TL, Sage RF.** 2009 The functional anatomy of rice leaves: Implications for refixation of photorespiratory CO₂ and efforts to engineer C₄ photosynthesis into rice. *Plant and Cell Physiology* **50**, 756–772.
- Sanchez PL, Wing RA, Brar DS.** 2013. The wild relative of Rice: Genomes and Genomics. *Genetics and Genomics of Rice* **5**, 9-25.
- Scafaro AP, von Caemmerer S, Evans JR, Atwell BJ.** 2011. Temperature response of mesophyll conductance in cultivated and wild *Oryza* species with contrasting mesophyll cell wall thickness. *Plant, Cell and Environment* **34**, 1999-2008.
- Sharkey TD.** 1985. Photosynthesis in intact leaves of C₃ plants: Physics, physiology and rate limitations. *Botanical Review* **51**, 53-105.
- Sharkey TD, Bernacchi CJ, Farquhar GD, Singsaas EL.** 2007. Fitting photosynthetic carbon dioxide response curves for C₃ leaves. *Plant, Cell and Environment* **30**, 1035–1040.
- Sharkey TD.** 2016. What gas exchange data can tell us about photosynthesis. *Plant, Cell and Environment* **39**, 1161–1163.

- Sharwood RE, Sonawane BV, Ghannoum O, Whitney SP.** 2016. Improved analysis of C₄ and C₃ photosynthesis via refined *in vitro* assays of their carbon fixation biochemistry. *Journal of Experimental Botany*, **67**, 3137–3148.
- Sharwood RE.** 2017. Engineering chloroplasts to improve Rubisco catalysis: prospects for translating improvements into food and fiber crops. *New Phytologist* **213**, 494–510.
- Simkin AJ, López-calcano PE, Raines CA.** 2019. Feeding the world□: improving photosynthetic efficiency for sustainable crop production. *Journal of Experimental Botany* **70**, 1119–1140.
- Takai T, Adachi S, Taguchi-Shiobara F, Sanoh-Arai Y, Iwasawa N, Yoshinaga S, Hirose S, Taniguchi Y, Yamanouchi U, Wu J, Matsumoto T, Sugimoto K, Kondo K, Ikka T, Ando T, Kono I, Ito S, Shomura A, Ookawa T, Hirasawa T, Yano M, Kondo M, Yamamoto T.** 2013. A natural variant of NAL1, selected in high-yield rice breeding programs, pleiotropically increases photosynthesis rate. *Scientific Reports* **3**, 2149.
- Terashima I, Hanba YT, Tholen D, Niinemets Ü.** 2011. Leaf functional anatomy in relation to photosynthesis. *Plant Physiology* **155**, 108–16.
- Tholen D, Boom C, Zhu XG.** 2012. Opinion: prospects for improving photosynthesis by altering leaf anatomy. *Plant Science* **197**, 92–101.
- Traw MB, Ackerly DD.** 1995. Leaf Position, Light Levels, and Nitrogen Allocation in Five Species of Rain Forest Pioneer Trees. *American Journal of Botany* **82**, 1137– 1143.
- Wing RA, Purugganan MD, Zhang Q.** 2018. The rice genome revolution: from an ancient grain to Green Super Rice. *Nature Reviews Genetics* **19**, 505–517.
- Wu A, Hammer GL, Doherty A, von Caemmerer S, Farquhar GD.** 2019. Quantifying impacts of enhancing photosynthesis on crop yield. *Nature plants* **5**, 380–388.
- Yeo ME, Yeo AR, Flowers TJ.** 1994. Photosynthesis and photorespiration in the genus *Oryza*. *Journal of Experimental Botany* **45**, 553–560.
- Zhao M, Lafitte HR, Sacks E, Dimayuga G, Acun TLB.** 2008. Perennial *O. sativa* × *O. rufipogon* interspecific hybrids, I. Photosynthetic characteristics and their inheritance. *Field Crops Research* **106**, 203–213.
- Zhao M, Ding Z, Lafitte R, Sacks E, Dimayuga G, Holt D.** 2010. Photosynthetic characteristics in *Oryza* species. *Photosynthetica* **48**, 234–240.
- Zhu XG, Long SP, Ort DR.** 2010. Improving photosynthetic efficiency for greater yield. *Annual Review of Plant Biology* **61**, 235–61.

Table 1. Leaf photosynthesis rate and related physiological traits of the selected cultivated and wild rice accessions under controlled growth conditions. Each value represents mean $\pm$ SE, where n = five data points from different plants. Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Genotypes	Cultivated/ Wild	A ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)	C_i ($\mu\text{mol CO}_2 \text{ mol}^{-1}$)	g_s ($\text{mol m}^{-2} \text{ s}^{-1}$)	CE ($\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)
<i>O. sativa</i> cv. Nipponbare	Cultivated	11.15 $\pm$ 1.2a	309.0 $\pm$ 40.5a	0.22 $\pm$ 0.05a	0.03 $\pm$ 0.005a
<i>O. sativa</i> cv. IR 64	Cultivated	11.22 $\pm$ 1.1a	300.9 $\pm$ 14.5a	0.21 $\pm$ 0.01a	0.03 $\pm$ 0.008a
<i>O. glaberrima</i> (IR 102925)	Cultivated	13.79 $\pm$ 0.4b	282.6 $\pm$ 12.9a	0.28 $\pm$ 0.02b	0.04 $\pm$ 0.003a
<i>O. latifolia</i> (IRGC 99596)	Wild	18.38 $\pm$ 0.1c	306.2 $\pm$ 13.9a	0.40 $\pm$ 0.03c	0.06 $\pm$ 0.001b
<i>O. australiensis</i> (IRGC 105272)	Wild	19.18 $\pm$ 0.1c	279.8 $\pm$ 21.2a	0.35 $\pm$ 0.02c	0.06 $\pm$ 0.001b

Table 2. Mitochondrial respiration rate in the light (R_d), CO_2 compensation point related to C_i (Γ^*), and mesophyll conductance (g_m) for the selected cultivated and wild rice genotypes. Each value represents mean $\pm$ SE, where n = three data points from different plants. Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Genotypes	Cultivated/ Wild	R_d ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)	Γ^* ($\mu\text{mol CO}_2 \text{ mol}^{-1}$)	g_m ($\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$)
<i>O. sativa</i> cv. Nipponbare	Cultivated	$1.05 \pm 0.15\text{a}$	$43.8 \pm 1.7\text{a}$	$0.12 \pm 0.01\text{a}$
<i>O. sativa</i> cv. IR 64	Cultivated	$1.03 \pm 0.10\text{a}$	$42.3 \pm 3.0\text{a}$	$0.11 \pm 0.01\text{a}$
<i>O. glaberrima</i> (IR 102925)	Cultivated	$1.15 \pm 0.16\text{a}$	$44.5 \pm 2.0\text{a}$	$0.26 \pm 0.03\text{b}$
<i>O. latifolia</i> (IRGC 99596)	Wild	$1.37 \pm 0.19\text{b}$	$45.4 \pm 0.7\text{a}$	$0.33 \pm 0.02\text{c}$
<i>O. australiensis</i> (IRGC 105272)	Wild	$1.39 \pm 0.18\text{b}$	$45.5 \pm 0.8\text{a}$	$0.30 \pm 0.05\text{c}$

Table 3. CO₂-saturated maximum photosynthesis (A_{max}), maximum *in vivo* Rubisco activity (V_{cmax}), maximum electron transport rate (J_{max}), and maximum rate of triose phosphate utilization (TPU) through A/C_i curve analysis. Each value represents mean $\pm$ SE, where n = five data points from different plants. Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Genotypes	Cultivated/ Wild	A_{max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	V_{cmax} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	J_{max} ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	TPU ($\mu\text{mol m}^{-2} \text{s}^{-1}$)
<i>O. sativa</i> cv. Nipponbare	Cultivated	23.99 $\pm$ 0.40b	62.75 $\pm$ 0.92b	124.86 $\pm$ 1.37b	9.10 $\pm$ 0.06b
<i>O. sativa</i> cv. IR 64	Cultivated	21.37 $\pm$ 0.88a	51.48 $\pm$ 2.69a	103.06 $\pm$ 3.19a	8.39 $\pm$ 0.19a
<i>O. glaberrima</i> (IR 102925)	Cultivated	30.41 $\pm$ 0.14c	70.04 $\pm$ 0.79cd	149.21 $\pm$ 1.80c	10.28 $\pm$ 0.10c
<i>O. latifolia</i> (IRGC 99596)	Wild	34.32 $\pm$ 0.39d	76.75 $\pm$ 1.24de	166.17 $\pm$ 2.86d	11.72 $\pm$ 0.27d
<i>O. australiensis</i> (IRGC 105272)	Wild	36.81 $\pm$ 0.37e	81.25 $\pm$ 1.08e	181.93 $\pm$ 0.77e	12.27 $\pm$ 0.12d

Table 4. *in vitro* RuBP-use and Rubisco carboxylation rate (V_{cmax}) for selected rice genotypes. Each value represents mean $\pm$ SE, where n = three data points from different plants. The significant difference among the accessions is indicated by different letters (one-way ANOVA followed by Tukey's HSD, $P < 0.05$).

Genotypes	Cultivated/ Wild	<i>RuBP-use</i> (nmol s ⁻¹)			V_{cmax} (μmol m ⁻² s ⁻¹)		
		120 s	180 s	240 s	120 s	180 s	240 s
<i>O. sativa</i> cv. Nipponbare	Cultivated	0.198 $\pm$ 0.01b	0.200 $\pm$ 0.01b	0.195 $\pm$ 0.01b	43.52 $\pm$ 1.50b	43.98 $\pm$ 1.43b	42.71 $\pm$ 1.40b
<i>O. sativa</i> cv. IR 64	Cultivated	0.166 $\pm$ 0.01a	0.168 $\pm$ 0.01a	0.165 $\pm$ 0.01a	36.37 $\pm$ 3.00a	36.91 $\pm$ 1.00a	36.22 $\pm$ 1.22a
<i>O. glaberrima</i> (IR 102925)	Cultivated	0.231 $\pm$ 0.01cd	0.236 $\pm$ 0.01c	0.223 $\pm$ 0.01c	50.67 $\pm$ 1.71c	51.87 $\pm$ 0.87c	49.01 $\pm$ 0.85c
<i>O. latifolia</i> (IRGC 99596)	Wild	0.256 $\pm$ 0.01d	0.261 $\pm$ 0.01d	0.245 $\pm$ 0.01cd	56.10 $\pm$ 1.30d	57.29 $\pm$ 1.35d	53.69 $\pm$ 1.39d
<i>O. australiensis</i> (IRGC 105272)	Wild	0.282 $\pm$ 0.01d	0.284 $\pm$ 0.01d	0.260 $\pm$ 0.01d	61.99 $\pm$ 0.48d	62.22 $\pm$ 1.35d	57.02 $\pm$ 0.65d

Figure legends

Fig. 1. Variations in leaf photosynthesis and related physiological features, as well as morphological differences across the selected cultivated and wild rice accessions.

(A-E) Quantification of net photosynthesis per unit area of flag leaf, A (A), stomatal conductance to water, g_s (B), intercellular CO_2 concentration, C_i (C), carboxylation efficiency, CE (D), and Total chlorophyll content, TCC (E) for six cultivated rice and three wild rice species. Cultivated rice varieties and wild species are represented with yellow- and green-shaded boxes, respectively. Each box plot shows the median and interquartile range with minimum and maximum values of fifteen data points from different plants. Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

(F) Representative images of the selected cultivated and wild rice accessions that captured the entire variation range for the leaf photosynthesis per unit area in the experiment. Shown are the field-grown plants of the selected genotypes. Scale bars represent 30 cm.

Fig. 2. Photochemical features of the selected cultivated and wild rice accessions.

Quantification of electron transport rate, J_{PSII} (A), the efficiency of PSII, Φ_{PSII} (B), and photochemical quenching, qP (C) of the selected accessions at varying CO_2 concentrations. Each value represents mean $\pm$ SE, where $n =$ five data points from different plants. Different letters at each CO_2 concentration indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Fig. 3. RuBP-use, Rubisco activity, and leaf nitrogen content across the selected accessions.

(A) NADH-linked spectrophotometric measurement of RuBP-use (nmol) at different time points from 10 to 1800 sec.

(B) Quantification of V_{max} for the selected accessions at different time points up to saturation.

(C) Leaf nitrogen content/unit leaf area (N_{area}) in the flag leaves of the selected accessions. Values represent mean $\pm$ SE (three data points from different plants). Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Fig. 4. Expression pattern of photosynthetic and chlorophyll biosynthetic genes across the selected accessions.

(A) Expression levels of genes encoding a key component of the light-harvesting complex, Rubisco, and Triose phosphate/phosphate translocator.

(B) Expression levels of key chlorophyll biosynthetic genes.

Relative expressions of the target genes are normalized using rice actin (LOC_Os03g50885.1) as the internal control. Values represent mean $\pm$ SE (three data points from different plants). Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Gene abbreviations: LHCI, Light-harvesting chlorophyll protein complex I; Rubisco SS, Ribulose biphosphate carboxylase small subunit; Rubisco LS, Ribulose biphosphate carboxylase large subunit; TPT, Triose phosphate/phosphate translocator; GluTR, Glutamyl-tRNA reductase; PPOX, Protoporphyrinogen oxidase; Mg-CHLD subunit ChlD, Magnesium chelatase; CHLSYN, Chlorophyll synthase; CAO, Chlorophyllide a oxygenase.

Fig. 5. Leaf developmental traits affecting photosynthesis rate and related physiological traits of rice.

(A) Correlation of leaf photosynthesis rate and related physiological traits with leaf developmental features. The red shade represents a significant positive correlation at $P \leq 0.05$.

(B) Bright-field images of a cross-section of a flag leaf of the selected genotypes showing variations in mesophyll cells and vasculature. Arrows indicate bulliform cells (bc), bundle sheath (bs), epidermis (ep), mesophyll cells (mc), and minor veins (miv). Scale bars represent 100 μ m.

(C) Bright-field microscopy images showing variations in mesophyll cell features, including lobbing. Scale bar represents 10 μ m.

(D-F) Quantification of mesophyll cell number per unit (cm^2) leaf area (D), individual mesophyll cell area (E), and individual mesophyll cell surface area exposed to intercellular space (S_{mes} , F) of the selected cultivated and wild rice accessions.

(G-I) Quantification of chloroplast number per unit (cm^2) mesophyll cell area (G), individual chloroplast area (H), and individual chloroplast area exposed to intercellular space (S_c , I) of the selected cultivated and wild rice accessions.

Cultivated rice accessions and wild species are represented with yellow- and green-shaded boxes, respectively. Each box plot shows the median and interquartile range with minimum

and maximum values. Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Fig. 6. Leaf developmental, photochemical, and biochemical traits contributing towards photosynthetic differences between two extreme genotypes: *Oryza sativa* ssp. *indica* cv. IR 64 and *Oryza australiensis*. Wider leaf, larger mesophyll cells, larger veins, fewer number of mesophyll cells between two consecutive veins, more chloroplasts, efficient Electron Transport Rate along with higher leaf nitrogen content and Rubisco activity facilitate higher leaf photosynthesis per unit area in the wild rice species *Oryza australiensis* compared to the cultivated variety *Oryza sativa* ssp. *indica* cv. IR 64.

F

Fig. 1. Variations in leaf photosynthesis and related physiological features, as well as morphological differences across the selected cultivated and wild rice accessions.

(A-E) Quantification of net photosynthesis per unit area of flag leaf, A (A), stomatal conductance to water, g_s (B), intercellular CO_2 concentration, C_i (C), carboxylation efficiency, CE (D), and Total chlorophyll content, TCC (E) for six cultivated rice and three wild rice species. Cultivated rice varieties and wild species are represented with yellow- and green-shaded boxes, respectively. Each box plot shows the median and interquartile range with minimum and maximum values of fifteen data points from different plants. Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

(F) Representative images of the selected cultivated and wild rice accessions that captured the entire variation range for the leaf photosynthesis per unit area in the experiment. Shown are the field-grown plants of the selected genotypes. Scale bars represent 30 cm.

Fig. 2. Photochemical features of the selected cultivated and wild rice accessions.

Quantification of electron transport rate, J_{PSII} (A), the efficiency of PSII, ϕ_{PSII} (B), and photochemical quenching, qP (C) of the selected accessions at varying CO₂ concentrations. Each value represents mean $\pm$ SE, where n = five data points from different plants. Different letters at each CO₂ concentration indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Fig. 3. RuBP-use, Rubisco activity, and leaf nitrogen content across the selected accessions.

(A) NADH-linked spectrophotometric measurement of RuBP-use (nmol) at different time points from 10 to 1800 sec.

(B) Quantification of V_{max} for the selected accessions at different time points up to saturation.

(C) Leaf nitrogen content/unit leaf area (N_{area}) in the flag leaves of the selected accessions.

Values represent mean $\pm$ SE (three data points from different plants). Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Fig. 4. Expression pattern of photosynthetic and chlorophyll biosynthetic genes across the selected accessions.

(A) Expression levels of genes encoding a key component of the light-harvesting complex, Rubisco, and Triose phosphate/phosphate translocator.

(B) Expression levels of key chlorophyll biosynthetic genes.

Relative expressions of the target genes are normalized using rice actin (LOC_Os03g50885.1) as the internal control. Values represent mean $\pm$ SE (three data points from different plants). Different letters indicate statistical significance according to one-way ANOVA followed by post-hoc Tukey HSD calculation at $P \leq 0.05$.

Gene abbreviations: LHCI, Light-harvesting chlorophyll protein complex I; Rubisco SS, Ribulose biphosphate carboxylase small subunit; Rubisco LS, Ribulose biphosphate carboxylase large subunit; TPT, Triose phosphate/phosphate translocator; GluTR, Glutamyl-tRNA reductase; PPOX, Protoporphyrinogen oxidase; Mg-CHLD subunit ChlD, Magnesium chelatase; CHLSYN, Chlorophyll synthase; CAO, Chlorophyllide a oxygenase.

Cultivated rice
Oryza sativa ssp. *indica* cv. IR 64

Less Biomass
Low Photosynthesis Rate

Wild rice
Oryza australiensis

More Biomass
High Photosynthesis Rate

Smaller mesophyll cells
Fewer chloroplasts

Larger mesophyll cells
More chloroplasts

More mesophyll
cells between
consecutive veins

Fewer mesophyll
cells between
consecutive veins

Light

Leaf Nitrogen content, Rubisco activity, and
RuBP-regeneration/Electron Transport Rate –
limited photosynthesis

Light

Higher Leaf Nitrogen content, Rubisco
activity, and Efficient Photosystem

Fig.6. Leaf developmental, photochemical, and biochemical traits contributing towards photosynthetic differences between two extreme genotypes: *Oryza sativa* ssp. *indica* cv. IR 64 and *Oryza australiensis*. Wider leaf, larger mesophyll cells, larger veins, fewer number of mesophyll cells between two consecutive veins, more chloroplasts, efficient Electron Transport Rate along with higher leaf nitrogen content and Rubisco activity facilitate higher leaf photosynthesis per unit area in the wild rice species *Oryza australiensis* compared to the cultivated variety *Oryza sativa* ssp. *indica* cv. IR 64.