Comparative whole-genome approach to identify bacterial traits for microbial interactions

Luca Zoccarato^{*a}, Daniel Sher^{*b}, Takeshi Miki^c, Daniel Segre^{d,e}, Hans-Peter Grossart^{*a,f,g}

(a) Department Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), 16775 Stechlin, Germany

(b) Department of Marine Biology, Leon H. Charney School of Marine Sciences, University of Haifa, 3498838 Haifa, Israel

(c) Department of Environmental Solution Technology, Ryukoku University, 612-8577 Kyoto, Japan

(d) Departments of Biology, Biomedical Engineering, Physics, Boston University, 02215 Boston, MA

(e) Bioinformatics Program & Biological Design Center, Boston University, 02215 Boston, MA

(f) Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), 14195 Berlin, Germany

(g) Institute of Biochemistry and Biology, Potsdam University, 14476 Potsdam, Germany

Corresponding authors(*)

Luca Zoccarato, zoccarato@igb-berlin.de; Daniel Sher, dsher@univ.haifa.ac.il; Hans-Peter Grossart, hgrossart@igb-berlin.de

Abstract

Microbial interactions shape the structure and function of microbial communities with profound consequences for biogeochemical cycles and ecosystem health. Yet, most interaction mechanisms are studied only in model systems and their prevalence is unknown. To systematically explore the functional and interaction potential of sequenced marine bacteria, we developed a new trait-based approach, and applied it to 473 complete genomes (248 genera), representing a significant fraction of marine microbial communities. We identified genome functional clusters (GFCs) which condense bacterial diversity into groups with potentially common ecology and life history. Most

GFCs revealed unique combinations of interaction traits, some widely distributed (e.g., antimicrobial activity), others less common (e.g., production of siderophores and phytohormones, biosynthesis and export of specific B vitamins). Specific GFCs, comprising Alpha- and Gammaproteobacteria, are predicted to preferentially interact synergistically and/or antagonistically with bacteria and phytoplankton. Moreover, linked trait clusters (LTCs) identify traits that may have evolved together and point to specific modes of interactions. Our approach translates multidimensional genomic information into an atlas of marine bacteria and their ecosystem functions, relevant for understanding the fundamental rules that govern community assembly and dynamics.

Keywords

marine bacteria; functional clusters; genetic traits; interaction mechanisms, B vitamins; siderophores; phytohormones; motility and chemotaxis; secretion systems; antimicrobial compounds

Introduction

Interactions among aquatic microorganisms such as symbiosis, parasitism, predation and competition, greatly shape the composition and activity of microbial communities¹⁻³. In particular, interactions between heterotrophic bacteria and primary producers (phytoplankton) influence the growth of both organisms^{4,5} with consequences for the ecosystem functioning and the biogeochemical cycles^{6,7}. For instance, heterotrophic bacteria consume up to 50% of the organic matter released by phytoplankton, significantly affecting the dynamics of the huge pool of dissolved organic carbon in the oceans⁸. Thus, if and how a bacterium can interact with other microorganisms may have important consequences for the biological carbon pump in the current and future oceans^{9,10}.

54 Studies using specific model organisms in binary co-cultures have started to elucidate mechanisms underlying
 55 ing marine microbial interactions (mostly between bacteria and phytoplankton) ^{1,2,5}. Although these results
 56 do not reflect the complexity of natural environments and the potential for higher-order effects ¹¹, they allow
 57 low to identify the chemical signals and resulting changes in gene expression and physiology that underlie
 58 these interactions. For example, bacteria associated with phytoplankton (e.g. within the phycosphere, ^{5,12})
 59 gain access to labile organic carbon released by the primary producers, e.g. amino acids and small sulfur-
 60 containing compounds ^{13–18}. In return, phytoplankton benefit from an increased accessibility to nutrients via
 61 bacteria-mediated processes, e.g. nitrogen and phosphorus remineralization ¹⁹, vitamin supply ^{14,20} and iron
 62 scavenging via formation of siderophores ^{21,22}. In addition to such metabolic interactions, direct signalling
 63 may also occur between bacteria and phytoplankton, with heterotrophic bacteria directly controlling the
 64 phytoplankton cell cycle through phytohormones ^{13,23} or harming it via toxins ^{18,24}. Through such specific in-
 65 fochemical-mediated interactions, bacteria may also directly affect the release rate of organic carbon from
 66 phytoplankton, as well as rates of mortality and aggregation ^{18,23,25}.

67 While much is known about microbial interactions involving model organisms such as specific strains of *Ros-*
 68 *eobacter* ^{13,18–20,24}, *Alteromonas* ^{26–28}, Cyanobacteria ^{19,29} or *Vibrio* ^{30,31}, relatively little is known regarding how
 69 widely distributed such interactions are across microbial diversity. The few experimental studies that measure
 70 microbial interactions across diversity (e.g. ^{32–34}) are usually limited in their phylogenetic scope and are
 71 performed under conditions different from natural marine environments. However, the knowledge obtained
 72 from model organisms on the molecular mechanisms underlying microbial interactions and the increasing availability
 73 of high-quality genomes present an opportunity to map known interaction mechanisms to a large set of bacterial
 74 species from various taxa. Here, we re-analyse 421 previously published genomes of diverse marine bacteria
 75 that represent a significant fraction of marine microbial communities (213 genera), providing an “atlas” of
 76 their functional metabolic capacity. The atlas includes also 52 bacteria isolated from extreme marine habitats,
 77 humans and plant roots which serve as functional out-groups and represent well known symbiotic bacteria.
 78 Several previous studies have aimed to characterize and cluster genomes based on their predicted functional
 79 similarity, defined usually using individual genes (e.g. ^{35–38}) or coarse

functional categories (e.g. COGs,^{37,39,40})(Supplementary Table 1, Supplementary text). We chose to take a trait-based approach rather than a gene-based one, which is an intermediate level of resolution between individual genes and coarse functions. Trait-based approaches offer a new perspective to investigating microbial diversity with a more mechanistic understanding⁴¹ but have been used only in few specific cases to highlight putative bacterial interactions (e.g.,⁴²). We focused on the following traits: 1) the metabolic capacity to synthesize or degrade specific biomolecules and to regulate such processes (KEGG modules); 2) specific functions related to microbial interactions, such as motility, chemotaxis and the ability to produce molecules such as siderophores, toxins and antibiotics. This allows to classify genomes into coherent functional units, some of which recapitulate known bacterial groups with well-defined ecological roles, while others refer to potential new groups. Furthermore, genetic traits can be grouped into linked trait clusters, representing functions that likely evolved together and may be functionally connected (i.e. participating in the same process). Our approach maps the mechanisms of microbial interactions elucidated in model organisms across bacterial diversity, suggests specific groups of bacteria likely to interact in a similar manner, and identifies trait combinations that likely mediate these interactions.

Results and Discussion

Genome functional clusters (GFCs) group genomes with similar ecology

To obtain an overview on the functional capabilities of marine bacteria, we re-annotated a set of 473 high-quality genomes and analysed them using a trait-based workflow, which focuses on the detection of complete genetic traits rather than on the presence of individual genes (Supplementary Fig. 1 and Supplementary text). Genetic traits were represented by metabolic KEGG modules, secondary metabolite pathways, transporters, phytohormones and siderophores production, as well as the degradation of specific sulfur metabolites. Among the identified genetic traits, those known to mediate cell-cell interactions in bacterial model systems (see^{4,5,43}) were flagged as “interaction traits” (e.g. production of certain vitamins, vibrioferrin or a specific secretion systems; Supplementary Table 2).

Based on the occurrence patterns of all traits (Fig. 1), we could cluster the genomes into 47 genome functional clusters (GFCs; Supplementary Table 3). In each GFC, genomes encode similar genetic traits, and thus GFCs are expected to be coherent in terms of functional and metabolic capacity, including the ways that the grouped bacteria respond to abiotic cues and interact with other microbes. Previous genome comparison approaches have identified genome clusters that match to ecologically relevant groups (e.g. ecotypes, as defined for *Bacillus pumilus*³⁶ and *Prochlorococcus*²⁹) or lifestyles (e.g. oligotrophic and copiotrophic species³⁹). Similarly, in our analysis we found GFCs that represent group of organisms with a defined ecology and life history, such as the Pelagibacterales group (GFC 2), different ecotypes of Cyanobacteria (GFCs 15 and 36), or *Vibrio* groups, characterized by different host-specificity and pathogenicity (GFCs 25 and 47)(see Supplementary text). Specific GFCs were also identified for each of three groups of Gammaproteobacteria (*Alteromonas*, *Marinobacter* and *Pseudoalteromonas*) which are typically considered as “copiotrophs”, often associated with organic particles or phytoplankton^{44–47}. A detailed analysis of the traits found in each of the respective GFCs (Supplementary Fig. 2) suggested that *Pseudoalteromonas* and *Alteromonas* bore more genetic traits involved in the resistance against antimicrobial compounds, as well as regulation for osmotic and redox stresses. They also had similar vitamin B1 and siderophore transporters, which are different from those encoded by *Marinobacter*. *Marinobacter* possessed several more transporters for phosphonate and amino acids, as well as specific regulatory systems for adhesion (e.g. alginate and type 4 fimbriae production) and chemotaxis. These patterns advocated that there might be coherent physiological and/or ecological differences between these three groups. Overall, our GFC framework recapitulates previous knowledge on bacterial groups with defined ecology and life history, and provides a way to delineate and characterize potentially new ecological groups.

The correlation between functional and taxonomic classifications still represents an open question and, in marine environments, recent studies have provided both supporting^{35,48} and disproving argumentations about the strength of such correlation^{37,49}. Therefore, we sought to understand the extent to which our retrieved GFCs overlapped with the genomes’ taxonomy. In our analysis, we defined a GFC as taxonomically coherent (i.e. monophyletic) when all grouped genomes belonged to the same taxon and all genomes of

that taxon were grouped in that GFC (see Supplementary Fig. 3 and Supplementary text for more information). Monophyletic GFCs imply that the taxonomic affiliation of these bacteria can predict the traits encoded in their genomes. According to this metric, 23 out of the 47 GFCs were taxonomically coherent, most of them at the genus level including all Firmicutes and half of the Alpha- and Gammaproteobacteria GFCs. Of the remaining 24 GFCs, 5 were polyphyletic (i.e. include genomes of different phyla) and 19 were paraphyletic. Paraphyletic GFCs contained genomes of multiple taxa (differing at genus, family or even order rank) whereby these taxa were also partitioned among multiple GFCs. These GFC comprised genomes of Cyanobacteria, Bacteroidota and the remaining half of the Alpha- and Gammaproteobacteria.

Overall, half of the detected GFCs were monophyletic and support the existence of a strong correlation between taxonomy and functionality in marine bacteria, while the remaining paraphyletic GFCs highlight that, in some cases, the taxonomic partitioning (based on the Genome Database Taxonomy⁵⁰) do not completely reflect the functional differentiation. At the whole-community level, it has been shown that taxonomically distinct communities exhibit similar functional profiles, which led to the suggestion that some bacterial clades have similar genetic capacity, and can replace each other while maintaining unchanged the community functioning^{51,52}. The polyphyletic GFCs may group such taxonomically different but functionally similar organisms, and this is supported by examples in GFCs 33 and 41 (grouping thermo- and halotolerant bacteria), or GFC 17 (grouping sulfur-oxidizing and facultative anaerobe bacteria). One of the main biological processes that mediates such functional “homogenisation” is horizontal gene transfer^{53,54} and, indeed, most of the paraphyletic GFCs grouped with each other known copiotrophs or particle/host associated bacteria (e.g. Rhodobacteraceae in GFCs 9, 30 and 40, Vibrionaceae in GFCs 25 and 47, Alteromonadaceae in GFCs 21 and 24, Oleiphilaceae in GFC 35, or Halomonadaceae in GFC 43;^{13,18,45,55,56}), whose lifestyles favour events of genetic exchange (e.g.⁵⁷). Conversely, another study suggested that this perceived similarity in community function reflects only known metabolic pathways, and it is therefore possible that adding to our analysis also unknown genes might separate these GFCs into monophyletic ones⁵⁸. Although hypothetical genes would be of no use (not informative) in a trait-based approach, we hypothesize that they could com-

156 plement the horizontal gene transfer hypothesis in explaining the blurred taxonomic profiles of the para-
 157 phyletic GFCs.
 158

Fig. 1: **Atlas of Marine Microbial Functional Traits** showing presence/absence of genetic traits across all analysed genomes. Columns represent genomes grouped into genome functional clusters (GFCs) as shown by the horizontal grey bar. The horizontal colour bar represents the taxonomic affiliation of genomes (mainly phyla, with the exception of Proteobacteria that are represented at the class level). The number next to each taxon in the legend represents the percent from the total genomes analysed. Rows represent the genetic traits grouped into linked trait clusters (LTCs) as shown by the vertical grey bar. LTCs discussed

in the text are labelled with the related number alongside the grey bar. Left-side row annotations show the frequency of each genetic trait across all tested genomes and the annotation tool (colour coded bar) with which they have been identified. Both dendrograms are computed using the *aggExCluster* function (r package *apcluster*) that generates hierarchal clustering from an affinity propagation result. An interactive version of this figure is available as Supplementary file 1.

159

160 **GFCs are ecologically-relevant entities in natural communities**

161 To what extent do the GFCs represent the diversity of natural communities, or display dynamics expected
162 from ecological units in the oceans? To test this, we mapped the 16S rDNA reads from a natural coastal
163 community that was sampled at high temporal frequency ⁵⁹ to the GFCs (Supplementary text and Supple-
164 mentary Fig 4). Firstly, a significant fraction of the natural community was represented with high fidelity in
165 the GFCs (mean 22.9% of the 16S rDNA reads, range 12.7-44.3%). Thus, despite the inherent bias derived
166 from using only high-quality, closed genomes available (mainly) from cultured bacteria (legend of Figure 1),
167 the GFCs represented a significant fraction of bacterial diversity. Similar results were obtained with an open-
168 ocean community from the Eastern Mediterranean sampled each season for 3 years ⁶⁰ (mean 13.9% of the
169 16S rDNA reads, range 0.5-60.0%), where the GFCs represented a significantly higher fraction of the micro-
170 bial community on particles >11 µm compared to free-living bacteria (5-0.22 µm; Supplementary Fig. 5).
171 Secondly, using a temporal deconvolution analysis of the coastal site ⁵⁹ we found that individual 16S phylo-
172 types belonging to the same GFC displayed significantly more synchronous temporal trends (p-value <
173 0.001) than phylotypes belonging to different GFCs (Supplementary text and Supplementary Fig. 6). Assum-
174 ing that similar temporal trends suggest similar ecological niches, these results suggest that (at least some
175 of) the GFCs represent bona-fide ecological units.

176

177 **Specific GFCs are enriched in “interaction traits”**

178 We next focused on selected traits potentially involved in microbial interactions – vitamin exchange,
 179 siderophore and phytohormone production and antibiosis – asking whether we could observe patterns in
 180 their distribution across the genome dataset. As shown in Figure 2, these traits are not equally distributed
 181 among the GFCs – rather, some GFCs were significantly enriched in “interaction traits” (Supplementary Fig.
 182 7a-b). While the number of interaction traits depended to some extent on genome size, as expected from
 183 previous studies ^{61,62}, Gammaproteobacteria and several Alphaproteobacteria encoded more interaction
 184 traits than expected by their genome size, while Bacteroidota encoded fewer (Supplementary Fig. 7c-d).
 185 Overall, GFCs grouping genomes of typical host- or particle-associated bacteria, such as of Alpha- and
 186 Gammaproteobacteria ⁵⁶, are predicted to bear almost the full combination of these traits to sense (chemo-
 187 taxis, quorum sensing), reach (motility and adhesion) and fight (production/resistance towards antimicro-
 188 bial compounds, secretion systems) for a targeted hotspot. Contrary, some ubiquitous copiotrophs (e.g.
 189 Bacteroidota) ⁵⁶ and known free-living taxa (e.g. pico-Cyanobacteria and Pelagibacterales) ^{29,61} possess only a
 190 scarce and scattered equipment of such traits. Below, we describe in more detail some of the main observa-
 191 tions on the distribution of interaction traits across diversity.

192

Fig. 2: Distribution across genome functional clusters (GFCs) of interaction traits known to mediate cell-cell interactions in bacterial model systems. Each slice shows the interaction traits present in a GFC and, as dendrogram, the functional similarity of genetic traits between the grouped genomes (hierarchical clustering of r-correlation matrix with complete agglomeration algorithm). The dark bars show the number of interaction traits annotated in each genome. Genomes belonging to model bacteria used to discover some of these traits are highlighted in blue if the interaction was positive (e.g. enhancing phytoplankton growth), in red if it was negative (e.g. kill the host) or in grey if the interaction shifted from positive to negative.

193

194 Many bacteria need “to shop” for their vitamins

195 Vitamins B₁, B₇ and B₁₂ are essential cofactors for microbes. Some microorganisms (including abundant
196 phytoplankton) are auxotrophic for these vitamins and need to obtain them from co-occurring bacteria^{63,64}.

197 Vitamins are found at low concentrations in aquatic ecosystems^{65,66} and their supply can limit biogeochem-
 198 ical cycles, e.g. through limiting primary productivity in the Southern Ocean⁶⁷. Less than half of all the gen-
 199 omes in our dataset were predicted to produce all three vitamins (~39%, including all pico-Cyanobacteria,
 200 Actinobacteriota and many Gammaproteobacteria; Fig. 3a). Of the rest, ~29% synthesized at least two B vit-
 201 amins (e.g. some Alphaproteobacteria, Bacteroidota and the rest of Gammaproteobacteria, which could
 202 produce vitamin B₁ and B₇) and ~23% could produce only one type of B vitamin (or ~9% none at all). This
 203 suggests that there is a major “market” for B vitamins, and indeed almost all genomes (~83%) encoded
 204 transporters for at least one vitamin.

205 A more detailed analysis of the genomes suggested that marine bacteria could be divided into three main
 206 groups based on their predicted strategy for B vitamins acquisition: (1) “consumers”, which lacked the bio-
 207 synthetic genes but harboured the vitamin transporters (we assumed transporters were for uptake, see
 208 Supplementary text for details on transporters’ directionality); (2) “independents”, which encoded the bio-
 209 synthetic pathways but not the relevant transporters; (3) “flexibles”, which encoded both the biosynthetic
 210 pathways and transporters for a specific vitamin. Bacteria possessing the latter strategy can potentially
 211 switch from being consumers to independent or vice-versa, according to what is more efficient given the
 212 surrounding conditions (e.g. availability of extracellular vitamins). The proportion of these three groups
 213 changed with the B vitamin studied and the taxonomy of the genomes (Fig. 3b). Very few genomes were
 214 flexibles for all three vitamins (~2%), and these were mostly Actinobacteriota (Fig. 3c). There were almost
 215 equal proportions of flexibles and independents for vitamin B₁ (Gammaproteobacteria and Cyanobacteria,
 216 respectively), whereas the most common bacterial strategies for vitamin B₇ and B₁₂ were independent (Fig.
 217 3b). Many different combinations of synthesis and uptake of the three vitamins were represented in the
 218 genomes (51 out of 64 possible combinations, Fig. 3c and Supplementary Fig 8a), several being enriched in
 219 specific GFCs (Supplementary Fig 9 and Supplementary text) and, notably, most strategies required the exo-
 220 genous uptake of at least one vitamin. While the perceived lack of biosynthetic capacity could be due to the
 221 utilization of precursors or to gaps in the pathway annotations (see Supplementary Fig 10 and Supplement-
 222 ary text), we speculate that this could be a potential manifestation of the “Black Queen” hypothesis, which

223 stipulates that bacteria “outsource” critical functions to the surrounding community, enabling a reduction of
 224 their metabolic cost ⁶⁸. In our dataset, the highest fraction of B-vitamin consumers, and hence putative
 225 auxotrophs, was observed for vitamin B₁, followed by B₁₂ and B₇. This order, however, does not reflect the
 226 metabolic costs of producing such vitamins, as B₁₂ would be the most “expensive” with about 20 genes in-
 227 volved ⁶⁹, whereas only 4 genes are required to synthesize B₇ ⁶³ and 5 genes for B₁ ^{70,71}. Therefore, we hypo-
 228 thesized that vitamin B₁ supplies might be more stable or frequent (e.g. as a result of higher export or lysis
 229 of producing bacteria) than that of vitamin B₁₂. Nevertheless, very few organisms were predicted to be
 230 auxotrophic for all three vitamins, suggesting that completely relying on exogenous sources for vitamins
 231 represents a risky strategy in marine pelagic environments. Taken together, these data provide a compre-
 232 hensive overview of the market for B vitamins in marine environments by defining specific roles (e.g. con-
 233 sumer, independent, flexible/source) and identifying which bacteria (taxon and GFC) fulfil each role.

234

Fig. 3: Plots of intersecting sets showing different genomic configurations of traits for the biosynthesis and transport of vitamins B₁, B₁₂, and B₇: (a) capabilities to produce or transport these vitamins, (b) different configurations to either produce and/or transport each of these vitamins, and (c) the most abundant configurations to produce and/or transport these vitamins across genomes; the remaining combinations are shown in Supplementary Fig. 8a. Overall, the left bar chart indicates the total number of genomes for each trait, the dark connected dots indicate the different configurations of traits and the waffle bar chart indicates the number (and percentage) of genomes provided with such a configuration; each piece of a waffle bar represents a genome and it is coloured according to the taxon.

Production of siderophores and phytohormones – key mechanisms of synergistic microbial interactions

The production and exchange of “common goods” such as siderophores⁷², as well as of specific phytohormones like auxin⁷³ represent traits that may mediate synergistic microbial interactions. As shown in Fig. 2, approximately 10% of the genomes have the capacity to produce siderophores (mainly Actinobacteriota and Gammaproteobacteria), while almost half of the genomes, from multiple taxa, encoded siderophore transporters (45% of the genomes). The occurrence of siderophores biosynthetic traits was partially following the GFC clustering (e.g. GFC 8, Actinobacteriota and GFC 25, Gammaproteobacteria) and partially scattered across single genomes in different taxa. In contrast, the distribution of the transporters mainly followed the GFC grouping (Fig. 2). Furthermore, microorganisms can utilize siderophore-bound iron also without the need for siderophore transporters, e.g. using ferric reductases located on the plasma membrane⁷⁴ or via direct endocytosis⁷⁵. In this regard, 5% of the genomes encoded the capacity to produce vibrioferrin (Supplementary Fig. 11), which is available to a wide range of organisms upon photolysis²¹. Field studies revealed that siderophore biosynthesis is widespread in the ocean⁷⁶, and that bacteria producing e.g. vibrioferrin can represent a relevant percentage of the total bacterial communities⁷⁷. Thus, siderophores can be considered as “keystone molecules” (*sensu*⁷⁸), produced by a limited subset of organisms but utilizable by a wide range of bacteria^{79–81}.

Several recent studies have shown that bacteria can influence the growth of phytoplankton through the production of phytohormones^{13,18,24}, and indeed the auxin hormone indole-3-acetic acid (IAA) has been identified in natural marine samples¹³. Nearly all genomes (~92%) in our dataset are predicted to produce IAA. Four pathways for the production of IAA were identified, with some organisms encoding more than one pathway. The tryptamine pathway was the most common one and was present in nearly all GFCs comprising genomes of Alphaproteobacteria, Gammaproteobacteria, Cyanobacteria and Actinobacteriota (Fig. 2). The indole-3-pyruvate pathway was the second most common with an almost identical distribution to the tryptamine pathway (missing in GFCs 36 and 15, pico-Cyanobacteria, and 28, Alphaproteobacteria), whereas the last two pathways were rarer (<10% of genomes) and limited to Alphaproteobacteria (indole-3-

acetonitrile) and some genomes of Cyanobacteria and Actinobacteriota (indole-3-acetamide). It is tempting to speculate that the widespread distribution of the capacity to produce IAA, and the diversity of biosynthetic pathways, suggest that many heterotrophic bacteria can directly increase phytoplankton growth through specific signalling (e.g. ^{13,18}). However, all pathways for IAA production are tightly intertwined with the metabolism of tryptophan, either involved in tryptophan catabolism (to cleave the amino group for nitrogen metabolism) or as a “release valve” to avoid the accumulation of toxic intermediates (e.g. α -keto acid indolepyruvate and indoleacetaldehyde). Additionally, IAA can be catabolized as a carbon source for growth (see ⁸² and references therein). Given the wide distribution of the tryptamine and indole-3-pyruvate pathways, we hypothesize that they might be more linked to the metabolism/catabolism of tryptophan, whereas the indole-3-acetonitrile and indole-3-acetamide pathways could be responsible for the production of IAA involved in phytoplankton-bacteria interactions. This hypothesis is supported by the presence of the latter traits in GFCs 9 and 40 that group model organisms known to interact through auxin with phytoplankton (Fig. 2) ^{13,18,24}, and by the fact that genes specifically related to these two pathways were found to be up-regulated in one of those studies ¹³.

Traits underlying potential antagonistic interactions

Experimental measurements of interactions among marine bacteria suggest that antagonism is common (>50% of the tested isolates) ^{32,34}, but in most cases the mechanisms behind antagonistic interactions are unclear. Antimicrobial compounds may underlie many antagonistic interactions in marine environments (e.g. ^{83–85}), and indeed genes encoding for the production of such compounds were found in several bacteria in our dataset (Supplementary Tables 5 and 6). Interestingly, antimicrobial compounds were predicted to be produced in ~30% of genomes (inner ring in Fig. 2), including also GFCs poor in other interaction traits. The most abundant traits across GFCs were bacteriocin and betalactone production ^{86,87} (Supplementary Fig. 12). Genes involved in the resistance to antimicrobial compounds were also relatively common (78% of genomes; Fig. 2), however, along with “specific” resistance traits (e.g., efflux pumps specific for antibiotics), we

noticed many KEGG modules that were annotated as “antimicrobial compounds resistance” but were also involved in other cellular functions (e.g. cell division, protein quality control and transport of other compounds; Supplementary Table 5)^{88,89}. All GFCs which grouped genomes of Cyanobacteria, Actinobacteriota, and Bacteroidota possessed only these “generic” resistance traits (Supplementary Fig. 12), suggesting that such clades are less efficient in resisting microbial “chemical warfare”. In support of this hypothesis, some Cyanobacteria strains are indeed used as markers for antibiotic contamination because of their sensitivity (e.g., references^{90,91}), and Bacteroidota are often inhibited when co-cultured with other bacteria that express antagonistic behaviour^{32,34}. Overall, these genome-based predictions are in agreement with previous experimental results^{32,34}, which suggested that Alpha- and Gammaproteobacteria commonly inhibited other bacteria, whereas Bacteroidota had a low inhibitory capacity and were the most sensitive to inhibition by other bacteria.

Antimicrobial compounds or toxins often need to be delivered into the target organism, e.g. using type IV or type VI secretion systems (T4SS and T6SS, respectively). Approximately 24% of the strains encoded T4SS or T6SS, and these were found primarily in GFCs containing Alpha- and Gammaproteobacteria (Fig. 2). The two secretion systems had different distributions among the GFCs, with only GFC 25 and 31 (comprising *Vibrio* and *Burkholderia*, genera of Gammaproteobacteria) bearing both systems. The T4SS system can perform multiple roles, including conjugation, DNA exchange and toxin delivery in bacteria-bacteria or bacteria-eukaryote interactions⁹². T4SSs were detected more frequently in Alphaproteobacteria (5 out of 8 GFCs). To date, T6SSs are known to be involved only in antagonistic interactions, including among marine bacteria⁹³, suggesting that the presence of this trait is a high-confidence predictor of the ability to directly inhibit other cells⁹⁴ and references therein). In our dataset, T6SSs occurred almost exclusively in GFCs comprising Gammaproteobacteria, specifically in *Marinobacter* and *Vibrio*, suggesting a strong capacity for contact-mediated antagonistic interactions in these taxa. Type III secretion systems (T3SS), which deliver effector molecules that maintain the bacterial association with the host⁹⁵, were found only in a few genomes as the *Vibrio* clustered in GFC 25. This GFC grouped known zooplanktonic hosts⁵⁵, suggesting a more specific role for T3SS in metazoan host-microbe interactions.

313

314 **Linked trait clusters (LTCs) delineate functional connectivity between in-** 315 **dividual interaction traits**

316 While individual traits may be important in determining the outcome of microbial interactions, such interac-
317 tions are often highly complex, requiring multiple traits such as motility, signalling and metabolic interac-
318 tions to operate together (e.g. ^{13,18,20,23,24}). If these interaction mechanisms are evolutionarily conserved,
319 traits that are functioning together to mediate such interactions should co-evolve, meaning that selection
320 would favour maintaining all relevant traits in the same genome ^{96,97}. To identify cases of co-evolving traits,
321 we used linkage disequilibrium analysis and clustered traits which were found together more often than ex-
322 pected by chance (adjusted p-value < 0.05) into Linked Trait Clusters (LTCs; Figure 1, Supplementary Table 7
323 and Supplementary Fig 13). For example, LTC 10 includes pathways for assimilatory sulfate reduction, siro-
324 heme and heme biosynthesis, as well as vitamins B₁ and B₇ biosynthesis. These traits appeared together
325 more often than random pairs of traits (mean r within this LTC is 0.38, compared to 0.09 among all traits;
326 Supplementary Fig. 13b) and they are also functionally linked. In fact, siroheme is a prosthetic group for as-
327 similatory sulfite reductases ^{98,99} and, in sulfate-reducing bacteria, siroheme can be "hijacked" for the bio-
328 synthesis of heme ¹⁰⁰. Finally, once reduced, sulphur can be incorporated into essential molecules such as
329 amino acids (methionine and cysteine) and membrane lipids, as well as into vitamins B₁ and B₇ ¹⁰¹.

330 We focus first on LTCs grouping traits not directly involved in interactions. Two LTCs, 3 and 5 (mean r of 0.30
331 and 0.35), occurred in nearly all genomes (>93%; Fig. 1) and, as expected, they linked traits which mediate
332 for core metabolic functions, common to any cell. These include biosynthesis of nucleotide (DNA and RNA)
333 and amino acids, as well as core metabolic pathways (glycolysis, pentose phosphate pathway and the first
334 three reactions of the TCA cycle). In contrast, other LTCs (i.e. 2, 4 and 7) highlighted cases in which major
335 metabolic pathways such as the TCA cycle or pathways for the cell wall assembly were missing in specific
336 bacterial clades. These absence-patterns were consistent with previous studies (see Supplementary Text for

337 detailed description). The LTC concept can, therefore, be used to identify traits that may function together,
338 providing hypotheses as to new modes of interaction that can be tested experimentally.

339 Analysis of LTCs which include interaction traits such as secretion systems suggested that these are linked
340 also to traits encoding for chemotaxis, motility and adhesion, a typical set of traits a bacterium would need
341 to locate, reach and settle on an organic matter particle or eukaryotic hosts (phytoplankton, zooplankton,
342 fish). Moreover, other genetic traits (e.g. metabolic or regulatory) were linked within these LTCs and they
343 may also be involved in microbial interactions (Fig. 4). For example, LTC 11 (mean $r = 0.40$) included, in addi-
344 tion to T6SS, traits for adhesion, a flagellar regulatory system, quorum sensing (controlling for swarming and
345 biofilm formation), chemotaxis and a nitrogen transporter with the regulation system. The same LTC also
346 encoded for the biosynthesis of ubiquinone, vitamins B₆ (pyridoxal) and B₇, and for two regulatory systems
347 (*BarA-UvrY*, *RstB-RstA*), which are known to regulate virulence, metabolism, biofilm formation, stress resist-
348 ance, quorum sensing and secretion systems (see ^{102,103}). This LTC was common in the GFCs grouping
349 Gammaproteobacteria such as *Pseudalteromonas* (GFC 21) *Alteromonas* (GFC 24), *Marinobacter* (GFC 35),
350 *Shewanella* (GFC 37) and *Vibrio* (GFCs 25 and 47; Supplementary Tables 3 and 7). All of these organisms are
351 known as particle and phytoplankton associated bacteria (e.g. ^{44–47,56}), and in such micro-environments they
352 can engage in microbiome interactions using these linked traits (i.e. biosynthesis of B vitamins, quorum
353 sensing and T6SS; ^{93,104}). Interestingly, the other two secretion systems, T4SS and T3SS, were also linked with
354 the regulation of nitrogen metabolism and with vitamin B₇ or B₁₂ transport as part of LTC 25 and LTC 17, re-
355 spectively. We propose that this provides further support for a putative functional link between microbial
356 interactions (injection of effector molecules), nitrogen and vitamin metabolism.

357 Notably, LTC 17 (which encodes the T3SS; mean $r = 0.47$) included also amino acid and sugar transporters,
358 and two regulation systems (*UhpB-UhpA*, *CitA-CitB*). The *UhpB-UhpA* genes control the motility and coloniz-
359 ation of fish pathogens ¹⁰⁵ and, not surprisingly, the LTC was found in GFCs 46 and 47 which included known
360 pathogenic bacteria (grouping *Aeromonas* and *V. natriegensis* genomes; Supplementary Tables 3 and 7)
361 ^{106,107}. LTC 17 was also found in GFCs that grouped non-pathogenic but still host associated taxa (GFC 25,
362 which includes *V. alginolyticus* and GFC 13 grouping Enterobacteriaceae) ¹⁰⁸, supporting the role of this LTC

363 (as well as of the T3SS it encodes) in interactions with eukaryotic hosts. Moreover, *CitA-CitB* regulation sys-
364 tem controls the citrate metabolism in response to changes of amino acid concentration or pH ¹⁰⁹. We spec-
365 ulate that the link between microbial interaction traits, citrate metabolism and protein and amino acid
366 transporters within LTC 17 may be relevant for a host-associated bacteria during the dispersal stage (lower
367 amino acid concentration and change in pH changes).

368 Finally, the analysis of LTC 19 (mean $r = 0.34$) lent further support for the hypothesis that some IAA produc-
369 tion pathways are involved in phytoplankton-bacteria interactions and not just in the tryptophan catabolism
370 (see above). This LTC included the indole-3-acetonitrile and indole-3-acetamide pathways, along with other
371 key microbial-phytoplankton interaction traits such as quorum sensing ^{24,110} and taurine degradation ^{13,16}.
372 Other linked genetic traits hint to additional molecular mechanisms: the manganese/iron transporter sug-
373 gests a micronutrient-dependent response, the transport of capsular polysaccharide may be involved in res-
374 istance to host defence and pathogenicity ¹¹¹, and the biosynthesis of putrescine can stimulate phytoplank-
375 ton growth, productivity and stress tolerance, as shown previously in plants ¹¹². Moreover, the LTC was
376 found complete in GFCs 9, 30 and 40 (Supplementary Table 7) which grouped genomes of bacteria known
377 to interact via IAA with phytoplankton (e.g. *Dinoroseobacter*, *Sulfitobacter* and *Phaeobacter*) ^{13,18,24,113}. We
378 propose that possessing this LTC indicates that the relevant bacterium is capable of affecting the growth of
379 phytoplankton through a combination of specific hormone signalling pathways and metabolic interactions.

380

20

pathogenicity). A solid arrow is used when the mediated mechanism has already been described in literature, while a dashed arrow indicates a potential new mechanism.

381

382 **Summary and outlook**

383 We present a framework that extrapolates from studies of specific model organisms to predict the interac-
 384 tion potential of other bacteria based on the traits encoded in their genomes. By focusing on biologically
 385 relevant traits (including specific interaction traits), we reduced a highly complex genomic dataset to a tract-
 386 able matrix of organisms by functions. By organizing the ensuing genomic information into GFCs, we further
 387 simplified the interpretation of complex genomic datasets, while at the same time highlighting the non-
 388 trivial grouping of organisms by phenotypic traits, sometimes irrespective of taxonomic boundaries. The
 389 LTCs provided evidence for the functional and evolutionary linkage between traits, raising hypotheses as to
 390 how these traits act together in the context of complex processes such as microbial interactions. This ap-
 391 proach can be easily scaled to different ecosystems (e.g. freshwater or terrestrial) and expanded to include
 392 information from additional data sources (e.g. metabolomics or high-throughput functional assays).
 393 Applying this approach to a wide diversity of bacterial taxa, we showed that marine bacteria encode differ-
 394 ent configurations of interaction traits and can, therefore, engage in different interaction types. Known
 395 particle associated taxa of Alpha- and Gammaproteobacteria possessed the full set of traits to interact with
 396 particles and living hosts, while Bacteroidota, a known ubiquitous copiotroph taxon, did not have this capa-
 397 city. Actinobacteriota and Cyanobacteria represented potential sources for B₇ in the B vitamin market, while
 398 most Alphaproteobacteria appeared as obligate customers. We suggest that siderophores, and vibrioferrin
 399 in particular, are keystone molecules being produced by only a few bacteria (Actinobacteriota and
 400 Gammaproteobacteria) but affecting a much larger diversity of potential users, in agreement with the Black
 401 Queen Hypothesis⁶⁸. Finally, the production of IAA might be more common than expected, and in some
 402 cases (e.g. GFCs 9, 30 and 40 encoding LTC 19) this may be linked with other traits involved in phytoplank-
 403 ton-bacteria interactions.

404 As with any bioinformatic approach, also our workflow aimed to identifying functional traits on genome se-
 405 quences, has inherent limitations. Firstly, on average, only 63% of genes were functionally annotated across
 406 genomes (range ~40-80%), with the remaining genes either annotated as hypothetical or not annotated at
 407 all. This is a strong reminder of the limitations of current genomic and metabolic knowledge. Secondly, only
 408 a fraction of the natural microbial diversity was represented by high-quality sequenced genomes, leading to
 409 a bias in the representation of different clades in the analysed dataset (Figure 1). While the analysed gen-
 410 omes did represent a significant fraction of natural diversity (Supplementary Figs 4 and 5), a future work,
 411 both culture dependent and independent, is needed to close these gaps and obtain an unbiased view of the
 412 traits encoded by marine microorganisms. Finally, the GFC and LTC concepts are both statistical in nature,
 413 representing the probability of bacteria having similar functional capacity, and of traits being functionally
 414 and/or evolutionarily linked. In support of the GFC concept, a study of marine Vibrionaceae suggested that
 415 ecologically-cohesive unit (similar to GFCs) are likely to interact in a similar way ³⁰. However, some interac-
 416 tion phenotypes, such as the ability to inhibit multiple target bacteria (“super-killers”), are not phylogenetic-
 417 ally conserved ^{30,114}. Experimental studies using both established and new model systems across multiple
 418 scales of diversity (e.g. between and within GFCs) are needed to test the GFC framework, whereas genetic
 419 manipulation of linked traits to test their effect on microbial interactions will be required to determine to
 420 what extent traits within LTCs are functionally linked. Nevertheless, GFCs and LTCs describe how putative in-
 421 teraction traits vary across microbial diversity, and thus can be used to quantify how the fraction of natural
 422 communities potentially “ready to interact” changes over space and time. In turn, this should help to elucid-
 423 ate fundamental rules that govern community dynamics and assembly in the oceans, and the roles played
 424 by microbial interactions will in global ecosystem-level processes and biogeochemistry.

425

426

427 **Materials and Methods**

428 **Genome selection**

429 A dataset of complete and high-quality draft genomes of marine bacteria was compiled performing an ex-
 430 tensive research on metadata available from NCBI (<http://www.ncbi.nlm.nih.gov/genome>), JGI (<https://img.jgi.doe.gov/cgi-bin/m/main.cgi?section=FindGenomes&page=genomeSearch>), and MegX (<https://mb3is.megx.net/browse/genomes>) websites. Although the focus of the analysis was on bacteria inhabiting
 433 the marine pelagic environment, some genomes from organisms isolated in extreme marine environments
 434 (i.e. thermal vents, saline and hypersaline environments, estuaries) and sediment, as well as from human
 435 and plant symbionts (*Sinorhizobium* and *Mesorhizobium*) were kept for comparison. The final list of 473
 436 genomes includes all of the genomes that, using CheckM ¹¹⁵, were defined “closed” (i.e. each DNA molecule,
 437 such as chromosome and plasmids, was represented as a single sequence in the fasta file) or high-quality
 438 draft genomes (>90% of completeness, <10% of contaminations, >18 tRNA genes and all 3 rDNAs present)
 439 ¹¹⁶. The final dataset included 473 complete genomes with 117 closed genomes and with >81% genomes
 440 that were >99% complete. Of these 473 genomes, 421 were isolated in marine pelagic and coastal zones, 34
 441 in extreme environments (e.g. salt marsh or hydrothermal vent), 6 in marine sediment and, of the remain-
 442 ing, 8 were human associated and 4 plant roots associated (Supplementary Table 3).

443 **Genome annotation**

444 Genome taxonomic classification was obtained using the protein phylogeny workflow implemented in
 445 GTDB-tk v1.4.0 ⁵⁰ with the command *classify_wf* (standard settings), and all retrieved genomes were func-
 446 tionally re-annotated using a standardized pipeline. In brief, gene calling and first raw annotation steps were
 447 performed with Prokka v1.14.5 (standard settings and ‘--rnammer’ for rRNA prediction) ¹¹⁷. The amino acid
 448 sequences translated from the identified coding DNA sequences of each genome were annotated against
 449 pre-computed hidden Markov model profiles of KEGG Ortholog (KEGG database v94.0) using kofamscan
 450 v1.2.0 (standard settings) ¹¹⁸. Additional targeted analyses were performed to annotate secondary metabol-
 451 ites, phytohormones, specific transporters and utilization of sulfur-metabolites. The genbank files generated

by Prokka were submitted to a local version of Anti-SMASH v5.1.2 (--clusterblast --subclusterblast --known-clusterblast --smcogs --inclusive --borderpredict --full-hmmer --asf --tta), which generated a list of predicted secondary metabolite biosynthesis gene clusters ¹¹⁹. Different pathways for the biosynthesis of the phytohormone indole-3-acetic acid (IAA) were manually identified by annotating (blastp v2.9.0 best hit, e-value <1e-5 and bit score > 60 ¹²⁰) the translated amino acid sequences against the KEGG orthologies required to generate IAA from tryptophan in the KEGG map01070 (Supplementary Table 8; Supplementary file 3). Translated amino acid sequences were also used as input for a GBlast search (BioV suite; standard settings) ¹²¹ to identify trans-membrane proteins mediating the transport of B vitamins and siderophores (Supplementary Table 4). For downstream analysis, only annotations with a trans-membrane alpha-helical overlap score >1 and a blast e-value <1e-6 were retained. Production and transport of photoactive siderophores (i.e. vibrioferrin) ²¹ were identified with functional annotations from the most similar (BLAST best hit) sequences. The predicted protein sequences were blasted (blastp v2.9.0; e-value <1e-5 and amino acid similarity >30%) ¹²² against a reference dataset assembled using all available sequences of related genes (pvsABCDE and pvuBCDE operons) ¹²³ available in UniProt (Supplementary Table 9 and Supplementary file 4). The same approach was carried out to annotate the dimethylsulfoniopropionate (DMSP) degradation pathways of demethylation and cleavage blasting (e-value <1e-70) against a reference dataset that contained all Uniprot sequences of the genes listed in ^{124,125} (Supplementary Table 9 and Supplementary file 5). Catabolic pathways of other two sulfur-metabolites, 2,3-dihydroxypropane-1-sulfonate (DHPS) and taurine, were identified checking for the presence of key reactions. The known routes to degrade DHPS share all a sulfopropanediol 3-dehydrogenase (hpsN gene; K15509) ¹⁴, while taurine can enter the TCA cycle either via taurine-pyruvate aminotransferase (tpa gene; K03851) or via taurine dehydrogenase (tauXY genes; K07255+K07256), followed by a sulfoacetaldehyde acetyltransferase (xsc gene; K03852) ¹²⁶.

KEGG module reconstruction

KEGG orthologies (KOs) annotations generated by kofamscan were recombined in KEGG modules (KMs) using an in-house R script. The KMs represent minimal functional units describing either pathways, structural

477 complexes (e.g. transmembrane pump or ribosome), functional sets (essential sets as Aminoacyl-tRNA syn-
 478 thases or nucleotide sugar biosynthesis) or signaling modules (phenotypic markers as pathogenicity). Briefly,
 479 using the R implementation of KEGG REST API ¹²⁷, the script fetches the diagrams of all KMs from the KEGG
 480 website. Each diagram represents a reactions' scheme of a KM listing all known KOs that can perform each
 481 of the reactions necessary to complete that scheme (Supplementary Fig. 1b). The completeness of a KM in
 482 a genome was assessed as the number of required reactions for which at least one KOs was annotated and
 483 only complete KMs were retained in downstream analyses (e.g., a KM with 7 out of 8 reactions is incom-
 484 plete and would be discarded). However, to partially compensate for possible annotation issues, one miss-
 485 ing reaction was allowed in KMs with ≥ 3 reactions (i.e., a KM with 7 out of 8 annotated reactions is con-
 486 sidered complete; see Supplementary text and Supplementary Table 10 for more details).

487 Genetic and interaction traits identification

488 The annotated complete KMs, secondary metabolites, phytohormones and transporters represent the ge-
 489 netic traits identified in the genomes (556 genetic traits). From this list, the subset of interaction traits was
 490 manually extracted based on current knowledge about processes that likely play a role in microbial interac-
 491 tions (list of picked interaction traits in Supplementary Table 2). Within the KMs we identified traits related
 492 to vitamin biosynthetic pathways, quorum sensing, chemotaxis, antimicrobial resistance, motility and adhe-
 493 sion (Supplementary Table 5). Since the ecological role of most secondary metabolites is still unclear, a care-
 494 ful literature search was performed to identify and retain only the secondary metabolite clusters with a pro-
 495 posed function which can be linked to microbial interaction processes, such as siderophore production,
 496 quorum sensing and antimicrobial compound biosynthesis (Supplementary Table 6). The phytohormone an-
 497 notations revealed the capability of producing indoleacetic acid (auxin) through four different pathways
 498 (Supplementary Table 8). Vitamin and siderophore transporters were identified in the transporter annota-
 499 tions looking for the related transporter families (e.g. TonB, Btu) and the substrate information (Supple-
 500 mentary Table 4).

Mapping to environmental datasets

The selected datasets represent amplicon time series generated with Illumina sequencing of the V4 region of the 16S rDNA of bacterial communities sampled in a coastal (between 1-5m of depth, Canoe Cove, Nahant, MA, USA)⁵⁹ and a pelagic (between 10-500m of depth, n-1200 station, Easter Mediterranean sea)⁶⁰ site. Mapping between the full 16S rDNAs extracted from genomes and amplicon sequences was performed with BLAST (blastn v2.9.0, e-value <1e-5) using different identity threshold to filter the best hits: 100%, 97%, 94.5% and 86.5%. The first two values represent a new proposed threshold and the most commonly used threshold to define operational taxonomic units (OTUs)¹²⁸, while the last two values were suggested as threshold to classify OTUs at genus and family levels¹²⁹. In addition, for each identity threshold we inspected the top blast hits (up to 20) and calculate a mapping specificity index as the number of hits assigned to a same GFC over the total number of hits (Supplementary Fig.s 4a and 5a). To avoid spurious mapping, we only retained in downstream analyses sequences with a specificity index = 1 (i.e. all best blast hits belonged to the same GFC).

The evaluation of the GFC concept was performed by using the coastal time series, because it offers a high (daily) temporal resolution of changes in the bacterial community composition and the authors applied a time deconvolution analysis to characterize the OTU temporal dynamics⁵⁹. The OTUs included in the deconvolution analysis represented ~97% of the total sequences in the dataset and for each pair of OTUs, the authors calculated the frequency interaction score (the higher the score, the more synchronous were the temporal dynamics of both OTUs, and vice-versa). From the list of all pairs, we only considered OTU pairs for which at least one of the two OTUs mapped to a GFC. We wanted to exclude cases of OTUs pairs that could belong to a same, but yet unknown, GFC. To test for the GFC concept, we compared the frequency interaction score between OTU pairs mapped to the same OTU (i.e. both OTUs mapped to the same GFC) versus OTU pairs mapped to different GFCs (e.g. the two OTUs mapped to different GFCs, or one of the OTU was unmapped). For each identity threshold, the normal distribution of the frequency interaction score was assessed with the Shapiro test (r's function *shapiro.test*; p.values << 0.001), and t-test (r's function *t.test*) was performed to test for a significant difference in the mean ranks between the two groups of OTU pairs. We

repeated the test by randomly assigning OTU pairs to the “same GFC” and “different GFC” groups by keeping the same group sizes and by creating two groups of equal size.

Statistical analyses

The presence/absence matrix of genetic traits across genomes served as basis to cluster the former into linked trait clusters (LTCs) and the latter into genome functional clusters (GFCs). Both clustering approaches implemented the Pearson coefficient r (also known as phi coefficient when applied to dichotomous variables) to calculate the genome and trait correlation matrices. While no pair of genomes scored a negative r value (as they all shared some “core” functional traits), negative correlations between pairs of genetic traits were threshold to zeros to ensure that the trait clustering was only driven by positive correlation. Moreover, only pairwise correlations with a FDR-corrected p -value < 0.05 (chi-square test, $df = 1$) were retained. The parsed correlation matrices were fed as similarity matrices into the affinity propagation algorithm implemented with the *apcluster* function ($q=0.5$; *r* package *apcluster*)¹³⁰. This machine learning algorithm was chosen because it does not require the number of clusters to be determined *a priori*, allowing instead this feature to emerge from the data¹³¹. Briefly, a functional similarity matrix is used to construct a network where nodes and edges are known to be genomes (or genetic traits) and their pairwise r correlation, respectively. Starting from a random set of exemplar nodes, clusters are created by expansion towards the adjoining, most similar, nodes. Through iterations of this procedure, the algorithm tries to maximize the total similarity between nodes within each cluster, eventually converging towards the best set of clusters. Clustering robustness and accuracy of both GFCs and LTCs were tested by performing a sensitivity analysis of the ‘ q ’ parameter which controls the clustering sensitivity in the *apcluster* function, and by down-sampling overrepresented taxa at 80%, 60% and 40% of their genome coverage (see Supplementary information and Supplementary Fig. 15).

For the LTC delineation, genetic traits were pre-filtered to remove noisy signals and only traits found in $\geq 3\%$ of the genomes (≥ 14 genomes) were used in the clustering (i.e. 379 genetic traits out of 578 in total). Similar to the context of linkage disequilibrium¹³², r indicates non-random association between genetic traits

because those traits are interactively linked to fitness, or simply because they are closely located on the chromosome (i.e. lower chances of recombination). However, as the genetic traits analysed commonly involve multiple genes, the second possibility is less likely. While exploring the functional potential, a LTC was considered complete in a genome when >50% of the grouped genetic traits were present and it was considered complete in a GFC when the average completeness of the included genomes was >50%. The genetic traits belonging to LTCs that were never found to be complete in at least one GFC were considered as unclustered.

All analyses were performed in R 4.0.4¹³³.

Authors' contributions

LZ, DS and H-PG designed the research. LZ and DS analyzed the data. TM contributed with analytic tools. LZ wrote the paper with substantial revisions from DS, DSegrè and H-PG. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Availability of data and material

All genomes are available in the online repositories of NCBI and JGI under the accession codes listed in Supplementary Table 3. The codes for functional annotation, reconstruction of KEGG modules and statistical analysis are available on GitHub (https://github.com/lucaz88/genome_comparison_code).

Acknowledgements

This work was supported by the Human Frontier Science Program (HFSP) through the grant number RGB 0020/2016. We thank Anna Godhe, Mats Töpel and Oskar Johansson for providing early access to some of the genomes included in the analysis. We thank the entire IAMM team (Dikla Aharonovich, David Bernstein,

575 Falk Eigemann, Elena Forchielli, Tal Luzzatto-Knaan, Shany Ofaim, Melissa Osborne, Solvig Pinnow, Dalit
576 Roth-Rosenberg, Angela Vogts and Maren Voss) for the constructive discussions and ideas that led us to
577 shape this manuscript. A sincere thank you to Silvia Munari for her diligent proofreading of this manuscript.

578

579 References

- 580 1. Hibbing, M. E., Fuqua, C., Parsek, M. R. & Peterson, S. B. Bacterial competition: Surviving and thriving
581 in the microbial jungle. *Nature Reviews Microbiology* vol. 8 15–25 (2010).
- 582 2. Zimmerman, A. E. *et al.* Metabolic and biogeochemical consequences of viral infection in aquatic
583 ecosystems. *Nat. Rev. Microbiol.* **18**, 21–34 (2020).
- 584 3. Bjorbækmo, M. F. M., Evenstad, A., Røsæg, L. L., Krabberød, A. K. & Logares, R. The planktonic protist
585 interactome: where do we stand after a century of research? *ISME J.* **14**, 544–559 (2020).
- 586 4. Amin, S. A., Parker, M. S. & Armbrust, E. V. Interactions between diatoms and bacteria. *Microbiol.*
587 *Mol. Biol. Rev.* **76**, 667–84 (2012).
- 588 5. Seymour, J. R., Amin, S. A., Raina, J.-B. & Stocker, R. Zooming in on the phycosphere: the ecological
589 interface for phytoplankton–bacteria relationships. *Nat. Microbiol.* **2**, 17065 (2017).
- 590 6. Farooq Azam and Francesca Malfatti. Microbial structuring of marine ecosystems. *Nat. Rev. Micro-*
591 *biol.* **5**, 782–791 (2007).
- 592 7. Zoccarato, L. & Grossart, H.-P. Relationship Between Lifestyle and Structure of Bacterial Communities
593 and Their Functionality in Aquatic Systems. in *Advances in Environmental Microbiology - The Struc-*
594 *ture and Function of Aquatic Microbial Communities* (ed. Hurst, C. J.) 13–52 (Springer Nature Switzer-
595 land AG 2019, 2019). doi:10.1007/978-3-030-16775-2_2.
- 596 8. Kirchman, D. L. *Processes in Microbial Ecology*. (Oxford University Press, 2012). doi:10.1093/
597 acprof:oso/9780199586936.001.0001.
- 598 9. Worden, A. Z. *et al.* Rethinking the marine carbon cycle: Factoring in the multifarious lifestyles of mi-
599 crobes. *Science (80-.)*. **347**, 1257594 (2015).
- 600 10. Gibert, J. P. Temperature directly and indirectly influences food web structure. *Sci. Rep.* **9**, 5312
601 (2019).
- 602 11. Levine, J. M., Bascompte, J., Adler, P. B. & Allesina, S. Beyond pairwise mechanisms of species coex-
603 istence in complex communities. *Nature* **546**, 56–64 (2017).
- 604 12. Bell, W. & Mitchell, R. Chemotactic and Growth Responses of Marine Bacteria to Algal Extracellular
605 Products. *Biol. Bull.* **143**, 265–277 (1972).
- 606 13. Amin, S. A. *et al.* Interaction and signalling between a cosmopolitan phytoplankton and associated
607 bacteria. *Nature* **522**, 98–101 (2015).

- 608 14. Durham, B. P. *et al.* Cryptic carbon and sulfur cycling between surface ocean plankton. *Proc. Natl.*
609 *Acad. Sci.* **112**, 453–457 (2015).
- 610 15. Durham, B. P. *et al.* Sulfonate-based networks between eukaryotic phytoplankton and heterotrophic
611 bacteria in the surface ocean. *Nat. Microbiol.* **4**, 1706–1715 (2019).
- 612 16. Moran, M. A. & Durham, B. P. Sulfur metabolites in the pelagic ocean. *Nature Reviews Microbiology*
613 vol. 17 665–678 (2019).
- 614 17. Paul, C., Mausz, M. A. & Pohnert, G. A co-culturing/metabolomics approach to investigate chemically
615 mediated interactions of planktonic organisms reveals influence of bacteria on diatom metabolism.
616 *Metabolomics* **9**, 349–359 (2013).
- 617 18. Segev, E. *et al.* Dynamic metabolic exchange governs a marine algal-bacterial interaction. *Elife* **5**,
618 e17473 (2016).
- 619 19. Christie-Oleza, J. A., Sousoni, D., Lloyd, M., Armengaud, J. & Scanlan, D. J. Nutrient recycling facilit-
620 ates long-term stability of marine microbial phototroph–heterotroph interactions. *Nat. Microbiol.* **2**,
621 17100 (2017).
- 622 20. Wang, H., Tomasch, J., Jarek, M. & Wagner-Döbler, I. A dual-species co-cultivation system to study
623 the interactions between Roseobacters and dinoflagellates. *Front. Microbiol.* **5**, 311 (2014).
- 624 21. Amin, S. A. *et al.* Photolysis of iron-siderophore chelates promotes bacterial-algal mutualism. *Proc.*
625 *Natl. Acad. Sci.* **106**, 17071–17076 (2009).
- 626 22. Keshtacher-Liebso, E., Hadar, Y. & Chen, Y. Oligotrophic Bacteria Enhance Algal Growth under Iron-
627 Deficient Conditions. *Appl. Environ. Microbiol.* **61**, 2439–2441 (1995).
- 628 23. van Tol, H. M., Amin, S. A. & Armbrust, E. V. Ubiquitous marine bacterium inhibits diatom cell divi-
629 sion. *ISME J.* 1–12 (2016) doi:10.1038/ismej.2016.112.
- 630 24. Seyedsayamdost, M. R., Case, R. J., Kolter, R. & Clardy, J. The Jekyll-and-Hyde chemistry of *Phaeobac-*
631 *ter gallaeciensis*. *Nat. Chem.* **3**, 331–335 (2011).
- 632 25. Grossart, H.-P. *et al.* Interactions between marine snow and heterotrophic bacteria: Aggregate form-
633 ation and microbial dynamics. *Aquat. Microb. Ecol.* **42**, 19–26 (2006).
- 634 26. Aharonovich, D. & Sher, D. Transcriptional response of *Prochlorococcus* to co-culture with a marine
635 *Alteromonas*: differences between strains and the involvement of putative infochemicals. *ISME J.* 1–
636 15 (2016) doi:10.1038/ismej.2016.70.
- 637 27. Coe, A. *et al.* Survival of *Prochlorococcus* in extended darkness. *Limnol. Oceanogr.* **61**, 1375–1388
638 (2016).
- 639 28. Hou, S. *et al.* Benefit from decline: The primary transcriptome of *Alteromonas macleodii* str. *Te101*
640 during *Trichodesmium* demise. *ISME J.* **12**, 981–996 (2018).
- 641 29. Biller, S. J., Berube, P. M., Lindell, D. & Chisholm, S. W. *Prochlorococcus*: the structure and function of
642 collective diversity. *Nat. Rev. Microbiol.* **13**, 13–27 (2015).

- 643 30. Cordero, O. X. *et al.* Population Genomics of Early Events in the Ecological Differentiation of Bacteria.
644 *Science* (80-.). **336**, 48–51 (2012).
- 645 31. Tai, V., Paulsen, I. T., Phillippy, K., Johnson, D. A. & Palenik, B. Whole-genome microarray analyses of
646 *Synechococcus-Vibrio* interactions. *Environ. Microbiol.* **11**, 2698–2709 (2009).
- 647 32. Long, R. A. & Azam, F. Antagonistic interactions among marine bacteria. *Appl. Environ. Microbiol.* **67**,
648 4875–4983 (2001).
- 649 33. Sher, D., Thompson, J. W., Kashtan, N., Croal, L. & Chisholm, S. W. Response of *Prochlorococcus* eco-
650 types to co-culture with diverse marine bacteria. *ISME J.* **5**, 1125–1132 (2011).
- 651 34. Grossart, H.-P., Schlingloff, A., Bernhard, M., Simon, M. & Brinkhoff, T. Antagonistic activity of bac-
652 teria isolated from organic aggregates of the German Wadden Sea. *FEMS Microbiol. Ecol.* **47**, 387–
653 396 (2004).
- 654 35. Snel, B., Bork, P. & Huynen, M. A. Genome phylogeny based on gene content. *Nat. Genet.* **21**, 108–
655 110 (1999).
- 656 36. Espariz, M., Zuljan, F. A., Esteban, L. & Magni, C. Taxonomic identity resolution of highly phylogenet-
657 ically related strains and selection of phylogenetic markers by using genome-scale methods: The ba-
658 cillus pumilus group case. *PLoS One* **11**, 1–17 (2016).
- 659 37. Hernández-González, I. L., Moreno-Hagelsieb, G. & Olmedo-Álvarez, G. Environmentally-driven gene
660 content convergence and the *Bacillus* phylogeny. *BMC Evol. Biol.* **18**, 1–15 (2018).
- 661 38. Pachiadaki, M. G. *et al.* Charting the Complexity of the Marine Microbiome through Single-Cell Gen-
662 omics. *Cell* **179**, 1623–1635 (2019).
- 663 39. Lauro, F. M. *et al.* The genomic basis of trophic strategy in marine bacteria. *Proc. Natl. Acad. Sci.* **106**,
664 15527–33 (2009).
- 665 40. Newton, R. J. *et al.* Genome characteristics of a generalist marine bacterial lineage. *ISME J.* **4**, 784–
666 798 (2010).
- 667 41. Krause, S. *et al.* Trait-based approaches for understanding microbial biodiversity and ecosystem func-
668 tioning. *Frontiers in Microbiology* vol. 5 251 (2014).
- 669 42. Bordron, P. *et al.* Putative bacterial interactions from metagenomic knowledge with an integrative
670 systems ecology approach. *Microbiologyopen* **5**, 106–117 (2016).
- 671 43. Raina, J.-B., Fernandez, V., Lambert, B., Stocker, R. & Seymour, J. R. The role of microbial motility and
672 chemotaxis in symbiosis. *Nat. Rev. Microbiol.* (2019) doi:10.1038/s41579-019-0182-9.
- 673 44. Holmström, C. & Kjelleberg, S. Marine *Pseudoalteromonas* species are associated with higher organ-
674 isms and produce biologically active extracellular agents. *FEMS Microbiol. Ecol.* **30**, 285–293 (1999).
- 675 45. Eilers, H., Pernthaler, J., Glöckner, F. O. & Amann, R. Culturability and in situ abundance of pelagic
676 Bacteria from the North Sea. *Appl. Environ. Microbiol.* **66**, 3044–3051 (2000).
- 677 46. Lupette, J. *et al.* *Marinobacter* Dominates the Bacterial Community of the *Ostreococcus tauri* Phyco-
678 sphere in Culture. *Front. Microbiol.* **7**, 1–14 (2016).

- 679 47. Sonnenschein, E. C., Syit, D. A., Grossart, H.-P. & Ullrich, M. S. Chemotaxis of *Marinobacter adhaer-*
680 *ens* and its impact on attachment to the diatom *Thalassiosira weissflogii*. *Appl. Environ. Microbiol.* **78**,
681 6900–6907 (2012).
- 682 48. Langille, M. G. I. *et al.* Predictive functional profiling of microbial communities using 16S rRNA
683 marker gene sequences. *Nat. Biotechnol.* **31**, 814–821 (2013).
- 684 49. Maistrenko, O. M. *et al.* Disentangling the impact of environmental and phylogenetic constraints on
685 prokaryotic within-species diversity. *ISME J.* **14**, 1247–1259 (2020).
- 686 50. Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: A toolkit to classify genomes
687 with the genome taxonomy database. *Bioinformatics* **36**, 1925–1927 (2020).
- 688 51. Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science* **348**, 1261359
689 (2015).
- 690 52. Louca, S., Parfrey, L. W. & Doebeli, M. Decoupling function and taxonomy in the global ocean micro-
691 biome. *Science (80-.).* **353**, 1272–1277 (2016).
- 692 53. Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive earth’s biogeochemical
693 cycles. *Science (80-.).* **320**, 1034–1039 (2008).
- 694 54. Martiny, J. B. H., Jones, S. E., Lennon, J. T. & Martiny, A. C. Microbiomes in light of traits: A phylogen-
695 etic perspective. *Science (80-.).* **350**, aac9323 (2015).
- 696 55. Hunt, D. E. *et al.* Resource Partitioning and Sympatric Differentiation Among Closely Related Bac-
697 terioplankton. *Science (80-.).* **320**, (2008).
- 698 56. Teeling, H. *et al.* Recurring patterns in bacterioplankton dynamics during coastal spring algae blooms.
699 *Elife* **5**, (2016).
- 700 57. Arias-Andres, M., Klümper, U., Rojas-Jimenez, K. & Grossart, H. P. Microplastic pollution increases
701 gene exchange in aquatic ecosystems. *Environ. Pollut.* **237**, 253–261 (2018).
- 702 58. Galand, P. E., Pereira, O., Hochart, C., Auguet, J. C. & Debross, D. A strong link between marine mi-
703 crobial community composition and function challenges the idea of functional redundancy. *ISME J.*
704 **12**, 2470–2478 (2018).
- 705 59. Martin-Platero, A. M. *et al.* High resolution time series reveals cohesive but short-lived communities
706 in coastal plankton. *Nat. Commun.* **9**, 1–11 (2018).
- 707 60. Rosenberg, D. R. *et al.* Particle-associated and free-living bacterial communities in an oligotrophic
708 sea are affected by different environmental and anthropogenic factors. *bioRxiv* 2020.04.20.051391
709 (2020) doi:10.1101/2020.04.20.051391.
- 710 61. Grote, J. *et al.* Streamlining and core genome conservation among highly divergent members of the
711 SAR11 clade. *MBio* **3**, e00252-12 (2012).
- 712 62. Shih, P. M. *et al.* Improving the coverage of the cyanobacterial phylum using diversity-driven genome
713 sequencing. *Proc. Natl. Acad. Sci.* **110**, 1053–1058 (2013).

- 714 63. Croft, M. T., Warren, M. J. & Smith, A. G. Algae Need Their Vitamins. *Eukaryot. Cell* **5**, 1175–1183
715 (2006).
- 716 64. Romine, M. F., Rodionov, D. A., Maezato, Y., Osterman, A. L. & Nelson, W. C. Underlying mechanisms
717 for syntrophic metabolism of essential enzyme cofactors in microbial communities. *ISME J.* **11**, 1434–
718 1446 (2017).
- 719 65. Suffridge, C. P. *et al.* B Vitamins and Their Congeners as Potential Drivers of Microbial Community
720 Composition in an Oligotrophic Marine Ecosystem. *J. Geophys. Res. Biogeosciences* **123**, 2890–2907
721 (2018).
- 722 66. Sañudo-Wilhelmy, S. A. *et al.* Multiple B-vitamin depletion in large areas of the coastal ocean. *Proc.*
723 *Natl. Acad. Sci.* **109**, 14041–14045 (2012).
- 724 67. Browning, T. J. *et al.* Nutrient co-limitation at the boundary of an oceanic gyre. *Nature* **551**, 242–246
725 (2017).
- 726 68. Morris, J. *et al.* The Black Queen Hypothesis: Evolution of Dependencies through Adaptive Gene
727 Loss. *MBio* **3**, e00036 (2012).
- 728 69. Fang, H., Kang, J. & Zhang, D. Microbial production of vitamin B12: A review and future perspectives.
729 *Microb. Cell Fact.* **16**, 1–14 (2017).
- 730 70. Jurgenson, C. T., Begley, T. P. & Ealick, S. E. The Structural and Biochemical Foundations of Thiamin
731 Biosynthesis. *Annu. Rev. Biochem.* **78**, 569–603 (2009).
- 732 71. McRose, D. *et al.* Alternatives to vitamin B1 uptake revealed with discovery of riboswitches in mul-
733 tiple marine eukaryotic lineages. *ISME J.* **8**, 2517–2529 (2014).
- 734 72. Vraspir, J. M. & Butler, A. Chemistry of Marine Ligands and Siderophores. *Ann. Rev. Mar. Sci.* **1**, 43–63
735 (2009).
- 736 73. De Smet, I. *et al.* Unraveling the evolution of auxin signaling. *Plant Physiol.* **155**, 209–21 (2011).
- 737 74. Morrissey, J. & Bowler, C. Iron utilization in marine cyanobacteria and eukaryotic algae. *Front. Micro-*
738 *biol.* **3**, 43 (2012).
- 739 75. Kazamia, E. *et al.* Endocytosis-mediated siderophore uptake as a strategy for Fe acquisition in diat-
740 oms. *Sci. Adv.* **4**, eaar4536 (2018).
- 741 76. Gärdes, A. *et al.* Detection of photoactive siderophore biosynthetic genes in the marine environ-
742 ment. *BioMetals* **26**, 507–516 (2013).
- 743 77. Yarimizu, K. *et al.* Distribution of dissolved iron and bacteria producing the photoactive siderophore,
744 vibrioferrin, in waters off Southern California and Northern Baja. *BioMetals* **32**, 139–154 (2019).
- 745 78. Madhusoodanan, J. Can single molecules bind together entire ecosystems? *Proc. Natl. Acad. Sci. U.*
746 *S. A.* **113**, 10451–10452 (2016).
- 747 79. Coale, T. H. *et al.* Reduction-dependent siderophore assimilation in a model pennate diatom. *Proc.*
748 *Natl. Acad. Sci.* **116**, 23609–23617 (2019).

- 749 80. Basu, S., Gledhill, M., de Beer, D., Prabhu Matondkar, S. G. & Shaked, Y. Colonies of marine cyanobac-
750 teria *Trichodesmium* interact with associated bacteria to acquire iron from dust. *Commun. Biol.* **2**, 1–
751 8 (2019).
- 752 81. Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host interactions.
753 *Nature Reviews Microbiology* vol. 18 152–163 (2020).
- 754 82. Patten, C. L., Blakney, A. J. C. & Coulson, T. J. D. Activity, distribution and function of indole-3-acetic
755 acid biosynthetic pathways in bacteria. *Crit. Rev. Microbiol.* **39**, 395–415 (2012).
- 756 83. Long, R. A. *et al.* Antagonistic interactions among marine bacteria impede the proliferation of *Vibrio*
757 *cholerae*. *Appl. Environ. Microbiol.* **71**, 8531–8536 (2005).
- 758 84. Long, R. A., Qureshi, A., Faulkner, D. J. & Azam, F. 2-n-pentyl-4-quinolinol produced by a marine *Al-*
759 *teromonas* sp. and its potential ecological and biogeochemical roles. *Appl. Environ. Microbiol.* **69**,
760 568–576 (2003).
- 761 85. Paz-Yepes, J., Brahamsha, B. & Palenik, B. Role of a Microcin-C-like biosynthetic gene cluster in allelo-
762 pathic interactions in marine *Synechococcus*. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 12030–12035 (2013).
- 763 86. Robinson, S. L., Christenson, J. K. & Wackett, L. P. Biosynthesis and chemical diversity of β -lactone
764 natural products. *Nat. Prod. Rep.* **36**, 458–475 (2019).
- 765 87. Cotter, P. D., Ross, R. P. & Hill, C. Bacteriocins-a viable alternative to antibiotics? *Nature Reviews Mi-*
766 *crobiology* vol. 11 95–105 (2013).
- 767 88. Guo, L. *et al.* Lipid A Acylation and Bacterial Resistance against Vertebrate Antimicrobial Peptides sys-
768 temic illnesses termed enteric fevers, which are char-acterized by microorganism colonization of the
769 intes-tine, followed by systemic spread to tissues rich in. *Cell* **95**, 189–198 (1998).
- 770 89. Hinz, A., Lee, S., Jacoby, K. & Manoil, C. Membrane proteases and aminoglycoside antibiotic resist-
771 ance. *J. Bacteriol.* **193**, 4790–4797 (2011).
- 772 90. Yasser, E.-N. & Adli, A. Toxicity of Single and Mixtures of Antibiotics to Cyanobacteria. *J. Environ.*
773 *Anal. Toxicol.* **05**, 274 (2014).
- 774 91. EMEA. *European Medicines Agency. Doc ref. EMEA/CHMP/SWP/4447/00. http://*
775 *www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/10/*
776 *WC500003978.pdf. http://www.emea.eu.int* (2006).
- 777 92. Grohmann, E., Christie, P. J., Waksman, G. & Backert, S. Type IV secretion in Gram-negative and
778 Gram-positive bacteria. *Molecular Microbiology* vol. 107 455–471 (2018).
- 779 93. Guillemette, R., Ushijima, B., Jalan, M., Häse, C. C. & Azam, F. Insight into the resilience and suscept-
780 ibility of marine bacteria to T6SS attack by *Vibrio cholerae* and *Vibrio coralliilyticus*. *PLoS One* **15**, 1–
781 19 (2020).
- 782 94. Cianfanelli, F. R., Monlezun, L. & Coulthurst, S. J. Aim, Load, Fire: The Type VI Secretion System, a
783 Bacterial Nanoweapon. *Trends in Microbiology* vol. 24 51–62 (2016).
- 784 95. Macho, A. P. & Zipfel, C. Targeting of plant pattern recognition receptor-triggered immunity by bac-
785 terial type-III secretion system effectors. *Current Opinion in Microbiology* vol. 23 14–22 (2015).

- 786 96. Pellegrini, M., Marcotte, E. M., Thompson, M. J., Eisenberg, D. & Yeates, T. O. Assigning protein func-
787 tions by comparative genome analysis: Protein phylogenetic profiles. *Proc. Natl. Acad. Sci. U. S. A.* **96**,
788 4285–4288 (1999).
- 789 97. Liu, C., Wright, B., Allen-Vercoe, E., Gu, H. & Beiko, R. Phylogenetic Clustering of Genes Reveals
790 Shared Evolutionary Trajectories and Putative Gene Functions. *Genome Biol. Evol.* **10**, 2255–2265
791 (2018).
- 792 98. Siegel, L. M., Murphy, M. J. & Kamin, H. Reduced nicotinamide adenine dinucleotide phosphate-
793 sulfite reductase of enterobacteria. I. The Escherichia coli hemoflavoprotein: molecular parameters
794 and prosthetic groups. *J. Biol. Chem.* **248**, 251–264 (1973).
- 795 99. Scott, A. I., Irwin, A. J., Siegel, L. M. & Shoolery, J. N. Sirohydrochlorin. Prosthetic Group of a Sulfite
796 Reductase Enzyme and Its Role in the Biosynthesis of Vitamin B12. *Journal of the American Chemical*
797 *Society* vol. 100 316–318 (1978).
- 798 100. Bali, S. *et al.* Molecular hijacking of siroheme for the synthesis of heme and d1 heme. *Proc. Natl.*
799 *Acad. Sci.* **108**, 18260–18265 (2011).
- 800 101. Cook, A. M., Smits, T. H. M. & Denger, K. Sulfonates and Organotrophic Sulfite Metabolism. in *Micro-*
801 *bial Sulfur Metabolism* 170–183 (Springer Berlin Heidelberg, 2008). doi:10.1007/978-3-540-72682-
802 1_14.
- 803 102. Zere, T. R. *et al.* Genomic targets and features of BarA-UvrY (-SirA) signal transduction systems. *PLoS*
804 *One* **10**, e0145035 (2015).
- 805 103. Huang, L., Xu, W., Su, Y., Zhao, L. & Yan, Q. Regulatory role of the RstB-RstA system in adhesion,
806 biofilm production, motility, and hemolysis. *Microbiologyopen* **7**, e00599 (2018).
- 807 104. Rolland, J. L., Stien, D., Sanchez-Ferandin, S. & Lami, R. Quorum Sensing and Quorum Quenching in
808 the Phycosphere of Phytoplankton: a Case of Chemical Interactions in Ecology. *J. Chem. Ecol.* 1–11
809 (2016) doi:10.1007/s10886-016-0791-y.
- 810 105. Chen, J. *et al.* The UhpA mutant of Edwardsiella piscicida enhanced its motility and the colonization
811 in the intestine of tilapia. *Fish Shellfish Immunol.* **104**, 587–591 (2020).
- 812 106. Janda, J. M. & Abbott, S. L. The genus Aeromonas: Taxonomy, pathogenicity, and infection. *Clin. Mi-*
813 *crobiol. Rev.* **23**, 35–73 (2010).
- 814 107. Darshanee Ruwandeepika, H. A. *et al.* Pathogenesis, virulence factors and virulence regulation of vi-
815 brios belonging to the Harveyi clade. *Rev. Aquac.* **4**, 59–74 (2012).
- 816 108. Murthy, K. N., Mohanraju, R., Karthick, P. & Ramesh, C. Phenotypic and molecular characterization of
817 epiphytic vibrios from the marine macro algae of Andaman Islands, India. *Indian J. Geo-Marine Sci.*
818 **45**, 304–309 (2016).
- 819 109. van Mastrigt, O., Mager, E. E., Jamin, C., Abee, T. & Smid, E. J. Citrate, low pH and amino acid limita-
820 tion induce citrate utilization in Lactococcus lactis biovar diacetylactis. *Microb. Biotechnol.* **11**, 369–
821 380 (2018).

- 822 110. Wang, H. *et al.* Identification of Genetic Modules Mediating the Jekyll and Hyde Interaction of *Dinor-*
823 *oseobacter shibae* with the Dinoflagellate *Prorocentrum minimum*. *Front. Microbiol.* **6**, 1262 (2015).
- 824 111. Frosch, M., Edwards, U., Bousset, K., Krauße, B. & Weisgerber, C. Evidence for a common molecular
825 origin of the capsule gene loci in Gram-negative bacteria expressing group II capsular polysacchar-
826 ides. *Mol. Microbiol.* **5**, 1251–1263 (1991).
- 827 112. Chen, D., Shao, Q., Yin, L., Younis, A. & Zheng, B. Polyamine function in plants: Metabolism, regula-
828 tion on development, and roles in abiotic stress responses. *Frontiers in Plant Science* vol. 9 1945
829 (2019).
- 830 113. Wienhausen, G., Noriega-Ortega, B. E., Niggemann, J., Dittmar, T. & Simon, M. The exometabolome
831 of two model strains of the Roseobacter group: A marketplace of microbial metabolites. *Front. Mi-*
832 *crobiol.* **8**, 1–15 (2017).
- 833 114. Russel, J., Røder, H. L., Madsen, J. S., Burmølle, M. & Sørensen, S. J. Antagonism correlates with
834 metabolic similarity in diverse bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 10684–10688 (2017).
- 835 115. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing the
836 quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.*
837 **25**, 1043–1055 (2015).
- 838 116. Bowers, R. M. *et al.* Minimum information about a single amplified genome (MISAG) and a metagen-
839 ome-assembled genome (MIMAG) of bacteria and archaea. *Nat. Biotechnol.* **35**, 725–731 (2017).
- 840 117. Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).
- 841 118. Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive
842 score threshold. *Bioinformatics* **36**, 2251–2252 (2020).
- 843 119. Blin, K. *et al.* AntiSMASH 5.0: Updates to the secondary metabolite genome mining pipeline. *Nucleic*
844 *Acids Res.* **47**, W81–W87 (2019).
- 845 120. Zhang, P. *et al.* The distribution of tryptophan-dependent indole-3-acetic acid synthesis pathways in
846 bacteria unraveled by large-scale genomic analysis. *Molecules* **24**, 1–14 (2019).
- 847 121. Reddy, V. S. & Saier, M. H. BioV Suite - A collection of programs for the study of transport protein
848 evolution. *FEBS J.* **279**, 2036–2046 (2012).
- 849 122. Amin, S. A., Green, D. H., Waheeb, D. Al, Gärdes, A. & Carrano, C. J. Iron transport in the genus
850 *Marinobacter*. *BioMetals* **25**, 135–147 (2012).
- 851 123. Tanabe, T. *et al.* Identification and Characterization of Genes Required for Biosynthesis and Transport
852 of the Siderophore Vibrioferri in *Vibrio parahaemolyticus*. *J. Bacteriol.* **185**, 6938–6949 (2003).
- 853 124. Moran, M. A., Reisch, C. R., Kiene, R. P. & Whitman, W. B. Genomic Insights into Bacterial DMSP
854 Transformations. *Ann. Rev. Mar. Sci.* **4**, 523–542 (2012).
- 855 125. Bullock, H. A., Luo, H. & Whitman, W. B. Evolution of dimethylsulfoniopropionate metabolism in mar-
856 ine phytoplankton and bacteria. *Front. Microbiol.* **8**, 637 (2017).

857 126. Cook, A. M. & Denger, K. Metabolism of taurine in microorganisms: a primer in molecular biod-
858 iversity? in *Advances in Experimental Medicine and Biology* (ed. Oja, S. S.) 3–13 (Springer New York,
859 2006). doi:10.1007/978-0-387-33504-9_1.

860 127. Tenenbaum, D. KEGGREST: Client-side REST access to KEGG. R package version 1.29.0. (2020).

861 128. Edgar, R. C. Updating the 97% identity threshold for 16S ribosomal RNA OTUs. *Bioinformatics* **34**,
862 2371–2375 (2018).

863 129. Yarza, P. *et al.* Uniting the classification of cultured and uncultured bacteria and archaea using 16S
864 rRNA gene sequences. *Nat. Rev. Microbiol.* **12**, 635–645 (2014).

865 130. Bodenhofer, U., Kothmeier, A. & Hochreiter, S. Apcluster: an R package for affinity propagation clus-
866 tering. *Bioinformatics* **27**, 2463–2464 (2011).

867 131. Frey, B. J. & Dueck, D. Clustering by passing messages between data points. *Science (80-.).* **315**, 972–
868 976 (2007).

869 132. Kimura, M. Theoretical foundation of population genetics at the molecular level. *Theor. Popul. Biol.*
870 **2**, 174–208 (1971).

871 133. R Core Team. R: A Language and Environment for Statistical Computing. (2021).
872